

RELIABLE CLUSTERING OF BERNOULLI MIXTURE MODELS

BY AMIR NAJAFI^{*}, ABOLFAZL MOTAHARI^{*} AND HAMID R. RABIEE[†]

^{*}*Bioinformatics Research Laboratory (BRL)*

[†]*Digital Media Laboratory (DML)*

*Computer Engineering Department,
Sharif University of Technology, Tehran, Iran*

A Bernoulli Mixture Model (BMM) is a finite mixture of random binary vectors with independent Bernoulli dimensions. The problem of clustering BMM data arises in a variety of real-world applications, ranging from population genetics to activity analysis in social networks. In this paper, we have analyzed the information-theoretic PAC-learnability of BMMs, when the number of clusters is unknown. In particular, we stipulate certain conditions on both sample complexity and the dimension of the model in order to guarantee the Probably Approximately Correct (PAC)-clusterability of a given dataset. To the best of our knowledge, these findings are the first non-asymptotic (PAC) bounds on the sample complexity of learning BMMs.

1. Introduction. Demixing data samples from mixture models, also called model-based clustering, has long been studied by statisticians and computer scientists. Although, plenty of promising algorithms have been introduced in this area, c.f. [1, 2, 3, 4], fewer efforts have been focused on deriving theoretical guarantees on the reliable clustering of data samples. The aim of this paper is to elaborate on this shortcoming by deriving analytic guarantees on the clustering accuracy of a particular case of interest: Bernoulli Mixture Models (BMM).

A Bernoulli Model (BM) refers to a random binary vector of the form $\mathbf{X} = [\mathbf{x}_1, \dots, \mathbf{x}_L] \in \{0, 1\}^L$ with dimension L and independent random components \mathbf{x}_i , where each \mathbf{x}_i is a Bernoulli random variable with probability of success p_i , i.e. $\mathbf{x}_i \sim \text{Bern}(p_i)$. Let us denote the frequency vector as $\mathbf{p} \triangleq [p_1, \dots, p_L] \in [0, 1]^L$. Then, $\mathbb{P}_{\text{BM}}(\mathbf{X}; \mathbf{p})$ denotes a Bernoulli model with

MSC 2010 subject classifications: Primary 68Q32; secondary 62B10

Keywords and phrases: PAC-Learnability, Sample Complexity, Mixture Model Analysis, High-dimensional Statistics

frequency vector \mathbf{p} :

$$\mathbb{P}_{\text{BM}}(\mathbf{X}; \mathbf{p}) \triangleq \prod_{\ell=1}^L p_{\ell}^{X_{\ell}} (1 - p_{\ell})^{1-X_{\ell}}.$$

In this regard, a BMM is defined as a mixture of a finite number of Bernoulli models [5]. Mathematically, it can be expressed as

$$(1.1) \quad \mathbb{P}_{\text{BMM}}(\mathbf{X}; K, \mathbf{p}^{(1)}, \dots, \mathbf{p}^{(K)}, \mathbf{w}) = \sum_{k=1}^K w_k \mathbb{P}_{\text{BM}}(\mathbf{X}; \mathbf{p}^{(k)}),$$

where $K \in \mathbb{N}$ denotes the number of clusters (or mixture components), $\{\mathbf{p}^{(1)}, \dots, \mathbf{p}^{(K)}\}$ is the set of frequency vectors associated to the clusters, and $\mathbf{w} = (w_1, \dots, w_K)$ is the cluster probability vector with $\sum_k w_k = 1$, and $w_k \geq 0$. Let \mathbf{P} to be a $K \times L$ matrix with $\mathbf{p}^{(k)}$ as its k th row. For simplicity, we denote $\mathcal{B} = \mathcal{B}(K, \mathbf{P}, \mathbf{w})$ as the BMM with the above-mentioned specifications. Assume $\mathbf{X}_1, \dots, \mathbf{X}_n \in \{0, 1\}^L$ to be n i.i.d. sample vectors drawn from \mathcal{B} . We define \mathcal{X} as the matrix $[\mathbf{X}_1 | \dots | \mathbf{X}_n]^T \in \{0, 1\}^{n \times L}$ to represent the given dataset.

The problem that we have tackled in this paper is the PAC-clustering of samples in \mathcal{X} , such that clusters with probability at least $1 - \zeta$ are approximately (up to an ϵ fraction of mis-clustering) correct, for arbitrarily small $\epsilon, \zeta > 0$. In order to guarantee the information-theoretic possibility of such clustering, we establish novel lower-bounds in the form of $n \geq \text{poly}(\epsilon^{-1}, \zeta^{-1})$ and $L > \text{poly}(\epsilon^{-1})$, where poly refers to a polynomial function. It should be noted that K , \mathbf{P} and \mathbf{w} are all assumed to be unknown. Statisticians have been studying BMMs for a long time [6, 7, 8, 9]. However, PAC-learnability (or PAC-clusterability)¹ of BMMs in terms of the minimum required sample size and/or model dimension has remained an open problem. To the best of our knowledge, this paper is the first attempt toward this goal by deriving a set of non-asymptotic conditions under which reliable clustering is possible.

The paper is organized as follows: In Section 2, related works are discussed. Section 3 formally presents our main results, where proofs and further discussions are given in Section 4. Finally, conclusions are made in Section 5.

¹In this paper, the notion of PAC-learnability is used in the information-theoretic sense, and to address those learning tasks that can be learned with a polynomial sample complexity w.r.t. ϵ and ζ . This notation is consistent with that of [10]. For those cases where the learning algorithm has also a polynomial computational complexity, the term “efficiently PAC-learnable” has been used.

2. Related Works. Employment of BMMs in order to model multi-dimensional categorical data goes back to [11], while more detailed mathematical and historical explanations can be found, for example, in [12, 13, 14]. In two classic works [9] and [15], a series of heuristic measures have been introduced to assess the number of mixture components in a BMM; However, their performance is validated only through experimental investigations. Authors in these papers have *conjectured* that learnability is possible as long as independence holds between cluster parameters, while their studies lack a rigorous sufficiency analysis. From an algorithmic point of view, Expectation-Maximization (EM) is the most widely used framework for statistical inference in BMMs; See [14] and [16]. In [17], a popular EM-based technique for unsupervised learning of finite mixture models (including BMMs) is introduced, which makes no assumption on the number of mixture components. Also, see [18] for another well-cited paper on model-based clustering of mixture model data. In [19], EM is employed to tune proper initialization for existing inference algorithms on BMMs. From a theoretical perspective, a set of statistical guarantees on the convergence of EM algorithm in mixture model problems has been recently given in [20], however, authors have mainly focused on Gaussian distributions rather than Bernoulli models.

Our work is also related to Bayesian non-parametric approaches in the sense that the number of clusters is open-ended, and will be inferred based on the observed data. Some good reviews on non-parametric approaches in statistics can be found, for example, in [21, 22, 23]. In particular, [24] has proposed a unified non-parametric framework for model-based clustering with the use of hierarchical Dirichlet mixtures. Almost all of the studies reviewed so far share a common property: at their best, authors have only proved convergence to a sub-optimal likelihood value, rather than providing guarantees on the accuracy of the final clustering/learning.

From a geneticist point of view, this paper basically builds upon the statistical model presented in [25]. More discussions on the biological aspects of this model can be found, for example, in [26], [27] and [28]. In [29], authors have performed a simulation study based on [25], in order to assess the number of clusters in a given population. A number of popular softwares for computational population analysis can be found in [28, 30, 31, 32], which mainly focus on binary datasets, the same configuration we have considered in this paper. The problem setting which is tackled by the current work encompasses both models described in [25] and [33], since we have made no restrictive assumptions, such as *independence*, on latent allele frequencies. The role of population stratification in genome-wide association studies

(GWAS), an important application of genetics research, are discussed in [26] and [34]. For more research on the employment of BMMs in GWA studies, see [35, 36, 37, 38].

A critical issue that needs to be discussed here is the following fundamental question: when is a BMM guaranteed to be identifiable? We call a BMM identifiable whenever there is a unique set of parameters $(K, \mathbf{P}, \mathbf{w})$ that corresponds to its probability measure. Reliable clustering of BMMs seems meaningless if there exist more than one true generative models for the samples. Identifiability of BMMs has been addressed in [39], where authors have shown that BMMs cannot be *strictly* identifiable regardless of their dimension, meaning that there always exist some sets of parameters which result in the same probability distribution. However, it does not mean that for every setting $(K, \mathbf{P}, \mathbf{w})$ there must exist another parameter set to produce the same model. Motivated by this idea, in [40] authors have extensively analyzed practical identifiability of BMMs via computer simulations. In [41], it has been proved that for $L \geq 2 \lceil \log_2 K \rceil + 1$, BMMs become *generically* identifiable meaning that sets of parameters with the same probability distributions have a zero Lebesgue measure. According to preceding discussions, we only focus on identifiable cases and this does not reduce the generality of our results as long as the above condition holds.

3. Main Result. In this section, we state our main result and explain its implications and consequences. Recall that the problem is to reliably cluster a set of i.i.d. samples which are drawn from a BMM with unknown parameters (the number of clusters is also assumed to be unknown). The samples are embedded as rows of the matrix \mathcal{X} . In fact, we design an algorithm which outputs a vector $\mathbf{Z} \in \{1, 2, \dots\}^n$ in which the i th element Z_i represents the cluster index of the data sample \mathbf{X}_i . Ultimately, we would like to compare the output of the algorithm with the true clustering which is denoted by $\mathbf{Z}_T \in \{1, 2, \dots, K\}^n$. Let us state two definitions in order to make the comparison mathematically concrete.

DEFINITION 3.1 (ϵ -purity). *A selected row sub-matrix of \mathcal{X} is ϵ -pure if at least $1 - \epsilon$ fraction of its rows have the same index in \mathbf{Z}_T .*

DEFINITION 3.2 (ϵ -correctness). *A clustering algorithm is ϵ -correct on \mathcal{X} if all the output clusters are ϵ -pure.*

It is obvious that the mixture components of the underlying BMM need to be sufficiently far apart from each other, such that a reliable clustering is feasible. For instance, if a BMM contains two mixture components with

exactly the same frequency vectors, no algorithm can index the samples correctly. Therefore, we impose a mild restriction on the parameters of the BMM which makes the clustering a feasible task.

DEFINITION 3.3 ((\mathcal{L}, δ) -separability). *A frequency matrix $\mathbf{P} \in [0, 1]^{K \times L}$ is said to be (\mathcal{L}, δ) -separable, if for each pair of rows of \mathbf{P} , there exist at least $\mathcal{L} \leq L$ columns such that the differences between frequencies exceed δ .*

We may now present our main result in the form of the following theorem which provides a sufficient sample complexity for reliable clustering of BMMs.

THEOREM 3.1 (Non-asymptotic (PAC) Bounds). *Assume $\mathcal{B} = \mathcal{B}(K, \mathbf{P}, \mathbf{w})$ to be a BMM where parameters K , \mathbf{P} and \mathbf{w} are all unknown. However, \mathbf{P} is assumed to be (\mathcal{L}, δ) -separable for some $\mathcal{L} \leq L$ and $\delta > 0$, and there exists $0 < \alpha \leq 1$ such that $w_k \geq \alpha$ for all k . Let $\mathcal{X} = [\mathbf{X}_1 | \mathbf{X}_2 | \dots | \mathbf{X}_n]^T$ to be a dataset including n i.i.d. samples drawn from \mathcal{B} . Assume $\epsilon, \zeta > 0$, such that*

$$\mathcal{L} \geq \frac{B \log^3(1/\epsilon)}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} \quad \text{and} \quad n \geq \frac{C \log^3(1/\epsilon)}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} \log \frac{L}{\zeta},$$

where B and C do not depend on ϵ and ζ . Then, there exists a clustering algorithm $\mathcal{A} : \{0, 1\}^{n \times L} \rightarrow \mathbb{N}^n$, such that \mathcal{A} is ϵ -correct on \mathcal{X} with probability at least $1 - \zeta$.

Proof of Theorem 3.1 with the mathematical formulation of the constants B and C are given in Section 4. Theorem 3.1 is the first result regarding information-theoretic PAC learn-ability of BMMs. It shows the feasibility of the PAC-clustering of samples in \mathcal{X} , such that clusters with probability at least $1 - \zeta$ are ϵ -pure, for arbitrarily small $\epsilon, \zeta > 0$. On the other hand, the imposed lower bounds on sample size n and the number of *informative dimensions* \mathcal{L} are of the form of $n \geq \text{poly}(\epsilon^{-1}, \zeta^{-1})$ and $\mathcal{L} > \text{poly}(\epsilon^{-1})$, respectively.

3.1. Algorithm. The proof of theorem 3.1 is based on an algorithm which employs a novel *pureness check* measure. Called maximal Mutual Dependence Measure (m-MDM), it can reliably test whether a given clustering of the dataset \mathcal{X} includes any non ϵ -pure clusters or not. The core idea for proposing such a measure is the following interesting property of BMMs: In a BMM, unlike a single Bernoulli model, different dimensions of the random binary vector are not statistically independent, and thus can have positive Mutual Information (MI) among each other [14].

DEFINITION 3.4 (Mutual Dependence Measure). Assume $\mathbf{Q} \in \{0, 1\}^{m \times d}$ to be a row/column sub-matrix of \mathcal{X} (with $m \leq n$ and $d \leq L$). The Mutual Dependence Measure (MDM) of \mathbf{Q} , denoted by $\mathcal{D}(\mathbf{Q})$, is defined as

$$(3.1) \quad \mathcal{D}(\mathbf{Q}) \triangleq \mathcal{D}_{\text{KL}}\left(\hat{\mathbb{P}}_{1,\mathbf{Q}} \parallel \hat{\mathbb{P}}_{2,\mathbf{Q}}\right),$$

where $\hat{\mathbb{P}}_{1,\mathbf{Q}}$ denotes the empirical probability distribution of the d -dimensional rows of \mathbf{Q} , while $\hat{\mathbb{P}}_{2,\mathbf{Q}}$ is defined as:

$$(3.2) \quad \hat{\mathbb{P}}_{2,\mathbf{Q}}(\mathbf{X}) \triangleq \prod_{\ell=1}^d \hat{p}_{\ell}^{X_{\ell}} (1 - \hat{p}_{\ell})^{1-X_{\ell}}, \quad \forall \mathbf{X} \in \{0, 1\}^d,$$

with \hat{p}_{ℓ} being the empirical frequency of the ℓ th column of \mathbf{Q} , i.e.

$$\hat{p}_{\ell} \triangleq \frac{1}{n} \sum_{i=1}^n Q_{i,\ell}, \quad \ell = 1, 2, \dots, d.$$

In fact, $\mathcal{D}(\mathbf{Q})$ is the Kullback-Leibler divergence between two distributions obtained under two separate assumptions. Under the first assumption, no restriction is imposed on the origin of the samples and the empirical distribution $\hat{\mathbb{P}}_{1,\mathbf{Q}}$ is simply an estimate of the true underlying distribution of the rows of \mathbf{Q} . Under the second assumption, samples are drawn from a single Bernoulli model and $\hat{\mathbb{P}}_{2,\mathbf{Q}}$ can be used as another estimate of the true distribution. Therefore, if the second assumption does hold, then the two distributions become equal as n goes to infinity. In other words, based on the law of large numbers, we have:

$$\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) = \mathcal{D}_{\text{KL}}\left(\lim_{n \rightarrow \infty} \hat{\mathbb{P}}_{1,\mathbf{Q}} \parallel \lim_{n \rightarrow \infty} \hat{\mathbb{P}}_{2,\mathbf{Q}}\right) = 0.$$

On the other hand, if \mathbf{Q} consists of samples from a BMM with a sufficient level of contributions from different mixture components, then $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ is proved to be strictly positive.

Having defined $\mathcal{D}(\mathbf{Q})$, one needs to move forward and check whether a subset of samples (a row sub-matrix of \mathcal{X}) is ϵ -pure or not. The maximal Mutual Dependence Measure (m-MDM) defined next is a tool to achieve this goal.

DEFINITION 3.5 (maximal Mutual Dependence Measure). The maximal Mutual Dependence Measure (m-MDM) of $\mathbf{Y} \in \{0, 1\}^{m \times L}$, a row sub-matrix of \mathcal{X} , for a length $d \leq L$ is defined as

$$\mathcal{D}_{\text{max}}(\mathbf{Y}; d) \triangleq \max_{\mathbf{Q} \in \text{Col}(\mathbf{Y}; d)} \mathcal{D}(\mathbf{Q}),$$

Algorithm 1 : BMM clustering via Exhaustive Search

 Inputs: Dataset \mathcal{X} , and parameters $(\delta, \epsilon, \alpha)$,

 Set $d \leftarrow \frac{1-\alpha}{\alpha\delta^2(1-\epsilon)} (1 + \log \frac{1}{\alpha\epsilon})$ (Sub-matrix column size)

 Set $\tau \leftarrow \frac{\epsilon}{2} (1 + \log \frac{1}{\alpha\epsilon})$ (Purity test threshold)

 Set $\kappa \leftarrow 1$ (Cluster number)

```

while  $\kappa < \lceil \frac{1}{\alpha} \rceil$  do
  for  $\forall \mathbf{Z} \in \{1, \dots, \kappa\}^n$ , where the size of each cluster is more than  $\alpha n/2$  do
    Set  $\mathbf{Y}_1 \dots, \mathbf{Y}_\kappa \leftarrow$  The clustered row sub-matrices of  $\mathcal{X}$  based on  $\mathbf{Z}$ .
    if  $\mathcal{D}_{\max}(\mathbf{Y}_k; d) \leq \tau$  for  $\forall k = 1, \dots, \kappa$  then
      Set  $\mathbf{Z}^* \leftarrow \mathbf{Z}$ , and
      Terminate the program.
    end if
  end for
  Set  $\kappa \leftarrow \kappa + 1$ 
end while

```

 Output: \mathbf{Z}^* , a clustering of data in \mathcal{X} .

where maximization is taken over $\text{Col}(\mathbf{Y}; d)$ which consists of all $\binom{L}{d}$ column sub-matrices of \mathbf{Y} with size $m \times d$.

The m-MDM measure is the main tool used in our proposed clustering strategy which is presented in Algorithm 1. In fact, Algorithm 1 works by searching over all possible clusterings of the dataset \mathcal{X} , which have the following two properties: (i) the number of clusters does not exceed $\lceil \frac{1}{\alpha} \rceil$, where α is a lower-bound on the probability of the smallest cluster in \mathcal{B} ; and (ii) the smallest cluster has at least $\alpha n/2$ members. We start with a single cluster and then increase the number of clusters one by one. Given the conditions of Theorem 3.1, the true clustering \mathbf{Z}_T would be in this search space with probability at least $1 - \zeta/3$.

For any given clustering, we check to see whether all the corresponding clusters are ϵ -pure or not. We can do this by evaluating the m-MDM over each clustered row sub-matrix of \mathcal{X} . If \mathcal{D}_{\max} has negligible values (smaller than a pre-defined threshold τ) in all the clusters, then the current clustering is accepted and the program terminates. We show that under the constraints of Theorem 3.1, the probability of accepting a clustering with even one non ϵ -pure cluster is less than $\zeta/3$. On the other hand, the algorithm eventually reaches the true clustering \mathbf{Z}_T by assuming that we have not accepted any other candidates up to that point. Again, we show that the probability of rejecting the true clustering \mathbf{Z}_T is no more than $\zeta/3$.

Therefore, one can deduce that with probability at least $1 - \zeta$, Algorithm 1

either outputs an ϵ -correct clustering on \mathcal{X} or the true clustering \mathbf{Z}_T (which is of course ϵ -correct as well). The key property in our analysis that has made it possible to draw the above-mentioned strategy is that our proposed m-MDM measure can detect the impurity of data clusters with a decision error which decays exponentially w.r.t. both n and \mathcal{L} .

3.2. Discussions. The m-MDM and its parent MDM are very powerful in differentiating between pure and non-pure group of samples. In this section, we elaborate on this fact. Assume a random vector $\mathbf{X} \in \{0, 1\}^L$ with $\mathbf{X} \sim \mathcal{B}(K, \mathbf{P}, \mathbf{w})$. If we condition on \mathbf{X} to be drawn from a particular mixture component of \mathcal{B} , say the k th one with $k \in \{1, 2, \dots, K\}$, then the probability distribution of \mathbf{X} would be

$$(3.3) \quad \mathbb{P}(\mathbf{X}|k) = \prod_{\ell=1}^L \mathbb{P}(X_{\ell}|k).$$

However, according to (1.1), the distribution of \mathbf{X} without this assumption is $\mathbb{P}(\mathbf{X}) = \sum_k w_k \mathbb{P}(\mathbf{X}|k)$. A more subtle comparison of $\mathbb{P}(\mathbf{X})$ and $\mathbb{P}(\mathbf{X}|k)$ simply reveals that in a mixture model, unlike the case of a single Bernoulli model, different dimensions of the vector are not necessarily independent from each other. This argument can be qualitatively justified as follows: a group of observed dimensions can convey information about the origin of the whole vector \mathbf{X} , which then impacts the distribution of any other group of dimensions. However, the above-mentioned flow of information does not occur when \mathbf{X} is known to be generated from a single Bernoulli model.

Based on the above argument, we have introduced the MDM in Definition 3.4, in order to quantify whether a selected row/column subset of samples in \mathcal{X} are more likely to be drawn from a single model, or a mixture of various models with different parameters.

When n is finite, MDM only works on a relatively small $m \times d$ sub-matrix of \mathcal{X} . This is due to the fact that the number of data samples m which are required to make reliable assessments regarding the ϵ -purity of the sub-matrix grows exponentially with respect to d (see Lemmas 4.1 and 4.2 in the next section). This fact should not be surprising since reliable computation of MDM is subject to having a relatively close estimation of a d -dimensional binary distribution. Hence large values of d are unsuitable for estimating $\mathcal{D}(\cdot)$. On the other hand, by choosing a small d , we are ignoring a huge amount of valuable information in the dataset.

To exploit all the information embedded in \mathcal{X} , the m-MDM is introduced. It computes numerous MDMs over various subsets of dimensions, and aggregates all these values to form a more informative measure.

4. Proof of Theorem 3.1. We first start by some lemmas which indicate the goodness of our proposed *purity check* measures. The following lemma shows that $\mathcal{D}(\mathbf{Q})$ deviates from zero with high probability whenever \mathbf{Q} is generated by a BMM with $K \geq 2$ mixture components.

LEMMA 4.1. *Assume \mathcal{B} to be a BMM with K clusters, d Bernoulli trials per sample, frequency matrix of $\mathbf{P} \in [0, 1]^{K \times d}$ and cluster probability vector of $\mathbf{w} = (w_1, w_2, \dots, w_K)$. Let \mathbf{P} to be (\mathcal{L}, δ) -separable for some $\mathcal{L} \leq d$ and $\delta > 0$. Also, assume $w_k \leq 1 - \epsilon, \forall k = 1, \dots, K$ and some $\epsilon > 0$. Consider $\mathbf{Q}_1, \dots, \mathbf{Q}_n$ to be n i.i.d. samples drawn from \mathcal{B} , and let $\mathbf{Q} = [\mathbf{Q}_1 | \dots | \mathbf{Q}_n]^T \in \{0, 1\}^{n \times d}$. Then, if $\mathcal{L} > \frac{1 + \log \frac{K}{\epsilon}}{(1 - \epsilon)\delta^2}$, we have*

$$\mathbb{P}\{\mathcal{D}(\mathbf{Q}) \leq \tau\} \leq 2^{d+1} e^{-\beta n},$$

where $\tau \triangleq \frac{\epsilon}{2} \left(1 + \log \frac{K}{\epsilon}\right)$ and $\beta \triangleq \frac{\tau^2}{d^{2d+1}}$.

Proof of Lemma 4.1 is given in Appendix A. The assumption of $w_k \leq 1 - \epsilon$ for all k yields that for $n \rightarrow \infty$, the set of observations $\mathbf{Q}_1, \dots, \mathbf{Q}_n$ is not ϵ -pure, almost surely. Hence, for an asymptotically large non ϵ -pure set of observations, we have $\mathcal{D}(\mathbf{Q}) \stackrel{a.s.}{\geq} \tau$. On the other hand, we have already discussed that for a completely pure set, i.e. when samples are drawn from a single Bernoulli model ($K = 1$), we have $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \stackrel{a.s.}{=} 0$. The following lemma will provide concentration bounds on the deviation of $\mathcal{D}(\mathbf{Q})$ when rows of \mathbf{Q} are drawn from a single Bernoulli model.

LEMMA 4.2. *Assume \mathcal{B} to be a single Bernoulli model ($K = 1$) with d Bernoulli trials per samples and an arbitrary frequency vector. Consider $\mathbf{Q}_1, \dots, \mathbf{Q}_n$ to be n i.i.d. samples drawn from \mathcal{B} , and let $\mathbf{Q} = [\mathbf{Q}_1 | \dots | \mathbf{Q}_n]^T$. Then, we have*

$$\mathbb{P}\{\mathcal{D}(\mathbf{Q}) \geq \tau\} \leq 2^{d+1} e^{-\beta d^2 n},$$

where τ and β are the same as in Lemma 4.1.

The proof for Lemma 4.2 is also given in Appendix A. And finally, the following lemma shows that the error rate in detecting improper clustering of samples, i.e. any clustering with at least one non ϵ -pure cluster, drops exponentially with respect to both n and \mathcal{L} .

LEMMA 4.3. *Assume \mathcal{B} to be a BMM with K clusters, L dimensions, \mathbf{P} as the frequency matrix and $\mathbf{w} = \{w_1, \dots, w_K\}$ as the cluster probability vector. Consider $\mathbf{Y}_1, \dots, \mathbf{Y}_n$ to be n i.i.d. samples drawn from \mathcal{B} and let*

$\mathbf{Y} = [\mathbf{Y}_1 | \dots | \mathbf{Y}_n]^T \in \{0, 1\}^{n \times L}$. For $K \geq 2$, assume \mathbf{P} to be (\mathcal{L}, δ) -separable for some $\mathcal{L} \leq L$ and $\delta > 0$. Also, assume $w_k \leq 1 - \epsilon$ for some $\epsilon > 0$ and all k . Assume $\mathcal{L} > d \triangleq \frac{K-1}{(1-\epsilon)\delta^2} (1 + \log \frac{K}{\epsilon})$ and let $\tau \triangleq \frac{\epsilon}{2} (1 + \log \frac{K}{\epsilon})$. Then,

$$\mathbb{P} \{ \mathcal{D}_{\max}(\mathbf{Y}; d) \leq \tau \} \leq 4^{\mathcal{L}} \exp \left(\frac{-\tau^2 n \mathcal{L}}{d^5 2^{d+1}} \right).$$

On the other hand, when $K = 1$

$$\mathbb{P} \{ \mathcal{D}_{\max}(\mathbf{Y}; d) \geq \tau \} \leq \binom{L}{d} 2^{d+1} \exp \left(\frac{-\tau^2 n}{d^2 2^{d+1}} \right).$$

Based on Lemma 4.3, for a BMM with sufficiently large n and \mathcal{L} , the probability of mis-detection between an ϵ -pure subset of samples in the dataset \mathcal{X} and a non ϵ -pure one is strictly bounded. In fact, Lemma 4.3 provides a mathematically rigor and reliable criterion to distinguish between “good” and “bad” clusterings of samples in a finite dataset.

The parameter α is user-defined. As long as for a BMM $\mathcal{B} = \mathcal{B}(K, \mathbf{p}, \mathbf{w})$ we have $\min_k w_k \geq \alpha$, K cannot not exceed $\lceil \frac{1}{\alpha} \rceil$. Given that the conditions in Theorem 3.1 are satisfied, with probability at least $1 - \zeta$ Algorithm 1 terminates before passing $\lceil \frac{1}{\alpha} \rceil$ clusters and as soon as it finds an ϵ -correct clustering of \mathbf{X} . Otherwise, the algorithm just outputs a null clustering. In the following, we use results from Lemma 4.3 to prove Theorem 3.1, which is also the mathematical analysis of Algorithm 1.

PROOF OF THEOREM 3.1. Algorithm 1 checks all possible cluster numbers κ (starting from $\kappa = 1$). Let us denote the number of clusterings that need to be checked before reaching the correct partitioning, i.e. \mathbf{Z}_T , by N . Then, obviously N satisfies the following inequality:

$$C \leq 1^n + 2^n + \dots + K^n \leq K^{n+1}.$$

In this regard, one can consider the following error events during the execution of Algorithm 1:

- \mathcal{E}_1 : Accepting a non ϵ -correct clustering of dataset \mathbf{X} , before reaching the true clustering \mathbf{Z}_T .
- \mathcal{E}_2 : Eventually reaching to the true clustering \mathbf{Z}_T , and denying it.
- \mathcal{E}_3 : The smallest true cluster in the dataset \mathbf{X} has less than $\alpha n/2$ members.

Obviously, probability of the algorithm failure, denoted by P_E , can be upper-bounded as

$$P_E = \mathbb{P} \{ \mathcal{E}_1 \cup \mathcal{E}_2 \cup \mathcal{E}_3 \} \leq \mathbb{P} \{ \mathcal{E}_1 \} + \mathbb{P} \{ \mathcal{E}_2 \} + \mathbb{P} \{ \mathcal{E}_3 \}.$$

In the following, we show that given the condition of Theorem 3.1, we have $\mathbb{P}\{\mathcal{E}_i\} \leq \zeta/3$ for $i = 1, 2, 3$.

From Lemma 4.3, we know that both probabilities of accepting a non ϵ -pure partitioning, and rejecting the correct one are bounded and decrease exponentially w.r.t. n . Recalling

$$d \triangleq \frac{1-\alpha}{\alpha\delta^2(1-\epsilon)} \left(1 + \log \frac{1}{\alpha\epsilon}\right) = O\left(\log \frac{1}{\epsilon}\right)$$

and

$$\tau \triangleq \frac{\epsilon}{2} \left(1 + \log \frac{1}{\alpha\epsilon}\right) = O\left(\epsilon \log \frac{1}{\epsilon}\right)$$

from Algorithm 1, and by using the union bound over all non ϵ -pure clusterings in the first N steps of the algorithm, one can see

$$\begin{aligned} \mathbb{P}\{\mathcal{E}_1\} &\leq N \mathbb{P}\{\mathcal{E}_1^{(1)}\} \leq K^{n+1} 4^{\mathcal{L}} \exp\left(\frac{-\alpha\tau^2 n \mathcal{L}}{d^5 2^{d+2}}\right) \\ (4.1) \quad &= \exp\left(O(n) + O(\mathcal{L}) - n \mathcal{L} \cdot O\left(\frac{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}}{\log^3(1/\epsilon)}\right)\right), \end{aligned}$$

where $\mathcal{E}_1^{(1)}$ represents the error event corresponding to the acceptance of a single non ϵ -pure clustering. The last inequality in (4.1) is a direct consequence of Lemma 4.3, considering the fact that for a non ϵ -correct clustering at least one cluster must not be ϵ -pure. We have also used the fact that each cluster has at least $\alpha n/2$ samples. From (4.1), one can deduce that by choosing

$$(4.2) \quad \mathcal{L} \geq \frac{B \log^3(1/\epsilon)}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} \quad , \quad n \geq n_{\min}^{(1)} \triangleq C^{(1)} \left(\frac{\log^3(1/\epsilon)}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} + \log \frac{1}{\zeta}\right),$$

where coefficients B and $C^{(1)}$ do not depend on ϵ or ζ , we can achieve $\mathbb{P}\{\mathcal{E}_1\} \leq \zeta/3$ for arbitrary small $\epsilon, \zeta > 0$.

A similar argument can be used to obtain an upper-bound on $\mathbb{P}\{\mathcal{E}_2\}$. It should be noted that for \mathcal{E}_2 to occur, at least one of the true clusters in \mathbf{Z}_T must have $\mathcal{D}_{\max} > \tau$. Since the number of clusters at that step of the algorithm is K , one can use the union bound over all K clusters each of which has at least $\alpha n/2$ members. Also, we aim to use the following inequality:

$$\log \binom{L}{d} \leq d \log \frac{Le}{d}.$$

In this regard, by using the second inequality in Lemma 4.3, it can be shown that

$$\begin{aligned}
 \mathbb{P}\{\mathcal{E}_2\} &\leq K \binom{L}{d} 2^{d+1} \exp\left(\frac{-\alpha\tau^2 n}{d^2 2^{d+2}}\right) \\
 (4.3) \quad &= \exp\left(O\left(\log \frac{1}{\epsilon} \left[1 + \log \frac{L}{\log(1/\epsilon)}\right]\right) - nO\left(\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}\right)\right).
 \end{aligned}$$

Thus, by choosing

$$(4.4) \quad n \geq n_{\min}^{(2)} \triangleq C^{(2)} \left(\frac{\log(1/\epsilon) \log L}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} + \log \frac{1}{\zeta} \right)$$

with $C^{(2)}$ being a coefficient that does not depend on ϵ or ζ , one can simply guarantee that $\mathbb{P}\{\mathcal{E}_2\} \leq \zeta/3$. Finally, according to Lemma A.3 in Appendix A, by choosing $n \geq n_{\min}^{(3)}$ which is defined as

$$n_{\min}^{(3)} \triangleq C^{(3)} \log \frac{1}{\zeta},$$

where again coefficient $C^{(3)}$ is independent of ϵ or ζ , one can guarantee that the probability of occurring \mathcal{E}_3 is less than $\zeta/3$.

Therefore, by aggregating the results by choosing \mathcal{L} to satisfy the inequality of (4.2), and the sample size n to be lower-bounded by

$$n \geq \frac{C \log^3(1/\epsilon)}{\epsilon^{2+\frac{1-\alpha}{\alpha\delta^2}}} \log \frac{L}{\zeta} \geq \max \left\{ n_{\min}^{(1)}, n_{\min}^{(2)}, n_{\min}^{(3)} \right\}$$

for some constant C , Algorithm 1 is guaranteed to output an ϵ -correct clustering on dataset \mathcal{X} with probability at least $1 - \zeta$, which completes the proof. \square

In an asymptotic regime, when the overall dimension L becomes very large while the number of informative dimensions \mathcal{L} is fixed, the minimum required dataset size should grow logarithmically w.r.t. L . This analytic observation makes sense since in finite n regimes, addition of more *non-informative* dimensions, i.e. those dimensions that have the same frequency values between all or at least some of the clusters only adds extra noise to the dataset \mathcal{X} which makes any decision-making strategy to be more challenging.

5. Conclusions. This paper aims to find the first non-asymptotic PAC bounds on the sample complexity of reliable clustering in BMMs, when the number of clusters is unknown. In order to achieve this goal, a novel statistic called maximal Mutual Dependence Measure (m-MDM) is proposed which can be computed over any selected subset of samples in the training dataset. It has been proved that m-MDM is capable of finding a reliable clustering with an arbitrary small error as long as sample complexity n and the number of informative dimensions \mathcal{L} are polynomially large w.r.t. error values. No restrictive assumptions have been made in our model, except those required for the identifiability and/or learnability of the task, such as: existence of sufficient deviations among the frequency vectors of different mixture components, and lower-bounding the size of the smallest cluster in the model. As a result, our findings encapsulate many classes of BMM inference problems. The proposed Algorithm 1 is NP, which means efficient PAC-learnability of BMMs still remains an open problem. Looking for polynomial-time algorithms for this problem should be the focus of future researches in this area. Also, deriving converse conditions for Theorem 3.1 is of paramount interest.

APPENDIX A: AUXILIARY LEMMAS AND PROOFS

PROOF OF LEMMA 4.1. Proof consists of two consecutive parts. First, we show that $\mathcal{D}(\mathbf{Q})$ is lower-bounded by a positive value in the asymptotic case, i.e.

$$(A.1) \quad \mathbb{P} \left\{ \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \leq 2\tau \right\} = 0.$$

Second, we prove that the probability of the event $|\mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})| > \tau$ decays exponentially w.r.t. n , which would complete the proof.

For the sake of simplicity in notations, let $\mathbb{P}_{\mathbf{p}}$ to represent the probability distribution of a Bernoulli model with frequency vector $\mathbf{p} \in [0, 1]^d$ in this proof. This way, one can write

$$(A.2) \quad \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) = \mathcal{D}_{\text{KL}} \left(\sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}} \left\| \mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}} \right. \right),$$

where $\mathbf{p}^{(k)}$ denotes the k th row of frequency matrix \mathbf{P} , and $\mathbb{E}_k \mathbf{p}^{(k)} \triangleq \sum_k w_k \mathbf{p}^{(k)}$. Here, $\sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}}$ indicates a mixture of Bernoulli models (a BMM), while $\mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}}$ denotes a single Bernoulli model with a frequency vector which is a weighted average of the K frequency vectors in \mathbf{P} .

It can be verified that when \mathbf{w} is a one-hot vector, i.e. one component is 1 and the rest are 0, then the two probability distributions $\sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}}$

and $\mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}}$ become equal and $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \stackrel{a.s.}{=} 0$. However, if for some $\epsilon > 0$ we have $w_k \leq 1 - \epsilon$, $\forall k$, and \mathcal{L} is *sufficiently* large, we can prove that $\sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}}$ may not be consistently approximated by a single Bernoulli model whose frequency vector is $\mathbb{E}_k \mathbf{p}^{(k)}$.

Based to the definition of the Kullback-Liebler divergence, r.h.s. of (A.2) can be expanded as follows which helps us to find a proper lower-bound for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$:

$$\begin{aligned}
 \mathcal{D}_{\text{KL}} \left(\sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}} \middle\| \mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}} \right) &= \sum_{\forall \mathbf{X}} \sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}}(\mathbf{X}) \log \left(\frac{\sum_u w_u \mathbb{P}_{\mathbf{p}^{(u)}}(\mathbf{X})}{\mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}}(\mathbf{X})} \right) \\
 &\geq \sum_{\forall \mathbf{X}} \sum_k w_k \mathbb{P}_{\mathbf{p}^{(k)}}(\mathbf{X}) \log \left(\frac{w_k \mathbb{P}_{\mathbf{p}^{(k)}}(\mathbf{X})}{\mathbb{P}_{\mathbb{E}_k \mathbf{p}^{(k)}}(\mathbf{X})} \right) \\
 &= \sum_k w_k \sum_{\ell=1}^L \left(\sum_{X_\ell \in \{0,1\}} \mathbb{P}_{p_\ell^{(k)}}(X_\ell) \log \left(\frac{\mathbb{P}_{p_\ell^{(k)}}(X_\ell)}{\mathbb{P}_{\mathbb{E}_k p_\ell^{(k)}}(X_\ell)} \right) \right) - \mathbb{H}(\mathbf{w}) \\
 (A.3) \quad &= \sum_{\ell=1}^L \sum_{k=1}^K w_k \mathcal{D}_{\text{KL}} \left(\mathbb{P}_{p_\ell^{(k)}} \middle\| \mathbb{P}_{\mathbb{E}_k p_\ell^{(k)}} \right) - \mathbb{H}(\mathbf{w}),
 \end{aligned}$$

where $\mathbb{H}(\mathbf{w}) \triangleq -\sum_k w_k \log w_k$ denotes the Shannon entropy of the discrete distribution \mathbf{w} , and summation over \mathbf{X} refers to $\forall \mathbf{X} \in \{0,1\}^d$. Moreover, it is easy to show that

$$\sum_{k=1}^K w_k \mathcal{D}_{\text{KL}} \left(\mathbb{P}_{p_\ell^{(k)}} \middle\| \mathbb{P}_{\mathbb{E}_k p_\ell^{(k)}} \right) = H \left(\sum_{k=1}^K w_k p_\ell^{(k)} \right) - \sum_{k=1}^K w_k H(p_\ell^{(k)}).$$

Here, and for the ease of notations $H(p)$ for $0 \leq p \leq 1$ refers to $H(p) \triangleq \mathbb{H}(\text{Bern}(p)) = -p \log p - (1-p) \log(1-p)$. Since $H(\cdot)$ is a strictly concave function, and considering the fact that $\mathbb{H}(\mathbf{w})$ is always upper-bounded by $\log K$ regardless of d , one can conclude that the lower-bound for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ in (A.3) becomes strictly positive when i) d is sufficiently large, and ii) frequency vectors $\mathbf{p}^{(k)}$ for $k = 1, \dots, K$ are sufficiently far apart from each other.

In order to simplify the lower-bound in (A.3), let us assume a random variable $\mathbf{A} \in [0,1]$, and define $\mathbf{a} \triangleq \mathbf{A} - \mathbb{E}\mathbf{A}$. According to the Taylor's series expansion theorem one can write

$$\begin{aligned}
 H(\mathbb{E}\mathbf{A}) - \mathbb{E}H(\mathbf{A}) &= H(\mathbb{E}\mathbf{A}) - \mathbb{E} \left\{ H(\mathbb{E}\mathbf{A}) + H'(\mathbb{E}\mathbf{A}) \mathbf{a} + \frac{1}{2} H''(\mathbb{E}\mathbf{A} + \xi) \mathbf{a}^2 \right\} \\
 &\geq \frac{\mathbb{E}\mathbf{a}^2}{2} \inf_{0 \leq p \leq 1} |H''(p)| = \inf_{0 \leq p \leq 1} \frac{\mathbb{E}\mathbf{a}^2}{2p(1-p)} = 2\mathbb{E}\mathbf{a}^2,
 \end{aligned}$$

where ξ is a random variable depending on \mathbf{A} [42]. Using this inequality, the lower-bound for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ can be written as

$$(A.4) \quad \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \geq 2 \sum_{\ell=1}^d \sum_{k=1}^K w_k \left(p_{\ell}^{(k)} - \sum_{u=1}^K w_u p_{\ell}^{(u)} \right)^2 - \mathbb{H}(\mathbf{w}).$$

We have already assumed that $w_k \leq 1 - \epsilon, \forall k$. Also, due to the (\mathcal{L}, δ) -separability assumption, for all pairs of rows in \mathbf{P} , say i and j , there exists a subset of columns $\mathcal{C}_{i,j} \subseteq \{1, 2, \dots, L\}$ where

$$\left| p_{\ell}^{(i)} - p_{\ell}^{(j)} \right| \geq \delta, \quad \ell \in \mathcal{C}_{i,j},$$

and $|\mathcal{C}_{i,j}| \geq \lceil \cdot \rceil$. Lemma A.1 shows that the lower-bound in (A.4) can be further simplified as

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) &\geq 2\mathcal{L}\epsilon(1-\epsilon)\delta^2 - \mathbb{H}\left(\left[1-\epsilon, \frac{\epsilon}{K-1}, \dots, \frac{\epsilon}{K-1}\right]\right) \\ &\geq 2\epsilon(1-\epsilon)\delta^2 \left(\mathcal{L} - \frac{1 + \log \frac{K}{\epsilon}}{2(1-\epsilon)\delta^2} \right), \end{aligned}$$

where by choosing \mathcal{L} to be greater or equal to $\frac{1 + \log \frac{K}{\epsilon}}{(1-\epsilon)\delta^2}$, τ becomes strictly greater than 2τ .

So far, we have shown that $\mathcal{D}(\mathbf{Q})$ becomes greater than 2τ when n goes to infinity. However, $\mathcal{D}(\mathbf{Q})$ is supposed to be computed over a finite sample size of n , thus it is necessary to show that $|\mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})|$ concentrates around zero w.r.t. n . In fact, Lemma A.2 proves that the above error decays exponentially with respect to n . Based on this result, the probability $\mathbb{P}\{\mathcal{D}(\mathbf{Q}) \leq \tau\}$ can be upper-bounded by

$$\begin{aligned} \mathbb{P}\{\mathcal{D}(\mathbf{Q}) \leq \tau\} &\leq 2^{L+1} \exp\left(\frac{-n\epsilon^2(1-\epsilon)^2\delta^4}{L^4 2^{L+1}} \left(\mathcal{L} - \frac{1 + \log \frac{K}{\epsilon}}{2(1-\epsilon)\delta^2}\right)^2\right) \\ &= 2^{d+1} \exp\left(\frac{-n\tau^2}{d^4 2^{d+1}}\right), \end{aligned}$$

which completes the proof. \square

LEMMA A.1. *The lower-bound for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ given in (A.4), subject to $w_k \leq 1 - \epsilon, \forall k$ is as follows:*

$$\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \geq 2\epsilon(1-\epsilon)\delta^2 \left(\mathcal{L} - \frac{1 + \log \frac{K}{\epsilon}}{2(1-\epsilon)\delta^2} \right).$$

PROOF. Based on the assumption made in Lemma 4.1 with respect to the non ϵ -purity of \mathbf{Q} , let us define a subset of \mathbb{R}^K denoted by $\Phi(x)$ for $\epsilon \leq x \leq 1 - 1/K$ as

$$\mathbf{W}(x) \triangleq \left\{ \mathbf{w} \in \mathbb{R}^K \mid \forall w_k \geq 0, \sum_k w_k = 1, \max_k w_k = 1 - x \right\}.$$

Hence, according to (A.4) the lower-bound for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ (for a non ϵ -pure \mathbf{Q}) can be written as

(A.5)

$$\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \geq \inf_{\epsilon \leq x \leq 1 - \frac{1}{K}} \left\{ \inf_{\mathbf{w} \in \mathbf{W}(x)} 2 \sum_{\ell=1}^d \sum_{k=1}^K w_k \left(p_{\ell}^{(k)} - \mu_{\ell}(\mathbf{w}) \right)^2 - \mathbb{H}(\mathbf{w}) \right\},$$

with $\mu_{\ell}(\mathbf{w}) \triangleq \sum_{u=1}^K w_u p_{\ell}^{(u)}$. In fact, (A.5) indicates minimization of the lower-bound over all asymptotically large non ϵ -pure matrices \mathbf{Q} . In order to further simplify the problem, minimization over $\mathbf{w} \in \mathbf{W}(x)$ can be carried out for the two terms in the r.h.s. of (A.5), separately. In this regard, it is easy to see that

(A.6)

$$\sup_{\mathbf{w} \in \mathbf{W}(x)} \mathbb{H}(\mathbf{w}) = (1-x) \log \frac{1}{1-x} + \sum_{k=2}^K \frac{x}{K-1} \log \frac{K-1}{x} \leq x \left(1 + \log \frac{K}{x} \right).$$

And for the first term in r.h.s of (A.5), the following lower-bound can be obtained:

$$\begin{aligned} & \inf_{\mathbf{w} \in \mathbf{W}(x)} 2 \sum_{\ell=1}^d \sum_{k=1}^K w_k \left(p_{\ell}^{(k)} - \mu_{\ell}(\mathbf{w}) \right)^2 \\ & \geq \inf_{\mathbf{w} \in \mathbf{W}(x)} 2 \sum_{\ell=1}^d \inf_{\forall \mu_{\ell} \in \mathbb{R}} \left\{ \sum_{k=1}^K w_k \left(p_{\ell}^{(k)} - \mu_{\ell} \right)^2 \right\} \\ & \geq 2 \min_{k_0=1, \dots, K} \sum_{\ell=1}^d \inf_{\mu_{\ell} \in \mathbb{R}} \left\{ (1-x) \left(p_{\ell}^{(k_0)} - \mu_{\ell} \right)^2 + \min_{k|k \neq k_0} x \left(p_{\ell}^{(k)} - \mu_{\ell} \right)^2 \right\} \\ & \geq 2 \min_{k_0=1, \dots, K} \min_{k|k \neq k_0} \sum_{\ell=1}^d x (1-x) \left(p_{\ell}^{(k)} - p_{\ell}^{(k_0)} \right)^2. \end{aligned}$$

Since for every pair (k, k_0) and at least \mathcal{L} dimensions out of $\ell = 1, 2, \dots, d$, the inequality $\left| p_{\ell}^{(k)} - p_{\ell}^{(k_0)} \right| \geq \delta$ holds, one can write

$$(A.7) \quad \inf_{\mathbf{w} \in \mathbf{W}(x)} 2 \sum_{\ell=1}^d \sum_{k=1}^K w_k \left(p_{\ell}^{(k)} - \mu_{\ell}(\mathbf{w}) \right)^2 \geq 2\mathcal{L}x(1-x)\delta^2.$$

Combining (A.6) with (A.7), the following lower-bound can be achieved for $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$, which is in terms of the free parameter $\epsilon \leq x \leq 1 - 1/K$:

$$\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \geq \inf_{\epsilon \leq x \leq 1 - \frac{1}{K}} 2x(1-x)\delta^2 \left(\mathcal{L} - \frac{1 + \log \frac{K}{x}}{2(1-x)\delta^2} \right).$$

Taking derivatives w.r.t. x from the above lower-bound, it can be easily verified that for $\mathcal{L} > \frac{1 + \log \frac{K}{\epsilon}}{2(1-\epsilon)\delta^2}$ it remains a monotonically increasing function w.r.t. x . Hence, its minimum occurs for $x = \epsilon$, which completes the proof. \square

LEMMA A.2. *The estimation error in computing $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q})$ of Lemma 4.1 over a finite sample size of n can be upper-bounded as*

$$\mathcal{P} \left\{ \left| \mathcal{D}(\mathbf{Y}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Y}) \right| > \epsilon \right\} \leq 2^{d+1} \exp \left(\frac{-n\epsilon^2}{d^4 2^{d+1}} \right).$$

PROOF. For the ease of notation, let us define $g(\hat{\mathbb{P}}_{\mathbf{Q}}) \triangleq \mathcal{D}(\mathbf{Q})$, where $\hat{\mathbb{P}}_{\mathbf{Q}}$ is a 2^d -dimensional random variable which represents the empirical distribution of the rows of the random matrix \mathbf{Q} , and $g : \mathbb{R}^{2^d} \rightarrow \mathbb{R}^+$ to be a corresponding function that maps $\hat{\mathbb{P}}_{\mathbf{Q}}$ to $\mathcal{D}(\mathbf{Q})$. According to Definition 3.4, it can be readily verified that

$$\begin{aligned} \mathcal{D}(\mathbf{Q}) &= \sum_{\forall \mathbf{X}} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) \log \left(\frac{\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X})}{\prod_{\ell=1}^d \hat{p}_{\ell}^{X_{\ell}} (1 - \hat{p}_{\ell})^{1-X_{\ell}}} \right) \\ &= \sum_{\forall \mathbf{X}} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) \left[\log \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \sum_{\ell=1}^d \mathbf{1}_{X_{\ell}} \log \hat{p}_{\ell} + \mathbf{1}_{1-X_{\ell}} \log (1 - \hat{p}_{\ell}) \right] \\ \text{(A.8)} \quad &= \sum_{\forall \mathbf{X}} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) \left[\log \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \sum_{\ell=1}^d \log \left(\sum_{\forall \mathbf{X}' | X'_{\ell} = X_{\ell}} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}') \right) \right], \end{aligned}$$

where $\mathbf{1}_X$ denotes the indicator function which returns 1 if $X = 1$ and zero otherwise. In derivation of (A.8), we have used the facts that

$$\hat{p}_{\ell} = \sum_{\forall \mathbf{X}' | X'_{\ell} = 1} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}') \quad \text{and} \quad 1 - \hat{p}_{\ell} = \sum_{\forall \mathbf{X}' | X'_{\ell} = 0} \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}').$$

Since g is a continuous function, the law of large numbers implies that

$$\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) = g \left(\lim_{n \rightarrow \infty} \hat{\mathbb{P}}_{\mathbf{Q}} \right) = g(\mathbb{P}_{\text{BMM}}),$$

where \mathbb{P}_{BMM} represents the true distribution of the BMM \mathcal{B} , which generates the rows of \mathbf{Q} . Obviously, unlike the empirical measure $\hat{\mathbb{P}}_{\mathbf{Q}}$, \mathbb{P}_{BMM} is a deterministic distribution which can be quantified given that parameters of \mathcal{B} are known. In this regard, using differential calculus implies the following equalities:

$$\left| \mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \right| = \left| g(\hat{\mathbb{P}}_{\mathbf{Q}}) - g(\mathbb{P}_{\text{BMM}}) \right| = \left| \int_{\mathcal{P}} \nabla g \cdot d\mathcal{P} \right|,$$

where ∇ denotes the gradient operator, and \mathcal{P} is any continuous path in \mathbb{R}^{2^d} that starts from $\hat{\mathbb{P}}_{\mathbf{Q}}$ and ends in \mathbb{P}_{BMM} . This way, by employing and the *Mean Value Theorem (MVT)*, one can write:

$$\begin{aligned} \left| \mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \right| &\leq \sum_{\mathbf{X} \in \{0,1\}^d} \left| \int_{\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X})}^{\mathbb{P}_{\text{BMM}}(\mathbf{X})} \nabla_{\mathbf{X}} g \right| \\ &\stackrel{(\text{MVT})}{\leq} \sum_{\forall \mathbf{X}} \left(\sup_{x \in \mathcal{N}(\mathbf{X})} |\nabla_{\mathbf{X}} g(x)| \right) \left| \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \mathbb{P}_{\text{BMM}}(\mathbf{X}) \right|, \end{aligned}$$

where $0 \leq x \leq 1$ denotes the dummy variable representing the probability of vector \mathbf{X} , and $\mathcal{N}(\mathbf{X})$ the interval between $\mathbb{P}_{\text{BMM}}(\mathbf{X})$ and $\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X})$. According to (A.8), it is easy to show that partial derivatives of g can be exactly computed at the true distribution \mathbb{P}_{BMM} through the following formula:

$$\nabla_{\mathbf{X}} g = \log \left(\frac{\mathbb{P}_{\text{BMM}}(\mathbf{X})}{\prod_{\ell=1}^d \left(\sum_{\forall \mathbf{X}' | X'_\ell = X_\ell} \mathbb{P}_{\text{BMM}}(\mathbf{X}') \right)} \right) - (d-1), \quad \mathbf{X} \in \{0,1\}^d.$$

Hence, the following upper-bound holds for the partial derivatives of g for all binary vectors \mathbf{X} for sufficiently large n :

$$\begin{aligned} \sup_{x \in \mathcal{N}(\mathbf{X})} |\nabla_{\mathbf{X}} g(x)| &\leq \left| \log \left(\frac{\mathbb{P}_{\text{BMM}}(\mathbf{X})}{\prod_{\ell=1}^d \mathbb{P}_{\text{BMM}}(\mathbf{X})} \right) \right| + (d-1) \\ &\leq d \left(\log \frac{1}{\mathbb{P}_{\text{BMM}}(\mathbf{X})} + 1 \right). \end{aligned}$$

So far, we have managed to upper-bound the estimation error in the current

lemma by the following summation:

$$\begin{aligned}
& \left| \mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \right| \\
& \leq d \sum_{\forall \mathbf{X}} \left(\log \frac{1}{\mathbb{P}_{\text{BMM}}(\mathbf{X})} + 1 \right) \left| \hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \mathbb{P}_{\text{BMM}}(\mathbf{X}) \right| \\
& = d \sum_{\forall \mathbf{X}} \left(\log \frac{1}{\mathbb{P}_{\text{BMM}}(\mathbf{X})} + 1 \right) \sigma(\mathbf{X}) \left| \frac{\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \mathbb{P}_{\text{BMM}}(\mathbf{X})}{\sigma(\mathbf{X})} \right| \\
& \leq d \left(\max_{\mathbf{X}} \left| \frac{\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \mathbb{P}_{\text{BMM}}(\mathbf{X})}{\sigma(\mathbf{X})} \right| \right) \sum_{\forall \mathbf{X}} \left(\log \frac{1}{\mathbb{P}_{\text{BMM}}(\mathbf{X})} + 1 \right) \sigma(\mathbf{X}),
\end{aligned}$$

with $\sigma(\mathbf{X}) \triangleq \sqrt{\mathbb{P}_{\text{BMM}}(\mathbf{X})(1 - \mathbb{P}_{\text{BMM}}(\mathbf{X}))}$.

An important issue that needs to be noted by a picky reader is that for those cases where $\mathbb{P}_{\text{BMM}}(\mathbf{X}) = 0$ or 1, the empirical probabilities always coincide with the true ones, hence corresponding error terms in the summation become exactly zero. As a result, such cases are implicitly omitted from all the summations.

It is easy to show that the summation over $\forall \mathbf{X}$ in the above inequality reaches its maximum at $\mathbb{P}_{\text{BMM}}(\mathbf{X}) = 2^{-d}$ for all \mathbf{X} , i.e.

$$\sum_{\forall \mathbf{X}} \left(\log \frac{1}{\mathbb{P}_{\text{BMM}}(\mathbf{X})} + 1 \right) \sigma(\mathbf{X}) \leq d 2^{d/2}.$$

What remains to do is to bound the deviations between true distribution \mathbb{P}_{BMM} and the empirical one $\hat{\mathbb{P}}_{\mathbf{Q}}$. Let us define the set of events $A_{\mathbf{X}}$, $\forall \mathbf{X} \in \{0, 1\}^d$ as

$$A_{\mathbf{X}} : \left| \frac{\hat{\mathbb{P}}_{\mathbf{Q}}(\mathbf{X}) - \mathbb{P}_{\text{BMM}}(\mathbf{X})}{\sigma(\mathbf{X})} \right| > \frac{\varepsilon}{d^2 2^{d/2}} \triangleq \delta.$$

By using the union and Chernoff bounds, one can have

$$\begin{aligned}
\mathbb{P} \left\{ \left| \mathcal{D}(\mathbf{Q}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Q}) \right| > \varepsilon \right\} & \leq \mathbb{P} \left\{ \bigcup_{\forall \mathbf{X}} A_{\mathbf{X}} \right\} \leq \sum_{\forall \mathbf{X}} \mathbb{P} \{A_{\mathbf{X}}\} \\
& \leq \sum_{\forall \mathbf{X}} e^{-n \mathcal{D}_{\text{KL}}(\mathbb{P}_{\text{BMM}}(\mathbf{X}) + \delta \sigma(\mathbf{X}) \parallel \mathbb{P}_{\text{BMM}}(\mathbf{X}))} \mathbb{I} \left(\delta \leq \frac{1 - \mathbb{P}_{\text{BMM}}(\mathbf{X})}{\sigma(\mathbf{X})} \right) \\
& + \sum_{\forall \mathbf{X}} e^{-n \mathcal{D}_{\text{KL}}(\mathbb{P}_{\text{BMM}}(\mathbf{X}) - \delta \sigma(\mathbf{X}) \parallel \mathbb{P}_{\text{BMM}}(\mathbf{X}))} \mathbb{I} \left(\delta \leq \frac{\mathbb{P}_{\text{BMM}}(\mathbf{X})}{\sigma(\mathbf{X})} \right),
\end{aligned}$$

where $\mathbb{I}(\cdot)$ is the indicator function. $\mathcal{D}_{\text{KL}}(\cdot \parallel \cdot)$ represents the Kullback-Leibler divergence, where by $\mathcal{D}_{\text{KL}}(x \parallel y)$ for $x, y \in [0, 1]$ we simply mean

$$x \log \frac{x}{y} + (1 - x) \log \frac{1 - x}{1 - y}.$$

KL divergence measures in the above upper-bounds can be lower-bounded according to Chernoff's theorem. Thus, the probability of observing a deviation greater than ε in estimating $\lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Y})$ can be upper-bounded as

$$\mathcal{P} \left\{ \left| \mathcal{D}(\mathbf{Y}) - \lim_{n \rightarrow \infty} \mathcal{D}(\mathbf{Y}) \right| > \varepsilon \right\} \leq 2^{d+1} \exp \left(\frac{-n\varepsilon^2}{d^4 2^{d+1}} \right),$$

which completes the proof. \square

PROOF OF LEMMA 4.2. Proof is highly similar to that of Lemma 4.1 and A.2. The main difference lies in the fact that when $K = 1$, i.e. a single Bernoulli model, one can easily verify

$$\begin{aligned} \nabla_{\mathbf{X}} g &= \log \left(\frac{\mathbb{P}_{\text{BMM}}(\mathbf{X})}{\prod_{\ell=1}^d \left(\sum_{\forall \mathbf{X}' | X'_\ell = X_\ell} \mathbb{P}_{\text{BMM}}(\mathbf{X}') \right)} \right) - (d - 1) = 1 - d, \\ &\quad \forall \mathbf{X} \in \{0, 1\}^d, \end{aligned}$$

and therefore $|\nabla_{\mathbf{X}} g| \leq d$.

Then, following the same steps as shown in proof of Lemma A.2 achieves the desired result and complete the proof. \square

PROOF OF THEOREM 4.3. The first inequality states that the probability of all $\binom{L}{d}$ sub-matrices of \mathbf{Y} with d columns having $\mathcal{D}(\mathbf{Q}) < \tau$ is strictly bounded from above. In order to show that, one should note the frequency matrix $\mathbf{P} \in [0, 1]^{K \times L}$ is assumed to be (\mathcal{L}, δ) -separable, which means for every pair of rows in \mathbf{P} there are at least $\mathcal{L} \leq L$ Bernoulli trials where their frequencies differ in a value greater than or equal to $\delta > 0$.

It can be easily seen that by examining all $\binom{L}{d}$ subsets of dimensions with size d , there are at least $h \triangleq \lfloor \mathcal{L}/d \rfloor$ non-overlapping sub-matrices of \mathbf{Y} , denoted by $\mathbf{Q}_1, \dots, \mathbf{Q}_h$, which are $(d/(K-1), \delta)$ -separable². Since $\mathbf{Q}_1, \dots, \mathbf{Q}_h$

²In practice, since *informative* dimensions are dispersed randomly among all Bernoulli trials, (d, δ) -separability holds almost surely for sufficiently large d . However, we have considered the improbable worst case where only $(d/(K-1), \delta)$ -separability is guaranteed.

do not overlap with each other, they are statistically independent which implies

$$\begin{aligned}\mathbb{P}\{\mathcal{D}_{\max}(\mathbf{Y}, d) \leq \tau\} &\leq \mathbb{P}\{\mathcal{D}(\mathbf{Q}_i) \leq \tau, \forall i\} \\ &= \prod_{i=1}^h \mathbb{P}\{\mathcal{D}(\mathbf{Q}_i) \leq \tau\}\end{aligned}$$

Using the upper-bound for $\mathbb{P}\{\mathcal{D}(\mathbf{Q}_i) \leq \tau\}$, $\forall i$ which is derived in Lemma 4.1 and approximating h with \mathcal{L}/d provide us with the claimed inequality.

For the second inequality, one can simply employ the union bound as

$$\begin{aligned}\mathbb{P}\{\mathcal{D}_{\max}(\mathbf{Y}; L_0) > \tau\} &\leq \mathbb{P}\left\{\max_{\mathbf{Q} \in \text{Col}(\mathbf{Y}; d)} \mathcal{D}(\mathbf{Q}) > \tau\right\} \\ (A.9) \quad &\leq \sum_{\mathbf{Q} \in \text{Col}(\mathbf{Y}; d)} \mathbb{P}\{\mathcal{D}(\mathbf{Q}) > \tau\},\end{aligned}$$

where $\text{Col}(\mathbf{Y}; d)$ represents the set of all $\binom{L}{d}$ sub-matrices of \mathbf{Y} with d columns. Again, substitution of $\mathbb{P}\{\mathcal{D}(\mathbf{Q}) > \tau\}$ with the upper-bound derived in Lemma A.2 gives us the claimed inequality and completes the proof. \square

LEMMA A.3. *Consider $\mathcal{M} = \mathcal{M}(K, \mathbf{w})$ to be a multinomial distribution with K mutually exclusive outcomes and corresponding probability vector $\mathbf{w} = (w_1, \dots, w_K)$. Assume there exists $\alpha > 0$ such that $\min_k w_k \geq \alpha$. Let $\mathbf{D} \triangleq \{X_1, \dots, X_n\}$ to be n i.i.d. samples drawn from \mathcal{M} . For $\zeta > 0$, assume*

$$n \geq \frac{2}{\alpha^2} \log \frac{K}{\zeta}.$$

Then, with probability at least $1 - \zeta$, the smallest cluster in \mathbf{D} has at least $\alpha n/2$ members.

PROOF. We denote the probability of the smallest cluster in \mathbf{D} having less than $\alpha n/2$ members by P_E . Let $\mathcal{A}_1, \dots, \mathcal{A}_K$ to denote the following events: for $k = 1, \dots, K$, \mathcal{A}_k represents the event that the k th cluster in \mathbf{D} (corresponding to probability component w_k) has less than $nw_k/2$ members. Then, the following holds according to union bound:

$$(A.10) \quad P_E \leq \sum_{k=1}^K \mathbb{P}\{\mathcal{A}_k\}.$$

Now, for each $k = 1, \dots, K$, consider the binomial random variable Y_k as follows:

$$(A.11) \quad \mathbb{P}(Y_k) \triangleq \begin{cases} w_k & Y_k = 1 \\ 1 - w_k & Y_k = 0 \\ 0 & \text{O.W.} \end{cases},$$

with $\mathbb{E}Y_k = w_k$. Let y_1, \dots, y_n to be n i.i.d. samples of Y_k . Define $S_k \triangleq y_1 + \dots + y_n$, while obviously we have $\mathbb{E}S_k = nw_k$. This way, one can easily verify the following chain of relations:

$$(A.12) \quad \begin{aligned} \mathbb{P}\{\mathcal{A}_k\} &= \mathbb{P}\{S_k < nw_k/2\} = \mathbb{P}\{S_k - \mathbb{E}S_k < -nw_k/2\} \\ &\leq \exp\left(\frac{-2n^2w_k^2}{4\sum_{i=1}^n (\max Y_k - \min Y_k)^2}\right) \\ &= \exp\left(\frac{-nw_k^2}{2}\right). \end{aligned}$$

Recall that we have $w_k \geq \alpha$ for all $k = 1, \dots, K$, thus one can write

$$(A.13) \quad P_E \leq \sum_{k=1}^K \exp\left(\frac{-nw_k^2}{2}\right) \leq K \exp\left(\frac{-n\alpha^2}{2}\right),$$

which given the condition on n in the lemma, results into $P_E \leq \zeta/3$ and completes the proof. \square

REFERENCES

- [1] M. Hollander, D. A. Wolfe, and E. Chicken, *Nonparametric statistical methods*. John Wiley & Sons, 2013.
- [2] C. Bouveyron and C. Brunet-Saumard, “Model-based clustering of high-dimensional data: A review,” *Computational Statistics & Data Analysis*, vol. 71, pp. 52–78, 2014.
- [3] P. D. McNicholas, “Model-based clustering,” *Journal of Classification*, vol. 33, no. 3, pp. 331–373, 2016.
- [4] P. Müller, F. A. Quintana, A. Jara, and T. Hanson, *Bayesian nonparametric data analysis*. Springer, 2015.
- [5] A. Juan and E. Vidal, “Bernoulli mixture models for binary images,” in *Pattern Recognition, 2004. ICPR 2004. Proceedings of the 17th International Conference on*, vol. 3. IEEE, 2004, pp. 367–370.
- [6] D. Tiedeman, “On the study of types,” in *Symposium on pattern analysis*, 1955, pp. 1–14.
- [7] J. H. Wolfe, “Pattern clustering by multivariate mixture analysis,” *Multivariate Behavioral Research*, vol. 5, no. 3, pp. 329–350, 1970.
- [8] L. D. Baker and A. K. McCallum, “Distributional clustering of words for text classification,” in *Proceedings of the 21st annual international ACM SIGIR conference on Research and development in information retrieval*. ACM, 1998, pp. 96–103.

- [9] C. Biernacki, G. Celeux, and G. Govaert, "An improvement of the nec criterion for assessing the number of clusters in a mixture model," *Pattern Recognition Letters*, vol. 20, no. 3, pp. 267–272, 1999.
- [10] M. Mohri, A. Rostamizadeh, and A. Talwalkar, *Foundations of machine learning*. MIT press, 2012.
- [11] P. F. Lazarsfeld, N. W. Henry, and T. W. Anderson, *Latent structure analysis*. Houghton Mifflin Boston, 1968, vol. 109.
- [12] C. M. Bishop, "Pattern recognition," *Machine Learning*, vol. 128, pp. 1–58, 2006.
- [13] G. McLachlan and D. Peel, *Finite mixture models*. John Wiley & Sons, 2004.
- [14] C. Li, B. Wang, V. Pavlu, and J. Aslam, "Conditional bernoulli mixtures for multi-label classification," in *Proceedings of The 33rd International Conference on Machine Learning*, 2016, pp. 2482–2491.
- [15] G. Celeux and G. Soromenho, "An entropy criterion for assessing the number of clusters in a mixture model," *Journal of classification*, vol. 13, no. 2, pp. 195–212, 1996.
- [16] J. A. Palmer, K. Kreutz-Delgado, and S. Makeig, "A generalized multivariate logistic model and em algorithm based on the normal variance mean mixture representation," in *Statistical Signal Processing Workshop (SSP), 2016 IEEE*. IEEE, 2016, pp. 1–5.
- [17] M. A. T. Figueiredo and A. K. Jain, "Unsupervised learning of finite mixture models," *IEEE Transactions on pattern analysis and machine intelligence*, vol. 24, no. 3, pp. 381–396, 2002.
- [18] C. Fraley and A. E. Raftery, "Model-based clustering, discriminant analysis, and density estimation," *Journal of the American statistical Association*, vol. 97, no. 458, pp. 611–631, 2002.
- [19] A. Juan, J. García-Hernández, and E. Vidal, "Em initialisation for bernoulli mixture learning," *Structural, Syntactic, and Statistical Pattern Recognition*, pp. 635–643, 2004.
- [20] S. Balakrishnan, M. J. Wainwright, B. Yu *et al.*, "Statistical guarantees for the em algorithm: From population to sample-based analysis," *The Annals of Statistics*, vol. 45, no. 1, pp. 77–120, 2017.
- [21] J. Rousseau, "On the frequentist properties of bayesian nonparametric methods," *Annual Review of Statistics and Its Application*, vol. 3, pp. 211–231, 2016.
- [22] P. Orbanz and Y. W. Teh, "Bayesian nonparametric models," in *Encyclopedia of Machine Learning*. Springer, 2011, pp. 81–89.
- [23] S. J. Gershman and D. M. Blei, "A tutorial on bayesian nonparametric models," *Journal of Mathematical Psychology*, vol. 56, no. 1, pp. 1–12, 2012.
- [24] Y. W. Teh, M. I. Jordan, M. J. Beal, and D. M. Blei, "Sharing clusters among related groups: Hierarchical dirichlet processes," in *Advances in neural information processing systems*, 2005, pp. 1385–1392.
- [25] J. K. Pritchard, M. Stephens, and P. Donnelly, "Inference of population structure using multilocus genotype data," *Genetics*, vol. 155, no. 2, pp. 945–959, 2000.
- [26] P. M. Visscher, M. A. Brown, M. I. McCarthy, and J. Yang, "Five years of gwas discovery," *The American Journal of Human Genetics*, vol. 90, no. 1, pp. 7–24, 2012.
- [27] J. Pella and M. Masuda, "The gibbs and split merge sampler for population mixture analysis from genetic data with incomplete baselines," *Canadian Journal of Fisheries and Aquatic Sciences*, vol. 63, no. 3, pp. 576–596, 2006.
- [28] N. M. Kopelman, J. Mayzel, M. Jakobsson, N. A. Rosenberg, and I. Mayrose, "Clumpak: a program for identifying clustering modes and packaging population structure inferences across k," *Molecular ecology resources*, vol. 15, no. 5, pp. 1179–1191, 2015.

- [29] G. Evanno, S. Regnaut, and J. Goudet, "Detecting the number of clusters of individuals using the software structure: a simulation study," *Molecular ecology*, vol. 14, no. 8, pp. 2611–2620, 2005.
- [30] J. Catchen, P. A. Hohenlohe, S. Bassham, A. Amores, and W. A. Cresko, "Stacks: an analysis tool set for population genomics," *Molecular ecology*, vol. 22, no. 11, pp. 3124–3140, 2013.
- [31] R. Peakall and P. E. Smouse, "Genalex 6: genetic analysis in excel. population genetic software for teaching and research," *Molecular ecology notes*, vol. 6, no. 1, pp. 288–295, 2006.
- [32] S. Purcell, B. Neale, K. Todd-Brown, L. Thomas, M. A. Ferreira, D. Bender, J. Maller, P. Sklar, P. I. De Bakker, M. J. Daly *et al.*, "Plink: a tool set for whole-genome association and population-based linkage analyses," *The American Journal of Human Genetics*, vol. 81, no. 3, pp. 559–575, 2007.
- [33] D. Falush, M. Stephens, and J. K. Pritchard, "Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies," *Genetics*, vol. 164, no. 4, pp. 1567–1587, 2003.
- [34] J. K. Pritchard, M. Stephens, N. A. Rosenberg, and P. Donnelly, "Association mapping in structured populations," *The American Journal of Human Genetics*, vol. 67, no. 1, pp. 170–181, 2000.
- [35] J. Yu, G. Pressoir, W. H. Briggs, I. V. Bi, M. Yamasaki, J. F. Doebley, M. D. McMullen, B. S. Gaut, D. M. Nielsen, J. B. Holland *et al.*, "A unified mixed-model method for association mapping that accounts for multiple levels of relatedness," *Nature genetics*, vol. 38, no. 2, pp. 203–208, 2006.
- [36] H. Zhou, J. Blangero, T. D. Dyer, K.-h. K. Chan, K. Lange, and E. M. Sobel, "Fast genome-wide qtl association mapping on pedigree and population data," *Genetic epidemiology*, vol. 41, no. 3, pp. 174–186, 2017.
- [37] A. Najafi, S. Janghorbani, S. A. Motahari, and E. Fatemizadeh, "Statistical association mapping of population-structured genetic data," *IEEE/ACM Transactions on Computational Biology and Bioinformatics*, 2017.
- [38] A. L. Price, N. J. Patterson, R. M. Plenge, M. E. Weinblatt, N. A. Shadick, and D. Reich, "Principal components analysis corrects for stratification in genome-wide association studies," *Nature genetics*, vol. 38, no. 8, pp. 904–909, 2006.
- [39] M. Gyllenberg, T. Koski, E. Reilink, and M. Verlaan, "Non-uniqueness in probabilistic numerical identification of bacteria," *Journal of Applied Probability*, vol. 31, no. 2, pp. 542–548, 1994.
- [40] M. A. Carreira-Perpinán and S. Renals, "Practical identifiability of finite mixtures of multivariate bernoulli distributions," *Neural Computation*, vol. 12, no. 1, pp. 141–152, 2000.
- [41] E. S. Allman, C. Matias, and J. A. Rhodes, "Identifiability of parameters in latent structure models with many observed variables," *The Annals of Statistics*, pp. 3099–3132, 2009.
- [42] R. Courant, *Differential and integral calculus*. John Wiley & Sons, 2011, vol. 2.

ROOM 401, COMPUTER ENGINEERING DEPT.,
 SHARIF UNIVERSITY OF TECHNOLOGY,
 AZADI AVE., TEHRAN, IRAN
 E-MAIL: najafy@ce.sharif.edu
motahari@sharif.edu

ROOM 803, COMPUTER ENGINEERING DEPT.,
 SHARIF UNIVERSITY OF TECHNOLOGY,
 AZADI AVE., TEHRAN, IRAN
 E-MAIL: rabiee@sharif.edu
 URL: