

# SCHRAMM–LOEWNER-EVOLUTION-TYPE GROWTH PROCESSES CORRESPONDING TO WESS–ZUMINO–WITTEN THEORIES

SHINJI KOSHIDA

ABSTRACT. A group theoretical formulation of Schramm–Loewner-evolution-type growth processes corresponding to Wess–Zumino–Witten theories is developed that makes it possible to construct stochastic differential equations associated with more general null vectors than the ones considered in the most fundamental example in [Alekseev *et al.*, *Lett. Math. Phys.* **97**, 243-261 (2011)]. Also given are examples of Schramm–Loewner-evolution-type growth processes associated with null vectors of conformal weight 4 in the basic representations of  $\widehat{\mathfrak{sl}}_2$  and  $\widehat{\mathfrak{sl}}_3$ .

## 1. INTRODUCTION

Two-dimensional conformal field theories (CFTs) [BPZ84] have been proven to be powerful tools for analyzing a vast variety of models in statistical physics at criticality [DFMS97]. CFTs are distinguished from other quantum field theories by the property that several correlation functions in a CFT can be computed exactly owing to its infinite-dimensional symmetry. Not only can a bulk system without boundaries be probed but so can one under certain boundary conditions by means of the framework of boundary conformal field theories (BCFTs) [Car86, Car89]. In a BCFT, a partition function under given boundary conditions is defined by a correlation function of boundary operators, each of which changes boundary conditions. In this picture, a partition function of a BCFT gives a prediction for a cluster interface evolving from one boundary point to another. One of the milestones is Cardy’s formula [Car92], later proved by Smirnov [Smi01], which gives the crossing probability in two-dimensional critical percolation computed as partition functions of a BCFT.

Another framework describing a cluster interface is the notion of Schramm–Loewner evolution (SLE), which was introduced in [Sch00] as a subsequential scaling limit of loop-erased random walks and uniform spanning trees. Schramm defined in [Sch00] two types of SLEs, radial and chordal, but in this paper, we only consider chordal SLE and simply call it SLE. SLE labeled by a positive real number  $\kappa > 0$  is a collection of properly normalized conformal maps  $\{g_t(z)\}_{t \geq 0}$  satisfying the following ordinary differential equation:

$$(1.1) \quad \partial_t g_t(z) = \frac{2}{g_t(z) - B_t}, \quad g_0(z) = z.$$

Here  $B_t$  is Brownian motion of variance  $\kappa$  on the real axis. Following its introduction, many properties of SLE have been unveiled mainly in a probability theoretical manner [RS05, Law04]. In particular, it has been found that SLE emerges in scaling limits of

---

DEPARTMENT OF BASIC SCIENCE, THE UNIVERSITY OF TOKYO  
E-mail address: koshida@vortex.c.u-tokyo.ac.jp.

several two-dimensional lattice models including the spin Ising model [CDCH<sup>+</sup>14, Law05, Smi06].

Because CFT and SLE are two different frameworks used to treat the same critical systems, they should be connected in some sense. There are several approaches to relate these two notions under the name of SLE/CFT correspondence. In works by Friedrich, Kalkkinen, and Kontsevich [FK04, Fri04, Kon03], the SLE-type probability measure was constructed as a section of the determinant bundle over the moduli space of Riemann surfaces. Recently, Dubédat [Dub15b, Dub15a] also constructed SLE-type probability measures by means of the localization technique and identified its partition function with a highest weight vector in a highest weight representation of the Virasoro algebra.

In this paper, we follow the *group theoretical* formulation of SLE/CFT correspondence originated by Bauer and Bernard [BB02, BB03a, BB03b]. They established the relation between SLE and CFT via a random process on an infinite-dimensional Lie group. Their strategy was to consider a random process  $\mathcal{G}_t$  on the *lower Borel subgroup* of the Virasoro group, of which the Lie algebra is the Virasoro algebra, under the initial condition that  $\mathcal{G}_0$  is the unit element. Through transformation of a primary field, the random process  $\mathcal{G}_t$  gives a one-parameter family of germs  $\{g_t(z)\}_{t \geq 0}$  of holomorphic functions at infinity. At the same time, each  $\mathcal{G}_t$  acts on a highest weight representation of the Virasoro algebra. In particular,  $\mathcal{G}_t|h, c\rangle$  for a certain highest weight vector  $|h, c\rangle$  of conformal weight  $h$  and central charge  $c$  is a random process on the corresponding highest weight Verma module. The SLE/CFT correspondence in the sense of Bauer and Bernard states that, if the family of germs  $\{g_t(z)\}_{t \geq 0}$  satisfies Eq. (1.1), then the random process  $\mathcal{G}_t|h, c\rangle$  is a martingale for a certain choice of the highest weight because of the existence of a null vector of conformal weight  $h + 2$  in the irreducible representation  $L(h, c)$ . The group theoretical formulation of SLE/CFT correspondence has been convenient for generalizing SLE by constructing stochastic differential equations associated with more general null vectors in highest weight representations of the Virasoro algebra [LR04], ones for the  $\mathcal{N} = 1$  superconformal algebra [Ras04b, NR05], and ones in logarithmic representations of the Virasoro algebra [Ras04a, MARR04].

Aside from the group theoretical formulation, there is a *correlation function* formulation, in which one directly investigates martingale conditions on correlation functions to discover the relation between SLE and CFT. By the correlation function formulation, SLE/CFT correspondence for multiple SLEs has also been established in [BBK05] and for further variants of SLE in [Kyt07].

CFTs to be considered next to the Virasoro algebra include Wess–Zumino–Witten (WZW) theories associated with the representation theory of affine Lie algebras. SLEs corresponding to WZW theories have been considered group theoretically in [Ras07] for the  $\mathfrak{sl}_2$  case and in the correlation function formulation in [BGLW05, ABI11], which is extended to multiple SLEs in [Sak13] and to coset WZW theories in [Naz12, Fuk17]. The purpose of this paper is to better understand the construction of SLE-type processes in [Ras07] to extend it to general WZW theories and to generalize SLE-type growth processes corresponding to WZW theories. To this purpose, we imitate the construction in [BB03a] by considering a random process on the negative part of the affine Virasoro group, instead of the Virasoro group in [BB03a]. This enables us to obtain an SLE-type growth process associated with a null vector of a certain type in the state space of a WZW theory, which reduces to the stochastic process constructed in [ABI11] in a particular case. We also propose new SLE-type growth processes associated with

null vectors in the basic representations of  $\widehat{\mathfrak{sl}}_2$  and  $\widehat{\mathfrak{sl}}_3$  and conjecture that a similar construction is possible for all  $\widehat{\mathfrak{sl}}_n$ .

Besides the construction of new stochastic processes, a possible application of our formulation is to establish the module structure of the space of SLE martingales over an affine Lie algebra. Indeed, consideration on a random process on the Virasoro group was essential to define the Virasoro module structure on the space of SLE martingales in [BB04, Kyt07]. Similarly, a random process on an affine Virasoro group helps in finding the affine Lie algebra module structure on the space of SLE martingales, which we believe is one of completed formulations of SLE/CFT correspondence. Studies in this direction will be reported elsewhere [Kos18].

This paper is organized as follows. In Sect. 2, we give an introduction to the representation theory of affine Lie algebras in a somewhat vertex operator algebraic way. In Sect. 3, we formally introduce a random process on a Lie group, which is a fundamental object in the group theoretical formulation of SLE/CFT correspondence, and explain our strategy for constructing SLE-type growth processes. In Sect. 4, we derive stochastic differential equations from a given random process on an infinite-dimensional Lie group, and in Sect. 5, we give examples of null vectors, which lead to SLE-type growth processes.

## 2. AFFINE LIE ALGEBRAS AND WZW THEORIES

Let  $\mathfrak{g}$  be a finite-dimensional simple Lie algebra. We denote a symmetric invariant bilinear form on  $\mathfrak{g}$  by  $(\cdot|\cdot) : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathbb{C}$ . The affine Lie algebra of  $\mathfrak{g}$  is a Lie algebra

$$(2.1) \quad \widehat{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}] \oplus \mathbb{C}K$$

with Lie brackets being defined by

$$(2.2) \quad [X \otimes t^m, Y \otimes t^n] = [X, Y] \otimes t^{m+n} + m(X|Y)\delta_{m+n,0}K, \quad X, Y \in \mathfrak{g},$$

$$(2.3) \quad [K, \widehat{\mathfrak{g}}] = \{0\}.$$

As usual, we write  $X_m$  for  $X \otimes t^m$ .

The state space of the WZW theory associated with  $\mathfrak{g}$  is a representation of the affine Lie algebra  $\widehat{\mathfrak{g}}$ . Let us fix a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$ , and, correspondingly, let  $\Delta$  and  $\Pi^\vee = \{\alpha_1^\vee, \dots, \alpha_\ell^\vee\}$  be the set of roots and the set of simple coroots, respectively. The weight lattice is denoted by  $P = \bigoplus_{i=1}^\ell \mathbb{Z}\Lambda_i$ , where  $\Lambda_i$  are fundamental weights defined by  $\langle \Lambda_i, \alpha_j^\vee \rangle = \delta_{ij}$ . The set of dominant weights is  $P_+ = \{\Lambda \in P \mid \langle \Lambda, \alpha_i^\vee \rangle \in \mathbb{Z}_{\geq 0}\}$ . We denote by  $L(\Lambda)$  the finite-dimensional irreducible representation of  $\mathfrak{g}$  of highest weight  $\Lambda \in P_+$ . Given a finite-dimensional representation of  $\mathfrak{g}$ , we can construct a representation of the affine Lie algebra  $\widehat{\mathfrak{g}}$  by means of parabolic induction. Let  $M$  be a representation of  $\mathfrak{g}$ . Then the representation  $M_k$  of a parabolic subalgebra  $\widehat{\mathfrak{p}} := \mathfrak{g} \otimes \mathbb{C}[t] \oplus \mathbb{C}K$  of the affine Lie algebra is  $M$  as a vector space, on which  $\mathfrak{g} \otimes t^0$  acts naturally,  $\mathfrak{g} \otimes t\mathbb{C}[t]$  acts trivially, and  $K$  acts as the multiplication by  $k \in \mathbb{C}$ . Here the scalar  $k$  is called the level. Then a representation  $\widehat{M}_k$  of the affine Lie algebra is obtained by induction:

$$(2.4) \quad \widehat{M}_k := \text{Ind}_{\widehat{\mathfrak{p}}}^{\widehat{\mathfrak{g}}} M_k = U(\widehat{\mathfrak{g}}) \otimes_{U(\widehat{\mathfrak{p}})} M_k.$$

By the Poincaré–Birkhoff–Witt theorem, it is isomorphic to  $U(\mathfrak{g} \otimes t^{-1}\mathbb{C}[t^{-1}]) \otimes_{\mathbb{C}} M_k$  as a vector space. For a finite-dimensional irreducible representation  $L(\Lambda)$  of  $\mathfrak{g}$ ,  $\widehat{L(\Lambda)}_k$ ,

called a Weyl module, may be reducible for specific levels. We denote by  $L_{\mathfrak{g},k}(\Lambda)$  its irreducible quotient.

A WZW theory is specified by a pair  $(\mathfrak{g}, k)$  of a finite-dimensional simple Lie algebra  $\mathfrak{g}$  and a positive integer level  $k \in \mathbb{Z}_{>0}$ . The state space of the WZW theory of  $(\mathfrak{g}, k)$  is identified with

$$(2.5) \quad V_{\mathfrak{g},k} = \bigoplus_{\Lambda \in P_+^k} L_{\mathfrak{g},k}(\Lambda),$$

where  $P_+^k = \{\Lambda \in P_+ \mid \langle \Lambda, \theta \rangle \leq k\}$  is the set of dominant weights of level  $k$ , where  $\theta$  is the highest root of  $\mathfrak{g}$  [FZ92].

To explain the Sugawara construction, we introduce the notion of field operators. For  $X \in \mathfrak{g}$ , the corresponding field  $X(z)$  is an  $\text{End}(V_{\mathfrak{g},k})$ -valued formal power series in a formal variable  $z$ :

$$(2.6) \quad X(z) = \sum_{n \in \mathbb{Z}} X_n z^{-n-1},$$

where the coefficients  $X_n \in \text{End}(V_{\mathfrak{g},k})$  are thought of as the action of  $X_n \in \widehat{\mathfrak{g}}$  on  $V_{\mathfrak{g},k}$ . From the definition of the affine Lie algebra, Eq. (2.2), two fields satisfy the commutation relation

$$(2.7) \quad [X(z), Y(w)] = [X, Y](w)\delta(z-w) + k(X|Y)\partial_w\delta(z-w),$$

where  $\delta(z-w) = \sum_{n \in \mathbb{Z}} z^{-n-1}w^n$  is the formal delta distribution. This commutation relation is often expressed equivalently in the form of operator product expansion (OPE):

$$(2.8) \quad X(z)Y(w) \sim \frac{[X, Y](w)}{z-w} + \frac{k(X|Y)}{(z-w)^2}.$$

For two fields  $X(z)$  and  $Y(z)$ , their naive product  $X(z)Y(z)$  is not well-defined because of the singularity at  $z=w$  in the above OPE. Instead, we define the normal ordered product of two fields. For a field  $X(z)$ , its positive and negative power parts  $X(z)_\pm$  are defined by

$$(2.9) \quad X(z)_+ = \sum_{n=-\infty}^{-1} X_n z^{-n-1}, \quad X(z)_- = \sum_{n=0}^{\infty} X_n z^{-n-1}.$$

Then the normal ordered product of  $X(z)$  and  $Y(z)$  is defined by

$$(2.10) \quad :X(z)Y(z): = X(z)_+Y(z) + Y(z)X(z)_-.$$

Let  $\{X_i\}_{i=1}^{\dim \mathfrak{g}}$  be an orthonormal basis of  $\mathfrak{g}$  with respect to the symmetric invariant bilinear form  $(\cdot|\cdot)$ . We also assume that the bilinear form is normalized so that  $(\theta|\theta) = 2$ . We consider the field

$$(2.11) \quad L(z) = \frac{1}{2(k+h^\vee)} \sum_{i=1}^{\dim \mathfrak{g}} :X_i(z)^2:,$$

where  $h^\vee$  is the dual Coxeter number of  $\mathfrak{g}$ . Then it satisfies

$$(2.12) \quad [L(z), L(w)] = \frac{c_k}{2} \partial_w^{(3)} \delta(z-w) + 2L(w) \partial_w \delta(z-w) + \partial L(w) \delta(z-w)$$

with  $c_k = k \dim \mathfrak{g} / (k+h^\vee)$ , which implies that the coefficients of the expansion  $L(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2}$  give a representation of the Virasoro algebra of the central charge  $c_k$  on

$V_{\mathfrak{g},k}$ . Moreover, each direct summand  $L_{\mathfrak{g},k}(\Lambda)$  of  $V_{\mathfrak{g},k}$ , and thus  $V_{\mathfrak{g},k}$  itself, decomposes into a direct sum of eigenspaces of  $L_0$  so that  $L_{\mathfrak{g},k}(\Lambda) = \bigoplus_{n \in \mathbb{Z}_{\geq 0}} L_{\mathfrak{g},k}(\Lambda)_{h_\Lambda + n}$ , where

$$(2.13) \quad L_{\mathfrak{g},k}(\Lambda)_h := \{v \in L_{\mathfrak{g},k}(\Lambda) \mid L_0 v = h v\}$$

is a finite-dimensional eigenspace of  $L_0$  and the constant  $h_\Lambda$  is given by

$$(2.14) \quad h_\Lambda = \frac{(\Lambda \mid \Lambda + 2\rho)}{2(k + h^\vee)}$$

with  $\rho = \sum_{i=1}^{\ell} \Lambda_i \in \mathfrak{h}^*$  being the Weyl vector. From direct computation, we find that each  $X(z)$  for  $X \in \mathfrak{g}$  is a primary field with respect to  $L(z)$ :

$$(2.15) \quad [L(z), X(w)] = X(w) \partial_w \delta(z-w) + \partial X(w) \delta(z-w),$$

which, in particular, implies that the mode  $X_{-n}$  raises the eigenvalue of  $L_0$  by  $n$ . We also remark that the *top space*  $L_{\mathfrak{g},k}(\Lambda)_{h_\Lambda}$  is isomorphic to the irreducible representation  $L(\Lambda)$  of  $\mathfrak{g}$ .

In the rest of this section, we give a vertex operator algebra (VOA) structure on  $V_{\mathfrak{g},k}$ , which is convenient in the following computation. To be precise, only the space  $L_{\mathfrak{g},k} := L_{\mathfrak{g},k}(0) \subset V_{\mathfrak{g},k}$  carries a VOA structure, and the other direct summands are regarded as modules over this VOA. In this paper, however, we sacrifice the preciseness and introduce the standard computational tools in the theory of VOAs, without distinguishing *intertwining operators* from the state-field correspondence map. For precise formulation, see [Kac98, FBZ04, FZ92]. The fundamental object in the theory of VOAs is the state-field correspondence map

$$(2.16) \quad \mathcal{Y}(-, z) : V_{\mathfrak{g},k} \rightarrow \text{End}(V_{\mathfrak{g},k})\{z\} = \left\{ \sum_{a \in \mathbb{C}} T_a z^a \mid T_a \in \text{End}(V_{\mathfrak{g},k}) \right\},$$

which satisfies

$$(2.17) \quad \mathcal{Y}(v, z) |0\rangle \Big|_{z=0} = v$$

for an arbitrary  $v \in V_{\mathfrak{g},k}$ . Here  $|0\rangle \in (L_{\mathfrak{g},k})_0 = \mathbb{C}|0\rangle$  is a fixed vector called the vacuum. Another significant property of  $\mathcal{Y}(-, z)$  is that its value is a field; *i.e.*, for arbitrary  $v, w \in V_{\mathfrak{g},k}$ , there are finitely many weights  $h_i \in \mathbb{R}$  ( $i = 1, \dots, n$ ) such that

$$(2.18) \quad \mathcal{Y}(v, z)w \in \bigoplus_{i=1}^n V_{\mathfrak{g},k}((z))z^{h_i}.$$

In this notation, the fields  $X(z)$  for  $X \in \mathfrak{g}$  and  $L(z)$  are described as

$$(2.19) \quad X(z) = \mathcal{Y}(X_{-1}|0\rangle, z),$$

$$(2.20) \quad L(z) = \mathcal{Y}(L_{-2}|0\rangle, z).$$

Other important fields include one corresponding to  $v \in L_{\mathfrak{g},k}(\Lambda)_{h_\Lambda}$  in the top space of a direct summand of  $V_{\mathfrak{g},k}$ . This kind of field is often called a *primary* field and is characterized by the following properties:

$$(2.21) \quad [L(z), \mathcal{Y}(v, w)] = h_\Lambda \mathcal{Y}(v, w) \partial_w \delta(z-w) + \partial \mathcal{Y}(v, w) \delta(z-w),$$

$$(2.22) \quad [X(z), \mathcal{Y}(v, w)] = \mathcal{Y}(Xv, w) \delta(z-w).$$

Notice that, for each  $\Lambda \in P_+^k$ , there are  $\dim L(\Lambda)$  linearly independent primary fields. The multiplet of them is regarded as an  $L(\Lambda)^* \otimes \text{End}(V_{\mathfrak{g},k})$ -valued field.

## 3. ITO PROCESS ON A LIE GROUP

In this section, we formally introduce an Ito process on a Lie group, which is needed in the following. For a precise definition, see [App14]. Let  $\mathfrak{g}$  be a finite-dimensional complex Lie algebra that is faithfully represented on a finite-dimensional vector space  $V$ . There is a Lie group  $G$  realized as a Lie subgroup of  $\mathrm{GL}(V)$  such that the Lie algebra of  $G$  is  $\mathfrak{g}$ . We take independent Brownian motions  $B_t^{(i)}$ ,  $i = 1, \dots, \dim \mathfrak{g}$ , and let  $\kappa_i > 0$  be the variances of  $B_t^{(i)}$ ; *i.e.*, we have

$$(3.1) \quad dB_t^{(i)} \cdot dB_t^{(j)} = \delta_{i,j} \kappa_i dt.$$

An Ito process  $\mathcal{G}_t$  on the Lie group  $G$  is described by a stochastic differential equation

$$(3.2) \quad \mathcal{G}_t^{-1} d\mathcal{G}_t = \left( \sum_i \mu_i X_i + \frac{1}{2} \sum_i \kappa_i \sigma_i^2 X_i^2 \right) dt + \sum_i \sigma_i X_i dB_t^{(i)},$$

where  $\{X_i\}_{i=1}^{\dim \mathfrak{g}}$  is a certain basis of  $\mathfrak{g}$ , and  $\mu_i$  and  $\sigma_i$  are random processes satisfying some finiteness conditions. Note that, in this description, terms  $X_i^2$  do not lie in the Lie algebra  $\mathfrak{g}$  but are understood as elements in  $\mathrm{End}(V)$ .

Although we assumed that the Lie group  $G$  is finite dimensional above, an Ito process on an infinite-dimensional Lie group is significant in the context of SLE/CFT correspondence. We assume that the expression in Eq. (3.2) makes sense on some representations even if the Lie group is infinite dimensional in the following discussion.

To construct an SLE-type growth process corresponding to a WZW theory, we consider an Ito process on the subgroup  $\widehat{G}_{\mathrm{Vir}}^{<0}$  of the affine Virasoro group. It is formally generated by exponentials of the negative part  $\widehat{\mathfrak{g}}_{\mathrm{Vir}}^{<0}$  of the affine Virasoro algebra, which is defined by  $\widehat{\mathfrak{g}}_{\mathrm{Vir}}^{<0} = \mathfrak{g} \otimes \mathbb{C}[t^{-1}]t^{-1} \oplus \bigoplus_{n \in \mathbb{Z}_{<0}} \mathbb{C}L_n$  with Lie brackets

$$(3.3) \quad [X \otimes t^m, Y \otimes t^n] = [X, Y] \otimes t^{m+n}, \quad X, Y \in \mathfrak{g},$$

$$(3.4) \quad [L_m, L_n] = (m - n)L_{m+n},$$

$$(3.5) \quad [L_m, X_n] = -nX_{m+n},$$

for  $m, n \in \mathbb{Z}_{<0}$ . The commutation relation in Eq. (2.15) suggests that the state space  $V_{\mathfrak{g},k}$  is indeed a representation of this Lie algebra.

We assume that an Ito process  $\mathcal{G}_t$  on  $\widehat{G}_{\mathrm{Vir}}^{<0}$  depends on finitely many mutually independent Brownian motions  $\{B_t^{(i)}\}_{i \in I_B}$  labeled by a certain finite set  $I_B$  and that the variance of each  $B_t^{(i)}$  is  $\kappa_i > 0$ . Then, analogously to the expression in Eq. (3.2), the increment of  $\mathcal{G}_t$  can be written in the form

$$(3.6) \quad \mathcal{G}_t^{-1} d\mathcal{G}_t = \left( \mu + \frac{1}{2} \sum_{i \in I_B} \kappa_i \sigma_i^2 \right) dt + \sum_{i \in I_B} \sigma_i dB_t^{(i)},$$

where  $\mu$  and  $\sigma_i$  for  $i \in I_B$  are  $\widehat{\mathfrak{g}}_{\mathrm{Vir}}^{<0}$ -valued random processes. We also impose the initial condition so that  $\mathcal{G}_0$  is the unit element in  $\widehat{G}_{\mathrm{Vir}}^{<0}$ .

As is explained in the previous section, for a vector  $v \in (L_{\mathfrak{g},k}(\Lambda))_{h_\Lambda}$  in the top space of  $L_{\mathfrak{g},k}(\Lambda)$ , the corresponding field  $\mathcal{Y}(v, z)$  satisfies the primary field conditions given by Eqs. (2.21) and (2.22). This implies that the adjoint action of  $\mathcal{G} \in \widehat{G}_{\mathrm{Vir}}^{<0}$  transforms the primary field  $\mathcal{Y}(v, z)$  so that

$$(3.7) \quad \mathcal{G}^{-1} \mathcal{Y}(v, z) \mathcal{G} = (f'(z))^{h_\Lambda} \mathcal{Y}(\Theta(z)v, f(z)).$$

Here  $f(z) \in \mathbb{C}[[z^{-1}]]z$  is a formal Laurent series in  $z^{-1}$  with a nonzero coefficient of  $z$ , and  $\Theta(z) \in G[[z^{-1}]]$  is a  $\mathbb{C}[[z^{-1}]]$ -valued point in the finite-dimensional Lie group  $G$  such that  $\text{Lie}(G) = \mathfrak{g}$ , which naturally defines a homomorphism  $L(\Lambda) \rightarrow L(\Lambda)[[z^{-1}]]$ . We have to comment that  $\mathcal{Y}(\Theta(z)v, f(z))$  is not a field, thus we make  $\mathcal{Y}(\Theta(z)v, f(z))$  meaningful in the following way. Let  $V_{\mathfrak{g},k} = \bigoplus_{h \in \mathcal{P}} (V_{\mathfrak{g},k})_h$  be the eigenspace decomposition with respect to  $L_0$ , where  $\mathcal{P} = \bigcup_{\Lambda \in P_+^k} (\mathbb{Z}_{\geq 0} + h_\Lambda)$  is the set of eigenvalues. Then the restricted dual space of  $V_{\mathfrak{g},k}$  is defined by  $V_{\mathfrak{g},k}^\dagger = \bigoplus_{h \in \mathcal{P}} (V_{\mathfrak{g},k})_h^*$ . Although  $\mathcal{Y}(\Theta(z)v, f(z))$  is not a field, for a vector  $u^* \in V_{\mathfrak{g},k}^\dagger$ , there are finitely many weights  $h_i \in \mathbb{R}$  ( $i = 1, \dots, n$ ), such that  $\langle u^* | \mathcal{Y}(\Theta(z)v, f(z)) \rangle \in V_{\mathfrak{g},k}^\dagger((z^{-1}))z^{h_i}$ . We understand  $\mathcal{Y}(\Theta(z)v, f(z))$  as such an object in the following.

Replacing  $\mathcal{G}$  by the Ito process  $\mathcal{G}_t$  on  $\widehat{G}_{\text{Vir}}^{<0}$ , we obtain

$$(3.8) \quad \mathcal{G}_t^{-1} \mathcal{Y}(v, z) \mathcal{G}_t = (f'_t(z))^{h_\Lambda} \mathcal{Y}(\Theta_t(z)v, f_t(z)),$$

where  $f_t(z)$  is a  $\mathbb{C}[[z^{-1}]]z$ -valued Ito process and  $\Theta_t(z)$  is a  $G[[z^{-1}]]$ -valued one. Namely, an Ito process  $\mathcal{G}_t$  on  $\widehat{G}_{\text{Vir}}^{<0}$  induces a pair of, in general, interacting random processes  $(f_t(z), \Theta_t(z))$  on  $\mathbb{C}[[z^{-1}]]z$  and  $G[[z^{-1}]]$ . We suppose that the increment of  $f_t(z)$  is written as

$$(3.9) \quad df_t(z) = \bar{f}_t(z) dt + \sum_{i \in I_B} f_t^i(z) dB_t^{(i)}$$

with  $\bar{f}_t(z)$  and  $f_t^i(z)$  being some  $\mathbb{C}[[z^{-1}]]z$ -valued random processes. To describe the increment of  $\Theta_t(z)$ , we take a basis  $\{X_r\}_{r \in I_{\mathfrak{g}}}$  of  $\mathfrak{g}$  with the index set  $I_{\mathfrak{g}}$ . Then we can express the increment of  $\Theta_t(z)$  as

$$(3.10) \quad d\Theta_t(z) \Theta_t(z)^{-1} = \sum_{r \in I_{\mathfrak{g}}} d\theta_{r,t}(z) X_r + \frac{1}{2} \sum_{r,s \in I_{\mathfrak{g}}} d\theta_{r,t}(z) d\theta_{s,t}(z) X_r X_s.$$

We also suppose that the increment of each  $\theta_{r,t}(z)$  is written as

$$(3.11) \quad d\theta_{r,t}(z) = \bar{\theta}_{r,t}(z) dt + \sum_{i \in I_B} \theta_{r,t}^i(z) dB_t^{(i)},$$

with  $\bar{\theta}_{r,t}(z)$  and  $\theta_{r,t}^i(z)$  being  $\mathbb{C}[[z^{-1}]]z$ -valued random processes.

Our strategy is to compare the increments of both sides of Eq. (3.8) and to establish stochastic differential equations on  $f_t(z)$  and  $\theta_{r,t}(z)$  for a given Ito process  $\mathcal{G}_t$  on  $\widehat{G}_{\text{Vir}}^{<0}$ .

Notice that, from  $d(\mathcal{G}_t^{-1} \mathcal{G}_t) = 0$ , we have

$$(3.12) \quad (d\mathcal{G}_t^{-1}) \mathcal{G}_t = \left( -\mu + \frac{1}{2} \sum_{i \in I_B} \kappa_i \sigma_i^2 \right) dt - \sum_{i \in I_B} \sigma_i dB_t^{(i)}.$$

Using this formula, we find that the increment of the left-hand side of Eq. (3.8) yields

$$(3.13) \quad \begin{aligned} & d(\mathcal{G}_t^{-1} \mathcal{Y}(v, z) \mathcal{G}_t) \\ &= (f'_t(z))^{h_\Lambda} \left( -[\mu, \mathcal{Y}(\Theta_t(z)v, f_t(z))] + \frac{1}{2} \sum_{i \in I_B} \kappa_i [\sigma_i, [\sigma_i, \mathcal{Y}(\Theta_t(z)v, f_t(z))]] \right) dt \\ & \quad - (f'_t(z))^{h_\Lambda} \sum_{i \in I_B} [\sigma_i, \mathcal{Y}(\Theta_t(z)v, f_t(z))] dB_t^{(i)}. \end{aligned}$$

The increment of the right-hand side of Eq. (3.8) is computed by standard Ito calculus as

$$\begin{aligned}
(3.14) \quad & d\left((f'_t(z))^{h_\Lambda} \mathcal{Y}(\Theta_t(z)v, f(z))\right) \\
&= (f'_t(z))^{h_\Lambda} \left\{ \left( h_\Lambda \frac{\bar{f}'_t(z)}{f'_t(z)} + \frac{1}{2} h_\Lambda (h_\Lambda - 1) \sum_{i \in I_B} \kappa_i \left( \frac{f''_t(z)}{f'_t(z)} \right)^2 \right. \right. \\
&\quad \left. \left. + \bar{f}_t(z) \frac{\partial}{\partial f_t(z)} + h_\Lambda \sum_{i \in I_B} \kappa_i \frac{f''_t(z)}{f'_t(z)} f_t^i(z) \frac{\partial}{\partial f_t(z)} \right. \right. \\
&\quad \left. \left. + \frac{1}{2} \sum_{i \in I_B} \kappa_i f_t^i(z)^2 \frac{\partial^2}{\partial f_t(z)^2} \right) \mathcal{Y}(\Theta_t(z)v, f_t(z)) \right. \\
&\quad \left. + \sum_{r \in I_{\mathfrak{g}}} \left( \bar{\theta}_{r,t}(z) + h_\Lambda \sum_{i \in I_B} \kappa_i \frac{f''_t(z)}{f'_t(z)} \theta_{r,t}^i(z) \right. \right. \\
&\quad \left. \left. + \sum_{i \in I_B} \kappa_i \theta_{r,t}^i(z) f_t^i(z) \frac{\partial}{\partial f_t(z)} \right) \mathcal{Y}(X_r \Theta_t(z)v, f_t(z)) \right. \\
&\quad \left. + \frac{1}{2} \sum_{r,s \in I_{\mathfrak{g}}} \sum_{i \in I_B} \kappa_i \theta_{r,t}^i(z) \theta_{s,t}^i(z) \mathcal{Y}(X_r X_s \Theta_t(z)v, f_t(z)) \right\} dt \\
&\quad + (f'_t(z))^{h_\Lambda} \sum_{i \in I_B} \left\{ \left( h_\Lambda \frac{f''_t(z)}{f'_t(z)} + f_t^i(z) \frac{\partial}{\partial f_t(z)} \right) \mathcal{Y}(\Theta_t(z)v, f_t(z)) \right. \\
&\quad \left. + \sum_{r \in I_{\mathfrak{g}}} \theta_{r,t}^i(z) \mathcal{Y}(X_r \Theta_t(z)v, f_t(z)) \right\} dB_t^{(i)}.
\end{aligned}$$

Comparing the coefficients of each  $dB_t^{(i)}$  in Eqs. (3.13) and (3.14), we find

$$\begin{aligned}
(3.15) \quad & [\sigma_i, \mathcal{Y}(\Theta_t(z)v, f_t(z))] = - \left( h_\Lambda \frac{f''_t(z)}{f'_t(z)} + f_t^i(z) \frac{\partial}{\partial f_t(z)} \right) \mathcal{Y}(\Theta_t(z)v, f_t(z)) \\
&\quad - \sum_{r \in I_{\mathfrak{g}}} \theta_{r,t}^i(z) \mathcal{Y}(X_r \Theta_t(z)v, f_t(z)).
\end{aligned}$$

The primary field conditions, Eqs. (2.21) and (2.22), help us determine  $f_t^i(z)$  and  $\theta_{r,t}^i(z)$  for a given  $\sigma_i$ . For example, if  $\sigma_i = L_n$ , we have

$$(3.16) \quad f_t^i(z) = -(f_t(z))^{n+1},$$

$$(3.17) \quad \theta_{r,t}^i(z) = 0, \quad \text{for all } r \in I_{\mathfrak{g}}.$$

Also, if  $\sigma_i = -(X_r)_n$ , we have

$$(3.18) \quad f_t^i(z) = 0,$$

$$(3.19) \quad \theta_{r,t}^i(z) = (f_t(z))^n,$$

$$(3.20) \quad \theta_{s,t}^i(z) = 0, \quad \text{for } s \neq r.$$

These give part of stochastic differential equations on  $f_t(z)$  and  $\theta_{r,t}(z)$ . Comparing the coefficients of  $dt$  in Eqs. (3.13) and (3.14), we can fully determine these stochastic

differential equations, in principle. In the next section, we complete this program for a certain choice of  $\mu, \sigma_i$ .

#### 4. DERIVATIONS OF STOCHASTIC DIFFERENTIAL EQUATIONS

Let  $\{X_r\}_{r \in I_{\mathfrak{g}}}$  be an orthonormal basis of  $\mathfrak{g}$  with respect to the symmetric invariant bilinear form  $(\cdot|\cdot)$ . We prepare mutually independent Brownian motions  $B_t^{(i)}$  labeled by  $i \in I_B := \{0\} \cup I_{\mathfrak{g}}$ , identifying  $I_{\mathfrak{g}}$  with a subset of  $I_B$ , and let  $\kappa_i > 0$  for  $i \in I_B$  be the variance of  $B_t^{(i)}$ .

We also assume that  $\mu = -2L_{-2n}$ ,  $\sigma_0 = L_{-n}$ , and  $\sigma_r = -(X_r)_{-n}$  for  $i \in I_{\mathfrak{g}}$  with  $n \in \mathbb{Z}_{>0}$ . Then the Ito process  $\mathcal{G}_t$  on  $\widehat{G}_{\text{vir}}^{\leq 0}$  is described by

$$(4.1) \quad \mathcal{G}_t^{-1} d\mathcal{G}_t = \left[ -2L_{-2n} + \frac{1}{2}\kappa_0 L_{-n}^2 + \frac{1}{2} \sum_{r \in I_{\mathfrak{g}}} \kappa_r (X_r)_{-n}^2 \right] dt + L_{-n} dB_t^{(0)} - \sum_{r \in I_{\mathfrak{g}}} (X_r)_{-n} dB_t^{(r)}.$$

We remark that, even if  $n = 1$ , the above Ito process  $\mathcal{G}_t$  is different from the one considered in [Ras07], which does not contain Brownian motion along  $L_{-1}$ . The last discussion in the previous section suggests that

$$(4.2) \quad f_t^0(z) = -(f_t(z))^{-n+1}, \quad \theta_{r,t}^0(z) = 0, \quad r \in I_{\mathfrak{g}},$$

$$(4.3) \quad f_t^r(z) = 0, \quad \theta_{r,t}^r(z) = (f_t(z))^{-n}, \quad \theta_{s,t}^r(z) = 0, \quad r \in I_{\mathfrak{g}}, \quad s \neq r.$$

These results simplify Eq. (3.14) to

$$(4.4) \quad d\left( (f_t'(z))^{h_{\Lambda}} \mathcal{Y}(\Theta_t(z)v, f_t(z)) \right) = (f_t'(z))^{h_{\Lambda}} \left\{ \left( h_{\Lambda} \frac{\bar{f}_t'(z)}{f_t'(z)} + \frac{1}{2} h_{\Lambda} (h_{\Lambda} - 1) (n-1)^2 \kappa_0 (f_t(z))^{-2n} + (\bar{f}_t(z) + h_{\Lambda} \kappa_0 (-n+1) (f_t(z))^{-2n+1}) \frac{\partial}{\partial f_t(z)} + \frac{1}{2} \kappa_0 (f_t(z))^{-2n+2} \frac{\partial^2}{\partial f_t(z)^2} \right) \mathcal{Y}(\Theta_t(z)v, f_t(z)) + \sum_{r \in I_{\mathfrak{g}}} \bar{\theta}_{r,t}(z) \mathcal{Y}(X_r \Theta_t(z)v, f_t(z)) + \frac{1}{2} \sum_{r \in I_{\mathfrak{g}}} \kappa_r (f_t(z))^{-2n} \mathcal{Y}(X_r^2 \Theta_t(z)v, f_t(z)) \right\} dt + (\text{terms proportional to } dB_t^{(i)}).$$

The increment in Eq. (3.13) becomes

$$\begin{aligned}
(4.5) \quad & d(\mathcal{G}_t^{-1}\mathcal{Y}(v, z)\mathcal{G}_t) \\
&= (f'_t(z))^{h_\Lambda} \left\{ \left( 2 \left( h_\Lambda(-2n+1) + \frac{\kappa_0}{2}(h_\Lambda^2(n-1)^2 + h_\Lambda n(n-1)) \right) (f_t(z))^{-2n} \right. \right. \\
&\quad \left. \left. + \left( 2 + \frac{\kappa_0}{2}(-n+1) + h_\Lambda \kappa_0(-n+1) \right) f_t(z)^{-2n+1} \frac{\partial}{\partial f_t(z)} \right. \right. \\
&\quad \left. \left. + \frac{1}{2} \kappa_0 (f_t(z))^{-2n+2} \frac{\partial^2}{\partial f_t(z)^2} \right) \mathcal{Y}(\Theta_t(z)v, f_t(z)) \right. \\
&\quad \left. + \frac{1}{2} \sum_{r \in I_{\mathfrak{g}}} \kappa_r (f_t(z))^{-2n} \mathcal{Y}(X_r^2 \Theta_t(z)v, f_t(z)) \right\} dt \\
&\quad + (\text{terms proportional to } dB_t^{(i)}).
\end{aligned}$$

Then we compare the coefficients of  $dt$  to obtain

$$(4.6) \quad \bar{f}_t(z) = \left( 2 + \frac{1}{2} \kappa_0(-n+1) \right) (f_t(z))^{-2n+1},$$

$$(4.7) \quad \bar{\theta}_{r,t}(z) = 0, \quad r \in I_{\mathfrak{g}}.$$

Therefore, the Ito processes  $f_t(z)$  and  $\theta_{k,t}(z)$  satisfy the following stochastic differential equations:

$$(4.8) \quad df_t(z) = \left( 2 - \frac{1}{2} \kappa_0(n-1) \right) (f_t(z))^{-2n+1} dt - (f_t(z))^{-n+1} dB_t^{(0)},$$

$$(4.9) \quad d\theta_{r,t}(z) = (f_t(z))^{-n} dB_t^{(r)}, \quad r \in I_{\mathfrak{g}}.$$

If  $n = 1$ , the differential equations (4.8) and (4.9) reduce to the ones considered in [ABI11]. Therefore, one can regard the set of equations (4.8) and (4.9) as generalization of the SLE-type growth process obtained in [ABI11].

If we define an Ito process  $g_t(z)$  by

$$(4.10) \quad g_t(z) = f_t(z)^n + nB_t^{(0)},$$

we obtain the following set of stochastic differential equations:

$$(4.11) \quad dg_t(z) = \frac{2n}{g_t(z) - nB_t^{(0)}} dt,$$

$$(4.12) \quad d\theta_{r,t}(z) = \frac{dB_t^{(r)}}{g_t(z) - nB_t^{(0)}}, \quad r \in I_{\mathfrak{g}}.$$

It is remarkable that Eq. (4.11) is nothing but the stochastic Loewner equation (1.1) with a rescaled time variable, but we have to comment that here the introduced  $g_t(z)$  cannot be a uniformizing map in the sense of [BB03a] unless  $n = 1$ , while  $f_t(z)$  can be. To see this, let us consider the automorphism group of an algebra of formal power series. Let  $\mathcal{O} := \mathbb{C}[[w]] = \varprojlim \mathbb{C}[w]/(w^n)$  be a complete  $\mathbb{C}$ -algebra. A continuous automorphism of  $\mathcal{O}$  is identified with the image of the topological generator  $w$  of  $\mathcal{O}$ . Under this identification, the group of continuous automorphisms  $\text{Aut}\mathcal{O}$  of  $\mathcal{O}$  is described as

$$(4.13) \quad \text{Aut}\mathcal{O} \simeq \{w \mapsto a_1 w + a_2 w^2 + \dots \mid a_1 \in \mathbb{C}^\times, a_i \in \mathbb{C} \ (i \geq 2)\}.$$

By setting  $w = 1/z$ , we regard the formal disk  $D = \text{Spec } \mathcal{O}$  as the formal neighborhood at infinity. Then  $f_t(z)$  constructed in our formulation defines a continuous automorphism of  $\mathcal{O}$  identified with

$$(4.14) \quad w \mapsto \frac{1}{f(1/w)},$$

while  $g_t(z)$  does not if  $n \geq 2$ . Indeed,  $1/(g_t(1/w))$  begins from  $w^n$  in  $\mathcal{O}$ , and, in particular, it is not invertible.

## 5. NULL VECTORS AND MARTINGALE CONDITIONS

For a certain vector  $w \in (L_{\mathfrak{g},k}(\Lambda))_{h_\Lambda}$ ,  $\mathcal{G}_t w$  is an Ito process on the representation  $L_{\mathfrak{g},k}(\Lambda)$ . It is an important problem when this random process is a martingale. To explain its importance, let us consider a correlation function or, precisely, a correlation-function-valued random process of the form

$$(5.1) \quad \langle 0 | \mathcal{Y}(v_1, z_1) \cdots \mathcal{Y}(v_N, z_N) \mathcal{G}_t w \rangle.$$

Here we take  $v_i \in (L_{\mathfrak{g},k}(\Lambda_i))_{h_{\Lambda_i}}$  and we let  $\langle 0 |$  be the dual of the vacuum vector  $|0\rangle \in L_{\mathfrak{g},k}$ . From the transformation rule of primary fields, Eqs. (2.21) and (2.22), and the translation invariance of the vacuum vector, the above function is the same as

$$(5.2) \quad \prod_{i=1}^N (f'_t(z_i))^{h_{\Lambda_i}} \langle 0 | \mathcal{Y}(\Theta_t(z) v_1, f_t(z_1)) \cdots \mathcal{Y}(\Theta_t(z) v_N, f_t(z_N)) w \rangle$$

As is demonstrated in [BB03a], the expectation value of Eq. (5.2) has a probability theoretical interpretation. If  $\mathcal{G}_t w$  is a martingale, the expectation value of Eq. (5.1) does not depend on time, yielding

$$(5.3) \quad \mathbb{E}[\langle 0 | \mathcal{Y}(v_1, z_1) \cdots \mathcal{Y}(v_N, z_N) \mathcal{G}_t w \rangle] = \langle 0 | \mathcal{Y}(v_1, z_1) \cdots \mathcal{Y}(v_N, z_N) w \rangle,$$

the right-hand side of which is a purely algebraic object computed in CFT. Therefore, the martingale property of  $\mathcal{G}_t$  is significant in bridging probability theory to CFT via SLE/CFT correspondence.

For our  $\mathcal{G}_t$  designed in Eq. (4.1), the random process  $\mathcal{G}_t w$  on  $L_{\mathfrak{g},k}(\Lambda)$  is a martingale if and only if the vector

$$(5.4) \quad \left[ -2L_{-2n} + \frac{1}{2} \kappa_0 L_{-n}^2 + \frac{1}{2} \sum_{r \in I_{\mathfrak{g}}} \kappa_r (X_r)_{-n}^2 \right] w$$

is a null vector in the corresponding Weyl module. For  $n = 1$ , the conditions for the above vector to be a singular vector have been analyzed in [ABI11] for the case when  $\kappa_r$  are the same constant for all  $r \in I_{\mathfrak{g}}$ .

Here, we give examples of null vectors of the above form for  $n = 2$  in the basic representations of  $\widehat{\mathfrak{sl}}_2$  and  $\widehat{\mathfrak{sl}}_3$ . We take an orthonormal basis of  $\mathfrak{sl}_2$  as

$$(5.5) \quad X_1 = \frac{1}{\sqrt{2}} H, \quad X_2 = \frac{1}{\sqrt{2}} (E + F), \quad X_3 = \frac{i}{\sqrt{2}} (E - F).$$

Here  $E$ ,  $H$ , and  $F$  are the standard basis

$$(5.6) \quad E = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad H = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad F = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix},$$

and the symmetric invariant bilinear form is given by  $(X|Y) = \text{Tr}(XY)$  for  $X, Y \in \mathfrak{sl}_2$ . Then the problem is whether there is a null vector in  $L_{\mathfrak{sl}_2,1}$  of the form

$$(5.7) \quad \left[ -2L_{-4} + \frac{1}{2}\kappa_0 L_{-2}^2 + \frac{1}{2} \sum_{r=1}^3 \kappa_r (X_r)_{-2}^2 \right] |0\rangle$$

with  $\kappa_i > 0$  for  $i = 0, 1, 2, 3$ .

Because the basic representation  $L_{\mathfrak{sl}_2,1}$  of  $\widehat{\mathfrak{sl}}_2$  is isomorphic to the lattice VOA  $V_Q$  associated with the root lattice  $Q = \mathbb{Z}\alpha$  with  $(\alpha|\alpha) = 2$  of  $\mathfrak{sl}_2$  [FK80], it is convenient to express the operation of  $\widehat{\mathfrak{sl}}_2$  in terms of the free Bose field and vertex operators. The isomorphism is described by the following assignment:

$$(5.8) \quad E(z) \mapsto \Gamma_\alpha(z), \quad H(z) \mapsto \alpha(z), \quad F(z) \mapsto \Gamma_{-\alpha}(z).$$

Here  $\alpha(z)$  is the free Bose field, and  $\Gamma_{\pm\alpha}(z)$  are vertex operators of charge  $\pm\alpha \in Q$ . Then the Virasoro field  $L(z)$  is given by the Sugawara construction, which yields

$$(5.9) \quad L(z) = \frac{1}{4} : \alpha(z)^2 : .$$

In the notion of VOA, the Virasoro field is  $\mathcal{Y}(L_{-2}|0\rangle, z)$  corresponding to the Virasoro vector  $L_{-2}|0\rangle$ . By the translation covariance of fields we have

$$(5.10) \quad \mathcal{Y}(L_{-4}|0\rangle, z) = \frac{1}{2} \partial^2 L(z) = \frac{1}{4} ( : \partial^2 \alpha(z) \alpha(z) : + : \partial \alpha(z)^2 : ).$$

By computation of the OPE, we can get

$$(5.11) \quad \begin{aligned} \mathcal{Y}(L_{-2}^2|0\rangle, z) &= : L(z)^2 : \\ &= \frac{1}{4} : \partial^2 \alpha(z) \alpha(z) : + \frac{1}{16} : \alpha(z)^4 : , \end{aligned}$$

$$(5.12) \quad \begin{aligned} \mathcal{Y}(E_{-2}F_{-2}|0\rangle, z) &= : \partial \Gamma_\alpha(z) \partial \Gamma_{-\alpha}(z) : \\ &= \frac{1}{12} ( \partial^3 \alpha(z) + 2 : \partial^2 \alpha(z) \alpha(z) : + 3 : \partial \alpha(z)^2 : - : \alpha(z)^4 : ) \end{aligned}$$

$$(5.13) \quad \begin{aligned} \mathcal{Y}(F_{-2}E_{-2}|0\rangle, z) &= : \partial \Gamma_{-\alpha}(z) \partial \Gamma_\alpha(z) : \\ &= \frac{1}{12} ( -\partial^3 \alpha(z) + 2 : \partial^2 \alpha(z) \alpha(z) : + 3 : \partial \alpha(z)^2 : - : \alpha(z)^4 : ) \end{aligned}$$

$$(5.14) \quad \mathcal{Y}(H_{-2}^2|0\rangle, z) = : \partial \alpha(z)^2 : .$$

Combining these, we can see that

$$(5.15) \quad \left( -2L_{-4} + \frac{4}{3}L_{-2}^2 + \frac{1}{2} \sum_{r=1}^3 (X_r)_{-2}^2 \right) |0\rangle = 0$$

in  $L_{\mathfrak{sl}_2,1}$ . Therefore,  $\mathcal{G}_t|0\rangle$  corresponding to  $n = 2$ ,  $\kappa_0 = \frac{8}{3}$ , and  $\kappa_1 = \kappa_2 = \kappa_3 = 1$  is a martingale.

We can also look for a null vector of the form of Eq. (5.4) in  $L_{\mathfrak{sl}_3,1}$  by similar computation. An orthonormal basis of  $\mathfrak{sl}_3$  is taken as

$$(5.16) \quad X_1 = \frac{1}{\sqrt{2}}H_1, \quad X_2 = \frac{1}{\sqrt{6}}(H_1 + 2H_2),$$

$$(5.17) \quad X_3 = \frac{1}{\sqrt{2}}(E_1 + F_1), \quad X_4 = \frac{i}{\sqrt{2}}(E_1 - F_1),$$

$$(5.18) \quad X_5 = \frac{1}{\sqrt{2}}(E_2 + F_2), \quad X_6 = \frac{i}{\sqrt{2}}(E_2 - F_2),$$

$$(5.19) \quad X_7 = \frac{1}{\sqrt{2}}([E_1, E_2] + [F_2, F_1]) \quad X_8 = \frac{i}{\sqrt{2}}([E_1, E_2] - [F_2, F_1]).$$

Here  $E_i$ ,  $H_i$ , and  $F_i$  are defined by

$$(5.20) \quad E_1 = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad H_1 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \quad F_1 = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$(5.21) \quad E_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}, \quad H_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}, \quad F_2 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix},$$

and the symmetric invariant bilinear form is given by the same formula  $(X|Y) = \text{Tr}(XY)$  for  $X, Y \in \mathfrak{sl}_3$  as in the case of  $\mathfrak{sl}_2$ . Again,  $L_{\mathfrak{sl}_3,1}$  is isomorphic to the lattice VOA associated to the root lattice  $Q = \mathbb{Z}\alpha_1 + \mathbb{Z}\alpha_2$  of type  $A_2$ . The action of  $\widehat{\mathfrak{sl}}_3$  is described by the assignment

$$(5.22) \quad E_i(z) \mapsto \Gamma_{\alpha_i}(z), \quad H_i(z) \mapsto \alpha_i(z), \quad F_i(z) \mapsto \Gamma_{-\alpha_i}(z)$$

for  $i = 1, 2$ . By similar but more complicated computation than in the case of  $\widehat{\mathfrak{sl}}_2$ , we find that

$$(5.23) \quad \left[ -2L_{-4} + \frac{6}{5}L_{-2}^2 + \frac{2}{5} \sum_{r=1}^8 (X_r)_{-2}^2 \right] |0\rangle = 0$$

in  $L_{\mathfrak{sl}_3,1}$ . Therefore, the random process  $\mathcal{G}_t|0\rangle$  in  $L_{\mathfrak{sl}_3,1}$  is a martingale if  $\kappa_0 = \frac{12}{5}$  and  $\kappa_i = \frac{4}{5}$  for  $i = 1, \dots, 8$ .

From the above observations for  $\widehat{\mathfrak{sl}}_2$  and  $\widehat{\mathfrak{sl}}_3$ , we make a conjecture that there is a null vector of the form

$$(5.24) \quad \left( -2L_{-4} + \frac{\kappa}{2}L_{-2}^2 + \frac{\tau}{2} \sum_{r=1}^{n^2-1} (X_r)_{-2}^2 \right) |0\rangle$$

in the basic representation  $L_{\mathfrak{sl}_n,1}$  of  $\widehat{\mathfrak{sl}}_n$ . Here  $\{X_r\}_{r=1}^{n^2-1}$  is an orthonormal basis of  $\mathfrak{sl}_n$  with respect to the trace form  $(X|Y) = \text{Tr}(XY)$  for  $X, Y \in \mathfrak{sl}_n$ .

The situation is more subtle for a spin representation of level 1. It may be impossible to construct a null vector of the form of Eq. (5.4) in  $L_{\mathfrak{g},1}(\Lambda)$  other than  $\Lambda = 0$ . Indeed, we have confirmed that there does not exist such a null vector in the spin- $\frac{1}{2}$  representation  $L_{\mathfrak{sl}_2,1}(\Lambda_1)$  of  $\widehat{\mathfrak{sl}}_2$ .

**Acknowledgements.** The author is grateful to K. Sakai and R. Sato for fruitful discussions. This work was supported by a Grant-in-Aid for JSPS Fellows (Grant No. 17J09658).

## REFERENCES

- [ABI11] A. Alekseev, A. Bytsko, and K. Izyurov. On SLE martingales in boundary WZW models. *Lett. Math. Phys.*, 97:243–261, 2011.
- [App14] David Applebaum. *Probability on Compact Lie Groups*, volume 70 of *Probability Theory and Stochastic Modeling*. Springer, 2014.
- [BB02] M. Bauer and D. Bernard. SLE $_{\kappa}$  growth processes and conformal field theories. *Phys. Lett. B*, 543:135–138, 2002.
- [BB03a] M. Bauer and D. Bernard. Conformal field theories of stochastic Loewner evolutions. *Commun. Math. Phys.*, 239:493–521, 2003.
- [BB03b] M. Bauer and D. Bernard. SLE martingales and the Virasoro algebra. *Phys. Lett. B*, 557:309–316, 2003.
- [BB04] Michel Bauer and Denis Bernard. Conformal transformations and the SLE partition function martingale. *Ann. Henri Poincaré*, 5:289–326, 2004.
- [BBK05] M. Bauer, D. Bernard, and K. Kytölä. Multiple Schramm-Loewner evolutions and statistical mechanics martingales. *J. Stat. Phys.*, 120:1125–1163, 2005.
- [BGLW05] E. Bettelheim, I. A. Gruzberg, A. W. W. Ludwig, and P. Wiegmann. Stochastic Loewner evolution for conformal field theories with Lie group symmetries. *Phys. Rev. Lett.*, 95:251601, 2005.
- [BPZ84] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov. Infinite conformal symmetry in two-dimensional quantum field theory. *Nucl. Phys. B*, 241:333–380, 1984.
- [Car86] J. L. Cardy. Effect of boundary conditions on the operator center of two-dimensional conformally invariant theories. *Nucl. Phys. B*, 275:200–218, 1986.
- [Car89] J. L. Cardy. Boundary conditions, fusion rules and the Verlinde formula. *Nucl. Phys. B*, 324:581–596, 1989.
- [Car92] J. L. Cardy. Critical percolation in finite geometries. *J. Phys. A: Math. Gen.*, 25:L201–L206, 1992.
- [CDCH<sup>+</sup>14] D. Chelkak, H. Duminil-Copin, C. Hongler, A. Kemppainen, and S. Smirnov. Convergence of Ising interfaces to Schramm’s SLE curves. *Comptes Rendus Mathématique*, 352:157–161, 2014.
- [DFMS97] Philippe Di Francesco, Pierre Mathieu, and David Sénéchal. *Conformal Field Theory*. Graduate Texts in Contemporary Physics. Springer-Verlag New York, Inc., 1997.
- [Dub15a] J. Dubédat. SLE and Virasoro representations: Fusion. *Commun. Math. Phys.*, 336:761–809, 2015.
- [Dub15b] J. Dubédat. SLE and Virasoro representations: Localization. *Commun. Math. Phys.*, 336:695–760, 2015.
- [FBZ04] E. Frenkel and D. Ben-Zvi. *Vertex Algebras and Algebraic Curves*, volume 88 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2nd edition, 2004.
- [FK80] I. B. Frenkel and V. G. Kac. Basic representations of affine Lie algebras and dual resonance models. *Invent. Math.*, 62:23–66, 1980.
- [FK04] R. Friedrich and J. Kalkkinen. On conformal field theory and stochastic Loewner evolution. *Nucl. Phys. B*, 687:279–302, 2004.
- [Fri04] R. Friedrich. On connections of conformal field theory and stochastic Loewner evolution, 2004. arXiv:math-ph/0410029.
- [Fuk17] Yoshiki Fukusumi. Multiple Schramm-Loewner evolutions for coset Wess-Zumino-Witten models, 2017. arXiv:1704.06006.
- [FZ92] I. B. Frenkel and Y. Zhu. Vertex operator algebras associated to representations of affine and Virasoro algebras. *Duke Math. J.*, 66:123–168, 1992.
- [Kac98] V. Kac. *Vertex Algebras for Beginners*, volume 10 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2nd edition, 1998.
- [Kon03] M. Kontsevich. CFT, SLE and phase boundaries, 2003. Oberwolfach Arbeitstagung.

- [Kos18] S. Koshida. Local martingales associated with Schramm-Loewner evolutions with internal symmetry. *J. Math. Phys.*, 59:101703, 2018. arXiv:1803.06808.
- [Kyt07] K. Kytölä. Virasoro module structure of local martingales of SLE variants. *Rev. Math. Phys.*, 5:455–509, 2007.
- [Law04] G. F. Lawler. An introduction to the stochastic Loewner evolution. In *Random Walks and Geometry*. De Gruyter, 2004.
- [Law05] Gregory F. Lawler. *Conformally Invariant Processes in the Plane*, volume 114 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2005.
- [LR04] F. Lesage and J. Rasmussen. SLE-type growth processes and the Yang-Lee singularity. *J. Math. Phys.*, 45:3040–3048, 2004.
- [MARR04] A. Moghimi-Araghi, M. A. Rajabpour, and S. Rouhani. Logarithmic conformal null vectors and SLE. *Phys. Lett. B*, 600:298–301, 2004.
- [Naz12] A. Nazarov. Schramm-Loewner evolution martingales in coset conformal field theory. *JETP Letters*, 96:90–93, 2012.
- [NR05] J. Nagi and J. Rasmussen. On stochastic evolutions and superconformal field theory. *Nucl. Phys. B*, 704:475–489, 2005.
- [Ras04a] J. Rasmussen. Note on stochastic Löwner evolutions and logarithmic conformal field theory. *J. Stat. Mech.*, page P09007, 2004.
- [Ras04b] J. Rasmussen. Stochastic evolutions in superspace and superconformal field theory. *Lett. Math. Phys.*, 68:41–52, 2004.
- [Ras07] J. Rasmussen. On  $SU(2)$  Wess-Zumino-Witten models and stochastic evolutions. *Afr. J. Math. Phys.*, 4:1–9, 2007.
- [RS05] S. Rohde and O. Schramm. Basic properties of SLE. *Ann. Math.*, 161:883–924, 2005.
- [Sak13] K. Sakai. Multiple Schramm-Loewner evolutions for conformal field theories with Lie algebra symmetries. *Nucl. Phys. B*, 867:429–447, 2013.
- [Sch00] O. Schramm. Scaling limits of loop-erased random walks and uniform spanning trees. *Israel J. Math.*, 118:221–288, 2000.
- [Smi01] S. Smirnov. Critical percolation in the plane: conformal invariance, Cardy’s formula, scaling limits. *C. R. Acad. Sci. Paris*, 333:239–244, 2001.
- [Smi06] Stanislav Smirnov. Towards conformal invariance of 2D lattice models. In *Proceedings of the International Congress of Mathematicians (Madrid, August 22-30, 2006)*, pages 1421–1451. Eur. Math. Soc., Zülich, 2006.