

On Estimation of L_r -Norms in Gaussian White Noise Models

Yanjun Han, Jiantao Jiao, and Rajarshi Mukherjee*

November 30, 2021

Abstract

We provide a complete picture of asymptotically minimax estimation of L_r -norms (for any $r \geq 1$) of the mean in Gaussian white noise model over Nikolskii-Besov spaces. In this regard, we complement the work of [LNS99], who considered the cases of $r = 1$ (with poly-logarithmic gap between upper and lower bounds) and r even (with asymptotically sharp upper and lower bounds) over Hölder spaces. We additionally consider the case of asymptotically adaptive minimax estimation and demonstrate a difference between even and non-even r in terms of an investigator's ability to produce asymptotically adaptive minimax estimators without paying a penalty.

Contents

1	Introduction	2
1.1	Organization	3
1.2	Notation	4
2	Function Spaces and Approximation	4
3	Main Results	5
3.1	Non-even r	6
3.1.1	$r = 1$	7
3.1.2	$r > 1$, non-even	8
3.1.3	Adaptive Estimation	10
3.2	Even r	11
4	Discussion and Open Problems	12
5	Proof of Main Results	13
5.1	Proof of Theorem 3.1	13
5.2	Proof of Theorem 3.3	14
5.3	Proof of Theorem 3.2 and Theorem 3.4	16

*Yanjun Han is with the Department of Electrical Engineering, Stanford University, 350 Serra Mall, Stanford, CA 94305, USA. Email: yjhan@stanford.edu. Jiantao Jiao is with the Department of Electrical Engineering and Computer Sciences, University of California, Berkeley, 257M Cory Hall, Berkeley, CA 94720, USA. Email: jiantao@eecs.berkeley.edu. Rajarshi Mukherjee is with the Department of Biostatistics, Harvard T. H. Chan School of Public Health, Harvard University, 655 Huntington Avenue, Boston, MA 02115, USA. Email: ram521@mail.harvard.edu. Yanjun Han and Jiantao Jiao were partially supported by the NSF Center for Science of Information under Grant CCF-0939370, and Jiantao Jiao was partially supported by NSF Grant IIS-1901252.

5.3.1	Reduction to Parametric Submodel	16
5.3.2	Construction of Two Priors	18
5.3.3	Minimax Lower Bound in the Parametric Submodel	20
5.4	Proof of Theorem 3.5	22
6	Technical Lemmas	26
6.1	Proof of Lemma 5.1	31
6.2	Proof of Lemma 5.2	34
6.3	Proof of Lemma 5.7	38
6.4	Proof of Lemma 5.5	39
6.5	Proof of Lemma 5.6	40
7	Acknowledgement	42

1 Introduction

Estimation of functionals of data generating distributions is a fundamental problem in statistics. Whereas relevant issues in finite dimensional parametric models are comparatively well understood [BKB⁺93, VdV00], corresponding nonparametric analogues are often much more challenging and have attracted tremendous interest over the last two decades. In this regard, initial efforts have focused on the inference of linear and quadratic functionals in Gaussian white noise and density models and have contributed immensely to the foundations of ensuing research. We do not attempt to survey the extensive literature in this area. However, the interested reader can find a comprehensive snapshot of the literature in [HM87, BR88, DLM90, DN90, Fan91, BM95, KP96, Lau96, Nem00, CL03, CL04, CL05, TLRvdV08] and the references therein.

For treatment of more general smooth functionals in Gaussian White Noise model (for smoothness measured in terms of differentiability in L_2), the excellent monograph of [Nem00] provides detailed analyses and references of cases where efficient parametric rate of estimation is possible. Further, in recent times, some progress has also been made towards the understanding of more complex nonparametric functionals over substantially more general observational models. These include causal effect functionals in observational studies and mean functionals in missing data models. For more details, we refer to [RLTvdV08, RLM⁺16, MNR17], which considers a general recipe to yield minimax estimation of a large class of nonparametric functionals common in statistical literature. However, apart from general theory of estimating linear functionals, most of the research endeavors, at least from the point of view of optimality, have focused on “smooth functionals” (see [RLTvdV08] for a more discussions on general classes of “smooth functionals”).

In contrast, the results on the asymptotically minimax estimation of non-smooth functionals have been comparatively sporadic ([IK81, Kor91, KT12]). The paradigm got an impetus from the seminal papers of [LNS99] and [CL11] which considered the estimating of L_r -norms in Gaussian mean models. Subsequently, significant progress has been made regarding inference of non-smooth functionals in discrete distribution settings ([VV11, JVHW15, WY16, HJW16, JHW16, JVHW17]). However, even in the simpler setting of Gaussian white noise model, a complete picture of minimax optimality for estimating integrated non-smooth functionals remain unexplored. This paper is motivated by taking a step in that direction by providing a complete description of asymptotically minimax estimation of L_r -norms (for $r \geq 1$) of the mean in Gaussian white noise model over Nikolskii-Besov spaces. We additionally consider the case of adaptive minimax estimation and demonstrate a difference between even and non-even r in terms of an investigator’s ability to produce asymptotically adaptive minimax estimators without paying a penalty.

More specifically, we consider noisy observation $\{Y(t)\}_{t \in [0,1]}$ in the Gaussian white noise model with known variance σ^2 as

$$dY(t) = f(t)dt + \frac{\sigma}{\sqrt{n}}dB(t), \quad (1.1)$$

where $f : [0, 1] \rightarrow \mathbb{R}$ is the unknown mean function and $\{B(t)\}_{t \in [0,1]}$ is the standard Brownian motion on $[0, 1]$. The main goal of this paper is to consider adaptive minimax estimation of the L_r -norm of the mean function f (i.e. $\|f\|_r \triangleq (\int_{[0,1]} |f(t)|^r dt)^{1/r}$ for $r \geq 1$) over Nikolskii-Besov spaces $\mathcal{B}_{p,\infty}^s(L)$ in $L_p[0, 1]$, $p \geq 1$ of smoothness $s > 0$ (defined in Section 2). It is worth noting here, that the specific cases of the mean function f being uniformly bounded away from 0 is significantly easier since in that case a natural plug-in principle yields asymptotic optimality.

As mentioned earlier, significant progress towards understanding these specific functionals has been made in [LNS99] and [CL11]. In particular, [LNS99] considers estimation of the L_r -norm over Hölder spaces of known smoothness and demonstrate rate optimal minimax estimation for r even positive integers. For $r = 1$ their results are suboptimal and leave a poly-logarithmic gap between the upper and lower bounds for the rate of estimation. Moreover, for general non-even r , [LNS99] provides no particular estimator. Finally, their results are non-adaptive in nature and requires explicit knowledge of the smoothness index of the underlying function class. Our main contribution is improving the lower bound argument (over function spaces similar to Hölder balls) to close the gap in non-adaptive minimax estimation of L_r -norm of the signal function. Moreover, for general non-even $r \geq 1$, the analysis extends further to demonstrate adaptive minimax estimators without logarithmic penalties which are typical in smooth functional estimation problems. However, the situation is different for even integers r where our results show that a poly-logarithmic penalty is necessary. In this effort, the fundamental work of [CL11], which considered estimating the L_1 -norm of the mean of an n -dimensional multivariate Gaussian vector, serves as a major motivation. In Section 3 we comment more on the main motivating ideas from [CL11] as well as the fundamental differences and innovations.

The main results of this paper are summarized below.

- (a) We produce minimax rate optimal estimator L_r norm of the unknown mean function f in Gaussian White noise model (1.1) with known variance.
- (b) For non-even r , an accompanying adaptive minimax optimal estimator is also provided. In contrast, for even integers r , we argue along the lines of standard results from [Ing87, IS12] that at least poly-logarithmic penalty needs to be paid for adaptation. The lower bound on this penalty is not sharp in this regard and only serves to demonstrate the lack of adaptation without paying a price.
- (c) Similar to [CL11], both our upper and lower bounds rely on best polynomial approximations of suitable functions on the unit interval, which might be of independent interest.

1.1 Organization

The rest of the paper is organized as follows. In Section 2 we discuss function spaces relevant to this paper as well as best polynomial approximations of continuous functions on compact intervals along with properties of Hermite polynomials, which are useful ingredients in the construction of our estimators. Section 3 contains the main results of the paper and is divided into two main subsections based on the non-even or even nature of r while estimating $\|f\|_r$. For non-even $r \geq 1$ (Section 3.1), we first lay down the basic principles for $r = 1$ (Section 3.1.1) followed by the general

r (Section 3.1.2). The case of even r is well understood from [LNS99] and is mostly presented here (Section 3.2) for completeness and discussing issues of adaptive estimation. In Section 4 we discuss remaining issues and future directions. Proofs of the main theorems are collected in Section 5 followed by proofs of several technical lemmas in Section 6.

1.2 Notation

In this paper, Poly_K denotes the set of all polynomials over $[-1, 1]$ with real coefficients and degree at most K . For any finite set S we denote its cardinality by $|S|$. For a function defined on $[0, 1]$, for $1 \leq q < \infty$ we let $\|h\|_q := (\int_{[0,1]} |h(x)|^q dx)^{1/q}$ denote the L_q semi-norm of h , $\|h\|_\infty := \sup_{x \in [0,1]} |h(x)|$ the L_∞ semi-norm of h . We say $h \in L_q[0, 1]$ for $q \in [1, \infty]$ if $\|h\|_q < \infty$. The results in this paper are mostly asymptotic (in n) in nature and thus requires some standard asymptotic notations. If a_n and b_n are two sequences of real numbers then $a_n \gg b_n$ (and $a_n \ll b_n$) implies that $a_n/b_n \rightarrow \infty$ (and $a_n/b_n \rightarrow 0$) as $n \rightarrow \infty$, respectively. Similarly $a_n \gtrsim b_n$ (and $a_n \lesssim b_n$) implies that $\liminf a_n/b_n = C$ for some $C \in (0, \infty]$ (and $\limsup a_n/b_n = C$ for some $C \in [0, \infty)$). Alternatively, $a_n = o(b_n)$ will also imply $a_n \ll b_n$ and $a_n = O(b_n)$ will imply that $\limsup a_n/b_n = C$ for some $C \in [0, \infty)$. Finally we comment briefly on the various constants appearing throughout the text and proofs. Given that our primary results concern convergence rates of various estimators, we will not emphasize the role of constants throughout and rely on fairly generic notation for such constants. In particular, for any fixed tuple v of real numbers, $C(v)$ will denote a positive real number which depends on elements of v only. Finally, whenever we use the symbol \lesssim in the asymptotic sense above, the hidden positive constant C will depend on the known parameters of the problem.

2 Function Spaces and Approximation

We begin with some standard definitions of function spaces [DL93, HKPT12] that we work with throughout. In the study of nonparametric functional estimation problem, many studies were devoted to the case where f is assumed to lie in a Hölder ball defined as

$$H(s, L) \triangleq \left\{ f \in L^2[0, 1] : \frac{|f^{(r)}(x) - f^{(r)}(y)|}{|x - y|^\alpha} \leq L, \forall x \neq y \in [0, 1] \right\},$$

where $s = r + \alpha > 0$ is the smoothness parameter, $r \in \mathbb{N}, \alpha \in (0, 1]$. In this paper, we consider another function class which is close but not identical to the Hölder ball where the dependence of the upper and lower bounds on n matches. The r -th symmetric difference operator Δ_h^r is defined as [DL93]

$$\Delta_h^r f(x) = \sum_{k=0}^r (-1)^{r-k} \binom{r}{k} f\left(x + \left(k - \frac{r}{2}\right)h\right),$$

with the agreement that $\Delta_h^r f(x) = 0$ when either $x + \frac{r}{2}h$ or $x - \frac{r}{2}h$ does not belong to $[0, 1]$. Then the r -th order modulus of smoothness is defined as [DL93]

$$\omega^r(f, t)_p = \sup_{0 < h \leq t} \|\Delta_h^r f\|_p, \quad (2.1)$$

with $p \in [1, \infty]$. Now define the Besov norm of a function f as [BI79]

$$\|f\|_{\mathcal{B}_{p,q}^s} = \|f\|_p + \begin{cases} \left[\int_0^\infty \left(\frac{\omega^r(f,t)_p}{t^s} \right)^q \cdot \frac{dt}{t} \right]^{\frac{1}{q}} & 1 \leq q < \infty, \\ \sup_{t>0} \frac{\omega^r(f,t)_p}{t^s} & q = \infty \end{cases}, \quad (2.2)$$

with parameters $s > 0, p, q \in [1, \infty]$, and $r = \lfloor s \rfloor + 1$. Then the corresponding Besov ball is defined by

$$\mathcal{B}_{p,q}^s(L) \triangleq \{f \in L^p[0, 1] : \|f\|_{\mathcal{B}_{p,q}^s} \leq L\}.$$

We note that the definition of Besov ball is a generalization of Hölder ball via the relationship $H(s, L') = \mathcal{B}_{\infty,\infty}^s(L)$ for non-integer s . Moreover, by the monotonicity of L_p norms on $[0, 1]$, we have $\mathcal{B}_{p,q}^s(L) \supseteq \mathcal{B}_{p',q}^s(L)$ for $p \leq p'$. As a result, for $1 \leq p < \infty$, $\mathcal{B}_{p,\infty}^s(L) \supseteq \mathcal{B}_{\infty,\infty}^s(L)$ is a function class slightly larger than the Hölder ball $H(s, L')$. In this paper we work with $\mathcal{B}_{p,\infty}^s(L)$ for $s \geq 0, p \geq 1$. These spaces are related to Nikolskii-spaces (see [Nik12] for relevant embeddings) and hence we shall refer them to as Nikolskii-Besov spaces throughout.

Polynomial approximations of continuous functions on compact intervals around the origin plays an important role in this paper. To introduce the basic ideas, consider the following best degree- K polynomial approximation of $|\mu|^r$ on $[-1, 1]$:

$$\sum_{k=0}^K g_{K,k}^{(r)} \mu^k \triangleq \arg \min_{Q \in \text{Poly}_K} \max_{\mu \in [-1, 1]} |Q(\mu) - |\mu|^r|.$$

In order to estimate such polynomials based on a sample $X \sim \mathcal{N}(\mu, 1)$, we will need the notion of Hermite polynomials. In particular, the Hermite polynomial of degree k defined by

$$H_k(x) \triangleq (-1)^k \exp\left(\frac{x^2}{2}\right) \cdot \frac{d^k}{dx^k} \left[\exp\left(-\frac{x^2}{2}\right) \right]. \quad (2.3)$$

The properties of the Hermite polynomials in the context of estimating moments of Gaussian random variables will be important for us and are summarized in Lemma 6.1.

3 Main Results

We divide our main results in two subsections based on the non-even or even nature of r . In particular, the construction of our estimator changes according to this distinction of r . However, before we go into the details of these constructions, we need a few definitions.

Consider the kernel projection $f_h(x)$ of $f(x)$ defined as

$$f_h(x) = \int_0^1 f(u) \cdot \frac{1}{h} K_M\left(\frac{x-u}{h}\right) du, \quad x \in [0, 1], \quad (3.1)$$

where $K_M(\cdot)$ is a kernel which maps all polynomials of degree at most M to themselves, and $\int_0^1 |K_M(u)|^{M+1} du < \infty$. The choice of M will be clear from the statements of the main results in Section 3. We assume that K_M is supported on $[-\frac{1}{2}, \frac{1}{2}]$, and the boundary of $f_h(x)$ (i.e., $x \in [0, h/2]$ or $x \in [1 - h/2, 1]$) is handled using the same way as [LNS99].

The corresponding unbiased kernel estimator of $f_h(x)$ defined as

$$\tilde{f}_h(x) = \int_0^1 \frac{1}{h} K_M\left(\frac{x-u}{h}\right) dY(u)$$

admits a usual decomposition into deterministic and stochastic components as follows:

$$\tilde{f}_h(x) = f_h(x) + \lambda_h \xi_h(x). \quad (3.2)$$

Above

$$\begin{aligned}\lambda_h &= \sqrt{\mathbb{E} \left(\int_0^1 \frac{1}{h} K_M \left(\frac{x-u}{h} \right) \cdot \frac{\sigma}{\sqrt{n}} dB(u) \right)^2} = \frac{\sigma \|K_M\|_2}{\sqrt{nh}}, \\ \xi_h(x) &= \frac{1}{h\lambda_h} \int_0^1 K_M \left(\frac{x-u}{h} \right) \cdot \frac{\sigma}{\sqrt{n}} dB(u).\end{aligned}\tag{3.3}$$

Clearly $\xi_h(x) \sim \mathcal{N}(0, 1)$ and random variables $\xi_h(x)$ and $\xi_h(y)$ are independent when $|x - y| > h$.

The reason for introducing the kernel projection estimator is simple and standard in nonparametric statistics. In particular, for a suitable chosen bandwidth h , $\|f_h - f\|_r$ is small, and it suffices to consider estimation of $\|f_h\|_r$ based on the Gaussian model (3.2). Indeed, a crucial part is to estimate $|f_h(x)|^r$. Whereas, for r an even positive integer, this task is relatively simpler [LNS99], the case of non-even r poses a more subtle problem due to non-differentiability of the function $u \mapsto |u|^r$ near the origin. Consequently, for such cases, recent techniques for estimating non-smooth functionals needs to be employed.

3.1 Non-even r

In this case, the construction depends on the best polynomial approximation of the function $u \mapsto |u|^r$ over the interval $[-1, 1]$ and borrows heavily from a recent line of work by [CL11, JVHW15, HJW16, WY16]. The general principle of the construction goes along the following heuristic steps.

- Approximate f by a kernel projection f_h (as in (3.1)) and consider estimating $\|f_h\|_r$ instead at a cost of incurring a truncation bias.
- A “large value” of the kernel estimator $|\tilde{f}_h(x)|$ (referred to as “smooth regime” hereafter) gives indication of a corresponding “large value” of $|f_h(x)|$ and a plug-in type estimator for $|f_h(x)|^r$ is reasonable.
- A relatively “small value” of $|\tilde{f}_h(x)|$ (referred to as “non-smooth regime” hereafter) gives indication of a correspondingly “small value” of $|f_h(x)|$ and a plug-in type estimator for $|f_h(x)|^r$ is no-longer reasonable owing to the non-differentiable nature of the absolute function near the origin. In this case, similar to [CL11, JVHW15, WY16, HJW16], an estimator based on the best polynomial approximation of the function $u \mapsto |u|^r$ is employed.
- The final estimator integrates over this two regimes of $|\tilde{f}_h(x)|$ followed by an optimal choice of h to trade off squared bias and variance.

Below we make the program laid down above more concrete and refer readers to [JVHW15, HJW16] for a detailed discussion of the general principle of estimating non-smooth functionals. The same procedure also works for estimating other non-smooth nonparametric functionals, e.g., the differential entropy [HJWW17]. It turns out that the treatment for $r = 1$ is easier, more transparent and slightly different than general non-even $r > 1$. Consequently, we will present the case for $r = 1$ first for the sake of clarity followed by the more general case.

Any candidate estimator below will be defined by the parameter tuple (h, c_1, c_2, ϵ) where h is the bandwidth of the kernel projection (3.1) and (c_1, c_2, ϵ) are constants depending on the known parameters of the problem (i.e. the variance σ^2 and radius of Besov balls L) to be specified later. More specifically, c_1 will be chosen as large as possible whereas c_2 and ϵ will be desirably small. In general we suppress the dependence of our estimators on (c_1, c_2, ϵ) . However, our adaptive estimator makes a data driven choice of the bandwidth h . Therefore we index our estimator by this bandwidth, namely, T_h .

3.1.1 $r = 1$

We follow the general principle laid down above. Recall that $\{g_{K,k}^{(1)}\}_{k=0}^K$ are the coefficients of the best polynomial approximation of $u \mapsto |u|$ of degree K on $[-1, 1]$ and H_k is the Hermite polynomial of degree k . With this in mind, the construction of our estimator T_h for every bandwidth h can be described the following steps.

- (I) Using the sample splitting technique for the Brownian motion [Nem00] to obtain two independent observations $\tilde{f}_{h,1}(x)$ and $\tilde{f}_{h,2}(x)$ for any $x \in [0, 1]$. This reduces the effective sample size n by half and for simplicity of notation we redefine $n/2$ as n .
- (II) For any x define an estimator of $|f_h(x)|$ as

$$T_h(x) = |\tilde{f}_{h,1}(x)| \mathbb{1}(|\tilde{f}_{h,2}(x)| \geq c_1 \lambda_h \sqrt{\ln n}) + \tilde{P}(\tilde{f}_{h,1}(x)) \mathbb{1}(|\tilde{f}_{h,2}(x)| < c_1 \lambda_h \sqrt{\ln n}),$$

where

$$\begin{aligned} \tilde{P}(u) &= \min\{\max\{P(u), -n^{2\epsilon} \lambda_h\}, n^{2\epsilon} \lambda_h\}, \quad \text{with} \\ P(u) &= \sum_{k=0}^K g_{K,k}^{(1)} \left(2c_1 \lambda_h \sqrt{\ln n}\right)^{1-k} \cdot \lambda_h^k H_k\left(\frac{u}{\lambda_h}\right), \end{aligned}$$

where $K = \lceil c_2 \ln n \rceil$.

- (III) Finally, an estimator for $\|f\|_1$ is defined as

$$T_h \triangleq \min \left\{ L, \max \left\{ 0, \int_0^1 T_h(x) dx \right\} \right\}.$$

The following Theorem describes the mean squared error of estimating $\|f\|_1$ by T_h over $\mathcal{B}_{p,\infty}^s(L)$.

Theorem 3.1. *Choose $M > \lceil s \rceil$ and consider the corresponding kernel projection f_h based on K_M defined by 3.1. Suppose (h, c_1, c_2, ϵ) satisfy $4c_1^2 \geq c_2, c_2 \ln n \geq 1, c_1 > 8, 7c_2 \ln 2 < \epsilon \in (0, 1)$. Then for any $p \in [1, \infty]$, we have*

$$\left(\sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{E}_f (T_h - \|f\|_1)^2 \right)^{\frac{1}{2}} \leq C \left(h^s + \frac{1}{\sqrt{nh \ln n}} + \frac{1}{n^{(1-\epsilon)/2}} \right),$$

for a constant $C > 0$ depending on s, p, L, σ .

As an immediate consequence of Theorem 3.1, by choosing $\epsilon > 0$ sufficiently small (i.e., choose a small c_2) and by setting

$$h \asymp (n \ln n)^{-\frac{1}{2s+1}},$$

we have the following result.

Corollary 3.1. *Under the assumptions of Theorem 3.1, for $h \asymp (n \ln n)^{-\frac{1}{2s+1}}$ we have*

$$\left(\sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{E}_f (T_h - \|f\|_1)^2 \right)^{\frac{1}{2}} \leq C (n \ln n)^{-\frac{s}{2s+1}},$$

for a constant $C > 0$ depending on s, p, L, σ .

The same asymptotic upper bound was demonstrated with a different estimator by [LNS99]. However, their results were over Hölder Balls corresponding to $p = \infty$ and require a uniform upper bound on $\|f\|_\infty$ for all f . In contrast, we do not require any such knowledge of upper bound and produce results for any $p \in [1, \infty]$. Moreover, the next theorem shows that our results are rate optimal in terms of matching lower bounds for any $1 \leq p < \infty$.

Theorem 3.2. *For any $1 \leq p < \infty$, we have*

$$\left(\inf_{\hat{T}} \sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{E}_f \left(\hat{T} - \|f\|_1 \right)^2 \right)^{\frac{1}{2}} \geq C' (n \ln n)^{-\frac{s}{2s+1}},$$

for a constant $C' > 0$ depending on s, p, L, σ and where the infimum above is taken over all measurable maps of $\{Y(t)\}_{t \in [0,1]}$.

Corollary 3.1 along with Theorem 3.2 provide a complete picture of the minimax rate of estimation of $\|f\|_1$ over $\mathcal{B}_{p,\infty}^s(L)$ for any $1 \leq p < \infty$. We remark that our lower bound proof does not provide matching results for $p = \infty$, and thus the gap in the exact framework considered by [LNS99] remains. However, our result provides strong moral evidence that upper bound of Theorem 2.1 of [LNS99] is rate optimal and it is the lower bound that stands to be improved.

3.1.2 $r > 1$, non-even

For $r > 1$ not an even integer, the general philosophy of the construction of a candidate estimator T_h is similar to the case of $r = 1$ in Section 3.1.1 and follows in three steps as before. However, the simple plug-in principle employed in the smooth regime (i.e., when $|\tilde{f}_h(x)|$ is large) incurs significant bias. As a consequence, Step II of the construction needs to be modified based on the following heuristics of Taylor expansion. The reason why we require r to be non-even is that the function $u \mapsto |u|^r$ is not an analytical function in this case.

Using the notation from Section 3.1.1, the heuristic Taylor expansion gives

$$f_h(x)^r \approx \sum_{k=0}^R \frac{r(r-1) \cdots (r-k+1)}{k!} (\tilde{f}_{h,1}(x))^{r-k} (f_h(x) - \tilde{f}_{h,1}(x))^k, \quad (3.4)$$

where we choose $R = \lfloor 2r \rfloor$ to reduce the approximation error of (3.4) to a desired order (cf. Lemma 6.14). Based on this approximate Taylor expansion, the right hand side of (3.4) is a natural candidate estimator for $f_h(x)^r$. However, such an estimator is infeasible due to its dependence on unknown $f_h(x)$. Consequently, we replace $(f_h(x) - \tilde{f}_{h,1}(x))^k$ in (3.4) by an unbiased estimator constructed as follows: let $\tilde{f}_{h,2}(x)$ be an independent copy of $\tilde{f}_{h,1}(x)$ obtained via the sample splitting technique of the Brownian motion [Nem00]. Then Lemma 6.1 gives

$$\begin{aligned} \mathbb{E}_{\tilde{f}_{h,2}} \left[\sum_{j=0}^k \binom{k}{j} \lambda_h^j H_j \left(\frac{\tilde{f}_{h,2}(x)}{\lambda_h} \right) (-\tilde{f}_{h,1}(x))^{k-j} \right] &= \sum_{j=0}^k \binom{k}{j} f_h(x)^j (-\tilde{f}_{h,1}(x))^{k-j} \\ &= (f_h(x) - \tilde{f}_{h,1}(x))^k, \end{aligned}$$

i.e., the estimator inside the expectation is an unbiased estimator of $(f_h(x) - \tilde{f}_{h,1}(x))^k$.

With the above intuition and notation in mind, the construction of our estimator T_h for every bandwidth h can be described the following steps.

(I) Using the sample splitting technique for the Brownian motion [Nem00] to obtain three independent observations $\tilde{f}_{h,1}(x)$, $\tilde{f}_{h,2}(x)$ and $\tilde{f}_{h,3}(x)$ for any $x \in [0, 1]$. Redefine $n/3$ as n for simplicity.

(II) For any $x \in [0, 1]$ define an estimator of $|f_h(x)|^r$ as

$$\begin{aligned} T_h(x) = & \tilde{P}_r(\tilde{f}_{h,1}(x)) \mathbb{1}(|\tilde{f}_{h,3}(x)| \leq c_1 \lambda_h \sqrt{\ln n}) \\ & + S_{\lambda_h}(\tilde{f}_{h,1}(x), \tilde{f}_{h,2}(x)) \mathbb{1}(\tilde{f}_{h,3}(x) > c_1 \lambda_h \sqrt{\ln n}) \\ & + S_{\lambda_h}(-\tilde{f}_{h,1}(x), -\tilde{f}_{h,2}(x)) \mathbb{1}(\tilde{f}_{h,3}(x) < -c_1 \lambda_h \sqrt{\ln n}), \end{aligned}$$

where

$$\begin{aligned} S_{\lambda_h}(u, v) = & \sum_{k=0}^R \frac{r(r-1) \cdots (r-k+1)}{k!} u^{r-k} \cdot \left(\sum_{j=0}^k \binom{k}{j} \lambda_h^j H_j \left(\frac{v}{\lambda_h} \right) (-u)^{k-j} \right) \\ & \cdot \mathbb{1} \left(u \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n} \right), \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} \tilde{P}_r(t) = & \min\{\max\{P_r(t), -\lambda_h^r n^{2\epsilon}\}, \lambda_h^r n^{2\epsilon}\}, \quad \text{with} \\ P_r(t) = & \sum_{k=0}^K g_{K,k}^{(r)} \left(2c_1 \lambda_h \sqrt{\ln n} \right)^{r-k} \cdot \lambda_h^k H_k \left(\frac{t}{\lambda_h} \right), \end{aligned}$$

where $K = \lceil c_2 \ln n \rceil$, H_k is the Hermite polynomial of degree k and $\{g_{K,k}^{(r)}\}_{k=0}^K$ is the coefficient of the best polynomial approximation of $u \rightarrow |u|^r$ on $[-1, 1]$.

(III) Finally, the overall estimator for $\|f\|_r$ is

$$T_h \triangleq \min \left\{ L, \left(\max \left\{ 0, \int_0^1 T_h(x) dx \right\} \right)^{\frac{1}{r}} \right\}.$$

The following Theorem describes the optimal mean squared error of estimating $\|f\|_r$ by T_h over $\mathcal{B}_{p,\infty}^s(L)$.

Theorem 3.3. *Let $r > 1$ be non-even and $p \in [r, \infty]$. Choose $M > \lceil s \rceil$ and consider the corresponding kernel projection f_h based on K_M defined by (3.1). Suppose that (h, c_1, c_2, ϵ) satisfy $c_1^2 \geq 16$, $c_2 \ln n \geq 1$, $4c_1^2 \geq c_2$, $7c_2 \ln 2 < \epsilon \in (0, 1)$ and $h = (n \ln n)^{-\frac{1}{2s+1}}$. Then*

$$\left(\sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{E}_f (T_h - \|f\|_r)^2 \right)^{\frac{1}{2}} \leq C (n \ln n)^{-\frac{s}{2s+1}},$$

for a constant $C > 0$ depending on s, p, r, L, σ .

The next theorem shows that the upper bounds in Theorem 3.3 are rate optimal in terms of matching lower bounds for any $1 \leq p < \infty$.

Theorem 3.4. For any non-even $r > 1$ and $r \leq p < \infty$,

$$\left(\inf_{\hat{T}} \sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{E}_f \left(\hat{T} - \|f\|_r \right)^2 \right)^{\frac{1}{2}} \geq C' (n \ln n)^{-\frac{s}{2s+1}},$$

for a constant $C' > 0$ depending on s, p, r, L, σ and where the infimum above is taken over all measurable maps of $\{Y(t)\}_{t \in [0,1]}$.

Theorem 3.3 along with Theorem 3.4 provides a complete picture of the minimax rate of estimation of $\|f\|_r$ over $\mathcal{B}_{p,\infty}^s(L)$ for any non-even $r > 1$ and $r \leq p < \infty$. This is a generalization of the result in [LNS99].

3.1.3 Adaptive Estimation

It is worth noting that the choice of h in Corollary 3.1 and Theorem 3.3 depends explicitly on the smoothness index s . Consequently, the resulting rate of estimation by T_h is non-adaptive over different possibilities of smoothness. However, the experienced reader will notice the structure of errors in Theorem 3.1 for a general T_h indicates a possible data driven adaptive choice of bandwidth h . In particular, a Lepski type argument [Lep91, Lep92a, Lep92b, Lep93] is standard in such situations and turns out to be sufficient for our purpose when $r = 1$. The similar construction for general non-even integer $r > 1$ is more subtle due to the unavailability of a transparent error analysis as in Theorem 3.1. Consequently, we describe the adaptive procedure slightly more generally below—without specifically alluding to the case of $r = 1$.

Let $r \geq 1$ be non-even, and $s \in (0, s_{\max}]$ for some given $0 < s_{\max} < \infty$ and consider adaptive estimation of $\|f\|_r$ over $\cup_{s \in (0, s_{\max}]} \mathcal{B}_{p,\infty}^s(L)$ with a known L . The knowledge of s_{\max} will be necessary for construction of kernels providing optimal approximations in Nikolskii-Besov spaces. Given access to an upper bound L_{\max} , our construction of the adaptive estimator can also adapt to the scaling parameter L , by noting that the construction of T_h does not require the knowledge of L (except for the final truncation step, where L can be replaced by L_{\max} without affecting the multiplicative constants in Theorems 3.1 and 3.3). However, since the dependence of the minimax risk on L is not the main focus of this paper, we assume that L is known and do not elaborate on the adaptation to L . In this framework, a Lepski type construction for an adaptive estimator can be achieved as follows.

(I) Let

$$\mathcal{H} = [h_{\min}, h_{\max}] \cap \left\{1, \frac{1}{2}, \frac{1}{3}, \dots\right\}, \quad \text{where} \\ h_{\min} = (n \ln n)^{-1}, \quad h_{\max} = (n \ln n)^{-\frac{1}{2s_{\max}+1}}.$$

(II) Define the data-driven bandwidth

$$\hat{h} \triangleq \max \left\{ h \in \mathcal{H} : (T_h - T_{h'})^2 \leq C^* \frac{\lambda_{h'}^2}{\ln n}, \quad \forall h' \in \mathcal{H}, h' \leq h \right\}. \quad (3.6)$$

(III) The final estimator is $\hat{T} = T_{\hat{h}}$.

Our next theorem justifies the adaptive nature of \hat{T} .

Theorem 3.5. *Let $r \geq 1$ be non-even and $p \in [r, \infty]$. Choose $M > \lceil s_{\max} \rceil$ and consider the corresponding kernel projection f_h based on K_M defined by (3.1). If $4r\epsilon(2s_{\max} + 1) < 1$ and the constant C^* is large enough (depending on $(p, r, s_{\max}, c_1, c_2, \sigma, L, K_M)$),*

$$\left(\sup_{f \in \mathcal{B}_{p, \infty}^s(L)} \mathbb{E}_f \left(\hat{T} - \|f\|_r \right)^2 \right)^{\frac{1}{2}} \leq C(n \ln n)^{-\frac{s}{2s+1}},$$

for a constant $C > 0$ depending on $s, s_{\max}, p, r, L, \sigma, c_1, \epsilon, C^*$.

Theorem 3.5 shows the existence of an adaptive minimax estimator for $\|f\|_r$ without any penalty on the minimax rate. One of the main challenges in the proof of Theorem 3.5 is to demonstrate desired tail bounds of $\{T_h\}_{h \in \mathcal{H}}$, for the candidate estimators in $\{T_h\}_{h \in \mathcal{H}}$ rely on truncated Hermite polynomials of high degrees evaluated at Gaussian random variables.

3.2 Even r

The case of non-adaptive minimax estimation of $\|f\|_r$ for r an even positive integer can be obtained by methods described in [LNS99]. Although their results were obtained over the Hölder Balls, the case of Nikolskii-Besov type spaces that we consider are very similar due to the same nature of approximation error of f by f_h . However, for the sake of exposition and completeness, we provide the details here again.

The crux of the construction on the fact that for r an even positive integer, the function $u \mapsto |u|^r = u^r$ is analytic. Consequently, it is possible to construct unbiased estimator of μ^r based on samples from $X \sim \mathcal{N}(\mu, \sigma^2)$. In particular, if $X \sim \mathcal{N}(\mu, \sigma^2)$ then arguing as Lemma 4.4 of [LNS99], $\mathbb{E}((X + i\sigma\xi)^r | X)$, with $\xi \sim \mathcal{N}(0, 1)$ independent of X , is an unbiased estimator of μ^r . As a result, a sequence estimator for $\|f\|_r$ indexed by bandwidth h can now be constructed in the following steps.

- (I) Approximate f by a kernel projection $f_h(x)$ (as in (3.1)) and consider estimating $\|f_h\|_r$ instead at a cost of incurring a truncation bias.
- (II) Fix $\xi \sim \mathcal{N}(0, 1)$ independent of $B(t)$. For every $x \in [0, 1]$ estimate $f_h(x)^r$ by $T_h(x) \triangleq \mathbb{E}_\xi[(f_h(x) + i\lambda\xi)^r]$.
- (III) Estimate $\|f_h\|^r = \int_0^1 f_h^r(x) dx$ by $\int_0^1 T_h(x) dx$.
- (IV) Finally, an estimator for $\|f\|_r$ is defined as

$$T_h \triangleq \left(\max \left\{ 0, \int_0^1 T_h(x) dx \right\} \right)^{1/r}.$$

Note that the construction changes from the construction of estimators when r is non-zero only in the definition of $T_h(x)$ and indeed this is due to the ability to produce estimate of analytic functions of Gaussian means. The following theorem describes the optimal mean squared error of estimating $\|f\|_r$ by T_h over $\mathcal{B}_{p, \infty}^s(L)$.

Theorem 3.6. *Let $2 \leq r \leq p$ and r be even. Choose $M > \lceil s \rceil$ and consider the corresponding kernel projection f_h based on K_M in (3.1). Letting $h = n^{-1/(2s+1-1/r)}$,*

$$\left(\sup_{f \in \mathcal{B}_{p, \infty}^s(L)} \mathbb{E}_f (T_h - \|f\|_r)^2 \right)^{\frac{1}{2}} \leq C n^{-\frac{s}{2s+1-1/r}},$$

for a constant $C > 0$ depending on s, p, r, L, σ .

Recall from Section 2 that the approximation error $\|f - f_h\|_r$ is always bounded by $C(L, K_M)h^s$ for any $f \in \mathcal{B}_{p,\infty}^s(L)$. Hence, the proof of Theorem 3.6 can be obtained verbatim from the proof of Theorem 2.3 of [LNS99] and is hence omitted. In fact, proof of the lower bound in Theorem 2.3 of [LNS99] implies that the proposed T_h with $h = n^{-1/(2s+1-1/r)}$ is in fact asymptotically minimax rate optimal over any $\mathcal{B}_{p,\infty}^s(L)$ as well. Therefore, as before, it remains to explore adaptive minimax estimation over a collection of smoothness classes. In this regard, one can argue along the lines of standard results from [Ing87, IS12], that unlike non-even r , adaptation over a range of smoothness indices is not possible without paying a poly-logarithmic penalty. In particular, consider testing $H_0 : f \equiv 0$ vs $H_1 : f \in \mathcal{B}_{r,\infty}^s(L), \|f\|_r \geq \rho_n$ with s varying over a range of values $[s_{\min}, s_{\max}] \subseteq (0, \infty)$. Whereas the minimax testing rate of separation for ρ_n with known s equals $n^{-s/(2s+1-1/r)}$ (See [IS12] and proof of [Car13, Theorem 3.4 (b)] for details), i.e., the minimax rate of estimation of $\|f\|_r$, the adaptation over $[s_{\min}, s_{\max}]$ needs an additional penalty for ρ_n which equals $(\ln \ln n)^{C(r,s)}$ for a constant $C(r, s) > 0$ depending on r and s . The proof of this additional poly-logarithmic penalty is proved for $r = 2$ in [Spo98] (see also proof of [GN15, Theorem 8.1.1]). The proof of this additional penalty needed for adaptive hypothesis testing builds on the usual second moment method type lower bound argument for non-adaptive testing and involves putting an additional uniform prior on a suitably discretized subset $[s_{\min}, s_{\max}]$. Using similar ideas, the proof of a required penalty for adaptation for general even r -can be obtained by combining proof technique of [Car13, Theorem 3.4 (b)] for non-adaptive testing and adaptive lower bound arguments as in [Spo98, GN15]. The details are omitted and we simply comment on the implications of the result. Indeed, such a poly-logarithmic penalty needed for adaptive hypothesis testing in L_r -norm, yields a penalty for adaptive estimation of $\|f\|_r$ over $\mathcal{B}_{r,\infty}^s(L)$. We believe that this poly-logarithmic penalty is not sharp for adaptive estimation of $\|f\|_r$ norms for even r and only serves to demonstrate the lack of adaptation without paying a price. In particular, we have shown elsewhere that a careful application of Lepski's method along with an involved computation of all central moments of T_h can yield a penalty which behaves $(\ln n)^{C'(r,s)}$ for a constant $C'(r, s) > 0$ depending on r and s . In future work, we plan to explore the exact nature of this poly-logarithmic penalty necessary for adaptation. Finally, careful readers will notice that, although the poly-logarithmic penalty on adaptive hypothesis testing is ubiquitous for any r , the result is interesting to us only for even values of r , since otherwise the minimax rate of estimation of $\|f\|_r$ is strictly slower than minimax rate of testing in L_r -norm.

4 Discussion and Open Problems

In this paper we complement the results in [LNS99] to provide a complete picture of asymptotically minimax estimation of L_r -norm of the mean in a Gaussian White Noise model. Unlike [LNS99], our results are rate optimal from both perspectives of upper and lower bounds. In this effort, best polynomial approximation of non-smooth functions plays a major role and might be of independent interest.

Several interesting questions remain open as challenging future directions. In particular, closing the lower bound gap over Hölder spaces, the actual premise considered by [LNS99], is definitely a question of interest. Understanding the sharp nature of the penalty required for adaptive estimation of $\|f\|_r$ when r is even is another question that remains unanswered. We plan to study these issues in detail in future work.

5 Proof of Main Results

5.1 Proof of Theorem 3.1

The proof of the Theorem hinges on the following lemma, the derivation of which can be found in Section 6. To state the lemma, consider

$$\xi(X, Y) \triangleq \tilde{P}(X) \mathbb{1}(|Y| < c_1 \lambda_h \sqrt{\ln n}) + |X| \mathbb{1}(|Y| \geq c_1 \lambda_h \sqrt{\ln n})$$

used for estimating μ with independent $X, Y \sim \mathcal{N}(\mu, \lambda_h^2)$ and λ_h defined by (3.3).

Lemma 5.1. *Under the assumptions of Theorem 3.1, for any fixed μ , $k \geq 2$, and $c_1 > \sqrt{8k}$, we have that there exists constants C_1 (depending on $(c_1, c_2, \epsilon, K_M)$) and C_2 (depending on $(c_1, c_2, \epsilon, K_M, k)$) such that*

$$\begin{aligned} |\mathbb{E}\xi(X, Y) - \mu| &\leq \frac{C_1}{\sqrt{nh \ln n}}, \\ \mathbb{E}|\xi(X, Y) - \mathbb{E}(\xi(X, Y))|^k &\leq C_2 (\lambda_h n^{2\epsilon})^k. \end{aligned}$$

Note that Lemma 5.1 yields bounds on all even central moments of $\xi(X, Y)$, a result that will be helpful in subsequent proof of adaptation. For now, we will only use the result for $k = 2$ which corresponds to bound on the variance of $\xi(X, Y)$.

Coming back to the proof of Theorem 3.1, note that there are three types of errors:

1. Approximation error incurred by replacing $\|f\|_1$ with $\|f_h\|_1$;
2. The bias of $\int_0^1 T_h(x) dx$ in estimating $\|f_h\|_1$;
3. The variance of $\int_0^1 T_h(x) dx$ in estimating $\|f_h\|_1$.

We bound these errors separately. For the approximation error, by an alternative characterization of Besov balls [HKPT12], for $f \in \mathcal{B}_{p,\infty}^s(L)$ we have that for a constant C_0 depending on L and K_M ,

$$\|f - f_h\|_p \leq C_0 h^s, \quad \forall h > 0.$$

As a result, the approximation error is upper bounded by

$$|\|f\|_1 - \|f_h\|_1| \leq \|f_h - f\|_1 \leq \|f_h - f\|_p \leq C_0 h^s.$$

Secondly we upper bound the bias. By Lemma 5.1, there exists a constant C_1 (depending on $(c_1, c_2, \epsilon, K_M)$) such that

$$\begin{aligned} \left| \mathbb{E} \int_0^1 T_h(x) dx - \|f_h\|_1 \right| &\leq \int_0^1 |\mathbb{E} T_h(x) - f_h(x)| dx \\ &\leq \int_0^1 \frac{C_1}{\sqrt{nh \ln n}} dx = \frac{C_1}{\sqrt{nh \ln n}}. \end{aligned}$$

Finally we upper bound the variance of T_h . Note that $T_h(x)$ and $T_h(y)$ are independent as long as $|x - y| > h$. Therefore by Lemma 5.1, there exists a constant C_2 (depending on $(c_1, c_2, \epsilon, K_M)$) such that

$$\text{Var} \left(\int_0^1 T_h(x) dx \right) = \int_0^1 \int_0^1 \text{Cov}(T_h(x), T_h(y)) dx dy$$

$$\begin{aligned}
&= \iint_{|x-y| \leq h} \text{Cov}(T_h(x), T_h(y)) dx dy \\
&\leq \iint_{|x-y| \leq h} \frac{\text{Var}(T_h(x)) + \text{Var}(T_h(y))}{2} dx dy \\
&= 2h \int_0^1 \text{Var}(T_h(x)) dx \\
&\leq C_2 h \int_0^1 \frac{1}{n^{1-\epsilon} h} dx \\
&= \frac{C_2}{n^{1-\epsilon}}.
\end{aligned}$$

Note that $0 \leq \|f\|_1 \leq L$ for any $f \in \mathcal{B}_{p,\infty}^s(L)$, and consequently

$$|T_h - \|f\|_1| \leq \left| \int_0^1 T_h(x) dx - \|f\|_1 \right|.$$

In summary, for any $f \in \mathcal{B}_{p,\infty}^s(L)$, by triangle inequality we have

$$\begin{aligned}
\left(\mathbb{E} (T_h - \|f\|_1)^2 \right)^{\frac{1}{2}} &\leq \left(\mathbb{E} \left(\int_0^1 T_h(x) dx - \|f\|_1 \right)^2 \right)^{\frac{1}{2}} \\
&\leq \sqrt{3} \left(\left| \|f\|_1 - \|f_h\|_1 \right| + \left| \mathbb{E} \int_0^1 T_h(x) dx - \|f_h\|_1 \right| + \sqrt{\text{Var} \left(\int_0^1 T_h(x) dx \right)} \right) \\
&\leq C^* \left(h^s + \frac{1}{\sqrt{nh \ln n}} + \frac{1}{n^{(1-\epsilon)/2}} \right),
\end{aligned}$$

where C^* is a constant depending on $(c_1, c_2, \sigma, \epsilon, L, K_M)$ which in turn satisfies the conditions of Theorem 3.1. This completes the proof of the theorem. \square

5.2 Proof of Theorem 3.3

The proof of the Theorem hinges on the following lemma.

Lemma 5.2. *Under the conditions of Theorem 3.3, the following hold for all $x \in [0, 1]$, $k \geq 2$, $c_1 > \sqrt{8k}$, and constants C_1 (depending on $(c_1, c_2, \epsilon, \sigma, K_M)$) and C_2 (depending on $(c_1, c_2, \epsilon, \sigma, K_M, k)$).*

$$\begin{aligned}
|\mathbb{E} T_h(x) - |f_h(x)||^r| &\leq C_1 (nh \ln n)^{-\frac{r}{2}}, \\
\mathbb{E} |T_h(x) - \mathbb{E} T_h(x)|^k &\leq C_2 \frac{n^{2k\epsilon}}{(nh)^{\frac{k}{2}}} \left(|f_h(x)|^{(r-1)k} + (nh)^{-\frac{(r-1)k}{2}} \right).
\end{aligned}$$

We postpone the proof of the lemma to Section 6 and complete the proof of Theorem 3.3 assuming its validity.

As is in the case of L_1 norm estimation, there are three types of error incurred by our estimator T_h , i.e., the approximation error, the bias and the variance. We analyze these errors separately.

For the approximation error, by the property of Besov spaces [HKPT12] we know that there exists a constant C_0 depending on L and K_M such that for all $f \in \mathcal{B}_{p,\infty}^s(L)$ the kernel approximation error satisfies $\|f - f_h\|_p \leq C_0 h^s$. Hence, by the monotonicity of L_p norms on $[0, 1]$ and $r \leq p$, we know that

$$\left| \|f\|_r - \|f_h\|_r \right| \leq \|f - f_h\|_r \leq \|f - f_h\|_p \leq C_0 h^s. \quad (5.1)$$

where $C_0 > 0$ is some universal constant which only depends on radius L and the kernel K_M .

For the bias and the variance, we look at the bias and variance of $\Phi_h = \int_0^1 T_h(x)dx$, which is the estimator for $\|f_h\|_r^r$. Similar to the proof of Theorem 3.1, one can show using Lemma 5.2 that for C_1 (depending on $(c_1, c_2, \epsilon, \sigma, K_M)$) and C_2 (depending on $(c_1, c_2, \epsilon, \sigma, K_M)$)

$$\begin{aligned} |\mathbb{E}\Phi_h - \|f_h\|_r^r| &\leq C_1(nh \ln n)^{-\frac{r}{2}}, \\ \text{Var}(\Phi_h) &\leq h \cdot \int_0^1 \frac{C_2}{n^{1-\epsilon}h} \left(|f_h(x)|^{2r-2} + \frac{1}{(nh)^{r-1}} \right) dx \\ &= \frac{C_2}{n^{1-\epsilon}} \left(\|f_h\|_{2r-2}^{2r-2} + \frac{1}{(nh)^{r-1}} \right). \end{aligned} \quad (5.2)$$

For the estimation performance of our final estimator T_h , first note that $\|f\|_p \leq L$ implies

$$|T_h - \|f\|_r| \leq |\max\{0, \Phi_h\}^{\frac{1}{r}} - \|f\|_r|.$$

Set $h = (n \ln n)^{-\frac{1}{2s+1}}$, and $\epsilon > 0$ sufficiently small satisfying the conditions of Theorem 3.3. We then divide our analysis into two cases.

First suppose that $\|f\|_r \leq 2C_0h^s$, with C_0 defined in (5.1). Then $\|f_h\|_r \leq \|f\|_r + \|f - f_h\|_r \leq 3C_0h^s$, and by the bias bound of Φ_h in (5.2), $|\mathbb{E}\Phi_h| \leq C_3h^{sr}$ for a constant C_3 depending on (C_0, C_1, r, s) . Hence, by Lemma 6.6,

$$\begin{aligned} &(\mathbb{E}(T_h - \|f\|_r)^2)^{\frac{1}{2}} \\ &\leq \left(\mathbb{E}(\max\{0, \Phi_h\}^{\frac{1}{r}} - \|f\|_r)^2 \right)^{\frac{1}{2}} \\ &\leq 2 \left(\sqrt{\mathbb{E}[\max\{0, \Phi_h\}^{\frac{2}{r}}]} + \|f\|_r \right) \\ &\leq 2(\mathbb{E}\Phi_h^2)^{\frac{1}{2r}} + 2C_0h^s \\ &\leq 2|\mathbb{E}\Phi_h|^{\frac{1}{r}} + 2(\text{Var}(\Phi_h))^{\frac{1}{2r}} + 2C_0h^s \\ &\leq 2(C_3^{\frac{1}{r}} + C_0)h^s + 2 \left[\frac{C_2}{n^{1-\epsilon}} \left(\|f_h\|_{2r-2}^{2r-2} + \frac{1}{(nh)^{r-1}} \right) \right]^{\frac{1}{2r}} \quad (\text{by (5.2)}) \\ &\leq 2(C_3^{\frac{1}{r}} + C_0)h^s + 2 \left[\frac{C_2}{n^{1-\epsilon}} \left(C(r, K_M)h^{-1+1/r} \|f\|_r^{r-1} \|f_h\|_r^{r-1} + \frac{1}{(nh)^{r-1}} \right) \right]^{\frac{1}{2r}} \quad (\text{by Lemma 6.6}) \\ &\leq C_4 \left[h^s + \left(\frac{h^{-1+1/r} \cdot h^{2s(r-1)}}{n^{1-\epsilon}} \right)^{\frac{1}{2r}} + \left(\frac{1}{n^{1-\epsilon}(nh)^{r-1}} \right)^{\frac{1}{2r}} \right] \leq 3C_4h^s, \end{aligned} \quad (5.3)$$

where C_4 is a constant depending on $C_0, C_1, C_2, C_3, r, K_M, s$.

Second suppose that $\|f\|_r > 2C_0h^s$, then $\|f_h\|_r \geq \|f\|_r - \|f - f_h\|_r > C_0h^s$. Using $|a^r - b^r| \geq b^{r-1}|a - b|$ for any $a, b \geq 0$ and $r \geq 1$, we have

$$\begin{aligned} |T_h - \|f\|_r| &\leq |T_h - \|f_h\|_r| + C_0h^s \\ &\leq \frac{|T_h^r - \|f_h\|_r^r|}{\|f_h\|_r^{r-1}} + C_0h^s \\ &\leq \frac{|\Phi_h - \|f_h\|_r^r|}{(C_0h^s)^{r-1}} + C_0h^s = C_0^{1-r}h^{(1-r)s}|\Phi_h - \|f_h\|_r^r| + C_0h^s. \end{aligned}$$

As a result, by triangle inequality we have

$$\begin{aligned}
(\mathbb{E}(T_h - \|f\|_r)^2)^{\frac{1}{2}} &\leq C_0^{1-r} h^{(1-r)s} (\mathbb{E}(\Phi_h - \|f\|_r^r)^2)^{\frac{1}{2}} + C_0 h^s \\
&\leq C_0^{1-r} h^{(1-r)s} |\mathbb{E}\Phi_h - \|f_h\|_r^r| + C_0^{1-r} h^{(1-r)s} \sqrt{\text{Var}(\Phi_h)} + C_0 h^s \\
&\leq C_4 \left[\begin{aligned} &h^{(1-r)s} \cdot h^{sr} + h^{(1-r)s} \sqrt{n^{\epsilon-1} h^{-1+1/r} h^{2s(r-1)}} \\ &+ h^{(1-r)s} \sqrt{n^{\epsilon-1} (nh)^{-(r-1)}} + h^s \end{aligned} \right] \leq C_5 h^s,
\end{aligned} \tag{5.4}$$

where the second inequality in the above display follows similar to before by equation (5.2) and Lemma 6.6 with C_4 being a constant depending on $C_0, C_1, C_2, C_3, r, K_M, s$. Combining inequalities (5.3) and (5.4) completes the proof of Theorem 3.3. \square

5.3 Proof of Theorem 3.2 and Theorem 3.4

The outline for the proof of lower bounds (i.e., Theorems 3.2 and 3.4) is as follows. In Section 5.3.1, we reduce the nonparametric problem to a parametric subproblem in the Gaussian location model. The minimax lower bound for the parametric submodel is proved by a generalized version of Le Cam's method involving a pair of priors, also known as the method of two fuzzy hypotheses [Tsy08]. In Section 5.3.2 we construct the priors using duality to best approximation and Section 5.3.3 finishes the proof.

In the sequel we assume that $r \geq 1$ is a fixed non-even real number.

5.3.1 Reduction to Parametric Submodel

Fix a smooth function $g(x)$ vanishing outside $[0, 1]$ with $\|g\|_{\mathcal{B}_{p,\infty}^s} = 1$, and $\|g\|_1 > 0$. Set

$$\begin{aligned}
h &= (n \ln n)^{-\frac{1}{2s+1}}, \\
N &= h^{-1},
\end{aligned}$$

and let I_1, \dots, I_N be the partition of the interval $[0, 1]$ into N subintervals of length h each (without loss of generality we assume that N is an integer), and let t_i be the left endpoint of subinterval I_i . With a point $\theta = (\theta_1, \dots, \theta_N) \in [-\sqrt{\ln N}, \sqrt{\ln N}]^N$ we associate the function

$$f_\theta(t) = L' \sum_{i=1}^N \theta_i \sqrt{\ln N} \cdot h^s g\left(\frac{t - t_i}{h}\right).$$

Lemma 5.3. *If*

$$\theta \in \Theta \triangleq [-\sqrt{\ln N}, \sqrt{\ln N}]^N \cap \left\{ \theta : \frac{1}{N} \sum_{i=1}^N |\theta_i|^p \leq \left(\frac{2}{\sqrt{\ln N}} \right)^p \right\}, \tag{5.5}$$

then for some constant $L' > 0$ independent of n , we have $f_\theta(t) \in \mathcal{B}_{p,\infty}^s(L)$.

Proof. Let $s_0 = \lfloor s \rfloor + 1$. Observe that

$$\|f_\theta\|_p = L' \|g\|_p h^s \cdot \sqrt{\ln N} \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^p \right)^{\frac{1}{p}},$$

the condition (5.5) ensures that $h^{-s}\|f_\theta\|_p$ is upper bounded by a numerical constant proportional to L' . By (6.10) in Lemma 6.10, this implies that there exists a constant C_0 independent of n such that $\omega_{s_0}(f, t^r)_p \leq C_0 L' t^{s_0}$ for any $t \geq h$. Moreover,

$$\|f_\theta^{(s_0)}\|_p = L' \|g^{(s_0)}\|_p h^{s-s_0} \cdot \sqrt{\ln N} \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^p \right)^{\frac{1}{p}},$$

the condition (5.5) ensures that $h^{s_0-s}\|f_\theta^{(s_0)}\|_p$ is upper bounded by a numerical constant proportional to L' . By (6.9) in Lemma 6.10, this implies that there exists a constant C_0 independent of n such that $\omega_{s_0}(f, t^r) \leq C_0 L' t^{s_0}$ for any $t \leq h$. Now by the definition of the Besov norm in (2.2), a suitable choice of the scale parameter L' ensures that $f_\theta(t) \in \mathcal{B}_{p,\infty}^s(L)$. \square

Fix any choice of L' given by Lemma 5.3. Note that for any $r \geq 1$, we have

$$\|f_\theta\|_r = L' \|g\|_r (n \ln n)^{-\frac{s}{2s+1}} \cdot \sqrt{\ln N} \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{\frac{1}{r}}. \quad (5.6)$$

Hence, a sufficient condition for Theorem 3.2 and Theorem 3.4 is that

$$\inf_{\hat{T}} \sup_{\theta \in \Theta} \mathbb{E}_\theta \left(\hat{T} - \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{\frac{1}{r}} \right)^2 \gtrsim \left(\frac{1}{\sqrt{\ln N}} \right)^2 = \frac{1}{\ln N}, \quad (5.7)$$

where the infimum is taken over all estimators \hat{T} which is a measurable real-valued function of $\{Y(t)\}_{t \in [0,1]}$.

Finally, we note that by the factorization theorem, to estimate $\left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{1/r}$ for $\theta \in \Theta$, the vector $y = (y_1, \dots, y_N)$ with

$$y_i \triangleq \frac{\sqrt{n}}{\sigma \|g\|_2 \sqrt{h}} \int_{I_i} g \left(\frac{t - t_i}{h} \right) dY(t), \quad i = 1, \dots, N,$$

constitute a sufficient statistic for the Gaussian white noise model. Note that

$$y_i = \alpha \theta_i + \xi_i, \quad i = 1, \dots, N, \quad (5.8)$$

with $\theta \in \Theta$, and

$$\begin{aligned} \alpha &\triangleq \sigma^{-1} L' \|g\|_2 n^{1/2} h^{s+1/2} \cdot \sqrt{\ln N} \asymp 1, \\ \xi_i &\triangleq \frac{1}{\|g\|_2 \sqrt{h}} \int_{I_i} g \left(\frac{t - t_i}{h} \right) dB_t. \end{aligned}$$

As a result, ξ_1, \dots, ξ_N are i.i.d. $\mathcal{N}(0, 1)$ random variables. Hence, we may further assume that our observation model is

$$y_i \stackrel{\text{ind}}{\sim} \mathcal{N}(\alpha \theta_i, 1), \quad i = 1, \dots, N, \quad (5.9)$$

which is a Gaussian location model with $\theta \in \Theta$, and the estimator \hat{T} in (5.7) is a function of (y_1, \dots, y_N) . Note that when $r = 1$, this parametric subproblem is similar to but very *different* from the problem considered in [CL11], where the authors did not have the second constraint in (5.5).

5.3.2 Construction of Two Priors

The minimax lower bound (5.7) follows from the generalized Le Cam's method involving two priors, which is known as the method of two fuzzy hypotheses presented in [Tsy08]. Suppose we observe a random vector $\mathbf{Z} \in (\mathcal{Z}, \mathcal{A})$ which has distribution P_θ where $\theta \in \Theta$. Let σ_0 and σ_1 be two prior distributions supported on Θ . Write F_i for the marginal distribution of \mathbf{Z} when the prior is σ_i for $i = 0, 1$. Let $\hat{T} = \hat{T}(\mathbf{Z})$ be an arbitrary estimator of a function $T(\theta)$ based on \mathbf{Z} , and $V(P, Q)$ be the total variation distance between two probability measures P, Q on the measurable space $(\mathcal{Z}, \mathcal{A})$. Concretely,

$$V(P, Q) \triangleq \sup_{A \in \mathcal{A}} |P(A) - Q(A)| = \frac{1}{2} \int |p - q| d\nu,$$

where $p = \frac{dP}{d\nu}$, $q = \frac{dQ}{d\nu}$, and ν is a dominating measure so that $P \ll \nu, Q \ll \nu$. We have the following general minimax lower bound.

Lemma 5.4. [Tsy08, Theorem 2.15] *Suppose that there exist $\zeta \in \mathbb{R}, \delta > 0, 0 \leq \beta_0, \beta_1 < 1$ such that*

$$\begin{aligned} \sigma_0(\theta : T(\theta) \leq \zeta - \delta) &\geq 1 - \beta_0, \\ \sigma_1(\theta : T(\theta) \geq \zeta + \delta) &\geq 1 - \beta_1. \end{aligned}$$

If $V(F_1, F_0) \leq \eta < 1$, then

$$\inf_{\hat{T}} \sup_{\theta \in \Theta} \mathbb{P}_\theta \left(|\hat{T} - T(\theta)| \geq \delta \right) \geq \frac{1 - \eta - \beta_0 - \beta_1}{2},$$

where $F_i, i = 0, 1$ are the marginal distributions of \mathbf{Z} when the priors are $\sigma_i, i = 0, 1$, respectively.

In the remainder of this section, we aim to construct two priors μ_0, μ_1 supported on $[-\sqrt{\ln N}, \sqrt{\ln N}]$ such that the following conditions hold (the numerical constant $d > 0$ is chosen later):

$$\int t^l \mu_1(dt) = \int t^l \mu_0(dt), \quad \text{for all } l = 0, 1, \dots, K \triangleq \lceil d \ln N \rceil, \quad (5.10)$$

$$\int |t|^r \mu_1(dt) - \int |t|^r \mu_0(dt) \gtrsim (\ln N)^{-\frac{r}{2}}, \quad (5.11)$$

$$\int |t|^p \mu_i(dt) \leq (\ln N)^{-\frac{p}{2}}, \quad \text{for } i = 0, 1. \quad (5.12)$$

In the next section we will choose the priors $\sigma_i, i = 0, 1$ in Lemma 5.4 to be close to the product measure $\mu_i^{\otimes N}, i = 0, 1$ with each copy given above. The condition (5.10) ensures a small total variation distance $V(F_1, F_0)$ in Lemma 5.4, the condition (5.11) ensures a large $\delta \asymp (\ln N)^{-\frac{r}{2}}$ in Lemma 5.4, and the condition (5.12) ensures that the support of $\mu_i^{\otimes N}$ is almost supported on Θ given in (5.5).

The following result is simply the duality between the problem of best uniform approximation and moment matching.

Lemma 5.5. *For any bounded interval $I \subseteq \mathbb{R}$ not containing zero, integers $q \geq 0, K > 0$ and continuous function f on I , let*

$$E_{q-1, K}(f; I) \triangleq \inf_{\{a_i\}} \sup_{x \in I} \left| \sum_{i=-q+1}^K a_i x^i - f(x) \right|$$

denote the best uniform approximation error of f by rational functions spanned by $\{x^{-q+1}, x^{-q+2}, \dots, x^K\}$. Then

$$\begin{aligned} 2E_{q-1,K}(f; I) &= \max \int f(t)\nu_1(dt) - \int f(t)\nu_0(dt), \\ \text{s.t. } \int t^l \nu_1(dt) &= \int t^l \nu_0(dt), \quad l = -q+1, \dots, K, \end{aligned} \quad (5.13)$$

where the maximum is taken over pairs of probability measures ν_0 and ν_1 supported on I .

Lemma 5.5 extends the duality results in [CL11, Lemma 1], [JVHW15, Lemma 10, Lemma 12] to rational functions. We list two possible proofs of Lemma 5.5. The first proof relies on the Hahn–Banach theorem and the Riesz representation theorem, where the essential argument is given in [LNS99]. The second proof makes use of the fact that the rational functions $\{x^{-q+1}, x^{-q+2}, \dots, x^K\}$ form a Chebyshev system in $C(I)$ and therefore the Chebyshev alternation theorem holds [DL93, Chapter 3, Theorem 5.1], so that the probability measures ν_0 and ν_1 can be explicitly constructed following the similar lines to [WY16, Appendix E]. For completeness we include the second proof in Section 6.4.

Here we apply this lemma to $f_q(t) = t^{-q+\frac{r}{2}}$ and

$$K = \lceil d \ln N \rceil, \quad I = \left[\frac{c}{(\ln N)^2}, 1 \right], \quad q = \left\lfloor \frac{p}{2} \right\rfloor,$$

where the constant $c \in (0, 1)$ appearing in the definition of I is given by the following lemma, which provides a lower bound for the approximation error of $f_q(t)$.

Lemma 5.6. *Fix a non-even real $r \geq 1$ and an integer $q \geq r/2$. For $f_q(t) = t^{-q+\frac{r}{2}}$, we have*

$$\liminf_{n \rightarrow \infty} n^{-(2q-r)} E_{q-1,n} \left(f; \left[\frac{c}{n^2}, 1 \right] \right) \geq c',$$

where constants $c \in (0, 1), c' > 0$ only depend on (q, r) .

By Lemma 5.6 and our definitions of f , I and K , we conclude that

$$E_{q-1,K}(f; I) \gtrsim (\ln N)^{2q-r}.$$

Let ν_0, ν_1 be the maximizers of (5.13). We define probability measures $\tilde{\nu}_0, \tilde{\nu}_1$ by

$$\tilde{\nu}_i(dx) = \left[1 - \mathbb{E}_{X \sim \nu_i} \left(\frac{c^q}{(\ln N)^{2q} X^q} \right) \right] \delta_0(dx) + \left(\frac{c}{(\ln N)^2 x} \right)^q \nu_i(dx) \quad i = 0, 1,$$

where $\delta_0(\cdot)$ is the delta measure at zero. It is straightforward to verify that $\tilde{\nu}_i$ forms a probability measure supported on $[0, 1]$, and

1. $\int t^l \tilde{\nu}_1(dt) = \int t^l \tilde{\nu}_0(dt)$, for all $l = 0, 1, \dots, q+K$;
2. $\int t^{\frac{r}{2}} \tilde{\nu}_1(dt) - \int t^{\frac{r}{2}} \tilde{\nu}_0(dt) \gtrsim (\ln N)^{-r}$;
3. $\int t^q \tilde{\nu}_i(dt) = c^q (\ln N)^{-2q} \leq (\ln N)^{-2q}$, for $i = 0, 1$.

Finally, we define the measures μ_0, μ_1 as follows. For $i = 0, 1$, let X_i follow the distribution $\tilde{\nu}_i$, the measure μ_i is defined as the probability distribution of $\epsilon_i \sqrt{X_i \cdot \ln N}$, where $\epsilon_i \sim \text{Unif}(\{\pm 1\})$ is independent of X_i . Clearly the measures μ_0, μ_1 are supported on $[-\sqrt{\ln N}, \sqrt{\ln N}]$, and it remains to check the conditions (5.10) to (5.12).

For (5.10), since μ_i is symmetric around zero, the condition clearly holds for odd l . For even $l = 2k$ with $0 \leq k \leq K$, by the first property of $\tilde{\nu}_i$ we have

$$\begin{aligned} \int t^l \mu_1(dt) &= \int (\epsilon_i \sqrt{t \cdot \ln N})^{2k} \mu_1(dt) = (\ln N)^k \int t^k \mu_1(dt) \\ &= (\ln N)^k \int t^k \mu_0(dt) = \int (\epsilon_i \sqrt{t \cdot \ln N})^{2k} \mu_0(dt) = \int t^l \mu_0(dt), \end{aligned}$$

i.e., (5.10) holds. Similarly, the condition (5.11) is checked via

$$\int |t|^r \mu_1(dt) - \int |t|^r \mu_0(dt) = (\ln N)^{\frac{r}{2}} \int t^{\frac{r}{2}} (\tilde{\nu}_1(dt) - \tilde{\nu}_2(dt)) \gtrsim (\ln N)^{-\frac{r}{2}}.$$

Finally, for (5.12), first note that

$$\int t^{2q} \mu_i(dt) = (\ln N)^q \int t^q \tilde{\nu}_i(dt) \leq (\ln N)^{-q}, \quad i = 0, 1.$$

Since $2q \geq p$, Hölder's inequality yields

$$\left(\int |t|^p \mu_i(dt) \right)^{\frac{1}{p}} \leq \left(\int |t|^{2q} \mu_i(dt) \right)^{\frac{1}{2q}} \leq \frac{1}{\sqrt{\ln N}},$$

i.e., (5.12) holds. Hence, the construction of μ_0, μ_1 satisfies all conditions in (5.10) to (5.12). This construction is partially inspired by [WY16]. We remark that the construction heavily relies on the fact that p is finite, where for $p = \infty$, Lemma 5.6 fails and the condition (5.12) would require that the priors μ_0, μ_1 be supported on a smaller interval $[-\frac{1}{\sqrt{\ln N}}, \frac{1}{\sqrt{\ln N}}]$.

5.3.3 Minimax Lower Bound in the Parametric Submodel

In this section we invoke Lemma 5.4 to finish the proof of (5.7), thereby proving the lower bound in Theorems 3.2 and 3.4. Consider the probability measures μ_0, μ_1 constructed in the Section 5.3.2, and define

$$\Delta = \int |t|^r \mu_1(dt) - \int |t|^r \mu_0(dt) \asymp (\ln N)^{-\frac{r}{2}}.$$

Denote by $\mu_i^{\otimes N}$ the N -fold product of μ_i . Consider the following event:

$$E_i \triangleq \{\theta : \theta \in \Theta\} \cap \left\{ \theta : \left| \frac{1}{N} \sum_{j=1}^N |\theta_j|^r - \mathbb{E}_{\mu_i} |\theta|^r \right| \leq \frac{\Delta}{4} \right\}, \quad i = 0, 1.$$

By Chebyshev's inequality it is easy to show that for $i = 0, 1$,

$$\begin{aligned} \mu_i^{\otimes N}(\{\theta : \theta \notin \Theta\}) &= \mu_i^{\otimes N} \left(\left\{ \theta : \frac{1}{N} \sum_{j=1}^N |\theta_j|^p > \left(\frac{2}{\sqrt{\ln N}} \right)^p \right\} \right) \\ &\leq \mu_i^{\otimes N} \left(\left\{ \theta : \frac{1}{N} \sum_{j=1}^N |\theta_j|^p - \mathbb{E}_{\mu_i} |\theta|^p > \left(\frac{1}{\sqrt{\ln N}} \right)^p \right\} \right) \end{aligned}$$

$$\begin{aligned}
&\leq \left(\frac{1}{\sqrt{\ln N}}\right)^{-2p} \cdot \text{Var}_{\mu_i^{\otimes N}} \left(\frac{1}{N} \sum_{j=1}^N |\theta_j|^p \right) \\
&= \frac{1}{N} \left(\frac{1}{\sqrt{\ln N}}\right)^{-2p} \cdot \text{Var}_{\mu_i}(|\theta|^p) \\
&\leq \frac{1}{N} \left(\frac{1}{\sqrt{\ln N}}\right)^{-2p} \cdot (\sqrt{\ln N})^{2p} \rightarrow 0,
\end{aligned}$$

and

$$\begin{aligned}
\mu_i^{\otimes N} \left(\left\{ \theta : \left| \frac{1}{N} \sum_{j=1}^N |\theta_j|^r - \mathbb{E}_{\mu_i} |\theta|^r \right| > \frac{\Delta}{4} \right\} \right) &\leq \frac{16}{\Delta^2} \text{Var}_{\mu_i^{\otimes N}} \left(\frac{1}{N} \sum_{j=1}^N |\theta_j|^r \right) \\
&\leq \frac{16}{N\Delta^2} \cdot (\sqrt{\ln N})^{2r} \rightarrow 0.
\end{aligned}$$

Hence, by the union bound, we have

$$\mu_i^{\otimes N}(E_i^c) \leq \mu_i^{\otimes N}(\{\theta : \theta \notin \Theta\}) + \mu_i^{\otimes N} \left(\left\{ \theta : \left| \frac{1}{N} \sum_{j=1}^N |\theta_j|^r - \mathbb{E}_{\mu_i} |\theta|^r \right| > \frac{\Delta}{4} \right\} \right) \rightarrow 0$$

for any $i = 0, 1$.

Now we are ready to apply Lemma 5.4 to

$$\begin{aligned}
T(\theta) &= \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{\frac{1}{r}}, \\
\zeta &= \left(\frac{\mathbb{E}_{\mu_1^{\otimes N}}[T(\theta)^r] + \mathbb{E}_{\mu_0^{\otimes N}}[T(\theta)^r]}{2} \right)^{\frac{1}{r}},
\end{aligned}$$

and let the prior σ_i be the conditional distribution of $\mu_i^{\otimes N}$ conditioning on E_i , i.e.,

$$\sigma_i(\cdot) = \frac{\mu_i^S(\cdot \cap E_i)}{\mu_i^S(E_i)}, \quad i = 0, 1.$$

By definition of E_i , the measure σ_i is a valid prior on Θ . Moreover, under σ_1 we have

$$T(\theta)^r - \zeta^r \geq \frac{\Delta}{4}.$$

By definition of E_1 and Θ , under σ_1 we have

$$\begin{aligned}
T(\theta) &= \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{\frac{1}{r}} \leq \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^p \right)^{\frac{1}{p}} \leq \frac{2}{\sqrt{\ln N}}, \\
\zeta &\leq \frac{1}{2} \left(\int |t|^r \mu_1(dt) \right)^{\frac{1}{r}} + \frac{1}{2} \left(\int |t|^r \mu_0(dt) \right)^{\frac{1}{r}} \\
&\leq \frac{1}{2} \left(\int |t|^p \mu_1(dt) \right)^{\frac{1}{p}} + \frac{1}{2} \left(\int |t|^p \mu_0(dt) \right)^{\frac{1}{p}} \leq \frac{1}{\sqrt{\ln N}}.
\end{aligned}$$

Hence, using the inequality $a^r - b^r \leq r(a^{r-1} + b^{r-1})(a - b)$ for any $a \geq b > 0$ and $r \geq 1$, the previous inequalities yield that under σ_1 ,

$$T(\theta) - \zeta \geq \frac{[T(\theta)]^r - \zeta^r}{r([T(\theta)]^{r-1} + \zeta^{r-1})} \gtrsim \frac{\Delta}{(\ln N)^{-\frac{r-1}{2}}} \gtrsim \frac{1}{\sqrt{\ln N}}.$$

Similarly, under σ_0 we have $T(\theta) - \zeta \lesssim -\frac{1}{\sqrt{\ln N}}$, hence in Lemma 5.4 we can set

$$\delta \gtrsim \frac{1}{\sqrt{\ln N}},$$

so that $\beta_0 = \beta_1 = 0$.

Now denote by F_0, F_1 the marginal distributions of \mathbf{Z} based on priors σ_0, σ_1 , and the counterparts G_0, G_1 based on priors $\mu_0^{\otimes N}, \mu_1^{\otimes N}$. By the data-processing property of the total variation distance, we have

$$V(F_i, G_i) \leq V(\sigma_i, \mu_i^{\otimes N}) = \mu_i^{\otimes N}(E_i^c) \rightarrow 0, \quad i = 0, 1.$$

Moreover, [CL11] shows that the χ^2 distance between G_0 and G_1 is upper bounded as

$$\chi^2(G_0, G_1) \leq \left(1 + e^{3\alpha^2 \ln N/2} \left(\frac{\alpha e \ln N}{d \ln N}\right)^{d \ln N}\right)^N - 1.$$

Hence, for choosing $d > 0$ large enough, $\chi^2(G_0, G_1)$ is upper bounded by a universal constant C . Now by Lemma 6.8, we have

$$V(G_0, G_1) \leq 1 - \frac{1}{2} \exp(-C).$$

In summary, the triangle inequality for total variation distance yields

$$V(F_0, F_1) \leq V(F_0, G_0) + V(G_0, G_1) + V(G_1, F_1) \rightarrow 1 - \frac{1}{2} \exp(-C) < 1,$$

and Lemma 5.4 together with Markov's inequality yields

$$\begin{aligned} \inf_{\hat{T}} \sup_{\theta \in \Theta} \mathbb{E}_{\theta} \left(\hat{T} - \left(\frac{1}{N} \sum_{i=1}^N |\theta_i|^r \right)^{\frac{1}{r}} \right)^2 &\gtrsim \delta^2 \cdot \inf_{\hat{T}} \sup_{\theta \in \Theta} \mathbb{P}(|\hat{T} - T(\theta)| \geq \delta) \\ &\gtrsim \frac{1}{\ln N} \cdot \frac{1}{4} \exp(-C) \gtrsim \frac{1}{\ln N}, \end{aligned}$$

which is (5.7), as desired. □

5.4 Proof of Theorem 3.5

For $s \in [0, s_{\max}]$ define the ideal bandwidth $h^* : [0, s_{\max}] \rightarrow \mathcal{H}$ by

$$h^*(s) \triangleq \frac{1}{\lfloor (n \ln n)^{\frac{1}{2s+1}} \rfloor}.$$

Then

$$\mathbb{E} \left(\hat{T} - \|f\|_r \right)^2$$

$$\begin{aligned}
&= \mathbb{E} \left[\left(\hat{T} - \|f\|_r \right)^2 \mathbb{1} \left(\hat{h} \geq h^*(s) \right) \right] + \mathbb{E} \left[\left(\hat{T} - \|f\|_r \right)^2 \mathbb{1} \left(\hat{h} < h^*(s) \right) \right] \\
&= \text{I} + \text{II}.
\end{aligned}$$

First note that

$$\begin{aligned}
\text{I} &\leq 2 \left\{ \mathbb{E} \left[\left(T_{h^*(s)} - \|f\|_r \right)^2 \right] + \mathbb{E} \left[\left(T_{\hat{h}} - T_{h^*(s)} \right)^2 \mathbb{1} \left(\hat{h} \geq h^*(s) \right) \right] \right\} \\
&\leq 2 \left\{ C_0 (n \ln n)^{-\frac{2s}{2s+1}} + C^* \frac{\lambda_{h^*(s)}^2}{\ln n} \right\} \leq C_0^* (n \ln n)^{-\frac{2s}{2s+1}},
\end{aligned} \tag{5.14}$$

where the last line follows from the definitions of $h^*(s)$ and \hat{h} respectively, and C_0 is a constant that depends on $(c_1, c_2, \epsilon, s_{\max}, \sigma, L)$.

To upper bound II, we have

$$\begin{aligned}
\text{II} &= \mathbb{E} \left[\left(\hat{T} - \|f\|_r \right)^2 \mathbb{1} \left(\hat{h} < h^*(s) \right) \right] \leq L^2 \cdot \mathbb{P}(\hat{h} < h^*(s)) \\
&\leq L^2 \cdot \sum_{h < h^*(s): h \in \mathcal{H}} \mathbb{P} \left(|T_h - T_{h^*(s)}| > \lambda_h \sqrt{\frac{C^*}{\ln n}} \right),
\end{aligned} \tag{5.15}$$

where the first inequality is due to that $\hat{T}, \|f\|_r \in [0, L]$, and the second inequality follows from the fact that $h^*(s)$ is not a feasible candidate in (3.6). Let

$$\gamma_n(h) \triangleq \lambda_h \sqrt{\frac{C^*}{\ln n}},$$

below we reduce the control of $\mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h))$ to suitable controls over

$$\mathbb{E} \left| \int_0^1 (T_h(x) - \mathbb{E}T_h(x)) dx \right|^k$$

for some k to be chosen a large enough constant depending on the tuple (s_{\max}, r, σ, p) . Consequently, in the following lemma we demonstrate the desired control over central moments of $\int_0^1 T_h(x) dx$ for every $h \in \mathcal{H}$.

Lemma 5.7. *Let $f \in \mathcal{B}_{p,\infty}^s(L)$ and $r \geq 1$ be non-even. Then, under the assumptions of Theorem 3.5, for any $h \in \mathcal{H}$ and integer $k \geq 2$,*

$$\begin{aligned}
&\mathbb{E} \left| \int_0^1 (T_h(x) - \mathbb{E}T_h(x)) dx \right|^k \\
&\leq C(r, k, c_1, \sigma) n^{2k\epsilon} \left(\begin{aligned} &\lambda_h^{kr} (h^{k-1} + h^{\frac{k}{2}}) \\ &+ \lambda_h^k (h^{k-1} \|f_h\|_{k(r-1)}^{k(r-1)} + h^{\frac{k}{2}} \|f_h\|_{2(r-1)}^{k(r-1)}) \end{aligned} \right),
\end{aligned}$$

where $C(r, k, c_1, \sigma)$ is a constant depending on (r, k, c_1, σ) .

By Lemma 6.6 and $h \in [0, 1]$, the following corollary is immediate.

Corollary 5.1. *Under the assumptions of Theorem 3.5, for any $h \in \mathcal{H}$ and integer $k \geq 2$,*

$$\mathbb{E} \left| \int_0^1 (T_h(x) - \mathbb{E}T_h(x)) dx \right|^k$$

$$\leq C'(r, k, c_1, \sigma) n^{2k\epsilon} \left(\lambda_h^{kr} h^{\frac{k}{2}} + \lambda_h^k (h^{\frac{k-1}{r}} \|f\|_r^{(k-1)(r-1)} \|f_h\|_r^{r-1} + h^{\frac{k}{2r}} \|f\|_r^{k(r-1)/2} \|f_h\|_r^{k(r-1)/2} \right),$$

where $C'(r, k, c_1, \sigma)$ is a constant depending on (r, k, c_1, σ) .

We defer the proof of Lemma 5.7 to Section 6 and complete the proof of Theorem 3.5 assuming its validity. Recall that $\|f - f_h\|_r \leq C_1 h^s$ for some $C_1 = C_1(s, r, L, K_M)$ for any $f \in \mathcal{B}_{p,\infty}^s(L)$. We divide the subsequent analysis into two cases. First consider the case when $\|f_h\|_r \leq \frac{C_1}{\sqrt{nh \ln n}}$, then

$$\begin{aligned} \mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h)) &\leq \mathbb{P}\left(T_h > \frac{\gamma_n(h)}{2}\right) + \mathbb{P}\left(T_{h^*(s)} > \frac{\gamma_n(h)}{2}\right) \\ &\leq \mathbb{P}\left(T_h > \frac{\gamma_n(h)}{2}\right) + \mathbb{P}\left(T_{h^*(s)} > \frac{\gamma_n(h^*(s))}{2}\right). \end{aligned} \quad (5.16)$$

where the last inequality uses the monotone decreasing nature of $h \mapsto \gamma_n(h)$. By (5.2) and the triangle inequality, we have

$$\int_0^1 \mathbb{E} T_h(x) dx \leq \|f_h\|_r^r + \left| \int_0^1 \mathbb{E} T_h(x) dx - \|f_h\|_r^r \right| \leq C_2 (nh \ln n)^{-\frac{r}{2}},$$

where $C_2 > 0$ is a constant depending on (c_1, c_2, r, C_1) . Choosing C^* large enough such that

$$\left(\frac{\gamma_n(h)}{4} \right)^r \geq C_2 (nh \ln n)^{-\frac{r}{2}},$$

for any $k \geq 2$ we have

$$\begin{aligned} \mathbb{P}\left(T_h > \frac{\gamma_n(h)}{2}\right) &\leq \mathbb{P}\left(\int_0^1 T_h(x) dx > \left(\frac{\gamma_n(h)}{2}\right)^r\right) \\ &\leq \mathbb{P}\left(\int_0^1 T_h(x) dx - \mathbb{E}\left(\int_0^1 T_h(x) dx\right) > \left(\frac{\gamma_n(h)}{2}\right)^r - C_2 (nh \ln n)^{-\frac{r}{2}}\right) \\ &\leq \mathbb{P}\left(\left|\int_0^1 T_h(x) dx - \mathbb{E}\left(\int_0^1 T_h(x) dx\right)\right| > \left(\frac{\gamma_n(h)}{4}\right)^r\right) \\ &\leq \frac{4^{rk} \mathbb{E}\left|\int_0^1 T_h(x) dx - \mathbb{E}\left(\int_0^1 T_h(x) dx\right)\right|^k}{(\gamma_n(h))^{kr}}. \end{aligned}$$

However, note that

$$\|f\|_r \leq \|f - f_h\|_r + \|f_h\|_r \leq C_1 h^s + \frac{C_1}{\sqrt{nh \ln n}} \leq C_3 \gamma_n(h),$$

where C_3 is a constant depending on (C_1, C^*) , Corollary 5.1 yields

$$\begin{aligned} &\mathbb{E}\left|\int_0^1 (T_h(x) - \mathbb{E} T_h(x)) dx\right|^k \\ &\leq C_4 n^{2k\epsilon} \left[\lambda_h^{kr} h^{\frac{k}{2}} + \lambda_h^k h^{\frac{k-1}{r}} \gamma_n(h)^{k(r-1)} + \lambda_h^k h^{\frac{k}{2r}} \gamma_n(h)^{k(r-1)} \right] \\ &\leq C_5 n^{2k\epsilon} (\ln n)^{\frac{k}{2}} \cdot h^{\frac{k}{2r}} \gamma_n(h)^{kr}, \end{aligned}$$

where C_4, C_5 are constants depending on $(r, k, c_1, \sigma, C_1, C_3, C^*)$. Consequently, we have shown that for any $h \leq h^*(s)$,

$$\mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h)) \leq 4^{kr} C_5 n^{2k\epsilon} (\ln n)^{\frac{k}{2}} \cdot h^{\frac{k}{2r}}. \quad (5.17)$$

Next consider the case when $\|f_h\|_r > \frac{C_1}{\sqrt{nh \ln n}}$. Note that

$$||f_{h^*(s)}\|_r - \|f_h\|_r \leq \|f_{h^*(s)} - f\|_r + \|f_h - f\|_r \leq 2C_1(h^*(s))^r,$$

thus if C^* is large enough such that $\gamma_n(h) \geq 6C_1(h^*(s))^r$, triangle inequality yields

$$\begin{aligned} & \mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h)) \\ & \leq \mathbb{P}\left(|T_h - \|f_h\|_r| > \frac{\gamma_n(h)}{3}\right) + \mathbb{P}\left(|T_{h^*(s)} - \|f_{h^*(s)}\|_r| > \frac{\gamma_n(h)}{3}\right). \end{aligned} \quad (5.18)$$

In this case, once again using $|a^r - b^r| \geq b^{r-1}|a - b|$ for any $a, b \geq 0$ and $r \geq 1$,

$$|T_h - \|f_h\|_r| \leq \frac{|T_h^r - \|f_h\|_r^r|}{\|f_h\|_r^{r-1}} \leq \frac{|\int_0^1 T_h(x) dx - \|f_h\|_r^r|}{\|f_h\|_r^{r-1}}, \quad (5.19)$$

where the last inequality follows from $0 \leq \|f_h\|_r \leq L$. Fix any $h \leq h^*(s)$, (5.19) yields

$$\begin{aligned} & \mathbb{P}\left(|T_h - \|f_h\|_r| > \frac{\gamma_n(h)}{3}\right) \\ & \leq \mathbb{P}\left(\left|\int_0^1 T_h(x) dx - \|f_h\|_r^r\right| > \frac{\gamma_n(h)}{3} \|f_h\|_r^{r-1}\right) \\ & \leq \mathbb{P}\left(\left|\int_0^1 T_h(x) dx - \mathbb{E}\left(\int_0^1 T_h(x) dx\right)\right| > \frac{\gamma_n(h)}{3} \|f_h\|_r^{r-1} - C_2(nh \ln n)^{-\frac{r}{2}}\right), \end{aligned}$$

where in the last line we have used (5.2) again. Choosing C^* large enough so that

$$\frac{\gamma_n(h)}{3} \|f_h\|_r^{r-1} - C_2(nh \ln n)^{-\frac{r}{2}} \geq \frac{\gamma_n(h)}{6} \|f_h\|_r^{r-1},$$

which is possible due to the assumption $\|f_h\|_r \geq \frac{C_1}{\sqrt{nh \ln n}}$, and noting that

$$\|f\|_r \leq \|f_h\|_r + \|f - f_h\|_r \leq \frac{C_1}{\sqrt{nh \ln n}} + C_1 h^s \leq 2\|f_h\|_r,$$

Corollary 5.1 yields that for integer $k \geq 2$,

$$\begin{aligned} & \mathbb{P}\left(|T_h - \|f_h\|_r| > \frac{\gamma_n(h)}{3}\right) \\ & \leq \left(\frac{6}{\gamma_n(h)\|f_h\|_r^{r-1}}\right)^k \cdot \mathbb{E}\left|\int_0^1 T_h(x) dx - \mathbb{E}\left(\int_0^1 T_h(x) dx\right)\right|^k \\ & \leq \frac{C_6}{\gamma_n(h)^k \|f_h\|_r^{k(r-1)}} \cdot n^{2k\epsilon} \left(\lambda_h^{kr} h^{\frac{k}{2}} + \lambda_h^k h^{\frac{k}{2r}} \|f_h\|_r^{k(r-1)}\right) \\ & \leq C_7 n^{2k\epsilon} (\ln n)^{\frac{k}{2}} \cdot \left(h^{\frac{k}{2}} \left(\frac{\lambda_h}{\|f_h\|_r}\right)^{k(r-1)} + h^{\frac{k}{2r}}\right) \end{aligned}$$

$$\leq C_8 n^{2k\epsilon} (\ln n)^{\frac{kr}{2}} \cdot h^{\frac{k}{2r}}, \quad (5.20)$$

where C_6, C_7, C_8 are constants depending on $(r, k, c_1, \sigma, K_M, C_1, C^*)$, and the last step follows from our assumption that $\|f_h\|_r \geq \frac{C_1}{\sqrt{nh \ln n}}$.

Combining (5.17), (5.18) and (5.20) we get that for any $h \leq h^*(s)$, $r \leq p$, integer $k \geq 2$,

$$\sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h)) \leq \max\{C_5, C_8\} 4^{kr} n^{2k\epsilon} (\ln n)^{\frac{kr}{2}} \cdot (h^{\frac{k}{2r}} + h^*(s)^{\frac{k}{2r}}), \quad (5.21)$$

given that the constant C^* is large enough. Note that $h \leq h^*(s) \leq (n \ln n)^{-\frac{1}{2s_{\max}+1}}$, the inequality (5.21) yields that as long as $4r\epsilon(2s_{\max}+1) < 1$, choosing k large enough will give

$$\sup_{f \in \mathcal{B}_{p,\infty}^s(L)} \mathbb{P}(|T_h - T_{h^*(s)}| > \gamma_n(h)) \leq \frac{C_9}{n^3}. \quad (5.22)$$

Plugging the previous tail bound into (5.15), we arrive at

$$\text{II} \leq \frac{C_9 L^2 |\mathcal{H}|}{n^3} \leq \frac{C_9 L^2 h_{\min}^{-1}}{n^3} = C_9 L^2 \cdot \frac{\ln n}{n^2} \leq C_{10} (n \ln n)^{-\frac{2s}{2s+1}},$$

and thereby complete the proof. \square

6 Technical Lemmas

In this section we collect some necessary technical lemmas necessary to prove the main results of this paper. We begin with a collection of lemmas available in literature which will serve as necessary tools to prove the other technical lemmas involved in the arguments laid down in Section 5.

Lemma 6.1. [*CL11*] For Hermite polynomial $H_k(x)$ of order k , if $X \sim \mathcal{N}(\mu, 1)$, we have

$$\mathbb{E}[H_k(X)] = \mu^k.$$

Moreover, if $|\mu| \leq M$ and $k \leq M^2$, we have

$$\mathbb{E}[H_k^2(X)] \leq (2M^2)^k.$$

Lemma 6.2. [*Ber12, VK87*] For any $r > 0$, the best polynomial approximation error of $|x|^r$ on $[-1, 1]$ satisfies

$$\inf_{Q \in \text{Poly}_n} \sup_{x \in [-1, 1]} |Q(x) - |x|^r| \leq \frac{\beta_r}{n^r},$$

where $\beta_r > 0$ is a universal constant depending on r only. Moreover, for n large enough, we can choose β_1 to be the Bernstein constant $\beta_* = 0.280169499$.

Lemma 6.3. [*BLM13*] For $X \sim \mathcal{N}(\mu, \sigma^2)$, we have

$$\mathbb{P}(|X - \mu| \geq t\sigma) \leq \exp\left(-\frac{t^2}{2}\right).$$

Lemma 6.4. *Let $X \sim \mathcal{N}(\mu, \sigma^2)$ with $\mu > c\sigma\sqrt{\ln n}$, where $c > 0, n \geq 2$. Then for any $\alpha \in \mathbb{R}$ and any integer $k \geq 2$, we have*

$$\begin{aligned} \mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha \right| &\leq D_\alpha \mu^\alpha, \\ \mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha - \mathbb{E} \left(\mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha \right) \right|^k &\leq D_{\alpha,k} \sigma^k \mu^{(\alpha-1)k}, \end{aligned}$$

where $D_\alpha, D_{\alpha,k} > 0$ are universal constants depending on α, k and c only.

Proof. Throughout the proof we use the asymptotic notation \lesssim to denote universal constants depending only on (α, c, k) .

For the first inequality, define

$$\begin{aligned} E_1 &\triangleq \{X : \frac{c\sigma\sqrt{\ln n}}{2} \leq X < \frac{\mu}{2}\}, \\ E_2 &\triangleq \{X : \frac{\mu}{2} \leq X \leq 2\mu\}, \quad \text{and} \\ E_3 &\triangleq \{X : X > 2\mu\}. \end{aligned}$$

By Lemma 6.3 and the triangle inequality, we have

$$\begin{aligned} \mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha \right| &\leq \sum_{i=1}^3 \mathbb{E} |\mathbb{1}(E_i) \cdot X^\alpha| \\ &\leq \max\{(\frac{c\sigma\sqrt{\ln n}}{2})^\alpha, (\frac{\mu}{2})^\alpha\} \cdot \mathbb{P}(E_1) + \max\{(\frac{\mu}{2})^\alpha, (2\mu)^\alpha\} + \mathbb{E} |X^\alpha \mathbb{1}(X \geq 2\mu)| \\ &\lesssim ((\sigma\sqrt{\ln n})^\alpha + \mu^\alpha) \cdot \exp(-\frac{\mu^2}{8\sigma^2}) + \mu^\alpha + \mathbb{E} |X^\alpha \mathbb{1}(X \geq 2\mu)|. \end{aligned}$$

For the last term, note that $X^\alpha \leq (1 + 2^\alpha)(X - \mu)^\alpha$ holds for any $\alpha \in \mathbb{R}$ when $X \geq 2\mu$, and

$$\mathbb{E}[(X - \mu)^{2\alpha} \mathbb{1}(X - \mu \geq c\sigma)] \lesssim \sigma^{2\alpha}$$

by the scaling property of the Gaussian random variable $X - \mu \sim \mathcal{N}(0, \sigma^2)$. Hence, by Lemma 6.3 and Cauchy-Schwartz,

$$\begin{aligned} \mathbb{E} |X^\alpha \mathbb{1}(X \geq 2\mu)| &\lesssim \mathbb{E} |(X - \mu)^\alpha \mathbb{1}(X \geq 2\mu)| \\ &\leq \mathbb{E}^{1/2} |X^{2\alpha} \mathbb{1}(X \geq 2\mu)| \cdot \mathbb{P}^{1/2}(X \geq 2\mu) \\ &\leq \mathbb{E}^{1/2} |(X - \mu)^{2\alpha} \mathbb{1}(X - \mu \geq c\sigma)| \exp(-\frac{\mu^2}{4\sigma^2}) \lesssim \sigma^\alpha \exp(-\frac{\mu^2}{4\sigma^2}). \end{aligned}$$

Combining the previous inequalities, we arrive at

$$\mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha \right| \lesssim (\sigma^\alpha + (\sigma\sqrt{\ln n})^\alpha + \mu^\alpha) \cdot \exp(-\frac{\mu^2}{8\sigma^2}) + \mu^\alpha. \quad (6.1)$$

If $\alpha \geq 0$, the desired inequality follows from $\sigma^\alpha \lesssim (\sigma\sqrt{\ln n})^\alpha \lesssim \mu^\alpha$ and $\exp(-\frac{\mu^2}{8\sigma^2}) \leq 1$. For $\alpha < 0$, the facts

$$(\sigma\sqrt{\ln n})^\alpha \lesssim \sigma^\alpha, \quad \exp(-\frac{\mu^2}{8\sigma^2}) \leq 1, \quad \frac{\sigma^\alpha}{\mu^\alpha} \exp(-\frac{\mu^2}{8\sigma^2}) \leq \max_{t \geq c\sqrt{\ln n}} t^{-\alpha} e^{-t^2/8} \lesssim 1$$

complete the proof of the desired inequality.

As for the second inequality, we first show the following chain of inequalities:

$$\begin{aligned} & \mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot X^\alpha - \mu^\alpha \right|^k \\ & \lesssim \mathbb{E} \left| \mathbb{1}(X \geq \frac{c\sigma\sqrt{\ln n}}{2}) \cdot (X^\alpha - \mu^\alpha) \right|^k + \mu^{k\alpha} \mathbb{P}(X < \frac{c\sigma\sqrt{\ln n}}{2}) \end{aligned} \quad (6.2)$$

$$\lesssim \sum_{i=1}^3 \mathbb{E}[\mathbb{1}(E_i) \cdot |X^\alpha - \mu^\alpha|^k] + \mu^{k\alpha} \exp(-\frac{\mu^2}{8\sigma^2}) \quad (6.3)$$

$$\lesssim \mathbb{E}[\mathbb{1}(E_2) \cdot |X^\alpha - \mu^\alpha|^k] + (\mu^{k\alpha} + (\sigma\sqrt{\ln n})^{k\alpha}) \exp(-\frac{\mu^2}{8\sigma^2}) \quad (6.4)$$

$$\lesssim \mathbb{E}[\mathbb{1}(E_2) \cdot |X^\alpha - \mu^\alpha|^k] + \sigma^k \mu^{(\alpha-1)k} \quad (6.5)$$

$$\lesssim \sup_{\xi \in [\mu/2, 2\mu]} |\xi|^{k(\alpha-1)} \mathbb{E}[\mathbb{1}(E_2) \cdot |X - \mu|^k] + \sigma^k \mu^{(\alpha-1)k} \quad (6.6)$$

$$\lesssim \sigma^k \mu^{(\alpha-1)k}. \quad (6.7)$$

We elaborate on the inequalities (6.2) to (6.7) here:

1. Inequality (6.2) follows from the triangle inequality $|a + b|^k \leq 2^{k-1}(|a|^k + |b|^k)$;
2. Inequality (6.3) follows from Lemma 6.3 and $c\sigma\sqrt{\ln n} \leq \mu$;
3. Inequality (6.4) follows from dealing with events E_1 and E_3 separately. For all $\alpha \in \mathbb{R}$, the condition $c\sigma\sqrt{\ln n}/2 \leq X \leq \mu/2$ implies that $|X^\alpha - \mu^\alpha| \lesssim (\sigma\sqrt{\ln n})^\alpha + \mu^\alpha$. Hence, by Lemma 6.3,

$$\begin{aligned} \mathbb{E}[\mathbb{1}(E_1) \cdot |X^\alpha - \mu^\alpha|^k] & \lesssim \mathbb{P}(E_1) \cdot ((\sigma\sqrt{\ln n})^\alpha + \mu^\alpha)^k \\ & \lesssim (\mu^{k\alpha} + (\sigma\sqrt{\ln n})^{k\alpha}) \exp(-\frac{\mu^2}{8\sigma^2}). \end{aligned}$$

As for the event E_3 , if $\alpha \leq 0$ we have $|X^\alpha - \mu^\alpha| \leq \mu^\alpha$ when $X \geq 2\mu$, and Lemma 6.3 gives

$$\mathbb{E}[\mathbb{1}(E_3) \cdot |X^\alpha - \mu^\alpha|^k] \leq \mu^{k\alpha} \mathbb{P}(E_3) \leq \mu^{k\alpha} \exp(-\frac{\mu^2}{2\sigma^2}).$$

If $\alpha > 0$, the condition $X \geq 2\mu$ ensures that

$$|X^\alpha - \mu^\alpha| \leq X^\alpha \leq 2^\alpha (X - \mu)^\alpha,$$

and thus Lemma 6.3 with Cauchy-Schwartz yields

$$\begin{aligned} \mathbb{E}[\mathbb{1}(E_3) |X^\alpha - \mu^\alpha|^k] & \lesssim \mathbb{E}[\mathbb{1}(E_3) (X - \mu)^{k\alpha}] \\ & \leq \mathbb{P}^{1/2}(E_3) \cdot \mathbb{E}^{1/2}[(X - \mu)^{2k\alpha}] \\ & \lesssim \sigma^{k\alpha} \exp(-\frac{\mu^2}{4\sigma^2}) \lesssim \mu^{k\alpha} \exp(-\frac{\mu^2}{4\sigma^2}), \end{aligned}$$

where the last step is due to the assumption $\sigma \lesssim \mu$;

4. If $\alpha \geq 0$, inequality (6.5) follows from $(\sigma\sqrt{\ln n})^{k\alpha} \lesssim \mu^{k\alpha}$ and

$$\frac{\mu^k}{\sigma^k} \cdot \exp\left(-\frac{\mu^2}{8\sigma^2}\right) \leq \max_{t \geq c\sqrt{\ln n}} t^k e^{-t^2/8} \lesssim 1$$

for any integer $k \geq 2$. If $\alpha < 0$, the desired inequality follows from

$$\mu^{k\alpha} + (\sigma\sqrt{\ln n})^{k\alpha} \lesssim \sigma^{k\alpha}$$

and

$$\frac{\mu^{k(1-\alpha)}}{\sigma^{k(1-\alpha)}} \cdot \exp\left(-\frac{\mu^2}{8\sigma^2}\right) \leq \max_{t \geq c\sqrt{\ln n}} t^{k(1-\alpha)} e^{-t^2/8} \lesssim 1$$

for any integer $k \geq 2$;

5. Inequality (6.6) follows from $X^\alpha - \mu^\alpha = \alpha\xi^{\alpha-1}(X - \mu)$ with some ξ lying between X and μ , as well as the definition of E_2 ;

6. Inequality (6.7) follows from $\mathbb{E}|X - \mu|^k \lesssim \sigma^k$ for $X \sim \mathcal{N}(\mu, \sigma^2)$.

Now observe that for any random variable Y and integer $k \geq 2$, by the triangle inequality and Jensen's inequality, the following inequality

$$\begin{aligned} \mathbb{E}|Y - \mathbb{E}Y|^k &\leq 2^{k-1} \left(\mathbb{E}|Y - c|^k + |\mathbb{E}Y - c|^k \right) \\ &\leq 2^{k-1} \left(\mathbb{E}|Y - c|^k + \mathbb{E}|Y - c|^k \right) \\ &= 2^k \mathbb{E}|Y - c|^k \end{aligned}$$

holds for any $c \in \mathbb{R}$. Hence, a combination of the previous two inequalities concludes the proof of the upper bound on k -th central moment. \square

Lemma 6.5. [QR07, Thm. E] Let $p_n(x) = \sum_{\nu=0}^n a_\nu x^\nu$ be a polynomial of degree at most n such that $|p_n(x)| \leq 1$ for $x \in [-1, 1]$. Then, $|a_{n-2\mu}|$ is bounded above by the modulus of the corresponding coefficient of T_n for $\mu = 0, 1, \dots, \lfloor n/2 \rfloor$, and $|a_{n-1-2\mu}|$ is bounded above by the modulus of the corresponding coefficient of T_{n-1} for $\mu = 0, 1, \dots, \lfloor (n-1)/2 \rfloor$. Here $T_n(x)$ is the n -th Chebyshev polynomials of the first kind.

Lemma 6.6. Let f and f_h be defined in Section 3, and let $r \geq 1$ and $k > 1$ be integers. There exists some universal constant c depending only on r, k and the kernel K such that

$$\|f_h\|_{k(r-1)}^{k(r-1)} \leq ch^{(k-1)(-1+1/r)} \|f\|_r^{(k-1)(r-1)} \|f_h\|_r^{r-1}.$$

Proof. The proof follows from the proof of [LNS99, Lemma 4.4] and is hence omitted. \square

Lemma 6.7. Suppose $\mathbb{1}(A)$ is an indicator random variable independent of X and Y and let $Z = X\mathbb{1}(A) + Y\mathbb{1}(A^c)$. Then

$$\text{Var}(Z) = \text{Var}(X)\mathbb{P}(A) + \text{Var}(Y)\mathbb{P}(A^c) + (\mathbb{E}X - \mathbb{E}Y)^2\mathbb{P}(A)\mathbb{P}(A^c).$$

In general for any integer $k \geq 2$,

$$\mathbb{E}|Z - \mathbb{E}Z|^k \leq 2^{k-1} \left(\mathbb{E}|X - \mathbb{E}X|^k \mathbb{P}(A) + \mathbb{E}|Y - \mathbb{E}Y|^k \mathbb{P}(A^c) + |\mathbb{E}X - \mathbb{E}Y|^k \mathbb{P}(A)\mathbb{P}(A^c) \right).$$

Proof. The identity for $\text{Var}(Z)$ follows from [CL11, Lemma 4]. For general k , by taking the expectation with respect to $\mathbb{1}(A)$ first, we have

$$\begin{aligned}\mathbb{E}|Z - \mathbb{E}Z|^k &= \mathbb{E}|X\mathbb{1}(A) + Y\mathbb{1}(A^c) - \mathbb{E}(X\mathbb{1}(A) + Y\mathbb{1}(A^c))|^k \\ &= \mathbb{P}(A)\mathbb{E}|X - \mathbb{E}X \cdot \mathbb{P}(A) - \mathbb{E}Y \cdot \mathbb{P}(A^c)|^k \\ &\quad + \mathbb{P}(A^c)\mathbb{E}|Y - \mathbb{E}X \cdot \mathbb{P}(A) - \mathbb{E}Y \cdot \mathbb{P}(A^c)|^k.\end{aligned}$$

By triangle inequality,

$$\begin{aligned}\mathbb{E}|X - \mathbb{E}X \cdot \mathbb{P}(A) - \mathbb{E}Y \cdot \mathbb{P}(A^c)|^k &= \mathbb{E}|(X - \mathbb{E}X) + (\mathbb{E}X - \mathbb{E}Y)\mathbb{P}(A^c)|^k \\ &\leq 2^{k-1} \left(\mathbb{E}|X - \mathbb{E}X|^k + |\mathbb{E}X - \mathbb{E}Y|^k \mathbb{P}(A^c)^k \right),\end{aligned}$$

and thus

$$\begin{aligned}\mathbb{E}|Z - \mathbb{E}Z|^k &\leq \mathbb{P}(A) \cdot 2^{k-1} (\mathbb{E}|X - \mathbb{E}X|^k + \mathbb{P}(A^c)^k |\mathbb{E}X - \mathbb{E}Y|^k) \\ &\quad + \mathbb{P}(A^c) \cdot 2^{k-1} (\mathbb{E}|Y - \mathbb{E}Y|^k + \mathbb{P}(A)^k |\mathbb{E}X - \mathbb{E}Y|^k) \\ &\leq 2^{k-1} \left(\mathbb{E}|X - \mathbb{E}X|^k \mathbb{P}(A) + \mathbb{E}|Y - \mathbb{E}Y|^k \mathbb{P}(A^c) + |\mathbb{E}X - \mathbb{E}Y|^k \mathbb{P}(A) \mathbb{P}(A^c) \right),\end{aligned}$$

where the last step uses $\mathbb{P}(A)^{k-1} + \mathbb{P}(A^c)^{k-1} \leq \mathbb{P}(A) + \mathbb{P}(A^c) = 1$ for $k \geq 2$. \square

Lemma 6.8. [Tsy08] *The total variation distance and the chi-squared distance are related via the following inequality:*

$$V(P, Q) \leq 1 - \frac{1}{2} \exp(-\chi^2(P, Q)).$$

Lemma 6.9. [HKPT12, Theorem C.2] *Let $q \geq 2$ and let X_1, \dots, X_n be independent random variables such that $\mathbb{E}(X_i) = 0$ and $\mathbb{E}|X_i|^q < \infty$.*

$$\mathbb{E} \left(\left| \sum_{i=1}^n X_i \right|^q \right) \leq C(q) \left[\sum_{i=1}^n \mathbb{E}|X_i|^q + \left(\sum_{i=1}^n \mathbb{E}|X_i|^2 \right)^{q/2} \right].$$

Finally, Lemma 6.10 presents the equivalence between Peetre's K -functional and modulus of smoothness on \mathbb{R} . For $p \in [1, \infty]$ and $r \in \mathbb{N}$, the Peetre's K -functional for f defined on \mathbb{R} is defined as

$$K_r(f, t^r)_p \triangleq \inf_g \|f - g\|_p + t^r \|g^{(r)}\|_p,$$

where the infimum is taken over all functions g defined on \mathbb{R} such that the derivative $g^{(r-1)}$ is locally absolutely continuous. Also recall the definition of the modulus of smoothness $\omega_r(f, t)_p$ in (2.1).

Lemma 6.10. *For any $p \in [1, \infty]$ and $r \in \mathbb{N}$, there exist universal constants $M = M(r, p)$ and $t_0 = t_0(r, p)$ such that for any $0 < t < t_0$ and f defined on \mathbb{R} ,*

$$M^{-1}K_r(f, t^r)_p \leq \omega_r(f, t)_p \leq MK_r(f, t^r)_p. \quad (6.8)$$

Furthermore,

$$\omega_r(f, t)_p \leq Mt^r \|f^{(r)}\|_p, \quad 0 < t < t_0, \quad (6.9)$$

and

$$\omega_r(f, t)_p \leq 2^r \|f\|_p, \quad t > 0. \quad (6.10)$$

Proof. The first inequality (6.8) is due to [DL93, Chapter 6, Theorem 2.4]. For the other inequalities, (6.9) follows from (6.8) by choosing $g = f$, and (6.10) follows from the definition (2.1) and the triangle inequality. \square

6.1 Proof of Lemma 5.1

The proof of Lemma 5.1 follows in turn from sequence of lemmas. We first consider the case where $|f_h(x)|$ is small for which the next lemma is crucial.

Lemma 6.11. *Let $|\mu| \leq 2c_1\lambda_h\sqrt{\ln n}$, and $X \sim \mathcal{N}(\mu, \lambda_h^2)$. Then for $c_2 \ln n \geq 1$, $4c_1^2 \geq c_2$, the bias and variance of $P(X)$ in estimating $|\mu|$ can be upper bounded as*

$$\begin{aligned} |\mathbb{E}P(X) - |\mu|| &\leq \frac{2c_1\beta_1}{c_2} \cdot \frac{\lambda_h}{\sqrt{\ln n}}, \\ \text{Var}(P(X)) &\leq 2^{7c_2 \ln n + 4} c_1^2 c_2^2 \lambda_h^2 (\ln n)^3, \end{aligned}$$

where the constant β_1 appears in Lemma 6.2.

Proof. By Lemma 6.1 we know that

$$\mathbb{E}P(X) = \sum_{k=0}^K g_{K,k} (2c_1\lambda_h\sqrt{\ln n})^{1-k} \cdot \mu^k.$$

By Lemma 6.2, we have

$$\sup_{x \in [-1, 1]} \left| \sum_{k=0}^K g_{K,k} x^k - |x| \right| \leq \frac{\beta_1}{K}.$$

By a variable substitution $x \mapsto \frac{\mu}{2c_1\lambda_h\sqrt{\ln n}}$, we obtain

$$\sup_{|\mu| \leq 2c_1\lambda_h\sqrt{\ln n}} \left| \sum_{k=0}^K g_{K,k} (2c_1\lambda_h\sqrt{\ln n})^{1-k} \cdot \mu^k - |\mu| \right| \leq \frac{\beta_1}{K} \cdot 2c_1\lambda_h\sqrt{\ln n}.$$

Hence, the bias of $P(X)$ is upper bounded by

$$|\mathbb{E}P(X) - |\mu|| \leq \frac{\beta_1}{K} \cdot 2c_1\lambda_h\sqrt{\ln n} = \frac{2c_1\beta_1}{c_2} \cdot \frac{\lambda_h}{\sqrt{\ln n}},$$

as desired.

As for the variance, first Lemma 6.5 tells us

$$|g_{K,k}| \leq 2^{3K}, \quad k = 0, 1, \dots, K.$$

Hence, with the help of Lemma 6.1, we know that

$$\begin{aligned} \text{Var}(P(X)) &\leq \mathbb{E}[P(X)^2] \\ &\leq (K+1) \sum_{k=0}^K |g_{K,k}|^2 (2c_1\lambda_h\sqrt{\ln n})^{2(1-k)} \cdot \lambda_h^{2k} \mathbb{E}[H_k(\frac{X}{\lambda_h})^2] \\ &\leq 2^{6K+1} K \sum_{k=0}^K (2c_1\lambda_h\sqrt{\ln n})^{2(1-k)} \cdot \lambda_h^{2k} [2(2c_1\sqrt{\ln n})^2]^k \\ &\leq 2^{7K+3} K \sum_{k=0}^K c_1^2 \lambda_h^2 \ln n \\ &\leq 2^{7K+4} c_1^2 K^2 \lambda_h^2 \ln n, \end{aligned}$$

where we have used the fact that $k \leq K \leq (2c_1\sqrt{\ln n})^2$. □

The next lemma is useful to analyze the plug-in approach where $f_h(x)$ is large. The key observation in this regime is that the plug-in approach is almost unbiased due to the measure concentration property of Gaussian distribution.

Lemma 6.12. *Let $|\mu| \geq \frac{c_1}{2}\lambda_h\sqrt{\ln n}$, $X \sim \mathcal{N}(\mu, \lambda_h^2)$, and $k \geq 2$ be any integer. Then*

$$\begin{aligned} |\mathbb{E}|X| - |\mu|| &\leq \frac{4\lambda_h}{c_1\sqrt{\ln n}} \cdot n^{-c_1^2/8}, \\ \mathbb{E}||X| - \mathbb{E}|X||^k &\leq C(c_1, k)\lambda_h^k, \end{aligned}$$

where $C(c_1, k)$ is a universal constant depending on k and c_1 only. In particular, when $k = 2$, we have

$$\text{Var}(|X|) \leq C(c_1, 2)\lambda_h^2.$$

Proof. By symmetry we can assume that $\mu \geq \frac{c_1}{2}\lambda_h\sqrt{\ln n}$, then the bias can be written as

$$|\mathbb{E}|X| - \mu| = |\mathbb{E}|X| - \mathbb{E}X| = 2\mathbb{E}|X|\mathbb{1}(X \leq 0).$$

With the help of the Gaussian tail bound (Lemma 6.3), we have

$$\begin{aligned} |\mathbb{E}|X| - \mu| &= 2 \int_{-\infty}^0 \mathbb{P}(X \leq t) dt \\ &= 2 \int_{-\infty}^0 \mathbb{P}\left(\frac{X - \mu}{\lambda_h} \leq \frac{t - \mu}{\lambda_h}\right) dt \\ &\leq 2 \int_{-\infty}^0 \exp\left(-\frac{(t - \mu)^2}{2\lambda_h^2}\right) dt \\ &\leq \frac{2\lambda_h^2}{\mu} \int_{-\infty}^0 \frac{\mu - t}{\lambda_h^2} \exp\left(-\frac{(t - \mu)^2}{2\lambda_h^2}\right) dt \\ &= \frac{2\lambda_h^2}{\mu} \exp\left(-\frac{\mu^2}{2\lambda_h^2}\right) \leq \frac{4\lambda_h}{c_1\sqrt{\ln n}} \cdot n^{-c_1^2/8}, \end{aligned}$$

which completes the proof of the bias bound.

As for the k -th central moment, we have

$$\begin{aligned} \mathbb{E}||X| - \mathbb{E}|X||^k &\leq 3^{k-1} \left(\mathbb{E}||X| - X|^k + \mathbb{E}|X - \mu|^k + |\mathbb{E}|X| - \mu|^k \right) \\ &\leq 3^{k-1} \left(2^k \mathbb{E}|X|^k \mathbb{1}(X \leq 0) + \lambda_h^k (k-1)!! + \left(\frac{4\lambda_h}{c_1\sqrt{\ln n}} \cdot n^{-c_1^2/8} \right)^k \right) \end{aligned}$$

where we have used the previous bias bound for the last term. Using the same technique as before,

$$\begin{aligned} \mathbb{E}|X|^k \mathbb{1}(X \leq 0) &= k \int_{-\infty}^0 |t|^{k-1} \mathbb{P}(X \leq t) dt \\ &\leq k \int_{-\infty}^0 |t|^{k-1} \exp\left(-\frac{(t - \mu)^2}{2\lambda_h^2}\right) dt \\ &\leq k\lambda_h^k \int_{-\infty}^0 \frac{1}{\lambda_h} \left(\frac{\mu - t}{\lambda_h}\right)^{k-1} \exp\left(-\frac{(t - \mu)^2}{2\lambda_h^2}\right) dt \end{aligned}$$

$$\begin{aligned}
&= k\lambda_h^k \cdot \sqrt{2\pi} \mathbb{E} \left(Z^{k-1} \mathbb{1} \left(Z \geq \frac{c_1}{2} \sqrt{\ln n} \right) \right) \\
&\leq C\lambda_h^k n^{-c_1^2/16},
\end{aligned}$$

where $Z = \frac{\mu - X}{\lambda_h} \sim \mathcal{N}(0, 1)$ is a standard normal random variable, the constant C depends on c_1 and k , and the last inequality is due to Cauchy–Schwartz

$$\mathbb{E} \left(Z^{k-1} \mathbb{1} \left(Z \geq \frac{c_1}{2} \sqrt{\ln n} \right) \right) \leq \mathbb{E}^{1/2}(Z^{2k-2}) \mathbb{P}^{1/2}(Z \geq \frac{c_1}{2} \sqrt{\ln n})$$

and Lemma 6.3. Hence the k -th central moment bound is proved. \square

Proof of Lemma 5.1. Throughout the proof, for any two sequences a_n, b_n , we will use the notation $a_n \lesssim b_n$ whenever $|\frac{a_n}{b_n}|$ is upper bounded by a universal constant which only depends on c_1, c_2, k .

Note that X, Y are independent, Lemma 6.7 can be employed here to establish upper bounds on k^{th} central moments of $\xi(X, Y)$. We distinguish into three cases:

1. Case I: $|\mu| \leq \frac{c_1}{2} \lambda_h \sqrt{\ln n}$. By Lemma 6.11 and Markov's inequality,

$$\begin{aligned}
|\mathbb{E}\tilde{P}(X) - |\mu|| &\leq |\mathbb{E}P(X) - |\mu|| + \mathbb{E}|P(X) - \tilde{P}(X)| \\
&= |\mathbb{E}P(X) - |\mu|| + \mathbb{E}|P(X) \mathbb{1}(|P(X)| \geq n^{2\epsilon} \lambda_h)| \\
&\lesssim \frac{\lambda_h}{\sqrt{\ln n}} + \frac{\mathbb{E}|P(X)|^2}{n^{2\epsilon} \lambda_h} \lesssim \frac{\lambda_h}{\sqrt{\ln n}} + \frac{\lambda_h}{n^\epsilon}.
\end{aligned} \tag{6.11}$$

Hence,

$$\begin{aligned}
|\mathbb{E}\xi(X, Y) - |\mu|| &\leq |\mathbb{E}\tilde{P}(X) - |\mu|| + (\mathbb{E}|\tilde{P}(X)| + \mathbb{E}|X|) \cdot \mathbb{P}(|Y| \geq c_1 \lambda_h \sqrt{\ln n}) \\
&\leq |\mathbb{E}\tilde{P}(X) - |\mu|| + (n^{2\epsilon} \lambda_h + |\mu| + \lambda_h) \cdot \mathbb{P} \left(\left| \frac{Y - \mu}{\lambda_h} \right| \geq \frac{c_1}{2} \sqrt{\ln n} \right) \\
&\lesssim \frac{\lambda_h}{\sqrt{\ln n}} + \frac{\lambda_h}{n^\epsilon} + (n^{2\epsilon} \lambda_h + \frac{c_1}{2} \lambda_h \sqrt{\ln n} + \lambda_h) \cdot n^{-c_1^2/8} \lesssim \frac{\lambda_h}{\sqrt{\ln n}},
\end{aligned}$$

where

- (a) the first inequality follows from the triangle inequality;
- (b) the second inequality follows from $\mathbb{E}|X| \leq \mu + \mathbb{E}|X - \mu| \leq \mu + \lambda_h$;
- (c) the third inequality follows from (6.11), the assumption $|\mu| \leq \frac{c_1}{2} \lambda_h \sqrt{\ln n}$ and Lemma 6.3;
- (d) the last inequality follows from $c_1 > \sqrt{8k}$.

As for the k -th central moment, Lemma 6.7 yields

$$\begin{aligned}
&\mathbb{E}|\xi(X, Y) - \mathbb{E}\xi(X, Y)|^k \\
&\lesssim \mathbb{E}|\tilde{P}(X) - \mathbb{E}\tilde{P}(X)|^k + (\mathbb{E}||X| - \mathbb{E}|X||^k + |\mathbb{E}\tilde{P}(X) - \mathbb{E}|X||^k) \cdot \mathbb{P}(|Y| \geq c_1 \lambda_h \sqrt{\ln n}) \\
&\lesssim \mathbb{E}|\tilde{P}(X)|^k + \left(\lambda_h^k + (n^{2\epsilon} \lambda_h + |\mu| + \lambda_h)^k \right) \cdot n^{-c_1^2/8} \lesssim (\lambda_h n^{2\epsilon})^k,
\end{aligned}$$

where the second inequality follows from the triangle inequality, Lemma 6.12 and $|\mathbb{E}\tilde{P}(X) - \mathbb{E}|X|| \leq |\mathbb{E}\tilde{P}(X)| + |\mu| + \mathbb{E}|X - \mu| \leq n^{2\epsilon} \lambda_h + |\mu| + \lambda_h$.

2. Case II: $\frac{c_1}{2}\lambda_h\sqrt{\ln n} < |\mu| < 2c_1\lambda_h\sqrt{\ln n}$. For the bias bound, we employ the triangle inequality, (6.11) and Lemma 6.12 to obtain

$$\begin{aligned} |\mathbb{E}\xi(X, Y) - |\mu|| &\leq |\mathbb{E}\tilde{P}(X) - |\mu|| + |\mathbb{E}|X| - |\mu|| \\ &\lesssim \frac{\lambda_h}{\sqrt{\ln n}} + \frac{\lambda_h}{n^\epsilon} + \frac{\lambda_h}{\ln n} \cdot n^{-c_1^2/8} \lesssim \frac{\lambda_h}{\sqrt{\ln n}}. \end{aligned}$$

As for the k -th central moment, by Lemma 6.7 we have

$$\begin{aligned} &|\mathbb{E}\xi(X, Y) - \mathbb{E}\xi(X, Y)|^k \\ &\lesssim \mathbb{E}|\tilde{P}(X) - \mathbb{E}\tilde{P}(X)|^k + \mathbb{E}||X| - \mathbb{E}|X||^k + |\mathbb{E}\tilde{P}(X) - \mathbb{E}|X||^k \\ &\lesssim \mathbb{E}|\tilde{P}(X)|^k + \mathbb{E}||X| - \mathbb{E}|X||^k + |\mathbb{E}\tilde{P}(X) - |\mu||^k + |\mathbb{E}|X| - |\mu||^k \\ &\lesssim (\lambda_h n^{2\epsilon})^k + \lambda_h^k + \left(\frac{\lambda_h}{\sqrt{\ln n}} + \frac{\lambda_h}{n^\epsilon}\right)^k + \left(\frac{\lambda_h}{\sqrt{\ln n}} \cdot n^{-c_1^2/8}\right)^k \lesssim (\lambda_h n^{2\epsilon})^k, \end{aligned}$$

where the third inequality follows from Lemma 6.12 and (6.11).

3. Case III: $|\mu| \geq 2c_1\lambda_h\sqrt{\ln n}$. By Lemma 6.12 and Lemma 6.3, the bias bound is given by

$$\begin{aligned} |\mathbb{E}\xi(X, Y) - |\mu|| &\leq |\mathbb{E}|X| - |\mu|| + (\mathbb{E}|X| + \mathbb{E}|\tilde{P}(X)|) \cdot \mathbb{P}(|Y| \leq c_1\lambda_h\sqrt{\ln n}) \\ &\leq |\mathbb{E}|X| - |\mu|| + (\mathbb{E}|X| + \mathbb{E}|\tilde{P}(X)|) \cdot \mathbb{P}\left(\left|\frac{Y - \mu}{\lambda_h}\right| \geq \frac{|\mu|}{2\lambda_h}\right) \\ &\lesssim \frac{\lambda_h}{\sqrt{\ln n}} \cdot n^{-c_1^2/8} + (|\mu| + \lambda_h + \lambda_h n^{2\epsilon}) \cdot \exp\left(-\frac{\mu^2}{8\lambda_h^2}\right) \lesssim \frac{\lambda_h}{\sqrt{\ln n}}, \end{aligned}$$

where the last inequality follows from $c_1 > \sqrt{8k}$ and

$$\sup_{|\mu| \geq 2c_1\lambda_h\sqrt{\ln n}} |\mu| \exp\left(-\frac{\mu^2}{8\lambda_h^2}\right) \lesssim \frac{1}{n}.$$

By Lemmas 6.3, 6.7 and 6.12, the k -th central moment can be upper bounded as

$$\begin{aligned} &|\mathbb{E}\xi(X, Y) - \mathbb{E}\xi(X, Y)|^k \\ &\lesssim \mathbb{E}(|X| - \mathbb{E}|X|)^k + (\mathbb{E}|\tilde{P}(X) - \mathbb{E}\tilde{P}(X)|^k + |\mathbb{E}|X| - \mathbb{E}\tilde{P}(X)|^k) \cdot \mathbb{P}(|Y| \leq c_1\lambda_h\sqrt{\ln n}) \\ &\lesssim \lambda_h^k + ((\lambda_h n^{2\epsilon})^k + (|\mu| + \lambda_h + \lambda_h n^{2\epsilon})^k) \cdot \exp\left(-\frac{\mu^2}{8\lambda_h^2}\right) \lesssim \lambda_h^k, \end{aligned}$$

where again the last step follows from taking supremum over $|\mu| \geq 2c_1\lambda_h\sqrt{\ln n}$.

Combining these three cases completes the proof of the lemma. \square

6.2 Proof of Lemma 5.2

The proof of Lemma 5.2 follows in turn from sequence of lemmas. We first consider the case where $|f_h(x)|$ is small for which the next lemma is crucial.

Lemma 6.13. *Let $|\mu| \leq 2c_1\lambda_h\sqrt{\ln n}$, and $X \sim \mathcal{N}(\mu, \lambda_h^2)$. Then for $c_2 \ln n \geq 1$, $4c_1^2 \geq c_2$, the bias and variance of $P_r(X)$ in estimating $|\mu|^r$ can be upper bounded as*

$$|\mathbb{E}P_r(X) - |\mu|^r| \leq \beta_r \cdot \left(\frac{2c_1\lambda_h}{c_2\sqrt{\ln n}}\right)^r,$$

$$\text{Var}(P_r(X)) \leq 2^{7c_2 \ln n + 2} c_2^2 (\ln n)^2 \cdot (2c_1 \lambda_h \sqrt{\ln n})^{2r},$$

where the constant β_r appears in Lemma 6.2.

Proof. By Lemma 6.1 we know that

$$\mathbb{E}P_r(X) = \sum_{k=0}^K g_{K,k}^{(r)} (2c_1 \lambda_h \sqrt{\ln n})^{r-k} \cdot \mu^k.$$

By Lemma 6.2, we have

$$\sup_{x \in [-1, 1]} \left| \sum_{k=0}^K g_{K,k}^{(r)} x^k - |x|^r \right| \leq \frac{\beta_r}{K^r}.$$

By a variable substitution $x \mapsto \frac{\mu}{2c_1 \lambda_h \sqrt{\ln n}}$, we obtain

$$\sup_{|\mu| \leq 2c_1 \lambda_h \sqrt{\ln n}} \left| \sum_{k=0}^K g_{K,k}^{(r)} (2c_1 \lambda_h \sqrt{\ln n})^{r-k} \cdot \mu^k - |\mu|^r \right| \leq \beta_r \cdot \left(\frac{2c_1 \lambda_h \sqrt{\ln n}}{K} \right)^r.$$

Hence, the bias of $P_r(X)$ is upper bounded by

$$|\mathbb{E}P_r(X) - |\mu|^r| \leq \beta_r \cdot \left(\frac{2c_1 \lambda_h \sqrt{\ln n}}{K} \right)^r = \beta_r \cdot \left(\frac{2c_1 \lambda_h}{c_2 \sqrt{\ln n}} \right)^r,$$

as desired.

As for the variance, first Lemma 6.5 tells us

$$|g_{K,k}| \leq 2^{3K}, \quad k = 0, 1, \dots, K.$$

Hence, with the help of Lemma 6.1, we know that

$$\begin{aligned} \text{Var}(P_r(X)) &\leq \mathbb{E}[P_r(X)^2] \\ &\leq (K+1) \sum_{k=0}^K |g_{K,k}^{(r)}|^2 (2c_1 \lambda_h \sqrt{\ln n})^{2(r-k)} \cdot \lambda_h^{2k} \mathbb{E} \left[H_k \left(\frac{X}{\lambda_h} \right)^2 \right] \\ &\leq 2^{6K+1} K \sum_{k=0}^K (2c_1 \lambda_h \sqrt{\ln n})^{2(r-k)} \cdot \lambda_h^{2k} [2(2c_1 \sqrt{\ln n})^2]^k \\ &\leq 2^{7K+1} K \sum_{k=0}^K (2c_1 \lambda_h \sqrt{\ln n})^{2r} \\ &\leq 2^{7K+2} K^2 (2c_1 \lambda_h \sqrt{\ln n})^{2r}, \end{aligned}$$

where we have used the fact that $k \leq K \leq (2c_1 \sqrt{\ln n})^2$. \square

Next we analyze the “smooth” regime where $|f_h(x)|$ is large. If $f_h(x) > 0$, the Taylor expansion based estimator $S_{\lambda_h}(\tilde{f}_{h,1}(x), \tilde{f}_{h,2}(x))$ is analyzed in detail in the following lemma. The analysis of the estimator $S_{\lambda_h}(-\tilde{f}_{h,1}(x), -\tilde{f}_{h,2}(x))$ in the case $f_h(x) < 0$ then follows by symmetry. Subsequently, we can take into account the sample splitting approach, and following the same approach as of the proof of Lemma 5.1 one can complete the proof of Lemma 5.2. We omit the details.

Lemma 6.14. Let $\mu \geq \frac{c_1}{2} \lambda_h \sqrt{\ln n}$, $k \geq 2$ be any integer, and $X_1, X_2 \sim \mathcal{N}(\mu, \lambda_h^2)$ be independent. The bias and k -th central moment of $S_{\lambda_h}(X_1, X_2)$ in estimating μ^r can be upper bounded as

$$\begin{aligned} |\mathbb{E} S_{\lambda_h}(X_1, X_2) - \mu^r| &\leq \lambda_h^r (\sqrt{\ln n})^{r-R-1} + (\lambda_h \sqrt{\ln n})^r n^{-c_1^2/32}, \\ \mathbb{E} |S_{\lambda_h}(X_1, X_2) - \mathbb{E} S_{\lambda_h}(X_1, X_2)|^k &\leq C(r, k) \lambda_h^k \mu^{(r-1)k}, \end{aligned}$$

where $C(r, k) > 0$ is a universal constant depending only on r, k and c_1 . In particular, for $k = 2$ we have

$$\text{Var}(S_{\lambda_h}(X_1, X_2)) \leq C(r, 2) \lambda_h^2 \mu^{2r-2}.$$

Proof. Throughout the proof we use the notation $a_n \lesssim b_n$ to show that $|\frac{a_n}{b_n}|$ is upper bounded by a universal constant which only depends on c_1, k and r .

First we analyze the bias. By Lemma 6.1 and independence of X_1 and X_2 ,

$$\begin{aligned} \mathbb{E} S_{\lambda_h}(X_1, X_2) &= \mathbb{E} \left[\mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}) \cdot \sum_{k=0}^R a_k X_1^{r-k} \sum_{j=0}^k \binom{k}{j} \mathbb{E}_{X_2} \left(\lambda_h^j H_j \left(\frac{X_2}{\lambda_h} \right) \right) (-X_1)^{k-j} \right] \\ &= \mathbb{E} \left[\mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}) \cdot \sum_{k=0}^R a_k X_1^{r-k} \sum_{j=0}^k \binom{k}{j} \mu^j (-X_1)^{k-j} \right] \\ &= \mathbb{E} \left[\mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}) \cdot \sum_{k=0}^R a_k X_1^{r-k} (\mu - X_1)^k \right], \end{aligned}$$

where $a_k \triangleq \frac{r(r-1)\cdots(r-k+1)}{k!}$ is the Taylor coefficient. Note that by the Taylor expansion with Lagrange remainder term, we have

$$\mu^r - \sum_{k=0}^R a_k X_1^{r-k} (\mu - X_1)^k = a_{R+1} \xi^{r-R-1} (\mu - X_1)^{R+1},$$

for some ξ lying between X_1 and μ . In view of $\mu \geq \frac{c_1}{2} \lambda_h \sqrt{\ln n}$ and $X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}$, we conclude that $\xi \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}$. Hence, the triangle inequality yields

$$\begin{aligned} &|\mathbb{E} S_{\lambda_h}(X_1, X_2) - \mu^r| \\ &\leq \mu^r \mathbb{P}(X_1 < \frac{c_1}{4} \lambda_h \sqrt{\ln n}) + \mathbb{E} \left| \mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}) \cdot \left(\sum_{k=0}^R a_k X_1^{r-k} (\mu - X_1)^k - \mu^r \right) \right| \\ &\leq \mu^r \mathbb{P}(|X_1 - \mu| \geq \frac{\mu}{2}) + \sup_{\xi \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}} \mathbb{E} |\mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}) \cdot a_{R+1} \xi^{r-R-1} (\mu - X_1)^{R+1}| \\ &\leq 2\mu^r \exp(-\frac{\mu^2}{8\lambda_h^2}) + \left(\frac{c_1}{4} \lambda_h \sqrt{\ln n} \right)^{r-R-1} |a_{R+1}| \cdot \mathbb{E} |\mu - X_1|^{R+1} \\ &\lesssim (\lambda_h \sqrt{\ln n})^r n^{-c_1^2/32} + \lambda_h^r (\sqrt{\ln n})^{r-R-1}, \end{aligned}$$

where we have used Lemma 6.3, $r - R - 1 < 0$ and

$$\max_{\mu \geq c_1 \lambda_h \sqrt{\ln n}/2} \mu^r \exp(-\frac{\mu^2}{8\lambda_h^2}) \lesssim (\lambda_h \sqrt{\ln n})^r n^{-c_1^2/32},$$

$$\mathbb{E}|X_1 - \mu|^k \leq \sqrt{\mathbb{E}|X_1 - \mu|^{2k}} \lesssim \lambda_h^k, \quad k = 0, 1, \dots$$

This proves the bias bound.

As for the k -th central moment, we may write

$$S_\lambda(X_1, X_2) = \sum_{\ell=0}^R \sum_{j=0}^{\ell} b_{j,\ell} \zeta_j \eta_j,$$

where

$$\begin{aligned} b_{j,\ell} &\triangleq \frac{r(r-1)\cdots(r-\ell+1)}{\ell!} \cdot (-1)^{\ell-j} \binom{\ell}{j}, \\ \zeta_j &\triangleq X_1^{r-j} \mathbb{1}(X_1 \geq \frac{c_1}{4} \lambda_h \sqrt{\ln n}), \\ \eta_j &\triangleq \lambda_h^j H_j\left(\frac{X_2}{\lambda_h}\right). \end{aligned}$$

By independence of X_1 and X_2 ,

$$S_\lambda(X_1, X_2) - \mathbb{E}[S_\lambda(X_1, X_2)] = \sum_{\ell=0}^R \sum_{j=0}^{\ell} b_{j,\ell} [(\zeta_j - \mathbb{E}[\zeta_j])\mathbb{E}[\eta_j] + \zeta_j(\eta_j - \mathbb{E}[\eta_j])].$$

Since the coefficients $b_{j,\ell}$ do not depend on μ or λ_h , by the triangle inequality it suffices to prove that for any $j = 0, 1, \dots, R$,

$$A_j \triangleq \mathbb{E}[|\zeta_j - \mathbb{E}[\zeta_j]|^k] \cdot |\mathbb{E}[\eta_j]|^k \lesssim \lambda_h^k \mu^{(r-1)k}, \quad (6.12)$$

$$B_j \triangleq \mathbb{E}[|\zeta_j|^k] \cdot \mathbb{E}[|\eta_j - \mathbb{E}[\eta_j]|^k] \lesssim \lambda_h^k \mu^{(r-1)k}. \quad (6.13)$$

For the first term A_j , the second inequality of Lemma 6.4 with $\sigma = \lambda_h, \alpha = r - j$ and $c = c_1/2$ gives

$$\mathbb{E}[|\zeta_j - \mathbb{E}[\zeta_j]|^k] \lesssim \lambda_h^k \mu^{(r-j-1)k}.$$

Moreover, Lemma 6.1 gives $\mathbb{E}[\eta_j] = \mu^j$, and therefore (6.12) holds.

For the second term B_j , we may assume that $j \geq 1$ since $B_0 = 0$. For the k -th moment of ζ_j , the first inequality of Lemma 6.4 with $\sigma = \lambda_h, \alpha = r - j$ and $c = c_1/2$ gives $\mathbb{E}[|\zeta_j|^k] \lesssim \mu^{(r-j)k}$. To upper bound the k -th central moment of η_j , we express the j -th Hermite polynomial H_j as $H_j(x) = \sum_{i=0}^j a_{i,j} x^i$, then

$$\eta_j - \mathbb{E}[\eta_j] = \sum_{i=0}^j a_{i,j} \lambda_h^{j-i} (X_2^i - \mathbb{E}[X_2^i]) = \sum_{i=1}^j a_{i,j} \lambda_h^{j-i} (X_2^i - \mathbb{E}[X_2^i]).$$

Write $X_2 = \mu + \lambda_h Z$, with $Z \sim \mathcal{N}(0, 1)$. Then for $i \geq 1$,

$$X_2^i - \mathbb{E}X_2^i = (\mu + \lambda_h Z)^i - \mathbb{E}(\mu + \lambda_h Z)^i = \sum_{i'=1}^i \binom{i}{i'} \mu^{i-i'} \lambda_h^{i'} (Z^{i'} - \mathbb{E}[Z^{i'}]).$$

Since all moments of Z are finite, the i' -th summand has k -th moment $\lesssim \mu^{(i-i')k} \lambda_h^{i'k}$, and the triangle inequality then yields

$$\mathbb{E}|X_2^i - \mathbb{E}X_2^i|^k \lesssim \sum_{i'=1}^i \mu^{(i-i')k} \lambda_h^{i'k} \lesssim \lambda_h^k (\mu^{(i-1)k} + \lambda_h^{(i-1)k}) \lesssim \lambda_h^k \mu^{(i-1)k},$$

where the last inequality follows from $i \geq 1$ and the assumption $\lambda_h \lesssim \mu$. Therefore, by a triangle inequality again,

$$\mathbb{E}[|\eta_j - \mathbb{E}[\eta_j]|^k] \lesssim \sum_{i=1}^j \lambda_h^{(j-i)k} \cdot \mathbb{E}|X_2^i - \mathbb{E}X_2^i|^k \lesssim \lambda_h^k \sum_{i=1}^j \lambda_h^{(j-i)k} \mu^{(i-1)k} \lesssim \lambda_h^k \mu^{(j-1)k}.$$

Finally, combining the previous upper bounds yields

$$B_j = \mathbb{E}[|\zeta_j|^k] \cdot \mathbb{E}[|\eta_j - \mathbb{E}[\eta_j]|^k] \lesssim \mu^{(r-j)k} \cdot \lambda_h^k \mu^{(j-1)k} = \lambda_h^k \mu^{(r-1)k},$$

which gives (6.13). Combining (6.12) and (6.13) completes the proof of the upper bound for the k -th central moment. \square

6.3 Proof of Lemma 5.7

For the simplicity of the proof notation, we shall assume that $J = 1/h$ is an integer. The more general proof follows with obvious modifications by working with $\lfloor 1/h \rfloor$. We only provide the proof here for the case when J is an even integer i.e. $J = 2L$ for some $L \geq 1$. The proof for J odd can be obtained similarly.

In particular, we consider the partition of $[0, 1]$ into $2L$ consecutive subintervals of length h each and for $l = 0, 1, \dots, 2L-1$ denote the l^{th} subinterval by I_l i.e. $I_l = [(l-1)h, lh)$ for $l = 0, \dots, 2L-2$ and $I_{2L-1} = [(2L-1)h, 2Lh]$. Let $I_1 = \bigcup_{l=0}^{L-1} I_{2l}$ and $I_2 = \bigcup_{l=0}^{L-1} I_{2l+1}$. Then

$$\begin{aligned} & \int_0^1 (T_h(x) - \mathbb{E}T_h(x)) dx \\ &= \int_{I_1} (T_h(x) - \mathbb{E}T_h(x)) dx + \int_{I_2} (T_h(x) - \mathbb{E}T_h(x)) dx \\ &= T_1 + T_2. \end{aligned}$$

Indeed,

$$\mathbb{E} \left| \int_0^1 (T_h(x) - \mathbb{E}T_h(x)) dx \right|^k \leq 2^{k-1} \left(\mathbb{E}|T_1|^k + \mathbb{E}|T_2|^k \right). \quad (6.14)$$

We now provide control over $\mathbb{E}|T_1|^k$. The bound over $\mathbb{E}|T_2|^k$ is similar and combining them shall yield the desired proof of the lemma. First note that

$$T_1 = \sum_{l=0}^{L-1} \int_{2lh}^{(2l+1)h} (T_h(x) - \mathbb{E}T_h(x)) dx = \sum_{l=0}^{L-1} \xi_{h,l},$$

where $\xi_{h,l}(x) = \int_{2lh}^{(2l+1)h} (T_h(x) - \mathbb{E}T_h(x)) dx$ are independent and zero-mean random variables for $l = 0, \dots, L-1$. Therefore, by Rosenthal's Inequality (Lemma 6.9) we have that

$$\mathbb{E}|T_1|^k = \mathbb{E} \left| \sum_{l=0}^{L-1} \xi_{h,l} \right|^k \leq C(k) \left[\sum_{l=0}^{L-1} \mathbb{E}|\xi_{h,l}|^k + \left(\sum_{l=0}^{L-1} \mathbb{E}|\xi_{h,l}|^2 \right)^{k/2} \right]. \quad (6.15)$$

Now, by Jensen's Inequality on the interval of length h

$$\mathbb{E}|\xi_{h,l}|^k = \mathbb{E} \left| \int_{2lh}^{(2l+1)h} (T_h(x) - \mathbb{E}(T_h(x))) dx \right|^k$$

$$\begin{aligned}
&\leq h^k \frac{1}{h} \int_{2lh}^{(2l+1)h} \mathbb{E}|T_h(x) - \mathbb{E}(T_h(x))|^k dx \\
&\leq h^k \frac{1}{h} \int_{2lh}^{(2l+1)h} C_2 n^{2k\epsilon} \left(\lambda_h^{kr} + \lambda_h^k |f_h(x)|^{k(r-1)} \right), \tag{6.16}
\end{aligned}$$

where the last inequality follows from Lemma 5.1 (for $r = 1$) and Lemma 5.2 (for $r > 1$) with C_2 a constant depending on $c_1, c_2, \epsilon, \sigma, K_M, k$. Plugging in the bound (6.16) into (6.15) and subsequently combining with (6.14) completes the proof of Lemma 5.7. \square

6.4 Proof of Lemma 5.5

Let S be the optimal value of the optimization program in (5.13), then the target is to prove that $S = 2E_{q-1,K}(f; I)$. We first show that $S \leq 2E_{q-1,K}(f; I)$. Let $P(x) = \sum_{k=-q+1}^K a_k x^k$ be the best approximating rational function attaining the approximation error $E_{q-1,K}(f; I)$, then for any feasible pair of probability measures (ν_0, ν_1) in (5.13), the triangle inequality gives

$$\begin{aligned}
\int f(x)(\nu_1(dx) - \nu_0(dx)) &= \int (f(x) - P(x))(\nu_1(dx) - \nu_0(dx)) \\
&\leq \int |f(x) - P(x)|(\nu_1(dx) + \nu_0(dx)) \\
&\leq \int E_{q-1,K}(f; I)(\nu_1(dx) + \nu_0(dx)) \\
&= 2E_{q-1,K}(f; I).
\end{aligned}$$

Consequently, $S \leq 2E_{q-1,K}(f; I)$. To show the other inequality, we construct the probability measures ν_0 and ν_1 explicitly. Since the interval I does not contain zero, the functions $\{x^{-q+1}, x^{-q+2}, \dots, x^K\}$ in $C(I)$ form a Chebyshev system [DL93, Section 3.3, Example 2], and the Chebyshev alternation theorem [DL93, Chapter 3, Theorem 5.1] shows that there exist a rational function $P(x) = \sum_{k=-q+1}^K a_k x^k$ and points $x_0 < x_1 < \dots < x_{q+K}$ in I with $f(x_i) - P(x_i) = \varepsilon \cdot (-1)^i E_{q-1,K}(f; I)$ for all $i = 0, 1, \dots, q+K$, and $\varepsilon \in \{\pm 1\}$. Construct the signed measure ν supported on $\{x_0, x_1, \dots, x_{q+K}\}$ with

$$\nu(\{x_i\}) = c_0 \cdot x_i^{q-1} \prod_{j \neq i} \frac{1}{x_i - x_j},$$

where $c_0 \in \mathbb{R}$ is a scaling factor such that $|\nu|(\mathbb{R}) = 1$ and $\nu(\{x_0\})$ has the same sign as ε . For each $k = -q+1, \dots, K$, using the divided difference of $x \mapsto x^{k+q-1}$, we have

$$\int x^k \nu(dx) = c_0 \cdot \sum_{i=0}^{q+K} x_i^{k+q-1} \prod_{j \neq i} \frac{1}{x_i - x_j} = 0.$$

Moreover, for each $i = 0, 1, \dots, q+K$, the difference $f(x_i) - P(x_i)$ has the same sign as $\nu(\{x_i\})$, for the signs of $\nu(\{x_i\})$ are also alternating. Therefore,

$$\begin{aligned}
\int f(x) \nu(dx) &= \int (f(x) - P(x)) \nu(dx) + \int P(x) \nu(dx) \\
&= \int |f(x) - P(x)| |\nu|(dx) + 0 \\
&= E_{q-1,K}(f; I) \cdot |\nu|(\mathbb{R}) = E_{q-1,K}(f; I).
\end{aligned}$$

Now the proof is completed by considering the Jordan decomposition of $\nu = \nu_+ - \nu_-$ and choosing $\nu_0 = 2\nu_+, \nu_1 = 2\nu_-$.

6.5 Proof of Lemma 5.6

We need some notions and results from approximation theory first. For functions defined on $[0, 1]$, define the r -th order Ditzian–Totik modulus of smoothness by [DT87]

$$\omega_\varphi^r(f, t)_\infty \triangleq \sup_{0 < h \leq t} \|\Delta_{h\varphi(x)}^r f(x)\|_\infty,$$

where $\varphi(x) \triangleq \sqrt{x(1-x)}$. This quantity is related to the polynomial approximation error via the following lemma.

Lemma 6.15. [DT87] *For an integer $u > 0$ and $n > u$, there exists some constant M_u which depends on u but not on $t \in (0, 1)$ nor f , such that*

$$\begin{aligned} E_{0,n}(f; [0, 1]) &\leq M_u \omega_\varphi^u(f, \frac{1}{n})_\infty, \\ \frac{M_u}{n^u} \sum_{k=0}^n (k+1)^{u-1} E_{0,k}(f; [0, 1]) &\geq \omega_\varphi^u(f, \frac{1}{n})_\infty. \end{aligned}$$

Take $u = 1$, the second inequality together with the monotonicity of $E_{0,n}(f; [0, 1])$ in n yields

$$\begin{aligned} E_{0,n}(f; [0, 1]) &\geq \frac{1}{Dn} \sum_{k=n}^{Dn} E_{0,k}(f; [0, 1]) \\ &\geq \frac{\omega_\varphi^1(f, \frac{1}{Dn})_\infty}{M_1} - \frac{1}{Dn} \sum_{k=0}^{n-1} E_{0,k}(f; [0, 1]) \\ &\geq \frac{\omega_\varphi^1(f, \frac{1}{Dn})_\infty}{M_1} - \frac{E_{0,0}(f; [0, 1])}{D}, \end{aligned}$$

where we choose $D = \lceil 1/\sqrt{c} \rceil$, with the constant $c \in (0, 1/2)$ is specified later. Since

$$E_{q-1,n}(f_q; I) = \inf_{a_1, \dots, a_{q-1}} E_{0,n} \left(x^{-q+\frac{r}{2}} + \sum_{k=1}^{q-1} \frac{a_k}{x^k}; I \right),$$

it suffices to obtain a lower bound independent of a_1, \dots, a_{q-1} for the polynomial approximation error $E_{0,n}(x^{-q+r/2} + \sum_{k=1}^{q-1} a_k x^{-k}; [cn^{-2}, 1])$. Define $g(x) = x^{-q+r/2} + \sum_{k=1}^{q-1} a_k x^{-k}$, and let $\tilde{g}(x) = g(cn^{-2} + (1 - cn^{-2})x)$ be the scaled version defined on $[0, 1]$. We distinguish into two cases.

First we consider the case where $E_{0,0}(\tilde{g}; [0, 1]) \leq C_1 n^{2q-r}$ for some fixed constant $C_1 = 2c^{-q+r/2}$. By the definition of $\omega_\varphi^1(f, t)_\infty$ and the choice $D = \lceil 1/\sqrt{c} \rceil$, there exists universal constants $0 < A < B$ (which are independent of everything) such that

$$\begin{aligned} \omega_\varphi^1(\tilde{g}, \frac{1}{Dn})_\infty &\geq \sup_{t \in [A, B]} \left| g\left(\frac{t+1}{(Dn)^2}\right) - g\left(\frac{t}{(Dn)^2}\right) \right| \\ &= (Dn)^{2q-r} \sup_{t \in [A, B]} \left| h_{q-\frac{r}{2}}(t) + \sum_{k=1}^{q-1} (Dn)^{2k-2q+r} a_k h_k(t) \right| \\ &\geq (Dn)^{2q-r} \inf_{b_1, \dots, b_{q-1}} \sup_{t \in [A, B]} \left| h_{q-\frac{r}{2}}(t) + \sum_{k=1}^{q-1} b_k h_k(t) \right|, \end{aligned}$$

where

$$h_k(t) \triangleq (t+1)^{-k} - t^{-k}.$$

Since r is not even, it is straightforward to verify that the functions $h_1, \dots, h_{q-1}, h_{q-\frac{r}{2}}$ are linearly independent in the interval $[A, B]$, we conclude that

$$\omega_\varphi^1(\tilde{g}, \frac{1}{Dn})_\infty \geq C_2(Dn)^{2q-r},$$

where the constant $C_2 > 0$ only depends on r, q, A, B but not on a_1, \dots, a_{q-1} or c, D . Hence, in this case we have

$$E_{0,n}(g; [cn^{-2}, 1]) \geq \frac{C_2(Dn)^{2q-r}}{M_1} - \frac{C_1 n^{2q-r}}{D}, \quad (6.17)$$

where neither of the constants depends on a_1, \dots, a_{q-1} .

Second we consider the case where $E_{0,0}(\tilde{g}; [0, 1]) > C_1 n^{2q-r}$. Since $2q - r > 0$, we have

$$\begin{aligned} E_{0,0}(\tilde{g}; [0, 1]) &\leq \max_{x \in [cn^{-2}, 1]} \left| x^{-q+r/2} + \sum_{k=1}^{q-1} a_k x^{-k} \right| \\ &\leq c^{-q+r/2} n^{2q-r} + \sum_{k=1}^{q-1} |a_k| c^{-k} n^{2k} \\ &\leq \frac{1}{2} E_{0,0}(\tilde{g}; [0, 1]) + \sum_{k=1}^{q-1} |a_k| c^{-k} n^{2k}, \end{aligned}$$

and thus there exists some $j \in \{1, \dots, q-1\}$ such that

$$|a_j| c^{-j} n^{2j} \geq C_3 E_{0,0}(\tilde{g}; [0, 1]),$$

where $C_3 = 1/(2q)$ is a numerical constant. Defining the interval $[A, B]$ as in the first case, we have

$$\begin{aligned} \omega_\varphi^1(\tilde{g}, \frac{1}{Dn})_\infty &\geq \sup_{t \in [A, B]} \left| g\left(\frac{t+1}{(Dn)^2}\right) - g\left(\frac{t}{(Dn)^2}\right) \right| \\ &= |a_j| (Dn)^{2j} \sup_{t \in [A, B]} \left| \frac{(Dn)^{2q-r-2j}}{a_j} h_{q-\frac{r}{2}}(t) + \sum_{k=1}^{q-1} (Dn)^{2k-2j} \frac{a_k}{a_j} h_k(t) \right| \\ &\geq |a_j| (Dn)^{2j} \inf_{b_1, \dots, b_{q-1}} \sup_{t \in [A, B]} |h_j(t) + b_j h_{q-\frac{r}{2}}(t) + \sum_{k \neq j} b_k h_k(t)| \\ &\geq C_4 |a_j| (Dn)^{2j} \\ &\geq C_3 C_4 \cdot E_{0,0}(\tilde{g}; [0, 1]), \end{aligned}$$

where the numerical constant $C_4 > 0$ (which only depends on r, q, A, B but not on a_1, \dots, a_{q-1} or c, D) again follows from the linear independence of the functions $h_{q-\frac{r}{2}}, h_1, \dots, h_{q-1}$ in $[A, B]$. Now in this case we have

$$E_{0,n}(g; [cn^{-2}, 1]) \geq \left(\frac{C_3 C_4}{M_1} - \frac{1}{D} \right) \cdot E_{0,0}(\tilde{g}; [0, 1])$$

$$\geq C_1 \left(\frac{C_3 C_4}{M_1} - \frac{1}{D} \right) \cdot n^{2q-r}, \quad (6.18)$$

where again none of the constants depends on a_1, \dots, a_{q-1} .

Now by (6.17) and (6.18), we plug in the definition $D = \lceil 1/\sqrt{c} \rceil$ and $C_1 = 2c^{-q+r/2}n^{2q-r}$, and use that the constants C_2, C_3, C_4 do not depend on a_1, \dots, a_{q-1} or c, D , to conclude that for $c > 0$ small enough, we have

$$\begin{aligned} E_{q,n}(x^{-q+r/2}; [cn^{-2}, 1]) &= \inf_{a_1, \dots, a_{q-1}} E_{0,n}(x^{-q+r/2} + \sum_{k=1}^{q-1} \frac{a_k}{x^k}; [cn^{-2}, 1]) \\ &\geq c' n^{2q-r}, \end{aligned}$$

which is the desired result. \square

7 Acknowledgement

We would like to thank Tsachy Weissman for the tremendous support and very helpful discussions. We are also grateful to an anonymous referee for detailed and helpful comments on improving this paper.

References

- [Ber12] Serge Bernstein. *Sur l'ordre de la meilleure approximation des fonctions continues par des polynomes de degré donné*, volume 4. Hayez, imprimeur des académies royales, 1912.
- [BI79] Oleg Vladimirovich Besov and Valentin Petrovich Ilyin. Integral representations of functions and imbedding theorems, v. 2. 1979.
- [BKB⁺93] Peter J Bickel, Chris AJ Klaassen, Peter J Bickel, Y Ritov, J Klaassen, Jon A Wellner, and YA'Acov Ritov. *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press Baltimore, 1993.
- [BLM13] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford University Press, 2013.
- [BM95] Lucien Birgé and Pascal Massart. Estimation of integral functionals of a density. *The Annals of Statistics*, pages 11–29, 1995.
- [BR88] Peter J Bickel and Ya'acov Ritov. Estimating integrated squared density derivatives: sharp best order of convergence estimates. *Sankhyā: The Indian Journal of Statistics, Series A*, pages 381–393, 1988.
- [Car13] Alexandra Carpentier. Honest and adaptive confidence sets in L_p . *Electronic Journal of Statistics*, 7:2875–2923, 2013.
- [CL03] T Tony Cai and Mark G Low. A note on nonparametric estimation of linear functionals. *Annals of statistics*, pages 1140–1153, 2003.
- [CL04] T Tony Cai and Mark G Low. Minimax estimation of linear functionals over nonconvex parameter spaces. *Annals of statistics*, pages 552–576, 2004.

- [CL05] T Tony Cai and Mark G Low. Nonquadratic estimators of a quadratic functional. *The Annals of Statistics*, pages 2930–2956, 2005.
- [CL11] T Tony Cai and Mark G Low. Testing composite hypotheses, hermite polynomials and optimal estimation of a nonsmooth functional. *The Annals of Statistics*, 39(2):1012–1041, 2011.
- [DL93] Ronald A DeVore and George G Lorentz. *Constructive approximation*, volume 303. Springer, 1993.
- [DLM90] David L Donoho, Richard C Liu, and Brenda MacGibbon. Minimax risk over hyperrectangles, and implications. *The Annals of Statistics*, pages 1416–1437, 1990.
- [DN90] David L Donoho and Michael Nussbaum. Minimax quadratic estimation of a quadratic functional. *Journal of Complexity*, 6(3):290–323, 1990.
- [DT87] Zeev Ditzian and Vilmos Totik. *Moduli of smoothness*. Springer, 1987.
- [Fan91] Jianqing Fan. On the estimation of quadratic functionals. *The Annals of Statistics*, pages 1273–1294, 1991.
- [GN15] Evarist Giné and Richard Nickl. *Mathematical foundations of infinite-dimensional statistical models*, volume 40. Cambridge University Press, 2015.
- [HJW16] Yanjun Han, Jiantao Jiao, and Tsachy Weissman. Minimax rate-optimal estimation of divergences between discrete distributions. *arXiv preprint arXiv:1605.09124*, 2016.
- [HJWW17] Yanjun Han, Jiantao Jiao, Tsachy Weissman, and Yihong Wu. Optimal rates of entropy estimation over Lipschitz balls. *arXiv preprint arXiv:1711.02141*, 2017.
- [HKPT12] Wolfgang Härdle, Gerard Kerkycharian, Dominique Picard, and Alexander Tsybakov. *Wavelets, approximation, and statistical applications*, volume 129. Springer Science & Business Media, 2012.
- [HM87] Peter Hall and James Stephen Marron. Estimation of integrated squared density derivatives. *Statistics & Probability Letters*, 6(2):109–115, 1987.
- [IK81] I.A. Ibragimov and R.Z. Khas’minskii. Estimates of the signal, its derivatives and point of maximum for Gaussian distributions. *Theory of Probability and Its Applications*, 25(4):703–720, 1981.
- [Ing87] Yu I Ingster. Minimax testing of nonparametric hypotheses on a distribution density in the L_p metrics. *Theory of Probability & Its Applications*, 31(2):333–337, 1987.
- [IS12] Yuri Ingster and Irina A Suslina. *Nonparametric goodness-of-fit testing under Gaussian models*, volume 169. Springer Science & Business Media, 2012.
- [JHW16] Jiantao Jiao, Yanjun Han, and Tsachy Weissman. Minimax estimation of the L_1 distance. In *2016 IEEE International Symposium on Information Theory (ISIT)*, pages 750–754. IEEE, 2016.
- [JVHW15] Jiantao Jiao, Kartik Venkat, Yanjun Han, and Tsachy Weissman. Minimax estimation of functionals of discrete distributions. *IEEE Transactions on Information Theory*, 61(5):2835–2885, 2015.

- [JVHW17] Jiantao Jiao, Kartik Venkat, Yanjun Han, and Tsachy Weissman. Maximum likelihood estimation of functionals of discrete distributions. *IEEE Transactions on Information Theory*, 63(10):6774–6798, 2017.
- [Kor91] A.P. Korostelevi. On estimation accuracy for nonsmooth functionals of regression. *Theory of Probability and Its Applications*, 35(4):784–787, 1991.
- [KP96] Gérard Kerkyacharian and Dominique Picard. Estimating nonquadratic functionals of a density using haar wavelets. *The Annals of Statistics*, 24(2):485–507, 1996.
- [KT12] Aleksandr Petrovich Korostelev and Alexandre B Tsybakov. *Minimax theory of image reconstruction*, volume 82. Springer Science & Business Media, 2012.
- [Lau96] Béatrice Laurent. Efficient estimation of integral functionals of a density. *The Annals of Statistics*, 24(2):659–681, 1996.
- [Lep91] OV Lepski. On a problem of adaptive estimation in Gaussian white noise. *Theory of Probability & Its Applications*, 35(3):454–466, 1991.
- [Lep92a] Oleg V Lepski. On problems of adaptive estimation in white Gaussian noise. *Topics in nonparametric estimation*, 12:87–106, 1992.
- [Lep92b] OV Lepski. Asymptotically minimax adaptive estimation. I: Upper bounds. Optimally adaptive estimates. *Theory of Probability & Its Applications*, 36(4):682–697, 1992.
- [Lep93] OV Lepski. Asymptotically minimax adaptive estimation. II. Schemes without optimal adaptation: Adaptive estimators. *Theory of Probability & Its Applications*, 37(3):433–448, 1993.
- [LNS99] Oleg Lepski, Arkady Nemirovski, and Vladimir Spokoiny. On estimation of the L_r norm of a regression function. *Probability theory and related fields*, 113(2):221–253, 1999.
- [MNR17] Rajarshi Mukherjee, Whitney K Newey, and James M Robins. Semiparametric efficient empirical higher order influence function estimators. *arXiv preprint arXiv:1705.07577*, 2017.
- [Nem00] Arkadi Nemirovski. Topics in non-parametric. *Ecole d’Eté de Probabilités de Saint-Flour*, 28:85, 2000.
- [Nik12] Sergei Mihailovic Nikol’skii. *Approximation of functions of several variables and embedding theorems*, volume 205. Springer Science & Business Media, 2012.
- [QR07] MA Qazi and QI Rahman. Some coefficient estimates for polynomials on the unit interval. *Serdica Math. J.*, 33:449–474, 2007.
- [RLM⁺16] James Robins, Lingling Li, Rajarshi Mukherjee, Eric Tchetgen Tchetgen, and Aad van der Vaart. Higher order estimating equations for high-dimensional models. *The Annals of Statistics (To Appear)*, 2016.
- [RLTvdV08] James Robins, Lingling Li, Eric Tchetgen, and Aad van der Vaart. Higher order influence functions and minimax estimation of nonlinear functionals. In *Probability and Statistics: Essays in Honor of David A. Freedman*, pages 335–421. Institute of Mathematical Statistics, 2008.

- [Spo98] Vladimir Spokoiny. Adaptive and spatially hypothesis testing of a nonparametric hypothesis. *Math. Methods Statist.*, 7:245–273, 1998.
- [TLRvdV08] Eric Tchetgen, Lingling Li, James Robins, and Aad van der Vaart. Minimax estimation of the integral of a power of a density. *Statistics & Probability Letters*, 78(18):3307–3311, 2008.
- [Tsy08] A. Tsybakov. *Introduction to Nonparametric Estimation*. Springer-Verlag, 2008.
- [VdV00] Aad W Van der Vaart. *Asymptotic statistics*, volume 3. Cambridge university press, 2000.
- [VK87] Richard S Varga and A Dzh Karpenter. On a conjecture of S. Bernstein in approximation theory. *Sbornik: Mathematics*, 57(2):547–560, 1987.
- [VV11] Gregory Valiant and Paul Valiant. The power of linear estimators. In *Foundations of Computer Science (FOCS), 2011 IEEE 52nd Annual Symposium on*, pages 403–412. IEEE, 2011.
- [WY16] Yihong Wu and Pengkun Yang. Minimax rates of entropy estimation on large alphabets via best polynomial approximation. *IEEE Transactions on Information Theory*, 62(6):3702–3720, 2016.