New series representations of π , π^3 and π^5 in terms of Euler numbers and π^2 , π^4 and π^6 in terms of Bernoulli numbers

H.C.Gupta Physics Department Indian Institute of Technology New Delhi-110016, INDIA <u>hcgupta@physics.iitd.ac.in</u>

ABSTRACT

New series representations for odd powers of π i.e. π , π^3 and π^5 in terms of Euler numbers and even powers of π i.e. π^2 , π^4 and π^6 in terms of Bernoulli numbers have been obtained empirically.

Keywords: Powers of π , Series representation, Bernoulli numbers, Euler numbers

Mathematics Subject Classification 2010: 11B68, 11B50

1. Introduction

The representation of the mathematical constants π and its powers by series, products and integrals have attracted the attention of researchers for a long time. The classic result for π , π^2 , π^3 , π^4 , π^5 , π^6 in terms of a series of powers of integers was presented by Leibnitz and Euler as

$$\pi = 4 \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)} \tag{1}$$

$$\pi^2 = 6 \ \sum_{k=1}^{\infty} \frac{1}{k^2} \tag{2}$$

$$\pi^{3} = 32 \sum_{k=0}^{\infty} \frac{(-1)^{k}}{(2k+1)^{3}} , \qquad \pi^{4} = 90 \sum_{k=1}^{\infty} \frac{1}{k^{4}}$$
(3)

$$\pi^{5} = \frac{1536}{5} \sum_{k=0}^{\infty} \frac{(-1)^{k}}{(2k+1)^{5}} , \qquad \pi^{6} = 945 \sum_{k=1}^{\infty} \frac{1}{k^{6}}$$
(4)

Borwein[5] had described very elegantly the history of π along with its selected useful formulas in terms of series representation. Alzer and Koumandos[2] presented a series representation for π as

$$\pi = 4 \sum_{k=0}^{\infty} \frac{1}{(1+\mu)^{k+1}} \sum_{m=0}^{k} \binom{k}{m} \frac{(-1)^m \ \mu^{k-m}}{(2m+1)}$$
(5)

where they introduced a positive parameter μ , on the lines of modifying the Amore's technique[3] to obtain one-parameter family of series representation of mathematical constants. Further, in case of mathematical constants involving π , series representations were given by Alzer etal.[1] in terms of digamma function. Recently, Blagouchine[4] obtained new series expansions for generalized Euler constants containing polynomials in π^{-2} with rational coefficients.

Alzer etal.[1] gave series representation for π^2 as

$$\pi^2 = 4 \, \sum_{k=1}^{\infty} \frac{\mu_k h_k}{k} \tag{6}$$

where μ_n denotes the normalized binomial mid-coefficient defined by

$$\mu_k = \frac{1.3.5.\dots(2k-1)}{2.4.6.\dots(2k-1)} \tag{7}$$

and
$$h_k = \sum_{n=1}^k \frac{1}{(2n-1)}$$
 (8)

Kolbig [6] gave an interesting relation as

$$\pi^2 = 2 \sum_{k=1}^{\infty} \frac{\sigma_k}{k} \tag{9}$$

where

$$\sigma_{n} = p_{n} \sum_{k=1}^{n} \frac{1}{(4k-1)} + q_{n} \sum_{k=1}^{n} \frac{1}{(4k-3)}$$
(10)

$$p_n = \prod_{k=1}^n \frac{(4k-1)}{4k}$$
, $q_n = \prod_{k=1}^n \frac{(4k-3)}{4k}$ (11)

Yet another series representation for π^2 were given by Alzer etal.[1] as

$$\pi^2 = 3 \sum_{k=1}^{\infty} \frac{\mu_k H_k}{k}$$
(12)

where

$$H_n = \sum_{k=1}^n \frac{1}{k}$$
(13)

In this paper, the series representations for π , π^3 and π^5 are presented where the series involves the application of Euler numbers E_{2k} , defined as

$$\sum_{m=1}^{\infty} \frac{(-1)^{m+1}}{(2m-1)^{2k+1}} = \frac{\pi^{2k+1}}{2^{2k+2} (2k)!} |E_{2k}|$$
(14)

with $E_0 = 1$ giving $E_2 = -1$, $E_4 = 5$, $E_6 = -61$, $E_8 = 1385$, $E_{10} = -50521$, $E_{12} = 2702765$, and the series $\sum_{j=0}^{k-1} \left[-\frac{1}{(2m-1)^2 \pi^2} \right]^j \frac{1}{(2k-2j-1)!}$ for integers j, $k \ge 0$.

Further, the series representations for π^2 , π^4 and π^6 are presented where the series involves the application of Bernoulli numbers B_{2k}, defined as

$$\sum_{m=1}^{\infty} \frac{1}{(m)^{2k}} = (-1)^{k-1} \frac{(2\pi)^{2k}}{2(2k)!} \quad B_{2k}, k = 1, 2, \dots$$
(15)

with $B_0 = 1$ giving $B_1 = -\frac{1}{2}$, $B_2 = \frac{1}{6}$, $B_4 = -\frac{1}{30}$, $B_6 = \frac{1}{42}$, $B_8 = -\frac{1}{30}$, $B_{10} = \frac{5}{66}$, $B_{12} = -\frac{691}{2730}$, and the series $\sum_{j=0}^{k-1} \left[-\frac{1}{(m)^2 \pi^2} \right]^j \frac{1}{(2k-2j-1)!}$ for integers j, $k \ge 0$.

2. Series representations for odd powers π , π^3 and π^5

2a. The series for π

The series for π is presented as

$$\pi = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^{2k+2} \quad (2k+1)! \quad \sum_{j=0}^{k} \left[-\frac{1}{(2n-1)^2 \pi^2} \right]^j \frac{1}{(2k-2j+1)!}$$
(16)

where j, k are integers and ≥ 0 .

Verification of the series for some k values

 $\begin{aligned} \mathbf{k} &= 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^2 \quad (1)! \quad \left[-\frac{1}{(2n-1)^2 \pi^2} \right]^0 \frac{1}{(1)!} = 4. \frac{\pi}{4} = \pi \\ \mathbf{k} &= 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^4 \quad (3)! \quad \left[\frac{1}{3!} - \frac{1}{(2n-1)^2 \pi^2} \right] = 16 \left[\frac{\pi}{4} - \frac{6}{\pi^2} \frac{\pi^3}{32} \right] = \pi \\ \mathbf{k} &= 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^6 \quad (5)! \quad \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{(2n-1)^4 \pi^4} \right] = 64 \left[\frac{\pi}{4} - \frac{20}{\pi^2} \frac{\pi^3}{32} + \frac{120}{\pi^4} \frac{5\pi^5}{1536} \right] = \pi \\ \mathbf{k} &= 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^8 \quad (7)! \quad \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{3!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{(2n-1)^6 \pi^6} \right] = \\ 256 \left[\frac{\pi}{4} - \frac{42}{\pi^2} \frac{\pi^3}{32} + \frac{840}{\pi^4} \frac{5\pi^5}{1536} - \frac{5040}{\pi^6} \frac{61\pi^7}{184320} \right] = \pi \\ \mathbf{k} &= 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)} 2^{10} \quad (9)! \quad \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{5!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{3!} \frac{1}{(2n-1)^6 \pi^6} + \frac{1}{(2n-1)^8 \pi^8} \right] = \\ 1024 \left[\frac{\pi}{4} - \frac{72}{\pi^2} \frac{\pi^3}{32} + \frac{3024}{\pi^4} \frac{5\pi^5}{1536} - \frac{60480}{\pi^6} \frac{61\pi^7}{184320} + \frac{362880}{\pi^8} \frac{1385\pi^9}{41287680} \right] = \pi \end{aligned}$

On similar lines, the series representation (16) can be proved for any integer k value.

<u>2b. The series for \pi^3</u>

The series for π^3 is presented as

$$\pi^{3} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^{3} (2^{2k+2}-1)} 2^{2k+4} \quad (2k+3)! \quad \sum_{j=0}^{k} \left[-\frac{1}{(2n-1)^{2} \pi^{2}} \right]^{j} \frac{1}{(2k-2j+1)!}$$
(17)

where j, k are integers and ≥ 0 .

Verification of the series for some k values

$$\begin{aligned} \mathbf{k} &= 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^4 \frac{1}{(2^2-1)} \quad (3)! \quad \left[-\frac{1}{(2n-1)^2 \pi^2} \right]^0 \frac{1}{(1!)} = 32. \frac{\pi^3}{32} = \pi^3 \\ \mathbf{k} &= 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^6 \frac{1}{(2^4-1)} \quad (5)! \quad \left[\frac{1}{3!} - \frac{1}{(2n-1)^2 \pi^2} \right] = 512 \left[\frac{1}{6} \frac{\pi^3}{32} - \frac{1}{\pi^2} \frac{5\pi^5}{1536} \right] = \pi^3 \\ \mathbf{k} &= 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^8 \frac{1}{(2^6-1)} \quad (7)! \quad \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{(2n-1)^4 \pi^4} \right] = \\ 20480 \left[\frac{1}{120} \frac{\pi^3}{32} - \frac{1}{6\pi^2} \frac{5\pi^5}{1536} + \frac{1}{\pi^4} \frac{61\pi^7}{184320} \right] = \pi^3 \\ \mathbf{k} &= 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^{10} \frac{1}{(2^8-1)} \quad (9)! \quad \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{3!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{(2n-1)^6 \pi^6} \right] = \\ \frac{371589120}{255} \left[\frac{1}{5040} \frac{\pi^3}{32} - \frac{1}{120\pi^2} \frac{5\pi^5}{1536} + \frac{1}{6\pi^4} \frac{61\pi^7}{184320} - \frac{1}{\pi^6} \frac{1385\pi^9}{41287680} \right] = \pi^3 \\ \mathbf{k} &= 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^{12} \frac{1}{(2^{10}-1)} \quad (11)! \quad \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{5!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{3!} \frac{1}{(2n-1)^6 \pi^6} + \frac{1}{(2n-1)^6 \pi^6} \right] \\ \mathbf{k} &= 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^3} 2^{12} \frac{1}{(2^{10}-1)} \quad (11)! \quad \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{5!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{3!} \frac{1}{(2n-1)^6 \pi^6} + \frac{1}{(2n-1)^6 \pi^6} + \frac{1}{(2n-1)^8 \pi^8} \right] \\ = \frac{4954521600}{31} \left[\frac{1}{362880} \frac{\pi^3}{32} - \frac{1}{5040\pi^2} \frac{5\pi^5}{1536} + \frac{1}{120\pi^4} \frac{61\pi^7}{184320} - \frac{1}{6\pi^6} \frac{61\pi^7}{184320} - \frac{1}{6\pi^6} \frac{1}{41287680} + \frac{1}{6\pi^6} \frac{1385\pi^9}{41287680} + \frac{1}{\pi^8} \frac{50521\pi^{11}}{14863564800} \right] = \pi^3 \end{aligned}$$

On similar lines, the series representation (17) can be proved for any integer k value.

<u>2c. The series for \pi^5</u>

The series for π^5 is presented as

$$\pi^{5} = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^{5} \{ [2^{2k+2}(2k^{2}+9k+6)]+1 \}} 2^{2k+6} \quad (2k+5)! \sum_{j=0}^{k} \left[-\frac{1}{(2n-1)^{2} \pi^{2}} \right]^{j} \frac{1}{(2k-2j+1)!}$$
(18)

where j, k are integers and ≥ 0 .

Verification of the series for some k values

$$\begin{aligned} \mathbf{k} &= 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^5} 2^6 \frac{1}{\{[4(6)]+1\}} \quad (5)! \quad \left[-\frac{1}{(2n-1)^2 \pi^2}\right]^0 \frac{1}{(1)!} = \frac{1536}{5} \cdot \frac{5\pi^5}{1536} = \pi^5 \\ \mathbf{k} &= 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^5} 2^8 \frac{1}{\{[16(17)]+1\}} \quad (7)! \quad \left[\frac{1}{3!} - \frac{1}{(2n-1)^2 \pi^2}\right] = \frac{256(5040)}{273} \left[\frac{1}{6} \frac{5\pi^5}{1536} - \frac{1}{\pi^2} \frac{61\pi^7}{184320}\right] = \pi^5 \\ \mathbf{k} &= 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^5} 2^{10} \frac{1}{\{[64(32)]+1\}} \quad (9)! \quad \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{(2n-1)^4 \pi^4}\right] = \\ \frac{1024(362880)}{2049} \left[\frac{1}{120} \frac{5\pi^5}{1536} - \frac{1}{6\pi^2} \frac{61\pi^7}{184320} + \frac{1}{\pi^4} \frac{1385\pi^9}{41287680}\right] = \pi^5 \\ \mathbf{k} &= 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^5} 2^{12} \frac{1}{\{[256(51)]+1\}} \quad (11)! \quad \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{3!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{(2n-1)^6 \pi^6}\right] = \\ \frac{4096(39916800)}{13057} \left[\frac{1}{5040} \frac{5\pi^5}{1536} - \frac{1}{120\pi^2} \frac{61\pi^7}{184320} + \frac{1}{6\pi^4} \frac{1385\pi^9}{41287680} - \frac{1}{\pi^6} \frac{50521\pi^{11}}{14863564800}\right] = \pi^5 \\ \mathbf{k} &= 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{(2n-1)^5} 2^{14} \frac{1}{\{[1024(74)]+1\}} \quad (13)! \quad \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(2n-1)^2 \pi^2} + \frac{1}{5!} \frac{1}{(2n-1)^4 \pi^4} - \frac{1}{3!} \frac{1}{(2n-1)^6 \pi^6} + \frac{1}{(2n-1)^6 \pi^6}\right] = \\ \frac{1}{\pi^6} \frac{1}{16384(6227020800)} = \pi^5 \end{aligned}$$

On similar lines, the series representation (18) can be proved for any integer k value.

3. Series representations for odd powers π^2 , π^4 and π^6

<u>3a. The series for \pi^2</u>

The series for π^2 is presented as

$$\pi^{2} = \sum_{n=1}^{\infty} \frac{1}{n^{2}} 2 \left(2k+3 \right) \quad \left(2k+1 \right)! \sum_{j=0}^{k} \left[-\frac{1}{(n)^{2} \pi^{2}} \right]^{j} \frac{1}{(2k-2j+1)!}$$
(19)

.

where j, k are integers and ≥ 0 .

Verification of the series for some k values

$$\begin{aligned} \mathbf{k} &= 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^2} 6 \quad (1)! \quad \left[-\frac{1}{(n)^2 \pi^2} \right]^0 \frac{1}{(1)!} = 6. \frac{\pi^2}{6} = \pi^2 \\ \mathbf{k} &= 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^2} 10 \quad (3)! \quad \left[\frac{1}{3!} - \frac{1}{(n)^2 \pi^2} \right] = 60 \left[\frac{1}{6} \frac{\pi^2}{6} - \frac{1}{\pi^2} \frac{\pi^4}{90} \right] = \pi^2 \\ \mathbf{k} &= 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^2} 14 \quad (5)! \quad \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(n)^2 \pi^2} + \frac{1}{(n)^4 \pi^4} \right] = 1680 \left[\frac{1}{120} \frac{\pi^2}{6} - \frac{1}{6} \frac{1}{\pi^2} \frac{\pi^4}{90} + \frac{1}{\pi^4} \frac{\pi^6}{945} \right] = \pi^2 \\ \mathbf{k} &= 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^2} 18 \quad (7)! \quad \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(n)^2 \pi^2} + \frac{1}{3!} \frac{1}{(n)^4 \pi^4} - \frac{1}{(n)^6 \pi^6} \right] = \\ 90720 \left[\frac{1}{5040} \frac{\pi^2}{6} - \frac{1}{120} \frac{1}{\pi^2} \frac{\pi^4}{90} + \frac{1}{6} \frac{1}{\pi^4} \frac{\pi^6}{945} - \frac{1}{\pi^6} \frac{\pi^6}{9450} \right] = \pi^2 \\ \mathbf{k} &= 4 \quad \text{gives series as } \sum_{n=1}^{\infty} \frac{1}{n^2} 22 \quad (9)! \quad \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(n)^2 \pi^2} + \frac{1}{5!} \frac{1}{(n)^4 \pi^4} - \frac{1}{3!} \frac{1}{(n)^6 \pi^6} + \frac{1}{(n)^8 \pi^8} \right] = \\ 7983360 \left[\frac{1}{362880} \frac{\pi^2}{6} - \frac{1}{5040} \frac{1}{\pi^2} \frac{\pi^4}{90} + \frac{1}{120} \frac{1}{\pi^4} \frac{\pi^6}{945} - \frac{1}{6} \frac{1}{\pi^6} \frac{\pi^6}{9450} + \frac{1}{\pi^6} \frac{\pi^8}{93555} \right] = \pi^2 \end{aligned}$$

On similar lines, the series representation (19) can be proved for any integer k value.

<u>3b. The series for \pi^4</u>

The series for π^4 is presented as

$$\pi^{4} = \sum_{n=1}^{\infty} \frac{1}{n^{4}} 6(2k+5) (2k+3) (2k+1)! \sum_{j=0}^{k} \left[-\frac{1}{(n)^{2} \pi^{2}} \right]^{j} \frac{1}{(2k-2j+1)!}$$
(20)

where j, k are integers and ≥ 0 .

Verification of the series for some k values

 $k = 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^4} 6 (5)(3) (1)! \left[-\frac{1}{(n)^2 \pi^2} \right]^0 \frac{1}{(1)!} = 90. \frac{\pi^4}{90} = \pi^4$ $k = 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^4} 6 (7)(5) (3)! \left[\frac{1}{3!} - \frac{1}{(n)^2 \pi^2} \right] = 1260 \left[\frac{1}{6} \frac{\pi^4}{90} - \frac{1}{\pi^2} \frac{\pi^6}{945} \right] = \pi^4$

$$k = 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^4} 6(9)(7) \quad (5)! \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(n)^2 \pi^2} + \frac{1}{(n)^4 \pi^4} \right] =$$

$$45360 \left[\frac{1}{120} \frac{\pi^4}{90} - \frac{1}{6} \frac{1}{\pi^2} \frac{\pi^6}{945} + \frac{1}{\pi^4} \frac{\pi^8}{9450} \right] = \pi^4$$

$$k = 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^4} 6(11)(9) \quad (7)! \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(n)^2 \pi^2} + \frac{1}{3!} \frac{1}{(n)^4 \pi^4} - \frac{1}{(n)^6 \pi^6} \right] =$$

$$2993760 \left[\frac{1}{5040} \frac{\pi^4}{90} - \frac{1}{120} \frac{1}{\pi^2} \frac{\pi^6}{945} + \frac{1}{6} \frac{1}{\pi^4} \frac{\pi^8}{9450} - \frac{1}{\pi^6} \frac{\pi^{10}}{93555} \right] = \pi^4$$

$$k = 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^4} 6(13)(11) \quad (9)! \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(n)^2 \pi^2} + \frac{1}{5!} \frac{1}{(n)^4 \pi^4} - \frac{1}{3!} \frac{1}{(n)^6 \pi^6} + \frac{1}{(n)^8 \pi^8} \right] =$$

$$311351040 \left[\frac{1}{362880} \frac{\pi^4}{90} - \frac{1}{5040} \frac{1}{\pi^2} \frac{\pi^6}{945} + \frac{1}{120} \frac{1}{\pi^4} \frac{\pi^8}{9450} - \frac{1}{6} \frac{1}{\pi^6} \frac{\pi^{10}}{93555} + \frac{1}{\pi^8} \frac{691\pi^{12}}{638512875} \right] = \pi^4$$

On similar lines, the series representation (20) can be proved for any integer k value.

<u>3c. The series for \pi^6</u>

The series for π^6 is presented as

$$\pi^{6} = \sum_{n=1}^{\infty} \frac{1}{n^{6}} \frac{45(2k+7)(2k+5)(2k+3)}{(k+5)} \quad (2k+1)! \quad \sum_{j=0}^{k} \left[-\frac{1}{(n)^{2} \pi^{2}} \right]^{j} \frac{1}{(2k-2j+1)!}$$
(21)

where j, k are integers and ≥ 0 .

Verification of the series for some k values

$$k = 0 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^6} 45 (7)(3) (1)! \left[-\frac{1}{(n)^2 \pi^2} \right]^0 \frac{1}{(1)!} = 945. \frac{\pi^6}{945} = \pi^6$$

$$k = 1 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^6} 45 (9)(7)(5) \frac{1}{6} (3)! \left[\frac{1}{3!} - \frac{1}{(n)^2 \pi^2} \right] = 14175 \left[\frac{1}{6} \frac{\pi^6}{945} - \frac{1}{\pi^2} \frac{\pi^8}{9450} \right] = \pi^6$$

$$k = 2 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^6} 45(11)(9) (5)! \left[\frac{1}{5!} - \frac{1}{3!} \frac{1}{(n)^2 \pi^2} + \frac{1}{(n)^4 \pi^4} \right] = 534600 \left[\frac{1}{120} \frac{\pi^6}{945} - \frac{1}{6} \frac{1}{\pi^2} \frac{\pi^8}{9450} + \frac{1}{\pi^4} \frac{\pi^{10}}{93555} \right] = \pi^6$$

$$k = 3 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^6} 45 (13)(11)(9) \frac{1}{8} (7)! \left[\frac{1}{7!} - \frac{1}{5!} \frac{1}{(n)^2 \pi^2} + \frac{1}{3!} \frac{1}{(n)^4 \pi^4} - \frac{1}{(n)^6 \pi^6} \right] = 36486450 \left[\frac{1}{5040} \frac{\pi^6}{945} - \frac{1}{120} \frac{1}{\pi^2} \frac{\pi^8}{9450} + \frac{1}{6} \frac{1}{\pi^4} \frac{\pi^{10}}{93555} - \frac{1}{\pi^6} \frac{691\pi^{12}}{638512875} \right] = \pi^6$$

$$k = 4 \text{ gives series as } \sum_{n=1}^{\infty} \frac{1}{n^6} 45(15)(13)(11) \frac{1}{9} (9)! \left[\frac{1}{9!} - \frac{1}{7!} \frac{1}{(n)^2 \pi^2} + \frac{1}{5!} \frac{1}{(n)^4 \pi^4} - \frac{1}{3!} \frac{1}{(n)^6 \pi^6} + \frac{1}{(n)^8 \pi^8} \right] = 3891888000 \left[\frac{1}{362880} \frac{\pi^6}{945} - \frac{1}{5040} \frac{1}{\pi^2} \frac{\pi^8}{9450} + \frac{1}{120} \frac{1}{\pi^4} \frac{\pi^{10}}{93555} - \frac{1}{6} \frac{1}{\pi^6} \frac{691\pi^{12}}{638512875} + \frac{1}{\pi^8} \frac{2\pi^{14}}{18243225} \right] = \pi^6$$

On similar lines, the series representation (21) can be proved for any integer k value.

References

[1] Alzer, H., Karayannakis, D. and Srivastava, H.M.: Series representations for some mathematical constants, J.Math. Anal. Appl. 320,145-162(2006).

[2] Alzer,H. and Koumandos,S.: Series and product representations for some mathematical constants, Periodica Mathematica Hungarica 58,71-82(2009).

[3] Amore, P.: Convergence acceleration of series through a variational approach, J.Math. Anal. Appl. 323,63-77(2006).

[4] Blagouchine, I.V.: Expansions of generalized Euler's constants into the series of polynomials in π^{-2} and into the formal enveloping series with rational coefficients only, J. Number Theory 158,365-396(2016).

[5] Borwein, P.: The amazing number π , Nieuw Arch. Wisk. (Ser.5) 1,254-258(2000).

[6] Kolbig,K.S.: The polygamma function $\psi^{(k)}(x)$ for $x = \frac{1}{4}$ and $x = \frac{3}{4}$, J. Comput. Appl. Math. 75(1996)43-46.