

The thermodynamics and phase transition of a rainbow black hole

Zhong-Wen Feng^{1,*}, De-Ling Tang^{2,†}, Dan-Dan Feng^{2,‡} and Shu-Zheng Yang^{1,§}

¹ College of Physics and Space Science, China West Normal University, Nanchong 637009, China

² School of Science and Engineering, Sichuan Minzu College, Kangding 626001, China

(Dated: December 28, 2021)

In this work, we construct a new kind of rainbow functions, which has generalized uncertainty principle parameter. Then, we investigate modified thermodynamic quantities and phase transition of rainbow Schwarzschild black hole by employing this new kind of rainbow functions. Our results demonstrate that the rainbow gravity and generalized uncertainty principle have a great effect on the picture of Hawking radiation. They prevent black holes from total evaporation and cause a remnant. In addition, after analyzing the modified local thermodynamic quantities, we find that the effect of rainbow gravity and the generalized uncertainty principle lead to one first-order phase transition, two second-order phase transitions, and two Hawking-Page-type phase transitions in the thermodynamic system of rainbow Schwarzschild black hole.

I. INTRODUCTION

In the theoretical physics, one of the biggest challenges is to combine the quantum theory with gravity theory. In order to achieve this purpose, various quantum gravity theories, such as the string theory, loop quantum gravity, M-theory, and non-commutative geometry have been proposed [1–5]. According to those theories, it is found that the existence of a minimum measurable length, which is expected to be of the order of the Planck scale. However, under the linear Lorentz transformations, one finds that the Planck length is variable, which is conflict with the previous conclusions. To resolve this contradiction, the standard energy-momentum dispersion relation needs to be modified [6–8]. Nowadays, the modifications of standard dispersion relation are called as modified dispersion relation (MDR) [9–11]. Interestingly, it should be noted that the MDR can modify the special relativity. In Ref. [12], this generalization of special relativity is called as double special relativity (DSR), which has two invariants: the velocity of light c , and the Planck length ℓ_p (or Planck energy E_p). However, DSR still two flaws in the theory: one is the DSR can only be used in flat spacetime, the other is the dual position space in this theory definition suffers a nonlinearity of the Lorentz transformation. Therefore, Magueijo and Smolin generalized the DSR into curved spacetime and proposed the rainbow gravity (RG) [13]. In the RG, the energy of a test particle affects the geometry. In other word, the spacetime cannot be described by a single metric, but a family of metrics (rainbow of metrics) parameterized by the ratio E/E_p . According to the RG, the deformed energy-momentum relation is given by

$$E^2 f^2(E/E_p) - p^2 g^2(E/E_p) = m^2, \quad (1)$$

where E_p , p and m represent the Planck energy, the momentum and mass of a test particle, respectively. The correction terms $f(E/E_p)$ and $g(E/E_p)$ are the rainbow functions, which satisfy the limits $\lim_{E/E_p \rightarrow 0} f(E/E_p) = 1$ and $\lim_{E/E_p \rightarrow 0} g(E/E_p) = 1$. Those relations show that the RG is constraint to reproduce the standard dispersion relation in low energy limit. According to energy dependent modification to the dispersion relation, metric in RG is given by $g(E) = \eta^{ab} e_a(E) \otimes e_b(E)$ with the energy dependence of the frame fields $e_0(E/E_{Pl}) = (1/f(E/E_{Pl})) \tilde{e}_0$ and $e_i(E/E_p) = (1/f(E/E_p)) \tilde{e}_i$.

In recent years, the implications of RG have been investigated in many contexts, such as cosmology, astrophysics, and black hole physics [14–18]. In particular, the RG has a very significant effect on the thermodynamics of black holes. In Refs. [19–35], the authors studied the modified thermodynamics of black holes by using the RG. These results demonstrated that the modifications are departed from the classical cases when the size of black holes approach the Planck scale. It indicates that the RG can stop the Hawking radiation in the last stages of black holes' evolution and naturally leads to remnants. Therefore, the RG can used to address the information loss and naked singularity problems of black holes. Recently, according to the rainbow functions that were proposed by Magueijo and Smolin, we discussed the thermodynamics and phase transition of a spherically symmetric black hole. Our calculations showed that the RG changes the picture of black hole's thermodynamic phase transition [36]. In addition, we found three Hawking-Page-type phase transitions in the framework of the RG.

On the other hand, the minimum measurable length can also change the conventional Heisenberg uncertainty principle (HUP) to the so-called generalized uncertainty principle (GUP). Since the GUP can be applied to many physical systems, it attracted people's attention now [37–44]. In Ref. [37], Adler, Chen and Santiago pointed out that uncertainties for position and momentum of the GUP (namely, Δx and Δp) can be considered as the temperature and radius of the black holes. With this heuristic theory, they studied the GUP corrected Hawking

* Email: zwfengphy@163.com

† Email: cwnutangdl@163.com

‡ Email: ncddfeng@163.com

§ Email: szyangphys@126.com

ing temperature of Schwarzschild (SC) black hole and showed that the modified Hawking temperature of SC black hole is higher than that of conventional case. However, in our previous work, by using the GUP together with Hamilton-Jacobi equation, the GUP corrected thermodynamics of SC black hole are similar to the results obtained in the framework of RG. This indicates that there has some connection between the GUP and RG [45]. From the above discussion, it is interesting to incorporate the GUP with RG to investigate the modified thermodynamics and phase transition of black holes, this may lead to some novel results. Following this line of research, we construct a new kind of rainbow functions, which contains the effect of GUP in this paper. Then, by using this new kind of rainbow functions, the modified thermodynamics, thermodynamic criticality and the phase transition of rainbow SC black hole will be analyzed.

The remainders of this paper are outlined as follows. In Section II, we derive a MDR via the GUP that proposed by Kempf, Mangano, and Mann. Then, comparing this MDR with the general form of RG, a new kind of rainbow functions, which contains the GUP parameter is obtained. In Section III, according to the new kind of rainbow functions, we calculate the modified thermodynamics of SC black hole in the context of RG. In Section IV, we study the modified local thermodynamic quantities of rainbow SC black hole, and discuss the phase transition between the hot space and the states of black holes. Finally, the discussion is given in Section V.

Throughout the paper, the natural units of $k_B = c = 1$ are used

II. THE RAINBOW FUNCTIONS WITH EFFECT OF GUP

To begin with, we need to recapitulate the way of getting a MDR, which influenced by the GUP. In Ref. [46], according to the GUP that proposed Kempf, Mangano, and Mann, we derive a kind of MDR, and used it to show that the GUP parameters can be constrained by the gravitational wave event GW150914 and GW151226. The expression of the GUP is given by

$$\Delta x \Delta p \geq \frac{\hbar}{2} \left[1 + \beta (\Delta p)^2 \right], \quad (2)$$

where Δx and Δp are the uncertainties for position and momentum, respectively. $\beta = \beta_0 \ell_p^2 = \beta_0 / M_p^2 = \beta_0 / E_p^2$ with the GUP parameters β_0 , and ℓ_p , M_p , and E_p are represent the Planck length, Planck mass, and Planck energy, respectively. [47]. Based on Eq. (2), one can easy obtain a nonzero minimal uncertainty $\Delta x_{\min} \approx \ell_p \sqrt{\beta_0}$. Furthermore, the operators for the modified position operator x_i and modified momentum p_i in Eq. (2) can be defined as

$$x_i = x_{0i}; \quad p_i = p_{0i} (1 + \beta \mathbf{k}^2), \quad (3)$$

where usual position operator x_{0i} and usual momentum p_{0i} are satisfying the canonical commutation relations $[x_{0i}, p_{0j}] = i\hbar \delta_{ij}$, $\mathbf{k}^2 = \delta_{ij} k^{0i} k^{0j}$, thus $\mathbf{k} = \sqrt{\delta_{ij} k^{0i} k^{0j}}$. When considering a general gravitational background metric $ds^2 = g_{ab} dx^a dx^b = g_{00} dt^2 + g_{ij} dx^i dx^j$ the modified square of the four-momentum can be expressed as [48]

$$p_a p^a = g_{ab} p^a p^b = g_{00} (p^{00})^2 + g_{ij} p^{0i} p^{0j} (1 + \beta \mathbf{k}^2)^2. \quad (4)$$

It is normally set that $g_{00} = -1$, in which case the space-time of MDR is flat. Next, by ignoring the higher order term $\mathcal{O}(\beta)$, the equation above can be rewritten as

$$p_a p^a = - (p^{00})^2 + \mathbf{k}^2 + 2\beta \mathbf{k}^2 \mathbf{k}^2. \quad (5)$$

In the right side of Eq. (5), the first two terms form the standard dispersion relation, that is $-(p^{00})^2 + \mathbf{k}^2 = -m^2$. Thus, Eq. (5) can be rewritten as $p_a p^a = -m^2 + 2\beta \mathbf{k}^2 \mathbf{k}^2$, and its time component of the momentum becomes

$$(p^{00})^2 = m^2 + \mathbf{k}^2 (1 - 2\beta \mathbf{k}^2). \quad (6)$$

Using the definition of a particle's energy $E = p^{00}$, the energy of a particle can be expressed in terms of the three spatial momentum and mass as follows

$$E^2 - \mathbf{k}^2 (1 - 2\beta \mathbf{k}^2) = m^2. \quad (7)$$

Now, for simplify the next research, we assume the mass of the test particle is zero, so that the momentum in Eq. (7) satisfying the standard massless energy-momentum dispersion relation, that is $E^2 = \mathbf{k}^2$. Furthermore, in order to obtain a MDR that consistent with Eq. (1), we need retained the form of $\mathbf{k}^2$ outside parentheses and replaced $\mathbf{k}^2$ in parentheses with the standard massless energy-momentum dispersion. Finally, the Eq. (7) can be rewritten as [31]

$$E^2 - p^2 (1 - 2\beta_0 E^2 / E_p^2) = 0. \quad (8)$$

Next, comparing Eq. (8) with Eq. (1), a new kind of rainbow functions is obtained as follows:

$$f(E/E_p) = 1, \quad g(E/E_p) = \sqrt{1 - 2\beta_0 E^2 / E_p^2}. \quad (9)$$

Now, by employing the GUP (2) and a general gravitational background metric, we construct a new kind of rainbow functions (9). It is obviously that Eq. (9) contains the GUP parameter β_0 and the ratio E/E_p , which indicates that it influenced by the effect of GUP and RG. In addition, Eq. (9) satisfying the conditions of $\lim_{E/E_p \rightarrow 0} f(E/E_p) = 1$ and $\lim_{E/E_p \rightarrow 0} g(E/E_p) = 1$ at low energies. In the subsequent discussions, by incorporating the rainbow functions (9) with a specific line element, we investigate the modified thermodynamics of a rainbow black hole.

III. THE MODIFIED THERMODYNAMICS OF SCHWARZSCHILD BLACK HOLE IN THE FRAME WORK OF RG

In this section, by utilizing the rainbow functions (9) with the definition of surface gravity and first law of thermodynamics, the modified thermodynamics of rainbow SC black hole are calculated. In Ref. [13], the line element of rainbow SC black hole is given by

$$\begin{aligned} ds^2 &= A(r, E, E_p) dt^2 + B(r, E, E_p) dr^2 + C(r, E, E_p) d\Omega^2 \\ &= -\frac{1 - (2GM/r)}{f^2(E/E_p)} dt^2 + \frac{[1 - (2GM/r)]^{-1}}{g^2(E/E_p)} dr^2 + \frac{r^2}{g^2(E/E_p)} d\Omega^2 \end{aligned} \quad (10)$$

where $A(r, E, E_p)$, $B(r, E, E_p)$ and $C(r, E, E_p)$ represent the geometric properties of the rainbow black hole, $d\Omega^2$ denotes the line elements of 2-dimensional hyper-surfaces, which can be expressed as $d\theta^2 + \sin^2\theta d\phi^2$ in this work. Obviously, metric (10) has only one root of $r_H = 2GM$ satisfying $A(r_H) = 0$. For $f(E/E_p) = g(E/E_p) = 1$, Eq. (10) reduces to the original metric of SC black hole. According to the line element of rainbow SC black hole, the surface gravity on the event horizon is given by

$$\kappa = -\frac{1}{2} \lim_{r \rightarrow r_H} \sqrt{-\frac{g^{11}}{g^{00}} \frac{\partial_r(g^{00})}{g^{00}}} = \frac{g(E/E_p)}{f(E/E_p)} \frac{1}{4GM}. \quad (11)$$

It is clear that the original surface gravity $\kappa_0 = 1/4GM$ is modified by the rainbow functions. Using Eq. (11), one can easily obtain the deformed Hawking temperature of SC black hole

$$T_H^{\text{modified}} = \frac{\kappa}{2\pi} = \frac{g(E/E_p)}{f(E/E_p)} T_H = \sqrt{1 - 2\beta_0 \frac{E^2}{E_p^2}} \frac{1}{8\pi GM}, \quad (12)$$

where $T_H = 1/8\pi GM$ is the original Hawking temperature of SC black hole. One can see that the modified Hawking temperature is depended on the GUP parameter β_0 , the ratio E/E_p and the mass of SC black hole, which means that the GUP and RG have a very significant effect on the thermodynamics of black hole. When $\beta_0 \rightarrow 0$ or $E/E_p \rightarrow 0$, the original Hawking temperature is recovered.

Here, we have two schemes to eliminate the energy ω dependence of the probe in Eq. (12). In scheme I, the HUP $\Delta x \geq 1/\Delta p$ still holds in the RG. Therefore, the HUP can be translated into a lower bound on the energy of a particle emitted in Hawking radiation, that is, $E \geq 1/\Delta x$ [37]. Then, in the vicinity of the black hole horizon, the uncertainty in the position can be taken to be the radius of the SC black hole $\Delta x \approx r_H = 1/2GM$. Thus, one has the following relation

$$E \geq \frac{1}{r_H} = \frac{1}{2GM}. \quad (13)$$

Substituting Eq. (13) into Eq. (12), the modified Hawking temperature becomes

$$T_H^{\text{modified}} = \sqrt{1 - \frac{\beta_0}{2G^2 M^2 E_p^2}} T_H = \sqrt{1 - \beta_0 \frac{M_p^2}{2M^2}} \frac{1}{8\pi GM}, \quad (14)$$

where in above equation the relation $E_p^{-2} = M_p^{-2} = G$ is used. It should be noted that the expression in the square root can not be less than zero, namely, $1 - \beta_0 M_p^2 / 2M^2 \geq 0$. Therefore, one can obtain a minimum mass of rainbow SC black hole, namely, $M_{\min} = M_p \sqrt{\beta_0/2}$. The minimum mass of rainbow SC black hole shows that the minimum mass of rainbow SC black hole is related to β_0 and M_p , it indicates that the rainbow SC black hole has a remnant, which is of the order of Planck scale and can be considered as a candidate of dark matter.

In scheme II, one can substitute the relation $\Delta p \sim 1/r_H$ and MDR (1) into Eq. (12) [23–35, 49, 50], the modified Hawking temperature becomes


$$\tilde{T}_H^{\text{modified}} = \sqrt{1 - \frac{2\beta_0 E_p^2}{2\beta_0 + 4G^2 M^2 E_p^2}} \frac{1}{8\pi GM}. \quad (15)$$

Now, by using Eq. (14) and Eq. (15), the behaviour of the original Hawking temperature T_H and modified Hawking temperature of SC black hole is depicted in Fig. 1.


In Fig. 1(a), the black dotted curve represents the original Hawking temperature of SC black hole for $\beta_0 = 0$, whereas the red solid curve illustrates the modified Hawking temperature of SC black hole for $\beta_0 = 1$. This diagram shows that the original Hawking temperature tends to infinity as $M \rightarrow 0$. On the contrary, the modified Hawking temperature shows another image of the black hole thermodynamics evolution. At the beginning of the evolution, the behaviour of modified Hawking temperature and that of original Hawking temperature look very much alike, since the effect of quantum gravity is weak at large scale. However, as the evolution progresses, the mass of the black hole gradually decrease, the behaviour modified Hawking temperature deviated from the original case. It rises at first, and then decreases rapidly after it reaches a maximum value. Finally, the modified temperature approaches zero when the mass tends to its minimum value $M_{\min}$. However, the behaviour of modified Hawking temperature (blue solid curve) in Fig. 1(b) is different from that in Fig. 1(a). In this case, $\tilde{T}_H^{\text{modified}}$ becomes finite when the mass of rainbow SC black hole goes to zero. Since the modified Hawking temperature $\tilde{T}_H^{\text{modified}}$ and its corresponding thermodynamic phase transition have been studied in detail in Refs. [49–51], we only focus on the Eq. (14) and its corresponding thermodynamic phase transition in upcoming contents.

Now, by using the area law [52, 53], the modified entropy is

$$S^{\text{modified}} = \frac{2\pi G}{M \sqrt{2 - \beta_0 M_p^2 / M^2}} \Xi, \quad (16)$$



(a) Scheme I



(b) Scheme II

FIG. 1. The $T - M$ diagram of original SC black hole and rainbow SC black hole, where G and M_p are equal to 1.

where $\Xi = \sqrt{2}M(2M^2 - \beta_0 M_p^2) + \beta_0 M_p^2 \sqrt{2M^2 - \beta_0 M_p^2}$. When $\beta_0 = 0$, the above equation becomes to the area law of the entropy of black hole $S = A/4 = 4\pi G M^2$ with the area of black hole A , this is consistent with the original entropy in Ref. [52]. Next, in order to determine whether the rainbow SC black hole exists a remnant, it is necessary to further analyze the specific heat of rainbow SC black. According to Ref. [45], the definition of specific heat is given by $\mathcal{C} = T_H (\partial S / \partial M) (\partial T_H / \partial M)^{-1}$. Therefore, the modified specific heat is

$$\mathcal{C}^{\text{modified}} = -\frac{4\pi G M^4}{M^2 - M_p^2 \beta_0} \sqrt{4 - \frac{2M_p^2 \beta_0}{M^2}}. \quad (17)$$

From the equation above, it is found that $\mathcal{C}^{\text{modified}} = 0$ at $M_{\min} = M_p \sqrt{\beta_0/2}$, which is equal to Eq. (15). For $\beta_0 = 0$, Eq. (17) recovers the original case $\mathcal{C}_0 = -8\pi M^2$. The associated $\mathcal{C} - M$ diagram is displayed in Fig. 2. One can see that the value of original specific heat (black




FIG. 2. The $\mathcal{C} - M$ diagram of original SC black hole and rainbow SC black hole, where we set $G = M_p = 1$.

dashed curve) is always negative, and it vanishes when $M \rightarrow 0$. Different from the original one, the modified specific heat (red solid curve) has a vertical asymptote at a certain point, where the temperature of rainbow SC black hole reaches its maximum value. It causes the value of specific heat to change from negative to positive, which implies a thermodynamic phase transition occurred at this certain location. The specific phase transition of rainbow SC black hole will be analyzed in the next section. In addition, it is clear that the modified specific heat vanishes when the mass of the reaches $M_{\min}$. Since the black hole does not exchange their heat with the surrounding spacetime when the specific heat is equal to zero, we can determine the rainbow SC black hole has a remnant at $M_{\min}$. Therefore, the remnant mass is $M_{\text{res}} = M_{\min}$.

IV. MODIFIED LOCAL THERMODYNAMIC QUANTITIES AND PHASE TRANSITION OF RAINBOW SC BLACK HOLE

It is well known that the black holes have rich phase structures and critical phenomena [54–63]. The phase structures and critical phenomena of black holes have been a subject of fascination for more than forty years since they do not only offer a deep insight into the quantum properties of gravity, but also allow for a deeper understanding of the thermal properties of conformal field theories. Therefore, in this section, for investigating the phase transition from the hot space to black hole, we need to further study the local thermodynamic quantities of the rainbow SC black hole. In order to obtain the local thermodynamic quantities, the rainbow SC black hole needs to be located at the center of a spherical cavity of radius $\mathcal{R}$ [64, 65]. Hence, the expression of local

temperature of rainbow SC black hole is given by

$$T_{\text{local}}^{\text{modified}} = \sqrt{1 - \beta_0 \frac{M_p^2}{2M^2}} \frac{1}{8\pi G M \sqrt{1 - \frac{2GM}{\mathcal{R}}}}. \quad (18)$$

As $\mathcal{R} \rightarrow \infty$, the above equation recovers the modified Hawking temperature. When considering the cavity of radius $\mathcal{R}$ as a invariable quantity, one can calculate the critical value of black hole's mass M_c , GUP parameter β_c , and local temperature T_c by the following equations [36]

$$\left(\frac{\partial T_{\text{local}}^{\text{modified}}}{\partial M} \right)_{\mathcal{R}} = 0, \quad \left(\frac{\partial^2 T_{\text{local}}^{\text{modified}}}{\partial M^2} \right)_{\mathcal{R}} = 0. \quad (19)$$

Substituting Eq. (18) into Eq. (19), one can obtain the critical mass, critical GUP parameter, and critical local temperature as follows

$$M_c = \frac{4\mathcal{R}}{15G}, \quad \beta_c = \frac{16\mathcal{R}^2}{375G^2M_p^2}, \quad T_c = \frac{15}{32\pi\mathcal{R}} \sqrt{\frac{3}{2}}. \quad (20)$$

For the convenience of discussion, here we setting $\mathcal{R} = 10$, $G = M_p = 1$ as an example. Thus, one obtain that $M_c = 2.66667$, $\beta_c = 4.26667$ and $T_c = 0.01827$. According to Eq. (18) and Eq. (20), the corresponding " $T_{\text{local}}^{\text{modified}} - M$ " diagram of rainbow SC black hole for different β_0 is depicted in Fig. 3.




FIG. 3. The $T_{\text{local}}^{\text{modified}} - M$ diagram of rainbow SC black hole for different β_0 with $\mathcal{R} = 10$ and $G = M_p = 1$.

Fig. 3 depicts $T_{\text{local}}^{\text{modified}} - M$ as a function of mass for fixed $G = M_p = 1$ and $\mathcal{R} = 10$. The value of β_0 increases from top to bottom. The red solid curve, black dashed curve and blue dotted curve correspond to the critical GUP parameter $\beta_0 = 0.234\beta_c \approx 1$, $\beta_0 = \beta_c = 4.26667$, and $\beta_0 = 1.17\beta_c \approx 5$, respectively. Obviously, one can observe a phase transition for $0 < \beta_0 < \beta_c$. Hence, in order to study the phase transition and critical phenomena

of rainbow SC black hole, we set $\beta_0 = 1$ in the following discussions.

Next, let us plot the original and rainbow local temperature versus mass for $G = M_p = 1$ and $\mathcal{R} = 10$ in Fig. 4. The black dashed curve represents the local temperature of original SC black hole, and the red solid curve represents the local temperature of rainbow black hole. The evaporation process of SC black hole in the framework of rainbow function is commonly organized in three stages: at the first stage ($M_2 \leq M \leq M_3$), the rainbow local temperature decreases through its evaporation process; at the second stage ($M_1 \leq M \leq M_2$), the rainbow local temperature increases as mass decreases, lasting up to the mass in which it comes near to a maximum value; at the final stage ($M_0 \leq M \leq M_1$), the mass of rainbow SC black hole cannot get smaller than M_0 since the negative temperature becomes violates the laws of thermodynamics, which indicates that the RG can stop the black hole evolution and leads to a remnant. Hence, we have $M_0 = M_{\text{res}}$. Employing the first law of thermody-




FIG. 4. The original and modified local temperature versus mass at $G = M_p = 1$ and $\mathcal{R} = 10$.

namics, expression of total internal energy is given by

$$\begin{aligned} E_{\text{local}}^{\text{modified}} &= \int_{M_0}^M T_{\text{local}}^{\text{modified}} dS \\ &= \frac{\mathcal{R}}{G} \left(\sqrt{1 - \frac{M_p G \sqrt{2\beta_0}}{\mathcal{R}}} - \sqrt{1 - \frac{2GM}{\mathcal{R}}} \right). \end{aligned} \quad (21)$$

When $\beta_0 \rightarrow 0$, the local energy of original SC black hole $E_{\text{local}} = \mathcal{R} \left(1 - \sqrt{1 - 2GM/\mathcal{R}} \right) / G$ is recovered. Then, using the modified local temperature Eq. (18) and the total internal energy Eq. (21), the modified heat capacity

at fixed $\mathcal{R}$ can be expressed as

$$\begin{aligned} C_{\text{local}}^{\text{modified}} &= \left(\frac{\partial E_{\text{local}}^{\text{modified}}}{\partial T_{\text{local}}^{\text{modified}}} \right)_{\mathcal{R}} \\ &= \frac{8\pi G M^3 (\mathcal{R} - 2GM) \sqrt{4M^2 - 2\beta_0 M_p^2}}{M (6GM^2 - 2M\mathcal{R} - 5\beta_0 G M_p^2) + 2\mathcal{R} M_p^2 \beta_0}. \end{aligned} \quad (22)$$

Clearly, when β_0 vanishes, the Eq. (22) reduces to the local heat capacity of original SC black hole $C_{\text{local}} = 8\pi G M^2 (\mathcal{R} - 2GM) / (3GM - \mathcal{R})$ [66]. According to Eq. (22), the variation of the heat capacity $C_{\text{local}}^{\text{modified}}$ with the mass M is plotted in Fig. 5.




FIG. 5. The original and modified local heat capacity versus mass with $G = M_p = 1$ and $\mathcal{R} = 10$.

As shown in Fig. 5, the rainbow heat capacity of SC black hole (red solid line) changes its sign at M_1 and M_2 for which the denominator in Eq. (22) vanishes. Meanwhile, the rainbow heat capacity C_{rainbow} goes to zero when mass approaches M_0 . When $M \rightarrow M_3$, the rainbow heat capacity tends to the original case (black dashed line) since the effect of GUP is negligible at this point. The rainbow heat capacity is divergent at the extreme points M_1 and M_2 . Hence, the phase transitions near there are second-orders. It is well known that the positive heat capacity means the thermally stable state, while the negative heat capacity is considered to be the unstable solution. So, it is easy to find that the rainbow black hole has one unstable region ($M_1 \leq M \leq M_2$), and two stable regions ($M_0 \leq M \leq M_1$ and $M_2 \leq M \leq M_3$). Now, the rainbow black hole can be classified into three types based on their local temperature and the heat capacity, namely, a large black hole (LBH) for $M_2 \leq M \leq M_3$, a small black hole (SBH) for $M_0 \leq M \leq M_1$, and an intermediate black hole (IBH) for $M_1 \leq M \leq M_2$, which never appears in the original case. Obviously, the small

black hole (SBH) and the large black hole (LBH) are stable, showing that they can survive for a long time within the frame of the RG theory. On the contrary, the IBH is unstable because its heat capacity is negative, which implies that it would decay into the SBH or LBH quickly. In addition, it should be noted that the stability of the black holes can be known from their free energy.

In order to obtain more details of the thermodynamic phase transition of the rainbow SC black hole, we use the three states of rainbow SC black hole to analyze the phase transition. In Refs. [67, 68], the free energy is defined as $F_{\text{on}} = E_{\text{local}} - T_{\text{local}} S$. Therefore, the free energy of rainbow SC black hole is given by

$$\begin{aligned} F_{\text{on}}^{\text{modified}} &= \frac{\mathcal{R}}{G} \left(\sqrt{1 - \frac{M_p G \sqrt{2\beta_0}}{\mathcal{R}}} - \sqrt{1 - \frac{2GM}{\mathcal{R}}} \right) \\ &\quad - \frac{\sqrt{2}M (2M^2 - M_p^2 \beta_0) + \Delta}{4M^2 \sqrt{2 - 4GM/\mathcal{R}}}. \end{aligned} \quad (23)$$

where $\Delta = M_p^2 \beta_0 \sqrt{2M^2 - \beta_0 M_p^2} \ln \left(2M + \sqrt{4M^2 - 2\beta_0 M_p^2} \right)$. Note that Eq. (23) is reduced to the original free energy for $\beta_0 = 0$. Meanwhile, as seen from Fig. 6 that the thermodynamic phase transition of the rainbow SC black hole occurs for $0 < \beta_0 < \beta_c$.




FIG. 6. Variation in free energy of rainbow SC black hole with local temperature for different β_0 with $G = M_p = 1$ and $\mathcal{R} = 10$.

From Fig. 6, we show F_{on} curves with T_{local} for different β_0 . The value of GUP parameter β_0 increases from top to bottom. It is clear that the swallow tail structure appears when the GUP parameter β_0 is smaller than the critical value β_c , which indicates there is a two-phase coexistence state. The results are consistent with the profile of $T_{\text{local}}^{\text{modified}} - M$ in Fig. 3. Meanwhile, Fig. 6 indicates a reverse van der Waals like behavior for $T > T_c$, which is completely different from the van der Waals one of physical thermodynamic systems in Ref. [69].

For further investigate the phase transition between the black holes and the hot space, we plot Fig. 7.




FIG. 7. (a) shows the original free energy of SC black hole as function of the local temperatures. (b) shows the modified free energy of SC black hole as function of the local temperatures. Here we choose $G = M_p = 1$ and $\mathcal{R} = 10$.

As seen from Fig. 7(a), it is found that the original free energy vanishes when $M \rightarrow 0$ because the vacuum state is Minkowski spacetime. Therefore, the $F_{\text{on}}^{\text{HFS}}$ represents the free energy of the hot flat space (HFS). In addition, the local temperature has a minimal value T'_1 , below which no black hole solution exists. The small-large black hole (Hawking-Page) transition occurs at T'_c . For $T < T'_c$, both the free energy of small and large black holes are higher than the $F_{\text{on}}^{\text{HFS}}$, it means that the HFS is more probable than the SBH and LBH. However, for $T > T'_c$, the free energy of SBH is higher than the $F_{\text{on}}^{\text{HFS}}$ while the LBH is lower than the $F_{\text{on}}^{\text{HFS}}$, which indicates that the LBH is more probable than the HFS. Therefore, one can find the radiation collapse to the LBH, and the SBH eventually decays into the LBH thermodynamically [36].

Next, let us focus on Fig. 7(b) for $G = \beta_0 = M_p = 1$ and $\mathcal{R} = 10$. In contrast to the conventional one, the rainbow free energy becomes to zero when $M \rightarrow M_{\text{res}} = M_0$, which implies the rainbow SC black hole only exists the hot curved space (HCS) [70]. Therefore, we study phase transition between various black hole states and the HCS via analyzing the free energies of the rainbow SC black hole.

(i) The SBH and IBH are degenerate at T_1 with the mass M_1 ; the intermediate-large black hole transition occurs at the inflection point T_2 corresponding to M_2 . Interestingly, it is found two intersections points (Hawking-Page-type critical points) between the line of free energy and the $F_{\text{on}}^{\text{HFS}}$ at $T_c^{(1)}$ and $T_c^{(2)}$ in Fig. 7(b), while only one Hawking-Page phase transition in the original case, as seen from Fig. 7(a).

(ii) There is a one-order phase transition at critical temperature $T_c^{\text{first-order}}$ since the images shows a characteristic swallow tail behavior.

(iii) For $T_0 < T < T_c^{(1)}$, the free energy of IBH and LBH are higher than the $F_{\text{on}}^{\text{HCS}}$, whereas the SBH is lower than $F_{\text{on}}^{\text{HCS}}$, it implies that the stable SBH is more probable than HFS. For $T_c^{(1)} < T < T_c^{\text{first-order}}$, both SBH and LBH are lower than the $F_{\text{on}}^{\text{HCS}}$. It should be noted that LBH and IBH should decay into SBH since the free energy of SBH is lower than those of IBH and LBH in this region. Consequently, for $T_0 < T < T_c^{\text{first-order}}$, SBH should undergoes a tunneling, which indicates that the HFS can collapse to SBH. As the local temperature increase, one finds that free energies of black holes satisfy the relation $F_{\text{on}}^{\text{LBH}} < F_{\text{on}}^{\text{SBH}} < F_{\text{on}}^{\text{HCS}} < F_{\text{on}}^{\text{IBH}}$ for $T_c^{\text{first-order}} < T < T_c^{(2)}$. Then, the relation becomes $F_{\text{on}}^{\text{LBH}} < F_{\text{on}}^{\text{SBH}} < F_{\text{on}}^{\text{IBH}} < F_{\text{on}}^{\text{HCS}}$ for $T_c^{(2)} < T < T_1$, it means that HCS does not only collapse into SBH but also into IBH and LBH. However, the IBH with negative heat capacity is unstable. Therefore, it would decay into the SBH. Meanwhile, it is obvious that the free energy of SBH is always higher than free energy of the LBH, it leads to the SBH eventually decays into the LBH thermodynamically. Therefore, for $T_c^{\text{first-order}} < T < T_1$, the HCS would eventually decays into the LBH via the quantum tunneling effect.

V. DISCUSSION AND CONCLUSION

In this paper, we first derived a new kind of rainbow functions, which contains the GUP parameter. Then, we considered this rainbow functions to investigate the modified thermodynamics and phase transitions the rainbow SC black hole. First of all, our results showed that the rainbow functions change the picture of Hawking radiation and stop the evaporation of a black hole as its size approaches the Planck scale. It naturally leads to a remnant of SC black holes, which can be considered as a candidate of dark matter. Second, as seen from Fig. 3, the phase transition occurs when the GUP pa-

parameter is smaller than the critical points. Therefore, we have set the rainbow parameters as $\beta_0 = 1$ in order for the study to be focused and feasible. Third, since the heat capacity enjoys two divergencies in Fig. 5, it demonstrated the thermodynamic system of rainbow SC black hole has two second order phase transitions at M_1 and M_2 . Fourth, according to the “ $F_{\text{on}} - T_{\text{local}}$ ” diagram, one can find an unstable black hole (IBH) interpolating between the stable SBH and LBH, which never exists in the original case. Fifth, the $F_{\text{on}}^{\text{modified}}$ surface showed the swallow tail shape, which indicates that the thermodynamic system of rainbow SC black hole exists a first-order transition. Sixth, it is found two Hawking-Page-type critical points from Fig. 7(b), while only one

Hawking-Page critical point in the original case. Finally, for $T_0 < T < T_c^{\text{First-order}}$, since the free energy of SBH is lower than those of the IBH and LBH, the unstable IBH and stable LBH decay into the stable SBH. However, for $T_c^{\text{First-order}} < T < T_1$, it is obviously the free energy of stable LBH is always lower than those of the SBH and IBH, this leads to the SBH and IBH eventually collapse into the LBH thermodynamically.

Acknowledgements

This work is supported in part by the National Natural Science Foundation of China (Grant Nos. 11847048 and 11573022) and the Fundamental Research Funds of China West Normal University (Grant Nos. 17E093, 17YC518 and 18Q067).

-
- [1] D. Amati, M. Ciafaloni, G. Veneziano, *Phys. Lett. B* **216**, 41 (1989).
 - [2] E. Witten, *Nucl. Phys. B*, **443**, 85 (1995).
 - [3] G. Amelino-Camelia, J. R. Ellis, N. E. Mavromatos, D. V. Nanopoulos and S. Sarkar, *Nature* **393**, 763 (1998).
 - [4] R. Gambini, J. Pullin, *Phys. Rev. D* **59**, 124021 (1999).
 - [5] F. Girelli, E. R. Livine, D. Oriti, *Nucl. Phys. B* **708**, 411 (2005).
 - [6] J. J. Peng, S. Q. Wu, *Gen. Relativ. Gravit.* **40**, 2619 (2008).
 - [7] G. 't Hooft, *Class. Quantum Gravity* **13**, 1023 (1996).
 - [8] V. A. Kosteleck'y, M. Mewes, *Phys. Rev. Lett.* **97**, 140401 (2006).
 - [9] W.-J. Pan, Y.-C. Huang, *Gen. Relativ. Gravit.* **48**, 144 (2016).
 - [10] L. Barcaroli, L. K. Brunkhorst, G. Gubitosi, N. Loret, C. Pfeifer, *Phys. Rev. D* **92**, 084053 (2015).
 - [11] S. Kiyota, K. Yamamoto, *Phys. Rev. D* **92**, 104036 (2015).
 - [12] G. Amelino-Camelia, J. Lukierski, A. Nowicki, *Int. J. Mod. Phys. A* **14**, 4575 (1999).
 - [13] J. Magueijo, L. Smolin, *Class. Quantum Gravity* **21**, 1725 (2004).
 - [14] A. Awad, A. F. Ali, B. Majumder, *JCAP* **10**, 052 (2013).
 - [15] A. S. Sefiedgar, *Int. J. Mod. Phys. D* **25**, 1650101 (2016).
 - [16] A. Chatrabhuti, V. Yingcharoenrat, P. Channuie, *Phys. Rev. D* **93**, 043515 (2016).
 - [17] S. H. Hendi, G. H. Bordbar, B. Eslam Panah, S. Panahiyan, *JCAP* **09**, 013 (2016).
 - [18] B. Eslam Panah, G. H. Bordbar, S. H. Hendi, R. Ruffini, Z. Rezaei, R. Moradi, *Astrophys. J.* **848**, 24 (2017).
 - [19] A. F. Ali, *Phys. Rev. D* **89**, 104040 (2014).
 - [20] A. F. Ali, M. Faizal, M. M. Khalil, *JHEP* **1412**, 159 (2014).
 - [21] A. F. Ali, M. Faizal and B. Majumder, *Europhys. Lett.* **109**, 20001 (2015).
 - [22] A. F. Ali, M. Faizal, M. M. Khalil, *Nucl. Phys. B* **894**, 341 (2015).
 - [23] S. H. Hendi, S. Panahiyan, B. E. Panah, *Adv. High Energy Phys.* **2015**, 743086 (2015).
 - [24] S. H. Hendi, G.-Q. Li, J.-X. Mo, S. Panahiyan, B. E. Panah, *Eur. Phys. J. C* **76**, 571 (2016).
 - [25] S. H. Hendi, S. Panahiyan, B. Eslam Panah, M. Momennia, *Eur. Phys. J. C* **76**, 150 (2016).
 - [26] S. H. Hendi, M. Momennia, B. E. Panah, M. Faizal, *Astrophys. J.* **827**, 153 (2016).
 - [27] S. H. Hendi, B. Eslam Panah, S. Panahiyan, M. Momennia, *Eur. Phys. J. C* **77**, 647 (2017).
 - [28] S. H. Hendi, S. Panahiyan, S. Upadhyay, B. Eslam Panah, *Phys. Rev. D* **95**, 084036 (2017).
 - [29] S. H. Hendi, B. Eslam Panah, S. Panahiyan, M. Momennia, *Eur. Phys. J. C* **77**, 647 (2017).
 - [30] S. H. Hendi, M. Momennia, B. E. Panah, S. Panahiyan, *Phys. Dark Universe* **16**, 26 (2017).
 - [31] S. H. Hendi, S. Panahiyan, B. E. Panah, M. Faizal, M. Momennia, *Phys. Rev. D* **94**, 024028 (2016).
 - [32] M. Dehghani, *Phys. Lett. B* **777**, 351 (2018).
 - [33] B. E. Panah, *Phys. Lett. B* **787**, 45 (2018).
 - [34] R. Mandal, S. Bhattacharyya, S. Gangopadhyay, *Gen. Relativ. Gravit.* **50**, 143 (2018).
 - [35] Z.-W. Feng, Q.-C. Ding, S.-Z. Yang, *Eur. Phys. J. C* **79**, 445 (2019).
 - [36] Z. W. Feng, S. Z. Yang, *Phys. Lett. B* **772**, 737 (2017).
 - [37] R. J. Adler, P. Chen, D. I. Santiago, *Gen. Relativ. Gravit.* **33**, 2101 (2001).
 - [38] A. J. M. Medved, E. C. Vagenas, *Phys. Rev. D* **70**, 124021 (2004).
 - [39] W. Kim, Y. Kim, *Phys. Lett. B* **718**, 687 (2012).
 - [40] M. I. Park, *Phys. Lett. B* **659**, 698 (2008).
 - [41] K. Nozari, S. H. Mehdipour, *Europhys. Lett.* **84**, 20008 (2008).
 - [42] B. Majumder, *Gen. Relativ. Gravit.* **45**, 2403 (2013).
 - [43] P. Bargueno, E. C. Vagenas, *Phys. Lett. B* **742**, 15 (2015).
 - [44] M. A. Anacleto, F. A. Brito, E. Passos, *Phys. Lett. B* **749**, 181 (2015).
 - [45] Z. W. Feng, H. L. Li, X. T. Zu, S. Z. Yang, *Eur. Phys. J. C* **76**, 212 (2016).
 - [46] Z. W. Feng, S. Z. Yang, H. L. Li, X. T. Zu, *Phys. Lett. B* **768**, 81 (2017).
 - [47] A. Kempf, G. Mangano, R. B. Mann, *Phys. Rev. D* **52**, 1108 (1995).
 - [48] B. R. Majhi, E. C. Vagenas, *Phys. Lett. B* **725**, 477 (2013).
 - [49] Y. Kim, W. Kim, *JCAP* **1410**, 003 (2014).
 - [50] M. Eune, Y. Gim, W. Kim, *Phys. Rev. D* **91**, 044037

- (2015).
- [51] S.-T. Hong, Y.-W. Kim, Y.-J. Park, *Phys. Rev. D* **99**, 024047 (2019).
 - [52] J. D. Bekenstein, *Phys. Rev. D* **7**, 2333 (1973).
 - [53] S. W. Hawking, D. N. Page, *Commun. Math. Phys.* **87**, 577 (1983).
 - [54] R. Banerjee, S. K. Modak, S. Samanta, *Phys. Rev. D* **84**, 064024 (2011).
 - [55] D. Kubizňák, R. B. Mann, *JHEP* **1207**, 033 (2012).
 - [56] S. Gunasekaran, R. B. Mann, *JHEP* **2012**, 110 (2012).
 - [57] S. W. Wei, Y. X. Liu, *Phys. Rev. D* **87**, 044014 (2013).
 - [58] R. G. Cai, Y. P. Hu, Q. Y. Pan, Y. L. Zhang, *Phys. Rev. D* **91**, 024032 (2015).
 - [59] M. H. Dehghani, S. Kamrani, A. Sheykhi, *Phys. Rev. D* **90**, 104020 (2014).
 - [60] S. W. Wei, Y. X. Liu, *Phys. Rev. Lett.* **115**, 111302 (2015).
 - [61] X. X. Zeng, H. B. Zhang, L. F. Li, *Phys. Lett. B* **756**, 170 (2016).
 - [62] X. X. Zeng, X. M. Liu, L. F. Li, *Eur. Phys. J. C* **76**, 616 (2016).
 - [63] X. X. Zeng, L. F. Li, *Adv. High Energy Phys.* **2016**, 6153435 (2016).
 - [64] R. C. Tolman, *Phys. Rev.* **35**, 904 (1930).
 - [65] J. Tao, P. Wang, H. T. Yang, *Phys. Rev. D* **94**, 064068 (2016).
 - [66] J. W. York Jr, *Phys. Rev. D* **33**, 2092 (1986).
 - [67] R. G. Cai, *Phys. Rev. D* **65**, 084014 (2002).
 - [68] Y. S. Myung, *Phys. Lett. B* **690**, 534 (2010).
 - [69] S. H. Hendi, A. Dehghani, *Eur. Phys. J. C* **79**, 223 (2019).
 - [70] W. Kim, Y. Kim, *Phys. Lett. B* **718**, (2012) 687.