arXiv:1710.04967v1 [math.CA] 6 Oct 2017

NEW ZEALAND JOURNAL OF MATHEMATICS
Volume ** (****), **–**

SOME GENERATING FUNCTIONS FOR q-POLYNOMIALS

Howard S. Cohl, Roberto S. Costas-Santos, and Tanay V. Wakhare

(Received 20 July, 2017)

Abstract. We obtain q-analogues of the Sylvester, Cesàro, Pasternack, and Bateman polynomials. We also derive generating functions for these polynomials.

1. Introduction

Trying to construct inverse Laplace transforms in order to better understand some radiation and conduction problems, H. Bateman introduced in [**1**] the polynomial

$$\mathscr{Z}_n(z) = {}_2F_2\left(\begin{matrix}-n, n+1\\ 1, 1\end{matrix}; z\right),$$

which we refer to as the Bateman-$\mathscr{Z}$ polynomial. By using [**7**, Theorem 48], we obtain the generating function

$$\sum_{n=0}^{\infty} \mathscr{Z}_n(z)\, t^n = \frac{1}{1-t}\, {}_1F_1\left(\begin{matrix}\frac{1}{2}\\ 1\end{matrix}; \frac{-4zt}{(1-t)^2}\right).$$

Using the above information, the Bateman polynomials are defined as [**2**, p. 25]

$$\mathscr{B}_n(z) = {}_3F_2\left(\begin{matrix}-n, n+1, \frac{z+1}{2}\\ 1, 1\end{matrix}; 1\right).$$

He also obtained the generating functions

$$\sum_{n=0}^{\infty} \mathscr{B}_n(z)\, t^n = \frac{1}{1-t}\, {}_2F_1\left(\begin{matrix}\frac{1}{2}, \frac{z+1}{2}\\ 1\end{matrix}; \frac{-4t}{(1-t)^2}\right),$$

$$\sum_{n=0}^{\infty} \left(\mathscr{B}_n(z-2) - \mathscr{B}_n(z)\right)\, t^n = \frac{2t}{(1-t)^3}\, {}_2F_1\left(\begin{matrix}\frac{3}{2}, \frac{z+1}{2}\\ 2\end{matrix}; \frac{-4t}{(1-t)^2}\right).$$

Lemma 1.1. *Let $t \in \mathbb{C}$. Then the following relation holds:*

$$\sum_{n=0}^{\infty} \mathscr{B}_m(-2n-1)\frac{t^n}{n!} = e^t\, \mathscr{Z}_m(-t). \tag{1.1}$$

The Bateman polynomial $\mathscr{B}_n$ was generalized by Pasternack in [**6**]. He defines the Pasternack polynomial $\mathscr{B}_n^m$ as

$$\mathscr{B}_n^m(z) = {}_3F_2\left(\begin{matrix}-n, n+1, \frac{z+m+1}{2}\\ 1, m+1\end{matrix}; 1\right),$$

2010 *Mathematics Subject Classification* 33D45; 33C20.
Key words and phrases: Basic Hypergeometric Functions, Generating Functions, q-Polynomials.

for $m \in \mathbb{C}\backslash\{-1\}$. The Pasternack polynomials reduce to the Bateman polynomials when $m = 0$.

Further information regarding such polynomials and their connection with (classical) orthogonal polynomials can be found in [**5**]. Indeed, we can write the Pasternack polynomials in terms of the continuous Hahn polynomials [**4**, (9.4.1)] as follows [**5**, p. 893]:

$$\mathscr{B}_n^m(z) = \frac{1}{i^n (m+1)_n}\, p_n\left(\frac{-iz}{2}; \frac{1+m}{2}, \frac{1-m}{2}, \frac{1-m}{2}, \frac{1+m}{2}\right).$$

We also consider the Sylvester polynomials, defined as (see [**8**, p. 185])

$$\varphi_n(z) = \frac{z^n}{n!}\, {}_2F_0\left(\begin{matrix} -n, z \\ - \end{matrix}; \frac{-1}{z}\right).$$

Notice that we also can write the Sylvester polynomials in terms of (classical) orthogonal polynomials [**4**]

$$\varphi_n(z) = (-1)^n L_n^{(-z-n)}(z) = \frac{z^n}{n!}\, C_n(-z; z).$$

Here $L_n^{(\alpha)}$ and C_n represent the Laguerre and Charlier polynomials, respectively. It is also known that the Sylvester polynomials satisfy the generating functions

$$\sum_{n=0}^{\infty} \varphi_n(z)\, t^n = \frac{e^{zt}}{(1-t)^z},$$

$$\sum_{n=0}^{\infty} (\lambda)_n\, \varphi_n(z)\, t^n = \frac{1}{(1-zt)^\lambda}\, {}_2F_0\left(\begin{matrix} \lambda, z \\ - \end{matrix}; \frac{t}{1-zt}\right).$$

The Cesàro polynomials are defined as [**8**, p. 449]

$$g_n^{(s)}(z) = \frac{(1+s)_n}{n!}\, {}_2F_1\left(\begin{matrix} -n, 1 \\ -s-n \end{matrix}; z\right).$$

Observe that this family can be written in terms of Jacobi polynomials [**8**, p. 449] as

$$g_n^{(s)}(z) = P_n^{(s+1,-s-n-1)}(2z-1).$$

Furthermore, they satisfy the generating functions

$$\sum_{n=0}^{\infty} g_n^{(s)}(z)\, t^n = (1-t)^{-s-1}(1-zt)^{-1},$$

$$\sum_{n=0}^{\infty} \binom{n+\ell}{\ell} g_{n+\ell}^{(s)}(z)\, t^n = (1-t)^{-s-1-\ell}(1-zt)^{-1} g_\ell^{(s)}\left(\frac{z(1-t)}{1-zt}\right),$$

where $\ell = 0, 1, 2, \ldots$.

The aim of this paper is to obtain q-analogues of all these families of polynomials as well as q-analogues of the generating functions stated above. The structure of this paper is as follows. In Section 2 we give some preliminaries on q-calculus and we define some q-analogues of Bateman-$\mathscr{Z}$, Bateman, Sylvester, Pasternack, and Cesàro polynomials. In Section 3 we state and derive q-analogues of the given generating functions associated with the q-Bateman-$\mathscr{Z}$, q-Bateman, q-Sylvester, q-Pasternack, and q-Cesàro polynomials.

2. Some q-Calculus. The Definitions.

Let $q \in \mathbb{C}$, $0 < |q| < 1$. A q-analogue of the hypergeometric series ${}_pF_r$ is the basic hypergeometric series [**3**]

$$ {}_r\phi_s\left(\begin{matrix} a_1, a_2, \dots, a_r \\ b_1, b_2, \dots, b_s \end{matrix}; q, z\right) = \sum_{k=0}^{\infty} \frac{(a_1;q)_k (a_2;q)_k \cdots (a_r;q)_k}{(b_1;q)_k (b_2;q)_k \cdots (b_s;q)_k} \left((-1)^k q^{\frac{k(k-1)}{2}}\right)^{1+s-r} \frac{z^k}{(q;q)_k}, $$

where $q \neq 0$ when $r > s+1$, and the (b_i) are such that the denominator never vanishes. We also need to define some other q-analogues, such as the q-analogue of a number $[a]_q$, factorial $[a]_q!$, and the Pochhammer symbol (rising factorial), $(a)_n$. These q-analogues are given as follows:

$$ [a]_q = \frac{1-q^a}{1-q}, $$

$$ [0]_q! = 1, \qquad [n]_q! = \prod_{k=1}^{n} [k]_q, \ n \in \mathbb{N}, $$

$$ (a;q)_0 = 1, \quad (a;q)_n = (1-a)(1-q)\cdots(1-aq^{n-1}), \quad n \in \mathbb{N}. $$

We also will need a q-analogue of the *binomial theorem* [**3**, (1.3.2)]

$$ {}_1\phi_0\left(\begin{matrix} a \\ - \end{matrix}; q, z\right) = \sum_{k=0}^{\infty} \frac{(a;q)_k}{(q;q)_k} z^k = \frac{(az;q)_\infty}{(z;q)_\infty}, \qquad |z| < 1, \ |q| < 1, \tag{2.1} $$

and the q-binomial coefficient

$$ \begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q;q)_n}{(q;q)_k (q;q)_{n-k}}. \tag{2.2} $$

Taking into account the previous definitions and results, we define q analogues of the above introduced polynomials. We define

$$ \mathscr{Z}_n(z;q) = {}_2\phi_2\left(\begin{matrix} q^{-n}, q^{n+1} \\ q, q \end{matrix}; q, q^n z\right), $$

and the q-Bateman polynomial as

$$ \mathscr{B}_n(z;q) = {}_3\phi_2\left(\begin{matrix} q^{-n}, q^{n+1}, q^{\frac{1+z}{2}} \\ q, q \end{matrix}; q, q^n\right), $$

the q-Pasternack polynomial as

$$ \mathscr{B}_n^m(z;q) = {}_3\phi_2\left(\begin{matrix} q^{-n}, q^{n+1}, q^{\frac{1+z+m}{2}} \\ q, q^{m+1} \end{matrix}; q, q^n\right), $$

the q-Sylvester polynomial as

$$ \varphi_n(z;q) = \frac{z^n}{(q;q)_n} {}_2\phi_0\left(\begin{matrix} q^{-n}, q^z \\ - \end{matrix}; q, q^n z^{-1}\right), $$

the q-Cesàro polynomial as

$$ g_n^{(s)}(z;q) = \frac{(q^{s+1};q)_n}{(q;q)_n} {}_2\phi_1\left(\begin{matrix} q^{-n}, q \\ q^{-s-n} \end{matrix}; q, z\right). $$

Remark 2.1. Note that $\mathscr{B}_n^0(z;q) = \mathscr{B}_n(z;q)$.

Lemma 2.2. *The q-Cesàro polynomial can be written as*

$$g_n^{(s)}(z;q) = \sum_{k=0}^{n} \begin{bmatrix} k+s \\ s \end{bmatrix}_q (zq^s)^{n-k}. \tag{2.3}$$

3. The Generating Functions

Theorem 3.1. *For any $t \in \mathbb{C}$ small enough, the q-Bateman-$\mathscr{Z}$ and the q-Bateman polynomials satisfy the following generating functions:*

$$\sum_{n=0}^{\infty} \mathscr{Z}_n(z;q)\, t^n = \frac{1}{1-t}\, {}_4\phi_5\left(\begin{matrix} -q, q^{\frac{1}{2}}, -q^{\frac{1}{2}}, 0, \\ q, qt^{\frac{1}{2}}, -qt^{\frac{1}{2}}, (qt)^{\frac{1}{2}}, -(qt)^{\frac{1}{2}} \end{matrix} ; q, zt \right), \tag{3.1}$$

$$\sum_{n=0}^{\infty} \mathscr{B}_n(z;q)\, t^n = \frac{1}{1-t}\, {}_5\phi_5\left(\begin{matrix} -q, q^{\frac{1}{2}}, -q^{\frac{1}{2}}, q^{\frac{z+1}{2}}, 0 \\ q, (qt)^{\frac{1}{2}}, -(qt)^{\frac{1}{2}}, qt^{\frac{1}{2}}, -qt^{\frac{1}{2}} \end{matrix} ; q, t \right), \tag{3.2}$$

$$\sum_{n=0}^{\infty} \Big(\mathscr{B}_n(z-2;q) - \mathscr{B}_n(z;q) \Big)\, t^n = \frac{q^{\frac{z-1}{2}}(1+q)t}{(t;q)_3}\, {}_5\phi_5\left(\begin{matrix} -q^2, q^{\frac{3}{2}}, -q^{\frac{3}{2}}, q^{\frac{z+1}{2}}, 0 \\ q^2, (q^3t)^{\frac{1}{2}}, -(q^3t)^{\frac{1}{2}}, q^2t^{\frac{1}{2}}, -q^2t^{\frac{1}{2}} \end{matrix} ; q, qt \right). \tag{3.3}$$

Proof. Let us start by proving identity (3.2), namely

$$\sum_{n=0}^{\infty} \mathscr{B}_n(z;q)\, t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k (q^{n+1};q)_k (q^{\frac{z+1}{2}};q)_k}{(q;q)_k^3} (q^k t)^n.$$

By using the identity

$$(q^{-n};q)_k = \frac{(q;q)_n}{(q;q)_{n-k}} (-1)^k q^{\frac{k(k-1)}{2} - nk}, \qquad k = 0, 1, \ldots, n,$$

we have

$$\sum_{n=0}^{\infty} \mathscr{B}_n(z;q)\, t^n = \sum_{n=0}^{\infty} \sum_{k=0}^{n} \frac{(q;q)_{n+k} (q^{\frac{z+1}{2}};q)_k}{(q;q)_{n-k} (q;q)_k^3} (-1)^k q^{\frac{k(k-1)}{2}} t^n.$$

Next, we rearrange the double summation and set $n \mapsto n+k$, obtaining

$$\begin{aligned} \sum_{n=0}^{\infty} \mathscr{B}_n(z;q)\, t^n &= \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \frac{(q;q)_{n+2k} (q^{\frac{z+1}{2}};q)_k}{(q;q)_n (q;q)_k^3} (-1)^k q^{\frac{k(k-1)}{2}} t^{n+k} \\ &= \sum_{k=0}^{\infty} \frac{(q;q)_{2k} (q^{\frac{z+1}{2}};q)_k}{(q;q)_k^3} (-t)^k q^{\frac{k(k-1)}{2}} \sum_{n=0}^{\infty} \frac{(q^{2k+1};q)_n}{(q;q)_n} t^n. \end{aligned}$$

By using the identity

$$(a;q)_{2n} = (a^{\frac{1}{2}};q)_n (-a^{\frac{1}{2}};q)_n ((aq)^{\frac{1}{2}};q)_n (-(aq)^{\frac{1}{2}};q)_n,$$

and (2.1), we obtain for the generating function

$$
\begin{aligned}
=&\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac12};q)_k(-q^{\frac12};q)_k(q^{\frac{z+1}{2}};q)_k}{(q;q)_k^2}(-t)^k q^{\frac{k(k-1)}{2}}\frac{(q^{2k+1}t;q)_\infty}{(t;q)_\infty}\\
=&\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac12};q)_k(-q^{\frac12};q)_k(q^{\frac{z+1}{2}};q)_k}{(t;q)_{2k+1}(q;q)_k^2}(-t)^k q^{\frac{k(k-1)}{2}}\\
=&\frac{1}{1-t}\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac12};q)_k(-q^{\frac12};q)_k(q^{\frac{z+1}{2}};q)_k}{(qt;q)_{2k}(q;q)_k^2}(-t)^k q^{\frac{k(k-1)}{2}}\\
=&\frac{1}{1-t}\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac12};q)_k(-q^{\frac12};q)_k(q^{\frac{z+1}{2}};q)_k}{(q;q)_k((qt)^{\frac12};q)_k(-(qt)^{\frac12};q)_k(qt^{\frac12};q)_k(-qt^{\frac12};q)_k}(-1)^k q^{\frac{k(k-1)}{2}}\frac{t^k}{(q;q)_k}.
\end{aligned}
$$

Hence the identity follows. In order to prove identity (3.3):

$$
\begin{aligned}
\sum_{n=0}^{\infty}\Big(\mathscr{B}_n(z-2;q)-\mathscr{B}_n(z;q)\Big)t^n=&\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q^{-n};q)_k(q^{n+1};q)_k}{(q;q)_k^3}q^{nk}\Big((q^{\frac{z-1}{2}};q)_k-(q^{\frac{z+1}{2}};q)_k\Big)t^n\\
=&\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q^{-n};q)_k(q^{n+1};q)_k}{(q;q)_k^3}q^{nk}(q^{\frac{z+1}{2}};q)_{k-1}(1-q^k)(-q^{\frac{z-1}{2}})t^n\\
=&\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q;q)_{n+k}(q^{\frac{z+1}{2}};q)_{k-1}(-1)^{k-1}q^{\frac{z-1}{2}}q^{\frac{k(k-1)}{2}}}{(q;q)_{n-k}(q;q)_{k-1}(q;q)_k^2}t^n.
\end{aligned}
$$

Taking into account that the above expression vanishes at $k=0$ we set $k\mapsto k+1$, rearrange the double sum, and set $n\mapsto n+k$, yielding

$$
=\sum_{k=0}^{\infty}\sum_{n=0}^{\infty}\frac{(q;q)_{n+2k+1}(q^{\frac{z+1}{2}};q)_k(-1)^k q^{\frac{z-1}{2}}q^{\frac{k(k+1)}{2}}}{(q;q)_{n-1}(q;q)_k(q;q)_{k+1}(q;q)_{k+1}}\,t^{n+k}.
$$

Here, again, the series vanishes at $n=0$ so we set $n\mapsto n+1$. Applying some basic identities of q-Pochhammer symbols, we obtain

$$
\begin{aligned}
=&(1+q)q^{\frac{z-1}{2}}t\sum_{k=0}^{\infty}\sum_{n=0}^{\infty}\frac{(q^3;q)_{n+2k}(q^{\frac{z+1}{2}};q)_k q^{\frac{k(k-1)}{2}}}{(q;q)_n(q;q)_k(q^2;q)_k(q^2;q)_k}\,(-qt)^k t^n\\
=&(1+q)q^{\frac{z-1}{2}}t\sum_{k=0}^{\infty}\frac{(q^3;q)_{2k}(q^{\frac{z+1}{2}};q)_k}{(q;q)_k(q^2;q)_k(q^2;q)_k}\,q^{\frac{k(k-1)}{2}}(-qt)^k\sum_{n=0}^{\infty}\frac{(q^{3+2k};q)_n}{(q;q)_n}\,t^n.
\end{aligned}
$$

Applying again (2.1) and simplifying one has

$$=(1+q)q^{\frac{z-1}{2}}t\sum_{k=0}^{\infty}\frac{(q^3;q)_{2k}(q^{\frac{z+1}{2}};q)_k}{(q;q)_k(q^2;q)_k(q^2;q)_k}\,q^{\frac{k(k-1)}{2}}(-qt)^k\frac{1}{(t;q)_{3+2k}}$$

$$=(1+q)q^{\frac{z-1}{2}}t\sum_{k=0}^{\infty}\frac{(q^{\frac{3}{2}};q)_k(-q^{\frac{3}{2}};q)_k(-q^2;q)_k(q^{\frac{z+1}{2}};q)_k}{(q;q)_k(q^2;q)_k}\,q^{\frac{k(k-1)}{2}}(-qt)^k\frac{1}{(t;q)_{3+2k}}$$

$$=\frac{(1+q)}{(t;q)_3}\,q^{\frac{z-1}{2}}t\sum_{k=0}^{\infty}\frac{(q^{\frac{3}{2}};q)_k(-q^{\frac{3}{2}};q)_k(-q^2;q)_k(q^{\frac{z+1}{2}};q)_kq^{\frac{k(k-1)}{2}}}{(q;q)_k(q^2;q)_k((tq^3)^{\frac{1}{2}};q)_k(-(tq^3)^{\frac{1}{2}};q)_k(q^2t^{\frac{1}{2}};q)_k(-q^2t^{\frac{1}{2}};q)_k}\,(-qt)^k.$$

Therefore the identity follows. Let us prove the generating function (3.3). We have

$$\sum_{n=0}^{\infty}\mathscr{Z}_n(z;q)\,t^n=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q^{-n};q)_k(q^{n+1};q)_k(-1)^kq^{\frac{k(k-1)}{2}}}{(q;q)_k^3}\,z^kq^{nk}t^n$$

$$=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q;q)_{n+k}\,q^{k(k-1)}}{(q;q)_{n-k}(q;q)_k^3}\,z^kt^n.$$

As in the previous identities, we rearrange the double sums and set $n\mapsto n+k$ obtaining

$$=\sum_{k=0}^{\infty}\sum_{n=0}^{\infty}\frac{(q;q)_{n+2k}\,q^{k(k-1)}}{(q;q)_n(q;q)_k^3}\,z^kt^{n+k}=\sum_{k=0}^{\infty}\frac{(q;q)_{2k}\,q^{k(k-1)}}{(q;q)_k^3}(zt)^k\sum_{n=0}^{\infty}\frac{(q^{2k+1};q)_n}{(q;q)_n}\,t^n$$

$$=\sum_{k=0}^{\infty}\frac{(q;q)_{2k}\,q^{k(k-1)}}{(q;q)_k^3}\,\frac{(zt)^k}{(t;q)_{2k+1}}=\frac{1}{1-t}\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac{1}{2}};q)_k(-q^{\frac{1}{2}};q)_k\,q^{k(k-1)}}{(q;q)_k(q;q)_k\ (tq;q)_{2k}}\,(zt)^k$$

$$=\frac{1}{1-t}\sum_{k=0}^{\infty}\frac{(-q;q)_k(q^{\frac{1}{2}};q)_k(-q^{\frac{1}{2}};q)_k\,q^{k(k-1)}}{(q;q)_k(q;q)_k((tq)^{\frac{1}{2}};q)_k(-(tq)^{\frac{1}{2}};q)_k(qt^{\frac{1}{2}};q)_k(-qt^{\frac{1}{2}};q)_k}\,(zt)^k.$$

Hence the identity follows. □

Theorem 3.2. *For any $t\in\mathbb{C}$ small enough, the following identities hold:*

$$\sum_{n=0}^{\infty}\mathscr{B}_m(-2n-1;q)\,\frac{t^n}{(q;q)_n}=\frac{1}{(t;q)_\infty}\mathscr{Z}_m(tq^{-n};q),\tag{3.4}$$

$$\sum_{n=0}^{\infty}\mathscr{B}_m^j(-2n-1-j;q)\,t^n=\frac{1}{1-t}\,{}_2\phi_2\left(\begin{matrix}q^{-m},q^{m+1}\\ q^{j+1},tq\end{matrix};q;t\right),\tag{3.5}$$

$$\sum_{n=0}^{\infty}\mathscr{B}_m^j(-2n-1-j;q)\,\frac{(q^\lambda;q)_n}{(q;q)_n}\,t^n=\frac{1}{(t;q)_\lambda}\,{}_3\phi_3\left(\begin{matrix}q^{-m},q^{m+1},q^\lambda\\ q,q^{j+1},tq^\lambda\end{matrix};q;t\right),\qquad\lambda\in\mathbb{C}.\tag{3.6}$$

Proof. Observe that if we set $\lambda = 1$ in (3.6) we obtain (3.5). Since $|q| < 1$, taking the limit $\lambda \to \infty$, setting $j = 0$, and taking into account the definition of the q-Bateman polynomials yields (3.4). Let us prove (3.6). Setting $z = -2n - 1 - j$ in the q-Pasternack polynomial, and using their basic hypergeometric expansion, we have

$$\begin{aligned}
\sum_{n=0}^{\infty} \mathscr{B}_m^j(-2n-1-j;q)\,\frac{(q^\lambda;q)_n}{(q;q)_n}\,t^n &= \sum_{n=0}^{\infty}\sum_{k=0}^{n} \frac{(q^{-m};q)_k (q^{m+1};q)_k (q^{-n};q)_k}{(q;q)_k^2 (q^{j+1};q)_k}\, q^{nk} t^n\, \frac{(q^\lambda;q)_n}{(q;q)_n} \\
&= \sum_{n=0}^{\infty}\sum_{k=0}^{n} \frac{(q^{-m};q)_k (q^{m+1};q)_k (-1)^k q^{\frac{k(k-1)}{2}} t^n (q^\lambda;q)_n}{(q;q)_k^2 (q^{j+1};q)_k (q;q)_{n-k}} \\
&= \sum_{k=0}^{\infty}\sum_{n=0}^{\infty} \frac{(q^{-m};q)_k (q^{m+1};q)_k (-t)^k q^{\frac{k(k-1)}{2}}}{(q;q)_k^2 (q^{j+1};q)_k}\, \frac{t^n (q^\lambda;q)_{n+k}}{(q;q)_n} \\
&= \frac{1}{(t;q)_{\lambda+k}} \sum_{k=0}^{\infty} \frac{(q^{-m};q)_k (q^{m+1};q)_k (q^\lambda;q)_k (-t)^k q^{\frac{k(k-1)}{2}}}{(q;q)_k (q;q)_k (q^{j+1};q)_k} \\
&= \frac{1}{(t;q)_\lambda}\, {}_3\phi_3\left(\begin{matrix} q^{-m}, q^{m+1}, q^\lambda \\ q, q^{j+1}, tq \end{matrix}; q; t\right).
\end{aligned}$$

□

Lemma 1.1 is straightforward as a limiting case of expression (3.4). The result follows by setting $t \mapsto t(1-q)$, taking the limit $q \uparrow 1$, and taking into account the limiting relations

$$\lim_{q\uparrow 1} \frac{(1-q)^n}{(q;q)_n} = \frac{1}{n!}, \qquad \lim_{q\uparrow 1} \frac{1}{(t(1-q);q)_\infty} = e^t,$$

and

$$\lim_{q\uparrow 1} {}_2\phi_2\left(\begin{matrix} q^a, q^b \\ q^c, q^d \end{matrix}; q; t(q-1)\right) = {}_2F_2\left(\begin{matrix} a, b \\ c, d \end{matrix}; t\right).$$

In fact, we obtain a new generating function for the Pasternack and Bateman polynomials by taking the same limit $q \uparrow 1$.

Corollary 3.3. *The following identities hold:*

$$\sum_{n=0}^{\infty} \mathscr{B}_m(-2n-1)\, t^n = \frac{1}{1-t}\, P_m\left(\frac{1+t}{1-t}\right),$$

$$\sum_{n=0}^{\infty} \mathscr{B}_m(-2n-1)\,\frac{(\lambda)_n}{n!}\, t^n = \frac{1}{(1-t)^\lambda}\, {}_3F_2\left(\begin{matrix} -m, m+1, \lambda \\ 1, 1 \end{matrix}; \frac{-t}{1-t}\right),$$

$$\sum_{n=0}^{\infty} \mathscr{B}_m^j(-2n-1-j)\, t^n = \frac{1}{1-t}\, {}_2F_1\left(\begin{matrix} -m, m+1 \\ j+1 \end{matrix}; \frac{-t}{1-t}\right),$$

$$\sum_{n=0}^{\infty} \mathscr{B}_m^j(-2n-1-j)\,\frac{(\lambda)_n}{n!}\, t^n = \frac{1}{(1-t)^\lambda}\, {}_3F_2\left(\begin{matrix} -m, m+1, \lambda \\ 1, j+1 \end{matrix}; \frac{-t}{1-t}\right),$$

where P_m is the Legendre polynomial.

Theorem 3.4. *For any $t \in \mathbb{C}$ small enough, the q-Pasternack polynomials satisfy the following generating function:*

$$\sum_{n=0}^{\infty} \mathscr{B}_n^m(z;q)\, t^n = \frac{1}{1-t}\, {}_5\phi_5\left(\begin{matrix} -q, q^{\frac{1}{2}}, -q^{\frac{1}{2}}, q^{\frac{z+m+1}{2}}, 0 \\ q^{m+1}, (qt)^{\frac{1}{2}}, -(qt)^{\frac{1}{2}}, qt^{\frac{1}{2}}, -qt^{\frac{1}{2}} \end{matrix} ; q, t \right). \tag{3.7}$$

Proof. Taking into account the expression of the basic hypergeometric series for these polynomials we have

$$\begin{aligned}
\sum_{n=0}^{\infty} \mathscr{B}_n^m(z;q)\, t^n &= \sum_{n=0}^{\infty} \sum_{k=0}^{n} \frac{(q^{-n};q)_k (q^{n+1};q)_k (q^{\frac{z+m+1}{2}};q)_k}{(q;q)_k^2 (q^{m+1};q)_k} \left(q^k t\right)^n \\
&= \sum_{n=0}^{\infty} \sum_{k=0}^{n} \frac{(q;q)_{n+k} (q^{\frac{z+m+1}{2}};q)_k}{(q;q)_{n-k} (q;q)_k^2 (q^{m+1};q)_k} (-1)^k q^{\frac{k(k-1)}{2}} t^n \\
&= \sum_{k=0}^{\infty} \sum_{n=0}^{\infty} \frac{(q;q)_{n+2k} (q^{\frac{z+m+1}{2}};q)_k}{(q;q)_n (q;q)_k^2 (q^{m+1};q)_k} (-1)^k q^{\frac{k(k-1)}{2}} t^{n+k} \\
&= \sum_{k=0}^{\infty} \frac{(q;q)_{2k} (q^{\frac{z+m+1}{2}};q)_k}{(q;q)_k^2 (q^{m+1};q)_k} (-1)^k q^{\frac{k(k-1)}{2}}\, t^k \frac{(q^{2k+1}t;q)_\infty}{(t;q)_\infty} \\
&= \frac{1}{1-t} \sum_{k=0}^{\infty} \frac{(q;q)_{2k} (q^{\frac{z+m+1}{2}};q)_k}{(q;q)_k^2 (q^{m+1};q)_k (qt;q)_{2k}} (-1)^k q^{\frac{k(k-1)}{2}}\, t^k \\
&= \frac{1}{1-t} \sum_{k=0}^{\infty} \frac{(q^{\frac{1}{2}};q)_k (-q^{\frac{1}{2}};q)_k, (-q;q)_k (q^{\frac{z+m+1}{2}};q)_k\, (-1)^k q^{\frac{k(k-1)}{2}}}{(q;q)_k (q^{m+1};q)_k ((qt)^{\frac{1}{2}};q)_k (-(qt)^{\frac{1}{2}};q)_k (qt^{\frac{1}{2}};q)_k (-qt^{\frac{1}{2}};q)_k}\, t^k.
\end{aligned}$$

□

Observe that we obtain (3.2) by setting $m = 0$ in (3.7).

Theorem 3.5. *The q-Sylvester polynomials satisfy the following generating functions:*

$$\sum_{n=0}^{\infty} \varphi_n(z;q)\, t^n = \frac{(q^z t;q)_\infty}{(t, zt;q)_\infty}, \tag{3.8}$$

$$\sum_{n=0}^{\infty} (q^\lambda;q)_n\, \varphi_n(z;q)\, t^n = \frac{1}{(zt;q)_\lambda}\, {}_2\phi_1\left(\begin{matrix} q^\lambda, q^z \\ ztq^\lambda \end{matrix} ; q, t \right), \tag{3.9}$$

where the principal values of q^z and q^λ are taken.

Proof. Let us prove the generating function (3.9) by using an analogous method as before, namely

$$\sum_{n=0}^{\infty}(q^{\lambda};q)_n\,\varphi_n(z;q)\,t^n=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q^{-n};q)_k(q^z;q)_k(-1)^k q^{-\frac{k(k-1)}{2}}(q^{\lambda};q)_n}{(q;q)_n(q;q)_k}\,q^{nk}z^{n-k}t^n$$

$$=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\frac{(q^z;q)_k(q^{\lambda};q)_n}{(q;q)_{n-k}(q;q)_k}\,z^{n-k}t^n=\sum_{k=0}^{\infty}\sum_{n=0}^{\infty}\frac{(q^z;q)_k(q^{\lambda};q)_{n+k}}{(q;q)_n(q;q)_k}\,z^n t^{n+k}$$

$$=\sum_{k=0}^{\infty}\frac{(q^z;q)_k(q^{\lambda};q)_k}{(q;q)_k}\,t^k\sum_{n=0}^{\infty}\frac{(q^{\lambda+k};q)_n(zt)^n}{(q;q)_n}=\sum_{k=0}^{\infty}\frac{(q^z;q)_k(q^{\lambda};q)_k}{(q;q)_k(zt;q)_{\lambda+k}}\,t^k$$

$$=\frac{1}{(zt;q)_{\lambda}}\sum_{k=0}^{\infty}\frac{(q^z;q)_k(q^{\lambda};q)_k}{(q;q)_k(ztq^{\lambda};q)_k}\,t^k.$$

Since $|q|<1$, (3.8) follows from taking $\lambda\to\infty$ and applying (2.1), both generating functions hold. □

Theorem 3.6. *For any $t\in\mathbb{C}$ small enough, the q-Cesàro polynomials satisfy the following generating function:*

$$\sum_{n=0}^{\infty}g_n^{(s)}(z;q)\,t^n=\frac{1}{(1-tzq^s)(t;q)_{s+1}}. \tag{3.10}$$

Proof. Let us prove this by using (2.3) and some basic properties of the q-Pochhammer symbol:

$$\sum_{n=0}^{\infty}g_n^{(s)}(z;q)\,t^n=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\begin{bmatrix}k+s\\ s\end{bmatrix}_q(qz^s)^{n-k}t^n=\sum_{n=0}^{\infty}\sum_{k=0}^{n}\begin{bmatrix}n-k+s\\ s\end{bmatrix}_q(qz^s)^k t^n$$

$$=\sum_{k=0}^{\infty}\sum_{n=0}^{\infty}\begin{bmatrix}n+s\\ s\end{bmatrix}_q(tqz^s)^k t^n=\frac{1}{(1-tzq^s)(t;q)_{s+1}}.$$

□

We leave it to the reader to check that by taking limit $q\uparrow 1$, we recover the original expressions by using the limit identities we gave before Theorem 3.4. In the future, it would be interesting to see if it is possible to use q-calculus to obtain q-analogues of the results obtained in [**1**].

Acknowledgments

The author R. S. Costas-Santos acknowledges financial support by Dirección General de Investigación, Ministerio de Economía y Competitividad of Spain, grant MTM2015-65888-C4-2-P.

References

[1] Bateman, H. Two systems of polynomials for the solution of Laplace integral equation. *Duke Math. Journal* **2** *(1936), 569–577.*

[2] Bateman, H. *Some properties of a certain set of polynomials. Tôhuko Math. Journal* **37** *(1933), 23–38.*

[3] Gasper, G.; Rahman, M. *Basic Hypergeometric Series.* Cambridge University Press, Cambridge, 2004.

[4] Koekoek, R.; Lesky, P. A; Swarttouw, R. F. *Hypergeometric orthogonal polynomials and their q-analogues.* Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2010. With a foreword by Tom H. Koornwinder.

[5] Koelink, H. T. On Jacobi and continuous Hahn polynomials. *Proc. Amer. Math. Soc.* **124***(3) (1996), 887–898.*

[6] Pasternack, S. A generalisation of the polynomial $F_n(x)$. *London, Edinburgh, Dublin Philosophical Magazine and J. Science* **28***, ser. 7 (1939), 209–226.*

[7] Rainville, E. D. *Special Functions.* Macmillan, New York, 1960.

[8] Srivastava, H. M.; Manocha, H. L. *A Treatise on Generating Functions.* Wiley, New York, 1984.

Howard S. Cohl
Applied and Computational Mathematics Division
National Institute of Standards and Technology
Mission Viejo, CA 92694
USA
howard.cohl@nist.gov

Roberto S. Costas-Santos
Dpto. Física y Matemáticas
Universidad de Alcalá
Alcalá de Henares, Madrid
SPAIN
rscosa@gmail.com

Tanay V. Wakhare
University of Maryland
College Park, MD 20742
USA
twakhare@gmail.com