

ROTA-BAXTER BISYSTEMS AND COVARIANT BIALGEBRAS

TIANSHUI MA, ABDENACER MAKHLOUF, AND SERGEI SILVESTROV

ABSTRACT. The aim of this paper is first to introduce and study Rota-Baxter cosystems and bisystems as generalization of Rota-Baxter coalgebras and bialgebras, respectively, with various examples. The second purpose is to provide an alternative definition of covariant bialgebras via coderivations. Furthermore, we consider coquasitriangular covariant bialgebras which are generalization of coquasitriangular infinitesimal bialgebras, coassociative Yang-Baxter pairs, coassociative Yang-Baxter equation, double coalgebras and dendriform coalgebras. We study some of their properties and relationships with Rota-Baxter cosystems and bisystems.

CONTENTS

1. Introduction	1
2. Preliminaries	2
3. Rota-Baxter cosystems, Rota-Baxter bisystems and Rota-Baxter bialgebras	3
4. Covariant bialgebras	10
5. Coquasitriangular covariant bialgebras and infinitesimal bialgebras	14
6. Examples of Rota-Baxter bialgebras and bisystems	22
6.1. 2-dimensional Examples	22
6.2. 3-dimensional and 4-dimensional Examples	24
References	29

1. INTRODUCTION

Rota-Baxter algebras were introduced in [19] in the context of differential operators on commutative Banach algebras and then led to an algebraic interpretation of Spitzer identity in fluctuation theory in probability and combinatorics [3]. Since then, they were intensively studied and appeared in various areas such as mathematical physics, mainly in Connes-Kreimer renormalization theory in quantum field theory (see [7, 8, 9, 12, 13]), Lie algebras (see [2]), multiple zeta functions (see [7, 15]), differential algebras (see [14]). One can refer to the book [11] for the detailed theory of Rota-Baxter algebras. In 2014, based on the dual method in the Hopf algebra theory, Jian and Zhang defined in [16] the notion of Rota-Baxter coalgebras and also provided

Date: August 17, 2017.

1991 Mathematics Subject Classification. 16T05, 16W99.

Key words and phrases. Rota-Baxter cosystem; Rota-Baxter bisystem; Covariant bialgebra; Yang-Baxter equation; Coquasitriangular covariant bialgebra; Coquasitriangular infinitesimal bialgebra.

Corresponding author: Sergei Silvestrov, sergei.silvestrov@mdh.se.

various examples. Therefore Rota-Baxter bialgebras were presented in [17] with examples constructed using Radford biproduct. In [4], T. Brzeziński introduced the notion of Rota-Baxter system, where Rota-Baxter algebras can be viewed as special case. Following the above idea, we aim in this paper to study the dual version and introduce Rota-Baxter co-(bi-)systems.

An associative (not necessarily unital) algebra A that admits a coassociative comultiplication which is a derivation is called infinitesimal bialgebra, see [1]. It is self-dual and can be viewed as a coalgebra that admits an associative multiplication which is a coderivation. In [4], T. Brzeziński proposed the concept of covariant bialgebra via two derivations. It is an extension of infinitesimal bialgebras and quasitriangular covariant bialgebras obtained by Yang-Baxter pairs. It turns out that covariant bialgebras are not self-dual. We provide another approach to study covariant bialgebras through two coderivations, and at the same time study similarly coquasitriangular infinitesimal bialgebras.

This paper is organized as follows. In Section 2, we review some preliminaries and in Section 3, we discuss the notions of Rota-Baxter cosystem, bisystem and relate them to Rota-Baxter coalgebras, pre-Lie coalgebras, weak copseudotwistors and Rota-Baxter bialgebras. Section 4 is devoted to study covariant bialgebras via coderivations. Firstly, we introduce the notion of coassociative Yang-Baxter pair (abbr. CYBP), and then investigate their relationships with Rota-Baxter cosystems and bisystems. Secondly, we give an alternative definition of covariant bialgebras through two coderivations and provide a characterization. In Section 5, as a special case, coquasitriangular covariant bialgebras are considered. They are a generalization coquasitriangular infinitesimal bialgebras. On the other hand, we introduce the concept of right (left) covariant modules. Furthermore, we consider coassociative Yang-Baxter equation, double coalgebras and dendriform coalgebras. We study some of their properties and relationships with cosystems and bisystems. In Section 6, we provide various examples of Rota-Baxter bialgebras and bisystems in dimensions 2, 3 and 4.

2. PRELIMINARIES

Throughout this paper, we assume that all vector spaces, algebras, coalgebras and tensor products are defined over a field K . An algebra is always assumed to be associative, but not necessarily unital. A coalgebra is always assumed to be coassociative, but not necessarily counital. Now, for a given a coalgebra C , we use Sweedler's notation for the comultiplication : $\Delta(c) = c_1 \otimes c_2$ for any $c \in C$, see [18]. We denote the category of left H -comodules (resp. right H -comodules) by ${}^H\text{Mod}$ (resp. Mod^H). For $(M, \rho_l) \in {}^H\text{Mod}$, we write: $\rho_l(x) = x_{(-1)} \otimes x_{(0)} \in H \otimes M$, for all $x \in M$. For $(M, \rho_r) \in \text{Mod}^H$, we write: $\rho_r(x) = x_{(0)} \otimes x_{(1)} \in M \otimes H$, for all $x \in M$. Given a K -space M , we write id_M for the identity map on M . We say that a coalgebra C is **non-degenerate** if provided with a linear map $f : C \rightarrow C$ such that for $\forall c \in C$, $c_1 \otimes f(c_2) = 0$ or $f(c_1) \otimes c_2 = 0$, it implies that $f = 0$. Obviously, any counital coalgebra is non-degenerate.

In what follows, we recall some useful definitions which will be used later (see [4, 11, 17]).

Definition 2.1. For a given field K and $\lambda \in K$, a **Rota-Baxter algebra of weight λ** is an algebra A together with a K -linear map $R : A \rightarrow A$ such that

$$R(a)R(b) = R(aR(b)) + R(R(a)b) + \lambda R(ab) \quad (2.1)$$

for all $a, b \in A$. Such a linear operator is called a **Rota-Baxter operator of weight λ on A** .

Definition 2.2. Let γ be an element in K . A pair (C, Q) is called a **Rota-Baxter coalgebra of weight γ** if C is a coalgebra and Q is a linear endomorphism of C satisfying

$$(Q \otimes Q)\Delta = (id \otimes Q)\Delta Q + (Q \otimes id)\Delta Q + \gamma \Delta Q,$$

that is, for all $c \in C$,

$$Q(c_1) \otimes Q(c_2) = Q(c)_1 \otimes Q(Q(c)_2) + Q(Q(c)_1) \otimes Q(c)_2 + \gamma Q(c)_1 \otimes Q(c)_2. \quad (2.2)$$

The map Q is called a **Rota-Baxter operator of weight γ on C** .

Definition 2.3. Let λ, γ be elements in K and H a bialgebra (not necessarily unital and counital). A triple (H, R, Q) is called a **Rota-Baxter bialgebra of weight (λ, γ)** if (H, R) is a Rota-Baxter algebra of weight λ and (H, Q) is a Rota-Baxter coalgebra of weight γ . We say that we have a (R, Q) -Rota-Baxter structure of weight (λ, γ) on the bialgebra H .

Definition 2.4. A triple (A, R, S) consisting of an algebra A and two K -linear operators $R, S : A \rightarrow A$ is called a **Rota-Baxter system** if, for all $a, b \in A$,

$$R(a)R(b) = R(R(a)b + aS(b)), \quad (2.3)$$

$$S(a)S(b) = S(R(a)b + aS(b)). \quad (2.4)$$

Definition 2.5. A **pre-Lie coalgebra** is a pair (C, Δ) consisting of a linear space C and a linear map $\Delta : C \rightarrow C \otimes C$ satisfying

$$\Delta_C - \Phi_{(12)}\Delta_C = 0, \quad (2.5)$$

where $\Delta_C = (\Delta \otimes id)\Delta - (id \otimes \Delta)\Delta$ and $\Phi_{(12)}(c_1 \otimes c_2 \otimes c_3) = c_2 \otimes c_1 \otimes c_3$.

3. ROTA-BAXTER COSYSTEMS, ROTA-BAXTER BISYSTEMS AND ROTA-BAXTER BIALGEBRAS

In this section we introduce Rota-Baxter cosystems, bisystems and relate them to Rota-Baxter coalgebras, bialgebras and pre-Lie coalgebras.

Definition 3.1. A triple (C, Q, T) consisting of a coalgebra C and two K -linear operators $Q, T : C \rightarrow C$ is called a **Rota-Baxter cosystem** if, for all $c \in C$,

$$Q(c_1) \otimes Q(c_2) = Q(Q(c)_1) \otimes Q(c)_2 + Q(c)_1 \otimes T(Q(c)_2), \quad (3.1)$$

$$T(c_1) \otimes T(c_2) = Q(T(c)_1) \otimes T(c)_2 + T(c)_1 \otimes T(T(c)_2). \quad (3.2)$$

A **morphism of Rota-Baxter cosystems between (C, Q_C, T_C) and (D, Q_D, T_D)** is a coalgebra map $f : C \rightarrow D$ such that $f \circ Q_C = Q_D \circ f$ and $f \circ T_C = T_D \circ f$.

Example 3.2. Let C be a coassociative coalgebra, $g : C \rightarrow C$ a comultiplicative map and $Q : C \rightarrow C$ a K -linear map such that for all $c \in C$,

$$Q(c_1) \otimes Q(c_2) = Q(Q(c)_1) \otimes Q(c)_2 + Q(c)_1 \otimes Q^g(Q(c)_2), \quad (3.3)$$

where $Q^g = Q \circ g$. Then (C, Q, Q^g) is a Rota-Baxter cosystem.

Proof. Eq.(3.3) is exactly Eq.(3.1) for (Q, Q^g) and Eq.(3.2) can be checked by Eq.(3.3) and $g : C \rightarrow C$ is a comultiplicative map. \square

The following proposition states that Rota-Baxter coalgebra can be viewed as a special case of Rota-Baxter cosystem.

Proposition 3.3. *Let C be a coalgebra. If (C, Q) is a Rota-Baxter coalgebra of weight γ , then $(C, Q, Q + \gamma id)$ and $(C, Q + \gamma id, Q)$ are Rota-Baxter cosystems.*

Proof. We only check that $(C, Q, Q + \gamma id)$ is a Rota-Baxter cosystem as follows:

$$\begin{aligned}
\text{RHS of Eq.(3.2)} &\stackrel{T=Q+\gamma id}{=} Q(Q(c)_1) \otimes Q(c)_2 + \gamma Q(c)_1 \otimes c_2 + Q(c)_1 \otimes Q(Q(c)_2) \\
&\quad + \gamma Q(c)_1 \otimes Q(c)_2 + \gamma c_1 \otimes Q(c_2) + \gamma^2 c_1 \otimes c_2 \\
&\stackrel{(2.2)}{=} Q(c_1) \otimes Q(c_2) + \gamma Q(c)_1 \otimes c_2 + \gamma c_1 \otimes Q(c_2) + \gamma^2 c_1 \otimes c_2 \\
&= (Q(c_1) + \gamma c_1) \otimes (Q(c_2) + \gamma c_2) \\
&= \text{LHS of Eq.(3.2)}.
\end{aligned}$$

It is obvious that Eq.(3.1) holds for $(C, Q, Q + \gamma id)$ by using Eq.(2.2). The proof for $(C, Q + \gamma id, Q)$ is similar. \square

Proposition 3.4. *Let C be a coalgebra, $Q : C \rightarrow C$ a left C -colinear map and let $T : C \rightarrow C$ be a right C -colinear map. Then (C, Q, T) is a Rota-Baxter cosystem if and only if, for all $c \in C$,*

$$c_1 \otimes (T \circ Q)(c_2) = (Q \circ T)(c_1) \otimes c_2 = 0. \quad (3.4)$$

In particular, if C is a non-degenerate coalgebra, then (C, Q, T) is a Rota-Baxter cosystem (with Q left and T right C -colinear) if and only if Q and T satisfy the orthogonality condition

$$T \circ Q = Q \circ T = 0. \quad (3.5)$$

Proof. Q is left C -colinear and T is right C -colinear, that is, $Q(c)_1 \otimes Q(c)_2 = c_1 \otimes Q(c_2)$ and $T(c)_1 \otimes T(c)_2 = T(c_1) \otimes c_2$. Thus Eq.(3.1) holds if and only if $c_1 \otimes (T \circ Q)(c_2) = 0$ and Eq.(3.2) holds if and only if $(Q \circ T)(c_1) \otimes c_2 = 0$. Since C is non-degenerate $Q \circ T = 0$. \square

Example 3.5. *Let C be a coalgebra. Assume that $\sigma, \tau \in C^*$ are such that for $\forall c \in C$, $\sigma(c_1)\tau(c_2) = 0$ and $\sigma(c_1)c_2 = c_1\sigma(c_2)$. Define $Q, T : C \rightarrow C$ by*

$$Q(c) = c_1\sigma(c_2), \quad T(c) = \tau(c_1)c_2$$

for all $c \in C$. Then it is obvious that Q is left C -colinear and T is right C -colinear since the comodule action is the coproduct in C . While

$$c_1 \otimes (T \circ Q)(c_2) = c_1 \otimes \tau(c_2)c_3\sigma(c_4) = c_1 \otimes \sigma(c_2)\tau(c_3)c_4 = 0$$

and

$$(Q \circ T)(c_1) \otimes c_2 = \tau(c_1)c_2\sigma(c_3) \otimes c_4 = \sigma(c_1)\tau(c_2)c_3 \otimes c_4 = 0.$$

Therefore (C, Q, T) is a Rota-Baxter cosystem.

Example 3.6. We construct Rota-Baxter bisystems for T_2 , the unital Taft-Sweedler algebra generated by g, x and the relations ($g^2 = 1$, $x^2 = 0$, $xg = -gx$). The comultiplication is defined by $\Delta(g) = g \otimes g$ and $\Delta(x) = x \otimes 1 + g \otimes x$, the counit is given by $\varepsilon(g) = 1$, $\varepsilon(x) = 0$.

The bialgebra T_2 is 4-dimensional, it is defined with respect to a basis $\{u_1 = 1, u_2 = g, u_3 = x, u_4 = gx\}$ by the following table which describes multiplying the i th row elements by the j th column elements.

	u_1	u_2	u_3	u_4
u_1	u_1	u_2	u_3	u_4
u_2	u_2	u_1	u_4	u_3
u_3	u_3	$-u_4$	0	0
u_4	u_4	$-u_3$	0	0

and

$$\Delta(u_1) = u_1 \otimes u_1, \Delta(u_2) = u_2 \otimes u_2, \Delta(u_3) = u_1 \otimes u_3 + u_3 \otimes u_2, \Delta(u_4) = u_2 \otimes u_4 + u_4 \otimes u_1.$$

$$\varepsilon(u_1) = \varepsilon(u_2) = 1, \quad \varepsilon(u_3) = \varepsilon(u_4) = 0.$$

A bisystem is given by a pair (R, S) and a pair (Q, T) taken from the following list for (R, S)

- $R(u_1) = 0, R(u_2) = 0, R(u_3) = 0, R(u_4) = pu_4,$
 $S(u_1) = -pu_1, S(u_2) = 0, S(u_3) = 0, S(u_4) = 0,$
- $R(u_1) = 0, R(u_2) = 0, R(u_3) = -pu_3, R(u_4) = -pu_4,$
 $S(u_1) = pu_1, S(u_2) = pu_2, S(u_3) = 0, S(u_4) = 0,$
- $R(u_1) = 0, R(u_2) = 0, R(u_3) = pu_3, R(u_4) = 0,$
 $S(u_1) = -pu_1, S(u_2) = 0, S(u_3) = 0, S(u_4) = 0,$
- $R(u_1) = -pu_1, R(u_2) = 0, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = 0, S(u_2) = 0, S(u_3) = pu_3, S(u_4) = 0,$
- $R(u_1) = -pu_1, R(u_2) = -pu_2, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = 0, S(u_2) = 0, S(u_3) = pu_3, S(u_4) = pu_3,$
- $R(u_1) = -pu_1, R(u_2) = -pu_2, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = 0, S(u_2) = 0, S(u_3) = pu_3, S(u_4) = -pu_3,$
- $R(u_1) = -pu_1, R(u_2) = 0, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = 0, S(u_2) = 0, S(u_3) = 0, S(u_4) = pu_4,$
- $R(u_1) = pu_1, R(u_2) = 0, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = 0, S(u_2) = 0, S(u_3) = 0, S(u_4) = 0,$

where p is a parameter, and the following list for (Q, T)

- $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = 0, Q(u_4) = -q_1u_4,$
 $T(u_1) = q_1u_1, T(u_2) = q_2u_2, T(u_3) = q_2u_3, T(u_4) = 0,$
- $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = 0, Q(u_4) = -q_1u_4,$
 $T(u_1) = q_1u_1, T(u_2) = q_2u_2, T(u_3) = 0, T(u_4) = 0,$

- $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = -q_1u_3, Q(u_4) = 0,$
 $T(u_1) = q_1u_1, T(u_2) = q_2u_2, T(u_3) = 0, T(u_4) = q_2u_4,$
- $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = -q_1u_3, Q(u_4) = 0,$
 $T(u_1) = q_1u_1, T(u_2) = q_2u_2, T(u_3) = 0, T(u_4) = 0,$
- $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = -q_1u_3, Q(u_4) = -q_1u_4,$
 $T(u_1) = q_1u_1, T(u_2) = q_2u_2, T(u_3) = 0, T(u_4) = 0,$
- $Q(u_1) = 0, Q(u_2) = -q_1u_2, Q(u_3) = 0, Q(u_4) = 0,$
 $T(u_1) = q_1u_1, T(u_2) = 0, T(u_3) = q_2u_3, T(u_4) = 0,$
- $Q(u_1) = 0, Q(u_2) = -q_1u_2, Q(u_3) = 0, Q(u_4) = 0,$
 $T(u_1) = q_2u_1, T(u_2) = 0, T(u_3) = q_1u_3, T(u_4) = q_1u_4,$
- $Q(u_1) = 0, Q(u_2) = -q_1u_2, Q(u_3) = 0, Q(u_4) = -q_2u_4,$
 $T(u_1) = q_2u_1, T(u_2) = 0, T(u_3) = q_1u_1, T(u_4) = 0,$

where q_1, q_2 are parameters.

Proposition 3.7. *Let (C, Q, T) be a Rota-Baxter cosystem. Then*

(1) (C, Δ_*) with $\Delta_* : C \rightarrow C \otimes C$, defined by

$$\Delta_*(c) = Q(c_1) \otimes c_2 + c_1 \otimes T(c_2) \quad (3.6)$$

for all $c \in C$, is a coassociative coalgebra.

(2) (C, Δ_\bullet) with $\Delta_\bullet : C \rightarrow C \otimes C$, defined by

$$\Delta_\bullet(c) = Q(c_1) \otimes c_2 - T(c_2) \otimes c_1 \quad (3.7)$$

for all $c \in C$, is a pre-Lie coalgebra.

Proof. (1) The coassociativity for (C, Δ_*) can be proved as follows. For all $c \in C$, we have

$$\begin{aligned}
(\Delta_* \otimes id) \circ \Delta_*(c) &= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 + Q(c_1)_1 \otimes T(Q(c_1)_2) \otimes c_2 \\
&\quad + Q(c_{11}) \otimes c_{12} \otimes T(c_2) + c_{11} \otimes T(c_{12} \otimes T(c_2)) \\
&\stackrel{(3.2)}{=} Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 + Q(c_1)_1 \otimes T(Q(c_1)_2) \otimes c_2 \\
&\quad + Q(c_{11}) \otimes c_{12} \otimes T(c_2) + c_1 \otimes Q(T(c_2)_1) \otimes T(c_2)_2 \\
&\quad + c_1 \otimes T(c_2)_1 \otimes T(T(c_2)_2) \\
&\stackrel{(3.1)}{=} Q(c_1) \otimes Q(c_{21}) \otimes c_{22} + Q(c_1) \otimes c_{21} \otimes T(c_{22}) \\
&\quad + c_1 \otimes Q(T(c_2)_1) \otimes T(c_2)_2 + c_1 \otimes T(c_2)_1 \otimes T(T(c_2)_2) \\
&= (id \otimes \Delta_*) \circ \Delta_*(c).
\end{aligned}$$

(2) Since

$$\begin{aligned}
\Delta_C(c) &= ((\Delta_\bullet \otimes id) \circ \Delta_\bullet - (id \otimes \Delta_\bullet) \circ \Delta_\bullet)(c) \\
&= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 - T(Q(c_1)_2) \otimes Q(c_1)_1 \otimes c_2 \\
&\quad - Q(T(c_2)_1) \otimes T(c_2)_2 \otimes c_1 + T(T(c_2)_2) \otimes T(c_2)_1 \otimes c_1
\end{aligned}$$

$$\begin{aligned}
 & -Q(c_1) \otimes Q(c_{21}) \otimes c_{22} + Q(c_1) \otimes T(c_{22}) \otimes c_{21} \\
 & +T(c_2) \otimes Q(c_{11}) \otimes c_{12} - T(c_2) \otimes T(c_{12}) \otimes c_{11} \\
 & \stackrel{(3.1)(3.2)}{=} -T(Q(c_1)_2) \otimes Q(c_1)_1 \otimes c_2 - Q(T(c_2)_1) \otimes T(c_2)_2 \otimes c_1 \\
 & -Q(c_1)_1 \otimes T(Q(c_1)_2) \otimes c_2 - Q(c_1) \otimes T(c_{22}) \otimes c_{21} \\
 & +T(c_2) \otimes Q(c_{11}) \otimes c_{12} - T(c_2)_2 \otimes Q(T(c_2)_1) \otimes c_1,
 \end{aligned}$$

we can get $(\Delta_C - \Phi_{(12)}\Delta_C)(c) = 0$. Therefore (C, Δ_\bullet) is a pre-Lie coalgebra. \square

Remark 3.8. Here the pre-Lie coalgebra is dual to pre-Lie algebra in [2].

Next we introduce the notion of weak copseudotwistor and relate it to Rota-Baxter cosystem.

Definition 3.9. Let C be a coalgebra with coassociative coproduct $\Delta : C \longrightarrow C \otimes C$. A K -linear map $F : C \otimes C \longrightarrow C \otimes C$ is called a **weak copseudotwistor** if there exists a K -linear map $\tilde{F} : C \otimes C \otimes C \longrightarrow C \otimes C \otimes C$, rendering commutative the following diagram:

$$\begin{array}{ccccc}
 & C \otimes C \otimes C & \xleftarrow{id \otimes \Delta} & C \otimes C & \xrightarrow{\Delta \otimes id} & C \otimes C \otimes C \\
 id \otimes F \swarrow & & & \uparrow F & & \searrow F \otimes id \\
 C \otimes C \otimes C & & & & & C \otimes C \otimes C \\
 \tilde{F} \swarrow & & & & & \searrow \tilde{F} \\
 & C \otimes C \otimes C & \xleftarrow{id \otimes \Delta} & C \otimes C & \xrightarrow{\Delta \otimes id} & C \otimes C \otimes C
 \end{array} \tag{3.8}$$

The map \tilde{F} is called a **weak companion of F** .

Proposition 3.10. If (C, Q, T) is a Rota-Baxter cosystem, then

$$F : C \otimes C \longrightarrow C \otimes C, \quad c \otimes d \mapsto Q(c) \otimes d + c \otimes T(d) \tag{3.9}$$

is a weak copseudotwistor.

Proof. Define K -linear map $\tilde{F} : C \otimes C \otimes C \longrightarrow C \otimes C \otimes C$,

$$\tilde{F}(c \otimes d \otimes e) = Q(c) \otimes Q(d) \otimes e + Q(c) \otimes d \otimes T(e) + c \otimes T(d) \otimes T(e), \tag{3.10}$$

where $c, d, e \in C$. Then we obtain

$$\begin{aligned}
 (\tilde{F} \circ (\Delta \otimes id))(c \otimes d) &= Q(c_1) \otimes Q(c_2) \otimes d + Q(c_1) \otimes c_2 \otimes T(d) + c_1 \otimes T(c_2) \otimes T(d) \\
 &\stackrel{(3.1)}{=} Q(Q(c)_1) \otimes Q(c)_2 \otimes d + Q(c)_1 \otimes T(Q(c)_2) \otimes d \\
 &\quad + Q(c_1) \otimes c_2 \otimes T(d) + c_1 \otimes T(c_2) \otimes T(d) \\
 &= ((F \otimes id) \circ (\Delta \otimes id) \circ F)(c \otimes d)
 \end{aligned}$$

and

$$\begin{aligned}
(\tilde{F} \circ (id \otimes \Delta))(c \otimes d) &= Q(c) \otimes Q(d_1) \otimes d_2 + Q(c) \otimes d_1 \otimes T(d_2) + c \otimes T(d_1) \otimes T(d_2) \\
&\stackrel{(3.1)}{=} Q(c) \otimes Q(d_1) \otimes d_2 + Q(c) \otimes d_1 \otimes T(d_2) \\
&\quad + c \otimes Q(T(d)_1) \otimes T(d)_2 + c \otimes T(d)_1 \otimes T(T(d)_2) \\
&= ((id \otimes F) \circ (id \otimes \Delta) \circ F)(c \otimes d).
\end{aligned}$$

Thus Eq.(3.8) holds. This finishes the proof. \square

Remark 3.11. (1) The coproduct Δ_* defined by Eq.(3.6) is simply equal to $F \circ \Delta$, where F is given in Eq.(3.9).

(2) If C is a coalgebra satisfying that for a map $f : C \rightarrow C$ such that for all $c, d \in C$, $c \otimes f(d) = f(c) \otimes d = 0$ implies that $f = 0$, then F given in Eq.(3.9) is a weak copseudotwistor with the companion defined in Eq.(3.10) if and only if (C, Q, T) is a Rota-Baxter cosystem.

Definition 3.12. A quintuple (H, R, S, Q, T) consisting of a bialgebra H and four K -linear maps $R, S, Q, T : H \rightarrow H$ is called a **Rota-Baxter bisystem** if (H, R, S) is a Rota-Baxter system and (H, Q, T) is a Rota-Baxter cosystem.

Theorem 3.13. Let $\lambda, \gamma \in K$ and H a bialgebra (not necessarily unital and counital). If (H, R, Q) is a Rota-Baxter bialgebra of weight (λ, γ) , then $(H, R, R + \lambda id, Q, Q + \gamma id)$, $(H, R + \lambda id, R, Q, Q + \gamma id)$, $(H, R, R + \lambda id, Q + \gamma id, Q)$, $(H, R + \lambda id, R, Q + \gamma id, Q)$ are Rota-Baxter bisystems.

Proof. It is obvious by [4, Lemma 2.2] and Proposition 3.3. \square

Proposition 3.14. Let H be a bialgebra, $R : H \rightarrow H$ a left H -linear map, $S : H \rightarrow H$ a right H -linear map, $Q : H \rightarrow H$ a left H -colinear map and $T : H \rightarrow H$ a right H -colinear map. Then (H, R, S, Q, T) is a Rota-Baxter bisystem if and only if, for all $x, y \in H$,

$$a(R \circ S)(b) = 0 = (S \circ R)(a)b$$

and

$$x_1 \otimes (T \circ Q)(x_2) = (Q \circ T)(x_1) \otimes x_2 = 0.$$

In particular, if H is non-degenerate as algebra and as coalgebra, respectively, then (H, R, S, Q, T) is a Rota-Baxter bisystem (with R left and S right H -linear, Q left and T right H -colinear) if and only if R and S , Q and T satisfy the orthogonality condition, respectively,

$$R \circ S = S \circ R = 0, \quad T \circ Q = Q \circ T = 0.$$

Proof. It is obvious by [4, Lemma 2.3] and Proposition 3.4. \square

Example 3.15. Let C be a bialgebra and H a Hopf algebra with the antipode S . Suppose that there are two bialgebra maps: $i : H \rightarrow C$ and $\pi : C \rightarrow H$ such that $\pi \circ i = id_H$, i.e., C is a

bialgebra with a projection (see [18]). Set $\Pi = \text{id}_C \star (i \circ S \circ \pi)$, where \star is the convolution product on $\text{End}(C)$. The right H -Hopf module structure is given by the following:

$$c \cdot h = ci(h),$$

$$\rho_R(c) = c_1 \otimes \pi(c_2),$$

for all $c \in C$ and $h \in H$. Then (C, Π, Π) is a Rota-Baxter bialgebra of weight $(-1, -1)$, and $(C, \Pi, \Pi - \text{id}, \Pi, \Pi - \text{id})$, $(C, \Pi - \text{id}, \Pi, \Pi, \Pi - \text{id})$, $(C, \Pi, \Pi - \text{id}, \Pi - \text{id}, \Pi)$, $(C, \Pi - \text{id}, \Pi, \Pi - \text{id}, \Pi)$ are Rota-Baxter bisystems.

Proof. It is a direct consequence by [17, Example 5.3] and Theorem 3.13. \square

Example 3.16. Let H be a bialgebra, $f : H \rightarrow H$ a multiplicative map, $g : H \rightarrow H$ a comultiplicative map and $R : C \rightarrow C$ a K -linear map such that for all $x, y \in H$,

$$R(x)R(y) = R(R(x)y + xR^f(y))$$

where $R^f = R \circ f$ and $Q : H \rightarrow H$ a K -linear map such that for all $x \in H$, (3.3) holds. Then (H, R, R^f, Q, Q^g) is a Rota-Baxter bisystem.

Proof. It is a direct consequence by [4, Lemma 4.2] and Example 3.2. \square

Example 3.17. Let $\{x, y, z\}$ be a basis of a 3-dimensional vector space H over K . We define multiplication and comultiplication over H :

$$x^2 = x, \quad y^2 = y, \quad z^2 = 0$$

$$xy = yx = y, \quad yz = z,$$

$$xz = zx = z, \quad zy = 0,$$

$$\Delta(x) = x \otimes x, \quad \Delta(y) = y \otimes y, \quad \Delta(z) = z \otimes z.$$

It is not hard to see that H is a bialgebra with the above constructions. Let R, S, Q, T be operators defined with respect to the basis $\{x, y, z\}$ by

$$\begin{aligned} R(x) &= \lambda_1 z, & R(y) &= \lambda_2 z, & R(z) &= 0, \\ S(x) &= -\lambda_3 y, & S(y) &= 0, & S(z) &= 0, \\ Q(x) &= \lambda_4 z, & Q(y) &= \lambda_4 z, & Q(z) &= 0, \\ T(x) &= \lambda_4(x + z), & T(y) &= \lambda_4(y + z), & T(z) &= \lambda_4 z, \end{aligned}$$

where $\lambda_i \in K, i = 1, 2, 3, 4$. Then (H, R, S, Q, T) is a Rota-Baxter bisystem.

4. COVARIANT BIALGEBRAS

In this section, we redefine covariant bialgebras via coderivations. Moreover, we consider coassociative Yang-Baxter pairs and coassociative Yang-Baxter equation.

Definition 4.1. Let C be a coassociative coalgebra. A coassociative Yang-Baxter pair (abbr. CYBP) on C is a pair of elements $\sigma, \tau \in (C \otimes C)^*$ that satisfy the following equations

$$\sigma(c_1, e)\sigma(c_2, d) - \sigma(c, d_1)\sigma(d_2, e) + \tau(d, e_1)\sigma(c, e_2) = 0, \quad (4.1)$$

$$\tau(c_1, e)\sigma(c_2, d) - \tau(c, d_1)\tau(d_2, e) + \tau(d, e_1)\tau(c, e_2) = 0, \quad (4.2)$$

where $\forall c, d, e \in C$.

Proposition 4.2. Let $f : C \longrightarrow D$ be a coalgebra map. If (σ, τ) is a CYBP on D , then

$$\sigma^f = \sigma \circ (f \otimes f) \text{ and } \tau^f = \tau \circ (f \otimes f) \quad (4.3)$$

form a CYBP on C .

Proof. We compute Eq.(4.1) for (σ^f, τ^f) as follows.

$$\begin{aligned} & \sigma^f(c_1, e)\sigma^f(c_2, d) - \sigma^f(c, d_1)\sigma^f(d_2, e) + \tau^f(d, e_1)\sigma^f(c, e_2) \\ &= \sigma(f(c)_1, f(e))\sigma(f(c)_2, f(d)) - \sigma(f(c), f(d)_1)\sigma(f(d)_2, f(e)) \\ & \quad + \tau(f(d), f(e)_1)\sigma(f(c), f(e)_2) \text{ by } f \text{ is coalgebra map} \\ & \stackrel{(4.1)}{=} 0. \end{aligned}$$

The proof for Eq.(4.2) is similar. □

Example 4.3. (1) The pair (σ, σ) is a CYBP if and only if σ is a solution of the coassociative Yang-Baxter equation

$$\sigma(c_1, e)\sigma(c_2, d) - \sigma(c, d_1)\sigma(d_2, e) + \sigma(d, e_1)\sigma(c, e_2) = 0.$$

(2) If for all $c, d, e \in C$, $\sigma \in (C \otimes C)^*$ satisfies

$$\sigma(c, d_1)\sigma(d_2, e) = \sigma(d, e_1)\sigma(c, e_2) = \sigma(c_1, e)\sigma(c_2, d), \quad (4.4)$$

the dual case of Frobenius-separability equation in [5], then $(\sigma, 0)$ and $(0, \sigma)$ are CYBPs.

(3) If for all $c, d, e \in C$, $\sigma, \tau \in (C \otimes C)^*$ satisfies (4.4), such that

$$\tau(d, e_1)\sigma(c, e_2) = \tau(c_1, e)\sigma(c_2, d) = 0, \quad (4.5)$$

then (σ, τ) is a CYBP.

(4) If for all $c, d, e \in C$, $\sigma, \tau \in (C \otimes C)^*$ satisfies

$$\sigma(c_1, e)\sigma(c_2, d) = \sigma(c, d_1)\sigma(d_2, e), \quad \tau(c, d_1)\tau(d_2, e) = \tau(d, e_1)\tau(c, e_2), \quad (4.6)$$

then $(\sigma, 0)$ and $(0, \tau)$ are CYBPs. If in addition, σ, τ satisfy (4.5), then (σ, τ) is a CYBP.

(5) Let C be a coalgebra. Suppose that $\xi, \zeta \in C^*$ such that for all $c \in C$, $\xi(c_1)\xi(c_2) = 0$ and $\xi(c_1)\zeta(c_2) = \zeta(c_1)\xi(c_2) = 0$, then $(\sigma = \xi \otimes \zeta, \tau = \zeta \otimes \xi)$ is a CYBP.

Proposition 4.4. *Let (σ, τ) be a CYBP on C . Define $Q, T : C \longrightarrow C$ by*

$$Q(c) = \sigma(c_1, c_3)c_2, \quad T(c) = \tau(c_1, c_3)c_2. \quad (4.7)$$

Then (C, Q, T) is a Rota-Baxter cosystem associated to (σ, τ) .

Furthermore, if $f : C \longrightarrow D$ is a coalgebra map and (σ^f, τ^f) is the CYBP induced by f as in Eq.(4.3), then f is a morphism of Rota-Baxter cosystems from the cosystem associated to (σ^f, τ^f) to the cosystem associated to (σ, τ) .

Proof. We only check that Eq.(3.1) holds as follows. For all $c \in C$, we have

$$\begin{aligned} \text{LHS of Eq.(3.1)} &= \sigma(c_1, c_{31})c_2 \otimes \sigma(c_{32}, c_5)c_4 \\ &\stackrel{(4.1)}{=} \sigma(c_{11}, c_5)\sigma(c_{12}, c_3)c_2 \otimes c_4 + \sigma(c_3, c_{51})\sigma(c_1, c_{52})c_2 \otimes c_4 \\ &= \sigma(c_1, c_3)\sigma(c_{211}, c_{213})c_{212} \otimes c_{22} + \sigma(c_1, c_3)\sigma(c_{221}, c_{223})c_{21} \otimes c_{222} \\ &= \text{RHS of Eq.(3.1)}, \end{aligned}$$

finishing the proof. \square

Remark 4.5. *By [4, Proposition 3.4] and Proposition 4.4, we obtain:*

Let H be a bialgebra. Suppose that (r, s) is an associative Yang-Baxter pair on H (see [4]) and (σ, τ) is a CYBP on H , then (H, R, S, Q, T) is a Rota-Baxter bisystem.

Example 4.6. (1) *The Rota-Baxter cosystem associated to (σ, τ) satisfying Eqs.(4.5) and (4.6) in Example 4.3 (4) will satisfy the separated equations*

$$Q(c_1) \otimes Q(c_2) = Q(Q(c)_1) \otimes Q(c)_2, \quad T(c_1) \otimes T(c_2) = T(c)_1 \otimes T(T(c)_2),$$

and

$$Q(c)_1 \otimes T(Q(c)_2) = Q(T(c)_1) \otimes T(c)_2 = 0.$$

For example,

$$Q(c)_1 \otimes T(Q(c)_2) = \sigma(c_1, c_3)\tau(c_{221}, c_{223})c_{21} \otimes T(c_{222}) \stackrel{(4.5)}{=} 0$$

(2) *The Rota-Baxter cosystem associated to (σ, τ) in Example 4.3 (5) is (C, Q, T) , where $Q(c) = \xi(c_1)c_2\zeta(c_3)$ and $T(c) = \zeta(c_1)c_2\xi(c_3)$.*

Definition 4.7. Let C be a coassociative coalgebra and $\delta_1, \delta_2 : C \otimes C \longrightarrow C$ (write $\delta_i(c \otimes d) = c \diamond_i d, i = 1, 2$) be coderivations (i.e. $(c \diamond_i d)_1 \otimes (c \diamond_i d)_2 = c_1 \otimes c_2 \diamond_i d + c \diamond_i d_1 \otimes d_2, i = 1, 2$) (see [6]).

(1) If M is a right C -comodule, then a K -linear map $\nabla_r : M \otimes C \longrightarrow M$ (write $\nabla_r(m \otimes c) = m \triangleleft c$) is called a **right covariant coderivation with respect to δ_1** if

$$(m \triangleleft c)_{(0)} \otimes (m \triangleleft c)_{(1)} = m \triangleleft c_1 \otimes c_2 + m_{(0)} \otimes m_{(1)} \diamond_1 c \quad (4.8)$$

for all $m \in M$ and $c \in C$.

(2) If M is a left C -comodule, then a K -linear map $\nabla_l : C \otimes M \rightarrow M$ (write $\nabla_r(c \otimes m) = c \triangleright m$) is called a **left covariant coderivation with respect to δ_2** if

$$(c \triangleright m)_{(-1)} \otimes (c \triangleright m)_{(0)} = c_1 \otimes c_2 \triangleright m + c \diamond_2 m_{(-1)} \otimes m_{(0)} \quad (4.9)$$

for all $m \in M$ and $c \in C$.

(3) A K -linear map $\nabla : C \otimes C \rightarrow C$ (write $\nabla(c \otimes d) = c \cdot d$) is called a **covariant coderivation with respect to (δ_1, δ_2)** if it is a right covariant coderivation with respect to δ_1 and left covariant coderivation with respect to δ_2 , i.e.

$$(c \cdot d)_1 \otimes (c \cdot d)_2 = c \cdot d_1 \otimes d_2 + c_1 \otimes c_2 \diamond_1 d = c_1 \otimes c_2 \cdot d + c \diamond_2 d_1 \otimes d_2 \quad (4.10)$$

for all $m \in M$ and $c \in C$.

Remark 4.8. (1) Obviously, any coderivation $\delta : C \otimes C \rightarrow C$ is a covariant coderivation with respect to (δ, δ) .

(2) A covariant coderivation with respect to $(0, 0)$ is the same as a C -bicomodule map $C \otimes C \rightarrow C$, where $\rho_r(c \otimes d) = c \otimes d_1 \otimes d_2$ and $\rho_l(c \otimes d) = c_1 \otimes c_2 \otimes d$.

In the following, we introduce a new algebraic structure using coderivations and covariant coderivations, close to covariant bialgebra which we still call covariant bialgebra.

Definition 4.9. A **covariant bialgebra** is a quadruple $(C, \delta_1, \delta_2, \mu)$, such that

- (1) C is a coassociative coalgebra,
- (2) $\delta_1, \delta_2 : C \otimes C \rightarrow C$ are coderivations,
- (3) (C, μ) is an associative algebra such that μ is a covariant coderivation with respect to (δ_1, δ_2) .

If C has a counit, then a covariant bialgebra $(C, \delta_1, \delta_2, \mu)$ is said to be **counital** if moreover we have $\varepsilon(cd) = \varepsilon(c)\varepsilon(d)$, here $cd = \mu(c \otimes d)$.

A **morphism of covariant bialgebras** is a K -linear map that is both an algebra and a coalgebra map.

Example 4.10. (C, μ, μ, μ) is a covariant bialgebra if and only if (c, Δ) is an infinitesimal bialgebra [1], i.e. an algebra equipped with a coassociative comultiplication which is also a derivation, because infinitesimal bialgebras are self-dual.

Theorem 4.11. Let C be a counital coalgebra, and let $\delta_1, \delta_2 : C \otimes C \rightarrow C$ be coderivations. Then there exists an associative covariant coderivation $\mu : C \otimes C \rightarrow C$ (write $\mu(c \otimes d) = cd$) with respect to (δ_1, δ_2) if and only if there exists $u \in (C \otimes C)^*$ such that, for all $c, d, e \in C$

$$c \diamond_1 d - c \diamond_2 d = c_1 u(c_2, d) - u(c, d_1) d_2, \quad (4.11)$$

$$(c \diamond_1 d) \diamond_1 e - c \diamond_1 (d \diamond_1 e) = u(d, e_1) c \diamond_1 e_2, \quad (4.12)$$

$$u(c \diamond_1 d, e) - u(c, d \diamond_1 e) = u(d, e_1) u(c, e_2) - u(c, d_1) u(d_2, e). \quad (4.13)$$

In this case,

$$cd = u(c, d_1) d_2 + c \diamond_1 d = c_1 u(c_2, d) + c \diamond_2 d. \quad (4.14)$$

Proof. (\implies) Set $u = \varepsilon \circ \mu \in (C \otimes C)^*$. Apply $\varepsilon \otimes id$ and $id \otimes \varepsilon$ to Eq.(4.10) for μ , respectively, we get Eq.(4.14), then Eq.(4.11) holds.

By the associativity of μ , we have

$$\mu \circ (\mu \otimes id) = \mu \circ (id \otimes \mu). \quad (4.15)$$

Then by Eq.(4.14),

$$\mu(u(c, d_1)d_2 \otimes e + c \diamond_1 d \otimes e) = \mu(c \otimes u(d, e_1)e_2 + c \otimes d \diamond_1 e).$$

Apply ε to both sides of above equality and by the definition of u , one obtain Eq.(4.13).

Since

$$\text{LHS of Eq.(4.15)} \stackrel{(4.14)}{=} u(c, d_1)u(d_2, e_1)e_2 + u(c, d_1)d_2 \diamond_1 e + u(c \diamond_1 d, e_1)e_2 + (c \diamond_1 d) \diamond_1 e,$$

and

$$\text{RHS of Eq.(4.15)} \stackrel{(4.14)}{=} u(d, e_1)u(c, e_{21})e_{22} + u(d, e_1)c \diamond_1 e_2 + u(c, (d \diamond_1 e)_1)(d \diamond_1 e)_2 + c \diamond_1 (d \diamond_1 e),$$

$$\begin{aligned} (c \diamond_1 d) \diamond_1 e - c \diamond_1 (d \diamond_1 e) &= ((u(c, d \diamond_1 e_1) - u(c \diamond_1 d, e_1)) - (u(c, d_1)u(d_2, e_1) \\ &\quad - u(d, e_{11})u(c, e_{12}))e_2 + u(d, e_1)c \diamond_1 e_2 \text{ (by coderivation)} \\ &\stackrel{(4.13)}{=} u(d, e_1)c \diamond_1 e_2, \end{aligned}$$

so Eq.(4.12) is proved.

(\Leftarrow) By Eq.(4.14), we have

$$\begin{aligned} (cd)_1 \otimes (cd)_2 &= u(c, d_1)d_2 \otimes d_3 + (c \diamond_1 d)_1 \otimes (c \diamond_1 d)_2 \\ &= u(c, d_1)d_2 \otimes d_3 + c_1 \otimes c_2 \diamond_1 d + c \diamond_1 d_1 \otimes d_2 \text{ (by coderivation)} \\ &= cd_1 \otimes d_2 + c_1 \otimes c_2 \diamond_1 d. \end{aligned}$$

Similarly, we get $(cd)_1 \otimes (cd)_2 = c_1 \otimes c_2 d + c \diamond_2 d_1 \otimes d_2$. The rest is obvious. \square

Corollary 4.12. *A quadruple $(C, \delta_1, \delta_2, \mu)$ consisting of a counital coalgebra C , coderivations $\delta_1, \delta_2 : C \otimes C \longrightarrow C$ and a K -linear map $\mu : C \otimes C \longrightarrow C$ is a counital covariant bialgebra if and only if, for all $c, d, e \in C$,*

$$c \diamond_1 d - c \diamond_2 d = c\varepsilon(d) - \varepsilon(c)d, \quad (4.16)$$

$$(c \diamond_1 d) \diamond_1 e - c \diamond_1 (d \diamond_1 e) = \varepsilon(d)c \diamond_1 e. \quad (4.17)$$

In this case,

$$cd = \varepsilon(c)d + c \diamond_1 d. \quad (4.18)$$

Proof. Assume $(C, \delta_1, \delta_2, \mu)$ to be a counital covariant bialgebra, then $u = \varepsilon \otimes \varepsilon$, and the rest is obvious. \square

Theorem 4.13. *Let C be a coalgebra and $\sigma, \tau \in C^*$. Define K -linear maps $\delta_\sigma, \delta_\tau, \mu : C \otimes C \longrightarrow C$ by*

$$\begin{aligned}\delta_\sigma(c \otimes d) &= c_1\sigma(c_2, d) - \sigma(c, d_1)d_2, \\ \delta_\tau(c \otimes d) &= c_1\tau(c_2, d) - \tau(c, d_1)d_2, \\ \mu(c \otimes d) &= c_1\sigma(c_2, d) - \tau(c, d_1)d_2.\end{aligned}\tag{4.19}$$

Then $(C, \delta_\sigma, \delta_\tau, \mu)$ is a covariant bialgebra if and only if, for all $c, d, e \in C$,

$$\begin{aligned}c_1(\sigma(c_{21}, e)\sigma(c_{22}, d) - \sigma(c_2, d_1)\sigma(d_2, e) + \tau(d, e_1)\sigma(c_2, e_2)) \\ = (\tau(c_1, e_1)\sigma(c_2, d) - \tau(c, d_1)\tau(d_2, e_1) + \tau(d, e_{11})\tau(c, e_{12}))e_2.\end{aligned}\tag{4.20}$$

Proof. It is straightforward to check that $\delta_\sigma, \delta_\tau$ are coderivations, and μ is a covariant coderivation with respect to $\delta_\sigma, \delta_\tau$. Thus in order to finish the proof we only need to prove that (C, μ) is an associative algebra if and only if Eq.(4.20) holds. While

$$\begin{aligned}(\mu \circ (id \otimes \mu))(c \otimes d \otimes e) &= c_1\sigma(c_2, d_1)\sigma(d_2, e) - c_1\tau(d, e_1)\sigma(c_2, e_2) \\ &\quad + \tau(d, e_{11})\tau(c, e_{12})e_2 - \tau(c, d_1)d_2\sigma(d_3, e).\end{aligned}$$

and

$$\begin{aligned}(\mu \circ (\mu \otimes id))(c \otimes d \otimes e) &= c_1\sigma(c_{21}, e)\sigma(c_{22}, d) - (\tau(c_1, e_1)\sigma(c_2, d)e_2 \\ &\quad + \tau(c, d_1)\tau(d_2, e_1)e_2 - \tau(c, d_1)d_2\sigma(d_3, e)),\end{aligned}$$

therefore we complete the proof. \square

Remark 4.14. *If C is a non-degenerate coalgebra, then the form of μ in Eq.(4.19) specifies the forms of δ_σ and δ_τ . Indeed, let us suppose that $\mu(c \otimes d) = c_1\sigma(c_2, d) - \tau(c, d_1)d_2$ is a (δ_1, δ_2) -covariant coderivation. Then, for all $c, d \in C$,*

$$c_1 \otimes c_2 \diamond_1 d = c_1 \otimes (c_{21}\sigma(c_{22}, d) - \sigma(c_2, d_1)d_2)$$

and

$$c \diamond_2 d_1 \otimes d_2 = (c_1\tau(c_2, d_1) - \tau(c, d_{11})d_{12}) \otimes d_2.$$

Since C is non-degenerate, these equations imply that $\delta_1 = \delta_\sigma$ and $\delta_2 = \delta_\tau$.

5. COQUASITRIANGULAR COVARIANT BIALGEBRAS AND INFINITESIMAL BIALGEBRAS

In this section, we study coquasitriangular covariant bialgebras generalizing coquasitriangular infinitesimal bialgebras, and corresponding to our covariant bialgebras.

Proposition 5.1. *Let C be a coalgebra and $\sigma, \tau \in C^*$. If (σ, τ) is a CYBP on C and $\delta_\sigma, \delta_\tau, \mu$ are defined by Eq.(4.19), then $(C, \delta_\sigma, \delta_\tau, \mu)$ is a covariant bialgebra. In this case, $(C, \delta_\sigma, \delta_\tau, \mu)$ is called a **coquasitriangular covariant bialgebra**.*

Furthermore, if $f : C \longrightarrow D$ is a coalgebra map and (σ^f, τ^f) is the CYBP induced by f as in Eq.(4.3), then f is a morphism of quasitriangular covariant bialgebras from the one associated to (σ^f, τ^f) to the one associated to (σ, τ) .

Proof. The first statement follows immediately from Theorem 4.13. The second one can be proved directly. \square

Remark 5.2. When $\sigma = \tau$ in Proposition 5.1, $\delta_\sigma = \delta_\tau = \mu$, then $(C, \delta_\sigma, \delta_\tau, \mu)$ is an infinitesimal bialgebra. In this case, we call $(C, \delta_\sigma, \delta_\tau, \mu)$ (abbr. (C, δ_σ)) a **coquasitriangular infinitesimal bialgebra**, which will be studied later.

Proposition 5.3. Let C be a coalgebra, $\sigma, \tau \in (C \otimes C)^*$, $\delta_\sigma, \delta_\tau, \mu$ are defined by Eq.(4.19). Then $(C, \delta_\sigma, \delta_\tau, \mu)$ is a coquasitriangular covariant bialgebra if and only if for all $c, d, e \in C$,

$$\sigma(c, de) = \sigma(c_1, e)\sigma(c_2, d), \quad \tau(cd, e) = -\tau(d, e_1)\tau(c, e_2). \quad (5.1)$$

Proof. It is straightforward to prove that Eq.(5.1) is equivalent to Eqs.(4.1) and (4.2), i.e. (σ, τ) is a CYBP. \square

Proposition 5.4. Counital coquasitriangular covariant bialgebra structures on C are determined by $\sigma \in (C \otimes C)^*$ such that for all $c, d, e \in C$,

$$\sigma(c, de) = \sigma(c_1, e)\sigma(c_2, d), \quad \tau(cd, e) = -\sigma(d, e_1)\sigma(c, e_2) + \varepsilon(c)\sigma(d, e) + \sigma(c, e)\varepsilon(d), \quad (5.2)$$

where $cd = \varepsilon(c)d + c_1\sigma(c_2, d) - \sigma(c, d_1)d_2$.

Proof. A coquasitriangular covariant bialgebra $(C, \delta_\sigma, \delta_\tau, \mu)$ is counital if and only if $\sigma(c, d) - \tau(c, d) = \varepsilon(c)\varepsilon(d)$, i.e., $\tau(c, d) = \sigma(c, d) - \varepsilon(c)\varepsilon(d)$. Thus we complete the proof by replacing τ by $\sigma - \varepsilon \otimes \varepsilon$ in Eq.(5.1). \square

Example 5.5. Let C be a counital coalgebra and $\alpha \in C^*$ such that $\alpha(c_1)\alpha(c_2) = \alpha(c)$, $\forall c \in C$, and $k \in \{0, 1\} \subset K$. Then $\sigma^\alpha(c \otimes d) = k\varepsilon(c)\alpha(d) + (1 - k)\alpha(c)\varepsilon(d)$ satisfies Eq.(5.2). Thus $(\sigma^\alpha, \tau^\alpha)$ is a CYBP, where

$$\tau^\alpha(c \otimes d) = \sigma^\alpha(c \otimes d) - \varepsilon(c)\varepsilon(d) = k\varepsilon(c)\alpha(d) + (1 - k)\alpha(c)\varepsilon(d) - \varepsilon(c)\varepsilon(d),$$

and

$$cd = \mu^\alpha(c \otimes d) = \varepsilon(c)d + k(c\alpha(d) - \varepsilon(c)\alpha(d_1)d_2) + (1 - k)(c_1\alpha(c_2)\varepsilon(d) - \alpha(c)\varepsilon(d)).$$

Then $(C, \sigma^\alpha, \tau^\alpha, \mu^\alpha)$ is a coquasitriangular covariant bialgebra.

By Proposition 4.4, $(C, Q^{\sigma^\alpha}, T^{\tau^\alpha})$ is a Rota-Baxter cosystem associated to $(\sigma^\alpha, \tau^\alpha)$, where

$$Q^{\sigma^\alpha}(c) = kc_1\alpha(c_2) + (1 - k)\alpha(c_1)c_2, \quad T^{\tau^\alpha}(c) = kc_1\alpha(c_2) + (1 - k)\alpha(c_1)c_2 - c.$$

By Proposition 3.7, (C, Δ_*) is a coassociative coalgebra, where

$$\Delta_*(c) = k(c_1\alpha(c_2) \otimes c_3 + c_1 \otimes c_2\alpha(c_3)) + (1 - k)(\alpha(c_1)c_2 \otimes c_3 + c_1 \otimes \alpha(c_2)c_3) - c_1 \otimes c_2.$$

And (C, Δ_\bullet) is a pre-Lie coalgebra, where

$$\Delta_\bullet(c) = k(c_1\alpha(c_2) \otimes c_3 - c_2 \otimes c_1\alpha(c_3)) + (1 - k)(\alpha(c_1)c_2 \otimes c_3 - c_3 \otimes \alpha(c_2)c_1) - c_2 \otimes c_1.$$

Define $\delta(c \otimes d) = \varepsilon(c)d - c\varepsilon(d)$. Then $(C, 0, -\delta, \mu^0)$ and $(C, \sigma, 0, \mu^\varepsilon)$ are counital covariant bialgebras associated to the coderivation δ .

Definition 5.6. A **coquasitriangular infinitesimal bialgebra** is a pair (C, σ) where C is a coassociative coalgebra and $\sigma \in (C \otimes C)^*$ is a solution to

$$\sigma(c_1, e)\sigma(c_2, d) - \sigma(c, d_1)\sigma(d_2, e) + \sigma(d, e_1)\sigma(c, e_2) = 0 \quad (5.3)$$

for all $c, d, e \in C$.

We call Eq.(5.3) a **coassociative Yang-Baxter equation (abbr. CYBA)** in C .

Remark 5.7. The linear form $\sigma \in (C \otimes C)^*$ is a solution to CAYB if and only if (C, μ_σ) is an infinitesimal bialgebra, where $\mu_\sigma(c \otimes d) = c_1\sigma(c_2, d) - \sigma(c, d_1)d_2$, $\forall c, d \in C$.

Example 5.8. Let C be any counital coalgebra possessing an element $\tau \in C^*$ such that $\tau(c_1)\tau(c_2) = 0$. Then $\sigma = \varepsilon \otimes \tau$ satisfies CAYB and the corresponding infinitesimal bialgebra structure is

$$\mu_\sigma(c \otimes d) = c\tau(d) - \varepsilon(c)\tau(d_1)d_2, \forall c, d \in C.$$

Proposition 5.9. A principal coderivation $\mu_\sigma : C \otimes C \longrightarrow C$ is associative if and only if, for all $c, d, e \in C$,

$$\begin{aligned} & c_1(\sigma(c_{21}, e)\sigma(c_{22}, d) - \sigma(c_2, d_1)\sigma(d_2, e) + \sigma(d, e_1)\sigma(c_2, e_2)) \\ &= (\sigma(c_1, e_1)\sigma(c_2, d) - \sigma(c, d_1)\sigma(d_2, e_1) + \sigma(d, e_{11})\sigma(c, e_{12}))e_2. \end{aligned}$$

Proposition 5.10. The pair (C, σ) is a coquasitriangular infinitesimal bialgebra if and only if for all $c, d, e \in C$,

$$\sigma(cd, e) = -\sigma(d, e_1)\sigma(c, e_2), \quad (5.4)$$

$$\sigma(c, de) = \sigma(c_1, e)\sigma(c_2, d). \quad (5.5)$$

Proof. Let $\sigma = \tau$ in Proposition 5.3. □

If (C, μ, Δ) is a finite dimensional infinitesimal bialgebra, then $C' = (C^*, \Delta^{*op}, -\mu^{*cop})$ and $'C = (C^*, -\Delta^{*op}, \mu^{*cop})$ are infinitesimal bialgebras (see [1]).

Proposition 5.11. Let (C, σ) be a finite dimensional coquasitriangular infinitesimal bialgebra. Then the maps

$$\zeta_\sigma : C \longrightarrow C', \quad \zeta_\sigma(c) = \sum e^i \sigma(e_i, c), \quad (5.6)$$

$$\xi_\sigma : C \longrightarrow 'C, \quad \xi_\sigma(c) = \sum e^i \sigma(c, e_i) \quad (5.7)$$

where $c \in C$, $\{e_i\}$ is the basis of C and $\{e^i\}$ is the dual basis, are morphisms of infinitesimal bialgebras.

Proof. We note that for all $b, c \in C$, $\langle \zeta_\sigma(c), b \rangle = \sigma(b, c)$ and $\langle \xi_\sigma(c), b \rangle = \sigma(c, b)$. For all $a, b, c \in C$,

$$\begin{aligned} \langle \zeta_\sigma(ab), c \rangle &= \sigma(c, ab) \stackrel{(5.5)}{=} \sigma(c_1, b)\sigma(c_2, a) \\ &= \langle \zeta_\sigma(b), c_1 \rangle \langle \zeta_\sigma(a), c_2 \rangle = \langle \zeta_\sigma(a)\zeta_\sigma(b), c \rangle, \end{aligned}$$

$$\langle \zeta_\sigma(a)_1 \otimes \zeta_\sigma(a)_2, b \otimes c \rangle = -\langle \zeta_\sigma(a), cb \rangle = -\sigma(cb, a)$$

$$\begin{aligned}
 & \stackrel{(5.4)}{=} \sigma(b, a_1)\sigma(c, a_2) = \langle \zeta_\sigma(a_1), b \rangle \langle \zeta_\sigma(a_2), c \rangle \\
 & = \langle \zeta_\sigma(a_1) \otimes \zeta_\sigma(a_2), b \otimes c \rangle, \\
 \langle \xi_\sigma(ab), c \rangle & = \sigma(ab, c) \stackrel{(5.4)}{=} -\sigma(b, c_1)\sigma(a, c_2) \\
 & = -\langle \xi_\sigma(b), c_1 \rangle \langle \xi_\sigma(a), c_2 \rangle = \langle \xi_\sigma(a)\xi_\sigma(b), c \rangle, \\
 \langle \xi_\sigma(a)_1 \otimes \xi_\sigma(a)_2, b \otimes c \rangle & = \langle \xi_\sigma(a), cb \rangle = \sigma(a, cb) \\
 & \stackrel{(5.5)}{=} \sigma(a_1, b)\sigma(a_2, c) = \langle \xi_\sigma(a_1), b \rangle \langle \xi_\sigma(a_2), c \rangle \\
 & = \langle \xi_\sigma(a_1) \otimes \xi_\sigma(a_2), b \otimes c \rangle,
 \end{aligned}$$

finishing the proof. \square

Corollary 5.12. *Let (C, σ) be a coquasitriangular infinitesimal Hopf algebra with bijective antipode S . Then*

$$\sigma(id \otimes S) = \sigma(S^{-1} \otimes id); \quad (5.8)$$

$$\sigma(S \otimes id) = \sigma(id \otimes S^{-1}); \quad (5.9)$$

$$\sigma(S \otimes S) = \sigma = \sigma(S^{-1} \otimes S^{-1}). \quad (5.10)$$

Proof. Since $\zeta_\sigma : C \rightarrow C'$ and $\xi_\sigma : C \rightarrow {}'C$ are infinitesimal Hopf algebra maps, then $\zeta_\sigma S = S^{-1*}\zeta_\sigma$ and $\xi_\sigma S = S^{-1*}\xi_\sigma$. For all $a, b \in C$,

$$\begin{aligned}
 \sigma(a, S(b)) & = \langle \zeta_\sigma(S(b)), a \rangle = \langle S^{-1*}(\zeta_\sigma(b)), a \rangle \\
 & = \langle \zeta_\sigma(b), S^{-1}(a) \rangle = \sigma(S^{-1}(a), b),
 \end{aligned}$$

$$\begin{aligned}
 \sigma(S(a), b) & = \langle \xi_\sigma(S(a)), b \rangle = \langle S^{-1*}(\xi_\sigma(a)), b \rangle \\
 & = \langle \xi_\sigma(a), S^{-1}(b) \rangle = \sigma(a, S^{-1}(b)),
 \end{aligned}$$

$$\sigma(S \otimes S) = \sigma(id \otimes S)(S \otimes id) = \sigma(S^{-1} \otimes id)(S \otimes id) = \sigma$$

and

$$\sigma(S^{-1} \otimes S^{-1}) = \sigma(S^{-1} \otimes id)(id \otimes S^{-1}) = \sigma(id \otimes S)(id \otimes S^{-1}) = \sigma,$$

finishing the proof. \square

Definition 5.13. A **double coalgebra** is a pair (C, D) of coassociative coalgebras together with a left D -comodule on C : $C \rightarrow D \otimes C$, $c \mapsto c_{-1} \otimes c_0$ and a right C -comodule on D : $D \rightarrow D \otimes C$, $d \mapsto d_{(0)} \otimes d_{(1)}$ such that

$$c_{-1} \otimes c_{01} \otimes c_{02} = c_{1-1} \otimes c_{10} \otimes c_2 + c_{-1(1)} \otimes c_{-1(0)} \otimes c_0, \quad (5.11)$$

$$d_{(0)1} \otimes d_{(0)2} \otimes d_{(1)} = d_1 \otimes d_{2(0)} \otimes d_{2(1)} + d_{(0)} \otimes d_{(1)-1} \otimes d_{(1)0}, \quad (5.12)$$

where $c \in C$ and $d \in D$.

Theorem 5.14. *Given a double coalgebra (C, D) . Define the comultiplication $\Delta : C \otimes D \longrightarrow C \otimes D \otimes C \otimes D$ by*

$$\Delta(c \otimes d) = c_1 \otimes c_{2-1} \otimes c_{20} \otimes d + c \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2. \quad (5.13)$$

Then $(C \otimes D, \Delta)$ is a coassociative coalgebra.

Proof. For all $c \in C$ and $d \in D$, we have

$$\begin{aligned} (\Delta \otimes id)\Delta(c \otimes d) &= c_{11} \otimes c_{12-1} \otimes c_{120} \otimes c_{2-1} \otimes c_{20} \otimes d \\ &\quad + c_1 \otimes c_{2-11(0)} \otimes c_{2-11(1)} \otimes c_{2-1(2)} \otimes c_{20} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \\ &\quad + c \otimes d_{1(0)1(0)} \otimes d_{1(0)1(1)} \otimes d_{1(0)2} \otimes d_{1(1)} \otimes d_2 \\ &= c_1 \otimes c_{2-1} \otimes c_{20} \otimes c_{3-1} \otimes c_{30} \otimes d \\ &\quad + c_1 \otimes c_{2-1(0)} \otimes c_{2-1(1)} \otimes c_{20-1} \otimes c_{200} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \\ &\quad + c \otimes d_{1(0)1(0)} \otimes d_{1(0)1(1)} \otimes d_{1(0)2} \otimes d_{1(1)} \otimes d_2 \\ &\stackrel{(5.12)}{=} c_1 \otimes c_{2-1} \otimes c_{20} \otimes c_{3-1} \otimes c_{30} \otimes d \\ &\quad + c_1 \otimes c_{2-1(0)} \otimes c_{2-1(1)} \otimes c_{20-1} \otimes c_{200} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \\ &\quad + c \otimes d_{11(0)} \otimes d_{11(1)} \otimes d_{12(0)} \otimes d_{12(1)} \otimes d_2 \\ &\quad + c \otimes d_{1(0)(0)} \otimes d_{1(0)(1)} \otimes d_{1(1)-1} \otimes d_{1(1)0} \otimes d_2 \\ &= c_1 \otimes c_{2-1} \otimes c_{20} \otimes c_{3-1} \otimes c_{30} \otimes d \\ &\quad + c_1 \otimes c_{2-1(0)} \otimes c_{2-1(1)} \otimes c_{20-1} \otimes c_{200} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \\ &\quad + c \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_{2(0)} \otimes d_{2(1)} \otimes d_3 \\ &\quad + c \otimes d_{1(0)} \otimes d_{1(0)1} \otimes d_{1(1)2-1} \otimes d_{1(1)20} \otimes d_2 \\ &= c_1 \otimes c_{21-1} \otimes c_{210} \otimes c_{22-1} \otimes c_{220} \otimes d \\ &\quad + c_1 \otimes c_{2-1(0)} \otimes c_{2-1(1)} \otimes c_{20-1} \otimes c_{200} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \\ &\quad + c \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_{2(0)} \otimes d_{2(1)} \otimes d_3 \\ &\quad + c \otimes d_{1(0)} \otimes d_{1(0)1} \otimes d_{1(1)2-1} \otimes d_{1(1)20} \otimes d_2 \\ &\stackrel{(5.11)}{=} c_1 \otimes c_{2-1} \otimes c_{201} \otimes c_{202-1} \otimes c_{2020} \otimes d \\ &\quad + c_1 \otimes c_{2-1(0)} \otimes c_{2-1(1)} \otimes c_{20-1} \otimes c_{200} \otimes d \\ &\quad + c_1 \otimes c_{2-1} \otimes c_{20} \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_2 \end{aligned}$$

$$\begin{aligned}
 & +c \otimes d_{1(0)} \otimes d_{1(1)} \otimes d_{2(0)} \otimes d_{2(1)} \otimes d_3 \\
 & +c \otimes d_{1(0)} \otimes d_{1(0)1} \otimes d_{1(1)2-1} \otimes d_{1(1)20} \otimes d_2 \\
 = & (id \otimes \Delta)\Delta(c \otimes d),
 \end{aligned}$$

finishing the proof. \square

Lemma 5.15. *Consider the maps:*

$$\rho_l : C \longrightarrow {}^lC \otimes C, \quad c \mapsto e^i \otimes e_i c, \quad (5.14)$$

$$\rho_r : {}^lC \longrightarrow {}^lC \otimes C, \quad c \mapsto -fe^i \otimes e_i, \quad (5.15)$$

where $c \in C, f \in {}^lC, \{e_i\}$ is the basis of C and $\{e^i\}$ is the dual basis. Then $(C, {}^lC)$ is a double coalgebra.

Proof. For all $c \in C$ and $f, g, h \in {}^lC$, we have

$$\begin{aligned}
 (id \otimes \rho_l)\rho_l(c) &= e^i \otimes e^j \otimes e_j e_i c \\
 &= e^i_2 \otimes e^i_1 \otimes e_i c = (\Delta \otimes id)\rho_l(c),
 \end{aligned}$$

so (C, ρ_l) is a left lC -comodule. Also

$$\begin{aligned}
 (\rho_r \otimes id)\rho_r(f) &= fe^i e^j \otimes e_j \otimes e_i \\
 &= -fe^i \otimes e_{i1} \otimes e_{i2} = (id \otimes \Delta)\rho_r(f),
 \end{aligned}$$

so $({}^lC, \rho_r)$ is a right C -comodule.

Applying $id \otimes g \otimes h$ to $fe^i e^j \otimes e_j \otimes e_i$,

$$\begin{aligned}
 fe^i e^j \langle g, e_j \rangle \langle h, e_i \rangle &= fhg \\
 &= fe^i \langle -\Delta^{*op}(h \otimes g), e_i \rangle \\
 &= -fe^i \langle \Delta^{*op}(h \otimes g), e_i \rangle \\
 &= -fe^i \langle g, e_{i1} \rangle \langle h, e_{i2} \rangle
 \end{aligned}$$

Since

$$\begin{aligned}
 c_{-1} \otimes c_{01} \otimes c_{02} &= e^i \otimes (e_i c)_1 \otimes (e_i c)_2 \\
 &= e^i \otimes e_i c_1 \otimes c_2 + e^i \otimes e_{i1} \otimes e_{i2} c
 \end{aligned}$$

and

$$\begin{aligned}
 c_{1-1} \otimes c_{10} \otimes c_2 + c_{-1(1)} \otimes c_{-1(0)} \otimes c_0 \\
 = e^i \otimes e_i c_1 \otimes c_2 - e^i e^j \otimes e_j \otimes e_i c,
 \end{aligned}$$

while for all $c' \in C$,

$$\begin{aligned}
 \langle e^i, c' \rangle e_{i1} \otimes e_{i2} c &= c'_1 \otimes c'_2 c \\
 &= \langle e^j, c'_1 \rangle \langle e^i, c'_2 \rangle e_j \otimes e_i c \\
 &= \langle -e^i e^j, c' \rangle e_j \otimes e_i c,
 \end{aligned}$$

thus Eq.(5.11) holds.

Since

$$\begin{aligned} d_{(0)1} \otimes d_{(0)2} \otimes d_{(1)} &= -(de^i)_1 \otimes (de^i)_2 \otimes e_i \\ &= -de^i_1 \otimes e^i_2 \otimes e_i - d_1 \otimes d_2 e^i \otimes e_i \end{aligned}$$

and

$$\begin{aligned} d_1 \otimes d_{2(0)} \otimes d_{2(1)} + d_{(0)} \otimes d_{(1)-1} \otimes d_{(1)0} \\ = -d_1 \otimes d_2 e^i \otimes e_i - de^i \otimes e^j \otimes e_j e_i \end{aligned}$$

while for all $c', d' \in C$,

$$\begin{aligned} \langle de^i_1, c' \rangle \langle e^i_2, d' \rangle e_i &= -\langle d, c'_2 \rangle \langle e^i_1, c'_1 \rangle \langle e^i_2, d' \rangle e_i \\ &= -\langle d, c'_2 \rangle \langle e^i, d' c'_1 \rangle e_i \\ &= -\langle d, c'_2 \rangle d' c'_1 \\ &= -\langle d, c'_2 \rangle \langle e^i, c'_1 \rangle d' e_i \\ &= \langle de^i, c' \rangle d' e_i \\ &= \langle de^i, c' \rangle \langle e^j, d' \rangle e_j e_i, \end{aligned}$$

so Eq.(5.12) holds. Therefore the proof is finished. \square

Theorem 5.16. *Let C be a finite dimensional infinitesimal bialgebra. Consider the linear space*

$$\widehat{D(C)} := C \otimes {}'C.$$

Let $\{e_i\}$ be the basis of C and $\{e^i\}$ the dual basis of $'C$. Define

$$\begin{aligned} \Delta(c \otimes f) &= c_1 \otimes c_{2-1} \otimes c_{20} \otimes f + c \otimes f_{1(0)} \otimes f_{1(1)} \otimes f_2 \\ &= c_1 \otimes e^i \otimes e_i c_2 \otimes f - c \otimes f_1 e^i \otimes e_i \otimes f_2. \end{aligned} \tag{5.16}$$

Then $\widehat{D(C)}$ is a coalgebra.

Furthermore, define $\sigma \in (\widehat{D(C)} \otimes \widehat{D(C)})^$ by*

$$\sigma(c \otimes f, d \otimes g) = \langle g, c \rangle \langle f, d \rangle. \tag{5.17}$$

Then σ is a solution of coassociative Yang-Baxter equation. i.e., $(\widehat{D(C)}, \sigma)$ is a coquasitriangular infinitesimal bialgebra. Moreover

$$\mu_\sigma(c \otimes f, d \otimes g) = -c \otimes f_1 \langle f_2, d \rangle g - c \langle f, d_1 \rangle d_2 \otimes g, \tag{5.18}$$

where $c, d \in C$ and $f, g \in {}'C$.

Proof. By Lemma 5.15, we get the first asseration. It is straightforward that Eq.(5.18) is a solution of CAYB in $\widehat{D(C)}$. \square

Definition 5.17. Let $(C, \delta_1, \delta_2, \mu)$ be a covariant bialgebra. A right C -module and a right C -comodule M is said to be a **right covariant C -module** if the action is a right covariant coderivation with respect to δ_1 . Symmetrically, a left C -module and a left C -comodule N is said to be a **left covariant C -module** if the action is a left covariant coderivation with respect to δ_2 . A **morphism of covariant modules** is a map that is both C -linear and C -colinear. The category of right (left) covariant C -modules is denoted by $\mathcal{M}_C^C(CM)$.

Example 5.18. Let $(C, \delta_\sigma, \delta_\tau, \mu)$ be a coquasitriangular covariant bialgebra associated to a CYBP (σ, τ) . Then every right C -comodule M is a right covariant C -module with the action

$$\triangleleft_M : M \otimes C \longrightarrow M, m \otimes c \mapsto m_{(0)}\sigma(m_{(1)}, c). \quad (5.19)$$

Similarly, every left C -comodule N is a left covariant C -module with the action

$$\triangleright_M : C \otimes N \longrightarrow N, c \otimes n \mapsto -\tau(c, m_{(-1)})m_{(0)}. \quad (5.20)$$

Proof. For all $m \in M$ and $c \in C$,

$$\begin{aligned} m \triangleleft_M c_1 \otimes c_2 + m_{(0)} \otimes m_{(1)} \triangleleft_1 c &= m_{(0)} \otimes m_{(1)1}\sigma(m_{(1)2}, c) \\ &= m_{(0)(0)} \otimes m_{(0)(1)}\sigma(m_{(1)}, c) \\ &= \rho(m \triangleleft_M c). \end{aligned}$$

So the action is a right covariant coderivation with respect to δ_σ . In what follows, we will show that (M, \triangleleft_M) is a right C -module. For all $c, d \in C$ and $m \in M$, we have

$$\begin{aligned} (\triangleleft_M \circ (id \otimes \mu))(m \otimes c \otimes d) &= m \triangleleft_M (cd) \\ &= m_{(0)}\sigma(m_{(1)}, cd) \\ &\stackrel{(5.1)}{=} m_{(0)}\sigma(m_{(1)1}, d)\sigma(m_{(1)2}, c) \\ &= m_{(0)(0)}\sigma(m_{(0)(1)}, d)\sigma(m_{(1)}, c) \\ &= (\triangleleft_M \circ (\triangleleft_M \otimes id))(m \otimes c \otimes d). \end{aligned}$$

Therefore, the first statement is verified. Similarly, we prove the second one. \square

Next, we describe the relation between Rota-Baxter cosystems and dendriform coalgebras.

Definition 5.19. ([10]) A triple $(C, \Delta_\prec, \Delta_\succ)$ consisting of a vector space C and two K -linear operators $\Delta_\prec, \Delta_\succ : C \longrightarrow C \otimes C$ is called a **dendriform coalgebra** if, for all $c \in C$,

$$(\Delta_\prec \otimes id) \circ \Delta_\prec(c) = (id \otimes \Delta_\prec + id \otimes \Delta_\succ) \circ \Delta_\prec(c), \quad (5.21)$$

$$(\Delta_\succ \otimes id) \circ \Delta_\prec(c) = (id \otimes \Delta_\prec) \circ \Delta_\succ(c), \quad (5.22)$$

$$(\Delta_\prec \otimes id + \Delta_\succ \otimes id) \circ \Delta_\succ(c) = (id \otimes \Delta_\succ) \circ \Delta_\succ(c). \quad (5.23)$$

Theorem 5.20. Let C be a coassociative coalgebra and $Q, T : C \longrightarrow C$ be K -linear maps. Define the K -linear maps $\Delta_\prec, \Delta_\succ : C \longrightarrow C \otimes C$ by

$$\Delta_\prec(c) = c_1 \otimes T(c_2), \quad \Delta_\succ(c) = Q(c_1) \otimes c_2 \quad (5.24)$$

for all $c \in C$. Then:

- (1) If (C, Q, T) is a Rota-Baxter cosystem, then $(C, \Delta_{\prec}, \Delta_{\succ})$ is a dendriform coalgebra.
- (2) If C is a non-degenerate coalgebra and $(C, \Delta_{\prec}, \Delta_{\succ})$ is a dendriform coalgebra, then (C, Q, T) is a Rota-Baxter cosystem.

Proof. (1) First, we prove that Eq.(5.21) holds as follows.

$$\begin{aligned}
 \text{RHS of Eq.(5.21)} &= c_1 \otimes T(c_2)_1 \otimes T(T(c_2)_2) + c_1 \otimes Q(T(c_2)_1) \otimes T(c_2)_2 \\
 &\stackrel{(3.2)}{=} c_1 \otimes T(c_{21}) \otimes T(c_{22}) \\
 &= c_{11} \otimes T(c_{12}) \otimes T(c_2) = \text{LHS of Eq.(5.21)}.
 \end{aligned}$$

Eq.(5.22) can be checked easily by the coassociativity.

$$\begin{aligned}
 \text{LHS of Eq.(5.23)} &= Q(Q(c_1)_1) \otimes Q(c_1)_2 \otimes c_2 + Q(c_1)_1 \otimes T(Q(c_1)_2) \otimes c_2 \\
 &\stackrel{(3.1)}{=} T(c_{11}) \otimes T(c_{12}) \otimes c_2 \\
 &= T(c_1) \otimes T(c_{21}) \otimes c_{22} = \text{RHS of Eq.(5.23)}.
 \end{aligned}$$

Thus $(C, \Delta_{\prec}, \Delta_{\succ})$ is a dendriform coalgebra.

- (2) It is straightforward by the proof of (1) and the non-degeneracy of C . \square

Remark 5.21. In [10], the notions of dendriform bialgebra, codendriform bialgebra and bidendriform bialgebra were introduced by combining dendriform algebra and dendriform coalgebra structures. As mentioned above, Rota-Baxter (co)algebra can produce dendriform (co)algebra. Then an interesting question is whether there is a kind of Rota-Baxter “bi”-algebras that lead to structures of dendriform bialgebra, codendriform bialgebra or bidendriform bialgebra.

6. EXAMPLES OF ROTA-BAXTER BIALGEBRAS AND BISYSTEMS

In this section we provide further examples of Rota-Baxter bialgebras and bisystems.

6.1. 2-dimensional Examples.

Example 6.1 (2-dimensional bialgebra). We consider the 2-dimensional bialgebra $K \times K$, where the multiplication is defined with respect to a basis $\{u_1, u_2\}$ by $u_1 \cdot u_i = u_i \cdot u_1 = u_i$ for $i = 1, 2$ and $u_2 \cdot u_2 = u_2$; and the comultiplication is defined by $\Delta(u_1) = u_1 \otimes u_1$, $\Delta(u_2) = u_2 \otimes u_2$. We set that u_1 is the unit and the counit is defined by $\varepsilon(u_1) = \varepsilon(u_2) = 1$.

- The (R, R) -Rota-Baxter structures of weight λ are given by
 - (a) $R(u_1) = -\lambda u_1$, $R(u_2) = 0$,
 - (b) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$,
 - (c) $R(u_1) = -\lambda u_1$, $R(u_2) = -\lambda u_2$.
- The (R, Q) -Rota-Baxter structures of weight λ and γ respectively are given by the pairs (R, Q) taken from the following list of R 's
 - (a) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$,

- (b) $R(u_1) = -\lambda u_2, R(u_2) = -\lambda u_2,$
- (c) $R(u_1) = \lambda u_2, R(u_2) = 0,$
- (d) $R(u_1) = 0, R(u_2) = \lambda u_1 - \lambda u_2,$
- (e) $R(u_1) = -2\lambda u_1 + \lambda u_2, R(u_2) = -\lambda u_1,$
- (f) $R(u_1) = -\lambda u_1, R(u_2) = 0,$
- (g) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_2,$
- (h) $R(u_1) = -\lambda u_1 - \lambda u_2, R(u_2) = -\lambda u_2,$
- (i) $R(u_1) = -\lambda u_1 + \lambda u_2, R(u_2) = 0,$
- (j) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_1,$
- (k) $R(u_1) = \lambda u_1 - \lambda u_2, R(u_2) = \lambda u_1 - \lambda u_2,$

and the following list of Q 's

- (a) $Q(u_1) = 0, Q(u_2) = -\gamma u_2,$
- (b) $Q(u_1) = -\gamma u_1, Q(u_2) = 0,$
- (c) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2.$
- The bisystems are given by the pairs (R, S) and pairs (Q, T) taken from the following list for (R, S)
 - (a) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1 + p_2 u_2, S(u_2) = p_2 u_2,$
 - (b) $R(u_1) = p_1 u_2, R(u_2) = p_1 u_2, S(u_1) = -p_2 u_1 + p_1 u_2, S(u_2) = 0,$
 - (c) $R(u_1) = p_1 u_2, R(u_2) = p_2 u_2, S(u_1) = (p_1 - p_2) u_1, S(u_2) = 0,$
 - (d) $R(u_1) = p_1 u_1, R(u_2) = p_1 u_1, S(u_1) = -p_2 u_1 + p_2 u_2, S(u_2) = p_1 u_1 - p_1 u_2,$
 - (e) $R(u_1) = -p_1 u_1 + p_1 u_2, R(u_2) = 0, S(u_1) = p_2 u_2, S(u_2) = p_2 u_2,$
 - (f) $R(u_1) = -p_1 u_1 + p_1 u_2, R(u_2) = -p_1 u_1 + p_1 u_2, S(u_1) = -(p_1 + p_2) u_1 + p_2 u_2, S(u_2) = -p_1 u_1,$
 - (g) $R(u_1) = -p_1 u_1 + p_1 u_2, R(u_2) = p_1 u_1 - p_1 u_2, S(u_1) = p_1 u_1, S(u_2) = p_1 u_1,$
 - (h) $R(u_1) = -(p_1 + p_2) u_1 + p_1 u_2, R(u_2) = -p_2 u_1, S(u_1) = -p_2 u_1 + p_2 u_2, S(u_2) = -p_2 u_1 + p_2 u_2,$
 - (i) $R(u_1) = p_1 u_1 + p_2 u_2, R(u_2) = p_2 u_2, S(u_1) = p_1 u_2, S(u_2) = 0,$
 - (j) $R(u_1) = p_1 u_1, R(u_2) = 0, S(u_1) = p_2 u_2, S(u_2) = (-p_1 + p_2) u_2,$
 - (k) $R(u_1) = 0, R(u_2) = -p_1 u_2, S(u_1) = p_1 u_1, S(u_2) = 0,$
 - (l) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = 0,$
 - (m) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = -p_1 u_2,$
 - (n) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1, S(u_2) = 0,$
 - (o) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = 0,$
 - (p) $R(u_1) = p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = -p_1 u_2,$
 - (q) $R(u_1) = p_1 u_2, R(u_2) = p_1 u_2, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = p_1 u_1 - p_1 u_2,$
 - (r) $R(u_1) = p_1 u_2, R(u_2) = p_1 u_2, S(u_1) = p_1 u_1 - p_1 u_2, S(u_2) = p_1 u_1 - p_1 u_2,$
 - (s) $R(u_1) = p_1 u_1 - p_1 u_2, R(u_2) = 0, S(u_1) = p_1 u_2, S(u_2) = 0,$
 - (t) $R(u_1) = p_1 u_1 - p_1 u_2, R(u_2) = -p_1 u_2, S(u_1) = p_1 u_2, S(u_2) = 0,$

- (u) $R(u_1) = p_1u_1$, $R(u_2) = 0$, $S(u_1) = p_1u_2$, $S(u_2) = 0$,
(v) $R(u_1) = p_1u_1 - p_1u_2$, $R(u_2) = 0$, $S(u_1) = p_1u_2$, $S(u_2) = 0$,
where p_1, p_2 are parameters, and the following list for (Q, T)
- (a) $Q(u_1) = 0$, $Q(u_2) = q_1u_2$, $T(u_1) = q_2u_1$, $T(u_2) = 0$,
(b) $Q(u_1) = q_1u_1$, $Q(u_2) = 0$, $T(u_1) = 0$, $T(u_2) = q_2u_2$,
(c) $Q(u_1) = q_1u_2$, $Q(u_2) = 0$, $T(u_1) = 0$, $T(u_2) = q_1u_2$,
(d) $Q(u_1) = 0$, $Q(u_2) = q_1u_1 + q_1u_2$, $T(u_1) = q_1u_1 + q_1u_2$, $T(u_2) = 0$,
(e) $Q(u_1) = 0$, $Q(u_2) = q_1u_1$, $T(u_1) = q_1u_1$, $T(u_2) = 0$,
(f) $Q(u_1) = q_1u_1 + q_1u_2$, $Q(u_2) = 0$, $T(u_1) = 0$, $T(u_2) = q_1u_1 + q_1u_2$,
where q_1, q_2 are parameters.

6.2. 3-dimensional and 4-dimensional Examples.

Example 6.2 (3-dimensional bialgebra). *We consider the 3-dimensional bialgebra $K \times K$, where the multiplication is defined with respect to a basis $\{u_1, u_2, u_3\}$ by $u_1 \cdot u_i = u_i \cdot u_1 = u_i$ for $i = 1, 2, 3$ and $u_2 \cdot u_2 = u_2$, $u_2 \cdot u_3 = u_3 \cdot u_2 = u_3$, $u_3 \cdot u_3 = 0$; and the comultiplication is defined by $\Delta(u_1) = u_1 \otimes u_1$, $\Delta(u_2) = u_1 \otimes u_2 + u_2 \otimes u_1 - u_2 \otimes u_2$, $\Delta(u_3) = u_1 \otimes u_3 + u_3 \otimes u_1 - u_3 \otimes u_2$. We set that u_1 is the unit and the counit is defined by $\varepsilon(u_1) = 1$, $\varepsilon(u_2) = \varepsilon(u_3) = 0$.*

- The (R, R) -Rota-Baxter structures of weight λ are given by
 - (a) $R(u_1) = -\lambda u_1$, $R(u_2) = 0$, $R(u_3) = 0$,
 - (b) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$, $R(u_3) = 0$,
 - (c) $R(u_1) = 0$, $R(u_2) = 0$, $R(u_3) = -\lambda u_3$,
 - (d) $R(u_1) = -\lambda u_1$, $R(u_2) = -\lambda u_2$, $R(u_3) = 0$,
 - (e) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$, $R(u_3) = -\lambda u_3$,
 - (f) $R(u_1) = -\lambda u_1$, $R(u_2) = 0$, $R(u_3) = -\lambda u_3$,
 - (g) $R(u_1) = -\lambda u_1$, $R(u_2) = -\lambda u_2$, $R(u_3) = -\lambda u_3$,
- The (R, Q) -Rota-Baxter structures of weight λ and γ respectively are given by the pairs (R, Q) taken from the following list of R 's
 - (a) $R(u_1) = 0$, $R(u_2) = 0$, $R(u_3) = -\lambda u_3$,
 - (b) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$, $R(u_3) = 0$,
 - (c) $R(u_1) = 0$, $R(u_2) = -\lambda u_2$, $R(u_3) = -\lambda u_3$,
 - (d) $R(u_1) = -\lambda u_2$, $R(u_2) = -\lambda u_2$, $R(u_3) = 0$,
 - (e) $R(u_1) = -\lambda u_2$, $R(u_2) = -\lambda u_2$, $R(u_3) = -\lambda u_3$,
 - (f) $R(u_1) = \lambda u_2$, $R(u_2) = 0$, $R(u_3) = 0$,
 - (g) $R(u_1) = \lambda u_2$, $R(u_2) = 0$, $R(u_3) = -\lambda u_3$,
 - (h) $R(u_1) = 0$, $R(u_2) = \lambda u_1 - \lambda u_2$, $R(u_3) = 0$,
 - (i) $R(u_1) = 0$, $R(u_2) = \lambda u_1 - \lambda u_2$, $R(u_3) = -\lambda u_3$,
 - (j) $R(u_1) = -2\lambda u_1 + \lambda u_2$, $R(u_2) = -\lambda u_1$, $R(u_3) = 0$,
 - (k) $R(u_1) = -2\lambda u_1 + \lambda u_2$, $R(u_2) = -\lambda u_1$, $R(u_3) = -\lambda u_3$,

- (l) $R(u_1) = -\lambda u_1, R(u_2) = 0, R(u_3) = 0,$
- (m) $R(u_1) = -\lambda u_1, R(u_2) = 0, R(u_3) = -\lambda u_3,$
- (n) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_2, R(u_3) = 0,$
- (o) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_2, R(u_3) = -\lambda u_3,$
- (p) $R(u_1) = -\lambda u_1 - \lambda u_2, R(u_2) = -\lambda u_2, R(u_3) = 0,$
- (q) $R(u_1) = -\lambda u_1 - \lambda u_2, R(u_2) = -\lambda u_2, R(u_3) = -\lambda u_3,$
- (r) $R(u_1) = -\lambda u_1 + \lambda u_2, R(u_2) = 0, R(u_3) = 0,$
- (s) $R(u_1) = -\lambda u_1 + \lambda u_2, R(u_2) = 0, R(u_3) = -\lambda u_3,$
- (t) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_1, R(u_3) = 0,$
- (u) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_1, R(u_3) = -\lambda u_3,$
- (v) $R(u_1) = \lambda u_1 - \lambda u_2, R(u_2) = \lambda u_1 - \lambda u_2, R(u_3) = 0,$
- (w) $R(u_1) = \lambda u_1 - \lambda u_2, R(u_2) = \lambda u_1 - \lambda u_2, R(u_3) = -\lambda u_3,$

and the following list of Q 's

- (a) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = -\gamma u_3.$
- (b) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = 0.$
- (c) $Q(u_1) = -\gamma u_1, Q(u_2) = 0, Q(u_3) = -\gamma u_3.$
- (d) $Q(u_1) = 0, Q(u_2) = -\gamma u_2, Q(u_3) = -\gamma u_3.$
- (e) $Q(u_1) = -\gamma u_1, Q(u_2) = 0, Q(u_3) = 0.$
- (f) $Q(u_1) = 0, Q(u_2) = -\gamma u_2, Q(u_3) = 0.$
- (g) $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = -\gamma u_3.$
- (h) $Q(u_1) = 0, Q(u_2) = pu_2 - \sqrt{-p(\gamma + p)}u_3, Q(u_3) = -\sqrt{-p(\gamma + p)}u_2 - (\gamma + p)u_3.$
- (i) $Q(u_1) = 0, Q(u_2) = pu_2 + \sqrt{-p(\gamma + p)}u_3, Q(u_3) = \sqrt{-p(\gamma + p)}u_2 - (\gamma + p)u_3.$
- (j) $Q(u_1) = -\gamma u_1, Q(u_2) = pu_2 - \sqrt{-p(\gamma + p)}u_3, Q(u_3) = -\sqrt{-p(\gamma + p)}u_2 - (\gamma + p)u_3.$
- (k) $Q(u_1) = -\gamma u_1, Q(u_2) = pu_2 + \sqrt{-p(\gamma + p)}u_3, Q(u_3) = \sqrt{-p(\gamma + p)}u_2 - (\gamma + p)u_3,$

where p is a parameter.

- Examples of bisystems are given by pairs (R, S) and pairs (Q, T) taken from the following list for (R, S)

- (a) $R(u_1) = p_1 u_3, R(u_2) = p_1 u_3, R(u_3) = -p_2 u_3,$
 $S(u_1) = (p_2 - p_3)u_1 + p_3 u_2, S(u_2) = p_2 u_2, S(u_3) = 0.$
- (b) $R(u_1) = p_1 u_3, R(u_2) = 0, R(u_3) = 0,$
 $S(u_1) = p_2 u_2 + (p_1 + p_3)u_3, S(u_2) = p_2 u_2 + p_3 u_3, S(u_3) = p_2 u_3.$
- (c) $R(u_1) = p_1 u_2 + p_2 u_3, R(u_2) = p_2 u_3, R(u_3) = -p_3 u_3,$
 $S(u_1) = p_1 u_1 + p_3 u_2, S(u_2) = p_3 u_2, S(u_3) = 0.$
- (d) $R(u_1) = -p_1 u_1 + p_1 u_2, R(u_2) = -p_2 u_1 + p_2 u_2, R(u_3) = 0,$
 $S(u_1) = -p_2 u_1 + p_3 u_3, S(u_2) = -p_2 u_1 + p_3 u_3, S(u_3) = -p_2 u_3.$
- (e) $R(u_1) = -p_1 u_1 + p_1 u_2, R(u_2) = -p_1 u_1 + p_1 u_2, R(u_3) = p_2 u_1 - p_2 u_2,$
 $S(u_1) = -(p_1 + p_3)u_1 + p_3 u_2, S(u_2) = -p_1 u_1, S(u_3) = p_2 u_1.$

- (f) $R(u_1) = -(p_1 + p_2)u_1 + p_1u_2$, $R(u_2) = -p_2u_1$, $R(u_3) = p_3u_1$,
 $S(u_1) = -p_2u_1 + p_2u_2$, $S(u_2) = -p_2u_1 + p_2u_2$, $S(u_3) = p_3u_1 - p_3u_2$.
- (g) $R(u_1) = p_1u_1 - p_1u_2 + p_2u_3$, $R(u_2) = p_1u_1 - p_1u_2 + p_2u_3$, $R(u_3) = -p_1u_3$,
 $S(u_1) = p_1u_2$, $S(u_2) = p_1u_2$, $S(u_3) = 0$.

where p_1, p_2, p_3 are parameters, an the following list for (Q, T)

- (a) $Q(u_1) = -q_1u_1$, $Q(u_2) = 0$, $Q(u_3) = -q_1u_3$,
 $T(u_1) = 0$, $T(u_2) = q_1u_2$, $T(u_3) = 0$.
- (b) $Q(u_1) = -q_1u_1 + q_1u_2$, $Q(u_2) = -q_1u_1 + q_1u_2$, $Q(u_3) = 0$,
 $T(u_1) = q_1u_2$, $T(u_2) = q_1u_1$, $T(u_3) = q_1u_3$.
- (c) $Q(u_1) = -q_1u_1 + q_1u_2$, $Q(u_2) = 0$, $Q(u_3) = -q_1u_3$,
 $T(u_1) = q_1u_1 + q_1u_2$, $T(u_2) = q_1u_1 + q_1u_2$, $T(u_3) = 0$.
- (d) $Q(u_1) = -\frac{q_1^2+q_2^2}{q_1}u_1$, $Q(u_2) = -\frac{q_1^2+q_2^2}{q_1}u_1$, $Q(u_3) = \frac{q_1^2+q_2^2}{q_2}u_1$,
 $T(u_1) = 0$, $T(u_2) = q_1u_2 + q_2u_3$, $T(u_3) = q_2u_2 + \frac{q_2^2}{q_1}u_3$.
- (e) $Q(u_1) = -\frac{q_1^2+q_2^2}{q_1}u_1 + q_1u_2 + q_2u_3$, $Q(u_2) = -\frac{2q_1^2+q_2^2}{q_1}u_1$, $Q(u_3) = \frac{2q_1^2+q_2^2}{q_2}u_1$,
 $T(u_1) = q_1u_1 + q_1u_2 + q_2u_3$, $T(u_2) = q_1u_1 + q_1u_2 + q_2u_3$, $T(u_3) = q_2u_1 + q_2u_2 + \frac{q_2^2}{q_1}u_3$.
- (f) $Q(u_1) = -q_1u_1 + q_1u_2$, $Q(u_2) = -q_1u_1 + q_1u_2$, $Q(u_3) = q_2u_1 - q_2u_2$,
 $T(u_1) = q_1u_2$, $T(u_2) = q_1u_1$, $T(u_3) = 0$.
- (g) $Q(u_1) = 0$, $Q(u_2) = q_1u_1 - \frac{q_1q_3+q_2^2}{q_3}u_2 + q_2u_3$, $Q(u_3) = \frac{q_1q_3}{q_2}u_1 + \frac{q_1q_3+q_2^2}{q_2}u_2 - q_3u_3$,
 $T(u_1) = \frac{q_3^2+q_2^2}{q_3}u_1$, $T(u_2) = q_3u_2 + q_2u_3$, $T(u_3) = q_2u_2 + \frac{q_2^2}{q_3}u_3$.

where q_1, q_2, q_3 are parameters.

Example 6.3. Now, we consider the unital Taft-Sweedler algebra generated by g, x and the relations $(g^2 = 1, x^2 = 0, xg = -gx)$. The comultiplication is defined by $\Delta(g) = g \otimes g$ and $\Delta(x) = x \otimes 1 + g \otimes x$, the counit is given by $\varepsilon(g) = 1$, $\varepsilon(x) = 0$.

The bialgebra T_2 is 4-dimensional, it is defined with respect to a basis $\{u_1 = 1, u_2 = g, u_3 = x, u_4 = gx\}$ by the following table which describes multiplying the i th row elements by the j th column elements.

	u_1	u_2	u_3	u_4
u_1	u_1	u_2	u_3	u_4
u_2	u_2	u_1	u_4	u_3
u_3	u_3	$-u_4$	0	0
u_4	u_4	$-u_3$	0	0

and

$$\Delta(u_1) = u_1 \otimes u_1, \Delta(u_2) = u_2 \otimes u_2, \Delta(u_3) = u_3 \otimes u_1 + u_2 \otimes u_3, \Delta(u_4) = u_4 \otimes u_2 + u_1 \otimes u_4.$$

$$\varepsilon(u_1) = \varepsilon(u_2) = 1, \quad \varepsilon(u_3) = \varepsilon(u_4) = 0.$$

- The (R, R) -Rota-Baxter structures of weight λ are given by

- (a) $R(u_1) = 0, R(u_2) = 0, R(u_3) = -\lambda u_3, R(u_4) = -\lambda u_4, ,$
- (b) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_2, R(u_3) = 0, R(u_4) = 0, ,$
- (c) $R(u_1) = -\lambda u_1, R(u_2) = -\lambda u_2, R(u_3) = -\lambda u_3, R(u_4) = -\lambda u_4, ,$
- The (R, Q) -Rota-Baxter structures of weight λ and γ respectively are given by pairs (R, Q) taken from the following list of R 's
 - (a) $R(u_1) = 0, R(u_2) = -p_1 u_1 + p_1 u_2 - \frac{(\lambda+p_1)(\lambda+p_1+p_2)}{p_3} u_3 + \frac{(\lambda+p_1)(\lambda+p_2)}{p_3} u_4,$
 $R(u_3) = -p_3 u_1 + p_3 u_2 - (2\lambda + p_1 + p_2) u_3 + (\lambda + p_2) u_4,$
 $R(u_4) = -p_3 u_1 + p_3 u_2 - (\lambda + p_1 + p_2) u_3 + p_2 u_4,$
 - (b) $R(u_1) = -\lambda u_1, R(u_2) = (\lambda + p_1) u_1 + p_1 u_2 - \frac{(\lambda+p_1)(\lambda+p_1+p_2)}{p_3} u_3 + \frac{(\lambda+p_1)(\lambda+p_2)}{p_3} u_4,$
 $R(u_3) = p_3 u_1 + p_3 u_2 - (2\lambda + p_1 + p_2) u_3 + (\lambda + p_2) u_4,$
 $R(u_4) = p_3 u_1 + p_3 u_2 - (\lambda + p_1 + p_2) u_3 + p_2 u_4,$
 - (c) $R(u_1) = -\lambda u_1, R(u_2) = \lambda u_1 + p_1 u_3 + \frac{p_1 p_2}{\lambda+p_2} u_4,$
 $R(u_3) = -(\lambda + p_2) u_3 - p_2 u_4,$
 $R(u_4) = (\lambda + p_2) u_3 + p_2 u_4,$
 - (d) $R(u_1) = -\lambda u_1, R(u_2) = \lambda u_1 + \frac{\lambda(\lambda+p_1)}{p_2} u_3 + \frac{\lambda(\lambda+p_1)}{p_2} u_4,$
 $R(u_3) = -p_2 u_1 - p_2 u_2 - (2\lambda + p_1) u_3 - (\lambda + p_1) u_4,$
 $R(u_4) = p_2 u_1 + p_2 u_2 + (\lambda + p_1) u_3 + p_1 u_4,$
 - (e) $R(u_1) = \frac{1}{2} \lambda u_1 - \frac{1}{2} \lambda u_2 + p_1 u_3 + p_2 u_4, R(u_2) = \frac{1}{2} \lambda u_1 - \frac{1}{2} \lambda u_2 - p_2 u_3 + p_1 u_4,$
 $R(u_3) = -\frac{1}{2} \lambda u_3 - \frac{1}{2} \lambda u_4,$
 $R(u_4) = -\frac{1}{2} \lambda u_3 - \frac{1}{2} \lambda u_4,$

where p_1, p_2, p_3 are parameters, and the following list of Q 's

- (a) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = -\gamma u_3, Q(u_4) = -\gamma u_4,$
- (b) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = -\gamma u_3, Q(u_4) = 0,$
- (c) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = 0, Q(u_4) = 0,$
- (d) $Q(u_1) = -\gamma u_1, Q(u_2) = 0, Q(u_3) = 0, Q(u_4) = 0,$
- (e) $Q(u_1) = -\gamma u_1, Q(u_2) = -\gamma u_2, Q(u_3) = 0, Q(u_4) = -\gamma u_4,$
- (f) $Q(u_1) = 0, Q(u_2) = 0, Q(u_3) = -\gamma u_3, Q(u_4) = -\gamma u_4.$
- (g) $Q(u_1) = 0, Q(u_2) = -\gamma u_2, Q(u_3) = 0, Q(u_4) = 0,$
- Examples of bisystems are given by a pair (R, S) and a pair (Q, T) taken from the following list for (R, S)
 - (a) $R(u_1) = -p_1 u_1, R(u_2) = 0, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = p_2 u_4, S(u_2) = p_3 u_4, S(u_3) = 0, S(u_4) = p_1 u_4,$
 - (b) $R(u_1) = -(p_1 + p_2) u_1, R(u_2) = -(p_1 + p_2) u_2, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = p_3 u_3 - \frac{p_3 p_2}{p_1} u_4, S(u_2) = -p_3 u_3 + \frac{p_3 p_2}{p_1} u_4, S(u_3) = p_1 u_3 - p_2 u_4, S(u_4) = -p_1 u_3 + p_2 u_4,$
 - (c) $R(u_1) = -(p_1 + p_2) u_1, R(u_2) = -(p_1 + p_2) u_2, R(u_3) = 0, R(u_4) = 0,$
 $S(u_1) = p_3 u_3 + \frac{p_3 p_2}{p_1} u_4, S(u_2) = p_3 u_3 + \frac{p_3 p_2}{p_1} u_4, S(u_3) = p_1 u_3 + p_2 u_4, S(u_4) = p_1 u_3 + p_2 u_4,$

- (d) $R(u_1) = -p_1u_1$, $R(u_2) = -p_1u_2$, $R(u_3) = 0$, $R(u_4) = 0$,
 $S(u_1) = p_3u_3 + p_4u_4$, $S(u_2) = p_4u_3 + p_3u_4$, $S(u_3) = p_1u_3$, $S(u_4) = p_1u_4$,
- (e) $R(u_1) = -p_1u_1$, $R(u_2) = -p_1u_2$, $R(u_3) = 0$, $R(u_4) = 0$,
 $S(u_1) = p_2u_4$, $S(u_2) = -p_2u_4$, $S(u_3) = -p_1u_4$, $S(u_4) = p_1u_4$,
- (f) $R(u_1) = -p_1u_1$, $R(u_2) = -p_1u_2$, $R(u_3) = 0$, $R(u_4) = 0$,
 $S(u_1) = p_2u_4$, $S(u_2) = p_2u_4$, $S(u_3) = p_1u_4$, $S(u_4) = p_1u_4$,
- (g) $R(u_1) = -(p_1 + p_2)u_1$, $R(u_2) = -(p_1 + p_2)u_2$, $R(u_3) = 0$, $R(u_4) = 0$,
 $S(u_1) = 0$, $S(u_2) = 0$, $S(u_3) = p_1u_3 - p_2u_4$, $S(u_4) = -p_1u_3 + p_2u_4$,
- (h) $R(u_1) = -2p_1u_1$, $R(u_2) = -2p_1u_2$, $R(u_3) = 0$, $R(u_4) = 0$,
 $S(u_1) = p_2u_3 - p_2u_4$, $S(u_2) = -p_2u_3 + p_2u_4$, $S(u_3) = p_1u_3 - p_1u_4$, $S(u_4) = -p_1u_3 + p_2u_4$,
- (i) $R(u_1) = 0$, $R(u_2) = 0$, $R(u_3) = -p_1u_3$, $R(u_4) = -p_1u_4$,
 $S(u_1) = p_1u_1$, $S(u_2) = p_1u_2$, $S(u_3) = 0$, $S(u_4) = 0$,
- (j) $R(u_1) = 0$, $R(u_2) = 0$, $R(u_3) = 0$, $R(u_4) = -p_1u_4$,
 $S(u_1) = p_1u_1$, $S(u_2) = 0$, $S(u_3) = 0$, $S(u_4) = 0$,

where p_1, p_2, p_3 are parameters, and the following list for (Q, T)

- (a) $Q(u_1) = 0$, $Q(u_2) = 0$, $Q(u_3) = q_1u_2 - \frac{q_1q_3}{q_2}$, $Q(u_4) = q_2u_2 - q_3u_4$,
 $T(u_1) = q_4u_1$, $T(u_2) = q_3u_2$, $T(u_3) = 0$, $T(u_4) = 0$,
- (b) $Q(u_1) = 0$, $Q(u_2) = q_1u_2 + q_2u_3$, $Q(u_3) = q_3u_3$, $Q(u_4) = \frac{q_1q_4}{q_2}u_2 + q_4u_4$,
 $T(u_1) = -q_3u_1$, $T(u_2) = 0$, $T(u_3) = 0$, $T(u_4) = 0$,
- (c) $Q(u_1) = q_1u_2 + q_2u_4$, $Q(u_2) = 0$, $Q(u_3) = q_3u_2 + \frac{q_3q_2}{q_1}u_4$, $Q(u_4) = q_4u_2 + \frac{q_2q_4}{q_1}u_4$,
 $T(u_1) = 0$, $T(u_2) = \frac{q_1^2 - q_4q_2}{q_1}u_2$, $T(u_3) = 0$, $T(u_4) = 0$,
- (d) $Q(u_1) = 0$, $Q(u_2) = q_1u_2 + q_2u_3$, $Q(u_3) = -q_3u_3$, $Q(u_4) = 0$,
 $T(u_1) = q_3u_1$, $T(u_2) = 0$, $T(u_3) = 0$, $T(u_4) = q_3u_4$,
- (e) $Q(u_1) = q_1u_2 + q_2u_4$, $Q(u_2) = 0$, $Q(u_3) = 0$, $Q(u_4) = \frac{q_1(q_1 - q_3)}{q_2}u_2 + (q_1 - q_3)u_4$,
 $T(u_1) = 0$, $T(u_2) = q_3u_2$, $T(u_3) = q_3u_3$, $T(u_4) = 0$,
- (f) $Q(u_1) = q_1u_2 + q_2u_4$, $Q(u_2) = -q_3u_2$, $Q(u_3) = 0$, $Q(u_4) = \frac{q_1(q_1 + q_3)}{q_2}u_2 + q_1u_4$,
 $T(u_1) = 0$, $T(u_2) = 0$, $T(u_3) = q_3u_3$, $T(u_4) = 0$,
- (g) $Q(u_1) = q_1u_1 + q_1u_2$, $Q(u_2) = 0$, $Q(u_3) = q_2u_2 + q_1u_3$, $Q(u_4) = -q_1u_4$,
 $T(u_1) = 0$, $T(u_2) = q_1u_1 + q_1u_2$, $T(u_3) = 0$, $T(u_4) = 0$,
- (h) $Q(u_1) = q_1u_1 + q_1u_2$, $Q(u_2) = 0$, $Q(u_3) = q_1u_3 - q_1u_4$, $Q(u_4) = 0$,
 $T(u_1) = 0$, $T(u_2) = q_1u_1 + q_1u_2$, $T(u_3) = 0$, $T(u_4) = q_1u_3 - q_1u_4$,
- (i) $Q(u_1) = q_1u_1 + q_1u_2$, $Q(u_2) = 0$, $Q(u_3) = q_1u_3 + q_1u_4$, $Q(u_4) = 0$,
 $T(u_1) = 0$, $T(u_2) = q_1u_1 + q_1u_2$, $T(u_3) = 0$, $T(u_4) = -q_1u_3 - q_1u_4$,
- (j) $Q(u_1) = q_1u_1$, $Q(u_2) = -q_2u_2$, $Q(u_3) = 0$, $Q(u_4) = -q_2u_4$,
 $T(u_1) = 0$, $T(u_2) = 0$, $T(u_3) = q_2u_3$, $T(u_4) = 0$,

where q_1, q_2, q_3 are parameters.

Acknowledgments This work was partially supported by China Postdoctoral Science Foundation (No.2017M611291), Foundation for Young Key Teacher by Henan Province (No. 2015GGJS-088) and Natural Science Foundation of Henan Province (No. 17A110007). T. Ma is grateful to the Erasmus Mundus project FUSION for supporting the postdoctoral fellowship visiting to Mälardalen University, Västerås, Sweden and to Division of Applied Mathematics at the School of Education, Culture and Communication for cordial hospitality.

REFERENCES

- [1] M. Aguiar, Infinitesimal Hopf algebras. Contemporary Mathematics 267, Amer. Math. Soc. (2000): 1-29.
- [2] H. H. An, C. M. Bai, From Rota-Baxter algebras to pre-Lie algebras. J. Phys. A. 41(1)(2008): 015201, 19 pp.
- [3] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity. Pacific J. Math. 10(1960): 731-742.
- [4] T. Brzeziński, Rota-Baxter systems, dendriform algebras and covariant bialgebras. J. Algebra 460(2016): 1-25.
- [5] S. Caenepeel, G. Militaru, S. Zhu, Frobenius and Separable Functors for Generalized Hopf Modules and Nonlinear Equations, Lecture Notes in Math., vol.1787, Springer, Berlin, 2002.
- [6] Y. Doi, Homological coalgebra. J. Math. Soc. Japan 33(1981): 31-50.
- [7] K. Ebrahimi-Fard, L. Guo, Multiple zeta values and Rota-Baxter algebras. Integers, 8(2)(2008): A4, 18 pp.
- [8] K. Ebrahimi-Fard, L. Guo, Free Rota-Baxter algebras and rooted trees. J. Algebra Appl. 7(2)(2008): 167-194.
- [9] K. Ebrahimi-Fard, L. Guo, Rota-Baxter algebras in renormalization of perturbative quantum field theory. Universality and renormalization, 47-105, Fields Inst. Commun., 50, Amer. Math. Soc., Providence, RI, 2007.
- [10] L. Foissy, Bidendriform bialgebras, trees, and free quasi-symmetric functions. J. Pure Appl. Algebra 209(2)(2007): 439-459.
- [11] L. Guo, An introduction to Rota-Baxter algebra. Surveys of Modern Mathematics, 4. International Press, Somerville, MA; Higher Education Press, Beijing, 2012. xii+226 pp.
- [12] L. Guo, Properties of free Baxter algebras. Adv. Math. 151(2000): 346-374.
- [13] L. Guo, W. Keigher, Baxter algebras and shuffle products. Adv. Math. 150(2000): 117-149.
- [14] L. Guo, W. Keigher, On differential Rota-Baxter algebras, J. Pure Appl. Algebra 212(2008) 522-540.
- [15] L. Guo, B. Zhang, Polylogarithms and multiple zeta values from free Rota-Baxter algebras. Sci. China Math. 2010, 53 (9): 2239-2258.
- [16] R. Q. Jian, J. Zhang, Rota-Baxter coalgebras, arXiv:1409.3052.
- [17] T. S. Ma, L. L. Liu, Rota-Baxter coalgebras and Rota-Baxter bialgebras. Linear Multilinear Algebra, 64(5)(2016): 968-979.
- [18] D. E. Radford, Hopf Algebras, KE Series on Knots and Everything, Vol. 49, World Scientific, New Jersey, 2012.
- [19] G. C. Rota, Baxter algebras and combinatorial identities. I, II, Bull. Amer. Math. Soc. 75(1969): 325-329, 330-334.

SCHOOL OF MATHEMATICS AND INFORMATION SCIENCE, HENAN NORMAL UNIVERSITY, XINXIANG 453007,
CHINA

E-mail address: `matianshui@yahoo.com`

UNIVERSITÉ DE HAUTE ALSACE, LABORATOIRE DE MATHÉMATIQUES, INFORMATIQUE ET APPLICATIONS, 4,
RUE DES FRÈRES LUMIÈRE F-68093 MULHOUSE, FRANCE

E-mail address: `abdenacer.makhlouf@uha.fr`

DIVISION OF APPLIED MATHEMATICS, SCHOOL OF EDUCATION, CULTURE AND COMMUNICATION, MÄLARDALEN
UNIVERSITY, 72123 VÄSTERÄS, SWEDEN

E-mail address: `sergei.silvestrov@mdh.se`