

COMMUTING GRAPHS OF BOUNDEDLY GENERATED SEMIGROUPS

TOMER BAUER AND BE'ERI GREENFELD

ABSTRACT. Araújo, Kinyon and Konieczny [3] pose several problems concerning the construction of arbitrary commuting graphs of semigroups.

We observe that every star-free graph is the commuting graph of some semigroup. Consequently, we suggest modifications for some of the original problems, generalized to the context of families of semigroups with a bounded number of generators, and pose related problems.

We construct monomial semigroups with a bounded number of generators, whose commuting graphs have an arbitrary clique number. In contrast to that, we show that the diameter of the commuting graphs of semigroups in a wider class (containing the class of nilpotent semigroups), is bounded by the minimal number of generators plus two.

We also address a problem concerning knit degree.

1. INTRODUCTION

We denote the center of a semigroup S by $Z(S)$. The *commuting graph* of S , denoted by $\Gamma(S)$, is the simple graph whose vertex set is $S - Z(S)$, where two vertices are connected by an edge if their corresponding elements in S commute. For example, if S is a commutative semigroup, then $\Gamma(S)$ is the empty graph. A study of these graphs in relation with certain algebraic structures can be found, for instance in [1, 2, 3, 4, 5].

Araújo, Kinyon and Konieczny [3] proved that for every natural number $n \geq 2$, there exists a semigroup whose commuting graph has diameter n . They also pose the following question:

Question 1 ([3], Section 6, (4)). *Can every number $n \geq 3$ be the clique number (girth, chromatic number) of the commuting graph of a semigroup?*

All semigroups constructed in this note are assumed to have a zero element, denoted 0. We call a semigroup *monomial* if it has a presentation with only monomial relations (i.e. relations of the form $w = 0$).

We now classify which graphs are realizable as commuting graphs of semigroups, thereby answering Question 1.

Definition 2. A vertex in a graph is called a *star* if it is connected to any other vertex. A graph is called *star-free* if none of its vertices is a star.

It is obvious that any commuting graph is star-free. The following result shows that the converse is true as well.

The first author is partially supported by the Bar-Ilan University President's Doctoral Fellowships of Excellence Program.

Proposition 3. *Let $\Gamma = (V, E)$ be a star-free graph. Then there exists a (monomial) semigroup S such that $\Gamma(S) = \Gamma$. Moreover, if Γ has n vertices, then there exists a (monomial) semigroup S , with $\Gamma(S) = \Gamma$, such that S is generated by n elements and its order is $n^2 + n + 1 - |E|$.*

Proof. Label the vertices of Γ to generate a free semigroup with zero. Consider its quotient semigroup

$$S = \left\langle v \in V \left| \begin{array}{l} v_1 v_2 v_3 = 0 \ \forall v_1, v_2, v_3 \\ vu = uv \ \forall (u, v) \in E \end{array} \right. \right\rangle.$$

It is clear that the center $Z(S)$ consists of all elements of the form vu . It is now evident why $\Gamma(S) = \Gamma$. Also, an easy calculation shows that if Γ has n vertices then this semigroup has precisely $n^2 + n + 1 - |E|$ elements.

Note we could set the relations $uv = vu = 0$ whenever $(u, v) \in E$, resulting in a monomial semigroup with the same commuting graph. \square

Let $\text{rank}(\Gamma)$ denote the minimal number of generators for a semigroup whose commuting graph is Γ . Note that the construction in Proposition 3 only shows that $\text{rank}(\Gamma) \leq |V|$. For a semigroup S , we call the minimal number of generators the *rank* of S .

Thus, one can modify Question 1 and ask for families of semigroups generated by a bounded number of generators, yet having arbitrary graph-theoretic invariants. We exhibit such families with arbitrary clique numbers (and leave open the same question about girth).

For a certain class of finite semigroups (which contains the class of nilpotent semigroups), we show that the diameter of the commuting graph is bounded by a linear function of the rank.

In Section 4 we provide a counterexample to another problem from [3] concerning the notion of knit degree (defined in the sequel).

2. COMMUTING GRAPHS WITH ARBITRARY CLIQUE NUMBER

Denote the clique number of a graph Γ by $\omega(\Gamma)$. In view of the previous section it is evident that there are semigroups whose commuting graphs have arbitrary clique number. However, the semigroups that arise from Proposition 3 have rank at least the prescribed clique number, thus unbounded.

To exhibit a family of boundedly generated semigroups with arbitrary clique number, we first introduce a family of monomial semigroups S_n having arbitrary odd clique number $2n - 1$, all of which are generated by two elements.

Commuting graphs with even clique number are then constructed by the simple operations of null union of semigroups and join of graphs:

Definition 4. Let S, T be two semigroups with zero. Their *null union* $S \bullet T$ is the semigroup $S \cup T$ after identifying together their zeros. For every pair of elements $s \in S$ and $t \in T$ their product in $S \bullet T$ is defined to be $st = ts = 0$.

Definition 5. The *join* of two simple graphs, denoted by \vee , is the graph union with all the edges that connect the vertices of the first graph with the vertices of the second graph.

Proposition 6. *Let n be a natural number. Then*


(1) *The semigroup*

$$S_n = \langle a, b \mid a^{n+1} = b^2 = 0, aba = ba^i b = 0 \forall i \geq 1 \rangle$$


satisfies $\omega(\Gamma(S_n)) = 2n - 1$.

(2) *The semigroup* $S_n \bullet S_1$ *satisfies* $\omega(\Gamma(S_n \bullet S_1)) = 2n$.

Example 7. The graph $\Gamma(S_2)$ is



and the graph $\Gamma(S_3)$ is



Proof of (1). Clearly $S_n = \{0, b, a, a^2, \dots, a^n, ab, a^2b, \dots, a^nb, ba, ba^2, \dots, ba^n\}$, so S_n has $3n + 2$ elements. We calculate $Z(S_n) = \{0, a^n b, ba^n\}$, so $|\Gamma(S_n)| = 3n - 1$. Take

$$C = \{a^i b\}_{1 \leq i \leq n-1} \cup \{ba^j\}_{1 \leq j \leq n-1} \cup \{b\}$$

which is clearly a clique of size $2n - 1$, and so $\omega(\Gamma(S_n)) \geq 2n - 1$.

Now, let C' be a clique of $\Gamma(S_n)$. If $b \in C'$, then $a^i \notin C'$ for all i so $|C'| \leq 2n - 1$. If $a^i \in C'$ with the minimal such i , then $b, a, \dots, a^{i-1}, a^j b, ba^j \notin C'$ for all $j \leq n - i$, which are together $2n - i$ vertices, so

$$|C'| \leq (3n - 1) - [1 + (i - 1) + 2(n - i)] = n + i - 1 \leq 2n - 1$$

If both $b \notin C'$ and $a^i \notin C'$ for all i , then $|C'| \leq 2n - 2$. This completes the proof of the first claim that $\omega(\Gamma(S_n)) = 2n - 1$. \square

Proof of (2). In order to construct commuting graphs with arbitrary even clique number, we shall need the following properties of join of graphs and null union of semigroups.

- (1) Let G, H be two graphs. Then $\omega(G \vee H) = \omega(G) + \omega(H)$.
- (2) Let S, T be two semigroups with zero. Then $\Gamma(S \bullet T) \cong \Gamma(S) \vee \Gamma(T)$ (naturally), as $Z(S \bullet T) = Z(S) \bullet Z(T)$.
- (3) Let S, T be two semigroups with zero. Then $\omega(\Gamma(S \bullet T)) = \omega(\Gamma(S)) + \omega(\Gamma(T))$.

Thus, to exhibit boundedly generated semigroups whose commuting graph has arbitrary even clique number, it is enough to consider

$$S_1 := \langle a, b \mid a^2 = b^2 = 0, aba = bab = 0 \rangle = \{0, a, b, ab, ba\}.$$

Clearly we have that $Z(S_1) = \{0, ab, ba\}$ and $\Gamma(S_1) \cong \overline{K}_2$, the edgeless graph with two vertices.

By the above properties and the fact that $\omega(\Gamma(S_1)) = 1$ one obtains that $\omega(\Gamma(S_n \bullet S_1)) = 2n$ and that $S_n \bullet S_1$ can be generated by 4 elements.

Notice that the commuting graph of the semigroup $S'_n := \underbrace{S_1 \bullet \cdots \bullet S_1}_n$ has clique number n . However, the semigroup S'_n can not be generated by less than $2n$ elements, which is obviously unbounded. \square

3. DIAMETER OF COMMUTING GRAPHS

Recall that the diameter of a (connected) graph is the maximal distance between two vertices. In [3], the authors introduce constructions of semigroups with connected commuting graphs having arbitrary diameter. The constructions there have unbounded rank that grows linearly with the diameter. By Proposition 3 it is again possible to obtain such examples, now with rank equal to the number of vertices (hence bigger than the prescribed diameter).

Proposition 6 exhibits commuting graphs of monomial semigroups with at most 4 generators having arbitrary clique number. For a wider family of semigroups, we show next that the diameter of their commuting graphs is effectively bounded by the rank of the semigroups.

We now show that there is a tight connection between the diameter and the rank of a semigroup in a wide class of semigroups. For a semigroup S and $m \in \mathbb{N}$ denote

$$S^m := \{a_1 a_2 \cdots a_m \mid a_i \in S \ \forall i\} \subseteq S$$

In particular we have $S^m \subseteq S^{m-1} \subseteq \cdots \subseteq S$.

Proposition 8. *Let S be a semigroup generated by d elements. Suppose it has a non-central ideal I such that $IS, SI \subseteq Z(S)$. If $\Gamma(S)$ is connected, then its diameter D satisfies $D \leq d + 2$.*

Proof. Pick $u \in I \setminus Z(S)$. We claim that for every $v \in S^2$, we have that $uv = vu$. Indeed, write $v = t_1 t_2$, and compute

$$uv = ut_1 t_2 = t_2 ut_1 = t_1 t_2 u = vu$$

since $ut_1 \in IS$ and $t_2 u \in SI$ are central.

Let $x, y \in \Gamma(S)$ be two vertices. If $x, y \in S^2$ then a path of length (at most) 2 can be found between them, as u commutes with both of them. Suppose only one of x, y , say x , is not contained in S^2 (but $y \in S^2$). Note that $|S \setminus S^2| \leq d$, because S is generated by d elements. Then by connectivity a shortest path can be found between x and some vertex $z \in S^2$. As the path is minimal, its length does not exceed d , which is the number of generators. We can now concatenate this path to $z - u - y$, resulting in a path of length at most $d + 2$, connecting x and y .

We now assume that both $x, y \notin S^2$. We can find shortest paths γ_x, γ_y from x, y to some $x', y' \in S^2$, respectively and denote the lengths of these paths by d_x, d_y . As before, minimality implies that γ_x consists of precisely d_x vertices from $S \setminus S^2$ (and likewise γ_y consists of d_y such vertices). But $d_x + d_y \leq d$, for otherwise the pigeonhole principle would imply that some vertex from $S \setminus S^2$ appears in both γ_x, γ_y , so a path between x and y can be found of length at most $d - 1$. Hence the path $x - \cdots - x' - u - y' - \cdots - y$ obtained by concatenating $\gamma_x, x' - u - y', \gamma_y$ has length at most $d + 2$. \square

Let S be a non-commutative nilpotent semigroup of nilpotency degree c , and let m be the minimal exponent for which $S^m \subseteq Z(S)$. Then a possible ideal satisfying the premise of the previous Proposition is $I = S^{m-1}$. The nilpotency of S guarantees $m \leq c$ because $S^c = \{0\} \subseteq Z(S)$. As mentioned, if S is commutative (e.g. $c \leq 2$), then its commuting graph is empty and the claim holds trivially.

4. A SEMIGROUP WITH KNIT DEGREE 3

Another question that was posed in [3] is related to the notion of knit degree:

Definition 9 ([3]). Let S be a semigroup. A path $a_1 - a_2 - \cdots - a_m$ in $\Gamma(S)$ is called a *left path* (abbreviated in [3] as *l-path*) if $a_1 \neq a_m$ and $a_1 a_i = a_m a_i$ for every $1 \leq i \leq m$. If $\Gamma(S)$ contains a left path, then the *knit degree* of S is defined to be the length of a shortest left path in $\Gamma(S)$.

For $n = 2$ and every $n \geq 4$ a band with knit degree n was constructed in [3]. In [3, Section 6(1)] it was guessed that a semigroup with knit degree 3 does not exist.

We now provide a counterexample.


Example 10. The following semigroup has knit degree 3:

$$S = \langle x_1, x_2, x_3, x_4 | R \rangle$$

with the relations

$$R = \left\{ \begin{array}{l} x_i x_j x_k = 0 \quad \forall i, j, k \\ x_1^2 = x_4 x_1 \\ x_4^2 = x_1 x_4 \\ x_1 x_2 = x_4 x_2 = x_1 x_3 = x_4 x_3 = x_2 x_1 = x_2 x_3 = x_3 x_2 = x_3 x_4 = 0 \end{array} \right\}$$

One may verify that the graph $\Gamma(S)$ is a path graph on four vertices:



The relations (except for $x_i x_j x_k = 0$) ensure that $x_1 - x_2 - x_3 - x_4$ is a left path.

We are left to show that a shorter left path does not exist in the graph. There is an automorphism transposing $x_1 \leftrightarrow x_4$ and $x_2 \leftrightarrow x_3$, so it suffices to check that $x_1 - x_2$, $x_2 - x_3$ and $x_1 - x_3$ are not left paths, which is an easy exercise.

5. FURTHER PROBLEMS

Question 11. Can $\Gamma(S)$ be connected with arbitrarily large diameter for S with bounded rank?

We would like to remark that in general, there is no uniform bound on the rank of finite graphs. In fact, we have:

Claim 12. Let G be a $(2n - 2)$ -regular graph on $2n$ vertices. Then $\text{rank}(G) = 2n$.

Proof. By the generic construction in Proposition 3, we have that $\text{rank}(G) \leq 2n$. To show that $\text{rank}(G) = 2n$, consider any subset X of at most $2n - 1$ vertices of G . Then X lacks some vertex v , so there exists a vertex u in G (the one not connected to v) which is connected to all vertices of X except u itself. Let S be a semigroup such that $\Gamma(S) = G$. Suppose that the elements of X plus some (perhaps none) elements of $Z(S)$ generate S . Then u must lie in $Z(S)$. We conclude that $\text{rank}(G) = 2n$. \square

Following that, we restate another question that was posed in [3].

Question 13. *Is there a family of graphs with bounded rank and unbounded girth?*

We remark that for the class of semigroups satisfying the condition in Proposition 8, the answer to the above question is negative.

It is natural to specialize and ask about C_n , the cyclic graph with n vertices.

Question 14. *What is the rank of C_n ?*

For example, here is an argument showing that the rank of C_4 is 4. Clearly every generating subset must contain at least 2 non-central generators, say x_1 and x_2 . They must lie on non-adjacent vertices of C_4 . If the rank were 2, then every other vertex would have corresponded to a central element (as it commutes with both x_1 and x_2), a contradiction. Hence, there must be at least 3 generators, but then the third generator again commutes with all generators, so 4 generators are needed. A similar reasoning shows that the rank of C_5 is at least 3.

Acknowledgement. We are grateful to Stuart Margolis, Michael Schein, Uzi Vishne and two anonymous referees for helpful comments.

REFERENCES

- [1] A. Abdollahi, S. Akbari, and H. R. Maimani. Non-commuting graph of a group. *J. Algebra*, 298(2):468–492, 2006.
- [2] S. Akbari, M. Ghandehari, M. Hadian, and A. Mohammadian. On commuting graphs of semisimple rings. *Linear Algebra Appl.*, 390:345–355, 2004.
- [3] J. Araújo, M. Kinyon, and J. Konieczny. Minimal paths in the commuting graphs of semigroups. *European J. Combin.*, 32(2):178–197, 2011.
- [4] M. Giudici and A. Pope. The diameters of commuting graphs of linear groups and matrix rings over the integers modulo m . *Australas. J. Combin.*, 48:221–230, 2010.
- [5] Y. Segev. On finite homomorphic images of the multiplicative group of a division algebra. *Ann. of Math. (2)*, 149(1):219–251, 1999.

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, RAMAT GAN 52900, ISRAEL