

PRODUCTS OF CW COMPLEXES

ANDREW D. BROOKE-TAYLOR

ABSTRACT. We provide for the first time a complete characterisation of when the product as topological spaces of two CW complexes is a CW complex, valid irrespective of set theory. The previous state-of-the-art was that, under set-theoretic assumptions like the Continuum Hypothesis, it is necessary and sufficient that a criterion of Whitehead or a criterion of Milnor holds. The general case without making such assumptions is not quite so simple, but we show that it still essentially reduces to counting the number of cells.

1. INTRODUCTION

CW complexes are considered to be well-behaved spaces that provide a convenient setting for the development of algebraic topology. Indeed, their dimension-by-dimension construction from Euclidean cells rules out many point-set-theoretic pathologies which are inessential to fundamental questions of homotopy theory, such as computation of the homotopy groups of spheres. Furthermore, since every topological space is weakly equivalent to a CW complex, no homotopy-theoretic information is lost by restricting attention to CW complexes.

The category of CW complexes does have drawbacks, however. A well-known quirk is that the inclusion of this category into the category of topological spaces does not preserve products: the product as topological spaces of two CW complexes X and Y need not be a CW complex. Given two CW complexes X and Y , there is a natural cell structure on their Cartesian product $X \times Y$ which satisfies closure-finiteness (the “C” of “CW”), but the compactly generated topology required for a CW complex is in general finer than the product topology. In the 1949 paper introducing CW complexes [9, Section 5], Whitehead showed that if one of the two CW complexes is locally finite (see Definition 7 below), the product is a CW complex, adding in a footnote “I do not know if this restriction [...] is necessary”. Soon afterwards Dowker [2] demonstrated that *some* assumption on X and Y was necessary for the product to be a CW complex: Dowker exhibited a CW complex X with countably many cells and a CW complex Y with continuum (2^{\aleph_0}) many cells such that $X \times Y$ is not a CW complex. On the other hand, Milnor [7] showed that Whitehead’s assumption itself was not necessary, as

Date: February 5, 2022.

the product of any two CW complexes each with countably many cells is a CW complex.

Subsequent advances have relied on making assumptions not just about the CW complexes X and Y , but about the whole domain of discourse: the model of set theory in which X and Y lie. In models of set theory in which the Continuum Hypothesis (the assumption that $2^{\aleph_0} = \aleph_1$) holds, Liu Ying-Ming [6] showed that the product of CW complexes X and Y will be a CW complex if and only if X or Y is locally finite or both are locally countable — that is, if and only if either Whitehead or Milnor’s criterion holds. Building on work of Gruenhage [4], Tanaka [8] showed that the validity of this characterisation is equivalent to a weaker set-theoretic assumption there denoted “not $\text{BF}(\omega_2)$ ”; in current set-theoretic notation, this is $\mathfrak{b} = \aleph_1$.

The present paper provides a complete answer to the question of when the product of a pair of CW complexes is again a CW complex. No extra set-theoretic assumptions are required for our characterisation, and if one does make the set-theoretic assumptions employed by Liu or Tanaka, our characterisation reduces to theirs.

Theorem 1. *Let X and Y be CW complexes. Then $X \times Y$ is a CW complex if and only if one of the following holds:*

- (1) *Either X or Y is locally finite.*
- (2) *Either X or Y has countably many cells in each connected component, and the other has fewer than \mathfrak{b} many cells in each connected component.*

The cardinal \mathfrak{b} in alternative (2) is the so-called *bounding number*: the least cardinality of a set of functions from \mathbb{N} to \mathbb{N} which is unbounded with respect to eventual domination (Definition 3). This relatively concrete definition for a cardinal has been extensively studied by set theorists, and \mathfrak{b} is a standard cardinal in the pantheon of *cardinal characteristics of the continuum* (for more on which see, for example, Blass [1]). The value of \mathfrak{b} in terms of Cantor’s \aleph hierarchy can vary between different models of set theory, much as the cardinality of the reals 2^{\aleph_0} can, so a corollary of our result is that no characterisation purely in terms of the \aleph hierarchy can be valid in all models of set theory. We will introduce \mathfrak{b} in more detail in Section 2.1 below; our proof and presentation require no background in set theory of the reader. We also review the basics of CW complexes in Section 2.2 for readers unfamiliar with them; further details, including a subsection on products where Dowker’s example is presented, may be found in the Appendix of the standard text of Hatcher [5].

The results of Tanaka [8] rely on reducing to the case where the CW complexes in question are “stars” consisting of a central vertex and edges emanating from it. In this one-dimensional setting, the “finitely many errors” aspect of eventual domination is unproblematic, thanks to local compactness away from the “bad” central point. For our full result no such reduction is

possible, and “finitely many errors” can multiply to infinitely many when cells of higher dimension are attached. It turns out that eventual domination is nevertheless the right notion in this case: whilst a naïve induction does not suffice to prove the theorem, the proof of our Lemma 18 shows that with sufficient care, an induction incorporating promises about the growth rate at later stages of the induction can be made to work. Interestingly, such promises are also used in *Hechler forcing*, the typical set-theoretic technique used for building models of set theory with a chosen value of \mathfrak{b} (although no knowledge of Hechler forcing is needed for our arguments).

2. PRELIMINARIES

2.1. The cardinal \mathfrak{b} . Recall that by definition \aleph_1 is the least uncountable cardinal, and 2^{\aleph_0} is the cardinality of the reals; Cantor showed that $2^{\aleph_0} \geq \aleph_1$. There are a number of well-studied definitions for cardinals that lie between these two bounds — so-called *cardinal characteristics of the continuum*. For an introduction to them see, for example, [1]. Of relevance to us is the bounding number \mathfrak{b} .

Definition 2. Given two functions f and g from \mathbb{N} to \mathbb{N} , say that f is *eventually dominated* by g , written $f \leq^* g$, if for all but finitely many n in \mathbb{N} , $f(n) \leq g(n)$.

We also write $f \leq g$ to mean that $f(n) \leq g(n)$ for all $n \in \mathbb{N}$.

Definition 3. The *bounding number* \mathfrak{b} is the least cardinality of a set of functions $\mathbb{N} \rightarrow \mathbb{N}$ that is unbounded with respect to eventual domination, that is,

$$\mathfrak{b} = \min\{|\mathcal{F}| : \mathcal{F} \subseteq \mathbb{N}^{\mathbb{N}} \wedge \forall g \in \mathbb{N}^{\mathbb{N}} \exists f \in \mathcal{F} \neg(f \leq^* g)\}.$$

It is a simple exercise to show that \mathfrak{b} is uncountable, and obviously \mathfrak{b} is at most the cardinality of $\mathbb{N}^{\mathbb{N}}$, so $\aleph_1 \leq \mathfrak{b} \leq 2^{\aleph_0}$. If the Continuum Hypothesis holds then of course $\aleph_1 = \mathfrak{b} = 2^{\aleph_0}$, but there are also models of set theory (that is, universes of sets satisfying the standard ZFC axioms) in which $\aleph_1 = \mathfrak{b} < 2^{\aleph_0}$, models in which $\aleph_1 < \mathfrak{b} = 2^{\aleph_0}$, and models in which $\aleph_1 < \mathfrak{b} < 2^{\aleph_0}$.

Recall that a cardinal κ is said to be *singular* if it can be expressed as a “small” union of “small” sets:

$$\kappa = \bigcup_{\alpha < \gamma} I_\alpha,$$

with $\gamma < \kappa$ and $|I_\alpha| < \kappa$ for each $\alpha < \gamma$. A cardinal is said to be *regular* otherwise.

Lemma 4 (Folklore). *The bounding number \mathfrak{b} is regular.*

Proof. Let X be a set of functions from \mathbb{N} to \mathbb{N} of cardinality \mathfrak{b} which is unbounded with respect to eventual domination; enumerate X as $X = \{f_\zeta : \zeta \in \mathfrak{b}\}$. Suppose for the sake of contradiction that \mathfrak{b} can be decomposed

as $\mathfrak{b} = \bigcup_{\alpha < \gamma} I_\alpha$ with $\gamma < \mathfrak{b}$ and $|I_\alpha| < \mathfrak{b}$ for every $\alpha < \gamma$. Then for each α there must be some $g_\alpha : \mathbb{N} \rightarrow \mathbb{N}$ eventually dominating each member of $\{f_\zeta : \zeta \in I_\alpha\}$. But then $\{g_\alpha : \alpha < \gamma\}$ would be an unbounded set of functions of cardinality $\gamma < \mathfrak{b}$, contradicting the minimality of \mathfrak{b} . \square

2.2. CW complexes. We review the basics of CW complexes here; for more detail see, for example, the Appendix of [5]. We also present a result (Proposition 8) that allows the formulation of our main theorem in terms of numbers of cells in connected components, as it is given in Theorem 1.

The n -disc D^n is the closed unit ball in \mathbb{R}^n . Its interior is here denoted $\overset{\circ}{D}^n$ and its boundary is the $(n-1)$ -sphere S^{n-1} . We denote the image of a function φ pointwise on a set X by $\varphi[X]$.

Definition 5. A Hausdorff space X is a *CW complex* if there exist continuous functions $\varphi_\alpha^n : D^n \rightarrow X$ (*characteristic maps*) for α in an arbitrary index set and $n \in \mathbb{N}$ a function of α , such that the following conditions hold.

- (1) The restriction $\varphi_\alpha^n \upharpoonright \overset{\circ}{D}^n$ of φ_α^n to the interior of D^n is a homeomorphism to its image, and X is the disjoint union as α varies of these homeomorphic images $\varphi_\alpha^n[\overset{\circ}{D}^n]$. We denote $\varphi_\alpha^n[D^n]$ by e_α^n and refer to it as an *n -dimensional cell*.
- (2) (*Closure-finiteness*) For each φ_α^n , the image $\varphi_\alpha^n[S^{n-1}]$ of the boundary of D^n is contained in finitely many cells all of dimension less than n .
- (3) The topology on X is the *weak topology*: a set is closed if and only if its intersection with each closed cell $\varphi_\alpha^n[D^n]$ is closed.¹

A compact subset of a CW complex is contained in finitely many cells e_α^n , and each closed cell $\bar{e}_\alpha^n = \varphi_\alpha^n[D^n]$ is compact, so (3) is equivalent to the statement that the topology is *compactly generated* — a set is closed if and only if its intersection with each compact set is closed. We can also restrict to compact sets of the form of a convergent sequence with its limit point: a space is said to be *sequential* if each subset Y is closed precisely when Y contains the limit point of every convergent countable sequence contained in Y . Since each $\varphi_\alpha^n[D^n]$ is sequential, a Hausdorff space satisfying criteria (1) and (2) of Definition 5 is a CW complex if and only if it is sequential.

Compactly generated spaces are sometimes referred to as k -spaces (for example in [4] and [8]), and compactly generated Hausdorff spaces as Kelley spaces (for example in [3]). We have elected to stick with the more descriptive “compactly generated” terminology used, for example, in [5].

A subcomplex A of a CW complex X is a subspace which is a union of cells of X , such that if e_α^n is contained in A then its closure $\varphi_\alpha^n[D^n]$ is contained in A . Clearly such an A forms a CW complex with those characteristic maps φ_α^n from the CW structure on X corresponding to the cells of A . An

¹That is, the topology which is finest possible while having all of the functions φ_α^n continuous.

important example of a subcomplex is the n -skeleton X^n of X for any given $n \in \mathbb{N}$: X^n is the union of all cells e_α^m of X of dimension $m \leq n$. Every subcomplex of X is closed in X .

Each cell e_α of a CW complex X , and hence each point x in X , is contained in a finite subcomplex. Indeed there is clearly a least such subcomplex, constructible by starting with e_α and working down through the dimensions adding only those cells necessary to contain the boundaries of those that came before.

Definition 6. For X a CW complex and $e_{X,\alpha}$ a cell of X , we denote by X_α^{\min} the minimal (with respect to inclusion) subcomplex of X containing e_α .

There is more variability, however, if we want x to be in the *interior* of the subcomplex.

Definition 7. Let κ be a cardinal. A CW complex X is said to be *locally less than κ* if for all x in X there is a subcomplex A of X with fewer than κ many cells such that x is in the interior of A . We write *locally finite* for locally less than \aleph_0 , and *locally countable* for locally less than \aleph_1 .

Proposition 8. Let κ be an uncountable regular cardinal. Then a CW complex X is locally less than κ if and only if each connected component of X contains fewer than κ many cells.

Proof. Clearly if each connected component of a CW complex X has fewer than κ many cells, then X is locally less than κ . For the converse, let κ be an uncountable regular cardinal, let X be a locally less than κ CW complex, and consider any point x in X ; we shall show that the connected component of X containing x contains fewer than κ many cells. The argument is by a recursive construction of the component.

For the base case, take a connected subcomplex A_1 of X with fewer than κ many cells which contains x in its interior. For notational convenience let A_0 be the empty space. Now suppose we have defined A_i a subcomplex of X with fewer than κ many cells, with x in its interior, and moreover with every point of A_{i-1} in its interior. Consider a cell e of A_i . Since X is locally less than κ , we may choose for every point y of \bar{e} a connected subcomplex A_y of X with fewer than κ many cells and an open set U_y of X such that $y \in U_y \subseteq A_y$. In particular, $U_y \cap \bar{e}$ is open in \bar{e} , and if $z \in U_y \cap \bar{e}$ then certainly z is in the interior of A_y . Since \bar{e} is compact, a finite set S_e of points y suffice for the sets $U_y \cap \bar{e}$ to cover \bar{e} . Take

$$A_{i+1} = \bigcup_{\substack{e \text{ a cell} \\ \text{of } A_i}} \bigcup_{y \in S_e} A_y.$$

Each A_y has fewer than κ many cells, and the union is over fewer than κ many indices, so by regularity of κ , A_{i+1} has fewer than κ many cells. Each A_y in the union is connected to A_i , so A_{i+1} is connected, and by construction A_i is contained in the interior of A_{i+1} , so the inductive step is complete.

Finally, let $A = \bigcup_{i \in \mathbb{N}} A_i$. Since κ is regular uncountable and each A_i has fewer than κ many cells, A has fewer than κ many cells. Since A is an increasing union of connected spaces it is connected; it is open by construction and closed as a subcomplex of X , so it is a connected component of X . \square

We note that Tanaka [8, Lemma 4], citing the paper [6] (in Chinese) of Liu, misstates this Proposition for local countability, with an erroneous point- and cell-wise version. As an example, take the closed unit interval I built as usual as an edge e connecting two vertices. Attach a 2-sphere to a point x in e (that is, attach a 2-disc using the constant characteristic map taking every point of its boundary circle to x), and attach a 3-sphere to every other point of the 2-sphere. Then every open neighbourhood in e of x intersects the boundary of only one other cell, but this CW complex is not locally countable at x . Fortunately the global version, Proposition 8 with $\kappa = \aleph_1$, suffices for the proofs of Tanaka's results.

We will also require the following basic result, a proof of which may be found in [5, Proposition A.3].

Lemma 9 (Whitehead [9, §5(G)]). *CW complexes are normal.*

3. THE CHARACTERISATION

One direction of Theorem 1 is immediate from the following theorem of Tanaka [8].

Theorem 10 (Tanaka). *The following are equivalent.*

- (1) $\kappa \geq \mathfrak{b}$
- (2) *If $X \times Y$ is a CW complex, then either*
 - (a) *X or Y is locally finite, or*
 - (b) *X or Y is locally countable and the other is locally less than κ .*

In light of Proposition 8, in order to prove Theorem 1 it remains to show that if $\kappa = \mathfrak{b}$, the disjunction ((2a) or (2b)) of statements from Theorem 10 implies that $X \times Y$ is a CW complex. The fact that if X or Y is locally finite then $X \times Y$ is a CW complex was shown by Whitehead [9, §5], and follows from the fact that the Cartesian product of a compactly generated space with a locally compact space is compactly generated (see, for example, [5, Proposition A.15]) — locally finite CW complexes are clearly locally compact.

It is therefore incumbent on us to show that if X is locally countable and Y is locally less than \mathfrak{b} then the product of X and Y is a CW complex. We may consider individual connected components, so for the rest of this section, we shall assume that X is a CW complex with countably many cells, and Y is a CW complex with fewer than \mathfrak{b} many cells.

We now give some definitions that will be useful for the proof. For the sake of expositional clarity various dependencies will be omitted from the notation.

We follow the standard notation from set theory, that when a natural number n is used in place of a set of natural numbers, it denotes the n -element set $\{0, \dots, n-1\}$. For a function $s : I \rightarrow K$, the function that extends s by taking value q on some $\alpha \notin I$ is denoted by $s \cup \{(\alpha, q)\}$.

We start by defining a descending sequence of neighbourhoods $B_n(x)$ open in a cell e that form a neighbourhood base in e of a point x .

Definition 11. Suppose x is a point in a CW complex X , with x lying in an open cell e of dimension d with characteristic map φ , and suppose n is a natural number. Let z be $\varphi^{-1}(x)$, and let $r \in \mathbb{R}$ be the minimum of $1/(n+1)$ and half the distance from z to the boundary of D^d . Then we define $B_n(x)$ to be the image under φ of the open ball of radius r about z in D^d .

The set $B_n(x)$ need not be open as a subset of X — to build an open neighbourhood in X we must also consider higher-dimensional cells whose boundaries intersect $B_n(x)$. For these cells we use the following “collar neighbourhoods”.

Definition 12. Let X be a CW complex, d a natural number, and $U \subseteq X^d$ a subset of X^d which is open in X^d . Let e be a $(d+1)$ -dimensional cell of X with characteristic map φ , and let n be a natural number. We define the open subset $C_n^e(U)$ of \bar{e} by

$$C_n^e(U) = \varphi \left[\left\{ t \cdot \vec{z} : t \in \left(\frac{n}{n+1}, 1 \right] \wedge \vec{z} \in \varphi^{-1}(U) \subseteq S^d \right\} \right],$$

where the \cdot denotes scalar multiplication in the vector space \mathbb{R}^{d+1} .

Note that if $\varphi^{-1}(U)$ is empty then $C_n^e(U)$ will also be empty, and that C_n^e distributes over unions: for any U and V , $C_n^e(U \cup V) = C_n^e(U) \cup C_n^e(V)$.

Definition 13. Suppose X is a CW complex with its cells enumerated as e_i for i in some index set I , and for each i in I let $d(i)$ be the dimension of e_i . Then for each $n \in \mathbb{N}$ we let

$$I^n = \{i \in I : d(i) \leq n\}.$$

Thus, for finite n the n -skeleton X^n is the union over i in I^n of the cells e_i . Using these notions, we may define an open neighbourhood of a point from a function to the naturals.

Definition 14. Let X be a CW complex with its cells enumerated as e_i for i in some index set I , and for each i let $d(i)$ be the dimension of e_i . Let x be a point of X , lying in cell e_{i_0} . Then for any function $f : I \rightarrow \mathbb{N}$ we define the open neighbourhood $U^X(x; f)$, or simply $U(x; f)$ when X is clear, of x in X recursively in dimension as follows.

- For all i in $I^{d(i_0)}$ other than i_0 , we take $U^X(x; f) \cap e_i = \emptyset$.
- For $i = i_0$, we take $U^X(x; f) \cap e_i = B_{f(i)}(x)$.
- If $U^X(x; f) \cap X^m$ has been defined for some $m \geq d(i_0)$, and $i \in I$ is such that $d(i) = m+1$, we set

$$U^X(x; f) \cap \bar{e}_i = C_{f(i)}^{e_i}(U^X(x; f) \cap X^m).$$

Clearly every such set $U^X(x; f)$ is open in X . Note also that if A is a subcomplex of X and $J \subseteq I$ is the set of indices of cells in A , $J = \{i \in I : e_i \subseteq A\}$, then $U^A(x; f \upharpoonright J) = U^X(x; f) \cap A$. We thus use the notation $U(x; f)$ omitting the superscript without fear of confusion, with the domain of f dictating the CW complex in which $U(x; f)$ is taken.

For functions $f: I \rightarrow \mathbb{N}$ we shall write $f \downarrow n$ as a shorthand for the restriction $f \upharpoonright I^n$; thus, $U(x; f \downarrow n) = U(x; f) \cap X^n$. In the arguments below we shall even use this notation when f has not yet been defined on $I \setminus I^n$. Also, as per the set-theoretic convention discussed above, $f \upharpoonright i$ denotes the restriction of f to natural numbers less than i , $f \upharpoonright i = f \upharpoonright \{0, \dots, i-1\}$.

Since each $U(x; f)$ for $f: I \rightarrow \mathbb{N}$ is open, it will suffice for our proof of Theorem 1 to produce sets of this form. In some sense this is also necessary:

Lemma 15. *For any CW complex X with cells e_i , $i \in I$, and for any x in X , the sets $U(x; f)$ as f varies over functions from I to \mathbb{N} form an open neighbourhood base at x .*

Proof. Given an open neighbourhood V of x , we construct recursively on dimension a function $f: I \rightarrow \mathbb{N}$ such that $\overline{U(x; f) \cap X^n} \subset V \cap X^n$ for every $n \in \mathbb{N}$. If x is in cell e_{i_0} of dimension $d(i_0)$, then as the base case we may choose $f(i_0)$ large enough that $B_{f(i_0)}(x)$ has closure contained in V , since $V \cap e_{i_0}$ is open in e_{i_0} , and set $f(i) = 0$ for every other i in $I^{d(i_0)}$. For the inductive step, suppose we have defined f on I^n in such a way that $\overline{U(x; f \downarrow n)} \subset V \cap X^n$, and suppose e_ℓ is an $(n+1)$ -cell of X with characteristic map φ_ℓ . Then $\varphi_\ell^{-1}(\overline{U(x; f \downarrow n)})$ is a compact subset of $\varphi_\ell^{-1}(V) \cap S^n$, and thus we may choose $f(\ell)$ sufficiently large that $C_{f(\ell)}^{e_\ell}(U(x; f \downarrow n))$ also has closure contained in $\varphi_\ell^{-1}(V)$. \square

We shall repeatedly require the following lemma allowing us to extend open sets on finite subcomplexes.

Lemma 16. *Suppose W and Z are CW complexes, W' is a finite subcomplex of W , Z' is a finite subcomplex of Z , U is a subset of W' that is open in W' , V is a subset of Z' that is open in Z' , and H is a sequentially closed subset of $W \times Z$ such that the closure of $U \times V$ is disjoint from H . Let e be a cell of Z whose boundary is contained in Z' . Then there is a $p \in \mathbb{N}$ such that $U \times (V \cup C_p^e(V))$ has closure disjoint from H .*

The point is that $V \cup C_p^e(V)$ is open in $Z' \cup e$, and we can build up open sets in the full CW complex Z in this way. Note also that apart from which CW complex e belongs to, Lemma 16 is symmetric in W and Z , so we will be able to use it to build up open sets of both X and Y in the proof of the main theorem.

Proof. Denote the subcomplex $Z' \cup e$ of Z by $Z'e$. The product $W' \times Z'e$ is a compact CW complex, and in particular normal and sequential. Thus $H \cap (W' \times Z'e)$ is a closed subset of $W' \times Z'e$ disjoint from $\overline{U \times V}$, and

so we may take disjoint open sets $\mathcal{O}_{U \times V}$ and \mathcal{O}_H in $W' \times Z'e$ such that $\overline{U \times V} \subseteq \mathcal{O}_{U \times V}$ and $H \cap (W' \times Z'e) \subseteq \mathcal{O}_H$. Now, for every point (u, v) of $\overline{U \times V}$, there is an open base set $R \times S$ of the product topology on $W' \times Z'e$ that contains (u, v) and is contained in $\mathcal{O}_{U \times V}$. By shrinking S if necessary, we may assume S is of the form $T \cup C_n^e(T)$ for some open subset T of Z' and some $n \in \mathbb{N}$ (recall that this also makes sense if $T \cap \bar{e}$ is empty, in which case n is arbitrary). Now, by compactness of $\overline{U \times V}$, finitely many such base sets $R \times S$ suffice to cover $\overline{U \times V}$, and we may choose $p \in \mathbb{N}$ to be strictly greater than all of the corresponding values n . Then $U \times (V \cup C_p^e(V))$ has closure contained in $\mathcal{O}_{U \times V}$, and hence disjoint from H , as required. \square

We are now ready to prove our main result. By Proposition 8, the formulation given here is equivalent to Theorem 1.

Theorem 17. *Let X and Y be CW complexes. Then $X \times Y$ is a CW complex if and only if one of the following holds:*

- (1) X or Y is locally finite.
- (2) One of X and Y is locally countable, and the other is locally less than \mathfrak{b} .

Proof. As discussed at the start of this section, it suffices to show that if X has countably many cells and Y has fewer than \mathfrak{b} many cells, then $X \times Y$ is a CW complex. So suppose X is a CW complex with countably many cells and Y is a CW complex with fewer than \mathfrak{b} many cells. We shall show that the product topology on $X \times Y$ is sequential, and so indeed makes $X \times Y$ a CW complex. To this end, let H be an arbitrary sequentially closed subset of $X \times Y$, and take $(x_0, y_0) \in X \times Y \setminus H$; we wish to construct an open neighbourhood of (x_0, y_0) disjoint from H .

Enumerate the cells of X as $e_{X,i}$ for i in \mathbb{N} , in such a way that for each i , the boundary of $e_{X,i}$ is contained in $\bigcup_{j < i} e_{X,j}$ — this is possible by closure-finiteness. We define the finite subcomplex X_i of X to be

$$X_i = \bigcup_{j \leq i} e_{X,j}.$$

Enumerate the cells of Y as $e_{Y,\alpha}$ for α in some index set J with cardinality $\mu < \mathfrak{b}$ (we leave J abstract rather than declaring $J = \mu$ so that the notation J^n of Definition 13 remains clear). Recall our notation Y_α^{\min} from Definition 6 for the minimal subcomplex of Y containing $e_{Y,\alpha}$. Let $m(i)$ be the dimension of cell $e_{X,i}$, and let $n(\alpha)$ be the dimension of cell $e_{Y,\alpha}$. Let e_{X,i_0} be the unique open cell of X containing x_0 , and e_{Y,α_0} the unique open cell of Y containing y_0 . We shall construct functions $f: \mathbb{N} \rightarrow \mathbb{N}$ and $g: J \rightarrow \mathbb{N}$ such that $U(x_0; f) \times U(y_0; g)$ is disjoint from H . As ever, the construction is by recursion, but we shall recurse over dimension on the Y side and over i on the X side, whilst also keeping track of a lower bound function for the X side. Specifically, we shall construct for each i in \mathbb{N} functions $f_i: \mathbb{N} \rightarrow \mathbb{N}$ and $g_i: J^{n(\alpha_0)+i} \rightarrow \mathbb{N}$ such that

- (a) $U(x_0; f_i) \times U(y_0; g_i)$ has closure disjoint from H ,
- (b) for all $j > i$, $g_j \downarrow n(\alpha_0) + i = g_i$, $f_j \upharpoonright i = f_i \upharpoonright i$, and for all $n \geq i$, $f_j(n) \geq f_i(n)$.

With such functions in hand we may define f and g by $f(i) = f_{i+1}(i)$ and $g(\alpha) = g_{n(\alpha)-n(\alpha_0)}(\alpha)$. Then

$$\begin{aligned} U(x_0; f) \times U(y_0; g) &= \bigcup_{i \in \mathbb{N}} U(x_0; f \upharpoonright i) \times U(y_0; g \downarrow n(\alpha_0) + i) \\ &= \bigcup_{i \in \mathbb{N}} U(x_0; f_i \upharpoonright i) \times U(y_0; g_i), \end{aligned}$$

each term of which will be disjoint from H by construction.

Aside. The reader familiar with Hechler forcing (the simplest forcing partial order used for constructing models of set theory with a specified value of \mathfrak{b}) will note that the requirements on the functions f_i in (b) are that they form a descending sequence of Hechler conditions, where the stem of f_i is $f_i \upharpoonright i$.

For the base case of the construction, consider $X \times Y_{\alpha_0}^{\min}$. Since $Y_{\alpha_0}^{\min}$ is a finite CW complex, $X \times Y_{\alpha_0}^{\min}$ is a CW complex, $(X \times Y_{\alpha_0}^{\min}) \cap H$ is closed, and we may choose a function $f_0: \mathbb{N} \rightarrow \mathbb{N}$ and a natural number number $g_0(\alpha_0)$ such that $U(x_0; f_0) \times B_{g_0(\alpha_0)}(y_0)$ has closure disjoint from H . For $\alpha \neq \alpha_0$ in $J^{n(\alpha_0)}$, set $g_0(\alpha) = 0$, so we have g_0 defined on all of $J^{n(\alpha_0)}$; since $U(y_0; g_0) = B_{g_0(\alpha_0)}(y_0)$, we have that $U(x_0; f_0) \times U(y_0; g_0)$ has closure disjoint from H .

Lemma 18. *Let Y' be a finite subcomplex of Y containing y_0 , let F be a function from \mathbb{N} to \mathbb{N} and s a function from the indices of Y' to \mathbb{N} such that $U(x_0; F) \times U(y_0; s) \subseteq X \times Y'$ has closure disjoint from H . Let i be a natural number and let Y'' be a subcomplex of Y that is a one cell extension of Y' , $Y'' = Y' \cup e_\alpha$. Then there is a function $f: \mathbb{N} \rightarrow \mathbb{N}$ such that*

- (1) $f(n) \geq F(n)$ for all n in \mathbb{N} , and $f(n) = F(n)$ for all $n < i$,
- (2) for every $f': \mathbb{N} \rightarrow \mathbb{N}$ such that $f' \geq^* f$ and $f' \geq F$, there is a $q \in \mathbb{N}$ such that $U(x_0; f') \times U(y_0; s \cup \{(\alpha, q)\})$ has closure disjoint from H .

Proof. The construction of f is by recursion on $n \geq i$, with repeated applications of Lemma 16. As the base case, set $f \upharpoonright i = F \upharpoonright i$.

Suppose we have constructed $f \upharpoonright n$. For every sequence $r: n \rightarrow \mathbb{N}$ such that $F(m) \leq r(m) \leq f(m)$ for all $m < n$, let $q(r)$ be the least $q \in \mathbb{N}$ such that $U(x_0; r) \times U(y_0; s \cup \{(\alpha, q)\})$ has closure disjoint from H ; such a q must exist by assumption on F and s and Lemma 16. Then let $p(r)$ be the least $p \in \mathbb{N}$ such that $U(x_0; r \cup \{(n, p)\}) \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H , again applying Lemma 16. Finally, we define $f(n)$ as

$$f(n) = \max(\{p(r) : F \upharpoonright n \leq r \leq f \upharpoonright n\} \cup \{F(n)\}).$$

We claim that this recursive construction yields a function $f: \mathbb{N} \rightarrow \mathbb{N}$ as per the statement of the lemma.

(1) is immediate from the construction. For (2), suppose $f': \mathbb{N} \rightarrow \mathbb{N}$ is such that $f' \geq^* f$ and $f' \geq F$. Let $n_0 \in \mathbb{N}$ be such that for all $n \geq n_0$, $f'(n) \geq f(n)$. Let r be the n_0 -tuple defined by $r(m) = \min(f(m), f'(m))$. Note that $r \upharpoonright i = f \upharpoonright i = F \upharpoonright i$. The natural number $q(r)$ is then a q as required by (2). Indeed we shall show by induction that, letting f'' be the function

$$f''(n) = \begin{cases} r(n) & \text{if } n < n_0 \\ f(n) & \text{if } n \geq n_0, \end{cases}$$

we obtain that $U(x_0; f'') \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H . The result will then follow, as $f' \geq f''$ and hence $U(x_0; f') \subseteq U(x_0; f'')$.

For the base case, $U(x_0; r) \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H by definition of $q(r)$. For $n \geq n_0$, suppose we have shown that $U(x_0; f'' \upharpoonright n) \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H . Then by minimality $q(f'' \upharpoonright n) \leq q(r)$. Also $f''(n) = f(n) \geq p(f'' \upharpoonright n)$; so $U(x_0; f'' \upharpoonright n+1) \times U(y_0; s \cup \{(\alpha, q(f'' \upharpoonright n))\})$ has closure disjoint from H , whence the possibly smaller set $U(x_0; f'' \upharpoonright n+1) \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H , as required for the inductive step. We therefore have that for every n , $U(x_0; f'' \upharpoonright n) \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H . Since

$$\bigcup_{n \in \mathbb{N}} \overline{U(x_0; f'' \upharpoonright n) \times U(y_0; s \cup \{(\alpha, q(r))\})}$$

is closed in every cell of $X \times Y''$, it is closed in $X \times Y''$, and so $U(x_0; f'') \times U(y_0; s \cup \{(\alpha, q(r))\})$ has closure disjoint from H , as required. \square

Returning to the construction of the functions f_i and g_i for i in \mathbb{N} , suppose that for all $j \leq k$ we have constructed the functions $f_j: \mathbb{N} \rightarrow \mathbb{N}$ and $g_j: J^{n(\alpha_0)+j} \rightarrow \mathbb{N}$ satisfying the requirements (a) and (b) above. Call $\alpha \in J$ *relevant* if $n(\alpha) = n(\alpha_0) + k + 1$; these are the indices we need to extend the definition of g to for the inductive step.

For relevant α , let Y'_α be $(Y_{\alpha_0}^{\min} \cup Y_\alpha^{\min}) \setminus e_\alpha$ and let J_α be the set of indices of cells in Y'_α . Apply Lemma 18 with f_k as F , $Y_{\alpha_0}^{\min} \cup Y_\alpha^{\min}$ as Y'' , Y'_α as Y' , $g_k \upharpoonright J_\alpha$ as s , and $k+1$ as i . The requirement of the lemma that $U(x_0; f_k) \times U(y_0; g_k \upharpoonright J_\alpha)$ have closure disjoint from H holds by the inductive hypothesis. We thus get for each relevant α a function $f_{k+1, \alpha}$ satisfying (1) and (2) of Lemma 18. Since there are fewer than \mathfrak{b} many members of J , there is a single function $f_{k+1}: \mathbb{N} \rightarrow \mathbb{N}$ that eventually dominates $f_{k+1, \alpha}$ for every relevant α . Taking f_{k+1} as f' in (2) of Lemma 18, we have that for each relevant α there is $q_\alpha \in \mathbb{N}$ such that the open subset $U(x_0; f_{k+1}) \times U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})$ of $X \times (Y_{\alpha_0}^{\min} \cup Y_\alpha^{\min})$ has closure disjoint from H . Products commute with closures in the product topology, so

$$\begin{aligned} & \overline{U(x_0; f_{k+1}) \times U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})} \\ &= \overline{U(x_0; f_{k+1})} \times \overline{U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})}, \end{aligned}$$

and since $Y^{n(\alpha_0)+k+1}$ has the weak topology,

$$\begin{aligned} \overline{\bigcup_{\alpha \text{ relevant}} U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})} \\ = \bigcup_{\alpha \text{ relevant}} \overline{U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})}. \end{aligned}$$

So $U(x_0; f_{k+1}) \times \bigcup_{\alpha \text{ relevant}} U(y_0; (g_k \upharpoonright J_\alpha) \cup \{(\alpha, q_\alpha)\})$ is an open subset of $X \times Y^{n(\alpha_0)+k+1}$ with closure disjoint from H . Since $f_{k+1} \geq f_k$, $U(x_0; f_{k+1}) \subseteq U(x_0; f_k)$, and so $U(x_0; f_{k+1}) \times U(y_0; g_k)$ has closure disjoint from H . Taking $g_{k+1} = g_k \cup \{(\alpha, q_\alpha) : \alpha \text{ is relevant}\}$ completes the inductive step.

We thus have a recursive construction of the functions f_i and g_i as required, which as discussed above allows us to form the functions f and g defining an open neighbourhood $U(x_0; f) \times U(y_0; g)$ of (x_0, y_0) disjoint from H . Since H was an arbitrary sequentially closed subset of $X \times Y$ and (x_0, y_0) was an arbitrary point in the complement of H in $X \times Y$, this shows that $X \times Y$ is sequential, and thus bears the weak topology. That is, $X \times Y$ is a CW complex. \square

ACKNOWLEDGEMENTS

The author would like to thank Jeremy Rickard for drawing his attention to this question, and Nicola Gambino, Dugald Macpherson and Severin Mejak for helpful comments on drafts of this paper. The author also gratefully acknowledges the support for this work of the EPSRC through Early Career Fellowship EP/K035703/2, *Bringing set theory and algebraic topology together*.

REFERENCES

- [1] Andreas Blass. Combinatorial cardinal characteristics of the continuum. In Matthew Foreman and Akihiro Kanamori, editors, *The Handbook of Set Theory*, volume 1, pages 395–489. Springer, 2010.
- [2] C. H. Dowker. Topology of metric complexes. *American Journal of Mathematics*, 74:557–577, 1952.
- [3] Peter Gabriel and Michel Zisman. *Calculus of Fractions and Homotopy Theory*. Number 35 in *Ergebnisse der Mathematik und ihrer Grenzgebiete*. Springer, Berlin, 1967.
- [4] Gary Gruenhage. k -spaces and products of closed images of metric spaces. *Proceedings of the American Mathematical Society*, 80(3):478–482, November 1980.
- [5] Allen Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, UK, 2002.
- [6] Liu Ying-Ming. A necessary and sufficient condition for the products of CW-complexes. *Acta Mathematica Sinica*, 21:171–175, 1978. Chinese.
- [7] John Milnor. Construction of universal bundles, I. *Annals of Mathematics*, 63(2):272–284, March 1956.
- [8] Yoshio Tanaka. Products of CW-complexes. *Proceedings of the American Mathematical Society*, 86(3):503–507, November 1982.
- [9] J. H. C. Whitehead. Combinatorial homotopy I. *Bulletin of the American Mathematical Society*, 55(3):213–245, 1949.

SCHOOL OF MATHEMATICS, UNIVERSITY OF LEEDS, LEEDS LS2 9JT, UNITED KING-
DOM