

ON THE ACTION OF THE STEENROD-MILNOR OPERATIONS ON THE INVARIANTS OF THE GENERAL LINEAR GROUPS

NGUYEN THAI HOA, PHAM THI KIM MINH, NGUYEN SUM

1. INTRODUCTION

Let p be an odd prime number. Denote by $GL_n = GL(n, \mathbb{F}_p)$ the general linear group over the prime field \mathbb{F}_p . Each subgroup of GL_n acts on the algebra $P_n = E(x_1, \dots, x_n) \otimes \mathbb{F}_p(y_1, \dots, y_n)$ in the usual manner. Here and in what follows, $E(\cdot, \dots, \cdot)$ and $\mathbb{F}_p(\cdot, \dots, \cdot)$ are the exterior and polynomial algebras over \mathbb{F}_p generated by the indicated variables. We grade P_n by assigning $\dim x_i = 1$ and $\dim y_i = 2$.

Dickson showed in [1] that the invariant algebra $\mathbb{F}_p(y_1, \dots, y_n)^{GL_n}$ is a polynomial algebra generated by the Dickson invariants $Q_{n,s}$, $0 \leq s < n$. Huỳnh Mùi [5, 6] computed the invariant algebras $P_n^{GL_n}$. He proved that $P_n^{GL_n}$ is generated by $Q_{n,s}$, $0 \leq s < n$, R_{n,s_1, \dots, s_k} , $0 \leq s_1 < \dots < s_k < n$. Here R_{n,s_1, \dots, s_k} are Mùi invariants and $Q_{n,s}$ are Dickson invariants (see Section 2).

Let \mathcal{A}_p be the mod p Steenrod algebra and let τ_s, ξ_i be the Milnor elements of dimensions $2p^s - 1$, $2p^i - 2$ respectively in the dual algebra \mathcal{A}_p^* of \mathcal{A}_p . In [4], Milnor showed that as an algebra,

$$\mathcal{A}_p^* = E(\tau_0, \tau_1, \dots) \otimes \mathbb{F}_p(\xi_1, \xi_2, \dots).$$

Then \mathcal{A}_p^* has a basis consisting of all monomials

$$\tau_S \xi^R = \tau_{s_1} \dots \tau_{s_k} \xi_1^{r_1} \dots \xi_m^{r_m},$$

with $S = (s_1, \dots, s_k)$, $0 \leq s_1 < \dots < s_k$, $R = (r_1, \dots, r_m)$, $r_i \geq 0$. Let $St^{S,R} \in \mathcal{A}_p$ denote the dual of $\tau_S \xi^R$ with respect to that basis. Then \mathcal{A}_p has a basis consisting of all operations $St^{S,R}$. For $S = \emptyset, R = (r)$, $St^{\emptyset, (r)}$ is nothing but the Steenrod operation P^r . So, we call $St^{S,R}$ the Steenrod-Milnor operation of type (S, R) .

We have the Cartan formula

$$St^{S,R}(uv) = \sum_{\substack{S_1 \cup S_2 = S \\ R_1 + R_2 = R}} St^{S_1, R_1}(u) St^{S_2, R_2}(v),$$

where $R_1 = (r_{1i})$, $R_2 = (r_{2i})$, $R_1 + R_2 = (r_{1i} + r_{2i})$, $S_1 \cap S_2 = \emptyset$, $u, v \in P_n$ (see Mùi [6]).

We denote $St_u = St^{(u), (0)}$, $St^{\Delta_i} = St^{\emptyset, \Delta_i}$, where $\Delta_i = (0, \dots, 1, \dots, 0)$ with 1 at the i -th place. In [6], Huỳnh Mùi proved that as a coalgebra,

$$\mathcal{A}_p = \Lambda(St_0, St_1, \dots) \otimes \Gamma(St^{\Delta_1}, St^{\Delta_1}, \dots).$$

Here, $\Lambda(St_0, St_1, \dots)$ (resp. $\Gamma(St^{\Delta_1}, St^{\Delta_2}, \dots)$) denotes the exterior (resp. polynomial) Hopf algebra with divided powers generated by the primitive Steenrod-Milnor operations St_0, St_1, \dots (resp. $St^{\Delta_1}, St^{\Delta_2}, \dots$).

The Steenrod algebra \mathcal{A}_p acts on P_n by means of the Cartan formula together with the relations

$$St^{S,R}x_k = \begin{cases} x_k, & S = \emptyset, R = (0), \\ y_k^{p^u}, & S = (u), R = (0), \\ 0, & \text{otherwise}, \end{cases}$$

$$St^{S,R}y_k = \begin{cases} y_k, & S = \emptyset, R = (0), \\ y_k^{p^i}, & S = \emptyset, R = \Delta_i, \\ 0, & \text{otherwise}, \end{cases}$$

for $k = 1, 2, \dots, n$ (see Steenrod-Epstein [8], Sum [11]). Since this action commutes with the action of GL_n , it induces an actions of \mathcal{A}_p on $P_n^{GL_n}$.

The action of $St^{S,R}$ on the modular invariants of subgroups of general linear group has partially been studied by many authors. This action for $S = \emptyset, R = (r)$ was explicitly determined by Smith-Switzer [7], Hung-Minh [2], Kechagias [3], Sum [11, 12, 13, 14]. Smith-Switzer [7], Wilkerson [15] have studied the action of St^{Δ_i} on the Dickson invariants.

The purpose of the paper is to compute the action of the Steenrod-Milnor operations on the generators of $P_2^{GL_2}$. More precisely, we explicitly determine the action of $St^{(i,j)}$ on the Dickson invariants $Q_{2,0}$ and $Q_{2,1}$.

The analogous results for $p = 2$ have been announced in [9].

The rest of the paper contains two sections. In Section 2, we recall some needed information on the invariant theory. In Section 3, we compute the action of the Steenrod-Milnor operations on the Dickson invariants.

2. PRELIMINARIES

Definition 2.1. Let (e_{k+1}, \dots, e_n) , $0 \leq k < n$, be a sequence of non-negative integers. Following Dickson [1], Mùì [5], we define

$$[k; e_{k+1}, \dots, e_n] = \frac{1}{k!} \begin{vmatrix} x_1 & \cdots & x_n \\ \vdots & \cdots & \vdots \\ x_1 & \cdots & x_n \\ y_1^{p^{e_{k+1}}} & \cdots & y_m^{p^{e_{k+1}}} \\ \vdots & \cdots & \vdots \\ y_1^{p^{e_n}} & \cdots & y_m^{p^{e_n}} \end{vmatrix},$$

in which there are exactly k rows of $(x_1 \dots x_n)$. (See the accurate meaning of the determinants in a commutative graded algebra in Mùì [5]). For $k = 0$, we write

$$[0; e_1, \dots, e_n] = [e_1, \dots, e_n] = \det(y_i^{p^{e_j}}).$$

In particular, we set

$$L_{n,s} = [0, 1, \dots, \hat{s}, \dots, n], \quad 0 \leq s \leq n,$$

$$L_n = L_{n,n} = [0, 1, \dots, n-1].$$

Each $[k; e_{k+1}, \dots, e_n]$ is an invariant of the special linear group SL_n and $[e_1, \dots, e_n]$ is divisible by L_n . Then, Dickson invariants $Q_{n,s}$, $0 \leq s < n$, are defined by

$$Q_{n,s} = L_{n,s}/L_n.$$

Here, by convention, $L_0 = [\emptyset] = 1$.

Now we recall the following which will be used in the next section.

Lemma 2.2 (Sum [12]). *The actions of St^R on $L_2, L_{2,0}, L_{2,1}$ for $\ell(R) = 2$ are respectively given in the following tables:*

(i, j)	$St^{(i,j)} L_2$	(i, j)	$St^{(i,j)} L_2$
$(0, 0)$	L_2	$(0, 1)$	$-L_2 Q_{2,0}$
$(0, p)$	$L_2(Q_{2,1}^{p+1} - Q_{2,0}^p)$	$(0, p+1)$	$L_2 Q_{2,0}^{p+1}$
$(1, p)$	$L_2 Q_{2,0} Q_{2,1}^p$	$(p, 0)$	$L_2 Q_{2,1}$
$(p+1, 0)$	$L_2 Q_{2,0}$	<i>otherwise</i>	$0,$

(i, j)	$St^{(i,j)} L_{2,0}$	(i, j)	$St^{(i,j)} L_{2,0}$
$(0, 0)$	$L_2 Q_{2,0}$	$(0, p)$	$-L_2 Q_{2,0}^{p+1}$
$(0, p^2)$	$L_2(Q_{2,0} Q_{2,1}^{p^2+p} - Q_{2,0}^{p^2+1})$	$(0, p^2 + p)$	$L_2 Q_{2,0}^{p^2+p+1}$
(p, p^2)	$L_2 Q_{2,0}^{p+1} Q_{2,1}^{p^2}$	$(p^2, 0)$	$L_2 Q_{2,0} Q_{2,1}^p$
$(p^2 + p, 0)$	$L_2 Q_{2,0}^{p+1}$	<i>otherwise</i>	$0,$

(i, j)	$St^{(i,j)} L_{2,1}$
$(0, 0)$	$L_2 Q_{2,1}$
$(0, p^2)$	$L_2(Q_{2,1}^{p^2+p+1} - Q_{2,0}^p Q_{2,1}^{p^2} - Q_{2,0}^{p^2} Q_{2,1})$
$(0, p^2 + 1)$	$L_2 Q_{2,0}^{p+1} Q_{2,1}^{p^2}$
$(1, 0)$	$L_2 Q_{2,0}$
$(1, p^2)$	$L_2(Q_{2,0} Q_{2,1}^{p^2+p} - Q_{2,0}^{p^2+1})$
$(p^2, 0)$	$L_2(Q_{2,1}^{p+1} - Q_{2,0}^p)$
$(p^2, 1)$	$L_2 Q_{2,0}^{p+1}$
$(p^2 + 1, 0)$	$L_2 Q_{2,0} Q_{2,1}^p$
<i>otherwise</i>	$0.$

3. MAIN RESULTS

First of all, we recall the following.

Proposition 3.1 (Hung-Minh [2], Sum [11]). *Let i be a nonnegative integer. We have*

$$St^{(i,0)}Q_{2,0} = \begin{cases} Q_{2,0}^p, & i = p^2 - 1, \\ (-1)^k \binom{k}{r} Q_{2,0}^{r+1} Q_{2,1}^{k-r}, & i = kp + r, \ 0 \leq r \leq k < p, \\ 0, & \text{otherwise}, \end{cases}$$

$$St^{(i,0)}Q_{2,1} = \begin{cases} Q_{2,1}^p, & i = p^2 - p, \\ (-1)^k \binom{k+1}{r} Q_{2,0}^r Q_{2,1}^{k+1-r}, & i = kp + r, \ r \leq k+1 \leq p, \\ 0, & \text{otherwise}. \end{cases}$$

Theorem 3.2 (Sum [12]). *Let j be a nonnegative integer and $s = 0, 1$. Then we have*

$$St^{(0,j)}Q_{2,s} = \begin{cases} Q_{2,0}^r Q_{2,s} \sum_{i=0}^k (-1)^i \binom{k+s}{i+s} \binom{r+i}{i} Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)}, & \text{if } j = kp + r \text{ with } 0 \leq k, \ r < p, \\ 0, & \text{otherwise}. \end{cases}$$

Our new results are the actions of $St^{(i,j)}$ on $Q_{2,0}$ and $Q_{2,1}$ with $i > 0$.

Theorem 3.3. *Let i, j be nonnegative integers. We have*

(i) *If $0 \leq k < i < p$ and $0 \leq r < p$ then*

$$St^{(i,kp+r)}Q_{2,0} = 0.$$

(ii) *If $0 \leq i < p$ then*

$$St^{(i,ip)}Q_{2,0} = (-1)^i Q_{2,0}^{i+1} Q_{2,1}^{ip}.$$

(iii)

$$St^{(1,j)}Q_{2,0} = \begin{cases} -Q_{2,0}^{r+2} Q_{2,1}^p \sum_{i=0}^{k-1} (-1)^i \left[k \binom{k-1}{i} \binom{r+i+1}{i+1} - \binom{k-2}{i-1} \binom{r+i}{i+1} \right] Q_{2,0}^{(k-1-i)p} Q_{2,1}^{i(p+1)}, & \text{if } j = kp + r \text{ with } 0 \leq k, \ r < p, \\ 0, & \text{otherwise}. \end{cases}$$

(iv) *If $k+1 < i < p$ and $0 \leq r < p$ then*

$$St^{(i,kp+r)}Q_{2,1} = 0.$$

(v) *If $0 \leq k < p$ then*

$$St^{(k+1,kp)}Q_{2,1} = (-1)^k Q_{2,0}^{k+1} Q_{2,1}^{kp}.$$

(vi)

$$St^{(1,j)}Q_{2,1} = \begin{cases} (k+1) Q_{2,0}^{r+1} \sum_{i=0}^k (-1)^i \binom{k}{i} \binom{r+i}{i} Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)} & \text{if } j = kp + r \text{ with } 0 \leq k, \ r < p, \\ 0, & \text{otherwise}. \end{cases}$$

Proof. We prove i) by induction on k . Let $k = 0$. For $r = 0$, using Cartan formula we have $St^{(i,0)}Q_{2,0} = 0$. For $r > 0$, applying the Cartan formula and the inductive hypothesis one gets

$$St^{(i,r)}Q_{2,0} = Q_{2,0}St^{(i,r-1)}Q_{2,0} = 0.$$

Part i) of the theorem holds for $k = 0$. Suppose it is true for $0 \leq k < p - 1$. For $i > k + 1$, we have

$$\begin{aligned} St^{(i,(k+1)p)}Q_{2,0} &= Q_{2,0}St^{(i,kp+p-1)}Q_{2,0} + (Q_{2,0}^p - Q_{2,1}^{p+1})St^{(i,kp)}Q_{2,0} \\ &\quad - Q_{2,0}^{p+1}St^{(i,(k-1)p+p-1)}Q_{2,0} - Q_{2,0}Q_{2,1}^pSt^{(i-1,kp)}Q_{2,0} \\ &= 0. \end{aligned}$$

Part i) of the theorem holds for $i > k + 1$ and $r = 0$. Suppose it is true for $i > k + 1$ and $0 \leq r < p - 1$. Then using the inductive hypothesis we have

$$\begin{aligned} St^{(i,(k+1)p+r+1)}Q_{2,0} &= Q_{2,0}St^{(i,(k+1)p+r)}Q_{2,0} + (Q_{2,0}^p - Q_{2,1}^{p+1})St^{(i,kp+r+1)}Q_{2,0} \\ &\quad - Q_{2,0}^{p+1}St^{(i,kp+r)}Q_{2,0} - Q_{2,0}Q_{2,1}^pSt^{(i-1,kp+r+1)}Q_{2,0} \\ &= 0. \end{aligned}$$

Part i) of the theorem is proved.

Now we prove Part ii). Obviously, it is true for $i = 0$. Suppose $i > 0$ and Part ii) holds for $i - 1$. Using the Cartan formula and the inductive hypothesis, we have

$$\begin{aligned} St^{(i,ip)}Q_{2,0} &= -Q_{2,0}Q_{2,1}^pSt^{(i-1,(i-1)p)}Q_{2,0} \\ &= -(-1)^{i-1}Q_{2,0}Q_{2,1}^pQ_{2,0}^iQ_{2,1}^{(i-1)p} \\ &= (-1)^iQ_{2,0}^{i+1}Q_{2,1}^{ip}. \end{aligned}$$

The proof of Part iii). It is well known that the Steenrod-Milnor operations are stable. That means $St^{(1,j)}x = 0$ for any homogeneous $x \in P_2$ with $\deg x < j$ (see Mui [5]). Since $\deg Q_{2,0} = p^2 - 1 < p^2$, we have $St^{(1,j)}Q_{2,0} = 0$ for $j \geq p^2$. Suppose that $j < p^2$. Then using the p -adic expansion of j , we have

$$j = kp + r \quad \text{for } 0 \leq k, r < p.$$

Now we prove the theorem by double induction on (k, r) .

If $k = 0$, then using Part i) of Theorem 3.3 we get $St^{(1,r)}Q_{2,0} = 0$, so the theorem is true for $k = 0$. If $r = p - 1$, then $\binom{r+i+1}{i+1} = \binom{r+i}{i+1} = 0$ in \mathbb{F}_p , so we obtain

$$St^{(1,kp+p-1)}Q_{2,0} = 0.$$

If $r = 0$, then the formula in the theorem becomes

$$St^{(1,kp)}Q_{2,0} = -kQ_{2,0}^2Q_{2,1}^p(Q_{2,0}^p - Q_{2,1}^{p+1})^{k-1}.$$

We prove this formula by induction on k . Suppose this formula holds for $0 \leq k < p - 1$. For $r = 0$, we have

$$\begin{aligned} St^{(1,(k+1)p)}Q_{2,0} &= (Q_{2,0}^p - Q_{2,1}^{p+1})St^{(1,kp)}Q_{2,0} - Q_{2,0}Q_{2,1}^pSt^{(0,kp)}Q_{2,0} \\ &= -kQ_{2,0}^2Q_{2,1}^p(Q_{2,0}^p - Q_{2,1}^{p+1})^k - Q_{2,0}^2Q_{2,1}^p(Q_{2,0}^p - Q_{2,1}^{p+1})^k \\ &= -(k+1)Q_{2,0}^2Q_{2,1}^p(Q_{2,0}^p - Q_{2,1}^{p+1})^k. \end{aligned}$$

So the formula holds for $k + 1$ and $r = 0$. Suppose $0 \leq r < p - 1$ and the formula holds for $j = (k + 1)p + r$. Applying the Cartan formula and the inductive hypothesis, one gets

$$\begin{aligned}
St^{(1, (k+1)p+r+1)} Q_{2,0} &= Q_{2,0} St^{(1, (k+1)p+r)} Q_{2,0} + (Q_{2,0}^p - Q_{2,1}^{p+1}) St^{(1, kp+r+1)} Q_{2,0} \\
&\quad - Q_{2,0}^{p+1} St^{(1, kp+r)} Q_{2,0} - Q_{2,0} Q_{2,1}^p St^{(0, kp+r+1)} Q_{2,0} \\
&= -Q_{2,0}^{r+3} Q_{2,1}^p \left(k \sum_{i=0}^k (-1)^i \left[\binom{k}{i} \binom{r+i+1}{i+1} - \binom{k-1}{i} \binom{r+i+2}{i+1} \right. \right. \\
&\quad \left. \left. + \binom{k-1}{i-1} \binom{r+i+1}{i} - \binom{k-1}{i} \binom{r+i+1}{i+1} \right] Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)} \right. \\
&\quad \left. + \sum_{i=0}^k (-1)^i \binom{k}{i} \left[\binom{r+i+1}{i+1} - \binom{r+i+1}{i} \right] Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)} \right. \\
&\quad \left. - \sum_{i=0}^k (-1)^i \left[\binom{k-1}{i-1} \binom{r+i}{i+1} + \binom{k-2}{i-1} \binom{r+i+1}{i+1} \right. \right. \\
&\quad \left. \left. + \binom{k-2}{i-2} \binom{r+i}{i} - \binom{k-2}{i-1} \binom{r+i}{i+1} \right] Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)} \right) \\
&= -Q_{2,0}^{r+3} Q_{2,1}^p \sum_{i=0}^k (-1)^i \left[(k+1) \binom{k}{i} \binom{r+i+2}{i+1} \right. \\
&\quad \left. - \binom{k-1}{i-1} \binom{r+i+1}{i+1} \right] Q_{2,0}^{(k-i)p} Q_{2,1}^{i(p+1)}.
\end{aligned}$$

Parts iv), v) and vi) are proved by the same argument as the previous one. \square

This work was supported in part by a grant of Quy Nhon University Research Project.

REFERENCES

- [1] L. E. Dickson, *A fundamental system of invariants of the general modular linear group with a solution of the form problem*, Trans. Amer. Math. Soc. **12** (1911), 75-98.
- [2] N.H.V. Hung and P.A. Minh, *The action of the mod p Steenrod operations on the modular invariants of linear groups*, Vietnam J. Math. **23** (1995), 39-56.
- [3] N.E. Kechagias, *The Steenrod algebra action on generators of rings of invariants of subgroups of $GL_n(\mathbb{Z}/p\mathbb{Z})$* , Proc. Amer. Math. Soc. **118** (1993), 943-952.
- [4] J. Milnor, *Steenrod algebra and its dual*, Ann. of Math. **67** (1958), 150-171.
- [5] H. Mui, *Modular invariant theory and the cohomology algebras of symmetric groups*, J. Fac. Sci. Univ. Tokyo Sec. IA Math. **22** (1975), 319-369.
- [6] H. Mui, *Cohomology operations derived from modular invariants*, Math. Z. **193** (1986), 151-163.
- [7] L. Smith and R. Switzer, *Realizability and non-realizability of Dickson algebras as cohomology rings*, Proc. Amer. Math. Soc. **89** (1983), 303-313.
- [8] N.E. Steenrod and D.B.A. Epstein, *Cohomology operations*, Ann. of Math. No. 50, Princeton University Press, 1962.
- [9] N. Sum, *On the action of the Steenrod-Milnor operations on the modular invariants of linear groups*, Japan. J. Math. **18** (1992), 115-137.

- [10] N. Sum, *On the action of the Steenrod algebra on the modular invariants of special linear group*, Acta Math. Vietnam. **18** (1993), 203-213.
- [11] N. Sum, *Steenrod operations on the modular invariants*, Kodai Math. J. **17** (1994), 585-595.
- [12] N. Sum, *On the action of the Steenrod-Milnor operations on the Dickson invariants*, Vinh University Journal of Science **33** (2004), 41-48.
- [13] N. Sum, *On the module structure over the Steenrod algebra of the Dickson algebra*, Quynhon University Journal of Science **1** (2007), 5-15.
- [14] N. Sum, *The action of the primitive Steenrod-Milnor operations on the modular invariants*, Geometry and Topology Monographs, Math. Sci. Publ., Coventry, **11** (2007), 349-367.
- [15] C. Wilkerson, *A primer on the Dickson invariants*, Contemporary Mathematics, Amer. Math. Soc. **19** (1983), 421-434.

Department of Mathematics, Quy Nhon University
170 An Duong Vuong, Quy Nhon, Binh Dinh, Viet Nam