# On generalized $\sigma$ -soluble groups\*

#### Jianhong Huang

School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou, 221116, P.R. China E-mail: jhh320@126.com

#### Bin Hu<sup>†</sup>

School of Mathematics and Statistics, Jiangsu Normal University, Xuzhou 221116, P. R. China E-mail: hubin118@126.com

#### Alexander N. Skiba

Department of Mathematics and Technologies of Programming, Francisk Skorina Gomel State University, Gomel 246019, Belarus

E-mail: alexander.skiba49@gmail.com

#### Abstract

Let  $\sigma = \{\sigma_i | i \in I\}$  be a partition of the set of all primes  $\mathbb{P}$  and G a finite group. Let  $\sigma(G) = \{\sigma_i : \sigma_i \cap \pi(G) \neq \emptyset$ . A set  $\mathcal{H}$  of subgroups of G is said to be a *complete Hall*  $\sigma$ -set of G if every member  $\neq 1$  of  $\mathcal{H}$  is a Hall  $\sigma_i$ -subgroup of G for some  $i \in I$  and  $\mathcal{H}$  contains exactly one Hall  $\sigma_i$ -subgroup of G for every i such that  $\sigma_i \in \sigma(G)$ . We say that G is  $\sigma$ -full if G possesses a complete Hall  $\sigma$ -set. A complete Hall  $\sigma$ -set  $\mathcal{H}$  of G is said to be a  $\sigma$ -basis of G if every two subgroups  $A, B \in \mathcal{H}$  are permutable, that is, AB = BA.

In this paper, we study properties of finite groups having a  $\sigma$ -basis. In particular, we prove that if G has a  $\sigma$ -basis, then G is generalized  $\sigma$ -soluble, that is, G has a complete Hall  $\sigma$ -set and for every chief factor H/K of G we have  $|\sigma(H/K)| \leq 2$ . Moreover, answering to Problem 8.28 in [3], we prove the following

**Theorem A.** Suppose that G is  $\sigma$ -full. Then every complete Hall  $\sigma$ -set of G forms a  $\sigma$ -basis of G if and only if G is generalized  $\sigma$ -soluble and for the automorphism group  $G/C_G(H/K)$ , induced by G on any its chief factor H/K, we have either  $\sigma(H/K) = \sigma(G/C_G(H/K))$  or  $\sigma(H/K) = \{\sigma_i\}$  and  $G/C_G(H/K)$  is a  $\sigma_i \cup \sigma_j$ -group for some  $i \neq j$ .

<sup>\*</sup>Research is supported by an NNSF grant of China (Grant No. 11401264) and a TAPP of Jiangsu Higher Education Institutions (PPZY 2015A013)

<sup>&</sup>lt;sup>†</sup>Corresponding author

 $<sup>^{0}</sup>$ Keywords: finite group, Hall subgroup,  $\sigma$ -semipermutable subgroup,  $\sigma$ -basis, generalized  $\sigma$ -soluble group.

<sup>&</sup>lt;sup>0</sup>Mathematics Subject Classification (2010): 20D10, 20D15

## 1 Introduction

Throughout this paper, all groups are finite and G always denotes a finite group. Moreover,  $\mathbb{P}$  is the set of all primes,  $\pi = \{p_1, \ldots, p_n\} \subseteq \mathbb{P}$  and  $\pi' = \mathbb{P} \setminus \pi$ . If n is an integer, the symbol  $\pi(n)$  denotes the set of all primes dividing n; as usual,  $\pi(G) = \pi(|G|)$ , the set of all primes dividing the order of G.

In what follows,  $\sigma$  is some partition of  $\mathbb{P}$ , that is,  $\sigma = \{\sigma_i | i \in I\}$ , where  $\mathbb{P} = \bigcup_{i \in I} \sigma_i$  and  $\sigma_i \cap \sigma_j = \emptyset$  for all  $i \neq j$ . By the analogy with the notation  $\pi(n)$ , we write  $\sigma(n)$  to denote the set  $\{\sigma_i | \sigma_i \cap \pi(n) \neq \emptyset\}$ ;  $\sigma(G) = \sigma(|G|)$ .

A complete set of Sylow subgroups S of G contains exactly one Sylow p-subgroup for each prime p, that is,  $S = \{1, P_1, \dots P_t\}$ , where  $P_i$  is a Sylow  $p_i$ -subgroup of G and  $\pi(G) = \{p_1, \dots, p_t\}$ . The set S is said to be a Sylow basis of G provided every two subgroups  $P, Q \in S$  are permutable, that is, PQ = QP.

In general, we say that a set  $\mathcal{H}$  of subgroups of G is a complete  $Hall\ \sigma$ -set of G [3] if every member  $\neq 1$  of  $\mathcal{H}$  is a Hall  $\sigma_i$ -subgroup of G for some  $\sigma_i \in \sigma$  and  $\mathcal{H}$  contains exactly one Hall  $\sigma_i$ -subgroup of G for every  $\sigma_i \in \sigma(G)$ . We say that G is  $\sigma$ -full [3] if G possesses a complete Hall  $\sigma$ -set. A complete Hall  $\sigma$ -set  $\mathcal{H}$  of G is said to be a  $\sigma$ -basis of G [2] if every two subgroups  $A, B \in \mathcal{H}$  are permutable, that is, AB = BA.

In this paper we deal with the following two generalizations of solubility.

#### **Definition 1.1.** We say that G is:

- (i)  $\sigma$ -soluble [2] if  $|\sigma(H/K)| = 1$  for every chief factor H/K of G.
- (i) generalized  $\sigma$ -soluble if G is  $\sigma$ -full and  $|\sigma(H/K)| \leq 2$  for every chief factor H/K of G.

Before continuing, consider three classical cases.

- **Example 1.2.** (i) In the classical case, when  $\sigma = \sigma^0 = \{\{2\}, \{3\}, \ldots\}$ , G is  $\sigma^0$ -soluble if and only if it is soluble. Note also that in view of the Burnside's  $p^a q^b$ -theorem, G is  $\sigma^0$ -soluble if and only if it is generalized  $\sigma^0$ -soluble.
- (ii) In the other classical case, when  $\sigma = \sigma^{\pi} = \{\pi, \pi'\}$  G is:  $\sigma^{\pi}$ -soluble if and only if G is  $\pi$ -separable; generalized  $\sigma^{\pi}$ -soluble if and only if G has both a Hall  $\pi$ -subgroup and a Hall  $\pi'$ -subgroup.
- (iii) In fact, in the theory of  $\pi$ -soluble groups  $(\pi = \{p_1, \ldots, p_n\})$  we deal with the partition  $\sigma = \sigma^{0\pi} = \{\{p_1\}, \ldots, \{p_n\}, \pi'\}$  of  $\mathbb{P}$ . Note that G is  $\sigma^{0\pi}$ -soluble if and only if it is  $\pi$ -soluble. Note also that G is generalized  $\sigma^{0\pi}$ -soluble if and only if the following hold: (a) G has both a Hall  $\pi$ -subgroup E and a Hall  $\pi'$ -subgroup, and (b)  $(H \cap E)K/K$  is a Sylow subgroup of H/K for every chief factor H/K of G.

The well-known Hall's theorem states that G is soluble if and only if it has a Sylow basis. This classical result makes natural to ask: Does it true that G is  $\sigma$ -soluble if and only if it has a  $\sigma$ -basis?

A partial confirmation of this hypothesis is given by the following

**Theorem 1.3** (See Theorem A in [2]). Every  $\sigma$ -soluble group possesses a  $\sigma$ -basis

Now consider the following

**Example 1.4.** Let  $\sigma = \{\{2,3\}, \{5\}, \{2,3,5\}'\}$ , and let  $G = A_5 \times (C_{11} \rtimes \operatorname{Aut}(C_{11}))$ , where  $A_5$  is the alternating group of degree 5 and  $C_{11}$  is a group of order 11. Let  $A_4 \simeq A \leq A_5$  and B a Sylow 5-subgroup of  $A_5$ . Let  $H_1 = AC_2$ ,  $H_2 = BC_5$  and  $H_3 = C_{11}$ , where  $C_5 \times C_2 = \operatorname{Aut}(C_{11})$ . Then the set  $\{H_1, H_2, H_3\}$  is a  $\sigma$ -basis of G. Nevertheless, G is not  $\sigma$ -soluble. This example shows that in general the answer to above question is negative.

Nevertheless, the following result is true.

**Theorem A.** Suppose that G possesses a complete Hall  $\sigma$ -set  $\mathcal{H}$ .

- (i) If  $\mathcal{H}$  is a  $\sigma$ -basis of G, then G is generalized  $\sigma$ -soluble.
- (ii) If  $H \leq A \in \mathcal{H}$  and  $HV^x = V^x H$  for all  $x \in G$  and all  $V \in \mathcal{H}$  such that (|H|, |V|) = 1, then  $H^G$  is generalized  $\sigma$ -soluble.

In view of the Burnside's  $p^a q^b$ -theorem, in the case where  $\sigma = \{\{2\}, \{3\}, \ldots\}$  we get from Theorem A(ii) the following

Corollary 1.5 (Isaaks [6]). If a p-subgroup H of G permutes with all Sylow subgroups P of G such that (p, |P|) = 1, then  $H^G$  is soluble.

Corollary 1.4. If  $H \in \mathcal{H}$  and  $HV^x = V^x H$  for all  $x \in G$  and all  $V \in \mathcal{H}$  such that (|H|, |V|) = 1, then G is  $\sigma'_i$ -soluble where  $\sigma(H) = {\sigma_i}$ .

Corollary 1.6 (Borovikov [7]). If a Sylow p-subgroup P of G is permutes with all Sylow subgroups Q of G such that (p, |Q|) = 1, then G is p'-soluble.

The integers n and m are called  $\sigma$ -coprime if  $\sigma(n) \cap \sigma(m) = \emptyset$ .

In fact, Theorem A(i) is a corollary of the following fact.

- **Theorem B.** Suppose that  $G = A_1A_2 = A_2A_3 = A_1A_3$ , where  $A_1$  is  $\sigma$ -soluble and  $A_2$  and  $A_3$  are generalized  $\sigma$ -soluble subgroups of G. If for some  $i, j, k \in I$  the three indices  $|G : N_G(O^{\sigma_i}(A_1))|$ ,  $|G : N_G(O^{\sigma_i}(A_2))|$ ,  $|G : N_G(O^{\sigma_k}(A_3))|$  are pairwise  $\sigma$ -coprime, then G is generalized  $\sigma$ -soluble.
- Corollary 1.7. Suppose that  $A_1$  is a  $\sigma$ -soluble subgroup and  $A_2$  and  $A_3$  are generalized  $\sigma$ -soluble subgroups of G. If the three indices  $|G:A_1|$ ,  $|G:A_2|$ ,  $|G:A_3|$  are pairwise  $\sigma$ -coprime, then G is generalized  $\sigma$ -soluble.
- Corollary 1.8. Suppose that  $G = A_1A_2 = A_2A_3 = A_1A_3$ , where  $A_1$ ,  $A_2$  and  $A_3$  are soluble subgroups of G. If the three indices  $|G:N_G(A_1)|$ ,  $|G:N_G(A_2)|$ ,  $|G:N_G(A_3)|$  are pairwise coprime, then G is soluble.
- **Corollary 1.8.** (H. Wielandt). If G has three soluble subgroups  $A_1$ ,  $A_2$  and  $A_3$  whose indices  $|G:A_1|$ ,  $|G:A_2|$ ,  $|G:A_3|$  are pairwise coprime, then G is itself soluble.

Following [3], we use  $\mathfrak{H}_{\sigma}$  to denote the class of all  $\sigma$ -full groups G such that every complete Hall  $\sigma$ -set of G forms a  $\sigma$ -basis of G.

In [5, VI, Section 3], Huppert described soluble groups G in which every complete Sylow set of G forms a Sylow basis of G. The results in [5, VI, Section 3] are motivations for the following two questions.

**Question 1.9** (See Problem 8.29 in [3]). Describe groups in  $\mathfrak{H}_{\sigma}$ .

**Question 1.10** (See Problem 8.30 in [3]). Describe  $\sigma$ -soluble groups groups in  $\mathfrak{H}_{\sigma} \cap \mathfrak{S}_{\sigma}$ 

As one more application of Theorem A, we prove the following result, which gives the answer to Question 1.9.

**Theorem C.** Suppose that G is  $\sigma$ -full. Then every complete Hall  $\sigma$ -set of G forms a  $\sigma$ -basis of G if and only if G is generalized  $\sigma$ -soluble and for the automorphism group  $G/C_G(H/K)$ , induced by G on any its chief factor H/K, we have  $|\sigma(G/C_G(H/K))| \leq 2$  and also  $\sigma(H/K) \subseteq \sigma(G/C_G(H/K))$  in the case  $|\sigma(G/C_G(H/K))| = 2$ .

ON the base of Theorem C we prove the following result which gives the answer to Question 1.10.

**Theorem D.** Suppose that G is  $\sigma$ -soluble. Then the following statements are equivalent:

- (i) Every complete Hall  $\sigma$ -set of G forms a  $\sigma$ -basis of G.
- (ii) The automorphism group  $G/C_G(H/K)$ , induced by G on any its chief factor H/K with  $\sigma(H/K) = {\sigma_i}$ , is either a  $\sigma_j$ -group or a  $\sigma_i \cup \sigma_j$ -group for some  $j \neq i$ .
  - (iii)  $G \simeq G^*/R$ , where  $G^* \leq A_1 \times \cdots \times A_t$  for some  $\sigma$ -biprimary  $\sigma$ -soluble groups  $A_1, \ldots, A_t$ .

Note that Satz 3.1 in [5, VI] can be obtained as a special case of Theorem B, when  $\sigma = \{\{2\}, \{3\}, \ldots\}.$ 

Finally, we prove the following

**Theorem E.** If G possesses a complete Hall  $\sigma$ -set  $\mathcal{H}$  with  $|G:N_G(H)|$  is  $\sigma$ -primary for all  $H \in \mathcal{H}$ , then G is generalized  $\sigma$ -soluble.

Corollary 1.11 (See Zhang [11] or Guo [12, Theorem 3]). If for every Sylow subgroup P of G the number  $|G:N_G(P)|$  is a prime power, then G is soluble.

## 2 Preliminaries

We use  $\mathfrak{S}_{g\sigma}$  to denote the class of all  $\sigma$ -soluble groups.

The direct calculations show that the following lemma is true

**Lemma 2.1.** (i) The class  $\mathfrak{S}_{g\sigma}$  is closed under taking products of normal subgroups, homomorphic images and subgroups. Moreover, any extension of the generalized  $\sigma$ -soluble group by a generalized  $\sigma$ -soluble group is generalized a  $\sigma$ -soluble group as well.

- (ii) If G/R,  $G/N \in \mathfrak{S}_{a\sigma}$ , then  $G/R \cap N \in \mathfrak{S}_{a\sigma}$ .
- (iii)  $\mathfrak{S}_{g\sigma} \subseteq \mathfrak{S}_{g\sigma^*}$  for any partition  $\sigma^* = \{\sigma_j^* \mid j \in J\}$  of  $\mathbb{P}$  such that  $J \subseteq I$  and  $\sigma_j \subseteq \sigma_j^*$  for all  $j \in J$ .

Recall that G is said to be: a  $D_{\pi}$ -group if G possesses a Hall  $\pi$ -subgroup E and every  $\pi$ -subgroup of G is contained in some conjugate of E; a  $\sigma$ -full group of Sylow type [1] if every subgroup of G is a  $D_{\sigma_i}$ -group for every  $\sigma_i \in \sigma$ .

In view of Theorem B in [2], the following fact is true.

**Lemma 2.1.** If G is  $\sigma$ -soluble, then G is a  $\sigma$ -full group of Sylow type.

Let  $\Pi \subseteq \sigma$ . A natural number n is said to be a  $\Pi$ -number if  $\sigma(n) \subseteq \Pi$ . A subgroup A of G is said to be: a  $\Pi$ -subgroup of G if  $\sigma(G) \subseteq \Pi$ ; a Hall  $\Pi$ -subgroup of G [1] if |A| is a  $\Pi$ -number and |G:A| is a  $\Pi'$ -number

We use  $\mathfrak{H}_{\sigma}$  to denote the class of all groups G such that G has a complete Hall  $\sigma$ -set  $\mathfrak{H} = \{H_1, \ldots, H_t\}$  satisfying the condition  $H_i^x H_j^y = H_j^y H_i^x$  for all  $x, y \in G$  and all  $i \neq j$ . Note that in view of Lemma 2.1, each  $\sigma$ -biprimary  $\sigma$ -soluble group belongs to the class  $\mathfrak{H}_{\sigma}$ .

**Lemma 2.2.** The class  $\mathfrak{H}_{\sigma}$  is closed under taking homomorphic images and and direct products. Moreover, if G is a  $\sigma$ -full group of Sylow type and  $E \leq G \in \mathfrak{H}_{\sigma}$ , then  $E \in \mathfrak{H}_{\sigma}$ .

**Proof.** Let R be a normal subgroup of  $G \in \mathfrak{H}_{\sigma}$ . By hypothesis, G has a complete Hall  $\sigma$ -set  $\mathcal{H} = \{H_1, \ldots, H_t\}$  satisfying the condition  $H_i^x H_j^y = H_j^y H_i^x$  for all  $x, y \in G$  and all  $i \neq j$ .

Then  $\mathcal{H}_0 = \{H_1R/R, \dots, H_tR/R\}$  is a complete Hall  $\sigma$ -set of G/R such that

$$(H_i R/R)^{xR} (H_j R/R)^{yR} = H_i^x H_j^y R/R = H_j^y H_i^x R/R$$
  
=  $(H_j R/R)^{yR} (H_i R/R)^{xR}$ .

Thus  $G/R \in \mathfrak{H}_{\sigma}$ . Therefore the class  $\mathfrak{H}_{\sigma}$  is closed under taking homomorphic images.

Now we show that the class  $\mathfrak{H}_{\sigma}$  is closed under taking direct products. It is enough to show that if  $A, B \in \mathfrak{H}_{\sigma}$ , then  $G = A \times B \in \mathfrak{H}_{\sigma}$ . Let  $\mathcal{A} = \{A_1, \dots, A_n\}$  be a complete Hall  $\sigma$ -set of A and  $\mathcal{B} = \{B_1, \dots, B_m\}$  be a complete Hall  $\sigma$ -set of B such that  $A_i^{a_1} A_j^{a_2} = A_j^{a_2} A_i^{a_1}$  for all  $a_1, a_2 \in A$  and all  $i \neq j$  and  $B_i^{b_1} B_j^{b_2} = B_j^{b_2} B_i^{b_1}$  for all  $b_1, b_2 \in B$  and all  $i \neq j$ . We can assume without loss of generality that  $1 \in \mathcal{A} \cap \mathcal{B}$  and that  $\sigma(G) = \{\sigma_1, \dots, \sigma_t\}$ . Therefore for every i there are indices  $a_i$  and  $b_i$  such that  $A_{a_i} \times B_{b_i}$  is a Hall  $\sigma_i$ -subgroup of G. Moreover, if  $x = a_1b_1$  and  $y = a_2b_2$ , where  $a_1, a_2 \in A$  and  $b_1, b_2 \in B$ , then

$$(A_{a_i} \times B_{b_i})^x (A_{a_j} \times B_{b_j})^y = (A_{a_i}^{a_1} \times B_{b_i}^{b_1}) (A_{a_j}^{a_2} \times B_{b_j}^{b_2})$$
$$= (A_{a_j}^{a_2} \times B_{b_j}^{b_2}) (A_{a_i}^{a_1} \times B_{b_i}^{b_1}) = (A_{a_j} \times B_{b_j})^y (A_{a_i} \times B_{b_i})^x.$$

Hence  $A_{a_1} \times B_{b_1}, \ldots, A_{a_t} \times B_{b_t}$  is a  $\sigma$ -basis of G. Thus  $G \in \mathfrak{H}_{\sigma}$ .

Finally, assume that G is a  $\sigma$ -full group of Sylow type. Then for every complete Hall  $\sigma$ -set  $\mathcal{E} = \{E_1, \ldots, E_r\}$  of E there is a complete Hall  $\sigma$ -set  $\mathcal{H} = \{H_1, \ldots, H_t\}$  of G such that  $E_i = H_i \cap E$  for all  $i = 1, \ldots t$ . We can assume without loss of generality that  $E_i$  is a  $\sigma_i$ -group.

Now let  $\Pi = \{\sigma_i, \sigma_j\}$ . Then for  $x, y \in E$  we have  $\langle E_i^x, E_j^y \rangle \leq E \cap H_i^x H_j^y$ , where  $E \cap H_i^x H_j^y$  is a Pi-subgroup of E and so  $|E \cap H_i^x H_j^y| \leq |E_i^x| |E_j^y|$ . Hence  $\langle E_i^x, E_j^y \rangle = E_i^x E_j^y$ . Thus  $E \in \mathfrak{H}_{\sigma}$ .

The lemma is proved.

The next lemma is evident.

**Lemma 2.3.** If the chief factors H/K and T/L of G are G-isomorphic, then  $C_G(H/K) = C_G(T/L)$ .

We use  $\mathfrak{X}_{\sigma}$  to denote the class of all generalized  $\sigma$ -soluble groups G such that for the automorphism group  $G/C_G(H/K)$ , induced by G on any its chief factor H/K, we have  $|\sigma(G/C_G(H/K))| \leq 2$  and also  $\sigma(H/K) \subseteq \sigma(G/C_G(H/K))$  in the case  $|\sigma(G/C_G(H/K))| = 2$ .

**Lemma 2.4.** The class  $\mathfrak{X}_{\sigma}$  is closed under taking homomorphic images, direct products and subgroups.

**Proof.** Let R be a normal subgroup of  $G \in \mathfrak{X}_{\sigma}$ . Then G/R is generalized  $\sigma$ -soluble and for any chief factor (H/R)/(K/R) of G/R we have  $C_{G/R}((H/R)/(K/R)) = C_G(H/K)/R$ , where H/K is a chief factor of G. Hence  $G/R \in \mathfrak{X}_{\sigma}$ , so the class  $\mathfrak{X}_{\sigma}$  is closed under taking homomorphic images.

Now let  $G = A_1 \times A_2$ , where  $A_1, A_2 \in \mathfrak{X}_{\sigma}$ . The Jordan-Hölder theorem for groups with operators [8, A, 3.2] implies that for every chief factor H/K of G, there is i such that some chief factor T/L of G below  $A_i$  is G-isomorphic to H/K. Moreover, T/L is a chief factor of  $A_i$  and  $G/C_G(T/L) \simeq A_i/C_{A_i}(T/L)$ . Hence, by Lemma 2.3,  $G \in \mathfrak{X}_{\sigma}$ , so the class  $\mathfrak{X}_{\sigma}$  is closed under taking direct products.

Finally, we show that if  $G \in \mathfrak{X}_{\sigma}$  and  $E \leq G$ , then  $E \in \mathfrak{X}_{\sigma}$ .

Let  $1 = G_0 < G_1 < \cdots < G_{t-1} < G_t = G$  be a chief series of G. Let H/K be any chief factor of E such that  $G_{l-1} \cap E \leq K < H \leq G_l \cap E$  for some l.

Since

$$C_G(G_l/G_{l-1}) \cap E \le C_E(G_l \cap E/G_{l-1} \cap E) \le C_E(H/K),$$

 $|\sigma(E/C_E(H/K))| \leq 2$ . Moreover, if  $|\sigma(E/C_E(H/K))| = 2$ , then  $|\sigma(G/C_G(G_l/G_{l-1}))| = 2$ ,  $\sigma(G/C_G(G_l/G_{l-1})) = \sigma(E/C_E(H/K))$  and  $\sigma(G_l/G_{l-1}) \subseteq \sigma(G/C_G(G_l/G_{l-1}))$ . Hence from the isomorphism  $E \cap G_l/E \cap G_{l-1} \simeq (E \cap G_l)G_{l-1}/G_{l-1}$  we get that  $\sigma(H/K) \subseteq \sigma(E/C_E(H/K))$ . Now applying the Jordan-Hölder theorem for groups with operators [8, A, 3.2] we get that  $E \in \mathfrak{X}_{\sigma}$ . Therefore the class  $\mathfrak{X}_{\sigma}$  is closed under taking subgroups.

The lemma is proved.

A class of groups is a collection  $\mathfrak{X}$  of groups with the property that if  $G \in \mathfrak{X}$  and if  $H \simeq G$ , then  $H \in \mathfrak{X}$ . The symbol  $(\mathfrak{Y})$  [8, p. 264] denotes the smallest class of groups containing  $\mathfrak{Y}$ .

For a class  $\mathfrak{X}$  of groups we define, following [8, p. 264]:

 $S(\mathfrak{X}) = (G : G \leq H \text{ for some } H \in \mathfrak{X});$ 

 $Q(\mathfrak{X}) = (G : G \text{ is an epimorphic image of some } H \in \mathfrak{X});$ 

 $D_0(\mathfrak{X}) = (G: G = H_1 \times \cdots \times H_r \text{ for some } H_1, \dots, H_r \in \mathfrak{X});$ 

 $R_0(\mathfrak{X}) = (G: G \text{ has normal subgroups } N_1, \ldots, N_r \text{ with all } G/N_i \in \mathfrak{X}) \text{ and } N_1 \cap \cdots \cap N_r = 1).$ 

We say, following [10], that a class  $\mathfrak{X}$  of groups is a semiformation if

$$S(\mathfrak{X}) = \mathfrak{X} = Q(\mathfrak{X}).$$

The following fact is well-known (see, for example, [13, p. 57]).

**Lemma 2.5.** If  $A \simeq B \leq G$ , then for some  $G^* \simeq G$  we have  $A \leq G^*$ .

**Lemma 2.6.** Let  $\mathfrak{X}$  be a class of groups.

- (1) If  $\mathfrak{X}$  is a semiformation, then  $S(D_0(\mathfrak{X}))=R_0(\mathfrak{X})$ .
- (2)  $Q(R_0(\mathfrak{X})) \subseteq R_0(Q(\mathfrak{X})).$
- (3)  $R_0(R_0(\mathfrak{X})) = R_0(\mathfrak{X}).$

**Proof.** (1) Let  $\mathfrak{M} = \mathrm{S}(\mathrm{D}_0(\mathfrak{X}))$  and  $\mathfrak{H} = \mathrm{R}_0(\mathfrak{X})$ . First suppose that  $G \in \mathfrak{M}$ . Then, in view of Lemma 2.5,  $G \leq H_1 \times \cdots \times H_r$  for some  $H_1, \ldots, H_r \in \mathfrak{X}$ . We show that  $G \in \mathfrak{H}$ . If r = 1, it is clear. Now assume that r > 1 and let  $N_i = H_1 \times \cdots H_{i-1}H_{i+1} \times \cdots \times H_r$  for all  $i = 1, \ldots, r$ . Then

$$G/G \cap N_i \simeq GN_i/N_i \leq (H_1 \times \cdots \times H_r)/N_i \simeq H_i \in \mathfrak{X}.$$

It is clear also that  $G \cap N_1, \ldots G \cap N_r$  are normal subgroups of G with  $(G \cap N_1) \cap \cdots \cap (G \cap N_r) = 1$ . Hence  $G \in \mathfrak{H}$ . Therefore  $\mathfrak{M} \subseteq \mathfrak{H}$ .

Now suppose that  $G \in \mathfrak{H}$ , that is, G has normal subgroups  $N_1, \ldots, N_r$  with all  $G/N_i \in \mathfrak{X}$  and  $N_1 \cap \cdots \cap N_r = 1$ . Then, by Lemma 4.17 in [8, A], G is isomorphic with a subgroup of  $(G/N_1) \times \cdots \times (G/N_r)$ . Hence  $G \in \mathfrak{M}$ , so  $\mathfrak{H} \subseteq \mathfrak{M}$ . Therefore we have (1).

(2), (3) See respectively Lemmas 1.18(b) and Lemma 1.6 in [8, II].

The lemma is proved.

**Lemma 2.7.** Let  $\mathfrak{X}$  be a semiformation of groups. Suppose that  $\mathfrak{F}$  is the class of groups A which can be represented in the form  $A \simeq A^*/R$ , where  $A^* \leq A_1 \times \cdots \times A_t$  for some  $A_1, \ldots, A_t \in \mathfrak{X}$ . If  $G/R, G/N \in \mathfrak{F}$ , then  $G/(R \cap N) \in \mathfrak{F}$ .

**Proof.** We can assume without loss of generality that  $R \cap N = 1$ . Then, by Lemma 2.5,

$$G \in \mathcal{R}_0(\mathcal{Q}(\mathcal{S}(\mathcal{D}_0(\mathfrak{X})))) = \mathcal{R}_0(\mathcal{Q}(\mathcal{R}_0(\mathfrak{X}))) \subseteq \mathcal{Q}(\mathcal{R}_0(\mathcal{R}_0(\mathfrak{X}))) = \mathcal{Q}(\mathcal{R}_0(\mathfrak{X})) = \mathcal{Q}(\mathcal{D}_0(\mathfrak{X}))) \subseteq \mathfrak{F}.$$

The lemma is proved.

**Lemma 2.8.** Let  $G = RH_1 \cdots H_n$ , where  $[H_i^G, H_j^G] = 1$  for all  $i \neq j$  and R is normal in G. Then  $G \simeq G^*$ , where  $G^* \leq (RH_1) \times \cdots \times (RH_n)$ .

**Proof.** See pages 671–672 in [5].

**Lemma 2.9** (See [14, 2.2.8]). If  $\mathfrak{F}$  is a non-empty formation and N, R be subgroups of G, where N is normal in G.

- (i)  $(G/N)^{\mathfrak{F}} = G^{\mathfrak{N}}N/N$ .
- (ii) If G = RN, then  $G^{\mathfrak{N}}N = R^{\mathfrak{N}}N$ .

**Lemma 2.10.** Suppose that G has a  $\sigma_i$ -subgroup  $A \neq 1$  and a  $\sigma_j$ -subgroup  $B \neq 1$  such that  $AB^x = B^x A$  for all  $x \in G$ . If  $O_{\sigma_i \cup \sigma_j}(G) = 1$ , then  $[A^G, B^G] = 1$ .

**Proof.** By hypothesis,  $A(B^x)^y = (B^x)^y A$  for all  $x, y \in G$  and

$$D = \langle A^{B^x} \rangle \cap \langle (B^x)^A \rangle \le AB^x,$$

where D is subnormal in G by [9, 1.1.9(2)]. Then  $D \leq O_{\sigma_i \cup \sigma_j}(G) = 1$ , so Then  $[A, B^x] \leq [\langle A^{B^x} \rangle, \langle B^x \rangle^A \rangle] \leq D = 1$ . Therefore  $[A^x, B^y] = 1$  for all  $x, y \in G$  and so  $[A^G, B^G] = 1$ . The lemma is proved.

# 3 Proofs of the results

**Proof of Theorem B.** Suppose that this theorem is false and let G be a counterexample with |G| minimal. We can assume without loss of generality that i = 1, j = 2 and k = 3.

(1) If R is a minimal normal subgroup of G, then G/R is generalized  $\sigma$ -soluble. Hence R is the unique minimal normal subgroup of G and R is not generalized  $\sigma$ -soluble. Thus  $C_G(R) = 1$ .

First note that if  $A_i \leq R$ , then for  $j \neq i$  we have  $G = A_i A_j = R A_j$ , so  $G/R \simeq A_j/A_j \cap R$  is generalized  $\sigma$ -soluble.

Now suppose that  $A_i \nleq R$  for all i = 1, 2, 3. For any group A and any  $l \in I$  we have  $O^{\sigma_l}(A) = A^{\mathfrak{G}_{\sigma_l}}$ , where  $\mathfrak{G}_{\sigma_l}$  is the class of all  $\sigma_l$ -groups. Therefore  $O^{\sigma_l}(A)R/R = O^{\sigma_l}(AR/R)$  by Lemma 2.10(ii). Hence

$$N_G(O^{\sigma_l}(A))R/R \le N_G(O^{\sigma_l}(A)R)/R = N_{G/R}(O^{\sigma_l}(AR/R).$$

Hence the three indices  $|(G/R): N_{G/R}((O^{\sigma_1}(A_1))|, |(G/R): N_{G/R}(O^{\sigma_2}(A_2))|, \text{ and } |(G/R): N_{G/R}(O^{\sigma_3}(A_3))|$  are pairwise  $\sigma$ -coprime. On the other hand, clearly,  $A_1R/R \simeq A_1/A_1 \cap R$  is  $\sigma$ -soluble and  $A_iR/R \simeq A_i/A_i \cap R$  is generalized  $\sigma$ -soluble for all i=2,3. Therefore the hypothesis holds for G/R, so G/R is generalized  $\sigma$ -soluble by the choice of G. Hence R is not generalized  $\sigma$ -soluble and R is the unique minimal normal subgroup of G by Lemma 2.1(ii). Therefore, since  $C_G(R)$  is normal in G,  $C_G(R) = 1$ .

(2) At least two of the subgroups  $A_1, A_2, A_3$  are not  $\sigma$ -primary.

Indeed, if  $A_i$  and  $A_j$  are  $\sigma$ -primary for some  $i \neq j$ , then  $G = A_i A_j$  is generalized  $\sigma$ -soluble, which contradicts the choice of G.

(3) If  $A_l$  is  $\sigma$ -soluble and L is a minimal normal subgroup of  $A_l$ , then for some  $t \neq l$  we have  $O^{\sigma_t}(A_t) = 1$ , so  $A_t$  is  $\sigma$ -primary.

We can assume without loss of generality that l=2. Let L be a minimal normal subgroup of  $A_l=A_2$ . Then L is a  $\sigma_i$ -group for some prime i since  $A_2$  is  $\sigma$ -soluble by hypothesis. On the other hand, again by hypothesis,  $|G:N_G(O^{\sigma_1}(A_1))|$  and  $|G:N_G(O^{\sigma_3}(A_3))|$  are  $\sigma$ -coprime, so at least one of the numbers,  $|G:N_G(O^{\sigma_3}(A_3))|$  say, is a  $\sigma'_i$ -number. But  $G=A_2A_3=A_2N_G(O^{\sigma_3}(A_3))$ , so  $|A_2:A_2\cap N_G(O^{\sigma_3}(A_3))|$  is a  $\sigma'_i$ -number. Hence

$$L \le A_2 \cap N_G(O^{\sigma_3}(A_3)) \le N_G(O^{\sigma_3}(A_3)).$$

Therefore

$$R \le L^G = L^{A_2 N_G(O^{\sigma_3}(A_3))} = L^{N_G(O^{\sigma_3}(A_3))} \le N_G(O^{\sigma_3}(A_3))$$

by Claim (1).

Suppose that  $O^{\sigma_3}(A_3) \neq 1$  and let  $H = O^{\sigma_3}(A_3) \cap R$ . If  $H \neq 1$ , then H is a non-identity normal generalized  $\sigma$ -soluble subgroup of R since  $A_3$  is generalized  $\sigma$ -soluble by hypothesis. Hence R is generalized  $\sigma$ -soluble, contrary to Claim (1). Thus H = 1, so  $O^{\sigma_3}(A_3) \leq C_G(R) \leq R$ , so  $O^{\sigma_3}(A_3) = 1$  by Claim (1). This contradiction completes the proof of (3).

Final contradiction. Since  $A_1$  is  $\sigma$ -soluble by hypothesis, Claim (3) implies that one of the subgroups  $A_2$  or  $A_3$ ,  $A_2$  say, is  $\sigma$ -primary. Then  $A_2$  is  $\sigma$ -soluble and so, again by Claim (3), one of the subgroups  $A_1$  or  $A_3$  is also  $\sigma$ -primary. But this is impossible by Claim (2). This contradiction completes the proof of the result.

**Proof of Theorem A.** We can assume without loss of generality that H is a  $\sigma_1$ -group,  $1 \in \sigma_1$  and  $\mathcal{H} = \{H_1, \dots, H_t\}$ , where  $H_i$  is a  $\sigma_i$ -group for all  $i = 1, \dots, t$ . Then t > 2 since otherwise G and  $H^G$  are generalized  $\sigma$ -soluble.

- (i) This assertion, in fact, is a corollary of Theorem D. Indeed, let  $E_i = H_1 \cdots H_{i-1} H_{i+1} \cdots H_t$  for all  $i = 1, \dots, t$ . Then, by induction,  $E_1, \dots, E_t$  are generalized  $\sigma$ -soluble and, by the Feit-Thompson theorem,  $E_1 = H_2 \cdots H_t$  is soluble. Since t > 2, and evidently, the indeces  $|G: E_1|$ ,  $|G: E_2|$ ,  $|G: E_3|$  are pairwise  $\sigma$ -coprime, G is generalized  $\sigma$ -soluble by Corollary 1.6.
- (ii) Assume that this assertion is falls and let G be a counterexample of minimal order. By hypothesis,  $HA^x = A^xH$  for all  $x \in G$  and all  $A \in \mathcal{H}$  such that (|H|, |A|) = 1.
  - (1) For some i > 1 and  $x \in G$  we have  $H_i^x \nleq N_G(H)$ .

Indeed, suppose that for all i > 1 and all  $x \in G$  we have  $H_i^x \leq N_G(H)$ . Then

$$E = (H_2)^G \cdots (H_t)^G \le N_G(H)$$

and so  $H^G = H^{H_1E} = H^{H_1} \le H_1$  is generalized  $\sigma$ -soluble, a contradiction. Hence we have (1).

(2)  $H^GR/R$  is generalized  $\sigma$ -soluble.

It is clear that  $\mathcal{H}_0 = \{H_1R/R, \dots, H_tR/R\}$  is a complete Hall  $\sigma$ -set of G/R and  $HR/R \leq H_1R/R$ . Moreover,

$$(HR/R)(H_iR/R)^{xR} = HH_i^xR/R = H_i^xHR/R = (H_iR/R)^{xR}(HR/R)$$

for all i > 1 and all  $xR \in G/R$ . Therefore HR/R is  $\sigma$ -semipermutable in G/R with respect to  $\mathcal{H}_0$ , so  $(HR/R)^G = H^GR/R$  is generalized  $\sigma$ -soluble by the choice of G.

(3) Every minimal normal subgroup of G is not generalized  $\sigma$ -soluble.

Indeed, if R is generalized  $\sigma$ -soluble, then from the isomorphism  $H^GR/R \simeq H^G/H^G \cap R$  and Claim (2) we get that  $H^G$  is generalized  $\sigma$ -soluble, contrary to the choice of G.

Final contradiction for (ii). Let i > 1. By hypothesis,  $HH_i^x = H_i^x H$  for all  $x \in G$ . Om the other hand,  $O_{\sigma_1 \cup \sigma_i}(G) = 1$  by Claim (3). Therefore  $[H^G, H_i^G] = 1$  by Lemma 2.11, contrary to Claim (1). Hence Assertion (ii) holds.

The theorem is proved.

In fact, Theorem C is a corollary of the following

**Proposition 3.1.** If G is  $\sigma$ -full, then  $\mathfrak{H}_{\sigma} = \mathfrak{X}_{\sigma}$ .

**Proof.** First we show  $\mathfrak{H}_{\sigma} \subseteq \mathfrak{X}_{\sigma}$ . Assume that this is false and let G be a group of minimal orderin  $\mathfrak{H}_{\sigma} \setminus \mathfrak{X}_{\sigma}$ . Then  $|\sigma(G)| > 2$ . By hypothesis, G has a complete Hall  $\sigma$ -set  $\mathcal{H} = \{H_1, \ldots, H_t\}$  satisfying the condition  $H_i^x H_j^y = H_j^y H_i^x$  for all  $x, y \in G$  and all  $i \neq j$ . We can assume without loss of generality that  $H_i$  is a  $\sigma_i$ -group for all  $i = 1, \ldots, t$ . Let R be a minimal normal subgroup of G

(1)  $G/R \in \mathfrak{X}_{\sigma}$ . Hence R is the only minimal normal subgroup of G.

First note that since  $G \in \mathfrak{H}_{\sigma}$ , we have  $G/R \in \mathfrak{H}_{\sigma}$  by Lemma 2.2 and so  $G/R \in \mathfrak{X}_{\sigma}$  by the choice of G. Now assume that G has a minimal normal subgroup  $N \neq R$ . Then  $G/N \in \mathfrak{X}_{\sigma}$ , so from the G-isomorphism  $RN/N \simeq R$ , Lemma 2.3 and the Jordan-Hölder theorem for groups with operators [8, A, 3.2] we get that  $G \in \mathfrak{X}_{\sigma}$ , a contradiction. Hence we have (1).

- (2) G is generalized  $\sigma$ -soluble. Hence R is a R is a  $\sigma_i \cup \sigma_j$ -group for some i, j (This follows from Theorem C(i)).
  - (3)  $O_{\sigma_l \cup \sigma_k}(G) = 1$  for all k, l such that  $\sigma(R) \not\subseteq O_{\sigma_i, \sigma_j}$ . (This follows from Claims (1) and (2)).
  - $(4) \mathfrak{H}_{\sigma} \subseteq \mathfrak{X}_{\sigma}.$

Let  $\sigma_k \in \sigma(G)$  such that  $i \neq k \neq j$ . First assume that  $\sigma(R) = \{\sigma_i, \sigma_j\}$ . Then  $R = (R \cap H_i)(R \cap H_j)$ . Claim (1) implies that  $O_{\sigma_i \cup \sigma_k}(G) = 1$  and  $O_{\sigma_j \cup \sigma_k}(G) = 1$ . Then, by Lemma 2.11,  $H_k^G \leq C_G(H_i)$  and  $H_k^G \leq C_G(H_j)$ . Therefore  $H_k^G \leq C_G(R)$ . Therefore  $G/C_G(R)$  is a  $\sigma_i \cup \sigma_j$ -group and so  $G \in \mathfrak{X}_\sigma$  since  $G/R \in \mathfrak{X}_\sigma$  by Claim (1), a contradiction. Hence we can assume that  $\sigma(R) = \{\sigma_i\}$ .

Let  $l \neq k$  and  $l \neq i \neq k$ . Then  $O_{\sigma_l \cup \sigma_k}(G) = 1$  by Claim (3). Hence  $[H_l^G, H_k^G] = 1$  by Lemma 2.11. Therefore from Claim (1) we get that  $H_l \leq C_G(R)$  for all  $l \neq i$  and so  $G/C_G(R)$  is a  $\sigma_i$ -group. Therefore we again get that  $G \in \mathfrak{X}_{\sigma}$ . This contradiction completes the proof of the inclusion  $\mathfrak{H}_{\sigma} \subseteq \mathfrak{X}_{\sigma}$ .

Now we show that if  $G \in \mathfrak{X}_{\sigma}$ , then VW = WV for every Hall  $\sigma_i$ -subgroup V, every Hall  $\sigma_j$ -subgroup W of G and all  $i \neq j$ . Assume that this is false and let G be a counterexample of minimal order. Then  $|\sigma(G)| > 2$ .

(a) G = RVW. Hence G/R is a  $\sigma_i \cup \sigma_i$ -group

Lemma 2.4 implies that the hypothesis holds for G/R, so the choice of G implies that

$$(VR/R)(WR/R) = (WR/R)(VR/R) = VWR/R.$$

Hence RVW is a subgroup of G. Suppose that RVW < G. For each  $k \neq i, j, H_k \cap R$  is a Hall  $\sigma_k$  subgroup of RVW. Therefore the hypothesis holds for RVW by Lemma 2.4 and so VW = WV, a contradiction. Thus we have (a).

(b) R is the unique minimal normal subgroup of G. Hence R is a p-group for some  $p \in \sigma_i$ .

Indeed, suppose that G has a minimal normal subgroup  $N \neq R$ . Then G/N is a  $\sigma_i \cup \sigma_j$ -group by Claim (1), so  $G \simeq G/R \cap N$  is a  $\sigma_i \cup \sigma_j$ -group. Hence  $|\sigma(G)| = 2$ , a contradiction. Therefore R is the unique minimal normal subgroup of G. Suppose that R is non-abelian. Then  $C_G(R) = 1$ , since  $C_G(R)$  is normal in G. Hence  $G \simeq G/C_G(R)$  is  $\sigma$ -biprimary, a contradiction. Thus we have (b).

(c) R is a Sylow p-subgroup of G.

Assume that R is not a Sylow p-subgroup of G. Since G = RVW, by Claim (5), but  $VW \neq WV$ , it follows that  $R \nleq VW$ . Hence for a Sylow p-subgroup P of G we have  $P \cap V = 1 = P \cap W$ . Hence  $P \nleq RVW = G$ , a contradiction. Thus we have (c).

(d)  $\mathfrak{X}_{\sigma} \subseteq \mathfrak{H}_{\sigma}$ .

In view of Claims (b) and (c), there is a maximal subgroup M of G such that G = RM and  $M_G = 1$ . Hence, for some k, we have  $R = C_G(R) = O_p = H_k$  by [8, A, 15.6]. Then  $\sigma_k \notin \sigma(G/C_G(R))$ , so  $G/C_G(R)) = G/R$  is a  $\sigma_l$ -group for some l since  $G \in \mathfrak{G}_{\sigma}$  by hypothesis. But then  $|\sigma(G)| \leq 2$ . This contradiction completes the proof of the inclusion  $\mathfrak{X}_{\sigma} \subseteq \mathfrak{H}_{\sigma}$ .

From Claims (c) and (d) we get that  $\mathfrak{X}_{\sigma} = \mathfrak{H}_{\sigma}$ . The theorem is proved.

**Proof of Theorem D.** The implications (i)  $\Rightarrow$  (ii) and (ii)  $\Rightarrow$  (i) directly follow from Theorem A.

Let  $\mathfrak{X}$  be the class of all  $\sigma$ -biprimary  $\sigma$ -soluble groups. Then from Lemma 2.2 it follows that if A is a group satisfying  $A \simeq A^*/R$  for some  $A^* \leq A_1 \times \cdots \times A_t$  and  $A_1, \ldots, A_t \in \mathfrak{X}$ , then A satisfies also Condition (i). Thus (iii)  $\Rightarrow$  (i).

Now we show that (i)  $\Rightarrow$  (iii). Assume that this is false and let G be a group of minimal order among the groups which satisfy Condition (i) but do not satisfy Condition (iii). Since G is  $\sigma$ -soluble, it has a complete Hall  $\sigma$ -set  $\mathcal{H} = \{H_1, \ldots, H_t\}$  and, by hypothesis,  $H_i^x H_j^y = H_j^y H_i^x$  for all  $x, y \in G$  and all  $i \neq j$ . We can assume without loss of generality that  $H_i$  is a  $\sigma_i$ -group for all  $i = 1, \ldots, t$ . Then t > 2 since otherwise Condition (iii) holds for G.

Let L be a minimal normal subgroup of G. Since G is  $\sigma$ -soluble, L is  $\sigma$ -primary, L is a  $\sigma_1$ -group say. Let  $\mathfrak{F}$  be the class of groups A which can be represented in the form  $A \simeq A^*/R$ , where  $A^* \leq A_1 \times \cdots \times A_t$  for some  $A_1, \ldots, A_t \in \mathfrak{X}$ . Lemma 2.2 implies that Condition (i) holds for G/L, so  $G/L \in \mathfrak{F}$  by the choice of G. If G has a minimal normal subgroup  $N \neq L$ , then also we have  $G/N \in \mathfrak{F}$ , so  $G \simeq G/1 = G/L \cap N \in \mathfrak{F}$  by Lemma 2.7. Thus L is the unique minimal normal subgroup of G.

Now let  $j \neq i \neq 1 \neq j$ , and let  $A = H_i$  and  $B = H_j$ . Then  $AB^x = B^xA$  for all  $x \in G$ . It is clear also that  $O_{\sigma_i \cup \sigma_j}(G) = 1$ . Then  $[A^G, B^G] = 1$  by Lemma 2.10. Now using Lemma 2.8, we get that  $G \in \mathfrak{F}$ . This contradiction completes the proof of the result.

**Proof of Theorem E.** Assume that this is false and let G be a counterexample of minimal order. Let  $\mathcal{H} = \{H_1, \dots, H_t\}$ . Then t > 2. We can assume without loss of generality that  $H_i$  is a non-identity  $\sigma_i$ -group for all  $i = 1, \dots, t$ . Let  $N_i = N_G(H_i)$  for all  $i = 1, \dots, t$ .

First we show that G is not simple. Assume that G is a simple non-abelian group. Then  $|G:N_i| \neq 1$  is  $\sigma$ -primary for all  $i=1,\ldots,t$ .

Let  $|G:N_1|$  be a  $\sigma_i$ -number and  $I_0 = \{1,\ldots,i\} \setminus \{1,i\}$ . Then  $I_0 \neq \emptyset$  since t > 2. We can assume without loss of generality that i = t. If for some  $k \in I_0$ ,  $|G:N_k|$  is not a  $\sigma_t$ -number, then  $G = N_1N_k$  and  $|G:N_1 \cap N_k| = |G:N_1||G:N_k|$  is a  $\sigma'_k$ -number, so  $H_k \leq N_1 \cap N_k$  since  $N_1 \cap N_k \leq N_G(H_k)$ . Therefore

$$H_k^G = H_k^{N_k N_1} = H_k^{N_1} \le N_1,$$

so G is not simple since  $N_1 \neq G$ , a contradiction. Therefore  $|G: N_k|$  is a  $\sigma_t$ -number for all  $k = 1, \ldots, t-1$ .

Let  $|G: N_t|$  be a  $\sigma_k$ -number and  $j \in \{1, \ldots, t\} \setminus \{k, t\}$ . Then  $G = N_j N_t$  and  $H_j \leq N_j \cap N_t$ , Hence  $H_i^G = H_i^{N_j N_t} = H_i^{N_t} \leq N_t$ , so G is not simple.

Now let R be any non-identity normal subgroup of G. Then  $\{H_1R/R, \ldots, H_tR/R\}$  is a complete Hall  $\sigma$ -set of G/R. On the other hand, since  $N_iR/R \leq N_{G/R}(H_iR/R)$  and  $|G:N_i|$  is  $\sigma$ -primary,  $N_{G/R}(H_iR/R)$  is  $\sigma$ -primary. Therefore the hypothesis holds on G/R for any minimal normal subgroup R of G. Hence the choice of G implies that G/R is generalized  $\sigma$ -soluble, and consequently R is not generalized  $\sigma$ -soluble.

It is clear that  $\{R_1, \ldots, R_t\}$  is a complete Hall  $\sigma$ -set of R, where  $R_i = H_i \cap R$  for all  $i = 1, \ldots, t$ . Moreover, we have also  $N_i \leq N_G(R_i)$  and so from  $|N_iR:N_i| = |R:R\cap N_i|$  we get that  $|R:N_R(R_i)|$  is  $\sigma$ -primary. Therefore the hypothesis holds for R and R < G since G is not simple. Therefore R is generalized  $\sigma$ -soluble by the choice of G. This contradiction completes the proof of the result.

## References

- [1] A. N. Skiba, On  $\sigma$ -subnormal and  $\sigma$ -permutable subgroups of finite groups, J. Algebra, **436** (2015), 1–16.
- [2] A. N. Skiba, A generalization of a Hall theorem, J. Algebra and its Application, 15(4) (2015), 21–36.
- [3] A. N. Skiba, On some results in the theory of finite partially soluble groups, *Commun. Math. Stat.*, 4(3) (2016), 281–309.
- [4] X. Yin, N. Yang, Finite groups with permutable Hall subgroups, Front. Math. China, (2017). https://doi.org/10.1007/s11464-017-0641-4.
- [5] B. Huppert, Endliche Gruppen I, Springer-Verlag, Berlin-Heidelberg-New York, 1967.
- [6] I.M. Isaacs, Semipermutable  $\pi$ -subgroups, Arch. Math., 102 (2014), 1–6.
- [7] M.T. Borovikov, Groups with permutable subgroups of mutually simple orders, *Questions of Algebra.* **5** (1990), 80–82.
- [8] K. Doerk, T. Hawkes, Finite Soluble Groups, Walter de Gruyter, Berlin, New York, 1992.
- [9] A. Ballester-Bolinches, R. Esteban-Romero, M. Asaad, *Products of Finite Groups*, Walter de Gruyter, Berlin-New York, 2010.
- [10] L.A. Shemetkov, A.N. Skiba, Formations of Algebraic Systems, Nauka, Main Editorial Board for Physical and Mathematical Literature, Moscow, 1989.
- [11] J. Zhang, Sylow numbers of finite groups, J. Algebra, 176 (1995), 111-123.
- [12] W. Guo, Finite groups with given indices of normalizers of Sylow subgroups, Siberian Math. J., 37 (1996), 207-214.
- [13] A.I. Mal'cev, *Algebraic Systems*, Nauka, Main Editorial Board for Physical and Mathematical Literature, Moscow, 1970.
- [14] A. Ballester-Bolinches, L.M. Ezquerro, Classes of Finite Groups, Springer-Verlag, Dordrecht, 2006.