

# DEPTH AND STANLEY DEPTH OF THE EDGE IDEALS OF THE POWERS OF PATHS AND CYCLES

ZAHID IQBAL AND MUHAMMAD ISHAQ

**ABSTRACT.** Let  $k$  be a positive integer. We compute depth and Stanley depth of the quotient ring of the edge ideal associated to the  $k^{th}$  power of a path on  $n$  vertices. We show that both depth and Stanley depth have the same values and can be given in terms of  $k$  and  $n$ . For  $n \geq 2k + 1$ , let  $n \equiv 0, k + 1, k + 2, \dots, n - 1 \pmod{2k + 1}$ . Then we give values of depth and Stanley depth of the quotient ring of the edge ideal associated to the  $k^{th}$  power of a cycle on  $n$  vertices and tight bounds otherwise, in terms of  $n$  and  $k$ . We also compute lower bounds for the Stanley depth of the edge ideals associated to the  $k^{th}$  power of a path and a cycle and prove a conjecture of Herzog presented in [7] for these ideals.

**Keywords:** Monomial ideal, edge ideal, depth, Stanley decomposition, Stanley depth,  $k^{th}$  power of a path,  $k^{th}$  power of a cycle.

**2000 Mathematics Subject Classification:** Primary: 13C15; Secondary: 13F20; 05C38; 05E99.

## 1. INTRODUCTION

Let  $K$  be a field and  $S := K[x_1, \dots, x_n]$  the polynomial ring over  $K$ . Let  $M$  be a finitely generated  $\mathbb{Z}^n$ -graded  $S$ -module. A Stanley decomposition of  $M$  is a presentation of  $K$ -vector space  $M$  as a finite direct sum

$$\mathcal{D} : M = \bigoplus_{i=1}^s v_i K[W_i],$$

where  $v_i \in M$ ,  $W_i \in \{x_1, \dots, x_n\}$  such that  $v_i K[W_i]$  denotes the  $K$ -subspace of  $M$ , which is generated by all elements  $v_i w$ , where  $w$  is a monomial in  $K[W_i]$ . The  $\mathbb{Z}^n$ -graded  $K$ -subspace  $v_i K[W_i] \subset M$  is called a Stanley space of dimension  $|W_i|$ , if  $v_i K[W_i]$  is a free  $K[W_i]$ -module, where  $|W_i|$  denotes the cardinality of  $W_i$ . Define

$$\text{sdepth}(\mathcal{D}) = \min\{|W_i| : i = 1, \dots, s\},$$

and

$$\text{sdepth}(M) = \max\{\text{sdepth}(\mathcal{D}) : \mathcal{D} \text{ is a Stanley decomposition of } M\}.$$

The number  $\text{sdepth}(\mathcal{D})$  is called the Stanley depth of decomposition  $\mathcal{D}$  and  $\text{sdepth}(M)$  is called the Stanley depth of  $M$ . Stanley conjectured in [22] that  $\text{sdepth}(M) \geq \text{depth}(M)$  for any  $\mathbb{Z}^n$ -graded  $S$ -module  $M$ . This conjecture was disproved by Duval et al. in [6] as was expected due to different natures of both the invariants. However, the relation between Stanley depth and some other invariants has already been established see [9, 10, 19, 24]. In [9], Herzog, Vladioiu and Zheng proved that the Stanley depth of  $M$  can be computed in a finite number of steps, if  $M = J/I$ , where  $I \subset J \subset S$  are monomial ideals. But practically it is too hard to compute Stanley depth by using this method see for instance, [1, 3, 13, 14]. For computing Stanley depth for some classes of modules we refer the reader to [12, 18, 20, 21]. In this paper we attempt to find values and reasonable bounds for depth and Stanley depth of  $I$  and  $S/I$ , where  $I$  is the edge ideal of a power of a path or a cycle. We also compare the values of  $\text{sdepth}(I)$  and  $\text{sdepth}(S/I)$  and give positive answers to the following conjecture of Herzog.

**Conjecture 1.1.** [7] *Let  $I \subset S$  be a monomial ideal then*

$$\text{sdepth}(I) \geq \text{sdepth}(S/I).$$

The above conjecture has been proved in some special cases see [11, 14, 18, 21]. The paper is organized as follows: First two sections are devoted to introduction, definitions, notations, and discussion of some known results. In third section, we compute depth and Stanley depth of  $S/I(P_n^k)$  as Theorems (3.7 and 3.11), where  $I(P_n^k)$  denotes the edge ideal of the  $k^{th}$  power of a path  $P_n$  on  $n$  vertices. These theorems prove that

$$\text{depth}(S/I(P_n^k)) = \text{sdepth}(S/I(P_n^k)) = \lceil \frac{n}{2k+1} \rceil.$$

Let  $I(C_n^k)$  be the edge ideal of the  $k^{th}$  power of a cycle  $C_n$  on  $n$  vertices. In fourth section we give some lower bounds for depth and Stanley depth of  $S/I(C_n^k)$  as Theorems (4.4 and 4.6). For  $n \geq 2k+1$ , our Corollaries (4.5 and 4.7) show that

$$\begin{aligned} \text{depth}(S/I(C_n^k)) &= \text{sdepth}(S/I(C_n^k)) = \lceil \frac{n}{2k+1} \rceil, \text{ if } n \equiv 0, k+1, \dots, n-1 \pmod{2k+1}; \\ \lceil \frac{n}{2k+1} \rceil - 1 &\leq \text{depth}(S/I(C_n^k)), \text{sdepth}(S/I(C_n^k)) \leq \lceil \frac{n}{2k+1} \rceil, \text{ if } n \equiv 1, 2, \dots, k \pmod{2k+1}. \end{aligned}$$

Last section is devoted to Conjecture 1.1 for  $I(P_n^k)$  and  $I(C_n^k)$ . By our Theorem 5.2 we have

$$\text{sdepth}(I(P_n^k)) \geq \lceil \frac{n}{2k+1} \rceil + 1,$$

which shows that  $I(P_n^k)$  satisfies Conjecture 1.1. Let  $n \geq 2k+1$ , Proposition 5.3 gives a lower bound for  $I(C_n^k)/I(P_n^k)$  that is

$$\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq \lceil \frac{n+k+1}{2k+1} \rceil.$$

Corollary 5.5 of this paper proves that  $I(C_n^k)$  satisfies Conjecture 1.1. The above results are proved in [11] when  $k=2$ .

## 2. DEFINITIONS AND NOTATION

Throughout this paper,  $\mathfrak{m}$  denotes the unique maximal graded ideal  $(x_1, x_2, \dots, x_n)$  of  $S$ . We set  $S_m := K[x_1, x_2, \dots, x_m]$ ,  $\text{supp}(v) := \{i : x_i | v\}$  and  $\text{supp}(I) := \{i : x_i | u, \text{ for some } u \in \mathcal{G}(I)\}$ , where  $\mathcal{G}(I)$  denotes the unique minimal set of monomial generators of the monomial ideal  $I$ . Let  $I \subset S$  be an ideal then we write  $I$  instead of  $IS$ . Thus every ideal will be considered an ideal of  $S$  unless otherwise stated. Let  $I$  and  $J$  be monomial ideals of  $S$ , then for  $I+J$  we write  $(I, J)$ .

**Definition 2.1.** Let  $G$  be a simple graph, for a positive integer  $k$  the  $k^{th}$  power of graph  $G$  is another graph  $G^k$  on the same set of vertices, such that two vertices are adjacent in  $G^k$  when their distance in  $G$  is at most  $k$ .

In the whole paper we label the vertices of the graph  $G$  by  $1, 2, \dots, n$ . We denote the set of vertices of  $G$  by  $[n] := \{1, 2, \dots, n\}$ . We assume that all graphs and their powers are simple graphs. We also assume that all graphs have at least two vertices and a non-empty edge set.

**Definition 2.2.** Let  $G$  be a graph then the edge ideal  $I(G)$  associated to  $G$  is the square free monomial ideal of  $S$  that is

$$I(G) = (x_i x_j : \{i, j\} \in E(G)).$$

**Definition 2.3.** Let  $G$  be a graph, then  $G$  is called a path if  $E(G) = \{\{i, i+1\} : i \in [n-1]\}$ . A path on  $n$  vertices is usually denoted by  $P_n$ .

**Definition 2.4.** Let  $n \geq 3$ , a graph  $G$  is called a cycle if  $E(G) = \{\{i, i+1\} : i \in [n-1]\} \cup \{1, n\}$ . A cycle on  $n$  vertices is usually denoted by  $C_n$ .

**Definition 2.5.** For  $n \geq 2$ , the  $k^{th}$  power of a path, denoted by  $P_n^k$ , is a graph such that for all  $1 \leq i < j \leq n$ ,  $\{i, j\} \in E(P_n^k)$  if and only if  $0 < j-i \leq k$ . That is the edge set of  $P_n^k$  is:

$$(1): \text{ If } n \leq k+1, \text{ then } E(P_n^k) = \{\{i, j\} : 1 \leq i < j \leq n\}.$$

(2): If  $n \geq k + 2$ , then

$$E(P_n^k) = \bigcup_{i=1}^{n-k} \{\{i, i+1\}, \{i, i+2\}, \dots, \{i, i+k\}\} \bigcup \bigcup_{j=n-k+1}^{n-1} \{\{j, j+1\}, \{j, j+2\}, \dots, \{j, n\}\}.$$

**Definition 2.6.** For  $n \geq 3$ , the  $k^{th}$  power of a cycle, denoted by  $C_n^k$ , is a graph such that for all vertices  $1 \leq i, j \leq n$ ,  $\{i, j\} \in E(C_n^k)$  if and only if  $|j - i| \leq k$  or  $|j - i| \geq n - k$ . The edge set of  $C_n^k$  is:

(1): If  $n \leq 2k + 1$ , then  $E(C_n^k) = \{\{i, j\} : 1 \leq i < j \leq n\}$ .

(2): If  $n \geq 2k + 2$ , then

$$E(C_n^k) = E(P_n^k) \bigcup \bigcup_{l=1}^k \{\{l, l+n-k\}, \{l, l+n-k+1\}, \{l, l+n-k+2\} \dots, \{l, n\}\}.$$

The minimal generating set for the edge ideal of  $P_n^k$  is:

(1): If  $n \leq k + 1$ , then  $\mathcal{G}(I(P_n^k)) = \{x_i x_j : 1 \leq i < j \leq n\}$ . That is  $I(P_n^k)$  is a square free Veronese ideal of degree 2 in variables  $x_1, x_2, \dots, x_n$ .

(2): If  $n \geq k + 2$ , then

$$\mathcal{G}(I(P_n^k)) = \bigcup_{i=1}^{n-k} \{x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{i+k}\} \bigcup \bigcup_{j=n-k+1}^{n-1} \{x_j x_{j+1}, x_j x_{j+2}, \dots, x_j x_n\}.$$

The minimal generating set for the edge ideal of  $C_n^k$  is:

(1): If  $n \leq 2k + 1$ , then  $\mathcal{G}(I(C_n^k)) = \{x_i x_j : 1 \leq i < j \leq n\}$ . That is  $I(C_n^k)$  is a square free Veronese ideal of degree 2 in variables  $x_1, x_2, \dots, x_n$ .

(2): If  $n \geq 2k + 2$ , then

$$\mathcal{G}(I(C_n^k)) = \mathcal{G}(I(P_n^k)) \bigcup \bigcup_{l=1}^k \{x_l x_{l+n-k}, x_l x_{l+n-k+1}, \dots, x_l x_n\}.$$

**Lemma 2.7.** [15, Lemma 3] For  $n \geq k + 1$ ,  $|\mathcal{G}(I(P_n^k))| = nk - \frac{k(k+1)}{2}$ .

**Remark 2.8.** If  $n \geq 2k + 1$ , then  $|\mathcal{G}(I(C_n^k))| = nk$ .

**Definition 2.9.** Let  $G$  be a graph and let  $i \in [n]$ , then we define

$$N_G(x_i) := \{x_j : x_i x_j \in \mathcal{G}(I(G))\},$$

where  $j \in [n] \setminus \{i\}$ .

Let  $k \geq 2$ ,  $0 \leq i \leq k - 1$  and  $n \geq 2k + 2$ , then in the following we introduce some particular monomial prime ideals of  $S$ .

$$A_{n-k+i} := (x_{n-k}, x_{n-k+1}, \dots, x_{n-k+i}).$$

(We set  $A_{n-k-1} = (0)$ ).

$$B_{n-k+i} := (N_{P_n^k}(x_{n-k+i})) = (x_{n-2k+i}, x_{n-2k+i+1}, \dots, x_{n-k+i-1}, x_{n-k+i+1}, \dots, x_n).$$

and

$$\begin{aligned} D_{n-k+i} &:= (N_{C_n^k}(x_{n-k+i})) \\ &= \begin{cases} (x_{n-2k}, x_{n-2k+1}, \dots, x_{n-k-1}, x_{n-k+1}, \dots, x_n), & \text{if } i = 0; \\ (x_{n-2k+i}, x_{n-2k+i+1}, \dots, x_{n-k+i-1}, x_{n-k+i+1}, \dots, x_n, x_1, x_2, \dots, x_i), & \text{if } 1 \leq i \leq k - 1. \end{cases} \end{aligned}$$

These monomial prime ideals and the following function play important role in the proof of our main theorems. Let  $k \geq 2$ ,  $0 \leq i \leq k - 1$  and  $2k + 2 \leq n \leq 3k + 1$ , we define a function

$$f : \{n - k, n - k + 1, \dots, n - k + i, \dots, n - 1\} \longrightarrow \mathbb{Z}^+ \cup \{0\}, \text{ by}$$

$$f(n-k+i) = \begin{cases} k, & \text{if } n-2k-1+i \geq k+1; \\ n-2k-2+i, & \text{if } 2 \leq n-2k-1+i < k+1. \end{cases}$$

In the following we recall some known results that we refer several times in this paper.

**Lemma 2.10.** (*Depth Lemma*) *If  $0 \rightarrow U \rightarrow M \rightarrow N \rightarrow 0$  is a short exact sequence of modules over a local ring  $S$ , or a Noetherian graded ring with  $S_0$  local, then*

- (1)  $\text{depth } M \geq \min\{\text{depth } N, \text{depth } U\}.$
- (2)  $\text{depth } U \geq \min\{\text{depth } M, \text{depth } N + 1\}.$
- (3)  $\text{depth } N \geq \min\{\text{depth } U - 1, \text{depth } M\}.$

**Lemma 2.11.** [21] *Let  $0 \rightarrow U \rightarrow V \rightarrow W \rightarrow 0$  be a short exact sequence of  $\mathbb{Z}^n$ -graded  $S$ -modules. Then*

$$\text{sdepth}(V) \geq \min\{\text{sdepth}(U), \text{sdepth}(W)\}.$$

**Lemma 2.12.** [11] *Let  $I \subset S$  be a squarefree monomial ideal with  $\text{supp}(I) = [n]$ , let  $v := x_{i_1}x_{i_2}\cdots x_{i_q} \in S/I$ , such that  $x_jv \in I$ , for all  $j \in [n] \setminus \text{supp}(v)$ . Then  $\text{sdepth}(S/I) \leq q$ .*

### 3. DEPTH AND STANLEY OF CYCLIC MODULES ASSOCIATED TO THE EDGE IDEALS OF THE POWERS OF A PATH

In this section, we compute depth and Stanley depth of  $S/I(P_n^k)$ . We start this section with the following elementary lemma:

**Lemma 3.1.** *Let  $a \geq 2$  be an integer,  $\{E_i : 1 \leq i \leq a\}$  and  $\{G_i : 0 \leq i \leq a\}$  be sequences of  $\mathbb{Z}^n$ -graded  $S$  modules, and we have the following short exact sequences:*

$$0 \rightarrow E_1 \rightarrow G_0 \rightarrow G_1 \rightarrow 0 \tag{1}$$

$$0 \rightarrow E_2 \rightarrow G_1 \rightarrow G_2 \rightarrow 0 \tag{2}$$

$$\vdots$$

$$0 \rightarrow E_{a-1} \rightarrow G_{a-2} \rightarrow G_{a-1} \rightarrow 0 \tag{a-1}$$

$$0 \rightarrow E_a \rightarrow G_{a-1} \rightarrow G_a \rightarrow 0 \tag{a}$$

*such that  $\text{depth}(G_a) \geq \text{depth}(E_a)$  and  $\text{depth}(E_i) \geq \text{depth}(E_{i-1})$ , for all  $2 \leq i \leq a$ , then  $\text{depth}(G_0) = \text{depth}(E_1)$ .*

*Proof.* By assumption  $\text{depth}(G_a) \geq \text{depth}(E_a)$ , applying Depth Lemma on the exact sequence (a) we get  $\text{depth}(G_{a-1}) = \text{depth}(E_a)$ . We also have by assumption

$$\text{depth}(G_{a-1}) = \text{depth}(E_a) \geq \text{depth}(E_{a-1}),$$

applying Depth Lemma to the exact sequence (a-1) we have  $\text{depth}(G_{a-2}) = \text{depth}(E_{a-1})$ . We repeat the same steps on all exact sequences one by one from bottom to top and we get  $\text{depth}(G_{i-1}) = \text{depth}(E_i)$  for all  $i$ . Thus if  $i = 1$  then we have  $\text{depth}(G_0) = \text{depth}(E_1)$ .  $\square$

**Lemma 3.2.** *Let  $k \geq 2$  and  $n \geq 2k + 2$ , then  $S/(I(P_n^k), A_{n-1}) \cong S_{n-k-1}/I(P_{n-k-1}^k)[x_n]$ .*

*Proof.* We have

$$\mathcal{G}(I(P_n^k)) = \bigcup_{i=1}^{n-k} \{x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{i+k}\} \bigcup \bigcup_{i=n-k+1}^{n-1} \{x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_n\}.$$

Then

$$\begin{aligned}
I(P_n^k) + A_{n-1} &= \left[ \sum_{i=1}^{n-2k-1} (x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{i+k}) + \sum_{i=n-2k}^{n-k} (x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{i+k}) + \right. \\
&\quad \left. \sum_{i=n-k+1}^{n-1} (x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_n) \right] + A_{n-1} \\
&= \sum_{i=1}^{n-2k-1} (x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{i+k}) + \\
&\quad \sum_{i=n-2k}^{n-k-2} (x_i x_{i+1}, x_i x_{i+2}, \dots, x_i x_{n-k-1}) + A_{n-1} = I(P_{n-k-1}^k) + A_{n-1}.
\end{aligned}$$

Thus the required result follows.  $\square$

**Lemma 3.3.** *Let  $k \geq 2$ ,  $0 \leq i \leq k-1$  and  $n \geq 3k+2$ , then*

$$S/(I(P_n^k) : x_{n-k+i}) \cong S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)[x_{n-k+i}].$$

*Proof.* It is enough to prove that  $(I(P_n^k) : x_{n-k+i}) = (I(P_{n-2k-1+i}^k), B_{n-k+i})$ . Clearly

$$I(P_{n-2k-1+i}^k) \subset I(P_n^k) \subset (I(P_n^k) : x_{n-k+i}).$$

Let  $u \in B_{n-k+i}$ , then by definition of  $I(P_n^k)$ ,  $u x_{n-k+i} \in I(P_n^k)$  that is  $u \in (I(P_n^k) : x_{n-k+i})$ . Thus  $B_{n-k+i} \subset (I(P_n^k) : x_{n-k+i})$  and we have  $(I(P_{n-2k-1+i}^k), B_{n-k+i}) \subset (I(P_n^k) : x_{n-k+i})$ . Now for the other inclusion, let  $w$  be a monomial generator of  $(I(P_n^k) : x_{n-k+i})$ , then  $w = \frac{v}{\gcd(v, x_{n-k+i})}$ , where  $v \in \mathcal{G}(I(P_n^k))$ . If  $\text{supp}(v) \cap \mathcal{G}(B_{n-k+i}) \neq \emptyset$ , then we have  $w \in \mathcal{G}(B_{n-k+i})$  and if  $\text{supp}(v) \cap \mathcal{G}(B_{n-k+i}) = \emptyset$  then

$$w \in \mathcal{G}(I(P_n^k)) \cap K[x_1, x_2, \dots, x_{n-2k-1+i}] = \mathcal{G}(I(P_{n-2k-1+i}^k)).$$

$\square$

**Lemma 3.4.** *Let  $n \geq 3k+2$  and  $0 \leq i \leq k-1$ , then we have*

$$S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \cong S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)[x_{n-k+i}].$$

*Proof.* Clearly  $((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) = ((I(P_n^k) : x_{n-k+i}), A_{n-k+(i-1)})$ . Now using the proof of Lemma 3.3

$((I(P_n^k) : x_{n-k+i}), A_{n-k+(i-1)}) = (I(P_{n-2k-1+i}^k), B_{n-k+i}, A_{n-k+(i-1)}) = (I(P_{n-2k-1+i}^k), B_{n-k+i})$  as  $A_{n-k+(i-1)} \subset B_{n-k+i}$ . Thus the required result follows by Lemma 3.3.  $\square$

**Remark 3.5.** Let  $m \geq 2$  and  $I(P_m^{m-1}) \subset S_m = K[x_1, x_2, \dots, x_m]$  be the edge ideal of the  $(m-1)^{th}$  power of path  $P_m$ , then  $I(P_m^{m-1})$  is a squarefree Veronese ideal of degree 2 in variables  $x_1, x_2, \dots, x_m$ . Thus by [8, Corollary 10.3.7] and [3, Theorem 1.1]

$$\text{depth}(S_m/I(P_m^{m-1})) = \text{sdepth}(S_m/I(P_m^{m-1})) = 1.$$

**Remark 3.6.** Let  $k \geq 2$  and  $2k+2 \leq n \leq 3k+1$ , then it is easy to see that

- (1): If  $n = 2k+2$ , then  $S/(I(P_n^k) : x_{n-k}) = S/(x_2, \dots, x_{n-k-1}, x_{n-k+1}, \dots, x_n) \cong K[x_1, x_{n-k}]$ .
- (2): If  $0 \leq i \leq k-1$  and  $n > 2k+2$ , then

$$\begin{aligned}
S/(I(P_n^k) : x_{n-k+i}) &= S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \\
&\cong S_{n-2k-1+i}/I(P_{n-2k-1+i}^{f(n-k+i)}) = \begin{cases} S_{n-2k-1+i}/I(P_{n-2k-1+i}^k), & \text{if } n-2k-1+i \geq k+1; \\ S_{n-2k-1+i}/I(P_{n-2k-1+i}^{n-2k-2+i}), & \text{otherwise.} \end{cases}
\end{aligned}$$

**Theorem 3.7.** *Let  $n \geq 2$ , then  $\text{depth}(S/I(P_n^k)) = \lceil \frac{n}{2k+1} \rceil$ .*

*Proof.* (a): Let  $n \leq k+1$ , then  $I(P_n^k)$  is a square free Veronese ideal thus by Remark 3.5,  $\text{depth}(S/I(P_n^k)) = 1 = \lceil \frac{n}{2k+1} \rceil$ .

(b): Let  $n \geq k+2$ , we consider the following cases:

(1): If  $k=1$ , then by [16, Lemma 2.8] we have  $\text{depth}(S/I(P_n^1)) = \lceil \frac{n}{3} \rceil = \lceil \frac{n}{2k+1} \rceil$ .

(2): If  $k \geq 2, k+2 \leq n \leq 2k+1$ , then we have  $\text{depth}(S/I(P_n^k)) \geq 1$  as  $\mathfrak{m} \notin \text{Ass}(S/I(P_n^k))$ . Since  $x_{k+1} \notin I(P_n^k)$  and  $x_s x_{k+1} \in \mathcal{G}(I(P_n^k))$  for all  $s \in \{1, \dots, k, k+2, \dots, n\}$ , therefore

$$(I(P_n^k) : x_{k+1}) = (x_1, \dots, x_k, x_{k+2}, \dots, x_n).$$

By [21, Corollary 1.3], we have

$$\text{depth}(S/I(P_n^k)) \leq \text{depth}(S/(I(P_n^k) : x_{k+1})) = \text{depth}(S/(x_1, \dots, x_k, x_{k+2}, \dots, x_n)) = 1.$$

Thus  $\text{depth}(S/I(P_n^k)) = 1 = \lceil \frac{n}{2k+1} \rceil$ .

(3): If  $k \geq 2, 2k+2 \leq n \leq 3k+1$ , now consider the following short exact sequences:

$$\begin{aligned} 0 &\longrightarrow S/(I(P_n^k) : x_{n-k}) \xrightarrow{\cdot x_{n-k}} S/I(P_n^k) \longrightarrow S/(I(P_n^k), x_{n-k}) \longrightarrow 0 \\ 0 &\longrightarrow S/((I(P_n^k), A_{n-k}) : x_{n-k+1}) \xrightarrow{\cdot x_{n-k+1}} S/(I(P_n^k), A_{n-k}) \longrightarrow S/(I(P_n^k), A_{n-k+1}) \longrightarrow 0 \\ &\quad \vdots \\ 0 &\longrightarrow S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \xrightarrow{\cdot x_{n-k+i}} S/(I(P_n^k), A_{n-k+(i-1)}) \longrightarrow S/(I(P_n^k), A_{n-k+i}) \longrightarrow 0 \\ &\quad \vdots \\ 0 &\longrightarrow S/((I(P_n^k), A_{n-2}) : x_{n-1}) \xrightarrow{\cdot x_{n-1}} S/(I(P_n^k), A_{n-2}) \longrightarrow S/(I(P_n^k), A_{n-1}) \longrightarrow 0. \end{aligned}$$

By Lemma 3.2,  $S/(I(P_n^k), A_{n-1}) \cong S_{n-k-1}/I(P_{n-k-1}^k)[x_n]$ . Since  $k+1 \leq n-k-1 \leq 2k$ , thus if  $n-k-1 = k+1$  then by case(a) and [9, Lemma 3.6], otherwise by case(b)(2) and [9, Lemma 3.6] we have

$$\text{depth}(S/(I(P_n^k), A_{n-1})) = 2.$$

Now we show that  $\text{depth}(S/(I(P_n^k) : x_{n-k})) = 2$ , for this we consider two cases:

If  $n = 2k+2$ , then by Remark 3.6

$$S/(I(P_n^k) : x_{n-k}) = S/(x_2, x_3, \dots, x_{n-k-1}, x_{n-k+1}, \dots, x_n) \cong K[x_1, x_{n-k}],$$

and thus  $\text{depth}(S/(I(P_n^k) : x_{n-k})) = 2$ .

If  $n > 2k+2$ , by Remark 3.6 we have

$$S/(I(P_n^k) : x_{n-k}) \cong S_{n-2k-1}/I(P_{n-2k-1}^{n-2k-2})[x_{n-k}],$$

where  $2 \leq n-2k-1 \leq k$ . Thus by Remark 3.5 and [9, Lemma 3.6] we get

$$\text{depth}(S/(I(P_n^k) : x_{n-k})) = 2.$$

Now let  $1 \leq i \leq k-1$ , by Remark 3.6

$$S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) = S/(I(P_n^k) : x_{n-k+i}) \cong S_{n-2k-1+i}/I(P_{n-2k-1+i}^{f(n-k+i)})[x_{n-k+i}].$$

Let  $T := S_{n-2k-1+i}/I(P_{n-2k-1+i}^{f(n-k+i)})[x_{n-k+i}]$ , we consider the following cases:

(i): If  $k+1 = n-2k-1+i$ , then  $T = S_{k+1}/I(P_{k+1}^k)[x_{n-k+i}]$ , thus by case(a) and [9, Lemma 3.6] we have  $\text{depth}(T) = 2$ .

(ii): If  $k+1 < n-2k-1+i$ , then  $T = S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)[x_{n-k+i}]$ . Since  $k+2 \leq n-2k-1+i \leq 2k-1$ , thus by case(b)(2) and [9, Lemma 3.6] we have  $\text{depth}(T) = 2$ .

(iii): If  $2 \leq n - 2k - 1 + i < k + 1$ , then  $T = S_{n-2k-1+i}/I(P_{n-2k-1+i}^{n-2k-2+i})[x_{n-k+i}]$ ,  
by Remark 3.5 and [9, Lemma 3.6] we have  $\text{depth}(T) = 2$ .

Thus by Lemma 3.1 we have  $\text{depth}(S/I(P_n^k)) = 2$ .

(4): If  $k \geq 2$ ,  $n \geq 3k + 2$ , then we consider the family of short exact sequences:

$$\begin{aligned} 0 \longrightarrow S/(I(P_n^k) : x_{n-k}) &= S/((I(P_n^k), A_{n-k-1}) : x_{n-k}) \xrightarrow{\cdot x_{n-k}} S/I(P_n^k) \longrightarrow S/(I(P_n^k), A_{n-k}) \longrightarrow 0 \\ 0 \longrightarrow S/((I(P_n^k), A_{n-k}) : x_{n-k+1}) &\xrightarrow{\cdot x_{n-k+1}} S/(I(P_n^k), A_{n-k}) \longrightarrow S/(I(P_n^k), A_{n-k+1}) \longrightarrow 0 \\ &\vdots \\ 0 \longrightarrow S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) &\xrightarrow{\cdot x_{n-k+i}} S/(I(P_n^k), A_{n-k+(i-1)}) \longrightarrow S/(I(P_n^k), A_{n-k+i}) \longrightarrow 0 \\ &\vdots \\ 0 \longrightarrow S/((I(P_n^k), A_{n-2}) : x_{n-1}) &\xrightarrow{\cdot x_{n-1}} S/(I(P_n^k), A_{n-2}) \longrightarrow S/(I(P_n^k), A_{n-1}) \longrightarrow 0. \end{aligned}$$

By Lemma 3.2,  $S/(I(P_n^k), A_{n-1}) \cong S_{n-k-1}/I(P_{n-k-1}^k)[x_n]$ . Thus by induction on  $n$  and [9, Lemma 3.6] we have

$$\text{depth}(S/(I(P_n^k), A_{n-1})) = \lceil \frac{n-k-1}{2k+1} \rceil + 1.$$

Let  $0 \leq i \leq k-1$ , by Lemma 3.4 we have

$$S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \cong S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)[x_{n-k+i}].$$

Thus by induction on  $n$  and [9, Lemma 3.6] we have

$$\text{depth}(S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i})) = \lceil \frac{n-2k-1+i}{2k+1} \rceil + 1.$$

We can see that

$$\text{depth}(S/(I(P_n^k), A_{n-1})) = \lceil \frac{n-k-1}{2k+1} \rceil + 1 \geq \lceil \frac{n-k-2}{2k+1} \rceil + 1 = \text{depth}(S/(I(P_n^k), A_{n-2}) : x_{n-1}).$$

And for all  $1 \leq i \leq k-1$ ,

$$\begin{aligned} \text{depth}(S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i})) &= \lceil \frac{n-2k-1+i}{2k+1} \rceil + 1 \geq \\ &\lceil \frac{n-2k-2+i}{2k+1} \rceil + 1 = \text{depth}(S/((I(P_n^k), A_{n-k+(i-2)}) : x_{n-k+(i-1)})). \end{aligned}$$

Thus by Lemma 3.1 we have

$$\text{depth}(S/I(P_n^k)) = \lceil \frac{n-2k-1}{2k+1} \rceil + 1 = \lceil \frac{n}{2k+1} \rceil.$$

□

**Lemma 3.8.** [23, Lemma 4] Let  $n \geq 2$ , then  $\text{sdepth}(S/I(P_n^1)) = \lceil \frac{n}{3} \rceil$ .

**Example 3.9.** Let  $n \geq 2$ , and  $n \leq 2k + 1$ , then  $\text{sdepth}(S/I(P_n^k)) = 1$ .

*Proof.* If  $n \leq k + 1$ , then by [3, Theorem 1.1]  $\text{sdepth}(S/I(P_n^k)) = 1$ . Now let  $k + 2 \leq n \leq 2k + 1$ , then  $\text{depth}(S/I(P_n^k)) \geq 1$  as  $\mathfrak{m} \notin \text{Ass}(S/I(P_n^k))$ , thus by [5, Theorem 1.4]  $\text{sdepth}(S/I(P_n^k)) \geq 1$ . Since  $x_{k+1} \notin I(P_n^k)$  and  $x_i x_{k+1} \in \mathcal{G}(I(P_n^k))$  for all  $i \in \{1, \dots, k, k+2, \dots, n\}$ , therefore

$$(I(P_n^k) : x_{k+1}) = (x_1, \dots, x_k, x_{k+2}, \dots, x_n).$$

Thus by [2, Proposition 2.7]

$$\text{sdepth}(S/I(P_n^k)) \leq \text{sdepth}(S/(I(P_n^k) : x_{k+1})) = \text{sdepth}(S/(x_1, \dots, x_k, x_{k+2}, \dots, x_n)) = 1.$$

□

**Proposition 3.10.** *Let  $k \geq 2$  and  $n \geq 2k + 2$ , then  $\text{sdepth}(S/I(P_n^k)) \geq \lceil \frac{n}{2k+1} \rceil$ .*

*Proof.* (1): If  $2k + 2 \leq n \leq 3k + 1$ , then we apply Lemma 2.11 on the exact sequences in case(b)(3) of Theorem 3.7 and we get  $\text{sdepth}(S/I(P_n^k)) \geq 2 = \lceil \frac{n}{2k+1} \rceil$ .

(2): If  $n \geq 3k + 2$ , then the proof is similar to Theorem 3.7. We apply Lemma 2.11 on the exact sequences in case(b)(4) of Theorem 3.7 we get

$$\text{sdepth}(S/I(P_n^k)) \geq \min \{ \text{sdepth}(S/(I(P_n^k) : x_{n-k})), \text{sdepth}(S/(I(P_n^k), A_{n-1})), \\ \min_{i=1}^{k-1} \{ \text{sdepth}(S/((I(P_n^k), A_{n-k+(i-1)}) : x_{n-k+i})) \} \} \geq \lceil \frac{n}{2k+1} \rceil.$$

□

**Theorem 3.11.** *Let  $n \geq 2$ , then  $\text{sdepth}(S/I(P_n^k)) = \lceil \frac{n}{2k+1} \rceil$ .*

*Proof.* By Lemma 3.8, Example 3.9 and Proposition 3.10, it is enough to prove, if  $k \geq 2$  then

$$\text{sdepth}(S/I(P_n^k)) \leq \lceil \frac{n}{2k+1} \rceil, \text{ for } n \geq 2k + 2.$$

We consider the following three cases:

(1): Let  $n = (2k + 1)l$ , where  $l \geq 1$ . We see that

$$v = x_{k+1}x_{3k+2}x_{5k+3} \cdots x_{(2k+1)l-k} \in S \setminus I(P_n^k),$$

$$\text{but } x_{t_1}v \in I(P_n^k), \text{ for all } t_1 \in [n] \setminus \text{supp}(v), \text{ by Lemma 2.12, } \text{sdepth}(S/I(P_n^k)) \leq l = \lceil \frac{n}{2k+1} \rceil.$$

(2): Let  $n = (2k + 1)l + r$ , where  $r \in \{1, 2, 3, \dots, k + 1\}$  and  $l \geq 1$ . We have

$$v = x_{k+1}x_{3k+2}x_{5k+3} \cdots x_{(2k+1)l-k}x_{(2k+1)l+r} \in S \setminus I(P_n^k),$$

$$\text{but } x_{t_2}v \in I(P_n^k), \text{ for all } t_2 \in [n] \setminus \text{supp}(v), \text{ by Lemma 2.12,}$$

$$\text{sdepth}(S/I(P_n^k)) \leq l + 1 = \lceil \frac{n}{2k+1} \rceil.$$

(3): Let  $n = (2k + 1)l + s$ , where  $s \in \{k + 2, k + 3, \dots, 2k\}$  and  $l \geq 1$ . Since

$$v = x_{k+1}x_{3k+2}x_{5k+3} \cdots x_{(2k+1)l+k+1} \in S \setminus I(P_n^k),$$

$$\text{but } x_{t_3}v \in I(P_n^k), \text{ for all } t_3 \in [n] \setminus \text{supp}(v), \text{ by Lemma 2.12,}$$

$$\text{sdepth}(S/I(P_n^k)) \leq l + 1 = \lceil \frac{n}{2k+1} \rceil.$$

□

#### 4. DEPT AND STANLEY DEPTH OF CYCLIC MODULES ASSOCIATED TO THE EDGE IDEALS OF THE POWERS OF A CYCLE

In this section, we compute bounds for depth and Stanley depth for cyclic modules associated to the edge ideals of powers of a cycle.

**Lemma 4.1.** *Let  $k \geq 2$  and  $n \geq 3k + 2$ , then  $S/(I(C_n^k), A_{n-1}) \cong S_{n-k}/I(P_{n-k}^k)$ .*

*Proof.* We have

$$\mathcal{G}(I(C_n^k)) = \mathcal{G}(I(P_n^k)) \bigcup \bigcup_{l=1}^{k-1} \{x_l x_{l+n-k}, x_l x_{l+n-k+1}, \dots, x_l x_{n-1}\} \bigcup \{x_1 x_n, x_2 x_n, \dots, x_k x_n\}.$$

Then

$$I(C_n^k) + A_{n-1} = I(P_n^k) + \sum_{l=1}^{k-1} (x_l x_{l+n-k}, x_l x_{l+n-k+1}, \dots, x_l x_{n-1}) + (x_1 x_n, x_2 x_n, \dots, x_k x_n) + A_{n-1}.$$



Thus by the proof of Lemma 3.2, we have

$$I(P_n^k) + A_{n-1} = I(P_{n-k-1}^k) + A_{n-1}.$$

And

$$\sum_{l=1}^{k-1} (x_l x_{l+n-k}, x_l x_{l+n-k+1}, \dots, x_l x_{n-1}) + A_{n-1} = A_{n-1}.$$

Therefore

$$\begin{aligned} S/(I(C_n^k), A_{n-1}) &= S/(I(P_{n-k-1}^k), A_{n-1}, (x_1 x_n, x_2 x_n, \dots, x_k x_n)) \\ &\cong K[x_1, x_2, \dots, x_{n-k-1}, x_n] / (I(P_{n-k-1}^k), (x_1 x_n, x_2 x_n, \dots, x_k x_n)). \end{aligned}$$

After renumbering the variables, we have

$$K[x_1, x_2, \dots, x_{n-k-1}, x_n] / (I(P_{n-k-1}^k), (x_1 x_n, x_2 x_n, \dots, x_k x_n)) \cong S_{n-k} / I(P_{n-k}^k).$$

□

**Lemma 4.2.** *Let  $k \geq 2$  and  $n \geq 3k + 2$  and  $0 \leq i \leq k - 1$ , then*

$$S/(I(C_n^k) : x_{n-k+i}) \cong S_{n-2k-1} / I(P_{n-2k-1}^k)[x_{n-k+i}].$$

*Proof.* Let  $w$  be a monomial generator of  $(I(C_n^k) : x_{n-k+i})$ , then  $w = \frac{v}{\gcd(v, x_{n-k+i})}$ , where  $v \in \mathcal{G}(I(C_n^k))$ . If  $\text{supp}(v) \cap \mathcal{G}(D_{n-k+i}) \neq \emptyset$ , then we have  $w \in \mathcal{G}(D_{n-k+i})$  and if  $\text{supp}(v) \cap \mathcal{G}(D_{n-k+i}) = \emptyset$  then

$$w \in E := \mathcal{G}(I(C_n^k)) \cap K[x_{i+1}, x_{i+2}, \dots, x_{n-2k-1+i}].$$

So we get

$$(I(C_n^k) : x_{n-k+i}) \subset E + D_{n-k+i},$$

the other inclusion being trivial we get that  $(I(C_n^k) : x_{n-k+i}) = E + D_{n-k+i}$ . Thus we have  $S/(I(C_n^k) : x_{n-k+i}) = S/(E + D_{n-k+i})$ . After renumbering the variables, we have

$$S/(I(C_n^k) : x_{n-k+i}) = S/(E, D_{n-k+i}) \cong S_{n-2k-1} / I(P_{n-2k-1}^k)[x_{n-k+i}].$$

□

**Lemma 4.3.** *Let  $k \geq 2$ ,  $n \geq 3k + 2$  and  $0 \leq i \leq k - 1$ , then we have*

$$S/((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \cong S_{n-2k-1} / I(P_{n-2k-1}^k)[x_{n-k+i}].$$

*Proof.* Clearly  $((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) = ((I(C_n^k) : x_{n-k+i}), A_{n-k+(i-1)})$ . By using the same arguments as in the proof of Lemma 4.2 we have

$$((I(C_n^k) : x_{n-k+i}), A_{n-k+(i-1)}) = (E, D_{n-k+i}, A_{n-k+(i-1)}) = (E, D_{n-k+i})$$

as  $A_{n-k+(i-1)} \subset D_{n-k+i}$ . Thus the required result follows by Lemma 4.2.

□

**Theorem 4.4.** *Let  $n \geq 3$ , then*

$$\begin{aligned} \text{depth}(S/I(C_n^k)) &= 1, \quad \text{if } n \leq 2k + 1; \\ \text{depth}(S/I(C_n^k)) &\geq \lceil \frac{n-k}{2k+1} \rceil, \quad \text{if } n \geq 2k + 2. \end{aligned}$$

*Proof.* (a): Let  $n \leq 2k + 1$ , then by [8, Corollary 10.3.7],  $\text{depth}(S/I(C_n^k)) = 1$ .

(b): Let  $n \geq 2k + 2$ , we consider the following cases:

(1): If  $k = 1$ , then by [4, Proposition 1.3]  $\text{depth}(S/I(C_n^1)) = \lceil \frac{n-1}{3} \rceil$ .

(2): If  $k \geq 2$  and  $2k + 2 \leq n \leq 3k + 1$ , then we have  $\text{depth}(S/I(C_n^k)) \geq 1 = \lceil \frac{n-k}{2k+1} \rceil$  as  $\mathfrak{m} \notin \text{Ass}(S/I(C_n^k))$ .

(3): If  $k \geq 2$  and  $n \geq 3k + 2$ , consider the family of short exact sequences:

$$\begin{aligned}
0 \longrightarrow S/(I(C_n^k) : x_{n-k}) &= S/((I(C_n^k), A_{n-k-1}) : x_{n-k}) \xrightarrow{\cdot x_{n-k}} S/I(C_n^k) \longrightarrow S/(I(C_n^k), A_{n-k}) \longrightarrow 0 \\
0 \longrightarrow S/((I(C_n^k), A_{n-k}) : x_{n-k+1}) &\xrightarrow{\cdot x_{n-k+1}} S/(I(C_n^k), A_{n-k}) \longrightarrow S/(I(C_n^k), A_{n-k+1}) \longrightarrow 0 \\
&\vdots \\
0 \longrightarrow S/((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) &\xrightarrow{\cdot x_{n-k+i}} S/(I(C_n^k), A_{n-k+(i-1)}) \longrightarrow S/(I(C_n^k), A_{n-k+i}) \longrightarrow 0 \\
&\vdots \\
0 \longrightarrow S/((I(C_n^k), A_{n-2}) : x_{n-1}) &\xrightarrow{\cdot x_{n-1}} S/(I(C_n^k), A_{n-2}) \longrightarrow S/(I(C_n^k), A_{n-1}) \longrightarrow 0
\end{aligned}$$

By Lemma 4.1 we have

$$S/(I(C_n^k), A_{n-1}) \cong S_{n-k}/I(P_{n-k}^k).$$

Let  $0 \leq i \leq k-1$ , then by Lemma 4.3, we have

$$S/((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i}) \cong S_{n-2k-1}/I(P_{n-2k-1}^k)[x_{n-k+i}].$$

By Theorem 3.7 and [9, Lemma 3.6], we have

$$\text{depth}(S/((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i})) = \lceil \frac{n-2k-1}{2k+1} \rceil + 1 = \lceil \frac{n}{2k+1} \rceil.$$

Again by Theorem 3.7, we have

$$\text{depth}(S/(I(C_n^k), A_{n-1})) = \lceil \frac{n-k}{2k+1} \rceil.$$

Thus by Depth Lemma  $\text{depth}(S/I(C_n^k)) \geq \lceil \frac{n-k}{2k+1} \rceil$ .

□

**Corollary 4.5.** *Let  $n \geq 3$ , if  $n \leq 2k+1$ , then  $\text{depth}(S/I(C_n^k)) = 1$ . If  $n \geq 2k+2$ , then*

$$\begin{aligned}
\text{depth}(S/I(C_n^k)) &= \lceil \frac{n}{2k+1} \rceil, & \text{if } n \equiv 0, k+1, \dots, n-1 \pmod{2k+1}; \\
\lceil \frac{n}{2k+1} \rceil - 1 &\leq \text{depth}(S/I(C_n^k)) \leq \lceil \frac{n}{2k+1} \rceil, & \text{if } n \equiv 1, \dots, k \pmod{2k+1}.
\end{aligned}$$

*Proof.* By Theorem 4.4, it is enough to prove that  $\text{depth}(S/I(C_n^k)) \leq \lceil \frac{n}{2k+1} \rceil$ , for  $k \geq 2$  and  $n \geq 2k+2$ . Since  $x_{n-k} \notin I(C_n^k)$ , thus by [21, Corollary 1.3] we have

$$\text{depth}(S/I(C_n^k)) \leq \text{depth}(S/(I(C_n^k) : x_{n-k})).$$

Now we consider two cases:

(1): Let  $2k+2 \leq n \leq 3k+1$ , then  $S/(I(C_n^k) : x_{n-k}) = S/(I(P_n^k) : x_{n-k})$  so by the proof of Theorem 3.7 we have  $\text{depth}(S/(I(P_n^k) : x_{n-k})) = 2 = \lceil \frac{n}{2k+1} \rceil$ . Therefore

$$\text{depth}(S/I(C_n^k)) \leq \text{depth}(S/(I(C_n^k) : x_{n-k})) = 2 = \lceil \frac{n}{2k+1} \rceil.$$

(2): Let  $n \geq 3k+2$ , then by Lemma 4.2,  $S/(I(C_n^k) : x_{n-k}) \cong S_{n-2k-1}/I(P_{n-2k-1}^k)[x_{n-k}]$ . By [9, Lemma 3.6] and Theorem 3.7,  $\text{depth}(S_{n-2k-1}/I(P_{n-2k-1}^k)[x_{n-k}]) = \lceil \frac{n}{2k+1} \rceil$ . Thus

$$\text{depth}(S/I(C_n^k)) \leq \text{depth}(S/(I(C_n^k) : x_{n-k})) = \lceil \frac{n}{2k+1} \rceil.$$

□

**Theorem 4.6.** *Let  $n \geq 3$ , then*

$$\begin{aligned}
\text{sdepth}(S/I(C_n^k)) &= 1, & \text{if } n \leq 2k+1; \\
\text{sdepth}(S/I(C_n^k)) &\geq \lceil \frac{n-k}{2k+1} \rceil, & \text{if } n \geq 2k+2.
\end{aligned}$$

*Proof.* (a): Let  $n \leq 2k+1$ , then  $\text{sdepth}(S/I(C_n^k)) = 1$  by [3, Theorem 1.1].

(b): Let  $n \geq 2k+2$ , consider the following cases:

- (1): If  $k = 1$ , then by [4, Proposition 1.8]  $\text{sdepth}(S/I(C_n^1)) \geq \lceil \frac{n-1}{3} \rceil$ .  
(2): If  $k \geq 2$ ,  $2k+2 \leq n \leq 3k+1$ , then  $\text{depth}(S/I(C_n^k)) \geq 1$  as  $\mathfrak{m} \notin \text{Ass}(S/I(C_n^k))$ , thus by [5, Theorem 1.4],  $\text{sdepth}(S/I(C_n^k)) \geq 1 = \lceil \frac{n-k}{2k+1} \rceil$ .  
(3): If  $k \geq 2$ ,  $n \geq 3k+2$ , this theorem can be proved by using similar arguments of Theorem 4.4. We apply Lemma 2.11 on the exact sequences of Theorem 4.4 we get

$$\text{sdepth}(S/I(C_n^k)) \geq \min \{ \text{sdepth}(S/(I(C_n^k) : x_{n-k})), \text{sdepth}(S/(I(C_n^k), A_{n-1})), \\ \min_{i=1}^{k-1} \{ \text{sdepth}(S/((I(C_n^k), A_{n-k+(i-1)}) : x_{n-k+i})) \} \} \geq \lceil \frac{n-k}{2k+1} \rceil.$$

□

**Corollary 4.7.** *Let  $n \geq 3$ , if  $n \leq 2k+1$ , then  $\text{sdepth}(S/I(C_n^k)) = 1$ . If  $n \geq 2k+2$ , then*

$$\text{sdepth}(S/I(C_n^k)) = \lceil \frac{n}{2k+1} \rceil, \quad \text{if } n \equiv 0, k+1, \dots, n-1 \pmod{2k+1}; \\ \lceil \frac{n}{2k+1} \rceil - 1 \leq \text{sdepth}(S/I(C_n^k)) \leq \lceil \frac{n}{2k+1} \rceil, \quad \text{if } n \equiv 1, \dots, k \pmod{2k+1}.$$

*Proof.* When  $k = 1$ , then by [4, Theorem 1.9],  $\text{sdepth}(S/I(C_n^k)) \leq \lceil \frac{n}{3} \rceil$ . By Theorem 4.6 it is enough to prove that  $\text{sdepth}(S/I(C_n^k)) \leq \lceil \frac{n}{2k+1} \rceil$ , for  $k \geq 2$  and  $n \geq 2k+2$ . The proof of Corollary 4.5 also works for Stanley depth by using [2, Proposition 2.7] instead of [21, Corollary 1.3] and Theorem 3.11 instead of Theorem 3.7. □

## 5. LOWER BOUNDS FOR STANLEY DEPTH OF EDGE IDEALS OF $k^{\text{th}}$ POWERS OF PATHS AND CYCLES AND A CONJECTURE OF HERZOG

In this section we compute some lower bounds for Stanley depth of  $I(P_n^k)$  and  $I(C_n^k)$ . These bounds are good enough to prove that Conjecture 1.1 is true for  $I(P_n^k)$  and  $I(C_n^k)$ . Let  $0 \leq i \leq k-1$ , define

$$R_{n-k+i} := K[\{x_1, x_2, \dots, x_n\} \setminus \{x_{n-k}, x_{n-k+1}, \dots, x_{n-k+i}\}]$$

and

$$B'_{n-k+i} := (N_{P_n^k}(x_{n-k+i}) \setminus \{x_{n-k}, x_{n-k+1}, \dots, x_{n-k+(i-1)}\}),$$

where  $R_{n-k+i}$  is a subring of  $S$  and  $B'_{n-k+i}$  is a monomial prime ideal of  $S$  generated by

$$N_{P_n^k}(x_{n-k+i}) \setminus \{x_{n-k}, x_{n-k+1}, \dots, x_{n-k+(i-1)}\}.$$

Let  $I \subset Z = K[x_{i_1}, x_{i_1+1}, \dots, x_{i_r}]$  be a monomial ideal and let  $Z' := Z[x_{i_r+1}]$ , then we write  $I Z' = I[x_{i_r+1}]$ . We recall in the following a useful remark of Cimpoeas.

**Remark 5.1.** [2, Remark 1.7] Let  $I$  be a monomial ideal of  $S$ , and  $I' = (I, x_{n+1}, x_{n+2}, \dots, x_{n+m})$  be a monomial ideal of  $S' = S[x_{n+1}, x_{n+2}, \dots, x_{n+m}]$ , then

$$\text{sdepth}_{S'}(I') \geq \min \{ \text{sdepth}_S(I) + m, \text{sdepth}_S(S/I) + \lceil \frac{m}{2} \rceil \}.$$

**Theorem 5.2.** *Let  $n \geq 2$ , then  $\text{sdepth}(I(P_n^k)) \geq \lceil \frac{n}{2k+1} \rceil + 1$ .*

*Proof.* (a): Let  $n \leq 2k+1$ , since the minimal generators of  $I(P_n^k)$  have degree 2, so by [13, Lemma 2.1]  $\text{sdepth}(I(P_n^k)) \geq 2 = \lceil \frac{n}{2k+1} \rceil + 1$ .

(b): Let  $n \geq 2k+2$ , if  $k = 1$ , then by [17, Theorem 2.3] we have  $\text{sdepth}(I(P_n^1)) \geq n - \lfloor \frac{n-1}{2} \rfloor = \lceil \frac{n-1}{2} \rceil + 1 \geq \lceil \frac{n}{3} \rceil + 1$ . Now let  $k \geq 2$ , we prove this result by induction on  $n$ . We consider the following decomposition of  $I(P_n^k)$  as a vector space:

$$I(P_n^k) = I(P_n^k) \cap R_{n-k} \bigoplus x_{n-k}(I(P_n^k) : x_{n-k})S.$$

Similarly, we can decompose  $I(P_n^k) \cap R_{n-k}$  by the following:

$$I(P_n^k) \cap R_{n-k} = I(P_n^k) \cap R_{n-k+1} \bigoplus x_{n-k+1}(I(P_n^k) \cap R_{n-k} : x_{n-k+1})R_{n-k}.$$

Continuing in the same way for  $1 \leq i \leq k-1$  we have

$$I(P_n^k) \cap R_{n-k+i} = I(P_n^k) \cap R_{n-k+(i+1)} \bigoplus x_{n-k+(i+1)}(I(P_n^k) \cap R_{n-k+i} : x_{n-k+(i+1)})R_{n-k+i}.$$

Finally we get the following decomposition of  $I(P_n^k)$ :

$$I(P_n^k) = I(P_n^k) \cap R_{n-1} \bigoplus_{i=1}^{k-1} \bigoplus x_{n-k+i}(I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i} \bigoplus x_{n-k}(I(P_n^k) : x_{n-k})S.$$

Therefore

$$\text{sdepth}(I(P_n^k)) \geq \min \left\{ \text{sdepth}(I(P_n^k) \cap R_{n-1}), \text{sdepth}((I(P_n^k) : x_{n-k})S), \min_{i=1}^{k-1} \{ \text{sdepth}((I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i}) \} \right\} \quad (5.1)$$

Now

$$I(P_n^k) \cap R_{n-1} = \mathcal{G}(I(P_{n-k-1}^k))[x_n],$$

thus by induction on  $n$  and [9, Lemma 3.6] we have

$$\text{sdepth}(I(P_n^k) \cap R_{n-1}) \geq \lceil \frac{n-k-1}{2k+1} \rceil + 1 + 1 \geq \lceil \frac{n}{2k+1} \rceil + 1.$$

Now we need to show that

$$\text{sdepth}((I(P_n^k) : x_{n-k})S) \geq \lceil \frac{n}{2k+1} \rceil + 1$$

and

$$\text{sdepth}((I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i}) \geq \lceil \frac{n}{2k+1} \rceil + 1.$$

For this we consider the following cases:

**(1):** Let  $2k+2 \leq n \leq 3k+1$ .

If  $n = 2k+2$ , then  $(I(P_n^k) : x_{n-k})S = (x_2, \dots, x_{n-k-1}, x_{n-k+1}, \dots, x_n)S$ , thus by [1, Theorem 2.2] and [9, Lemma 3.6] we have

$$\text{sdepth}((I(P_n^k) : x_{n-k})S) = \lceil \frac{n-2}{2} \rceil + 2 \geq \lceil \frac{n}{2k+1} \rceil + 1.$$

If  $2k+3 \leq n \leq 3k+1$ , then by Remark 3.6  $(I(P_n^k) : x_{n-k})S = (\mathcal{G}(I(P_{n-2k-1}^{f(n-k)})), B_{n-k})[x_{n-k}]$ . Since  $\text{sdepth}(I(P_{n-2k-1}^{f(n-k)})) + |\mathcal{G}(B_{n-k})| \geq 2$ , by Remark 3.5 we have

$$\text{sdepth}(S_{n-2k-1}/I(P_{n-2k-1}^{f(n-k)})) + \lceil \frac{|\mathcal{G}(B_{n-k})|}{2} \rceil \geq 2,$$

then by Remark 5.1,

$$\text{sdepth}(\mathcal{G}(I(P_{n-2k-1}^{f(n-k)})), B_{n-k}) \geq 2,$$

and by [9, Lemma 3.6] we have

$$\text{sdepth}((I(P_n^k) : x_{n-k})S) \geq 3 = \lceil \frac{n}{2k+1} \rceil + 1.$$

Now since

$$(I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i} = (\mathcal{G}(I(P_{n-2k-1+i}^{f(n-k+i)})), B'_{n-k+i})[x_{n-k+i}].$$

So by the same arguments we have

$$\text{sdepth}((I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i}) \geq 3 = \lceil \frac{n}{2k+1} \rceil + 1.$$

(2): Let  $n \geq 3k + 2$ , then by the proof of Lemma 3.3

$$(I(P_n^k) : x_{n-k})S = (\mathcal{G}(I(P_{n-2k-1}^k)), B_{n-k})[x_{n-k}]$$

and

$$(I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i} = (\mathcal{G}(I(P_{n-2k-1+i}^k)), B'_{n-k+i})[x_{n-k+i}].$$

By Remark 5.1 we have

$$\begin{aligned} \text{sdepth}(\mathcal{G}(I(P_{n-2k-1}^k)), B_{n-k}) &\geq \\ \min \left\{ \text{sdepth}(\mathcal{G}(I(P_{n-2k-1}^k))) + |\mathcal{G}(B_{n-k})|, \text{sdepth}(S_{n-2k-1}/I(P_{n-2k-1}^k)) + \lceil \frac{|\mathcal{G}(B_{n-k})|}{2} \rceil \right\}. \end{aligned}$$

By induction on  $n$  we have  $\text{sdepth}(\mathcal{G}(I(P_{n-2k-1}^k))) \geq \lceil \frac{n-2k-1}{2k+1} \rceil + 1 = \lceil \frac{n}{2k+1} \rceil$ , and by Theorem 3.11 we have  $\text{sdepth}(S_{n-2k-1}/I(P_{n-2k-1}^k)) = \lceil \frac{n}{2k+1} \rceil - 1$ . Therefore

$$\text{sdepth}(\mathcal{G}(I(P_{n-2k-1}^k)), B_{n-k}) \geq \lceil \frac{n}{2k+1} \rceil + 1.$$

Thus by [9, Lemma 3.6] we have  $\text{sdepth}((I(P_n^k) : x_{n-k})S) > \lceil \frac{n}{2k+1} \rceil + 1$ .

Now using Remark 5.1 again, we get

$$\begin{aligned} \text{sdepth}(\mathcal{G}(I(P_{n-2k-1+i}^k)), B'_{n-k+i}) &\geq \\ \min \left\{ \text{sdepth}(\mathcal{G}(I(P_{n-2k-1+i}^k))) + |\mathcal{G}(B'_{n-k+i})|, \text{sdepth}(S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)) + \lceil \frac{|\mathcal{G}(B'_{n-k+i})|}{2} \rceil \right\}. \end{aligned}$$

By induction on  $n$  we have  $\text{sdepth}(\mathcal{G}(I(P_{n-2k-1+i}^k))) \geq \lceil \frac{n-2k-1+i}{2k+1} \rceil + 1$ , and by Theorem 3.11 we have  $\text{sdepth}(S_{n-2k-1+i}/I(P_{n-2k-1+i}^k)) = \lceil \frac{n-2k-1+i}{2k+1} \rceil$ . Therefore

$$\text{sdepth}(\mathcal{G}(I(P_{n-2k-1+i}^k)), B'_{n-k+i}) \geq \lceil \frac{n-2k-1+i}{2k+1} \rceil + 1.$$

Thus by [9, Lemma 3.6],  $\text{sdepth}((I(P_n^k) \cap R_{n-k+(i-1)} : x_{n-k+i})R_{n-k+i}) \geq \lceil \frac{n}{2k+1} \rceil + 1$ . This completes the proof.  $\square$

**Proposition 5.3.** *Let  $n \geq 2k + 1$ , then  $\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq \lceil \frac{n+k+1}{2k+1} \rceil$ .*

*Proof.* When  $k = 1$ , then by [4, Proposition 1.10] we have the required result. Now assume that  $k \geq 2$  and consider the following cases:

- (1): If  $2k + 1 \leq n \leq 3k + 1$ . Since  $I(C_n^k)$  is a monomial ideal generated by degree 2 so by [9, Theorem 2.1]  $\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq 2 = \lceil \frac{n+k+1}{2k+1} \rceil$ .
- (2): When  $3k+2 \leq n \leq 4k+1$ , then we use [9] to show that there exist Stanley decompositions of desired Stanley depth. Let  $s \in \{1, 2, \dots, k\}$ ,  $j_s \in \{1, 2, \dots, k+1-s\}$  and

$$L := \bigoplus_{s=1}^k \left( \bigoplus_{j_s=1}^{k+1-s} x_{j_s} x_{n+1-s} K[x_{j_s}, x_{j_s+k+1}, x_{n+1-s}] \right).$$

It is easy to see that  $L \subset I(C_n^k) \setminus I(P_n^k)$ . Now let  $u_i \in I(C_n^k) \setminus I(P_n^k)$  be a squarefree monomial such that  $u_i \notin L$  then clearly  $\deg(u_i) \geq 3$ . Since

$$I(C_n^k)/I(P_n^k) \cong L \bigoplus_{u_i} u_i K[\text{supp}(u_i)]$$

Thus  $\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq 3 = \lceil \frac{n+k+1}{2k+1} \rceil$  as required.

- (3): Now, assume that  $n \geq 4k + 2$ . We have the following  $K$ -vector space isomorphism:

$$I(C_n^k)/I(P_n^k) \cong \bigoplus_{j_1=1}^k x_{j_1} x_n \frac{K[x_{j_1+k+1}, x_{j_1+k+2}, \dots, x_{n-k-1}]}{(x_{j_1+k+1}x_{j_1+k+2}, x_{j_1+k+1}x_{j_1+k+3}, \dots, x_{n-k-2}x_{n-k-1})} [x_{j_1}, x_n]$$

$$\begin{aligned}
& \bigoplus_{j_2=1}^{k-1} \bigoplus_{j_2=1}^{k-1} x_{j_2} x_{n-1} \frac{K[x_{j_2+k+1}, x_{j_2+k+2}, \dots, x_{n-k-2}]}{(x_{j_2+k+1}x_{j_2+k+2}, x_{j_2+k+1}x_{j_1+k+3}, \dots, x_{n-k-3}x_{n-k-2})} [x_{j_2}, x_{n-1}] \bigoplus \\
& \quad \dots \\
& \bigoplus_{j_{k-1}=1}^2 x_{j_{k-1}} x_{n-(k-2)} \frac{K[x_{j_{k-1}+k+1}, x_{j_{k-1}+k+2}, \dots, x_{n-2k+1}]}{(x_{j_{k-1}+k+1}x_{j_{k-1}+k+2}, x_{j_{k-1}+k+1}x_{j_{k-1}+k+3}, \dots, x_{n-2k}x_{n-2k+1})} [x_{j_{k-1}}, x_{n-k+2}] \\
& \quad \bigoplus x_1 x_{n-(k-1)} \frac{K[x_{k+2}, x_{k+3}, \dots, x_{n-2k}]}{(x_{k+2}x_{k+3}, x_{k+2}x_{k+4}, \dots, x_{n-2k-1}x_{n-2k})} [x_1, x_{n-(k-1)}].
\end{aligned}$$

Thus

$$I(C_n^k)/I(P_n^k) \cong \bigoplus_{s=1}^k \left( \bigoplus_{j_s=1}^{k+1-s} x_{j_s} x_{n+1-s} (S_{j_s+k+1, n-s-k} / (\mathcal{G}(I(P_n^k)) \cap S_{j_s+k+1, n-s-k}) [x_{j_s}, x_{n+1-s}]) \right),$$

where  $S_{j_s+k+1, n-s-k} = K[x_{j_s+k+1}, x_{j_s+k+2}, \dots, x_{n-s-k}]$ . Indeed, if  $u \in I(C_n^k)$  such that  $u \notin I(P_n^k)$  then  $(x_{j_s} x_{n+1-s})|u$ , for only one pair of  $s$  and  $j_s$ . If  $(x_{j_s} x_{n+1-s})|u$ , then  $u = x_{j_s}^\gamma x_{n+1-s}^\delta v$ ,  $v \in S_{j_s+k+1, n-s-k}$ , since  $v \notin I(P_n^k)$ , it follows that  $v \notin \mathcal{G}(I(P_n^k)) \cap S_{j_s+k+1, n-s-k}$ . Clearly,

$$S_{j_s+k+1, n-s-k} / \mathcal{G}(I(P_n^k)) \cap S_{j_s+k+1, n-s-k} \cong S_{n-(j_s+2k+s)} / I(P_{n-(j_s+2k+s)}^k).$$

Thus by Theorem 3.11 and [9, Lemma 3.6], we have

$$\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq \min_{s=1}^k \left\lceil \frac{n - (j_s + s + 2k)}{2k+1} \right\rceil + 2\}.$$

It is easy to see that  $\max\{j_s + s\} = k+1$ . Therefore

$$\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq \left\lceil \frac{n - (3k+1)}{2k+1} \right\rceil + 2 = \left\lceil \frac{n+k+1}{2k+1} \right\rceil.$$

□

**Theorem 5.4.** *Let  $n \geq 3$ , then*

$$\begin{aligned}
\text{sdepth}(I(C_n^k)) &\geq 2, \quad \text{if } n \leq 2k+1; \\
\text{sdepth}(I(C_n^k)) &\geq \left\lceil \frac{n-k}{2k+1} \right\rceil + 1, \quad \text{if } n \geq 2k+2.
\end{aligned}$$

*Proof.* (a): Let  $n \leq 2k+1$ , since the minimal generators of  $I(C_n^k)$  have degree 2, so by [13, Lemma 2.1]  $\text{sdepth}(I(C_n^k)) \geq 2$ .

(b): Let  $n \geq 2k+2$ , consider the short exact sequence

$$0 \longrightarrow I(P_n^k) \longrightarrow I(C_n^k) \longrightarrow I(C_n^k)/I(P_n^k) \longrightarrow 0,$$

then by Lemma 2.11,

$$\text{sdepth}(I(C_n^k)) \geq \min\{\text{sdepth}(I(P_n^k)), \text{sdepth}(I(C_n^k)/I(P_n^k))\}.$$

By Theorem 5.2,

$$\text{sdepth}(I(P_n^k)) \geq \left\lceil \frac{n}{2k+1} \right\rceil + 1,$$

and by Proposition 5.3, we have

$$\text{sdepth}(I(C_n^k)/I(P_n^k)) \geq \left\lceil \frac{n+k+1}{2k+1} \right\rceil = \left\lceil \frac{n-k}{2k+1} \right\rceil + 1,$$

this finishes the proof.

□

**Corollary 5.5.** *Let  $n \geq 3$ , if  $n \leq 2k + 1$ , then*

$$\text{sdepth}(I(C_n^k)) \geq 2 = \text{sdepth}(S/I(C_n^k)) + 1.$$

*If  $n \geq 2k + 2$ , then*

$$\begin{aligned} \text{sdepth}(I(C_n^k)) &\geq \text{sdepth}(S/I(C_n^k)), \text{ if } n \equiv 1, \dots, k \pmod{2k+1}; \\ \text{sdepth}(I(C_n^k)) &\geq \text{sdepth}(S/I(C_n^k)) + 1, \text{ if } n \equiv 0, k+1, \dots, n-1 \pmod{2k+1}. \end{aligned}$$

*Proof.* Proof follows by Corollary 4.7 and Theorem 5.4. □

## REFERENCES

- [1] C. Biro, D. M. Howard, M. T. Keller, W. T. Trotter, S. J. Young, *Interval partitions and Stanley depth*, Journal of Combinatorial Theory, Series A, 117 (2010), 475-482.
- [2] M. Cimpoeas, *Several inequalities regarding Stanley depth*, Romanian Journal of Mathematics and Computer Science, 2(2012), 28-40.
- [3] M. Cimpoeas, *Stanley depth of squarefree Veronese ideals*, An. St. Univ. Ovidius Constanta, 21(3)(2013), 67-71.
- [4] M. Cimpoeas, *On the Stanley depth of edge ideals of line and cyclic graphs*, Romanian Journal of Mathematics and Computer Science, 5(1)(2015), 70-75.
- [5] M. Cimpoeas, *Some remarks on the Stanley depth for multigraded modules*, Le Matematiche Vol. LXIII (2008) Fasc. II, 165-171.
- [6] A. M. Duval, B. Goeckneker, C. J. Klivans, J. L. Martine, *A non-partitionable Cohen-Macaulay simplicial complex*, Advances in Mathematics, 299(2016), 381-395.
- [7] J. Herzog, *A survey on Stanley depth*, In Monomial ideals, computations and applications, vol. 2083 of Lecture Notes in Math. Springer, Heidelberg, 2013, 3-45.
- [8] J. Herzog, T. Hibi, *Monomial Ideals*, Springer-Verlag London Limited, (2011).
- [9] J. Herzog, M. Vladiu, X. Zheng, *How to compute the Stanley depth of a monomial ideal*, J. Algebra, 322(9)(2009), 3151-3169.
- [10] J. Herzog, D. Popescu, M. Vladiu, *Stanley depth and size of a monomial ideal*, Proc. Amer. Math. Soc., 140(2)(2012), 493-504.
- [11] Z. Iqbal, M. Ishaq, M. Aamir, *Depth and Stanley depth of edge ideals of square paths and square cycles*, to appear in Comm. Algebra, <http://dx.doi.org/10.1080/00927872.2017.1339068>,
- [12] M. Ishaq, M. I. Qureshi, *Upper and lower bounds for the Stanley depth of certain classes of monomial ideals and their residue class rings*, Comm. Algebra, 41(3)(2013), 1107-1116.
- [13] M. T. Keller, Y. Shen, N. Streib, S. J. Young, *On the Stanley Depth of Squarefree Veronese Ideals*, Journal of Algebraic Combinatorics, 33(2)(2011), 313-324.
- [14] M. T. Keller, S. J. Young, *Combinatorial reductions for the Stanley depth of  $I$  and  $S/I$* , arXiv:1702.00781v1.
- [15] M. C. Lin, D. Rautenbach, F. J. Soulignac, J. L. Szwarcfiter, *Powers of cycles, powers of paths, and distance graphs*, Discrete Appl. Math., 159 (2011), 621-627.
- [16] S. Morey, *Depths of powers of the edge ideal of a tree*, Comm. Algebra, 38(11)(2010), 4042-4055.
- [17] R. Okazaki, *A lower bound of Stanley depth of monomial ideals*, J. Commut. Algebra, 3(1)(2011), 83-88.
- [18] D. Popescu, M. I. Qureshi, *Computing the Stanley depth*, J. Algebra, 323(10), 2943-2959.
- [19] D. Popescu, *Stanley conjecture on intersection of four monomial prime ideals*, Comm. Algebra, 41(11)(2013), 4351-4362.
- [20] M. R. Pournaki, S. A. Seyed Fakhari, S. Yassemi, *Stanley depth of powers of the edge ideals of a forest*, Proc. Amer. Math. Soc., 141(10)(2013), 3327-3336.
- [21] A. Rauf, *Depth and Stanley depth of multigraded modules*, Comm. Algebra, 38(2)(2010), 773-784.
- [22] R. P. Stanley, *Linear Diophantine equations and local cohomology*, Invent. Math. 68(2)(1982), 175-193.
- [23] A. Stefan, *Stanley depth of powers of path ideal*, <http://arxiv.org/pdf/1409.6072.pdf>.
- [24] Z. Tang, *Stanley depths of certain Stanley-Reisner rings*, J. Algebra, 409(1)(2014), 430-443.

ZAHID IQBAL, SCHOOL OF NATURAL SCIENCES, NATIONAL UNIVERSITY OF SCIENCES AND TECHNOLOGY ISLAMABAD, SECTOR H-12, ISLAMABAD PAKISTAN.

*E-mail address:* 786zahidwarraich@gmail.com

MUHAMMAD ISHAQ, SCHOOL OF NATURAL SCIENCES, NATIONAL UNIVERSITY OF SCIENCES AND TECHNOLOGY ISLAMABAD, SECTOR H-12, ISLAMABAD PAKISTAN.

*E-mail address:* ishaq-maths@yahoo.com