

KÄHLER-EINSTEIN METRICS

GÁBOR SZÉKELYHIDI

Dedicated to Sir Simon Donaldson on the occasion of his 60th birthday

ABSTRACT. We survey the theory of Kähler-Einstein metrics, with particular focus on the circle of ideas surrounding the Yau-Tian-Donaldson conjecture for Fano manifolds.

1. INTRODUCTION

A starting point in the study of Kähler-Einstein metrics is Kähler's observation [104], that for Hermitian metrics satisfying what is now known as the Kähler condition, the Einstein equations reduce to a scalar complex Monge-Ampère equation. Over the many decades since, the field has grown into a very rich subject with deep connections to nonlinear PDE, geometric analysis, complex algebraic geometry, string theory, and others. The goal of this survey is to give an overview of some of these developments and in particular to showcase the diverse ideas that have been brought to bear on the problem.

Let us start with Kähler's observation, and consider a Hermitian metric $g_{j\bar{k}}$ on a complex manifold M . The associated (1,1)-form, or Kähler form, is defined to be

$$\omega = \sqrt{-1}g_{j\bar{k}}dz^j \wedge d\bar{z}^k$$

in local coordinates, and the metric g is Kähler if $d\omega = 0$. Kähler showed that in this case we can locally write the metric g in terms of a potential function ϕ :

$$g_{j\bar{k}} = \frac{\partial^2 \phi}{\partial z^j \partial \bar{z}^k}.$$

The Ricci curvature of g is then given by

$$\text{Ric}_{j\bar{k}} = -\frac{\partial^2}{\partial z^j \partial \bar{z}^k} \log \det(g),$$

and so we can obtain solutions of the Einstein equation $\text{Ric} = \lambda g$, by solving the scalar equation

$$\det \left(\frac{\partial^2 \phi}{\partial z^j \partial \bar{z}^k} \right) = e^{-\lambda \phi}.$$

Under certain conditions Kähler potentials exist globally, not just locally. Let us suppose that M is compact. A Kähler form ω on M defines a cohomology class $[\omega] \in H^2(M)$, and it is natural to consider, as Calabi [28] did, the space of all Kähler forms on M in a fixed cohomology class. The $\partial\bar{\partial}$ -lemma states that any other Kähler form $\eta \in [\omega]$ can be written as

$$\eta = \omega + \sqrt{-1}\partial\bar{\partial}\phi$$

for a function $\phi : M \rightarrow \mathbf{R}$, and so the space of Kähler metrics in a fixed cohomology class are parametrized by scalar functions, in analogy with a conformal class in Riemannian geometry.

A further important observation is that for any Kähler metric g on M , its Ricci form

$$\text{Ric}(g) = \sqrt{-1} R_{j\bar{k}} dz^j \wedge d\bar{z}^k$$

is a closed form in the first Chern class $c_1(M)$. Calabi [28] conjectured that conversely any representative of $c_1(M)$ is the Ricci form of a unique Kähler metric in every Kähler class. This fundamental conjecture was proven by Yau [177], by solving the complex Monge-Ampère equation

$$(\omega + \sqrt{-1} \partial \bar{\partial} \phi)^n = e^{F+c} \omega^n$$

for ϕ and a constant c , given a Kähler form ω and function F . Perhaps the most important case, which has had an enormous impact, is when $c_1(M) = 0$. In this case Yau's result implies that every Kähler class on M admits a unique Ricci flat metric.

More generally, if we seek a Kähler-Einstein metric ω satisfying $\text{Ric}(\omega) = \lambda \omega$, then we must have $c_1(M) = \lambda[\omega]$. In particular if $\lambda \neq 0$, then either $c_1(M)$ or $-c_1(M)$ must be a Kähler class, and the cohomology class $[\omega]$ is uniquely determined. When $c_1(M)$ is negative, then the works of Yau [177] and Aubin [9] yield a Kähler-Einstein metric on M . It was already known by Matsushima [126], however, that when $c_1(M)$ is positive, i.e. M is Fano, then M can only admit a Kähler-Einstein metric if its holomorphic automorphism group is reductive. Later Futaki [93] found a different obstruction stemming from the automorphism group, showing that a certain numerical invariant $F(v)$ must vanish for all holomorphic vector fields v on M . These obstructions rule out the existence of a Kähler-Einstein metric on the blowup $\text{Bl}_p \mathbf{P}^2$ for instance. On the other hand, Tian [165] showed that in the case of Fano surfaces the reductivity of the automorphism group, or alternatively the vanishing of Futaki's obstruction, is actually sufficient for the existence of a Kähler-Einstein metric.

At this point let us digress briefly on parallel developments in the theory of holomorphic vector bundles. In algebraic geometry a basic problem is to construct moduli spaces of various objects, for instance vector bundles over a curve. It turns out that in general it is not possible to parametrize all vector bundles of a fixed topological type with a nice space, but rather we need to restrict ourselves to semistable bundles – a notion introduced by Mumford [127]. While this is a purely algebro-geometric notion, it was shown by Narasimhan-Seshadri [129], and later reproved by Donaldson [69], that stability has a differential geometric meaning: an indecomposable vector bundle over a curve is stable if and only if it admits a Hermitian metric with constant curvature. The Hitchin-Kobayashi correspondence, proved by Donaldson [70, 71] and Uhlenbeck-Yau [173], is the higher dimensional generalization of this, stating that an indecomposable vector bundle is stable if and only if it admits a Hermitian-Einstein metric. There is a particularly rich interplay between this result for complex surfaces and Donaldson theory [85] for smooth four-manifolds.

In analogy with these results on vector bundles, Yau [178] conjectured that the existence of a Kähler-Einstein metric on a Fano manifold M should be related to the stability of M in a suitable sense. This conjecture was made precise by Tian [166],

who generalized Futaki's obstruction [93] to the notion of K-stability: Tian showed (see also Ding-Tian [65]) that given any \mathbf{C}^* -equivariant family $\pi : X \rightarrow \mathbf{C}$ with generic fiber $X_t \cong M$, and \mathbf{Q} -Fano central fiber X_0 , the Futaki invariant $F(X)$ of the induced vector field on the central fiber can be defined. The \mathbf{Q} -Fano condition here means that X_0 is a normal variety with \mathbf{Q} -Cartier anticanonical divisor, and this assumption allows for a differential-geometric definition of $F(X)$. Tian showed moreover that if M admits a Kähler-Einstein metric, then $F(X) \geq 0$ for all such families, with equality only if X is a product. This obstruction is called K-stability, and it is a far reaching generalization of Futaki's obstruction. Indeed the latter can be viewed as a special case using only product families.

The Donaldson-Uhlenbeck-Yau theorem and Yau and Tian's conjectures on the existence of Kähler-Einstein metrics can be seen as two instances of a relationship between quotient constructions in symplectic and algebraic geometry, due to Kempf-Ness [106]. This is because in both settings the geometric structure we seek, a Hermitian-Einstein metric or a Kähler-Einstein metric, can be viewed as a zero of a moment map. This was discovered by Atiyah-Bott [8] for vector bundles over curves, and independently by Fujiki [90] and Donaldson [72] for Kähler-Einstein metrics. In fact even more generally, constant scalar curvature Kähler metrics, and the extremal Kähler metrics introduced by Calabi [29] fit into this framework.

Motivated by this, Donaldson [77] introduced a generalization of K-stability for any pair (M, L) of a projective manifold M equipped with an ample line bundle L . The definition is similar to Tian's notion, in that we need to consider \mathbf{C}^* -equivariant degenerations $\pi : X \rightarrow \mathbf{C}$ of M , compatible with the polarization L of M . The central fiber, however, is allowed to be a singular scheme, and the corresponding numerical invariant, the Donaldson-Futaki invariant $DF(X)$, is defined purely algebraically. In this generality we have

Conjecture 1 (Yau-Tian-Donaldson). *The manifold M admits a constant scalar curvature Kähler metric in $c_1(L)$, if and only if the pair (M, L) is K-stable.*

The conjecture can be extended [158] to characterize the existence of extremal metrics, and there are also variants for more general "twisted" equations by Derivan [62]. One direction of the conjecture is fairly well understood, namely that the existence of a constant scalar curvature metric implies K-stability (see e.g. Tian [166], Donaldson [75, 78] and Stoppa [155], Berman-Darvas-Lu [19]), however the converse in general is wide open at present.

The main subject of this survey is the case when M is a Fano manifold and $L = -K_M$, since then a constant scalar curvature metric in $c_1(L)$ is actually Kähler-Einstein. In this case Chen-Donaldson-Sun [41, 42, 43, 44] proved the following breakthrough result.

Theorem 2. *A Fano manifold M admits a Kähler-Einstein metric if and only if $(M, -K_M)$ is K-stable.*

Our aim in this survey is not so much to describe the proof of this result, but rather to highlight the diversity of ideas that are in some way related to the Yau-Tian-Donaldson conjecture. There are several other excellent surveys on the subject, such as those of Thomas [162], Phong-Sturm [139] and Eyssidieux [86], with a focus on different aspects of the theory.

The solution of the conjecture in the Fano case has certainly closed a chapter, but it has also set the scene for a great deal of further development, much of which is likely yet to come.

Acknowledgements. It is my pleasure to thank Simon Donaldson for his advice and support over the years – it would be hard to overstate the influence that his ideas and approach to mathematics have had on my interests.

I also thank Julius Ross and Valentino Tosatti for helpful comments on this survey. This work was supported in part by NSF grant DMS-1350696.

2. THE MOMENT MAP PICTURE

In this section we describe how the scalar curvature of a Kähler metric can be viewed as an infinite dimensional moment map, following Donaldson [72]. This point of view is invaluable in building intuition for the problem, and in retrospect it motivates many of the basic constructions and results that were known beforehand. We will keep the discussion at a formal level, and not delve into the precise definitions relating to infinite dimensional manifolds.

Let (X, ω) denote a compact symplectic manifold, such that $H^1(X) = 0$ for simplicity. Let \mathcal{J} be the space of almost complex structures on X , compatible with ω . The space \mathcal{J} has a natural complex structure, and each tangent space $T_J\mathcal{J}$ is equipped with the L^2 -inner product given by the metric $g_J(\cdot, \cdot) = \omega(\cdot, J\cdot)$. This structure turns \mathcal{J} into an infinite dimensional Kähler manifold, and the group $\mathcal{G} = \text{Ham}(X, \omega)$ of Hamiltonian symplectomorphisms acts on \mathcal{J} , preserving this Kähler structure. We identify the Lie algebra of \mathcal{G} with the functions $C_0^\infty(X)$ with zero mean on X with respect to the volume form ω^n , through the Hamiltonian construction. We further identify $C_0^\infty(X)$ with its dual using the L^2 inner product. The key calculation is the following.

Proposition 3 (Fujiki [90], Donaldson [72]). *A moment map for the action of \mathcal{G} on \mathcal{J} is given by*

$$\begin{aligned} \mu : \mathcal{J} &\rightarrow C_0^\infty(X) \\ J &\mapsto \overline{S} - S_J, \end{aligned}$$

where S_J is the scalar curvature of the metric g_J whenever J is integrable, and \overline{S} is its average, which is independent of J .

In particular an integrable complex structure J satisfies $\mu(J) = 0$ if and only if the Kähler metric g_J on X has constant scalar curvature. The precise meaning of this result is an identity relating the linearization of the scalar curvature S_J under varying the complex structure J , and the infinitesimal action of Hamiltonian symplectomorphisms on \mathcal{J} . Indeed, let $h \in C_0^\infty(M)$, and let $A \in T_J\mathcal{J}$ be an infinitesimal variation of J . The variation of J by the Hamiltonian vector field v_h is the Lie derivative $L_{v_h}J$, and we write $DS_J(A)$ for the variation of the scalar curvature S_J in the direction A . The content of Proposition 3 is the identity

$$\langle DS_J(A), h \rangle_{L^2} = \langle JA, L_{v_h}J \rangle_{L^2},$$

which can be checked by direct calculation. See [90, 72, 167, 94] for the details.

Suppose for the moment that instead of the infinite dimensional group \mathcal{G} acting on \mathcal{J} , we had a compact group G acting on a compact Kähler manifold (V, ω) , with moment map μ . Let us assume that ω is the curvature form of a line bundle

$L \rightarrow V$ endowed with a Hermitian metric, and so V is in fact a projective manifold. Let G^c denote the complexification of G , acting on V by biholomorphisms. The Kempf-Ness theorem [106] says, in this finite dimensional situation, that a G^c -orbit contains a zero of the moment map if and only if it is polystable. One way to define polystability, that is useful in the infinite dimensional setting as well, is that an orbit $G^c \cdot p$ is polystable if a suitable real valued G -invariant function is proper on the orbit. More precisely we consider the function

$$(2.1) \quad \begin{aligned} f : G^c/G &\rightarrow \mathbf{R} \\ [g] &\mapsto \log \|g \cdot \hat{p}\|, \end{aligned}$$

defined using a G -invariant norm on V . Here $\hat{p} \in L$ is a non-zero lift of p , and we need a lift of the G^c -action to the total space of L in a way compatible with the choice of moment map μ . The compatibility of the lift of the action with the moment map μ can be expressed by the formula

$$(2.2) \quad df_{g \cdot p}(\sqrt{-1}\xi) = \langle \mu(g \cdot p), \xi \rangle$$

for the variation of f , where $g \in G^c$, and $\xi \in \mathfrak{g}$. Note in particular that the critical points of f are precisely zeros of the moment map.

Since the function f also turns out to be convex along geodesics in the symmetric space G^c/G , it is clear that properness of f corresponds to the existence of a zero of the moment map in the orbit $G^c \cdot p$, at least if we ignore subtleties related to the possible stabilizer of p . What is less clear, however, is that to verify whether f is proper on G^c/G , it is enough to check whether f is proper along each geodesic ray in G^c/G obtained from one-parameter subgroups $\mathbf{C}^* \subset G^c$. In fact it is enough to consider only one-parameter subgroups of the form $t \mapsto e^{\sqrt{-1}t\xi}$ for $\xi \in \mathfrak{g}$ generating a circle subgroup. For such a one-parameter subgroup we can test the properness of f by computing the limit

$$(2.3) \quad \lim_{t \rightarrow \infty} f'(e^{\sqrt{-1}t\xi} \cdot p) = \langle \mu(q), \xi \rangle,$$

where $q = \lim_{t \rightarrow \infty} e^{\sqrt{-1}t\xi} \cdot p$. Properness of f is then equivalent to $\langle \mu(q), \xi \rangle > 0$ whenever $q \notin G^c \cdot p$. This is in essence the Hilbert-Mumford numerical criterion for stability, proved by Mumford [127], to which we refer the reader for the detailed development of this theory.

Let us return to the infinite dimensional setting of the action of \mathcal{G} on \mathcal{J} . A first issue is that the complexification \mathcal{G}^c does not exist, but we can still try to interpret what its orbits would be if it did. Indeed in each tangent space $T_J \mathcal{J}$ we have a subspace spanned by elements of the form $L_{v_h} J$ giving the infinitesimal action of Hamiltonian vector fields, and we can simply complexify this subspace. The orbits of \mathcal{G}^c then ought to be integral submanifolds of this distribution on \mathcal{J} .

Note that ultimately we are interested in the metrics g_J determined by the pairs (ω, J) , and for any diffeomorphism f the metric given by (ω, f^*J) is isometric to that given by $((f^{-1})^*\omega, J)$. We can therefore switch our point of view from studying different complex structures on a symplectic manifold (X, ω) to studying different Kähler forms on a complex manifold (X, J) , as is more standard in Kähler geometry. To see what this corresponds to in terms of the complexified orbits of \mathcal{G} , note that when J is integrable, then $JL_{v_h}J = L_{Jv_h}J$, and at the same time we have the formula $-L_{Jv_h}\omega = 2\sqrt{-1}\partial\bar{\partial}h$. Using the $\partial\bar{\partial}$ -lemma we can interpret this as saying that in our infinite dimensional setting the role of the symmetric space

G^c/G is played by the space of Kähler metrics in the Kähler class $[\omega]$ (see [73]). In conclusion the Kempf-Ness theorem suggests that the existence of a constant scalar curvature metric in the Kähler class $[\omega]$ is equivalent to stability of this class in a suitable sense.

Let us see how this formal discussion motivates several of the basic constructions in the field, which were actually discovered before the moment map picture was understood:

2.1. The Mabuchi metric. In finite dimensions the metric on the symmetric space G^c/G is given by an inner product on the Lie algebra \mathfrak{g} . In the infinite dimensional setting we chose the L^2 product on Hamiltonian functions, which by the above discussion correspond to variations in the Kähler potential. This leads to a very natural Riemannian structure on the space of Kähler metrics first introduced by Mabuchi [124] and later rediscovered by Semmes [147] and Donaldson [73]. For a compact Kähler manifold (M, ω) , Let us denote by

$$\mathcal{H} = \{\phi \in C^\infty(M) : \omega + \sqrt{-1}\partial\bar{\partial}\phi > 0\}$$

the space of Kähler potentials. For $\phi \in \mathcal{H}$ let $\omega_\phi = \omega + \sqrt{-1}\partial\bar{\partial}\phi$ be the corresponding Kähler metric. When $\omega \in c_1(L)$ for an ample line bundle L , then \mathcal{H} can also be thought of as the space of positively curved Hermitian metrics $e^{-\phi}$ on L .

Each tangent space $T_\phi\mathcal{H}$ can be identified with $C^\infty(M)$, and the Mabuchi metric is defined by simply taking the L^2 inner product:

$$\langle f, g \rangle_\phi = \int_M fg \omega_\phi^n.$$

One can show that this metric turns \mathcal{H} , at least formally, into a non-positively curved symmetric space.

Of great interest is the study of geodesics in \mathcal{H} . A calculation shows that a path $\phi_t \in \mathcal{H}$ is a geodesic, if it satisfies the equation

$$\ddot{\phi}_t - \frac{1}{2}|\nabla \dot{\phi}_t|_{\omega_{\phi_t}}^2 = 0.$$

An important observation due to Semmes and Donaldson, however, is that this geodesic equation is equivalent to a homogeneous complex Monge-Ampère equation. Indeed, let $A_{a,b} = S^1 \times (a, b)$ be a cylinder, and given a path $\phi_t \in \mathcal{H}$ for $a < t < b$, define the form

$$\Omega = \pi^*\omega + \sqrt{-1}\partial\bar{\partial}\phi_t$$

on the product $M \times A_{a,b}$. Here $\pi : M \times A_{a,b} \rightarrow M$ is the projection, and $\sqrt{-1}\partial\bar{\partial}$ involves the variables on $A_{a,b}$ as well. A calculation shows that ϕ_t is a geodesic if and only if Ω is non-negative and $\Omega^{n+1} = 0$, i.e. ϕ_t solves the homogeneous complex Monge-Ampère equation on $M \times A_{a,b}$. When a, b are finite we have geodesic segments, while if a or b is infinite, then we have geodesic rays.

Since the equation $\Omega^{n+1} = 0$ is degenerate elliptic, the regularity theory is very subtle. Chen [47] showed that any two potentials $\phi_0, \phi_1 \in \mathcal{H}$ can be connected by a unique weak geodesic ϕ_t , for which $\Delta\phi_t$ is bounded, using the Laplacian on $M \times A_{a,b}$ (see also Błocki [23]). This was improved to a bound on $|\nabla^2\phi_t|$ by Chu-Tosatti-Weinkove [50] (see also Berman [15] for a weaker result in the projective case). It turns out that these results are essentially optimal, since there are counterexamples to the existence of smooth geodesics, by Lempert-Vivas [110] and Darvas [53] (see

also Donaldson [76] and Ross-Witt Nyström [144]). However even weak geodesics are enough for many applications, as we will describe below.

2.2. The K-energy. In the finite dimensional setting we described how the existence of a zero of the moment map is related to properness of the log-norm functional f in (2.1). In infinite dimensions this translates to the K-energy, defined by Mabuchi [123]. The formula (2.2) for the variation of the log-norm functional suggests that the K-energy $K : \mathcal{H} \rightarrow \mathbf{R}$ can be defined through its variation. If $\phi_t \in \mathcal{H}$ is a path, then

$$\begin{aligned} \frac{d}{dt}K(\phi_t) &= \langle \bar{S} - S(\omega_{\phi_t}), \dot{\phi}_t \rangle_{L^2(\omega_{\phi_t})} \\ &= \int_M \dot{\phi}_t (\bar{S} - S(\omega_{\phi_t})) \omega_{\phi_t}^n. \end{aligned}$$

Mabuchi [124] showed that the K-energy is convex along smooth geodesics in \mathcal{H} , which we now see as a general result about the log-norm functionals. It is clear from the definition that the critical points are constant scalar curvature metrics, and Mabuchi also showed that if two critical points ϕ_0, ϕ_1 are connected by a smooth geodesic, then the metrics ω_{ϕ_0} and ω_{ϕ_1} are isometric by an automorphism of M .

It was only much later that Berman-Berndtsson [17] showed that convexity holds along weak geodesics, and as an application proved a general uniqueness result along these lines. Note that uniqueness in various degrees of generality has been proven previously using other methods, see for example [13, 75, 47, 22].

In the finite dimensional setting the existence of a critical point of the log-norm functional is equivalent to its properness. Tian [166] showed that the analogous result holds for Kähler-Einstein metrics, characterizing their existence in terms of properness of the K-energy in a suitable sense. See also Darvas-Rubinstein [55] for a more precise properness statement in the Kähler-Einstein case, in the presence of automorphisms. In the general constant scalar curvature case Berman-Darvas-Lu [19] showed one direction of this correspondence, namely that the existence of a cscK metric implies properness of the K-energy, as was conjectured by Tian [167].

2.3. The Futaki invariant. A construction that predates both of the previous ones is Futaki's obstruction [93] to the existence of a Kähler-Einstein metric on a Fano manifold M , analogous to the Kazdan-Warner obstruction [105] for the prescribed curvature problem on the 2-sphere.

In retrospect, Futaki's obstruction can be viewed as the first glimpse of the obstruction to Kähler-Einstein metrics given by K-stability. Recall that in the finite dimensional picture, polystability of p is related to the limit (2.3) of the derivative of the log-norm functional along the orbit $e^{\sqrt{-1}t\xi} \cdot \hat{p}$ of a one-parameter subgroup. The simplest example is if $\xi \in \mathfrak{g}$ is in the stabilizer of p , so that the one-parameter subgroup simply acts on the line L_p . The quantity $\langle \mu(p), \xi \rangle$ is then the weight of this action, and polystability requires that this weight vanishes, since otherwise the log-norm functional would not be bounded from below.

The infinite dimensional analog of this weight can be defined as follows. An element $\xi \in \mathfrak{g}$ in the stabilizer of a point p corresponds to a function h on M , whose Hamiltonian vector field v_h preserves the complex structure of M as well, i.e. v_h is a holomorphic Killing field on (M, ω) , and v_h generates a circle action on

M . The corresponding weight is then

$$(2.4) \quad F(v_h) = \int_M h(\bar{S} - S(\omega)) \omega^n.$$

Futaki used a different, but essentially equivalent definition, and showed that $F(v_h)$ only depends on the vector field v_h , and not on the metric in the class $[\omega]$ used in the formula. In addition the invariant can be defined for any holomorphic vector field, not just those that generate circle actions. If M admits a constant scalar curvature metric in the class $[\omega]$, then it is clear from the definition that $F(v) = 0$ for all holomorphic vector fields v .

Tian's definition [65, 166] of K-stability is motivated by probing the properness of the K-energy along more general families of metrics $\phi_t \in \mathcal{H}$. From the finite dimensional picture it is most natural to consider geodesic rays, however this is technically rather difficult. Instead Tian used families of metrics on M obtained from embedding $M \subset \mathbf{P}^N$ into a projective space, and then considering the restrictions of the Fubini-Study metrics $\sigma_t^* \omega_{FS}$ pulled back under a one-parameter family of automorphisms σ_t of \mathbf{P}^N . We will discuss this construction and others in more detail in Section 3.

2.4. The Ding functional. The constructions in the previous subsections apply to the general existence problem for constant scalar curvature metrics, not just Kähler-Einstein metrics. At the same time we will see that the Kähler-Einstein problem has several special features. One of these is an alternative variational description of Kähler-Einstein metrics as critical points of the Ding functional \mathcal{D} defined in [64]. Thinking of \mathcal{H} as the space of positively curved metrics $e^{-\phi}$ on $-K_M$, the variation of \mathcal{D} along a path ϕ_t is defined to be

$$\frac{d}{dt} \mathcal{D}(\phi_t) = -\frac{1}{V} \int_M \dot{\phi}_t \omega_{\phi_t}^n + \frac{\int_M \dot{\phi}_t e^{-\phi_t}}{\int_M e^{-\phi_t}},$$

where V is the volume with respect to ω^n , and we can naturally think of $e^{-\phi_t}$ as defining volume forms on M . The critical points of this functional satisfy $e^{-\phi} = C\omega_\phi^n$, so they are Kähler-Einstein metrics.

The Ding functional has many analogous properties to the K-energy, such as the convexity along weak geodesics proved by Berndtsson [22], but it has technical advantages over the K-energy, since defining it requires less regularity of ϕ . Recently Donaldson [68] gave a variation of the infinite dimensional moment map picture discussed above, in which Berndtsson's convexity result [21] gives rise to the Kähler structure on \mathcal{J} , and the Ding functional corresponds to the log-norm functional. The weight again recovers the Futaki invariant, and the existence of a Kähler-Einstein metric is related to properness of \mathcal{D} by Tian [166].

3. K-STABILITY

In this section we survey the concept of K-stability of a Fano manifold M , or more generally a projective manifold M with an ample line bundle L , from different points of view. We first discuss the original notion for Fano manifolds, due to Tian [166], which is fairly differential geometric. A much more algebro-geometric definition for general pairs (M, L) was given by Donaldson [77]. In both of these definitions one needs to consider \mathbf{C}^* -equivariant degenerations of M , and the main difference is that in Tian's definition the central fiber is required to be a \mathbf{Q} -Fano

variety, whereas it can be an arbitrary scheme in Donaldson's definition. This added flexibility is needed when dealing with general polarized manifolds, but Li-Xu [118] showed that in the Fano case the two notions of K-stability are equivalent. We will now consider these two notions in more detail, along with a more analytic approach through geodesic rays in \mathcal{H} .

3.1. Tian's definition. The first notion of K-stability was introduced by Tian [166], in the context of Fano manifolds. As we have discussed in the previous section, Tian showed that if a Fano manifold M admits a Kähler-Einstein metric, then the K-energy on the Kähler class $c_1(M)$ is proper, and K-stability can be thought of as probing this properness along certain families of metrics.

Suppose that we have a \mathbf{C}^* -equivariant family of varieties $\pi : X \rightarrow \mathbf{C}$, with generic fiber $\pi^{-1}(t) \cong M$ for $t \neq 0$. Assume in addition that the central fiber is normal, and that a power of the relative anticanonical line bundle on the regular locus extends to a relatively ample line bundle on X . In this situation we call the family X a special degeneration of M .

The \mathbf{C}^* -action on such a special degeneration induces a \mathbf{C}^* -action on the central fiber $X_0 = \pi^{-1}(0)$. Using that X_0 has relatively mild singularities, Tian (see also Ding-Tian [65]) showed that one can define the Futaki invariant of this \mathbf{C}^* -action on X_0 using a differential geometric formula similar to (2.4). This is then defined to be the Futaki invariant $F(X)$ of the special degeneration X . Note that any \mathbf{C}^* -action on the Fano manifold M gives rise to a product action on the trivial family $X = M \times \mathbf{C}$, and the Futaki invariant of this family is simply the Futaki invariant of the original \mathbf{C}^* -action. At the same time there are infinitely many special degenerations, even if M admits no \mathbf{C}^* -actions.

In order to relate special degenerations to families of metrics and properness of the K-energy, note that any special degeneration $\pi : X \rightarrow \mathbf{C}$ for M can be realized as a family in projective space. More precisely there is an embedding $X \subset \mathbf{P}^N \times \mathbf{C}$, such that the \mathbf{C}^* -action on X is induced by the action of a one-parameter subgroup $\sigma : \mathbf{C}^* \rightarrow SL(N+1)$ on \mathbf{P}^N . Here M is embedded in $\mathbf{P}^N \times \{1\}$ using a basis of sections of $-rK_M$ for a suitable $r > 0$. We can now define a family of metrics $\omega_t \in c_1(M)$ by restricting the Fubini-Study metric to the non-zero fibers of X . Equivalently we can write

$$(3.1) \quad \omega_t = \frac{1}{r} \sigma_{e^{-t}}^* \omega_{FS}|_M.$$

Ding-Tian [65] showed that with suitable normalizing factors which we omit,

$$(3.2) \quad \lim_{t \rightarrow \infty} \frac{d}{dt} K(\omega_t) = F(X),$$

as suggested by (2.3) in the finite dimensional picture. Note, however, that the family ω_t is usually not a geodesic ray.

With these results in mind we have the following definition, due to Tian [166].

Definition 4. A Fano manifold M is K-stable, if $F(X) \geq 0$ for all special degenerations X of M , with equality only for product degenerations.

In the same paper, Tian showed that if M admits a Kähler-Einstein metric then the K-energy is proper, and as a consequence he showed the following fundamental result (see also Berman [16] for the case when M admits holomorphic vector fields).

Theorem 5. *If a Fano manifold admits a Kähler-Einstein metric, then it is K-stable.*

3.2. The Donaldson-Futaki invariant. The central fiber of a special degeneration has fairly mild singularities, and so a differential geometric definition of the Futaki invariant was possible. On the other hand Donaldson [77] assigned an invariant, called the Donaldson-Futaki invariant, to essentially arbitrary \mathbf{C}^* -equivariant degenerations of a polarized manifold (M, L) , by giving a purely algebro-geometric definition of the Futaki invariant for a \mathbf{C}^* -action on any polarized scheme (V, L) .

To give the definition, note that a \mathbf{C}^* -action λ on (V, L) induces actions on the spaces of sections $H^0(V, kL)$, and so in particular for each k we have a total weight w_k . For large k we have expansions

$$(3.3) \quad \begin{aligned} \dim H^0(V, kL) &= a_0 k^n + a_1 k^{n-1} + \dots \\ w_k &= b_0 k^{n+1} + b_1 k^n + \dots, \end{aligned}$$

and the Donaldson-Futaki invariant of λ is defined to be

$$DF(\lambda) = \frac{a_1}{a_0} b_0 - b_1.$$

When V is smooth, the equivariant Riemann-Roch formula can be used to show that this coincides with Futaki's differential geometric definition.

Such polarized schemes with \mathbf{C}^* -actions arise naturally as the central fibers of test-configurations.

Definition 6. Let (M, L) be a polarized manifold. A test-configuration for (M, L) with exponent r is a \mathbf{C}^* -equivariant flat family $\pi : (X, \mathcal{L}) \rightarrow \mathbf{C}$, such that \mathcal{L} is relatively ample, and

$$(\pi^{-1}(t), \mathcal{L}|_{\pi^{-1}(t)}) \cong (M, rL),$$

for any $t \neq 0$. In addition it is natural to require that the total space X is normal (see Li-Xu [118] and Ross-Thomas [143]).

The Donaldson-Futaki invariant $DF(X, \mathcal{L})$ of the test-configuration is defined to be the Donaldson-Futaki invariant of the induced \mathbf{C}^* -action on the central fiber.

Given this definition, K-stability can be defined as follows, in analogy with Definition 4.

Definition 7. A polarized manifold (M, L) is K-stable, if $DF(X, \mathcal{L}) \geq 0$ for all test-configurations for (M, L) , with equality only if $X \cong M \times \mathbf{C}$.

Using this definition, the following result was shown by Stoppa [155], building on work by Donaldson [78] and Arezzo-Pacard [6].

Theorem 8. *Suppose that M admits a constant scalar curvature metric in $c_1(L)$, and it has no nonzero holomorphic vector fields. Then (M, L) is K-stable.*

The result can be extended to the case when M has holomorphic vector fields, and also to extremal metrics (see [156], [19]). As we stated in the introduction, the Yau-Tian-Donaldson conjecture is the converse of this result, saying that if (M, L) is K-stable, then there is a constant scalar curvature metric in $c_1(L)$. However it is likely that actually a stronger notion of stability is needed in general, in view of examples of Apostolov-Calderbank-Gauduchon-Tønnesen-Friedman [5], that are shown to be unstable in a suitable sense by Dervan [60]. One possibility for such a

stronger stability notion is provided by the formalism of filtrations [160, 176], while another is the concept of uniform K-stability [62, 25].

In order to compare Definitions 4 and 7, let us point out that it is fairly easy to construct interesting test-configurations, which are not special degenerations, using for instance deformation to the normal cone. This was explored in detail by Ross-Thomas [143, 142]. One can simply take any subscheme $Z \subset M$, and let $X = \text{Bl}_{Z \times \{0\}} M \times \mathbf{C}$, with a suitable relatively ample line bundle \mathcal{L} . For instance when Z is a smooth submanifold of M , then the central fiber of X will be isomorphic to $\text{Bl}_Z M \cup P_Z$, where $P_Z = \mathbf{P}(N_Z \oplus \underline{\mathbf{C}})$ is the projective completion of the normal bundle of Z in M , and P_Z is glued along its zero section to the blowup $\text{Bl}_Z M$ along its exceptional divisor. In fact Odaka [130] showed that by blowing up “flag ideals” of $M \times \mathbf{C}$ instead of just subschemes, one can essentially recover all test-configurations, and using this approach Odaka-Sano [133] and Dervan [61] were able to prove the K-stability of certain varieties.

With this in mind, it appears that in the Fano case Definition 7 is more restrictive than Tian’s Definition 4, since test-configurations are much more general than special degenerations. It is quite remarkable then that for Fano manifolds the two notions turn out to be equivalent. This was first proven by Li-Xu [118] purely algebro-geometrically, using the minimal model program. Roughly speaking the minimal model program allowed them to modify an arbitrary test-configuration into a special degeneration, while controlling the sign of the Donaldson-Futaki invariant at each step. A more differential geometric proof also follows from Chen-Donaldson-Sun’s proof [41] of the YTD-conjecture for Fano manifolds.

One suggestive example is to consider a polarized toric manifold (M, L) , with Delzant polytope P . It is natural in this case to only allow torus equivariant test-configurations. The only torus equivariant test-configurations with normal central fiber are product configurations induced by a \mathbf{C}^* -action on M and indeed, when M is Fano, then Wang-Zhu [175] showed that the only obstruction to the existence of a Kähler-Einstein metric is that given by the Futaki invariants of these \mathbf{C}^* -actions. On the other hand, as shown by Donaldson [77], any rational piecewise linear convex function on P gives rise to a test-configuration for (M, L) and there are (non-Fano) examples where these give an obstruction to the existence of a cscK metric, not detected by product configurations.

3.3. Intersection theoretic formula. An alternative formula for the Donaldson-Futaki invariant in terms of intersection products has been very useful in more algebro-geometric developments. It was observed by Wang [174], and it is also related to the CM-polarization of Tian [166, 136]. To explain it, note that any test-configuration (X, \mathcal{L}) can be extended trivially at infinity to obtain a \mathbf{C}^* -equivariant family

$$(\overline{X}, \overline{\mathcal{L}}) \rightarrow \mathbf{P}^1.$$

The line bundle $\overline{\mathcal{L}}$ is relatively ample, and by taking the tensor product with a line bundle pulled back from \mathbf{P}^1 we can assume that it is actually ample. A calculation shows that in terms of this family the Donaldson-Futaki invariant of a test-configuration of exponent r is

$$(3.4) \quad DF(X, \mathcal{L}) = \frac{n}{n+1} \mu(M, rL) (\overline{\mathcal{L}})^{n+1} + \overline{\mathcal{L}}^n \cdot K_{\overline{X}/\mathbf{P}^1},$$

using the intersection product on \overline{X} . Here for a polarized variety (M, L) , the “slope” $\mu(M, L)$ is defined by

$$\mu(M, L) = \frac{-K_M \cdot L^{n-1}}{L^n},$$

and up to a constant multiple is it the average scalar curvature \overline{S} of a Kähler metric in $c_1(L)$.

This reformulation of the Donaldson-Futaki invariant has various advantages, as shown for instance in the works of Li-Xu [118], Odaka [131] and others. Here we just mention one, namely the extension of K-stability to non-algebraic Kähler manifolds due to Dervan-Ross [63] and Sjöström Dyrefelt [150]. While the expansions (3.3) do not make sense in the absence of a line bundle, one can make sense of the intersection product (3.4) even in the Kähler case.

3.4. Geodesic rays. In Section 2 we described how in the finite dimensional moment map picture, stability can be tested using geodesic rays in the symmetric space G^c/G . Donaldson [73] formulated conjectures saying that in an analogous way geodesic rays in \mathcal{H} can be used to detect the existence of a constant scalar curvature metric. Since that time there has been enormous progress on our understanding of geodesic rays, although these conjectures are still mostly open except in the Fano case.

We have seen in (3.1) that a special degeneration or a test-configuration for M gives rise to a path in the space \mathcal{H} of Kähler potentials. Unless we have a product test-configuration, this path cannot be expected to be a geodesic ray in \mathcal{H} , but rather it is a geodesic in a finite dimensional space of *Bergman metrics*, i.e. those obtained by restricting the Fubini-Study metric. It turns out that the relation (3.2) between the Futaki invariant of a test-configuration and the asymptotic derivative of the K-energy along the corresponding Bergman geodesic does not hold for general test-configurations. The general formula for the limit has been obtained by Paul [135] in terms of hyperdiscriminant and Chow polytopes, leading to an alternative notion of stability.

To relate this to geodesic rays in \mathcal{H} , note that a given test-configuration X for M can be realized as a family in projective spaces of arbitrarily large dimension, and in this way we obtain not one, but a whole sequence of Bergman geodesics of metrics on M from X using the formula (3.1). Phong-Sturm [140] showed that one can pass to a limit, and obtain a geodesic ray in \mathcal{H} in a suitable weak sense, with an arbitrary initial point ϕ_0 .

One can also directly construct such a weak geodesic ray in \mathcal{H} from the test-configuration X in the following way (see [7], [48], [141], [16] for this in various degrees of generality). Let us denote by X_Δ the family X restricted to the unit disk $\Delta \subset \mathbb{C}$. We use the “initial point” ϕ_0 to define a metric $e^{-\phi_0}$ on the line bundle \mathcal{L} over ∂X_Δ . The geodesic ray is then obtained by finding an S^1 -invariant metric $e^{-\phi}$ on \mathcal{L} over X_Δ , with positive curvature current, and solving the homogeneous complex Monge-Ampère equation $(\sqrt{-1}\partial\bar{\partial}\phi)^{n+1} = 0$ on the interior of X_Δ , in the sense of pluripotential theory. The existence of such a solution, and its regularity properties are discussed by Phong-Sturm [141]. Over the punctured disk Δ^* the family X is biholomorphic to $M \times \Delta^*$, and so the metric $e^{-\phi}$ on \mathcal{L} induces a family of metrics on $L \rightarrow M$, which is the geodesic ray we were after.

There has been a lot of work relating the behavior of the K-energy along such a geodesic ray to the Donaldson-Futaki invariant of the test-configuration (see e.g. [48], [141]). The first sharp result in this direction is due to Berman [16], using the Ding functional instead of the K-energy, in the case when M is Fano. He shows that along a geodesic ray ϕ_t constructed from a test-configuration (X, \mathcal{L}) as above, we have

$$(3.5) \quad DF(X, \mathcal{L}) = \lim_{t \rightarrow \infty} \frac{d}{dt} \mathcal{D}(\phi_t) + q,$$

where $q \geq 0$ is a rational number determined by the central fiber of the test-configuration. An analogous formula for the K-energy, for test-configurations with smooth total space, was subsequently obtained by Sjöström Dyrefelt [150]. The relation (3.5) led Fujita [92] to study the notion of Ding stability, where the Donaldson-Futaki invariant is replaced by the asymptotic derivative of the Ding functional. As an important application, he showed that projective space has the maximal volume amongst Kähler-Einstein manifolds of a fixed dimension, using that Kähler-Einstein manifolds are stable in this sense. This direction has been taken much further in recent work of Fujita [91], Li [111], Li-Xu [116, 117], Liu-Xu [119], leading to new examples of K-stable varieties, as we mention in Section 5.1.

A different direction was pursued by Boucksom-Hisamoto-Jonsson [25], who introduced the point of view of thinking of a test-configuration for (M, L) as a non-Archimedean metric on L . From this point of view the asymptotic derivative of the Ding functional can be seen as a non-Archimedean Ding functional. Building on these ideas, and using techniques from non-Archimedean analysis [24], as well as advances on the geometry of the space of metrics \mathcal{H} due to Darvas [54], Berman-Boucksom-Jonsson [18] proposed a variational approach to proving Theorem 2. This approach relies far less on differential geometric methods than Chen-Donaldson-Sun's approach [41], and as such it may eventually lead to a solution of the Yau-Tian-Donaldson conjecture for singular Fano varieties. At present, however, even in the smooth case it yields a weaker result than [41] since it requires assuming uniform K-stability.

3.5. Finite dimensional approximation. In the previous section we mentioned the idea of realizing a geodesic in \mathcal{H} as a limit of finite dimensional Bergman geodesics. This idea of approximating the geometry of \mathcal{H} with that of larger and larger spaces of Bergman metrics is a fundamental one, going back at least to work of Tian [164] on the problem of approximating an arbitrary Kähler metric $\omega \in c_1(L)$ on M with a sequence of metrics

$$\omega_k = \frac{1}{k} \phi_k^* \omega_{FS}.$$

Here $\phi_k : M \rightarrow \mathbf{P}^{N_k}$ are projective embeddings given by bases of sections of kL , for sufficiently large k . Fixing a Hermitian metric on L with curvature form ω , we have induced L^2 inner products on each space of sections $H^0(kL)$. It is then natural to use an orthonormal basis $\{s_0^{(k)}, \dots, s_{N_k}^{(k)}\}$ of $H^0(kL)$ to define the embedding ϕ_k . A calculation shows that

$$(3.6) \quad \frac{1}{k} \phi_k^* \omega_{FS} = \omega + \sqrt{-1} \partial \bar{\partial} \log \rho_{\omega, k},$$

where the function

$$\rho_{\omega,k} = \sum_{i=0}^{N_k} |s_i^{(k)}|^2$$

is called the Bergman kernel. Tian gave an asymptotic expansion of this function as $k \rightarrow \infty$, which was later refined by Zelditch [179], Lu [120], Catlin [32] and others, showing that

$$(3.7) \quad \rho_k = 1 + \frac{S(\omega)}{2} k^{-1} + O(k^{-2}), \text{ as } k \rightarrow \infty.$$

The lower order terms can also be computed, and are all given in terms of covariant derivatives of the curvature tensor of ω (see Lu [120]). Note that normalising constants in the coefficients depend on conventions for defining the L^2 product on $H^0(kL)$. In addition, integrating the expansion over M recovers the Hirzebruch-Riemann-Roch formula. Using this expansion in (3.6) we see that $\omega_k - \omega = O(k^{-2})$, and so indeed we can approximate an arbitrary metric ω with Bergman metrics.

Donaldson [75, 79] took this point of view further, realizing that it is advantageous to approximate the entire moment map package described in Section 2 by a sequence of analogous finite dimensional problems, a process that can be thought of as quantization [74]. A starting point is the work of Zhang [180] and Luo [122], relating the GIT stability of the Chow point of a projective submanifold $V \subset \mathbf{P}^N$ under the action of $SL(N+1)$ to the existence of a special embedding of V . Letting n denote the dimension of V , we define the matrix

$$M(V)_{jk} = \sqrt{-1} \int_V \frac{Z_j \bar{Z}_k}{\sum_{i=0}^N |Z_i|^2} \omega_{FS}^n,$$

where the Z_i are the homogeneous coordinates on \mathbf{P}^N . Luo shows that GIT stability is related to the existence of *balanced embeddings*, for which the matrix $M(V)$ is a multiple of the identity matrix. In fact $M(V)$, or rather its projection to $\mathfrak{su}(N+1)$ can be viewed as a moment map, and zeroes of this moment map are the balanced embeddings.

Given any embedding $V \subset \mathbf{P}^N$, we can consider the restriction of the Fubini-Study metric on V , and then construct a new embedding using an L^2 orthonormal basis of sections of $O(1)|_V$ as above. It turns out that $V \subset \mathbf{P}^N$ is a balanced embedding if this new embedding coincides with the old one (up to the action of $SU(N+1)$), i.e. if the Bergman kernel is constant. The expansion (3.7) then suggests that for a given pair (M, L) , balanced embeddings using a basis of $H^0(kL)$ for very large k ought to be related to constant scalar curvature metrics in $c_1(L)$. Indeed, Donaldson [75] shows that if a cscK metric exists, and M admits no holomorphic vector fields, then M also admits balanced embeddings using a basis of $H^0(kL)$ for sufficiently large k . Since such balanced embeddings are unique up to the action of $SU(N+1)$, this implies in particular that if a cscK metric exists, it is unique. Moreover M satisfies an asymptotic GIT stability condition, which implies K-semistability.

By “quantizing” the K-energy as well, Donaldson [79] showed that if a cscK metric exists, and M has no holomorphic vector fields, then the K-energy is bounded below on \mathcal{H} , generalizing Bando-Mabuchi’s result [13] in the Kähler-Einstein case. Going further, using such finite dimensional approximations, Donaldson [78] showed that a destabilizing test-configuration leads to a nontrivial lower bound for the

Calabi functional: for any metric $\omega \in c_1(L)$, and any test-configuration (X, \mathcal{L}) for (M, L) we have

$$\|S(\omega) - \overline{S}\|_{L^2} \geq \frac{-DF(X, \mathcal{L})}{\|(X, \mathcal{L})\|_2},$$

where a suitable L^2 -type norm on test-configurations is used for normalizing the Donaldson-Futaki invariant. This result shows, in particular, that the existence of a cscK metric implies K-semistability, even when M has holomorphic vector fields.

These ideas have since been developed further in many papers. A very small sample of such developments is [125, 81, 151, 88, 145, 112, 89, 45, 146, 148].

4. GEOMETRIC AND ALGEBRAIC LIMITS

Many of the ideas that we have discussed so far apply not only to the problem of Kähler-Einstein metrics, but also to more general constant scalar curvature metrics. In this section this will no longer be the case, since it will be crucial that the Einstein equation controls the Ricci curvature of the metric. Our goal is to present some of the ideas involved in Chen-Donaldson-Sun's proof of Theorem 2 [41], that is the existence part of the Yau-Tian-Donaldson Conjecture in the Fano case.

Suppose that we have a Fano manifold M , and we are trying to find a metric $\omega \in c_1(M)$ satisfying $\text{Ric}(\omega) = \omega$. There are several natural "continuity methods" for deforming a given initial metric $\omega_0 \in c_1(M)$ to a Kähler-Einstein metric. One approach is to choose a reference metric $\alpha \in c_1(M)$, and then try to find a family of metrics ω_t for $t \in [0, 1]$ satisfying

$$(4.1) \quad \text{Ric}(\omega_t) = t\omega_t + (1-t)\alpha.$$

Using Yau's theorem [177] we can solve this equation for $t = 0$, and using the implicit function theorem Aubin [10] showed that the set of t for which a solution exists is open. The remaining difficulty is then to understand what can happen to a sequence of solutions ω_{t_k} as $t_k \rightarrow T$, for some $T \in (0, 1]$. This approach was used in [56], based on the techniques of Chen-Donaldson-Sun.

Chen-Donaldson-Sun [41] used a variant of this, proposed by Donaldson [80], where the ω_t are Kähler-Einstein metrics on the complement $M \setminus D$ of a suitable smooth divisor $D \subset M$, while along D the ω_t have singularities modeled on a two-dimensional cone with cone angle $2\pi t$. In analogy with (4.1), the metrics ω_t satisfy

$$(4.2) \quad \text{Ric}(\omega_t) = t\omega_t + (1-t)[D],$$

where $[D]$ is the current of integration along D . Here the Ricci curvature is defined as the curvature current of a metric on $-K_M$ induced by the volume form of ω_t . The advantage of this continuity path over (4.1) is that on $M \setminus D$ the metric ω_t is Einstein, at the expense of introducing singularities along D . At the same time in algebraic geometry it is natural to consider pairs (M, D) , and metrics with cone singularities along D are useful differential geometric counterparts (see e.g. [30]). The relevant linear theory for the implicit function theorem was developed by Donaldson [82], with further refinements by Jeffres-Mazzeo-Rubinstein [102], and as in the case of (4.1) one needs to understand the limiting behavior of a sequence ω_{t_k} as $t_k \rightarrow T$.

A somewhat different strategy is given by the Ricci flow, originally introduced by Hamilton [98]. On a Fano manifold M the (normalized) Kähler-Ricci flow is

given by the parabolic equation

$$\frac{\partial}{\partial t}\omega_t = \omega_t - \text{Ric}(\omega_t),$$

with initial metric $\omega_0 \in c_1(M)$. It was known for a long time that the flow exists for all time [31], and the main difficulty was understanding the behavior of ω_t as $t \rightarrow \infty$. Building on Perelman's deep results on the Ricci flow [137] as well as on ideas in Chen-Donaldson-Sun's proof, Chen-Wang [49] developed the necessary convergence theory for understanding this limiting behavior (see also Bamler [11]). This was then used by Chen-Sun-Wang [46] to give a proof of Theorem 2 using the Ricci flow.

We now return to the continuity method (4.1), and a sequence ω_{t_k} with $t_k \rightarrow T$. If such a sequence (or a subsequence) converges to a solution ω_T of (4.1), then either we can further increase t using the implicit function theorem, or $T = 1$ and we obtain the Kähler-Einstein metric that we seek. The key question is then to understand what happens when the sequence ω_{t_k} diverges. In this case we need to construct a special degeneration for M with non-positive Futaki invariant.

In broad outline the strategy is the following. We first define a certain limit space Z out of the sequence (M, ω_{t_k}) (up to choosing a subsequence), and show that this limit can be given the structure of an algebraic variety, and in addition Z can also be viewed as an algebro-geometric limit of the images of a sequence of projective embeddings $\phi_k : M \rightarrow \mathbf{P}^N$. For simplicity suppose that $T = 1$. Then the limit Z is shown to admit a possibly singular Kähler-Einstein metric. If Z is biholomorphic to M , then we obtain the Kähler-Einstein metric that we were after, but if not, then there is a special degeneration for M with central fiber Z . This necessarily has vanishing Futaki invariant, and so M is not K-stable. The argument is a little more involved when $T < 1$, since then the limit Z admits a "twisted" Kähler-Einstein metric, and we obtain a special degeneration with a strictly negative Futaki invariant.

In the next two sections we give some more details on the construction of the limit Z , and on relating the geometric and algebraic limits. It will be helpful to consider a more general sequence (M_k, ω_k) than that obtained from the equations (4.1), even allowing the manifolds M_k to vary. Some aspects of the problem, such as the convergence theory, are easier when we consider a sequence of Kähler-Einstein manifolds, however others, such as using the K-stability assumption, are easier along a continuity method such as (4.1) (see for instance Donaldson [84] for the difficulty in using a sequence of Kähler-Einstein metrics).

4.1. Gromov-Hausdorff limits. In this section we will consider a sequence (M_k, ω_k) that are either all Kähler-Einstein with volumes controlled from below, or they arise from (4.1) or another similar continuity method. From a geometric point of view the advantage of working with such metrics as opposed to those with just constant scalar curvature is that control of the Ricci curvature allows us to extract a Gromov-Hausdorff limit of the sequence (M_k, ω_k) . This is a consequence of the Gromov compactness theorem [96] and the Bishop-Gromov volume comparison theorem [37]. For simplicity let us denote by M_k the metric space (M_k, ω_k) . Up to choosing a subsequence of the M_k , the limit is a metric space (Z, d) , such that we have metrics (distance functions) d_k on the disjoint unions $M_k \sqcup Z$, extending those

on M_k and Z , satisfying the following: for all $\epsilon > 0$, the ϵ -neighborhoods of both M_k and Z cover all of $M_k \sqcup Z$ for sufficiently large k .

While at first (Z, d) is just a metric space, a series of works by Cheeger-Colding [34, 35, 36] developed a detailed structure theory (see also Cheeger [33] for an exposition). The general theory applies to Riemannian manifolds with only a lower bound on the Ricci curvature, and also allows collapsing phenomena, but our sequence M_k satisfies the non-collapsing condition

$$\text{Vol}(B(q_k, 1)) > c > 0,$$

for a constant $c > 0$, where $q_k \in M_k$ are some basepoints. This again uses the Bishop-Gromov inequality, together with Myers's Theorem and the fact that we control the volumes of the M_k .

The basic concept is that of a tangent cone. For any $p \in Z$, and a sequence $r_k \rightarrow \infty$ the Gromov compactness theorem can be applied to the sequence of pointed manifolds $(Z, r_k d, p)$ and, up to choosing a subsequence, we obtain a limit metric space Z_p , called a tangent cone of Z at p . A fundamental result of Cheeger-Colding is that in our situation these tangent cones are metric cones. More precisely for each tangent cone Z_p there is a length space Y of diameter at most π , such that Z_p is the completion of $Y \times (0, \infty)$ using the metric

$$d((y_1, r_1), (y_2, r_2)) = r_1^2 + r_2^2 - 2r_1 r_2 \cos d_Y(y_1, y_2).$$

We write this space as $C(Y)$.

A point $p \in Z$ is called regular, if a tangent cone Z_p is isometric to \mathbf{R}^m (in our situation $m = 2n$). In fact in this case each tangent cone at p is Euclidean. This means that if we take a sufficiently small ball centered at p , and scale it up to unit size, then it will be very close to the Euclidean ball in the Gromov-Hausdorff sense. But recall that for sufficiently large k , the metric space M_k is very close to Z . This implies that we can find points $p_k \in M_k$, and small balls centered at p_k which, scaled to unit size, are close to Euclidean in the Gromov-Hausdorff sense. If the metrics ω_k were Einstein, then results of Anderson [4] and Colding [51] would show that we can choose holomorphic coordinates centered at p_k with respect to which the components of ω_k are controlled in $C^{2,\alpha}$. If all points of Z were regular, this would mean that there is a uniform scale at which we have $C^{2,\alpha}$ control of the metrics ω_k . In particular we could take a limit of the Kähler structures (M, ω_k) , and turn Z into a smooth Kähler manifold. Note, however, that even in this case Z may not be biholomorphic to M .

When the ω_k are not actually Einstein, but rather arise from (4.1), then Anderson's result still applies at the regular points [161]. However, in general not all points of Z are regular, and instead Cheeger-Colding defined a stratification based on how close each tangent cone is to being Euclidean. More precisely, for $0 \leq k \leq m-1$, the set S_k is defined to be those points $p \in Z$ for which no tangent cone Z_p splits off an isometric factor of \mathbf{R}^{k+1} . They then showed that the Hausdorff dimension of S_k is at most k and in addition $S_{m-1} = S_{m-2}$, so Z is regular outside a codimension 2 set. A key technical step in Chen-Donaldson-Sun's work is to improve this dimension estimate of the singular set from the Hausdorff dimension to the Minkowski dimension - i.e. to show that there is a constant K for which the singular set S_{m-1} can be covered by Kr^{2-m} balls of radius r , for any $r \in (0, 1)$. For even better estimates on the Minkowski dimension of the singular set for Einstein manifolds, see Cheeger-Naber [38, 39].

This discussion applies also to the tangent cones Z_p , as well as to further iterated tangent cones. In our situation this implies that each tangent cone Z_p is a Ricci flat Kähler cone, at least outside of a codimension 2 set. In addition on the regular set in Z_p , the Ricci flat Kähler metric is given by $\sqrt{-1}\partial\bar{\partial}r^2$, where r is the distance from the vertex of Z_p .

4.2. The partial C^0 -estimate. The theory described in the previous section gives a good understanding of the “geometric” limit of the spaces (M_k, ω_k) . In this section we incorporate the holomorphic structure through the partial C^0 -estimate, which is an idea due to Tian [165] used in his work on Kähler-Einstein metrics on Fano surfaces. For simplicity we will assume here that $M_k = M$, and the sequence ω_k is obtained from (4.1).

Suppose that $-lK_M$ is very ample for some large $l > 0$. The metrics ω_k define Hermitian metrics h_k on $-lK_M$, with curvature forms $l\omega_k$. As in Section 3.5, we have a natural projective embedding $\phi_k : M \rightarrow \mathbf{P}^N$ obtained by using an orthonormal basis $\{s_i\}$ of $H^0(M, -lK_M)$ for the L^2 inner product

$$\langle s, t \rangle_{L^2} = \int_M \langle s, t \rangle_{h_k} (l\omega_k)^n,$$

defined by h_k . As before, the metric ω_k is related to the pullback of the Fubini-Study metric under ϕ_k by the Bergman kernel $\rho_{\omega_k, l}$.

In Section 3.5 we saw the expansion (3.7) of $\rho_{\omega_k, l}$ as $l \rightarrow \infty$ for a fixed metric ω_k , whereas here the main interest is in obtaining bounds for $\rho_{\omega_k, l}$ for sufficiently large l , which apply uniformly to a sequence of metrics ω_k . Tian [165] conjectured that if, as in our situation, we have a positive lower bound for the Ricci curvatures of ω_k , and a lower bound for the volumes, then for sufficiently large l we have a bound $\rho_{\omega_k, l} > d > 0$ from below. This is called the partial C^0 -estimate, and it is equivalent to showing that for any point $p \in M$, the bundle $-lK_M$ has a holomorphic section s satisfying $\|s\|_{L^2} = 1$, and $|s|_{h_k}^2(p) > d$. Note that if $-lK_M$ is very ample, then $\rho_{\omega, l}$ is positive for any metric $\omega \in c_1(M)$, and the key point is to have a uniform lower bound along the sequence of metrics ω_k .

For the details on how to use the partial C^0 -estimate to define the structure of an algebraic variety on Z , see Tian [165] and more generally Donaldson-Sun [67]. Here let us just note that a key consequence of the estimate is that there is a uniform bound on the derivatives of the embeddings $\phi_k : (M, \omega_k) \rightarrow \mathbf{P}^N$, independent of k . Up to further increasing the multiple of $-K_M$ that we use, this eventually leads to the result that the Gromov-Hausdorff limit Z is homeomorphic to the algebro-geometric limit of the images $\phi_k(M)$ in projective space.

Let us say a few words on the proof of the partial C^0 -estimate. Fixing a point $p \in M$, for each metric ω_k we must construct holomorphic sections s_k of $-lK_M$, for sufficiently large l , such that $\|s_k\|_{L^2, h_k} = 1$, and $|s_k|_{h_k}^2(p) > d > 0$ for some d independent of k . The simplest situation is when the geometry of each ω_k is very well controlled near p at a suitable scale. For instance this would be the case if ω_k did not actually depend on k (or more generally if the ω_k were to converge to a smooth metric ω on M). In this case, at a sufficiently small scale, the metrics ω_k appear to be very close to the Euclidean space $(\mathbf{C}^n, \omega_{Euc})$. The trivial line bundle over \mathbf{C}^n has metric $e^{-l|z|^2}$ with curvature $l\omega_{Euc}$, and so for sufficiently large l , there will be a holomorphic section s with $|s|^2(0) = 1$ and $\|s\|_{L^2} = 1$ but with s decaying rapidly away from the origin. Using cutoff functions this section can be glued onto

M to produce a smooth section s_k of $-lK_M$ satisfying

$$|s_k|_{h_k}^2(p) \sim 1, \quad \|s_k\|_{L^2, h_k}^2 \sim 1, \quad \text{and} \quad \|\bar{\partial}s_k\|_{L^2, h_k} \sim 0.$$

Using the Hörmander L^2 -technique, these sections can be perturbed slightly to obtain the required holomorphic sections of $-lK_M$. This method was used by Tian [164] to find a weak form of the asymptotic expansion (3.7) of the Bergman kernel as $l \rightarrow \infty$, for a fixed metric ω . As we have described in Section 3.5, this expansion has been very influential in the study of constant scalar curvature Kähler metrics.

In general we do not have such good uniform control of the local geometry of the metrics ω_k . The first more general situation that was understood was the case of Fano surfaces, by Tian [165]. He proved an orbifold compactness theorem for Kähler-Einstein surfaces, using techniques employed in the study of Yang-Mills connections by Uhlenbeck [172, 171] (see also Anderson [3] and Bando-Kasue-Nakajima [12]). In effect this implies that at a suitable scale, the local geometry of a non-collapsed sequence of Kähler-Einstein surfaces (M_k, ω_k) is modeled either on flat \mathbf{C}^2 , or its quotient by a finite group. Using this together with the L^2 technique, Tian showed the partial C^0 -estimate for such a sequence of Kähler-Einstein surfaces.

There was little further progress until the work of Donaldson-Sun [67], who proved the partial C^0 -estimate for non-collapsing sequences of Kähler-Einstein manifolds (M_k, ω_k) , generalizing the result of Tian. In the proof they combined the L^2 technique with the Cheeger-Colding structure theory that we described above. Very roughly the structure theory, as described above, implies that the local geometry of the (M_k, ω_k) near a point p_k is modeled by Ricci flat cones $C(Y)$. Although Y may have singularities, the estimate on the size of the singular set ensures that we can find a cutoff function η , supported on the regular part of $C(Y)$, for which $\|\nabla\eta\|_{L^2} \ll 1$. Taking l sufficiently large, the line bundle $-lK_{M_k}$ with its Hermitian metric h_k is modeled, at least on the regular part, by the trivial bundle over $C(Y)$, with trivializing section s whose norm decays exponentially fast. One can then use ηs to obtain a smooth section of $-lK_{M_k}$ which is approximately holomorphic in an L^2 -sense, and which can then be perturbed to a genuine holomorphic section s_k . A gradient estimate is then used to show that $|s_k|^2(p_k) > d$ for a controlled constant $d > 0$, while $\|s_k\|_{L^2} \sim 1$, as required.

Shortly after Donaldson-Sun's work, the partial C^0 -estimate was proven in various other settings. In their proof of the Yau-Tian-Donaldson conjecture, Chen-Donaldson-Sun [43, 44] proved it for Kähler-Einstein metrics with conical singularities, while Phong-Song-Sturm [138] extended Donaldson-Sun's work to Kähler-Ricci solitons using also some techniques of Tian-Zhang [168]. In [161] we showed that it holds along the continuity method (4.1). The ideas have also been applied to the Kähler-Ricci flow, by Tian-Zhang [169] in dimensions up to 3, and Chen-Wang [49] in general. These results on the Kähler-Ricci flow also led to a proof of Tian's general conjecture on the partial C^0 -estimate, for a non-collapsed sequence of Kähler manifolds with only a positive lower bound on the Ricci curvature (see Jiang [103] for dimensions up to 3, and Chen-Wang [49] in general).

4.3. The K-stability condition. In the previous sections we have described how out of a sequence (M, ω_k) along the continuity method (4.1), or from a sequence

of Kähler-Einstein metrics, or even from a sequence along a solution of the Kähler-Ricci flow one can construct a limit Z , which has the structure of an algebraic variety. This limit variety can be singular, although it is always normal with log terminal singularities. In addition one can show that Z admits a singular Kähler metric, which satisfies either an analog of the equation (4.1) with $t = T$ (where recall that $T = \lim_{k \rightarrow \infty} t_k$), is Kähler-Einstein, or is a Kähler-Ricci soliton in the case of the Ricci flow. In this section we will explain how the K-stability assumption on M can be used to show that in the limit we obtain a Kähler-Einstein metric on M . In addition we discuss some of the developments regarding singular Kähler-Einstein metrics, as they play an important role in the proof of the Yau-Tian-Donaldson conjecture.

Let us suppose that we are working with a continuity method such as (4.1), we have $T = 1$, and for simplicity assume that Z is smooth. Note that Z then admits a Kähler-Einstein metric, and the goal is to show that if M is K-stable, then actually $Z \cong M$. To see this, note first that Matsushima's Theorem [126] implies that the automorphism group of Z is reductive. On the other hand, Z can also be viewed as an algebraic limit of the projective varieties $\phi_k(M) \subset \mathbf{P}^N$, for a sequence of embeddings $\phi_k : M \rightarrow \mathbf{P}^N$, and so in a suitable Hilbert scheme, Z can be viewed as an element in the closure of the $GL(N+1)$ -orbit of M . From this point of view it is the stabilizer of Z under the $GL(N+1)$ action that is reductive, and then an appropriate version of the Luna slice theorem [121, 83] implies that there is a special degeneration for M , whose central fiber is Z . Futaki's result [93] furthermore implies that the Futaki invariant of this special degeneration must vanish. If M is K-stable, then necessarily $Z \cong M$, and so M admits a Kähler-Einstein metric.

Suppose now that $T < 1$, and that for simplicity Z is smooth. Under Chen-Donaldson-Sun's continuity method (4.2) one obtains a limiting Kähler-Einstein metric on Z , with cone singularities along a divisor D_∞ . The strategy is then to apply the argument used above, but to the pairs (M, D) and (Z, D_∞) . More precisely, one shows, using the equation on Z , that in a suitable Hilbert scheme of pairs the stabilizer of (Z, D_∞) is reductive, and from this one constructs a test-configuration X for (M, D) with central fiber (Z, D_∞) . The Donaldson-Futaki invariant has a natural generalization to pairs [82], called the log Futaki invariant $DF(X, (1-T)D)$, depending on the divisor D as well as the "angle" parameter T such that $DF(X, 0D) = DF(X)$. It is shown in [44] that $DF(X, (1-T)D) = 0$, since in fact the log Futaki invariant of any vector field on the pair $(Z, (1-T)D_\infty)$ vanishes. This uses the existence of a conical Kähler-Einstein metric on Z . In addition $DF(X, D) > 0$, which corresponds to an existence result for Kähler-Einstein metrics with cone singularities that have small cone angles. Since the log Futaki invariant is linear in the angle parameter, it follows that $DF(X) < 0$.

Using the continuity method (4.1) is a bit more cumbersome, because in that case instead of working with pairs of the form (M, D) that can be parametrized by a Hilbert scheme, we need to work with pairs (M, α) , where α is a $(1,1)$ -form, or even a current, and the Luna slice theorem does not apply directly in this infinite dimensional setting. In [56] we overcome this problem by an approximation argument, which we now outline.

The starting point is to show that out of a sequence of solutions to (4.1) with $t_i \rightarrow T$, we obtain a limiting metric ω_T on the space Z satisfying the equation

$$\text{Ric}(\omega_T) = T\omega_T + (1-T)\beta,$$

where β is a positive $(1,1)$ -current on Z . This can be used to show that the automorphism group of the pair (Z, β) , suitably defined, is reductive, and moreover a version of the log Futaki invariant of the pair $(Z, (1-T)\beta)$ vanishes for all vector fields. The missing piece for exploiting this as above is to find a test-configuration for M with central fiber Z , under which the form α on M converges to the current β on Z , but in the end we are not able to do this. Instead, in order to apply a result such as Luna's slice theorem, the idea is to approximate the currents α and β with currents of integration along divisors.

Note first that according to Shiffman-Zelditch [149], we can write α as an average of currents of integration $[M \cap H]$ for hyperplane sections of M , under a suitable projective embedding $M \subset \mathbf{P}^N$. As discussed above, the geometric limit space Z is also an algebraic limit $\lim_{k \rightarrow \infty} \rho_k(M)$ for automorphisms ρ_k of \mathbf{P}^N . We can arrange that the limits $\rho_\infty(H) = \lim_{k \rightarrow \infty} \rho_k(H)$ exist for all hyperplanes H and so β is an average of the hyperplane sections $[Z \cap \rho_\infty(H)]$ of Z . This can be used to show that a suitable tuple $(Z, \rho_\infty(H_1), \dots, \rho_\infty(H_l))$ has the same automorphism group as (Z, β) . One simple example to keep in mind is when Z is a line in \mathbf{P}^2 , and β is the Fubini-Study metric. In this case the automorphism group of (Z, β) is trivial under our definition, and so we must "mark" Z with at least 3 hyperplane sections to have a tuple with the same automorphism group.

We can argue as before with these tuples, and we obtain a test-configuration X for M with central fiber Z , such that in addition the limits of the hyperplanes H_1, \dots, H_l under this test-configuration are $\rho_\infty(H_1), \dots, \rho_\infty(H_l)$. However, since we do not know that the form α converges to β under this test-configuration, we cannot say anything about the Donaldson-Futaki invariant of X . To overcome this we choose enough hyperplane sections so that by an approximation argument the log Futaki invariant of the tuple is close to that of the pair (Z, β) . It remains then to use this to show that $DF(X) < 0$, contradicting K-stability of M , unless we actually have $Z \cong M$ and $\beta = \alpha$, so that the continuity method can be continued past $t = T$.

To conclude this section we make some remarks about the situation when the limit space Z is not smooth. The overall strategy remains the same, but there are substantial technical difficulties in carrying it through. To start with, one needs to make sense of the Kähler-Einstein equation (or more general equations such as (4.1)) on a singular variety Z . Working on a resolution of singularities $\pi : X \rightarrow Z$, the equation is equivalent to a complex Monge-Ampère equation

$$(\omega + \sqrt{-1}\partial\bar{\partial}\phi)^n = e^{F-\phi}dV,$$

on X , where dV is a smooth volume form, however the function F need not be bounded, and the form ω is only non-negative. More precisely we have $0 \leq e^F \in L^p$ for some $p > 1$, and ω is also big, i.e. $\int_X \omega^n > 0$. It cannot be expected that ϕ is twice differentiable, and so the equation needs to be defined in a weak sense. This was done for the local theory of the complex Monge-Ampère operator by Bedford-Taylor [14]. Further important progress in understanding such Monge-Ampère equations with right hand side in L^p was made by Kołodziej [109] using techniques of pluripotential theory. A detailed study of Kähler-Einstein metrics on singular varieties is given in Eyssidieux-Guedj-Zeriahi [87].

The lack of regularity of such a Kähler-Einstein metric at the singular points means that results such as Matsushima's Theorem on the automorphism group and the vanishing of the Futaki invariant cannot be generalized in a straightforward way.

Instead Chen-Donaldson-Sun [44] proved these results in the singular setting using the theory of geodesics and Berndtsson's convexity and uniqueness results [22].

It is natural to expect that Theorem 2 can be extended to \mathbf{Q} -Fano varieties, and indeed Berman [16] showed that \mathbf{Q} -Fano varieties admitting (singular) Kähler-Einstein metrics are K-stable. The converse, however, is so far only known for smoothable \mathbf{Q} -Fanos, by work of Spotti-Sun-Yao [154].

5. APPLICATIONS

In this final section we briefly survey some applications of the Yau-Tian-Donaldson conjecture for Fano manifolds, and the techniques involved in its solution. The first is perhaps the most direct type of application, namely obtaining new examples of Kähler-Einstein manifolds. The second and third are more theoretical, on understanding the moduli space of Fano Kähler-Einstein manifolds, and the behavior of singular Kähler-Einstein metrics at the singular points.

5.1. New examples. According to Theorem 2, K-stability is a necessary and sufficient condition for a Fano manifold M to admit a Kähler-Einstein metric. As such it is natural to expect that the result can be used to show the existence of new Kähler-Einstein metrics. Unfortunately for any given Fano manifold M of dimension at least two, there are infinitely many possible special degenerations, and so at present there is no general method for testing K-stability. On the other hand there are special circumstances in which one can show that certain varieties admit a Kähler-Einstein metric.

In the two-dimensional case, Tian [165] used the α -invariant [163] and its refinements, to show that any Fano surface with reductive automorphism group admits a Kähler-Einstein metric. More generally, calculations of the α -invariant, and related concepts such as Nadel's multiplier ideal sheaves [128], lead to many examples of Kähler-Einstein manifolds (see e.g. [40, 59]). Recently Fujita [91] showed that Fano n -folds M with α -invariant $\alpha(M) = n/(n+1)$, which are borderline for Tian's criterion [163], are still K-stable, and as such they admit Kähler-Einstein metrics.

A different source of examples where K-stability can be checked effectively is Fano manifolds M with large automorphism groups, since in [56] we showed that it is enough to check special degenerations that are compatible with the automorphism group. For instance if M is a toric manifold, then the only torus equivariant special degenerations for M are products, and so M admits a Kähler-Einstein metric whenever the Futaki invariants of all vector fields on M vanish (this result is originally due to Wang-Zhu [175], using different methods). This result has recently been generalized to reductive group compactifications by Delcroix [58] using analytic methods. Subsequently, by classifying equivariant special degenerations, Delcroix [57] generalized this result to spherical varieties, giving new examples of Kähler-Einstein manifolds.

A more general setting is complexity-one spherical varieties [170], for instance Fano n -folds with an effective action of an $(n-1)$ -dimensional torus. The equivariant special degenerations of these complexity-one T-varieties were classified by Ilten-Süß [101], using the theory of polyhedral divisors [2] and as an application they obtained several new examples of K-stable Fano 3-folds.

Theorem 2 was generalized to the setting of Kähler cones in [52] and, combined with the methods of Ilten-Süß [101], this lead to new examples of Ricci flat Kähler cones. As we discussed in Section 4.1, such Kähler cones arise as tangent cones

of Gromov-Hausdorff limits of Kähler-Einstein manifolds, but they have also been studied extensively from the point of view of Sasakian geometry [95, 26, 152]. To give a simple example, consider the affine variety $M_{p,q} \subset \mathbf{C}^4$, given by the equation $xy + z^p + w^q = 0$. This has a natural action of a 2-torus, acting diagonally on \mathbf{C}^4 . Using K-stability, it is shown in [52] that $M_{p,q}$ admits a Ricci flat Kähler cone metric compatible with the torus action, if and only if $2p > q$ and $2q > p$. These two conditions correspond to the calculation of a Donaldson-Futaki invariant for two torus equivariant degenerations of $M_{p,q}$, to the hypersurfaces given by the equations $xy + z^p = 0$ and $xy + w^q = 0$ respectively.

For manifolds without such large symmetry groups, the general approach of Tian [165] for Fano surfaces has recently been revisited in the work of Odaka-Spotti-Sun [134], Spotti-Sun [153] and Liu-Xu [119]. Here, given a family of Fano manifolds some elements of which are known to be Kähler-Einstein, one tries to find new Kähler-Einstein manifolds using a continuity method in the family. The key problem is to classify the possible Gromov-Hausdorff limits of Kähler-Einstein manifolds in the family. Using this method, and combined with work of Fujita [91], Liu-Xu [119] showed that all smooth 3-dimensional Fano hypersurfaces admit Kähler-Einstein metrics. Ultimately such a study can lead to a complete understanding of the Kähler-Einstein moduli space and its compactification, as we will discuss in the next section.

While a general algorithm for testing K-stability seems to be some ways off at present, it seems likely that these techniques will lead to many more examples.

5.2. The moduli space of Kähler-Einstein manifolds. A basic question in algebraic geometry is to understand the moduli space of a certain manifold, as well as its compactifications. Perhaps the most well known example is the Deligne-Mumford compactification of the moduli space of hyperbolic Riemann surfaces. This compactification of the moduli space of smooth Riemann surfaces is obtained by adding certain “stable” curves with nodal singularities, and the degeneration of smooth curves to these singular ones can in turn be modeled differential geometrically by degenerating families of hyperbolic metrics.

Canonically polarized manifolds are a higher dimensional generalization of hyperbolic Riemann surfaces and compact moduli spaces are obtained by adding the “KSBA-stable” varieties (see Kollár-Shepherd-Barron [108] Alexeev [1], and the survey [107]). It is interesting to note that the KSBA-stable varieties were subsequently found to coincide with the K-stable varieties by work of Odaka [131], and furthermore Berman-Guenancia [20] showed that these all admit (singular) Kähler-Einstein metrics. To complete the analogy with hyperbolic Riemann surfaces it remains to understand the convergence of smooth Kähler-Einstein metrics to the singular ones on a metric level. We emphasize that in the canonically polarized setting all smooth manifolds admit Kähler-Einstein metrics with negative Ricci curvature by Yau’s theorem [177], and K-stability only enters in picking out the right type of singular varieties to allow.

By contrast, in the Fano case one cannot form a Hausdorff moduli space containing all the smooth manifolds, since there are non-trivial families over \mathbf{C} that are isomorphic to a product over \mathbf{C}^* . For instance any non-trivial test-configuration for a Fano manifold with smooth central fiber gives such a family. One can hope, however, that K-stable Fano manifolds form good moduli spaces, and that a compactification is obtained by including certain singular K-stable varieties. Moreover

this compact moduli space should agree with the space of Kähler-Einstein Fano manifolds, compactified using the Gromov-Hausdorff topology. A first step is to understand the neighborhood of a Kähler-Einstein manifold in the moduli space. In [159] (see also Brönnle [27]) we proved that given a Kähler-Einstein manifold M , its small deformations that admit KE metrics are precisely the ones that are (poly)stable for the action of $\text{Aut}(M)$ on the deformation space of M . Thus a small neighborhood of M in the moduli space is modeled by a GIT quotient for the action of $\text{Aut}(M)$.

In the case of surfaces, a global study of the moduli space of Kähler-Einstein manifolds was achieved by Odaka-Spotti-Sun [134], who explicitly identified the compactified moduli space in terms of certain algebro-geometric compactifications. In higher dimensions Spotti-Sun-Yao [154] and Li-Wang-Xu [114] showed that smoothable K -stable \mathbf{Q} -Fano varieties, i.e. the ones that we expect to appear in the compactified moduli space, all admit singular Kähler-Einstein metrics, which in addition can be obtained as Gromov-Hausdorff limits of smooth KE metrics. Regarding the global structure of the moduli space, the smoothable K -stable \mathbf{Q} -Fano varieties are parametrized by a proper algebraic space by work of Odaka [132] and Li-Wang-Xu [114], and moreover the K -stable smooth Fanos form a quasi-projective variety by Li-Wang-Xu [115]. It remains to understand more precisely what varieties these moduli spaces parametrize in analogy with Odaka-Spotti-Sun's work [134] on surfaces, but as we mentioned above, there are special cases where progress has been made, such as in the case of degree 4 del Pezzo manifolds by Spotti-Sun [153] and cubic threefolds by Liu-Xu [119].

5.3. Asymptotics of singular Kähler-Einstein metrics. In Section 4.3 we discussed how singular Kähler-Einstein metrics play an important role in the proof of Theorem 2. Of particular interest are those singular Kähler-Einstein varieties, which arise as limits of smooth Kähler-Einstein manifolds, in view of the importance of forming complete moduli spaces of Fano manifolds [134, 154, 132, 114], as well as Calabi-Yau manifolds [181]. We do not yet have a detailed understanding of the behavior of such singular Kähler-Einstein metrics near the singular points, although much progress has been made. Such an understanding would be necessary in order to use differential geometric techniques on these singular spaces (see e.g. [30, 97] for such applications).

A first step in understanding the metric behavior near the singular points is to understand the metric tangent cone, as discussed in Section 4.1. In the general Riemannian context the tangent cone may depend on the sequence of rescalings chosen to construct it, but in the setting of Kähler-Einstein manifolds Donaldson-Sun [66] showed that the tangent cone is unique, and it has the structure of an affine variety. They also made progress towards an algebro-geometric description of the metric tangent cone, and conjectured that in general the tangent cone at a point $x \in M$ of a singular Kähler-Einstein metric on M can be determined from the germ of the singularity (M, x) using K -stability. Note that it is not always the case that the germ of the tangent cone at the vertex is biholomorphic to the germ (M, x) . For example (see [157], [99]) there is a Calabi-Yau metric on a neighborhood of the origin in the hypersurface $x_0^k + x_1^2 + x_2^2 + x_3^2 = 0$ in \mathbf{C}^4 whose tangent cone at the origin is given by $\mathbf{C} \times \mathbf{C}^2/\mathbf{Z}_2$, if $k > 4$.

An important special case of this conjecture for singular Calabi-Yau varieties was proven by Hein-Sun [100], namely when the germ (M, x) is isomorphic to

the germ $(C, 0)$ of a Ricci flat Kähler cone C at its vertex, and C satisfies some additional technical conditions. This uses some ideas of Li [111] and Li-Liu [113] on an alternative characterization of K-semistability based on the work of Fujita [92] we mentioned in Section 3.4. Using this, Hein-Sun showed that the singular Ricci flat metric on M near x is asymptotic, in a strong sense, to the Ricci flat metric on C . Recently, using more algebro-geometric techniques, Li-Xu [117] have also made significant progress towards resolving Donaldson-Sun's conjecture, and they have announced a complete solution.

REFERENCES

- [1] ALEXEEV, V. Log canonical singularities and complete moduli of stable pairs. *arXiv:9608013*.
- [2] ALTMANN, K., AND HAUSEN, J. Polyhedral divisors and algebraic torus actions. *Math. Ann.* 334 (2006), 557–607.
- [3] ANDERSON, M. Ricci curvature bounds and Einstein metrics on compact manifolds. *J. Amer. Math. Soc.* 2, 3 (1989), 455–490.
- [4] ANDERSON, M. Convergence and rigidity of manifolds under Ricci curvature bounds. *Invent. Math.* 97 (1990), 429–445.
- [5] APOSTOLOV, V., CALDERBANK, D. M. J., GAUDUCHON, P., AND TØNNESSEN-FRIEDMAN, C. W. Hamiltonian 2-forms in Kähler geometry III, extremal metrics and stability. *Invent. Math.* 173, 3 (2008), 547–601.
- [6] AREZZO, C., AND PACARD, F. Blowing up and desingularizing constant scalar curvature Kähler manifolds. *Acta Math.* 196, 2 (2006), 179–228.
- [7] AREZZO, C., AND TIAN, G. Infinite geodesic rays in the space of Kähler potentials. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)* 2, 4 (2003), 617–630.
- [8] ATIYAH, M. F., AND BOTT, R. The Yang-Mills equations over Riemann surfaces. *Philos. Trans. Roy. Soc. London Ser. A* 308 (1983), 523–615.
- [9] AUBIN, T. Équations du type Monge-Ampère sur les variétés kählériennes compactes. *Bull. Sci. Math. (2)* 102, 1 (1978), 63–95.
- [10] AUBIN, T. Réduction de cas positif de l'équation de Monge-Ampère sur les variétés kählériennes compactes à la démonstration d'une inégalité. *J. Funct. Anal.* 57, 2 (1984), 143–153.
- [11] BAMLER, R. Convergence of Ricci flows with bounded scalar curvature. *arXiv:1603.05235*.
- [12] BANDO, S., KASUE, A., AND NAKAJIMA, H. On a construction of coordinates at infinity on manifolds with fast curvature decay and maximal volume growth. *Invent. Math.* 97, 2 (1989), 313–349.
- [13] BANDO, S., AND MABUCHI, T. Uniqueness of Einstein Kähler metrics modulo connected group actions. In *Algebraic geometry, Sendai* (1985), vol. 10 of *Adv. Stud. Pure Math.*, pp. 11–40.
- [14] BEDFORD, E., AND TAYLOR, B. A. The Dirichlet problem for a complex Monge-Ampère equation. *Invent. Math.* 37, 1 (1976), 1–44.
- [15] BERMAN, R. On the optimal regularity of weak geodesics in the space of metrics on a polarized manifold. *arXiv:1405.6482*.
- [16] BERMAN, R. K-polystability of Q-Fano varieties admitting Kähler-Einstein metrics. *Invent. Math.* 203, 3 (2016), 973–1025.
- [17] BERMAN, R., AND BERNDTSSON, B. Convexity of the K-energy on the space of Kähler metrics and uniqueness of extremal metrics. *J. Amer. Math. Soc.* 30, 4 (2017), 1165–1196.
- [18] BERMAN, R., BOUCKSOM, S., AND JONSSON, M. A variational approach to the Yau-Tian-Donaldson conjecture. *arXiv:1509.04561*.
- [19] BERMAN, R., DARVAS, T., AND LU, C. H. Regularity of weak minimizers of the K-energy and applications to properness and K-stability. *arXiv:1602.03114*.
- [20] BERMAN, R., AND GUENANCIA, H. Kähler-Einstein metrics on stable varieties and log canonical pairs. *Geom. and Func. Anal.* 24, 6 (2014), 1683–1730.
- [21] BERNDTSSON, B. Curvature of vector bundles associated to holomorphic fibrations. *Ann. of Math. (2)* 169, 2 (2009), 531–560.

- [22] BERNDTSSON, B. A Brunn-Minkowski type inequality for Fano manifolds and some uniqueness theorems in Kähler geometry. *Invent. Math.* 200, 1 (2015), 149–200.
- [23] BLOCKI, Z. *On geodesics in the space of Kähler metrics*, vol. 21 of *Adv. Lect. Math. (ALM)*. Int. Press, Somerville, MA, 2012, pp. 3–19.
- [24] BOUCKSOM, S., FAVRE, C., AND JONSSON, M. Valuations and plurisubharmonic singularities. *Publ. RIMS* 44 (2008), 449–494.
- [25] BOUCKSOM, S., HISAMOTO, T., AND JONSSON, M. Uniform K-stability, Duistermaat-Heckman measures and singularities of pairs. *arXiv:1504.06568*.
- [26] BOYER, C., AND GALICKI, K. *Sasakian geometry*. Oxford Mathematical Monographs. OUP, 2008.
- [27] BRÖNNLE, T. A. *Deformation constructions of extremal metrics*. Phd Thesis, Imperial College London.
- [28] CALABI, E. The space of Kähler metrics. *Proc. Internat. Congress Math. Amsterdm*, 2 (1954), 206–207.
- [29] CALABI, E. Extremal Kähler metrics. In *Seminar on Differential Geometry*, S. T. Yau, Ed. Princeton, 1982.
- [30] CAMPANA, F., GUENANCIA, H., AND PĂUN, M. Metrics with cone singularities along normal crossing divisors and holomorphic tensor fields. *Ann. Sci. École Norm. Sup. (4)* 46 (2013), 879–916.
- [31] CAO, H. D. Deformation of Kähler metrics to Kähler-Einstein metrics on compact Kähler manifolds. *Invent. Math.* 81, 2 (1985), 359–372.
- [32] CATLIN, D. The Bergman kernel and a theorem of Tian. In *Analysis and geometry in several complex variables (Katata, 1997)* (1999), Trends Math., Birkhäuser Boston Inc., pp. 1–23.
- [33] CHEEGER, J. *Degeneration of Riemannian metrics under Ricci curvature bounds*. Lezioni Fermiane. Scuola Normale Superiore, Pisa, 2001.
- [34] CHEEGER, J., AND COLDING, T. On the structure of spaces with Ricci curvature bounded below. I. *J. Differential Geom.* 46, 3 (1997), 406–480.
- [35] CHEEGER, J., AND COLDING, T. On the structure of spaces with Ricci curvature bounded below. II. *J. Differential Geom.* 54, 1 (2000), 13–35.
- [36] CHEEGER, J., AND COLDING, T. On the structure of spaces with Ricci curvature bounded below. III. *J. Differential Geom.* 54, 1 (2000), 37–74.
- [37] CHEEGER, J., AND EBIN, D. G. *Comparison theorems in Riemannian geometry*. AMS Chelsea Publishing, Providence, RI, 2008.
- [38] CHEEGER, J., AND NABER, A. Lower bounds on Ricci curvature and quantitative behavior of singular sets. *Invent. Math.* 191, 2 (2013), 321–339.
- [39] CHEEGER, J., AND NABER, A. Regularity of Einstein manifolds and the codimension 4 conjecture. *Ann. of Math. (2)* 182, 3 (2015), 1093–1165.
- [40] CHELTSOV, I., AND PARK, J. Log canonical thresholds and generalized Eckardt points. *Sb. Math.* 193, 5–6 (2002), 779–789.
- [41] CHEN, X., DONALDSON, S., AND SUN, S. Kähler-Einstein metrics and stability. *Int. Math. Res. Not. IMRN*, 8 (2014), 2119–2125.
- [42] CHEN, X., DONALDSON, S., AND SUN, S. Kähler-Einstein metrics on Fano manifolds. I: Approximation of metrics with cone singularities. *J. Amer. Math. Soc.* 28, 1 (2015), 183–197.
- [43] CHEN, X., DONALDSON, S., AND SUN, S. Kähler-Einstein metrics on Fano manifolds. II: Limits with cone angle less than 2π . *J. Amer. Math. Soc.* 28, 1 (2015), 199–234.
- [44] CHEN, X., DONALDSON, S., AND SUN, S. Kähler-Einstein metrics on Fano manifolds. III: Limits as cone angle approaches 2π and completion of the main proof. *J. Amer. Math. Soc.* 28, 1 (2015), 235–278.
- [45] CHEN, X., AND SUN, S. Space of Kähler metrics V – Kähler quantization. In *Metric and differential geometry*, vol. 297 of *Progr. Math.* Birkhäuser Verlag, Basel, 2012, pp. 19–41.
- [46] CHEN, X., SUN, S., AND WANG, B. Kähler-Ricci flow, Kähler-Einstein metric, and K-stability. *arXiv:1508.04397*.
- [47] CHEN, X. X. The space of Kähler metrics. *J. Differential Geom.* 56, 2 (2000), 189–234.
- [48] CHEN, X. X., AND TANG, Y. Test configuration and geodesic rays. *Astérisque*, 321 (2008), 139–167.
- [49] CHEN, X. X., AND WANG, B. Space of Ricci flows (II). *arXiv:1405.6797*.

- [50] CHU, J., TOSATTI, V., AND WEINKOVE, B. On the $C^{1,1}$ regularity of geodesics in the space of Kähler metrics. *arXiv:1611.02390*.
- [51] COLDING, T. H. Ricci curvature and volume convergence. *Ann. of Math. (2)* 145, 3 (1997), 477–501.
- [52] COLLINS, T., AND SZÉKELYHIDI, G. Sasaki-Einstein metrics and K-stability. *arXiv:1512.07213*.
- [53] DARVAS, T. Morse theory and geodesics in the space of Kähler metrics. *Proc. Amer. Math. Soc.* 142, 8 (2014), 2775–2782.
- [54] DARVAS, T. The Mabuchi geometry of finite energy classes. *Adv. Math.* 285 (2015), 182–219.
- [55] DARVAS, T., AND RUBINSTEIN, Y. A. Tian’s properness conjectures and Finsler geometry of the space of Kähler metrics. *J. Amer. Math. Soc.* 30, 2 (2017), 347–387.
- [56] DATAR, V., AND SZÉKELYHIDI, G. Kähler-Einstein metrics along the smooth continuity method. *Geom. and Func. Anal.* 26, 4 (2016), 975–1010.
- [57] DELCROIX, T. K-stability of Fano spherical varieties. *arXiv:1608.01852*.
- [58] DELCROIX, T. Kähler-Einstein metrics on group compactifications. *Geom. and Func. Anal.* 27, 1 (2017), 78–129.
- [59] DEMAILLY, J.-P., AND KOLLÁR, J. Semi-continuity of complex singularity exponents and Kähler-Einstein metrics on Fano orbifolds. *Ann. Sci. Éc. Norm. Supér. (4)* 34 (2001), 525–556.
- [60] DERVAN, R. Relative K-stability for Kähler manifolds. *arXiv:1611.00569*.
- [61] DERVAN, R. Alpha invariants and K-stability for general polarizations of Fano varieties. *Int. Math. Res. Not. IMRN*, 16 (2015), 7162–7189.
- [62] DERVAN, R. Uniform stability of twisted constant scalar curvature Kähler metrics. *Int. Math. Res. Not.* 15 (2016), 4728–4783.
- [63] DERVAN, R., AND ROSS, J. K-stability for Kähler manifolds. *arXiv:1602.08983*.
- [64] DING, W. Y. Remarks on the existence problem of positive Kähler-Einstein metrics. *Math. Ann.* 282, 3 (1988), 463–471.
- [65] DING, W. Y., AND TIAN, G. Kähler-Einstein metrics and the generalized Futaki invariant. *Invent. Math.* 110, 2 (1992), 315–335.
- [66] DONALDSON, S., AND SUN, S. Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry, II. *arXiv:1507.05082*.
- [67] DONALDSON, S., AND SUN, S. Gromov-Hausdorff limits of Kähler manifolds and algebraic geometry. *Acta Math.* 213, 1 (2014), 63–106.
- [68] DONALDSON, S. K. The Ding functional, Berndtsson convexity and moment maps. *arXiv:1503.05173*.
- [69] DONALDSON, S. K. A new proof of a theorem of Narasimhan and Seshadri. *J. Differential Geom.* 18, 2 (1983), 269–277.
- [70] DONALDSON, S. K. Anti self-dual yang-mills connections over complex algebraic surfaces and stable vector bundles. *Proc. London Math. Soc.* 50 (1985), 1–26.
- [71] DONALDSON, S. K. Infinite determinants, stable bundles and curvature. *Duke Math. J.* 54, 1 (1987), 231–247.
- [72] DONALDSON, S. K. Remarks on gauge theory, complex geometry and four-manifold topology. In *Fields Medallists’ Lectures*, Atiyah and Iagolnitzer, Eds. World Scientific, 1997, pp. 384–403.
- [73] DONALDSON, S. K. Symmetric spaces, Kähler geometry and Hamiltonian dynamics. In *Northern California Symplectic Geometry Seminar*, vol. 196 of *Amer. Math. Soc. Transl. Ser. 2*. Amer. Math. Soc., Providence, RI, 1999, pp. 13–33.
- [74] DONALDSON, S. K. Planck’s constant in complex and almost-complex geometry. In *XIIIth International Congress on Mathematical Physics (London, 2000)*. Int. Press, Boston, MA, 2001, pp. 63–72.
- [75] DONALDSON, S. K. Scalar curvature and projective embeddings, I. *J. Differential Geom.* 59 (2001), 479–522.
- [76] DONALDSON, S. K. Holomorphic discs and the complex Monge-Ampère equation. *J. Symplectic Geom.* 1, 2 (2002), 171–196.
- [77] DONALDSON, S. K. Scalar curvature and stability of toric varieties. *J. Differential Geom.* 62 (2002), 289–349.
- [78] DONALDSON, S. K. Lower bounds on the Calabi functional. *J. Differential Geom.* 70, 3 (2005), 453–472.

- [79] DONALDSON, S. K. Scalar curvature and projective embeddings, II. *Q. J. Math.* 56, 3 (2005), 345–356.
- [80] DONALDSON, S. K. Discussion of the Kähler-Einstein problem. <http://www2.imperial.ac.uk/~skdona/KENOTES.PDF> (2009).
- [81] DONALDSON, S. K. Some numerical results in complex differential geometry. *Pure Appl. Math. Q.* 5, 2 (2009), 571–618.
- [82] DONALDSON, S. K. Kähler metrics with cone singularities along a divisor. In *Essays in mathematics and its applications*. Springer, Heilderlberg, 2012, pp. 49–79.
- [83] DONALDSON, S. K. *Stability, birational transformations and the Kähler-Einstein problem*, vol. 17 of *Surv. Differ. Geom.* Int. Press, Boston, MA, 2012, pp. 203–228.
- [84] DONALDSON, S. K. b-stability and blow-ups. *Proc. Edinb. Math. Soc. (2)* 57, 1 (2014), 125–137.
- [85] DONALDSON, S. K., AND KRONHEIMER, P. B. *The Geometry of Four-Manifolds*. OUP, 1990.
- [86] EYSSIDIEUX, P. Métriques de Kähler-Einstein sur les variétés de Fano [d’après Chen-Donaldson-Sun et Tian]. *Astérisque*, 380, Séminaire Bourbaki. Vol. 2014/2015 (2016).
- [87] EYSSIDIEUX, P., GUEDJ, V., AND ZERIAHI, A. Singular Kähler-Einstein metrics. *J. Amer. Math. Soc.* 22, 3 (2009), 607–639.
- [88] FINE, J. Calabi flow and projective embeddings. With an appendix by Kefeng Liu and Xiaonan Ma. *J. Differential Geom.* 84, 3 (2010), 489–523.
- [89] FINE, J. Quantisation and the Hessian of Mabuchi energy. *Duke Math. J.* 161, 14 (2012), 2753–2798.
- [90] FUJIKI, A. Moduli space of polarized algebraic manifolds and Kähler metrics [translation of *Sūgaku* 42 (1990), no. 3, 231–243;]. *Sugaku Expositions* 5, 2 (1992), 173–191.
- [91] FUJITA, K. K-stability of Fano manifolds with not small alpha invariants. *arXiv:1606.08261*.
- [92] FUJITA, K. Optimal bounds for the volumes of Kähler-Einstein Fano manifolds. *arXiv:1508.04578*.
- [93] FUTAKI, A. An obstruction to the existence of Einstein-Kähler metrics. *Invent. Math.* 73 (1983), 437–443.
- [94] GAUDUCHON, P. *Calabi’s extremal Kähler metrics: An elementary introduction*.
- [95] GAUNTLETT, J. P., MARTELLI, D., SPARKS, J., AND YAU, S.-T. Obstructions to the existence of Sasaki-Einstein metrics. *Comm. Math. Phys.* 273, 3 (2007), 803–827.
- [96] GROMOV, M. *Metric structures for Riemannian and non-Riemannian spaces*. Birkhäuser Boston Inc., 2007.
- [97] GUENANCIA, H., AND TAJI, B. Orbifold stability and Miyaoka-Yau inequality for minimal pairs. *arXiv:1611.0598*.
- [98] HAMILTON, R. Three-manifolds with positive Ricci curvature. *J. Differential Geom.* 17, 2 (1982), 255–306.
- [99] HEIN, H.-J., AND NABER, A. Isolated Einstein singularities with singular tangent cones. *in preparation*.
- [100] HEIN, H.-J., AND SUN, S. Calabi-Yau manifolds with isolated conical singularities. *arXiv:1607.02940*.
- [101] ILTEN, N., AND SÜSS, H. K-stability for varieties with torus action of complexity one. *Duke Math. J.* 166, 1 (2017), 177–204.
- [102] JEFFRES, T., MAZZEO, R., AND RUBINSTEIN, Y. A. Kähler-Einstein metrics with edge singularities. *Ann. of Math. (2)* 183, 1 (2016), 95–176.
- [103] JIANG, W. Bergman kernel along the Kähler Ricci flow and Tian’s conjecture. *J. Reine Angew. Math.* 717 (2016), 195–226.
- [104] KÄHLER, E. Über eine bemerkenswerte Hermitesche Metrik. *Abh. Math. Sem. Univ. Hamburg* 9, 1 (1933), 173–186.
- [105] KAZDAN, J. L., AND WARNER, F. W. Curvature functions for compact 2-manifolds. *Ann. of Math.* 99 (1974), 14–47.
- [106] KEMPF, G., AND NESS, L. The length of vectors in representation spaces. In *Algebraic geometry (Proc. Summer Meeting, Univ. Copenhagen, Copenhagen, 1978)*, vol. 732 of *Lecture Notes in Math.* Springer, Berlin, 1979, pp. 233–243.
- [107] KOLLÁR, J. Moduli of varieties of general type. In *Handbook of moduli. Vol. II*. 2013, pp. 131–157.
- [108] KOLLÁR, J., AND SHEPHERD-BARRON, N. I. Threefolds and deformations of surface singularities. *Invent. Math.* 91, 2 (1988), 299–338.

- [109] KOŁODZIEJ, S. The complex Monge-Ampère equation. *Acta Math.* 180, 1 (1998), 69–117.
- [110] LEMPert, L., AND VIVAS, L. Geodesics in the space of Kähler metrics. *Duke Math. J.* 162, 7 (2013), 1369–1381.
- [111] LI, C. K-semistability is equivariant volume minimization. *arXiv:1512.07205*.
- [112] LI, C. Constant scalar curvature Kähler metric obtains the minimum of K-energy. *Internat. Math. Res. Notices*, 9 (2011), 2161–2175.
- [113] LI, C., AND LIU, Y. Kähler-Einstein metrics and volume minimization. *arXiv:1602.05094*.
- [114] LI, C., WANG, X., AND XU, C. On proper moduli space of smoothable Kähler-Einstein Fano varieties. *arXiv:1411.0761*.
- [115] LI, C., WANG, X., AND XU, C. Quasi-projectivity of the moduli space of smooth Kähler-Einstein Fano manifolds. *arXiv:1502.06532*.
- [116] LI, C., AND XU, C. Stability of Valuations and Kollár Components. *arXiv:1604.05398*.
- [117] LI, C., AND XU, C. Stability of valuations: higher rational rank. *arXiv:1707.05561*.
- [118] LI, C., AND XU, C. Special test configurations and K-stability of Fano varieties. *Ann. of Math. (2)* 180, 2 (2014), 197–232.
- [119] LIU, Y., AND XU, C. K-stability of cubic threefolds. *arXiv:1706.01933*.
- [120] LU, Z. On the lower order terms of the asymptotic expansion of Tian-Yau-Zelditch. *Amer. J. Math.* 122, 2 (1998), 235–273.
- [121] LUNA, D. Slices étales. In *Sur les groupes algébriques*, Bull. Soc. Math. France, Paris, Mémoire 33. Soc. Math. France, Paris, 1973, pp. 81–105.
- [122] LUO, H. Geometric criterion for Gieseker-Mumford stability of polarized manifolds. *J. Differential Geom.* 49 (1998), 577–599.
- [123] MABUCHI, T. K-energy maps integrating Futaki invariants. *Tohoku Math. J.* 38, 4 (1986), 575–593.
- [124] MABUCHI, T. Some symplectic geometry on compact Kähler manifolds. *Osaka J. Math.* 24 (1987), 227–252.
- [125] MABUCHI, T. An energy-theoretic approach to the Hitchin-Kobayashi correspondence for manifolds, I. *Invent. Math.* 159, 2 (2005), 225–243.
- [126] MATSUSHIMA, Y. Sur la structure du groupe d'homéomorphismes analytiques d'une certaine variété kählérienne. *Nagoya Math. J.* 11 (1957), 145–150.
- [127] MUMFORD, D., FOGARTY, J., AND KIRWAN, F. *Geometric invariant theory*, third ed., vol. 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (2) [Results in Mathematics and Related Areas (2)]*. Springer-Verlag, Berlin, 1994.
- [128] NADEL, A. M. Multiplier ideal sheaves and Kähler-Einstein metrics of positive scalar curvature. *Ann. of Math. (2)* 132, 3 (1990), 549–596.
- [129] NARASIMHAN, M. S., AND SESHADRI, C. S. Stable and unitary vector bundles on a compact Riemann surface. *Ann. of Math. (2)* 82 (1965), 540–567.
- [130] ODAKA, Y. A generalization of the Ross-Thomas slope theory. *Osaka J. Math.* 50, 1 (2013), 171–185.
- [131] ODAKA, Y. The GIT stability of polarized varieties via discrepancy. *Ann. of Math. (2)* 177, 2 (2013), 645–661.
- [132] ODAKA, Y. Compact moduli spaces of Kähler-Einstein Fano varieties. *Publ. Res. Inst. Math. Sci.* 51, 3 (2015), 549–565.
- [133] ODAKA, Y., AND SANO, Y. Alpha invariant and K-stability of Q-Fano varieties. *Adv. Math.* 229, 5 (2012), 2818–2834.
- [134] ODAKA, Y., SPOTTI, C., AND SUN, S. Compact moduli spaces of Del Pezzo surfaces and Kähler-Einstein metrics. *J. Differential Geom.* 102, 1 (2016), 127–172.
- [135] PAUL, S. T. Hyperdiscriminant polytopes, Chow polytopes, and Mabuchi energy asymptotics. *Ann. of Math. (2)* 175, 1 (2012), 255–296.
- [136] PAUL, S. T., AND TIAN, G. CM stability and the generalized Futaki invariant, I. *arXiv:0605278*.
- [137] PERELMAN, G. The entropy formula for the Ricci flow and its geometric applications. *math.DG/0211159*.
- [138] PHONG, D. H., SONG, J., AND STURM, J. Degenerations of Kähler-Ricci solitons on Fano manifolds. *Univ. Iagel. Acta Math.* 52 (2015), 29–43.
- [139] PHONG, D. H., AND STURM, J. Lectures on stability and constant scalar curvature. In *Current Developments in Mathematics 2007*. International Press.

- [140] PHONG, D. H., AND STURM, J. Test-configurations for K-stability and geodesic rays. *Jour. Symplectic Geom.* 5, 2 (2007), 221–247.
- [141] PHONG, D. H., AND STURM, J. Regularity of geodesic rays and Monge-Ampère equations. *Proc. Amer. Math. Soc.* 138, 10 (2010), 3637–3650.
- [142] ROSS, J., AND THOMAS, R. P. An obstruction to the existence of constant scalar curvature Kähler metrics. *J. Differential Geom.* 72 (2006), 429–466.
- [143] ROSS, J., AND THOMAS, R. P. A study of the Hilbert-Mumford criterion for the stability of projective varieties. *J. Algebraic Geom.* 16, 2 (2007), 201–255.
- [144] ROSS, J., AND WITT NYSTRÖM, D. Harmonic discs of solutions to the complex homogeneous Monge-Ampère equation. *Publ. Math. Inst. Hautes Études Sci.* 122 (2015), 315–335.
- [145] RUBINSTEIN, Y. A., AND ZELDITCH, S. Bergman approximations of harmonic maps into the space of Kähler metrics on toric varieties. *J. Symplectic Geom.* 8, 3 (2010), 239–265.
- [146] SANO, Y., AND TIPLER, C. Extremal metrics and lower bound of the modified K-energy. *J. Eur. Math. Soc.* 17, 9 (2015), 2289–2310.
- [147] SEMMES, S. Complex Monge-Ampère equations and symplectic manifolds. *Amer. J. Math.* 114 (1992), 495–550.
- [148] SEYYEDALI, R. Relative Chow stability and extremal metrics. *arXiv:1610.07555*.
- [149] SHIFFMAN, B., AND ZELDITCH, S. Distribution of zeros of random and quantum chaotic sections of positive line bundles. *Comm. Math. Phys.* 200, 3 (1999), 661–683.
- [150] SJÖSTRÖM DYREFELT, Z. K-semistability of cscK manifolds with transcendental cohomology class. *arXiv:1601.07659*.
- [151] SONG, J., AND ZELDITCH, S. Bergman metrics and geodesics in the space of Kähler metrics on toric varieties. *Anal. PDE* 3, 3 (2010), 295–358.
- [152] SPARKS, J. New results in Sasaki-Einstein geometry. In *Riemannian topology and geometric structures on manifolds*, vol. 271 of *Progr. Math.* Birkhäuser Boston Inc., 2009, pp. 161–184.
- [153] SPOTTI, C., AND SUN, S. Explicit Gromov-Hausdorff compactifications of moduli spaces of Kähler-Einstein Fano manifolds. *arXiv:1705.00377*.
- [154] SPOTTI, C., SUN, S., AND YAO, C. Existence and deformations of Kahler-Einstein metrics on smoothable Q-Fano varieties. *Duke Math. J.* 165, 16 (2016), 3043–3083.
- [155] STOPPA, J. K-stability of constant scalar curvature Kähler manifolds. *Adv. Math.* 221, 4 (2009), 1397–1408.
- [156] STOPPA, J., AND SZÉKELYHIDI, G. Relative K-stability of extremal metrics. *J. Eur. Math. Soc.* 13, 4 (2011), 899–909.
- [157] SZÉKELYHIDI, G. Degenerations of \mathbf{C}^n and Calabi-Yau metrics. *arXiv:1706.00357*.
- [158] SZÉKELYHIDI, G. Extremal metrics and K-stability. *Bull. Lond. Math. Soc.* 39, 1 (2007), 76–84.
- [159] SZÉKELYHIDI, G. The Kähler-Ricci flow and K-polystability. *Amer. J. Math.* 132 (2010), 1077–1090.
- [160] SZÉKELYHIDI, G. Filtrations and test-configurations, with an appendix by S. Boucksom. *Math. Ann.* 362, 1-2 (2015), 451–484.
- [161] SZÉKELYHIDI, G. The partial C^0 -estimate along the continuity method. *J. Amer. Math. Soc.* 29, 2 (2016), 537–560.
- [162] THOMAS, R. P. Notes on GIT and symplectic reduction for bundles and varieties. In *Surveys in differential geometry. Vol. X*. Int. Press, Somerville, MA, 2006, pp. 221–273.
- [163] TIAN, G. On Kähler-Einstein metrics on certain Kähler manifolds with $c_1(M) > 0$. *Invent. Math.* 89 (1987), 225–246.
- [164] TIAN, G. On a set of polarized Kähler metrics on algebraic manifolds. *J. Differential Geom.* 32, 1 (1990), 99–130.
- [165] TIAN, G. On Calabi’s conjecture for complex surfaces with positive first Chern class. *Invent. Math.* 101, 1 (1990), 101–172.
- [166] TIAN, G. Kähler-Einstein metrics with positive scalar curvature. *Invent. Math.* 137 (1997), 1–37.
- [167] TIAN, G. *Canonical metrics in Kähler geometry*. Lectures in Mathematics ETH Zürich. Birkhäuser Verlag, Basel, 2000.
- [168] TIAN, G., AND ZHANG, Z. Degeneration of Kähler-Ricci solitons. *Int. Math. Res. Not. IMRN*, 5 (2012), 957–985.
- [169] TIAN, G., AND ZHANG, Z. Regularity of Kähler-Ricci flows on Fano manifolds. *Acta Math.* 216, 1 (2016), 127–176.

- [170] TIMASHEV, D. A. *Homogeneous spaces and equivariant embeddings*, vol. 138 of *Encyclopedia of Mathematical Sciences*. Springer, Heilderlberg, 2011. Invariant Theory and Algebraic Transformation Groups, 8.
- [171] UHLENBECK, K. Connections with L^p bounds on curvature. *Comm. Math. Phys.* 83 (1982), 31–42.
- [172] UHLENBECK, K. Removable singularities in Yang-Mills fields. *Comm. Math. Phys.* 83 (1982), 11–29.
- [173] UHLENBECK, K., AND YAU, S.-T. On the existence of Hermitian-Yang-Mills connections in stable vector bundles. *Comm. Pure Appl. Math.* 39 (1986), 257–293.
- [174] WANG, X. Height and GIT weight. *Math. Res. Lett.* 19, 4 (2012), 909–926.
- [175] WANG, X.-J., AND ZHU, X. Kähler-Ricci solitons on toric manifolds with positive first Chern class. *Adv. Math.* 188 (2004), 87–103.
- [176] WITT NYSTRÖM, D. Test configurations and Okounkov bodies. *Compos. Math.* 148, 6 (2012), 1736–1756.
- [177] YAU, S.-T. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation I. *Comm. Pure Appl. Math.* 31 (1978), 339–411.
- [178] YAU, S.-T. Open problems in geometry. *Proc. Symposia Pure Math.* 54 (1993), 1–28.
- [179] ZELDITCH, S. Szegő kernel and a theorem of Tian. *Int. Math. Res. Notices* 6 (1998), 317–331.
- [180] ZHANG, S. Heights and reductions of semi-stable varieties. *Compos. Math.* 104, 1 (1996), 77–105.
- [181] ZHANG, Y. Completion of the moduli space for polarized Calabi-Yau manifolds. *J. Differential Geom.* 103, 3 (2016), 521–544.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NOTRE DAME, NOTRE DAME, IN 46556
E-mail address: gszekely@nd.edu