

CONTINUANTS, RUN LENGTHS, AND BARRY'S MODIFIED PASCAL TRIANGLE

JEFFREY SHALLIT AND LUKAS SPIEGELHOFFER

ABSTRACT. We show that the n 'th diagonal sum of Barry's modified Pascal triangle can be described as the continuant of the run lengths of the binary representation of n . We also obtain an explicit description for the row sums.

1. INTRODUCTION

In 2006 in the *On-Line Encyclopedia of Integer Sequences* (OEIS) [8], sequence A119326, Paul Barry introduced a modified Pascal triangle, defined for integers $0 \leq k \leq n$, as follows:

$$T(n, k) = \sum_{\substack{0 \leq j \leq n-k \\ 2|j}} \binom{k}{j} \binom{n-k}{j}.$$

The first few rows of this triangle are as follows:

$$\begin{array}{cccccccc} & & & & & & & 1 \\ & & & & & & 1 & 1 \\ & & & & 1 & 1 & 1 \\ & & 1 & 1 & 1 & 1 \\ & 1 & 1 & 1 & 2 & 1 & 1 \\ & 1 & 1 & 4 & 4 & 1 & 1 \\ & 1 & 1 & 7 & 10 & 7 & 1 & 1 \\ 1 & 1 & 11 & 19 & 19 & 11 & 1 & 1 \\ 1 & 1 & 16 & 31 & 38 & 31 & 16 & 1 & 1 \end{array}$$

2010 *Mathematics Subject Classification*. Primary: 05A10; Secondary: 05A19, 11A55, 11A63.

Key words and phrases. modified Pascal triangle, continued fraction, continuant, Stern's diatomic sequence. The first author is supported by a grant from NSERC..

The second author acknowledges support by project F5502-N26 (FWF), which is a part of the Special Research Program "Quasi Monte Carlo Methods: Theory and Applications". Moreover, the second author thanks the Erwin Schrödinger Institute for Mathematics and Physics for providing the opportunity to carry out research work during his visit for the programme "Tractability of High Dimensional Problems and Discrepancy".

Similarly, one can consider $T(n, k) \bmod 2$, whose terms are given by sequence **A114213**:

$$\begin{array}{cccccccccc}
 & & & & & & & & & & \\
 & & & & & & & & & & 1 \\
 & & & & & & & & & 1 & 1 \\
 & & & & & & & & 1 & 1 & 1 \\
 & & & & & & 1 & 1 & 1 & 1 \\
 & & & 1 & 1 & 0 & 1 & 1 \\
 & 1 & 1 & 0 & 0 & 1 & 1 \\
 & 1 & 1 & 1 & 0 & 1 & 1 & 1 \\
 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\
 1 & 1 & 0 & 1 & 0 & 1 & 0 & 1 & 1
 \end{array}$$

Sequences **A114212** and **A114214**, respectively, are the row sums and diagonal sums of this latter triangle. We denote them by $r(n)$ and $d(n)$, respectively:

$$\begin{aligned}
 r(n) &= \sum_{k=0}^n (T(n, k) \bmod 2) \\
 d(n) &= \sum_{k=0}^{\lfloor n/2 \rfloor} (T(n-k, k) \bmod 2).
 \end{aligned}$$

In May 2016, the first author observed, empirically, a connection between $d(n)$ and the binary representation of n . In this note we prove this connection, and also prove a formula for $r(n)$. The connection involves Stern's "diatomic sequence" $s(n)$, defined by $s(0) = 0$, $s(1) = 1$, $s(2n) = s(n)$, and $s(2n+1) = s(n) + s(n+1)$; see [9].

2. THE DIAGONAL SUMS

Let the binary representation of n be denoted by $\sum_{i=0}^j \varepsilon_i(n) 2^i$. We show that the diagonal sum $d(n)$ can be expressed in terms of this representation. Given a string s of 0's and 1's, we consider its *run lengths*: the lengths of maximal blocks of consecutive identical elements. For example, if $s = 111000011111$, then the run lengths of s are $(3, 4, 5)$.

If m is a sequence of positive integers, we may associate an integer with it via the continued fraction expansion: if $m = (m_0, \dots, m_k)$, we say that the *continuant* of m is the numerator of the continued fraction $[m_0; m_1, \dots, m_k]$ (see [3, Ch. 34, §4]).

Theorem 2.1. *Let $n \geq 0$ be an integer and let m be the sequence of run lengths of the binary representation of n . Then $d(n)$ equals the continuant of m .*

We will use Lucas' famous congruence for binomial coefficients [7, p. 230]: if p is a prime number and $n = (n_\nu \cdots n_0)_p$ and $k = (k_\nu \cdots k_0)_p$, then

$$\binom{n}{k} \equiv \binom{n_\nu}{k_\nu} \cdots \binom{n_0}{k_0} \pmod{p}.$$

This implies that $\binom{n}{k}$ is not divisible by p if and only if $k_i \leq n_i$ for all i . Moreover, it follows that the number of odd binomial coefficients $\binom{n}{k}$ equals $2^{s_2(n)}$, where s_2 is the binary sum-of-digits function [4].

We prove the following statement, which reduces the problem to divisibility by 2 of binomial coefficients. We will derive Theorem 2.1 from it in a moment.

Proposition 2.2. *Let n and k be nonnegative integers such that $k \leq n$. If $2 \mid n+k$, then $T(n, k) \equiv \binom{n}{k} \pmod{2}$. Otherwise, $T(n, k) \equiv \binom{n-1}{k} \pmod{2}$.*

Proof. We prove the first statement. By replacing n with $n + k$ we get the equivalent assertion that if $2 \mid n$ or $2 \mid k$, then

$$(1) \quad \sum_{\substack{0 \leq j \leq n \\ 2 \mid j}} \binom{n}{j} \binom{k}{j} \equiv \binom{n+k}{k} \pmod{2}.$$

By Lucas' congruence the left-hand side is congruent to

$$\sum_{j=0}^n \binom{n}{j} \binom{k}{j} \equiv \sum_{j=0}^n \binom{n \wedge k}{j} \equiv 2^{s_2(n \wedge k)} \pmod{2},$$

where $n \wedge k$ is the integer whose binary digits satisfy $\varepsilon_i(n \wedge k) = \min(\varepsilon_i(n), \varepsilon_i(k))$. This expression is odd if and only if $s_2(n \wedge k) = 0$, which is the case if and only if the binary representations of n and k are disjoint. To handle the right-hand side of Eq. (1), we note that $\binom{n+k}{k}$ is odd if $n \wedge k = 1$. On the other hand, if the binary representations of n and k are not disjoint, then the condition $\varepsilon_i(k) \leq \varepsilon_i(n+k)$ is violated for $i = \min\{j : \varepsilon_j(n) = 1, \varepsilon_j(k) = 1\}$; therefore $\binom{n+k}{k}$ is even. This proves the first assertion.

For the second assertion, we use Lucas' congruence again: for $2 \mid j$ and $2 \mid m$ we have $\binom{m}{j} \equiv \binom{m+1}{j} \pmod{2}$. Since $2 \nmid n - k$, we obtain $\binom{n-k}{j} \equiv \binom{n-1-k}{j} \pmod{2}$. Moreover, by $2 \nmid n - k$ the ranges of summation in $T(n, k)$ and $T(n-1, k)$ are the same. $\square$

From this proposition we obtain in particular the identity

$$(2) \quad d(2n) = d(2n+1).$$

Carlitz [2] proved that Stern's diatomic sequence $s(n)$ satisfies $s(n+1) = \sum_{k=0}^{\lfloor n/2 \rfloor} \left(\binom{n-k}{k} \pmod{2} \right)$. By Proposition 2.2 and Eq. (2) we therefore have

$$(3) \quad d(2n) = d(2n+1) = s(2n+1).$$

It is well-known [5, 6] that if $m = (m_0, \dots, m_k)$ is the sequence of run-lengths of the binary representation of n and n is odd, then $s(n)$ is the continuant of m . Therefore $d(n)$ is the continuant of m . In order to complete the proof of the conjecture, we have to show that the same is true for even n . By Eq. (3) it is sufficient to prove the following lemma.

Lemma 2.3. *If n is even, then the continuant of the sequence of run-lengths of the binary representation of n is equal to the continuant corresponding to $n+1$.*

Proof. Let $n = 1^{m_0} 0^{m_1} \dots 1^{m_{k-1}} 0^{m_k}$. We distinguish between two cases. If $m_k = 1$, then $n+1 = (1^{m_0} 0^{m_1} \dots 0^{m_{k-2}} 1^{m_{k-1}+1})$ and the statement follows from the identity $[m_0; m_1, \dots, m_{k-1}, 1] = [m_0; m_1, \dots, m_{k-1} + 1]$. If $m_k \geq 2$, then $n+1 = (1^{m_0} 0^{m_1} \dots 0^{m_{k-2}} 1^{m_{k-1}} 0^{m_k-1} 1)$ and the statement follows from $[m_0; m_1, \dots, m_k] = [m_0; m_1, \dots, m_{k-1}, m_k - 1, 1]$. $\square$

Remark. The sequence $(d(n))_{n \geq 0}$ is a 2-regular sequence [1], as it satisfies the equalities

$$\begin{aligned} d(2n+1) &= d(2n) \\ d(4n+2) &= 3d(2n) - d(4n) \\ d(8n) &= -d(2n) + 2d(4n) \\ d(8n+4) &= 4d(2n) - d(4n). \end{aligned}$$

3. THE ROW SUMS

We will prove

Theorem 3.1.

$$r(n) = \begin{cases} 2^{s_2(n)}, & \text{if } n \text{ odd;} \\ 2^{s_2(n)} + 2^{s_2(n-2)}, & \text{if } n \text{ even.} \end{cases}$$

A similar characterization was stated, without proof or attribution, in the notes to **A114212** of the OEIS.

Proof. From Proposition 2.2 we get, for integers $n \geq k \geq 0$, that

$$\begin{aligned} T(2n, 2k) &\equiv T(2n+1, 2k) \equiv T(2n+1, 2k+1) \equiv \binom{n}{k} \pmod{2}; \\ T(2n, 2k+1) &\equiv \binom{n-1}{k} \pmod{2}. \end{aligned}$$

Then

$$\begin{aligned} r(2m) &= \sum_{k=0}^{2m} (T(2m, k) \pmod{2}) \\ &= \sum_{k=0}^m (T(2m, 2k) \pmod{2}) + \sum_{k=0}^{m-1} (T(2m, 2k+1) \pmod{2}) \\ &= \sum_{k=0}^m \left(\binom{m}{k} \pmod{2} \right) + \sum_{k=0}^{m-1} \left(\binom{m-1}{k} \pmod{2} \right) \\ &= 2^{s_2(m)} + 2^{s_2(m-1)} \\ &= 2^{s_2(2m)} + 2^{s_2(2m-2)}. \end{aligned}$$

Similarly,

$$\begin{aligned} r(2m+1) &= \sum_{k=0}^{2m+1} (T(2m+1, k) \pmod{2}) \\ &= \sum_{k=0}^m (T(2m+1, 2k) \pmod{2}) + \sum_{k=0}^m (T(2m+1, 2k+1) \pmod{2}) \\ &= \sum_{k=0}^m \left(\binom{m}{k} \pmod{2} \right) + \sum_{k=0}^m \left(\binom{m}{k} \pmod{2} \right) \\ &= 2^{s_2(m)} + 2^{s_2(m)} \\ &= 2^{s_2(2m+1)}. \end{aligned}$$

□

REFERENCES

- [1] J.-P. ALLOUCHE AND J. O. SHALLIT, *The ring of k -regular sequences*, Theoret. Comput. Sci., 98 (1992), pp. 163–197.
- [2] L. CARLITZ, *Single variable Bell polynomials*, Collect. Math., 14 (1962), pp. 13–25.
- [3] G. CHRYSTAL, *Algebra*, vol. 2, Adam and Charles Black, 1900.
- [4] J. W. L. GLAISHER, *On the residue of a binomial-theorem coefficient with respect to a prime modulus*, Quarterly J. Pure Appl. Math., 30 (1899), pp. 150–156.
- [5] D. H. LEHMER, *On Stern's diatomic series*, Amer. Math. Monthly, 36 (1929), pp. 59–67.

- [6] D. A. LIND, *An extension of Stern's diatomic series*, Duke Math. J., 36 (1969), pp. 55–60.
- [7] E. LUCAS, *Théorie des fonctions numériques simplement périodiques*, Amer. J. Math., 1 (1878), pp. 197–240.
- [8] N. J. A. SLOANE, *The On-Line Encyclopedia of Integer Sequences*, 2017. Published electronically at <https://oeis.org>.
- [9] M. A. STERN, *Über eine zahlentheoretische Funktion*, J. reine angew. Math., 55 (1858), pp. 193–220.

SCHOOL OF COMPUTER SCIENCE, UNIVERSITY OF WATERLOO, WATERLOO, ON N2L 3G1, CANADA
E-mail address: `shallit@cs.uwaterloo.ca`

INSTITUT FÜR DISKRETE MATHEMATIK UND GEOMETRIE, TECHNISCHE UNIVERSITÄT WIEN, WIEDNER HAUPT-
STRASSE 8–10, 1040 WIEN, AUSTRIA
E-mail address: `lukas.spiegelhofer@tuwien.ac.at`