arXiv:1710.06355v1 [math.PR] 17 Oct 2017

Spectra of Wishart Matrices with size-dependent entries.

Nathan Noiry*

Abstract

We prove the convergence of the empirical spectral measure of Wishart matrices with size-dependent entries and characterize the limiting law by its moments. We apply our result to the cases where the entries are Bernoulli variables with parameter c/n or truncated heavy-tailed random variables. In both cases, when c goes to infinity or when the truncation is small, the limiting spectrum is a perturbation of the Marchenko-Pastur distribution and we compute its leading term.

MSC 2010 Classification: 05C80; 60B20.
Keywords: Wishart matrices; Marchenko-Pastur distribution; Erdös-Rényi bipartite random graphs; heavy tailed random variables.

1 Introduction

Let X_n be a real random matrix of size $n \times m$ with i.i.d. entries. We define the Wishart matrix $W_n = \frac{1}{n} X_n X_n^T$, where X_n^T is the transpose of X_n. The spectral measure of W_n is the random probability law:

$$\mu_{W_n} = \frac{1}{n} \sum_{\lambda \in \mathrm{Spec}(W_n)} \delta_\lambda,$$

where $\mathrm{Spec}(W_n)$ is the spectrum of W_n and δ_λ the Dirac at λ. Since W_n is a positive symmetric matrix, its eigenvalues are nonnegative reals. The work of Marchenko and Pastur [11] implies that, when the entries have variance equal to 1 and finite moments of all order. Then, almost surely, μ_{W_n} weakly converges to a probability law μ_α as $n, m \to +\infty$ and $m/n \to \alpha > 0$. The law μ_α is given by:

$$\mu_\alpha(\mathrm{d}x) = \frac{\sqrt{(b-x)(x-a)}}{2\pi x} \mathrm{d}x + \mathbf{1}_{\alpha<1} (1-\alpha)\, \delta_0(\mathrm{d}x),$$

where $a = (1-\sqrt{\alpha})^2$ and $b = (1+\sqrt{\alpha})^2$.

The main issue of this paper is to let the law of the entries of X_n depend on n. Informally, our first result (Theorem 1) states that in that case, under some moment conditions, the measures μ_{W_n} converge weakly to a probability law which is characterized by its moments, for which we provide a formula. This is an analog for Wishart matrices of a result obtained by

*nathan.noiry@parisnanterre.fr

Zakharevich for Wigner matrices in [14]. The method here is based on a proof of the convergence of all the moments of the spectral measures μ_{W_n}. The k-th limiting moment will write:

$$\sum_{a=1}^{k}\sum_{l=1}^{a}\alpha^l \sum_{\substack{\mathbf{b}=(b_1,\dots,b_a)\\ b_1\geq b_2\geq\dots\geq b_a\geq 2\\ b_1+b_2+\cdots+b_a=2k}} |\mathscr{W}_k(a,a+1,l,\mathbf{b})|\prod_{i=1}^{a}A_{b_i}.$$

The set $\mathscr{W}_k(a,a+1,l,\mathbf{b})$ is a combinatorial object linked with closed words on planar rooted trees and encodes the combinatorics of moments. We give a precise definition in Section 2. Interestingly, the A_i's coefficients, given by the formula (1), are the only reminders of the laws of the entries of the matrices.

The convergence of the spectral measure was already proved in [3] by Benaych-Georges and Cabanal-Duvillard, using different arguments. See also Male in [10] for related work. However, the main advantage of our approach is the explicit formula we obtain for the moments, which is more amenable to analysis, as we will see in Sections 2 and 3.

In [13], Vengerovsky treated the particular case of diluted matrices $X_n(i,j)=a(i,j)d_n(i,j)$ where the $a(i,j)$'s are i.i.d. centered random variables and the $d_n(i,j)$'s are i.d.d. with Bernoulli law of parameter c/n. He derived a formula for the limiting moments, in terms of combinatorial quantities that admit a recursive formula.

In the second part of the paper, we will focus on this particular case and let the entries of X_n be i.i.d. Bernoulli laws with parameter c/n. In this setting, the Wishart matrices can be easily linked with the adjacency matrix of a bipartite random graph which admits a limit for the local weak topology. This convergence can be used to prove the convergence of the resolvent of the bipartite graph and therefore of μ_{W_n} itself, as explained in [6] by Bordenave and Lelarge. This is the content of Theorem 2. The limiting spectral measure $\mu_{\alpha,c}$ depends only on α and c and converges to the law μ_α as $c\to\infty$. In Theorem 3, we describe how $\mu_{\alpha,c}$ differs from its limit μ_α by giving an asymptotic expansion in $1/c$ of its moments. More precisely, we will obtain that, in the sense of moments convergence:

$$c\big(\mu_{\alpha,c}-\mu_\alpha\big)\underset{c\to+\infty}{\longrightarrow}\mu_\alpha^{(1)},$$

where $\mu_\alpha^{(1)}$ is a signed measure of total mass zero, see Theorem 3. The proof, based on a more careful analysis of the moment formula obtained in 1, is inspired by the computations made in [8] by Enriquez and Ménard. A natural extension would be to prove that the convergence holds in the sense of weak convergence, but it should involved new techniques since the moments of a signed measure of total mass zero do not characterized it. See Figure 5 for numerical simulations.

In the last part of this paper, we apply our results to heavy tailed random matrices. In that case, the entries of X_n do not have finite moments of all order so that our main result does not apply. Instead, we truncate the entries at a constant $B>0$ times the largest n-th quantile of the corresponding law. By Theorem 1, the spectral measures associated to the truncated random matrices converges to a deterministic probability law. The moments of this limiting law admit an asymptotic expansion involving the measures μ_α and $\mu_\alpha^{(1)}$, as $B\to 0$. See Theorem 4.




Figure 1: Numerical simulations for the spectrum of 100 Wishart matrices associated to random matrices of size $n \times \alpha n$ with i.i.d. entries with Bernoulli law of parameter c/n, with $c = 20$ and $n = 3000$. The theoritical densities of μ_α and $\mu_\alpha^{(1)}$ are drawn in blue. The top diagrams correspond to $\alpha = 2$ whereas the bottom diagrams correspond to $\alpha = 4$.

2 A generalized Marchenko-Pastur theorem

Let $\mathscr{P} = \{P_n\}_{n\geq 1}$ be a family of probability laws on $\mathbf{R}$ which have zero mean. For all $n \geq 1$, let $X_n = (X_n(i,j))_{1\leq i,j\leq n}$ be a random $n \times m$ matrix with i.i.d. entries with law P_n. We will make the hypothesis that the ratio m/n converges to a real $\alpha > 0$ and that for all $k \geq 1$, the following limit exists and is finite:

$$A_k := \lim_{n\to+\infty} \frac{M_k(P_n)}{n^{k/2-1} M_2(P_n)^{k/2}}, \tag{1}$$

where $M_k(P_n)$ is the k-th moment of P_n. Denote by $\mathscr{A}$ the sequence formed by the A_k's. We are interested in the behavior of the spectral measures of the sequence of random matrices

$$W_n := \frac{1}{nM_2(P_n)} X_n X_n^T.$$

In order to properly state our first result, we need to introduce the notion of word on a labeled graph. A labeled graph is a graph $\mathrm{G} = (\mathrm{V}, \mathrm{E})$ together with a labeling of the vertices, that is a one-to-one application from V to $\{1, \ldots, |\mathrm{V}|\}$. A relabeling of a labeled graph is a new

choice of bijection between V and $\{1,\dots,|\mathrm{V}|\}$. Note that there are $|\mathrm{V}|!$ choices of labelings for a given graph $\mathrm{G}=(\mathrm{V},\mathrm{E})$. A word of length $k\geq 1$ on a labeled graph G is a sequence of labels $i_1,i_2,\dots,i_k$ such that $\{i_j,i_{j+1}\}$ is a pair of adjacent labels (that is the associated vertices are neighbours in G) for all $1\leq j\leq k-1$. A word of length k is said to be closed if $i_1=i_k$. Let $\mathbf{i}=i_1,\dots,i_k$ and $\mathbf{i}'=i'_1,\dots,i'_k$ be two words of length k on two labeled graphs G and G$'$ having the same number of vertices. Then, $\mathbf{i}$ and $\mathbf{i}'$ are said to be equivalent if there exists a bijection σ of $\{1,\dots,|\mathrm{V}|\}$ such that $\sigma(i_j)=i'_j$ for all $1\leq j\leq k$. In words, $\mathbf{i}$ and $\mathbf{i}'$ are equivalents if there exists a relabeling of a G such that the word associated to $\mathbf{i}$ is exactly $\mathbf{i}'$. One can check that this defines an equivalence relation on words on labeled graphs.

Recall that a planar rooted tree is a connected graph without loop embedded in the plane, with a distinguished vertex called the root. A vertex at odd (resp. even) distance from the root will be called an odd (resp. even) vertex. An edge with an odd (resp. even) origin vertex will be called an odd (resp. even) edge.

Theorem 1. *Suppose that for some $\gamma>0$, $A_k=O(\gamma^k)$ as $k\to+\infty$. Then there exists a probability law $\mu_{\mathscr{A},\alpha}$ depending only on $\mathscr{A}$ and α, such that μ_{W_n} converges weakly to $\mu_{\mathscr{A},\alpha}$ in probability: for all $\varepsilon>0$ and all bounded continuous function $f:\mathbf{R}\to\mathbf{R}$,*

$$\mathbf{P}\left(\left|\int_{\mathbf{R}} f\mathrm{d}\mu_{W_n}-\int_{\mathbf{R}} f\mathrm{d}\mu_{\mathscr{A},\alpha}\right|>\varepsilon\right)\underset{n\to+\infty}{\longrightarrow}0.$$

Moreover, the measure $\mu_{\mathscr{A},\alpha}$ is characterized by its sequence of moments:

$$M_k(\mu_{\mathscr{A},\alpha})=\sum_{a=1}^{k}\sum_{l=1}^{a}\alpha^l\sum_{\substack{\mathbf{b}=(b_1,\dots,b_a)\\ b_1\geq b_2\geq\dots\geq b_a\geq 2\\ b_1+b_2+\dots+b_a=2k}}|\mathscr{W}_k(a,a+1,l,\mathbf{b})|\prod_{i=1}^{a}A_{b_i},\tag{2}$$

where $\mathscr{W}_k(a,a+1,l,\mathbf{b})$ is a set of representatives of the equivalence classes of closed words on labeled rooted planar trees having "a" edges, of which l are odd edges, starting from the root and such that for all $1\leq i\leq a$, one edge is browsed b_i times.

Remark 1. *The theorem can be thought as a universality result. Namely, if two sequences of probability law P_n and P'_n have the same asymptotic $\mathscr{A}$, the limiting spectral measures of W_n and W'_n are the same (in probability).*

Corollary 1. *If for all $k>2$, $A_k=0$, the measures $\mu_{\mathscr{A},\alpha}$ and μ_α coincide. For example, this is the case when the laws P_n are all equal.*

As the statement suggests, we are going to prove the result by the method of moments. Classically, we start with a computation of the average moments of μ_{W_n}. For $k\geq 1$, we can write:

$$\mathbf{E}M_k(\mu_{W_n})=\frac{1}{n^{k+1}M_2(P_n)^k}\sum_{\substack{1\leq i_1,\dots,i_k\leq n\\ 1\leq j_1,\dots j_k\leq m}}\mathbf{E}[X(i_1,j_1)X(i_2,j_1)\cdots X(i_k,j_k)X(i_1,j_k)].\tag{3}$$

Denote $(\mathbf{i},\mathbf{j})$ the generic word $i_1j_1i_2\dots i_1j_k$ appearing in (3). We define the bipartite graph $\mathrm{G}=(\mathrm{V},\mathrm{E})$ associated to the word $(\mathbf{i},\mathbf{j})$ by:

$$\mathrm{V}=\{(i_r,\mathbf{i}),(j_r,\mathbf{j});\,1\leq r\leq k\}\quad\text{and}\quad\mathrm{E}=\big\{\{(i_r,\mathbf{i}),(j_r,\mathbf{j})\},\{(i_{r+1},\mathbf{i}),(j_r,\mathbf{j})\};\,1\leq r\leq k\big\},$$

where we used the convention $k+1=1$. The abstract symbols $\mathbf{i}$ and $\mathbf{j}$ are needed to obtain a bipartite graph since the i_r's and j_r's can have common values (see Figure 2 for illustration). We will refer to $\mathbf{i}$ and $\mathbf{j}$ letters. In words, the vertices of G are the letters of the word $(\mathbf{i},\mathbf{j})$ and two vertices are linked by an edge when they are consecutive in $(\mathbf{i},\mathbf{j})$. Denote by s the number of vertices, a the number of edges, l the number of $\mathbf{j}$-vertices and $\bar{l}$ the number of $\mathbf{i}$-vertices in the word. Since G is connected, $s \leq a+1$. Moreover, since P_n has zero mean, each edge must appear at least twice in the word to give a non-zero contribution in (3). As a consequence we obtain the bound $a \leq k$ because $i_1 j_1 \dots j_k$ possesses $2k$ edges counted with multiplicity.




Figure 2: Example of a word $(\mathbf{i},\mathbf{j})$ with its associated graph and quantities.

Two words $(\mathbf{i},\mathbf{j})$ and $(\mathbf{i}',\mathbf{j}')$ are said equivalent if one can find a permutation σ of $\{1,\dots,n\}$ and another one τ of $\{1,\dots,m\}$ such that

$$\forall p \in \{1,\dots k\}, \quad \sigma(i_p) = i'_p \text{ and } \tau(j_p) = j'_p.$$

One can check that this is an equivalence relation on the words appearing in (3). Note that $(\mathbf{i},\mathbf{j})$ has

$$C(s,l) = n(n-1)\cdots(n-l+s+1) \times m(m-1)\cdots(m-l+1) \sim \alpha^l n^s$$

equivalents. Fix $a \in \{1,\dots,k\}$, $1 \leq s \leq a+1$ and $1 \leq l \leq a$. Let $\mathscr{B}_{a,k}$ be the set of a-tuples $\mathbf{b} = (b_1,\dots,b_a)$ of integers such that

1. $b_1 \geq b_2 \geq \cdots \geq b_a \geq 2$;
2. $b_1 + \cdots + b_a = 2k$.

For all $k \geq 1$ and $\mathbf{b} \in \mathscr{B}_{a,k}$, we introduce $\mathscr{W}_k(a,s,l,\mathbf{b})$ a set of representatives of the equivalence classes of words $(\mathbf{i},\mathbf{j})$ such that the associated graph has a edges, s vertices of which l are $\mathbf{j}$-vertices and such that for all $1 \leq i \leq a$ there is an edge in E which has multiplicity b_i in $(\mathbf{i},\mathbf{j})$. We can rewrite (3) as:

$$\sum_{a=1}^{k} \sum_{s=1}^{a+1} \sum_{l=1}^{s} \frac{C(s,l)}{n^{a+1}} \sum_{b \in \mathscr{B}_{a,k}} \sum_{(\mathbf{i},\mathbf{j}) \in \mathscr{W}(a,s,l,b)} \prod_{1 \leq i \leq a} \frac{M_{b_i}(P_n)}{n^{b_i/2-1} M_2(P_n)^{b_i/2}}. \tag{4}$$

From this equation we easily deduce the form of the limiting moments:

Lemma 1. *An asymptotic contribution arises only if $s = a+1$ that is when the graph associated to $(\mathbf{i},\mathbf{j})$ is a tree. More precisely the limit of* (4) *when $n \to +\infty$ is*

$$M_k := \lim_{n \to +\infty} \mathbf{E} M_k(\mu_{W_n}) = \sum_{a=1}^{k} \sum_{l=1}^{a} \alpha^l \sum_{b \in \mathscr{B}_{a,k}} |\mathscr{W}_k(a,a+1,l,b)| \prod_{i=1}^{a} A_{b_i}. \tag{5}$$

Proof. Since $C(s,l)n^{-a-1} \sim \alpha^l n^{s-a-1}$ when $n \to +\infty$ we deduce that when $s < a+1$ the asymptotic contribution is zero. Hence a possible non-zero contribution arises only when $s = a+1$. The formula is a consequence of (1). □

Remark 2. *The only non-zero contributions arise when* $(\mathbf{i},\mathbf{j})$ *is a walk on a tree that browses every edge and starts and finishes at the same vertex. Therefore each edge must be visited an even number of time: each* b_i *in the tuple* $\mathbf{b}$ *is even.*

Remark 3. *The set* $\mathscr{W}_k(a,a+1,l,\mathbf{b})$ *is also the set of closed words on rooted planar trees having a edges out of which* l *are odd edges, starting from the root and such that for all* $1 \le i \le a$*, one edge is browsed* b_i *times. Notice that the number of* $\mathbf{j}$*-vertices is equal to the number of vertices in odd generations.*

In view of Theorem 1 we have to prove that $M_k(\mu_{W_n})$ concentrates around its mean. Since we are looking for a convergence in probability, it is sufficient to show that its variance vanishes when n tends to infinity.

Lemma 2. *For all* $k \ge 1$, $\mathbf{Var}(M_k(P_n)) = O(n^{-1})$*. In particular* $M_k(P_n)$ *converges to* M_k *in probability.*

Proof. Let $k \ge 1$. We can write $\mathbf{E}[M_k(P_n)^2] - \mathbf{E}[M_k(P_n)]^2$ as

$$\frac{1}{n^{2(k+1)}M_2(P_n)^{2k}} \sum_{(\mathbf{i},\mathbf{j}),(\mathbf{i}',\mathbf{j}')} \Big(\mathbf{E}[P(\mathbf{i},\mathbf{j})P(\mathbf{i}',\mathbf{j}')] - \mathbf{E}[P(\mathbf{i},\mathbf{j})]\mathbf{E}[P(\mathbf{i}',\mathbf{j}')]\Big), \tag{6}$$

where $P(\mathbf{i},\mathbf{j})$ is the product $X(i_1,j_1)X(i_2,j_1)\cdots X(i_1,j_k)$. We note G (resp. G$'$) the graph associated to $(\mathbf{i},\mathbf{j})$ (resp. $(\mathbf{i}',\mathbf{j}')$), the corresponding quantities such as s and s' being defined as before. We also consider the graph $\mathrm{G}\cup\mathrm{G}'$ associated to $i_1j_1\ldots i_1j_ki'_1j'_1\ldots i'_1j'_k$, and introduce S its number of vertices, A its number of edges and L its number of $\mathbf{j}$-vertices and $\mathbf{j}'$-vertices. Note that if G and G$'$ have no edge in common, then the contribution is zero by independence of $P(\mathbf{i},\mathbf{j})$ and $P(\mathbf{i}',\mathbf{j}')$. We can therefore restrict the sum to pairs of words $\big((\mathbf{i},\mathbf{j}),(\mathbf{i}',\mathbf{j}')\big)$ sharing at least an edge. In this case $\mathrm{G}\cup\mathrm{G}'$ is connected, hence $A \ge S-1$. Moreover, each edge must appear at least twice otherwise the contribution is zero since P_n has mean zero. Therefore, the sum (6) rewrites

$$\frac{1}{n^{2(k+1)}M_2(P_n)^{2k}} \sum_{S=1}^{2k+1} \sum_{L=1}^{S} \sum_{A=S-1}^{2k} C(S,L) \times \sum_{\mathbf{B}\in\mathscr{B}_{A,2k}} \sum_{(\mathbf{i},\mathbf{j}),(\mathbf{i}',\mathbf{j}')\in\mathscr{W}_k(A,S,L,\mathbf{B})} \Big(\mathbf{E}[P(\mathbf{i},\mathbf{j})P(\mathbf{i}',\mathbf{j}')] - \mathbf{E}[P(\mathbf{i},\mathbf{j})]\mathbf{E}[P(\mathbf{i}',\mathbf{j}')]\Big). \tag{7}$$

Fix a generic couple $\big((\mathbf{i},\mathbf{j}),(\mathbf{i}',\mathbf{j}')\big)$. Let e be an edge of $\mathrm{G}\cup\mathrm{G}'$. The corresponding A-tuple $\mathbf{B} \in \mathscr{B}_{A,2k}$ possesses a coefficient B_i such that e has multiplicity B_i. Note b_i (resp. b'_i) the multiplicity of e in $(\mathbf{i},\mathbf{j})$ (resp. $(\mathbf{i}',\mathbf{j}')$). We have the relation $b_i + b'_i = B_i$. The contribution of this generic couple in (7) is therefore

$$\frac{C(S,L)}{n^{A+2}} \times \left(\frac{\mathbf{E}[P(\mathbf{i},\mathbf{j})P(\mathbf{i}',\mathbf{j}')]}{\prod_{1\le i\le A} n^{B_i/2-1}M_2^{B_i/2}} - \frac{\mathbf{E}[P(\mathbf{i},\mathbf{j})]}{\prod_{1\le i\le A} n^{b_i/2-1}M_2^{b_i/2}} \times \frac{\mathbf{E}[P(\mathbf{i}',\mathbf{j}')]}{\prod_{1\le i\le A} n^{b'_i/2-1}M_2^{b'_i/2}} \right).$$

By assumption (1), the absolute value of the difference inside the parentheses is bounded. This gives the conclusion since $C(S,L) \sim \alpha^L n^S$ and $S \leq A+1$. □

In order to obtain Theorem 1, it remains to show that the sequence $\{M_k\}_{k\geq 1}$ entirely determines a probability law. To that aim, it is enough to prove that M_k does not grow faster than k^{ck} for some positive constant c. First, remark that

$$|\mathscr{W}_k(a,a+1,l,\mathbf{b})| \leq \frac{(2k)^k}{k+1}\binom{2k}{k}.$$

Indeed there are $\frac{1}{a+1}\binom{2a}{a}$ rooted planar trees having a edges. Moreover, two elements $(\mathbf{i},\mathbf{j})$ and $(\mathbf{i}',\mathbf{j}')$ in $\mathscr{W}(a,a+1,l,b)$ inducing the same tree differ only by the order in which each edge is browsed in the reading of $(\mathbf{i},\mathbf{j})$ (resp. $(\mathbf{i}',\mathbf{j}')$). For a fixed multiplicity b_i and its associated edge e, there are at most $\binom{2k}{b_i}$ different possibilities to place the occurrences of e because a word has $2k$ edges counted with multiplicity. Therefore the number of $(\mathbf{i},\mathbf{j}) \in \mathscr{W}_k(a,s,l,\mathbf{b})$ associated to a fixed tree is bounded by

$$\prod_{1\leq i\leq a}\binom{2k}{b_i} \sim \prod_{1\leq i\leq a}\frac{(2k)^{b_i}}{b_i!} \leq (2k)^{2k}.$$

Using formula (5) and the assumption $A_i = O(\gamma^i)$, we obtain the estimation

$$M_k = O\left(\gamma^k\alpha^k k(2k)^{2k+1}\sum_{a=1}^{k}|\mathscr{B}_{a,k}|\right).$$

Finally the cardinality of $\mathscr{B}_{a,k}$ is bounded by the cardinality of the number of unsorted partitions of the integer $2k$. This last quantity is equal to

$$\sum_{i=1}^{2k}\binom{2k-1}{i-1} = 2^{2k-1},$$

where we summed over the number of partitions of $2k$ in i parts. As a result $M_k = O(k^{ck})$ for some constant $c > 0$. This concludes the proof of Theorem 1.

The proof of the almost sure convergence of μ_{W_n} would require a concentration result analogous for instance to [5, lemma 4.18]. Rather than proving this kind of result, which would be a technical task, we present an alternative approach, specific to the case where the entries have Bernoulli law, borrowed from Bordenave and Lelarge's paper [6].

3 The Bernoulli case

In this section, we study the particular case where P_n is the the centered Bernoulli law of parameter c/n, c being a positive number, that is:

$$P_n\left(-\frac{c}{n}\right) = 1-\frac{c}{n} \quad \text{and} \quad P_n\left(1-\frac{c}{n}\right) = \frac{c}{n}.$$

In this case, since the second moment of P_n verifies $M_2(P_n) = c/n + o(c/n)$ as $n \to \infty$, we set

$$W_n = \frac{1}{c}X_n X_n^T$$

to simplify notations. We first give another proof for the convergence of the μ_{W_n}, thanks to an interpretation of the hermitization of X as the adjacency matrix of a random bipartite graph $G_{n,m}$. This makes possible the use of the results of Bordenave and Lelarge in [6] after identifying the local limit of $G_{n,m}$. In a second part, we give an asymptotic expansion in $1/c$ for the moments of the limiting spectral measure, inspired by Enriquez and Ménard (see [8]).

3.1 Another proof of the convergence

To obtain the almost sure convergence of μ_{W_n}, we will rather study the convergence of $W_n' = c^{-1}A_nA_n^T$, where A_n is an $n \times m$ matrix having i.i.d. entries with (non-centered) Bernoulli law of parameter c/n. It is indeed sufficient because, denoting respectively F and F' the cumulative distribution functions of μ_{W_n} and $\mu_{W_n'}$, a consequence of Lidskii's inequalities is that:

$$||F - F'||_\infty \leq \frac{\mathrm{rk}(X_n - A_n)}{n},$$

where rk is the rank operator. As announced before, we have the following theorem.

Theorem 2. *There exists a probability law $\mu_{\alpha,c}$ depending only on α and c such that, almost surely, $\mu_{W_n'}$ converges weakly to $\mu_{\alpha,c}$. Hence, μ_{W_n} converges weakly to $\mu_{\alpha,c}$.*

Remark 4. *It can be proved that the set of atoms of $\mu_{\alpha,c}$ is dense in $\mathbf{R}_+$. More precisely, it is the image by $x \mapsto x^2$ of the set of totally real algebraic integers, which coincides with the set of eigenvalues of finite trees as proved in [12] by Salez. Besides, a consequence of the results of Bordenave, Sen and Virag in [7] is that $\mu_{\alpha,c}$ possesses a continuous part if and only if $c > 1$.*

Define the hermitization of A_n as the hermitian matrix:

$$H(A_n) = \begin{pmatrix} 0 & A_n \\ A_n^T & 0 \end{pmatrix}. \tag{8}$$

Remark that the spectrum of $H(A_n)$ is $\{\pm\sqrt{\lambda_i(A_nA_n^T)}\}_{1\leq i\leq n}$. Let f be the bijection of $\mathbf{R}_+$: $f(x) = x^2$. For a measure ν on $\mathbf{R}_+$ we define $\mathrm{Sym}(\nu)(\cdot) = (\nu(\cdot) + \nu(-\cdot))/2$ the symmetrized version of ν. Then

$$\mu_{H(A_n)} = (\mathrm{Sym} \circ f_*)\mu_{A_nA_n^T},$$

where $f_*\nu$ is the pushforward of a measure ν by f. Since $\mathrm{Sym} \circ f_*$ defines a bijection between the measures which are supported on $\mathbf{R}_+$ and the symmetric measures on $\mathbf{R}$, it suffices to show the convergence of $\mu_{H(A_n)}$ to obtain Theorem 2. To avoid some unpleasant confusions, we will add an apostrophe to the asymptotic measures involved in the proof.

Now, $H(A_n)$ can be interpreted as the adjacency matrix of a random bipartite graph. Let $K_{n,m}$ be the complete bipartite graph with n and m vertices of each color. The vertices of $K_{n,m}$ will be denoted $1,\dots,n+m$, two of them being linked by an edge if and only if one belongs to $\{1,\dots,n\}$ and the other to $\{n+1,\dots,n+m\}$. Perform a Bernoulli percolation with parameter c/n on $K_{n,m}$: keep (independently) each edge with probability c/n and remove it with probability $1 - c/n$. We denote by $G_{n,m}$ the resulting random graph. The adjacency matrix of $G_{n,m}$ has the same law as $H(A_n)$. In the setting of local convergence introduced by Benjamini and Schramm [4] and Aldous and Steele [1], $G_{n,m}$ converges in law to a random tree $T_{\alpha,c}$ for the local topology. To give a precise statement, we give some definitions in what follows.

For any connected, locally finite graph G and any vertex $v \in \mathrm{G}$ we will note (G, v) the class of pointed graphs isomorphic to the graph G pointed in v. For any $r \geq 0$, $[\mathrm{G}, v]_r$ will denote the ball of radius r around v in G for the graph distance. This induces a topology (called the local topology) on the set $\mathscr{G}^*$ of pointed graphs (up to isomorphism) which are locally finite and connected, making it a separable and complete space.

For all nonnegative real number x let $\mathscr{P}(x)$ denote the Poisson law with parameter x. Let $\mathrm{T}_{\alpha,c,1}$ be the random tree where each individual reproduces independently from each other and such that individuals of an even and an odd generation reproduce respectively according to the laws $\mathscr{P}(c)$ and $\mathscr{P}(\alpha c)$. Let $\mathrm{T}_{\alpha,c,2}$ be the random tree where each individual reproduces independently from each other and such that individuals of an even and an odd generation reproduce respectively according to the laws $\mathscr{P}(\alpha c)$ and $\mathscr{P}(c)$. Notice that $\mathrm{T}_{\alpha,c,2}$ has the same law as the random tree issued from a children of the root of $\mathrm{T}_{\alpha,c,1}$.




Figure 3: The recursive relation between $\mathrm{T}_{\alpha,c,1}$ and $\mathrm{T}_{\alpha,c,2}$. Here N_1 and N_2 are independent random variables with law $\mathscr{P}(c)$ and $\mathscr{P}(\alpha c)$; and the $\mathrm{T}^{(i)}_{\alpha,c,1}$ (resp. the $\mathrm{T}^{(i)}_{\alpha,c,2}$) i.i.d. copies of $\mathrm{T}_{\alpha,c,1}$ (resp. $\mathrm{T}_{\alpha,c,2}$).

Let B be Bernoulli random variable of parameter $1/(\alpha+1)$ independent of $\mathrm{T}_{\alpha,c,1}$ and $\mathrm{T}_{\alpha,c,2}$. We define $\mu'_{\alpha,c}$ as the law of the random tree $\mathrm{T}_{\alpha,c} := \mathbf{1}_{B=1}\mathrm{T}_{\alpha,c,1} + \mathbf{1}_{B=0}\mathrm{T}_{\alpha,c,2}$. Let o be a uniformly distributed vertex on $\mathrm{G}_{n,m}$. We define the random probability measure on $\mathscr{G}^*$

$$U_o(\mathrm{G}_{n,m}) := \delta_{(\mathrm{G}_{n,m}(o),o)} = \frac{1}{n+m}\sum_{i=1}^{n+m} \delta_{(\mathrm{G}_{n,m}(i),i)}.$$

Integrating with respect to the randomness of $\mathrm{G}_{n,m}$ gives a new measure $\mathbf{E}[U_o(\mathrm{G}_{n,m})]$ which is characterized by the relation $\mathbf{E}[U_o(\mathrm{G}_{n,m})](A) = \mathbf{P}((\mathrm{G}_{n,m}(o),o) \in A)$ for all measurable set $A \in \mathscr{B}(\mathscr{G}^*)$.

Proposition 1. *The deterministic probability measure $\mathbf{E}[U_o(G_{n,m})]$ converges weakly to $\mu'_{\alpha,c}$. Moreover, if o_1 and o_2 are two independent copies of o, the product $\mathbf{E}[U_{o_1}(G_{n,m})] \otimes \mathbf{E}[U_{o_2}(G_{n,m})]$ converges weakly to $\mu'_{\alpha,c} \otimes \mu'_{\alpha,c}$.*

Proof. The first part is a combinatorial argument that shows that $\mathbf{P}([\mathrm{G}_{n,m}, 1]_r \equiv t)$ converges to $\mathbf{P}([\mathrm{T}_{\alpha,c,1}, \rho]_r \equiv t)$ as $n \to +\infty$ for all $r \geq 1$ and all rooted planar tree t of depth r. For the second part, it suffices to remark that to independent uniform vertices o_1 and o_2 are almost surely at distance greater than r as $n \to +\infty$, for any $r \geq 1$. □

Let us discuss the consequences of this proposition. It implies the validity of the main assumptions of the convergence theorem of Bordenave and Lelarge [6, theorem 5], relative to

the empirical spectral measure of the adjacency matrix of large graphs having a local limit. What remains to check is the uniform integrability of the sequence of degrees $\{\deg_{\mathrm{G}_{n,m}}(o)\}_{n\geq 1}$, which holds. The existence of a probability law μ_c' such that almost surely $\mu_{H(A_n)}$ converges weakly to μ_c' is then a direct application of a result of Bordenave and Lelarge [6, theorem 5]. We also get a description of the Stieltjes transform of μ_c'. Indeed [6] shows that there exists a unique pair of probability laws $(\mathscr{L}_1, \mathscr{L}_2)$ on the set of analytic functions on $\mathbf{C}_+ = \{z \in \mathbf{C} : \mathrm{Im}(z) > 0\}$ such that for all $z \in \mathbf{C}_+$:

$$\begin{cases} X_1(z) \overset{(d)}{=} -\left(z + \sum_{i=1}^{N_1} X_2^{(i)}(z)\right)^{-1} \\ X_2(z) \overset{(d)}{=} -\left(z + \sum_{i=1}^{N_2} X_1^{(i)}(z)\right)^{-1}, \end{cases}$$

where X_1 and X_2 are independent random variables having laws $\mathscr{L}_1$ and $\mathscr{L}_2$, the $X_1^{(i)}$ and $X_2^{(i)}$ are i.i.d. copies of X_1 (resp. X_2), N_1 has law $\mathscr{P}(c)$ and N_2 has law $\mathscr{P}(\alpha c)$; each of these variables being independent. Then, the Stieltjes transform of μ_c' is given by:

$$\forall z \in \mathbf{C}_+, \quad m_{\mu_c'}(z) := \int_{\mathbf{R}} \frac{1}{x-z} \mathrm{d}\mu_c'(x) = \frac{1}{\alpha+1}\mathbf{E}[X_1(z)] + \frac{\alpha}{\alpha+1}\mathbf{E}[X_2(z)]. \tag{9}$$

This concludes the proof of theorem 2 the limiting law being $\mu_c = (\mathrm{Sym} \circ f_*)^{-1}\mu_c'$.

3.2 Asymptotic expansion of the moments

Combining Theorem 1 with Theorem 2, we deduce that, almost surely, μ_{W_n} converges weakly to a probability law $\mu_{\alpha,c}$ which is characterized by its sequence of moments. In that case, the asymptotic $\mathscr{A} = \{A_k\}_{k\geq 1}$ of the laws P_n which are Bernoulli laws of parameter c/n is given by:

$$A_k = \lim_{n\to+\infty} \frac{1}{n^{k/2-1}} \frac{(1-c/n)^k(c/n) + (-c/n)^k(1-c/n)}{[(1-c/n)^2(c/n) + (-c/n)^2(1-c/n)]^{k/2}} = c^{1-k/2}\mathbf{1}_{k>1}.$$

This leads to the following formula for the k-th moment of $\mu_{\alpha,c}$:

$$M_k(\mu_{\alpha,c}) = \sum_{a=1}^{k}\sum_{l=1}^{a} \alpha^l \sum_{\substack{b_1\geq b_2\geq\ldots\geq b_a\geq 2 \\ b_1+b_2+\cdots+b_a=2k}} |\mathscr{W}(a,a+1,l,b)| \prod_{i=1}^{a} c^{1-b_i/2}. \tag{10}$$

When $c \to +\infty$, we retrieve the moments of the Marchenko-Pastur law μ_α. It is therefore natural to try to understand how $\mu_{\alpha,c}$ differs from μ_α when c is large but finite. We give an answer to this question by giving an asymptotic expansion in $1/c$ of the moments of $\mu_{\alpha,c}$. This is done by a more careful treatment of equation (10), which is combinatorial in nature. The method is inspired by the paper [8] of Enriquez and Ménard, where the authors treated the case of adjacency matrices of Erdös-Rényi graphs with parameter c/n.

Theorem 3. *There exists a signed measure $\mu_\alpha^{(1)}$ such that for all $k \geq 1$, as $c \to +\infty$:*

$$M_k(\mu_{\alpha,c}) = M_k\left(\mu_\alpha + \frac{1}{c}\mu_\alpha^{(1)}\right) + o\left(\frac{1}{c}\right). \tag{11}$$

Moreover, the measure $\mu_\alpha^{(1)}$ has a total mass zero and the following density:

$$\frac{x^2 - 2x(\alpha+1) + (\alpha^2+1)}{2\alpha\sqrt{(b-x)(x-a)}}\mathbf{1}_{(a,b)}. \tag{12}$$

Proof. Fix an integer $k \geq 1$. First, Remark 2 ensures that all the b_i's in (10) are even. Therefore, we can rewrite:

$$M_k(\mu_{\alpha,c}) = \sum_{a=1}^{k} \sum_{l=1}^{a} \alpha^l \sum_{\substack{d_1 \geq d_2 \geq \ldots \geq d_a \geq 1 \\ d_1+d_2+\cdots+d_a=k}} |\mathscr{W}_k(a,a+1,l,2d)| \prod_{i=1}^{a} c^{1-d_i}. \tag{13}$$

As $c \to +\infty$, nonvanishing terms correspond to the case where all the d_i's are equal to 1. This forces a to be equal to k and leads to:

$$M_k(\mu_{\alpha,c}) = \sum_{l=1}^{a} \alpha^l |\mathscr{W}_k(k,k+1,l,(2,\ldots,2))| + o(1)$$

as $c \to +\infty$.

Recall that $\mathscr{W}_k(k,k+1,l,(2,\ldots,2))$ is a set of representatives of closed words starting at the root, of length $2k+1$ on labeled planar rooted trees having k edges, l of these being odd edges. This allows to write:

$$M_k(\mu_{\alpha,c}) = \sum_{\mathrm{T} \in \mathscr{T}_k} \alpha^{l(\mathrm{T})} + o(1)$$

as $c \to +\infty$, where $\mathscr{T}_k$ is the set of planar rooted trees having k edges and $l(\mathrm{T})$ the number of odd edges in a given tree $\mathrm{T} \in \mathscr{T}_k$. For convenience, we introduce the notations

$$a_k := \sum_{\mathrm{T} \in \mathscr{T}_k} \alpha^{l(\mathrm{T})} \quad \text{and} \quad b_k := \sum_{\mathrm{T} \in \mathscr{T}_k} \alpha^{\bar{l}(\mathrm{T})},$$

where $\bar{l}(\mathrm{T})$ is the number of even edges of a given tree $\mathrm{T} \in \mathscr{T}_k$.

It turns out that the a_k's are the moments of μ_α. To obtain the term of order $1/c$ we will need to compute the generating series of the a_k's and b_k's.

Let T be a planar tree having $k+1$ edges. Let T_1 be the tree induced by the first child of the root and T_2 the connected component of the root after removing the edge between the root and its first child (see Figure 4).




Figure 4: Decomposition of a planar tree.

Denoting p (resp. q) the number of edges of T_1 (resp. T_2), we have $p+q=k$. It is straightforward to obtain the relations $l(\mathrm{T}) = 1 + \bar{l}(\mathrm{T}_1) + l(\mathrm{T}_2)$ and $\bar{l}(\mathrm{T}) = l(\mathrm{T}_1) + \bar{l}(\mathrm{T}_2)$. Therefore

$$\begin{cases} a_{k+1} = \alpha \sum\limits_{p+q=k} a_p b_q \\ b_{k+1} = \sum\limits_{p+q=k} a_p b_q. \end{cases} \tag{14}$$

Denoting $A(z)=\sum_{k\geq 0}a_k z^k$ and $B(z)=\sum_{k\geq 0}b_k z^k$ the generating functions of the a_k's and the b_k's we obtain the functional relations:

$$\begin{cases} A = 1+\alpha zAB \\ B = 1+zAB. \end{cases} \tag{15}$$

It implies that $zA^2+(\alpha z-z-1)A+1=0$. If we denote $S(z):=-z^{-1}A(z^{-1})$ the Stieltjes transform of the measure with moments a_k's, then S satisfies the equation:

$$zS^2-(\alpha-z-1)S+1=0. \tag{16}$$

The function S of the variable $z\in\mathbf{C}_+$ is the limit of the Stieltjes transform of the μ_{W_n} when $c\to+\infty$. The imaginary part of a Stieltjes transform is positive: this allows us to choose the right solution for equation (16). For a complex z, if we denote $\sqrt{z}$ the square root having a positive imaginary part on the upper half plane:

$$S(z)=\frac{\alpha-z-1+\sqrt{(z-b)(z-a)}}{2z}, \tag{17}$$

where $a=(1-\sqrt{\alpha})^2$ and $b=(1+\sqrt{\alpha})^2$. This is the Stieltjes transform of the Marchenko-Pastur law μ_α, as announced.

Let us compute the perturbation of order $1/c$. It arises when all the d_i's are equal to 1 except one which is equal to 2 in (13). This forces a to be equal to $k-1$ and leads to the following expansion as $c\to+\infty$:

$$M_k(\mu_{\alpha,c})=M_k(\mu_\alpha)+\frac{1}{c}\sum_{l=1}^{k-1}\alpha^l|\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))|+o\left(\frac{1}{c}\right).$$

In that case $\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))$ is the set of equivalence classes of closed words of length $2k+1$ on labeled planar rooted tree having $k-1$ edges, starting at the root and such that each edge is browsed exactly two times except one which is browsed four times. Let us denote

$$a_k^{(1)}=\sum_{l=1}^{k-1}\alpha^l|\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))|,$$

and

$$b_k^{(1)}=\sum_{l=1}^{k-1}\alpha^l|\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))|.$$

The associated generating series will be denoted $A^{(1)}$ and $B^{(1)}$. Remark that by definition $a_0^{(1)}=a_1^{(1)}=b_0^{(1)}=b_1^{(1)}=0$. We are going to obtain a recursion linking the four generating series $A,B,A^{(1)}$ and $B^{(1)}$. The idea is to use a first generation decomposition of the planar rooted tree on which the words are written, and then to distinguished whether or not the quadruple edge is an edge of this generation. For all $k\geq 1$, we use the partition

$$\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))=\mathscr{W}_k^{(0)}(k-1,k,l,(4,2,\ldots,2))\bigsqcup\mathscr{W}_k^{(1)}(k-1,k,l,(4,2,\ldots,2)),$$

where $\mathscr{W}_k^{(0)}(k-1,k,l,(4,2,\ldots,2))$ is the set of representative belonging to $\mathscr{W}_k(k-1,k,l,(4,2,\ldots,2))$ such that the quadruple edge is not a first generation edge, and $\mathscr{W}_k^{(1)}(k-1,k,l,(4,2,\ldots,2))$

is the set of representatives belonging to $\mathscr{W}_k(k-1,k,l,(4,2,\dots,2))$ such that the quadruple edge is a first generation edge. The associated quantities will be denoted $a_k^{(1,0)}, a_k^{(1,1)}, A^{(1,0)}, \dots$ For example:

$$a_k^{(1,0)} = \sum_{l=1}^{k-1} \alpha^l |\mathscr{W}_k^{(0)}(k-1,k,l,(4,2,\dots,2))|.$$

A representative word $(\mathbf{i},\mathbf{j}) \in \mathscr{W}_k(k-1,k,l,(4,2,\dots,2))$ can be written:

$$(\mathbf{i},\mathbf{j}) = i_1 \mathbf{S}_1 \zeta \xi \mathbf{S}_2 \xi \zeta \mathbf{S}_3 \zeta \xi \mathbf{S}_4 \xi \zeta \mathbf{S}_5 i_1,$$

where:

1. $i_1 \mathbf{S}_1 \zeta \mathbf{S}_5 i_1$ is the contour of a planar tree having p_1 edges;
2. $\xi \mathbf{S}_2 \xi$ is the contour of a planar tree having p_2 edges;
3. $\zeta \mathbf{S}_3 \zeta$ is the contour of a planar tree having p_3 edges;
4. $\xi \mathbf{S}_4 \xi$ is the contour of a planar tree having p_4 edges;
5. $\xi \mathbf{S}_2 \xi \mathbf{S}_4 \xi$ is the contour of a planar tree having $p_2 + p_4$ edges.

The above integers satisfy $p_1 + p_2 + p_3 + p_4 = k-2$. See Figure 5 for an illustration.




Figure 5: The writing $(\mathbf{i},\mathbf{j})$ and its quadruple edge $\{\zeta,\xi\}$.

All of these conditions are sufficient to define a class of canonical representatives. Let T be the planar rooted tree on which a representative word $(\mathbf{i},\mathbf{j})$ is written. Denote e_4 the quadruple edge, $\mathrm{T} \setminus e_4$ the connected component of the root after removing e_4 and T^{e_4} the planar rooted tree formed by the descendants of e_4. Then, the above conditions ensures that $(\mathbf{i},\mathbf{j})$ is such that $\mathrm{T} \setminus e_4$ and T^{e_4} are respectively browsed in lexicographic order.

Let $(\mathbf{i},\mathbf{j}) \in \mathscr{W}_k^{(0)}(k-1,k,l,(4,2,\dots,2))$. The underlying tree can have $p \in \{1,\dots,k-2\}$ edges which are all browsed two times by $(\mathbf{i},\mathbf{j})$. One of the tree induced by the children of the root contains the quadruple edge, leading to p different choices. On another side, if $(\mathbf{i},\mathbf{j}) \in \mathscr{W}_k^{(1)}(k-1,k,l,(4,2,\dots,2))$ then the underlying tree can have $p \in \{1,\dots,k-1\}$ edges out of which one is the quadruple edge. There are $\binom{p+1}{2}$ choices for the locations of the the visits of the quadruple edge. See Figure 6 for an illustration.




Figure 6: First edge decomposition of a word respectively in $\mathscr{W}_k^{(0)}(k-1,k,l,(4,2,\dots,2))$ on the left and in $\mathscr{W}_k^{(1)}(k-1,k,l,(4,2,\dots,2))$ on the right, where the quadruple edge is in red.

As a consequence, we get the following recursions:

$$a_k^{(1,0)} = \sum_{p=1}^{k-2} \alpha^p p \sum_{q_1+\cdots+q_p=k-p-1} b^{(1)}_{q_1+1} b_{q_2} \cdots b_{q_p},$$

and

$$a_k^{(1,1)} = \sum_{p=1}^{k-2} \alpha^p \binom{p+1}{2} \sum_{q_1+\cdots+q_{p+1}=k-p-1} b_{q_1} b_{q_2} \cdots b_{q_{p+1}}.$$

This yields

$$A^{(1,0)} = \frac{\alpha z B^{(1)}}{(1-\alpha z B)^2} = \alpha z A^2 B^{(1)}$$

and

$$A^{(1,1)} = \frac{\alpha z^2 B^2}{(1-\alpha z B)^3} = \alpha z^2 A^3 B^2,$$

where we used equation (15). The same arguments and computations give $B^{(1,0)} = zA^{(1)}B^2$ and $B^{(1,1)} = z^2A^2B^3$, to finally obtain

$$\begin{cases} A^{(1)} = \alpha z A^2 B^{(1)} + \alpha z^2 A^3 B^2 \\ B^{(1)} = zA^{(1)} B^2 + z^2 A^2 B^3. \end{cases} \tag{18}$$

We deduce, using equation (15), that $A^{(1)}$ is given by:

$$A^{(1)} = \frac{\alpha (zAB)^2}{1-\alpha(zAB)^2}(zA^2B + A) = \frac{AB}{1-\alpha(zAB)^2}\alpha(zAB)^2. \tag{19}$$

To obtain a more explicit formula for $A^{(1)}$, one can compute $\alpha(zAB)^2$ using first that $B = (A+\alpha-1)/\alpha$ and then that $zA^2 = (1-(\alpha-1)z)A - 1$. After simplifications:

$$\alpha(zAB)^2 = \frac{(1-\alpha z - z)A + z - 1}{\alpha z} = \frac{(\alpha^2+1)z^2 - 2z(\alpha-1) + 1 - (1-\alpha z - z)\sqrt{\delta}}{2\alpha z^2}, \tag{20}$$

since $A = (2z)^{-1}(1-(\alpha-1)z - \sqrt{\delta})$. Using that $\sqrt{\delta} = -2zA - (\alpha-1)z + 1$, one can then check that $\sqrt{\delta}AB = 1 - \alpha(zAB)^2$. From (20), one can finally rewrite (19) as

$$A^{(1)} = \frac{1}{\sqrt{\delta}} \frac{(\alpha^2+1)z^2 - 2z(\alpha+1) + 1 - (1-\alpha z - z)\sqrt{\delta}}{2\alpha z^2}.$$

Therefore, the function $S^{(1)}(z) = -\frac{1}{z}A^{(1)}(\frac{1}{z})$ is given by

$$S^{(1)}(z) = -\frac{z^2 - 2z(\alpha+1) + (\alpha^2+1)}{2\alpha\sqrt{(z-b)(z-a)}} + \frac{z-\alpha-1}{2\alpha}. \tag{21}$$

It corresponds to the Stieltjes transform of the measure $\mu_\alpha^{(1)}$ with density:

$$-\frac{1}{\pi}\lim_{\varepsilon\to 0} \mathrm{Im}\big(S^{(1)}(x+i\varepsilon)\big) = \frac{x^2 - 2x(\alpha+1) + (\alpha^2+1)}{2\alpha\pi\sqrt{(b-x)(x-a)}} \mathbf{1}_{(a,b)}.$$

This concludes the proof of Theorem 3. □

The case $\alpha = 1$, which corresponds to asymptotic square matrices X_n, should be emphasized. In this setting the density is

$$\frac{1}{2\pi}\frac{x^2-4x+2}{\sqrt{x(4-x)}}\mathbf{1}_{[0,4]},$$

which corresponds to the pushforward by $x \mapsto x^2$ of the density obtained in [8] by Enriquez and Ménard for the Wigner case, as expected.

4 Heavy tailed random matrices

In this section we use Theorem 1 to study the spectral measure associated to heavy tailed Wishart matrices. For all $n \geq 1$, let X_n be a random matrix of size $n \times m$ having i.i.d. entries with heavy tailed law P. As before, we suppose that the ratio m/n converges to $\alpha > 0$. We will consider the case where P has density

$$\frac{C(\beta)}{1+|x|^\beta},$$

where $1 < \beta < 3$ and $C(\beta) = (\int_{\mathbf{R}}(1+|x|^\beta)^{-1}\mathrm{d}x)^{-1}$. Theorem 1.10 of Belinschi, Dembo and Guionnet in [2] ensures that, since P is in the domain of attraction of a $(\beta-1)$-stable law, the spectral measure of

$$n^{-\frac{2}{\beta-1}}X_nX_n^T$$

converges to a deterministic probability law $\mu_{\alpha,\beta}$ depending only on α and β.

To apply Theorem 1, let us consider the truncated version of X_n. For all $n \geq 1$, let P_n be the probability law given by

$$P_n(\mathrm{d}x) = \frac{C}{1+|x|^\beta}\mathbf{1}_{[-Bn^{1/(\beta-1)},Bn^{1/(\beta-1)}]}\mathrm{d}x + Z(B,\beta)\left(\delta_{-Bn^{1/(\beta-1)}}(\mathrm{d}x) + \delta_{Bn^{1/(\beta-1)}}(\mathrm{d}x)\right),$$

where $B > 0$ and $Z(B,\beta) = 2C(\beta)\int_{Bn^{1/(\beta-1)}}^{+\infty}(1+|x|^\beta)^{-1}\mathrm{d}x$. In words, P_n is the truncation of P at $-Bn^{1/(\beta-1)}$ and $Bn^{1/(\beta-1)}$. We will denote Y_n the random matrix of size $n \times m$ with i.i.d. entries having law P_n. Let us compute the asymptotic $\mathscr{A} = \{A_k\}_{k\geq 2}$ of the sequence $\{P_n\}_{n\geq 1}$. For all $k \geq 1$, as n tends to infinity:

$$\begin{aligned}
\frac{M_k(P_n)}{n^{k/2-1}M_2(P_n)^{k/2}} &\sim n^{1-k/2}(2C)^{1-k/2}\frac{\int_1^{Bn^{1/(\beta-1)}}x^{k-\beta}\mathrm{d}x + \frac{(Bn^{1/(\beta-1)})^{k+1-\beta}}{1-\beta}}{(\int_1^{Bn^{1/(\beta-1)}}x^{2-\beta}\mathrm{d}x + \frac{(Bn^{1/(\beta-1)})^{3-\beta}}{1-\beta})^{k/2}} \\
&\sim n^{1-k/2}(2C)^{1-k/2}\frac{\frac{(Bn^{1/(\beta-1)})^{k+1-\beta}}{k+1-\beta} + \frac{(Bn^{1/(\beta-1)})^{k+1-\beta}}{1-\beta}}{(\frac{(Bn^{1/(\beta-1)})^{3-\beta}}{3-\beta} + \frac{(Bn^{1/(\beta-1)})^{3-\beta}}{1-\beta})^{k/2}} \\
&\sim n^{1-k/2}(2C)^{1-k/2}\frac{\frac{1}{k+1-\beta}+\frac{1}{1-\beta}}{\frac{1}{3-\beta}+\frac{1}{1-\beta}}B^{1-\beta+\frac{k}{2}(\beta-1)}n^{\frac{1}{\beta-1}\left(k+1-\beta+\frac{k}{2}(\beta-3)\right)}.
\end{aligned}$$

We finally obtain:

$$\frac{M_k(P_n)}{n^{k/2-1}M_2(P_n)^{k/2}} \sim (2C)^{1-k/2}\frac{\frac{1}{k+1-\beta}+\frac{1}{1-\beta}}{\frac{1}{3-\beta}+\frac{1}{1-\beta}}B^{1-\beta+\frac{k}{2}(\beta-1)}.$$

The quantity $n^{1/(\beta-1)}$ corresponds to the largest n-th quantile of P. Therefore, our choice of law P_n can be interpreted as a truncation of the largest entries in each rows of X_n. If one had chosen an order of truncation smaller than $n^{1/(\beta-1)}$, the A_k's would have been all equal to zero which corresponds to the Marchenko-Pastur regime, meaning that the truncation is too large and leads to a non-heavy tailed behavior. On the contrary, if one had chosen an order of truncation larger than $n^{1/(\beta-1)}$, the A_k's would have been all infinite, meaning that the truncation is not large enough to apply Theorem 1. In this spirit, the parameter $B > 0$ can be seen as a finer adjustment of the truncation.

Theorem 1 ensures that there exists a probability law $\mu_{\mathscr{A},\alpha} = \mu_{\alpha,\beta,B}$ such that the spectral measures μ_n associated to the Wishart matrices $\frac{1}{nM_2(P_n)} Y_n Y_n^T$ converges weakly in probability to $\mu_{\alpha,\beta,B}$. Using equation (2), we obtain an asymptotic development of the moments of $\mu_{\alpha,\beta,B}$:

Theorem 4. *For all $k \geq 1$, as $B \to 0$:*

$$M_k(\mu_{\alpha,\beta,B}) = M_k(\mu_\alpha) + B^{\beta-1} \frac{1}{2C(\beta)} \cdot \frac{(3-\beta)^2}{(2-\beta)(5-\beta)} M_k\left(\mu_\alpha^{(1)}\right) + o\left(B^{\beta-1}\right).$$

Remark 5. *For simplicity we considered the explicit law $P(\mathrm{d}x) = \frac{C(\beta)}{1+|x|^\beta}$. However, using Karamata's estimates (Theorem 2, Section VIII.9 of [9]) on truncated moments of regularly varying functions, one could have studied in a similar way the case when P is in the domain of attraction of a $(\beta-1)$-stable law, for $1 < \beta < 3$.*

Acknowledgments. The author would like to warmly thank his advisors Nathanaël Enriquez and Laurent Ménard for many helpful discussions and suggestions about this work.

References

[1] David Aldous and J Michael Steele. The objective method: probabilistic combinatorial optimization and local weak convergence. In *Probability on discrete structures*, pages 1–72. Springer, 2004.

[2] Serban Belinschi, Amir Dembo, and Alice Guionnet. Spectral measure of heavy tailed band and covariance random matrices. *Communications in Mathematical Physics*, 289(3):1023–1055, 2009.

[3] Florent Benaych-Georges and Thierry Cabanal-Duvillard. Marchenko-pastur theorem and bercovici-pata bijections for heavy-tailed or localized vectors. *ALEA: Latin American Journal of Probability and Mathematical Statistics*, 9(2):685–715, 2012.

[4] Itai Benjamini and Oded Schramm. Recurrence of distributional limits of finite planar graphs. In *Selected Works of Oded Schramm*, pages 533–545. Springer, 2011.

[5] Charles Bordenave, Djalil Chafaï, et al. Around the circular law. *Probability surveys*, 9, 2012.

[6] Charles Bordenave and Marc Lelarge. Resolvent of large random graphs. *Random Structures & Algorithms*, 37(3):332–352, 2010.

[7] Charles Bordenave, Arnab Sen, and Balint Virag. Mean quantum percolation. *To appear in Journal of the European Mathematical Society (JEMS)*, 2015.

[8] Nathanaël Enriquez and Laurent Ménard. Spectra of large diluted but bushy random graphs. *Random Structures & Algorithms*, 2015.

[9] William Feller and T Teichmann. An introduction to probability theory and its applications, vol. 2, 1967.

[10] Camille Male. The limiting distributions of large heavy wigner and arbitrary random matrices. *Journal of Functional Analysis*, 272(1):1–46, 2017.

[11] Vladimir Alexandrovich Marchenko and Leonid Andreevich Pastur. Distribution of eigenvalues for some sets of random matrices. *Matematicheskii Sbornik*, 114(4):507–536, 1967.

[12] Justin Salez. Every totally real algebraic integer is a tree eigenvalue. *Journal of Combinatorial Theory, Series B*, 111:249–256, 2015.

[13] Valentin Vengerovsky. Eigenvalue distribution of a large weighted bipartite random graph. *Journal of Mathematical Physics, Analysis, Geometry*, 10(2):240–255, 2014.

[14] Inna Zakharevich. A generalization of wigner's law. *Communications in mathematical physics*, 268(2):403–414, 2006.

Nathan Noiry :
Laboratoire Modal'X,
UPL, Université Paris Nanterre,
F92000 Nanterre France