

# Heat trace asymptotics for wedge-like singularity.

Asilya Suleymanova

October 2017

## Abstract

In this note we consider a heat trace expansion on a manifold with wedge-like singularity. We show that there are two terms in the expansion that contain information about the presence of the singularity, namely the logarithmic term  $ct^{-1/2} \log t$  and the half power term  $+bt^{-1/2}$ . We also give a geometric expression for  $c$ .

## Contents

|          |   |          |
|----------|---|----------|
| <b>1</b> | <b>Geometric setup</b>                              | <b>1</b> |
| <b>2</b> | <b>Curvature tensors near the singularity</b>       | <b>2</b> |
| <b>3</b> | <b>Resolvent expansion</b>                          | <b>3</b> |
| <b>4</b> | <b>Heat trace expansion</b>                         | <b>6</b> |
| 4.1      | Logarithmic term in low dimensional cases . . . . . | 6        |

## 1 Geometric setup

Consider a Riemannian manifold,  $(M, g)$ , of dimension  $m$  such that  $M = M_1 \cup U$ , where  $M_1$  is a compact manifold with boundary and  $U = (0, 1) \times N$  has a fibration  $(0, 1) \times F \rightarrow U \rightarrow (0, 1) \times \Sigma$ , where  $N, \Sigma$  are compact. This fibration induces a vertical bundle of tangents to the fibres  $T_V N \subset TN$  and a horizontal cotangent subbundle  $T_H^* N$  of cotangents annihilating  $T_V N$ . We take a complementary tangent subbundle  $T_H N \subset TN$ , inducing a vertical  $T_V^* N$  that annihilates  $T_H N$  such that  $TN = T_V N \oplus T_H N$ .

The metric on  $U$  is

$$g_U = dr^2 + g_H + r^2 g_V(s), \quad r \in (0, 1),$$

where  $g_H$  is induced from a metric on  $\Sigma$  and  $g_V(s)$  is a family of metrics on the fibres,  $s \in \Sigma$ .

Assume now that  $\dim \Sigma = 1$ , i.e. it is a circle. Assume furthermore that metric on the fibres does not depend on  $s$ , i.e.  $g_V(s) = g_V$ .

$$g_{\text{wedge}} = dr^2 + d\theta^2 + r^2 g_V, \quad r \in (0, 1),$$

We call such singularity a wedge-like singularity.

## 2 Curvature tensors near the singularity

In this section we give explicit formulas for the curvature tensors in  $(U, g_{\text{wedge}})$  in terms of the curvature tensors on fibers. For the latter we use the classical tensor notations and for the tensors on  $(U, g_{\text{wedge}})$  we use classical notations but with tilde.

Let  $(x^1, \dots, x^{m-2})$  be local coordinates on a fiber  $F$  and  $p = (r, \theta, x^1, \dots, x^{m-2}) \in U$ . For  $i, j \in \{r, \theta, 1, \dots, m-2\}$  denote by  $\tilde{g}_{ij}$  the components of the metric tensor on  $g_{\text{wedge}}$ , and by  $g_{ij}$  for  $i, j \in \{1, \dots, m-2\}$  the components of the metric tensor  $g_V$ . Then

$$\tilde{g}_{rr} = \tilde{g}_{\theta\theta} = 1,$$

$$\tilde{g}_{ir} = \tilde{g}_{ri} = \tilde{g}_{j\theta} = \tilde{g}_{\theta j} = 0, \quad \text{for } i \in \{\theta, 1, \dots, m-2\} \text{ and } j \in \{r, 1, \dots, m-2\}$$

and

$$\tilde{g}_{ij} = r^2 g_{ij}, \quad \text{for } i, j \in \{1, \dots, m-2\}.$$

Then

$$\tilde{g}_{ij,k}(r, \theta, x) = \begin{cases} 2r g_{ij}(x), & \text{if } k = r, \\ 0 & \text{if } k = \theta, \\ r^2 g_{ij,k}(x) & \text{otherwise.} \end{cases}$$

The Christoffel symbols are of course

$$\tilde{\Gamma}_{jk}^i = \frac{1}{2} \tilde{g}^{im} (\tilde{g}_{mj,k} + \tilde{g}_{mk,j} - \tilde{g}_{jk,m}),$$

and now we express them in terms of the Christoffel symbols  $\Gamma_{jk}^i$  and the metric tensor  $g_{ij}$ .

Let  $i = r$ , then

$$\tilde{\Gamma}_{jk}^r = \begin{cases} 0, & \text{if } j = r \text{ or } k = r \text{ or } j = \theta \text{ or } k = \theta, \\ -rg_{jk} & \text{otherwise.} \end{cases}$$

Let  $i \notin \{r, \theta\}$  and  $j = r$ , then

$$\tilde{\Gamma}_{rk}^i = \tilde{\Gamma}_{kr}^i = \begin{cases} 0, & \text{if } k = r \text{ or } i = \theta \text{ or } k = \theta, \\ r^{-1}\delta_k^i & \text{otherwise.} \end{cases}$$

Assume that  $i, j, k \notin \{r, \theta\}$ , then  $\tilde{\Gamma}_{jk}^i = \Gamma_{jk}^i$ . Finally  $\tilde{\Gamma}_{jk}^\theta = \tilde{\Gamma}_{\theta k}^i = \tilde{\Gamma}_{k\theta}^i = 0$ . The scalar curvature  $\text{Scal}(p)$  can be expressed in the following way

$$\begin{aligned} \text{Scal}(p) &= \tilde{g}^{ij} \left( \tilde{\Gamma}_{ij,m}^m - \tilde{\Gamma}_{im,j}^m + \tilde{\Gamma}_{ij}^l \tilde{\Gamma}_{ml}^m - \tilde{\Gamma}_{im}^l \tilde{\Gamma}_{jl}^m \right) \\ &= r^{-2} (\text{Scal}(x) - (m-2)(m-3)), \end{aligned} \quad (2.1)$$

where  $p = (r, \theta, x) \in U$ ,  $x \in F$  and  $\text{Scal}(p)$  is the scalar curvature on  $(U, g_{\text{wedge}})$ , and  $\text{Scal}(x)$  is the scalar curvature on  $(F, g_V)$ .

If  $i, j, k, l \notin \{r, \theta\}$ , we obtain

$$\tilde{R}_{ijkl}(p) = r^{-2} (R_{ijkl}(x) - g_{ip}(x)g_{jm}(x)(\delta_k^p \delta_l^m - \delta_l^p \delta_k^m)). \quad (2.2)$$

Similarly, for the tensor Ricci

$$\tilde{\text{Ric}}_{ij}(p) = r^{-2} (\text{Ric}_{ij}(x) - (m-3)g_{ij}(x)). \quad (2.3)$$

### 3 Resolvent expansion

Let  $\Delta$  be the Friedrichs extension of the Laplace-Beltrami operator on  $U$

$$\Delta = -\partial_r^2 + \Delta_\Sigma + r^{-2}A(s),$$

where  $\Delta_\Sigma$  is the Laplace operator on the singular stratum  $\Sigma$ ,  $A(s), s \in \Sigma$  is a smooth family of elliptic operators on functions on fibers  $F$ . In the case of the wedge-like singularity, we have  $\Delta_\Sigma = -\partial_\theta^2$ , and  $A(s) = \Delta_F + \left(\frac{m-2}{2}\left(\frac{m-2}{2} - 1\right)\right)$ , where  $\Delta_F$  is the Laplace-Beltrami operator on the fiber  $F$ .

Let  $\varphi(r, \theta)$  be a cutoff function supported near  $r = 0$ , by [BS, Theorem 5.2], for  $d > m/2$  the resolvent  $(\Delta + z^2)^{-d}$  is trace class and the following is true

$$\text{tr}(\varphi(r, \theta)(\Delta + z^2)^{-d}) = \int_0^\infty \varphi(r, \theta)\sigma(r, rz)dr,$$

where  $\sigma(r, rz) := \sigma(r, \theta, rz) := \text{tr}_{L^2(F)}(\Delta + z^2)^{-d}$ .

From the local heat kernel expansion of heat kernel away from the singularity we know, see [S, (3.6), (3.17)], that

$$\begin{aligned} & \text{tr}_{L^2(F)}(\Delta + \zeta^2/r^2)^{-d} \sim_{\zeta \rightarrow \infty} \\ & (4\pi)^{-\frac{m}{2}} \sum_{j=0}^{\infty} (\zeta/r)^{-2d+m-2j} \frac{\Gamma(-\frac{m}{2} + d + j)}{(d-1)!} \int_F r^{m-2} u_j(p) \, \text{dvol}_F. \end{aligned}$$

Above  $u_j(p)$  are polynomials in the curvature tensors at  $p \in M$  that come from the local heat kernel expansion away from the singularity, in particular  $u_0(p) \equiv 1$ ,  $u_1(p) = \text{Scal}(p)$  and

$$u_2(p) = \frac{1}{360} (12\Delta \text{Scal}(p) + 5 \text{Scal}(p)^2 - 2|\text{Ric}(p)|^2 + 2|\text{R}(p)|^2).$$

Hence

$$\sigma(r, \zeta) \sim_{\zeta \rightarrow \infty} \sum_{j=0}^{\infty} \zeta^{-2d+m-2j} \sigma_j(r), \quad (3.1)$$

where

$$\begin{aligned} \sigma_j(r) &= (4\pi)^{-\frac{m}{2}} r^{2d-m+2j} \int_F r^{m-2} u_j(p) \, \text{dvol}_F \frac{\Gamma(-\frac{m}{2} + d + j)}{(d-1)!} \\ &= (4\pi)^{-\frac{m}{2}} r^{2d-2+2j} \int_F u_j(p) \, \text{dvol}_F \frac{\Gamma(-\frac{m}{2} + d + j)}{(d-1)!}. \end{aligned} \quad (3.2)$$

We can apply the Singular Asymptotics Lemma to obtain an asymptotic expansion of the resolvent trace, see [BS, p. 287],

**Proposition 3.1.**

$$\int_0^\infty \varphi(r, \theta) \sigma(r, \zeta) dr \sim \sum_{l=0}^{\infty} z^{-l-1} \frac{1}{l!} \int_0^\infty \int_\Sigma \zeta^l \partial_r^l \left( \sigma(r, \zeta) \varphi(r, \theta) \right) \Big|_{r=0} d\theta d\zeta \quad (3.3)$$

$$+ \sum_{j=0}^{\infty} \int_0^\infty \int_\Sigma \sigma_j(r) (rz)^{-2d+m-2j} \varphi(r) d\theta dr \quad (3.4)$$

$$+ \sum_{l=\frac{m}{2}-d+1}^{\infty} z^{-2d+m-2l} \log z \int_\Sigma \frac{\partial_r^{2d-m+2l-1} \left( \sigma_l(r) \varphi(r, \theta) \right) \Big|_{r=0}}{(2d-m+2l-1)!} d\theta. \quad (3.5)$$

Assume  $\varphi(r, \theta) \equiv 1$  near  $r = 0$  and consider the last sum in the expansion

$$\begin{aligned}
L &:= \sum_{l=\frac{m}{2}-d+1}^{\infty} z^{-2d+m-2l} \log z \int_{\Sigma} \frac{\partial_r^{2d-m+2l-1} \left( \sigma_l(r) \varphi(r, \theta) \right) |_{r=0}}{(2d-m+2l-1)!} d\theta \\
&= \sum_{j=\frac{m}{2}-d+1}^{\infty} z^{-2d+m-2j} \log z \frac{\Gamma(-\frac{m}{2} + d + j) \text{vol}(\Sigma)}{(d-1)!(2d-m+2j-1)!} \times \\
&\quad \times (4\pi)^{-\frac{m}{2}} \partial_r^{2d-m+2j-1} \left( r^{2d-2+2j} \int_F u_j(p) \, d\text{vol}_N \right) |_{r=0}.
\end{aligned}$$

To continue the computation we note that for  $j \geq m/2$  we have

$$\begin{aligned}
&\partial_r^{2d-m+2j-1} \left( r^{2d-2+2j} \int_F u_j(p) \, d\text{vol}_N \right) |_{r=0} \\
&= \partial_r^{2j-m+1} \left( r^{2j} \int_F u_j(p) \, d\text{vol}_N \right) |_{r=0},
\end{aligned}$$

this makes sense only for  $j \geq \frac{m-1}{2}$ , therefore

$$\begin{aligned}
L &= \sum_{j \geq \frac{m-1}{2}} z^{-2d+m-2j} \log z \frac{\Gamma(-\frac{m}{2} + d + j) \text{vol}(\Sigma)}{(d-1)!(2d-m+2j-1)!} \times \\
&\quad \times (4\pi)^{-\frac{m}{2}} \partial_r^{2j-m+1} \left( r^{2j} \int_F u_j(p) \, d\text{vol}_F \right) |_{r=0} \\
&= \sum_{l=0}^{\infty} z^{-2d-2l+1} \log z \frac{\Gamma(d+l-\frac{1}{2}) \text{vol}(\Sigma)}{(d-1)!(2d+2l-2)!} \times \\
&\quad \times (4\pi)^{-\frac{m}{2}} \partial_r^{2l} \left( r^{m+2l-1} \int_F u_{\frac{m-1}{2}+l}(p) \, d\text{vol}_F \right) |_{r=0} \\
&= z^{-2d+1} \log z \frac{\Gamma(d-\frac{1}{2}) \text{vol}(\Sigma)}{(d-1)!(2d-2)!} \times \\
&\quad \times (4\pi)^{-\frac{m}{2}} \left( r^{m-1} \int_F u_{\frac{m-1}{2}}(p) \, d\text{vol}_F \right) |_{r=0}.
\end{aligned}$$

The last equality is due to the fact that  $r^{m-1+2l} u_{\frac{m-1}{2}+l}(p)$  is a smooth function with respect to  $r$ .

By [S, p.47], the logarithmic part in the heat trace expansion coming from this term is

$$-(4\pi)^{-\frac{m}{2}} t^{-\frac{1}{2}} \log t \times \frac{1}{2} \text{vol}(\Sigma) \left( r^{m-1} \int_F u_{\frac{m-1}{2}}(p) \, d\text{vol}_F \right) \Big|_{r=0}. \quad (3.6)$$

## 4 Heat trace expansion

From Proposition 3.1 and (3.6) we obtain

$$\text{tr } e^{-t\Delta} \sim_{t \rightarrow 0+} (4\pi t)^{-\frac{m}{2}} \sum_{j=0}^{\infty} \tilde{a}_j t^j + b t^{-1/2} + c t^{-1/2} \log t, \quad (4.1)$$

where

$$\tilde{a}_j = \begin{cases} \int_M u_j \, d\text{vol}_M & \text{for } j \leq m/2 - 1, \\ \int_M u_j \, d\text{vol}_M & \text{for } j > m/2 - 1. \end{cases}$$

and

$$c = \frac{1}{2} \text{vol}(\Sigma) \left( r^{m-1} \int_F u_{\frac{m-1}{2}}(p) \, d\text{vol}_F \right) \Big|_{r=0},$$

$b$  is to be computed.

Note that if  $m$  is even, the logarithmic term is zero.

### 4.1 Logarithmic term in low dimensional cases

Let  $\dim M = 3$ , then  $c = \frac{1}{12} \text{vol}(\Sigma) \text{Scal}(x) = 0$ , so the logarithmic term is always zero in this case.

Let  $\dim M = 5$ , then

$$\begin{aligned} c &= \frac{1}{2} \text{vol}(\Sigma) \int_F \left( \frac{1}{360} (5 \text{Scal}(p)^2 - 2|\text{Ric}(p)|^2 + 2|\text{R}(p)|^2) \right) d\text{vol}_F \\ &= -\frac{\text{vol}(\Sigma)}{720} (4\pi)^{-2} \int_F \left( 3(\text{Scal}(x) - 6)^2 + 6(\text{Ric}_{ij}(x) - 2g_{ij})^2 \right) d\text{vol}_F. \end{aligned}$$

The above expression is equal to zero if and only if  $\text{Scal}(x) \equiv 6$  and  $\text{Ric}_{ij}(x) = 2g_{ij}(x)$ ; equivalently if and only if the sectional curvature of the fiber  $(F, g_V)$  is equal to one. Therefore the logarithmic term in this case is equal to zero if and only if the fiber  $(F, g_V)$  is isometric to a spherical space form.

## References

- [BS] J. Brüning, R. Seeley, *The expansion of the resolvent near a singular stratum of conical type*, J. Funct. Anal. 95 (1991), 255–290.
- [S] A. Suleymanova, On the spectral geometry of manifolds with conic singularities, PhD thesis, <https://edoc.hu-berlin.de/handle/18452/19097> (2017).