

A RECOGNITION ALGORITHM FOR SIMPLE-TRIANGLE GRAPHS

ASAHI TAKAOKA

ABSTRACT. A simple-triangle graph is the intersection graph of triangles that are defined by a point on a horizontal line and an interval on another horizontal line. The time complexity of the recognition problem for simple-triangle graphs was a longstanding open problem, which was recently settled. This paper provides a new recognition algorithm for simple-triangle graphs to improve the time bound from $O(n^2\bar{m})$ to $O(nm)$, where n , m , and \bar{m} are the number of vertices, edges, and non-edges of the graph, respectively. The algorithm uses the vertex ordering characterization that a graph is a simple-triangle graph if and only if there is a linear ordering of the vertices containing both an alternating orientation of the graph and a transitive orientation of the complement of the graph. We also show, as a byproduct, that an alternating orientation can be obtained in $O(nm)$ time for cocomparability graphs, and it is NP-complete to decide whether a graph has an orientation that is alternating and acyclic.

1. INTRODUCTION

A graph is an *intersection graph* if there is a set of objects such that each vertex corresponds to an object and two vertices are adjacent if and only if the corresponding objects have a nonempty intersection. Such a set of objects is a *representation* of the graph. See [3, 13, 23, 29] for survey. Let L_1 and L_2 be two horizontal lines in the plane with L_1 above L_2 . *Trapezoid graphs* are the intersection graphs of trapezoids that are defined by an interval on L_1 and an interval on L_2 . Trapezoid graphs have been introduced in [7, 8] as a generalization of both interval graphs and permutation graphs. Many recognition algorithms and structural characterizations are provided for trapezoid graphs [5, 6, 10, 15, 20, 21, 26].

We obtain some interesting subclasses of trapezoid graphs by restricting the trapezoids in the representation. A trapezoid graph is a *simple-triangle graph* if every trapezoid in the representation is a triangle with the apex on L_1 and the base on L_2 . Similarly, a trapezoid graph is a *triangle graph* if every trapezoid in the representation is a triangle, but there is no restriction on which line contains the apex and the base. Simple-triangle graphs and triangle graphs have been introduced in [7] and studied under the name of *PI graphs* and *PI* graphs*, respectively [3, 4, 7, 29], where *PI* stands for *Point-Interval*. Moreover, a trapezoid graph is a *parallelogram graph* if every trapezoid is a parallelogram, and parallelogram graphs coincide with *bounded tolerance graphs* [2, 14]. These three graph classes are proper subclasses of trapezoid graphs, and they contain both interval graphs and permutation graphs as proper subclasses.

The time complexity of the recognition problem of these three graph classes was a longstanding open problem [3, 29], which was recently settled. While the problem is NP-complete for triangle graphs [24] and parallelogram graphs [27], an $O(n^2\bar{m})$ -time recognition algorithm has been given for simple-triangle graphs [25], where n and \bar{m} are the number of vertices and non-edges of the graph, respectively. This algorithm reduces the recognition problem of simple-triangle graph to the *linear-interval cover problem*, a problem of covering an associated bipartite graph by two chain graphs satisfying additional conditions. In [31], we showed an alternative algorithm for the linear-interval cover problem, but this does not improve the running time. Meanwhile, we showed in [32] a vertex ordering characterization that a graph is a simple-triangle graph if and only if there is a linear ordering of the vertices containing both an alternating orientation of the graph and a transitive orientation of the complement of the graph. Using this vertex ordering characterization, we show in this paper an $O(nm)$ -time recognition algorithm for simple-triangle graphs, where m is the number of edges of the graph.

Date: January 14, 2022.

2010 Mathematics Subject Classification. 68R10, 05C75, 05C85, 05C62.

Key words and phrases. Alternately orientable graphs, Cocomparability graphs, PI graphs, Recognition algorithm, Simple-triangle graphs, Vertex ordering characterization.

Our algorithm is shown in the next section, and correctness of the algorithm is proved in Section 3. We finally discuss our results and further research in Section 4.

2. THE RECOGNITION ALGORITHM

2.1. Preliminaries.

Notation. In this paper, we will deal only with finite graphs having no loops and multiple edges. Unless stated otherwise, graphs are assumed to be undirected, but we also deal with graphs having directed edges. We write uv for the *undirected edge* joining a vertex u and a vertex v , and we write (u, v) for the *directed edge* from u to v . For a graph $G = (V, E)$, we sometimes write $V(G)$ for the vertex set V of G and write $E(G)$ for the edge set E of G .

Let $G = (V, E)$ be an undirected graph. The *complement* of G is the graph $\overline{G} = (V, \overline{E})$ such that $uv \in \overline{E}$ if and only if $uv \notin E$ for any two vertices $u, v \in V$. A sequence of distinct vertices (v_0, v_1, \dots, v_k) is a *path from v_0 to v_k* in G if $v_0v_1, v_1v_2, \dots, v_{k-1}v_k \in E$. These edges are *the edges on the path*. The *length* of the path is the number k of the edges on the path. A path (v_0, v_1, \dots, v_k) is a *cycle* in G if in addition $v_kv_0 \in E$. The edges $v_0v_1, v_1v_2, \dots, v_{k-1}v_k$ and v_kv_0 are *the edges on the cycle*. The *length* of the cycle is the number $k + 1$ of the edges on the cycle. A *chord* of a path (cycle) is an edge joining two vertices that are not consecutive on the path (cycle). A path (cycle) is *chordless* if it contains no chords.

Let $G = (V, E)$ be an undirected graph. An *orientation* of G is a graph obtained from G by orienting each edge in E , that is, replacing each edge $uv \in E$ with either (u, v) or (v, u) . An *oriented graph* is an orientation of some graph. A *partial orientation* of G is a graph obtained from G by orienting each edge in a subset of E . A *partially oriented graph* is a partial orientation of some graph. Notice that a (partially) oriented graph contains no pair of edges (u, v) and (v, u) for some vertices u, v . We will denote a (partial) orientation of a graph only by its edge set when the vertex set is clear from the context.

Let $H = (V, F)$ be a (partially) oriented graph. A sequence of distinct vertices (v_0, v_1, \dots, v_k) is a *directed path from v_0 to v_k* in H if $(v_0, v_1), (v_1, v_2), \dots, (v_{k-1}, v_k) \in F$. A directed path (v_0, v_1, \dots, v_k) is a *directed cycle* in H if in addition $(v_k, v_0) \in F$. The edges on the path (cycle) and the length of the path (cycle) are defined analogously to the undirected case. A (partial) orientation of a graph is called *acyclic* if it contains no directed cycles. A *linear extension* (or topological sort) of an acyclic (partial) orientation F is a linear ordering of the vertices such that $(u, v) \in F$ if and only if u is a predecessor of v in the ordering. Let $F_s \subseteq F$ be a set of directed edges in H . The *reversal* F_s^{-1} of F_s is the set of directed edges obtained from F_s by reversing all the edges in F_s , that is, $F_s^{-1} = \{(u, v) : (v, u) \in F_s\}$.

Comparability graphs. An orientation F of a graph G is a *transitive orientation* if $(u, v) \in F$ and $(v, w) \in F$ then $(u, w) \in F$. A graph is a *comparability graph* if it has a transitive orientation. The complement of a comparability graph is a *cocomparability graph*. The class of cocomparability graphs contains the class of trapezoid graphs as a proper subclass [7, 8], and hence the complement of any simple-triangle graph has a transitive orientation. Note that every cocomparability graph contains no chordless cycle of length greater than or equal to 5 (see [3, 11] for example), and thus every chordless cycle of a cocomparability graph has length at most 4.

An orientation F of a graph G is *quasi-transitive* if $(u, v) \in F$ and $(v, w) \in F$ then $uw \in E(G)$ and therefore either $(u, w) \in F$ or $(w, u) \in F$. In other words, an orientation F of G is quasi-transitive if for any path of three vertices (u, v, w) in G , either $(u, v), (w, v) \in F$ or $(v, u), (v, w) \in F$. We can see that an orientation is transitive if and only if it is quasi-transitive and acyclic. Trivially, a graph having a transitive orientation also has a quasi-transitive orientation. The converse is also known to be true [12, 17], that is, if a graph G has a quasi-transitive orientation F that is not acyclic, then G has another quasi-transitive orientation F' that is also acyclic.

There is a well-known algorithm for recognizing comparability graphs and producing a transitive orientation of a graph, which takes $O(nm)$ time (see [13] for example). A linear-time algorithm is also known for producing a transitive orientation of a comparability graph [22]. To be precise, the algorithm produces a linear extension of the transitive orientation. However, even if the given graph is not a comparability graph, the linear-time algorithm may produce an orientation that is not transitive. Hence, to recognize comparability graphs, we

must verify transitivity of the orientation. The best known method for verifying transitivity uses Boolean matrix multiplication to test whether $A^2 = A$, where A is the adjacency matrix of the orientation F (see [29] for example). An alternative method for verifying transitivity, which takes $O(m^{3/2})$ time, is discussed in [29, Section 11.1.4].

Let \overline{G} be the complement of a graph G . The linear-time algorithm of [22] also can produce a linear extension of an orientation \overline{F} of \overline{G} such that \overline{F} is transitive if and only if G is a cocomparability graph. We show a method for verifying transitivity of \overline{F} in $O(nm)$ time. Recall that an orientation is transitive if and only if it is quasi-transitive and acyclic. Obviously, \overline{F} is acyclic. We can test whether \overline{F} is quasi-transitive by checking for any path of three vertices (u, v, w) in \overline{G} , either $(u, v), (w, v) \in \overline{F}$ or $(v, u), (v, w) \in \overline{F}$. The number of paths in G of three vertices is at most nm , and they can be found in $O(nm)$ time. Thus we have the following.

Theorem 1. *Cocomparability graphs can be recognized in $O(nm)$ time.*

Alternately orientable graphs. An orientation of a graph is *alternating* if it is (quasi-)transitive on every chordless cycle of length greater than or equal to 4, that is, the directions of the edges alternate. A graph is an *alternately orientable graph* [18] if it has an alternating orientation. It is clear from the definition that alternately orientable graphs generalize comparability graphs. A polynomial-time recognition algorithm is known for alternately orientable graphs [18].

We say a graph is an *alternately orientable cocomparability graph* if it is a cocomparability graph that has an alternating orientation. Obviously, alternately orientable cocomparability graphs can be recognized in polynomial time. It is also known that every alternately orientable cocomparability graph is a trapezoid graph but the converse is not true [9].

We say an orientation is an *acyclic alternating orientation* if it is alternating and acyclic, and we say a graph is *acyclic alternately orientable* if it has an acyclic alternating orientation. We can see that an alternating orientation is acyclic if it contains no directed cycles of length 3. Recall that a graph has a quasi-transitive orientation if and only if it has an orientation that is quasi-transitive and acyclic. It was conjectured that a similar statement might hold for alternating orientation, that is, a graph is alternately orientable if and only if it has an acyclic alternating orientation [18]. Later, however, a counterexample was provided [19]. Thus, the class of acyclic alternately orientable graphs is a proper subclass of alternately orientable graphs. Moreover, the recognition problem is NP-complete for acyclic alternately orientable graphs.

Theorem 2. *The recognition problem is NP-complete for acyclic alternately orientable graphs.*

Proof. We can verify in polynomial time whether an orientation is alternating [18]. Since testing acyclicity of an orientation takes linear time, the recognition problem is in NP. We show a polynomial-time reduction from the non-betweenness problem, which is known to be NP-complete [16]. Given a finite set A and a collection C of ordered triples (a_i, a_j, a_k) of distinct elements of A , the non-betweenness problem is to decide whether there is a bijection $f : A \rightarrow \{1, 2, \dots, |A|\}$ such that for each $(a_i, a_j, a_k) \in C$, either $f(a_i), f(a_k) < f(a_j)$ or $f(a_j) < f(a_i), f(a_k)$. Let G be a graph such that

$$V(G) = \{v_i : a_i \in A\} \cup \{u_i, w_i : t_i \in C\} \text{ and}$$

$$E(G) = \{v_i v_j : 1 \leq i < j \leq |A|\} \cup \{u_h w_h, u_h v_j, w_h v_i, w_h v_k : t_h = (a_i, a_j, a_k) \in C\}.$$

Clearly, we can construct the graph G in time polynomial in n and m . We can also see from the construction that the set of chordless cycles $\{(u_h, w_h, v_i, v_j), (u_h, w_h, v_k, v_j) : t_h = (a_i, a_j, a_k) \in C\}$ contains all the chordless cycles of length greater than or equal to 4. Thus an orientation F of G is alternating if and only if for any three vertices v_i, v_j, v_k with $t_h = (a_i, a_j, a_k) \in C$, either $(v_i, v_j), (v_k, v_j) \in F$ or $(v_j, v_i), (v_j, v_k) \in F$. Therefore, G has an acyclic alternating orientation if and only if there is a bijection $f : A \rightarrow \{1, 2, \dots, |A|\}$ such that for each $(a_i, a_j, a_k) \in C$, either $f(a_i), f(a_k) < f(a_j)$ or $f(a_j) < f(a_i), f(a_k)$. Thus we have the theorem \square

The vertex ordering characterization of simple-triangle graphs. Now, we show the vertex ordering characterization of simple-triangle graphs.

Theorem 3 ([32]). *A graph G is a simple-triangle graph if and only if there is an (acyclic) alternating orientation F of G and a transitive orientation \overline{F} of the complement \overline{G} of G such that $F \cup \overline{F}$ is acyclic. Moreover, if G is a simple-triangle graph, then for any transitive orientation \overline{F} of \overline{G} , there is an (acyclic) alternating orientation F of G such that $F \cup \overline{F}$ is acyclic. An acyclic orientation of a complete graph is equivalent to the linear ordering of the vertices of the graph. The orientation $F \cup \overline{F}$ is called an apex ordering of a simple-triangle graph since it coincides with the ordering of the apices of the triangles in the representation.*

Theorem 3 implies that a simple-triangle graph is an (acyclic) alternately orientable cocomparability graph, but the converse is not known to be true [32]. We also note that Theorem 3 is similar to a well-known fact that a graph is a permutation graph if and only if it is both a comparability graph and a cocomparability graph [28].

2.2. The recognition algorithm. Let F be an alternating orientation of a graph G , and let \overline{F} be a transitive orientation of the complement \overline{G} of G . Suppose that $F \cup \overline{F}$ is not acyclic. It is well known that an orientation of a complete graph is not acyclic if and only if it contains a directed cycle of length 3. Each directed cycle of length 3 in $F \cup \overline{F}$ consists of either three edges in F or two edges in F with one edge in \overline{F} since \overline{F} is a transitive orientation. We will refer to a directed cycle (a, b, c) of length 3 in $F \cup \overline{F}$ as a Δ -obstruction if $(a, b), (b, c) \in F$ and $(c, a) \in \overline{F}$.

It is clear from Theorem 3 that if a graph G is a simple-triangle graph, then there is a pair of an acyclic alternating orientation F of G and a transitive orientation \overline{F} of \overline{G} such that $F \cup \overline{F}$ contains no Δ -obstructions. Conversely, a graph G is a simple-triangle graph if for some transitive orientation \overline{F} of \overline{G} , there is an acyclic alternating orientation F of G such that $F \cup \overline{F}$ contains no Δ -obstructions. We can also have the following.

Theorem 4. *Let \overline{F} be a transitive orientation of \overline{G} . If G has a (not necessarily acyclic) alternating orientation F such that $F \cup \overline{F}$ contains no Δ -obstructions, then G also has an acyclic alternating orientation F' of G such that $F' \cup \overline{F}$ contains no Δ -obstructions, that is, G is a simple-triangle graph.*

Our recognition algorithm is due to this structural characterization, and we will prove the theorem when we show the correctness of the algorithm.

Figure 1 shows our algorithm to recognize simple-triangle graphs. The algorithm finds an apex ordering of the given graph if it is a simple-triangle graph or report that the graph is not a simple-triangle graph.

| | |
|----------------|---|
| Input: | A graph G . |
| Output: | An apex ordering of G if G is a simple-triangle graph or report that G is not a simple-triangle graph. |
| Step 1: | Compute a transitive orientation \overline{F} of the complement \overline{G} of G . If G has no transitive orientations, then report that G is not a simple-triangle graph. |
| Step 2: | Compute a partial orientation F of G that satisfies the following three conditions: <ul style="list-style-type: none"> (1) $F \cup \overline{F}$ contains no Δ-obstructions, (2) The edges on every chordless cycle of length 4 are oriented so that the directions of the edges alternate (Notice that the length of every chordless cycle in G is at most 4 since G is a cocomparability graph), and (3) Every edge remains undirected if it is not on a chordless cycle of length 4. If G has no such orientations, then report that G is not a simple-triangle graph. |
| Step 3: | Choose a vertex v of G . Let F_v be the set of all the edges $(w, u) \in F$ such that (u, v, w) form a directed cycle in F . Reverse the orientation of all the edges in F_v to compute another partial orientation F' of G , that is, $F' = (F - F_v) \cup F_v^{-1}$. Repeat this procedure for all the vertices of G . We denote the resultant orientation by F'' . |
| Step 4: | Output a linear extension of $F'' \cup \overline{F}$. |

FIGURE 1. The recognition algorithm for simple-triangle graphs

We remark that in **Step 2** of the algorithm, we keep an edge undirected if it is not on a chordless cycle of length 4, since it is needed to do **Step 3** and **Step 4** correctly. If we do not need to find an apex ordering of the given graph, in **Step 2** we only have to compute an alternating orientation F of G such that $F \cup \overline{F}$ contains no Δ -obstructions.

We will prove the correctness of the algorithm in the next section. Notice that in order to prove the correctness, it suffices to show that when **Step 3** is finished, $F'' \cup \overline{F}$ is acyclic.

Details of the algorithm. In the rest of this section, we show that the algorithm runs in $O(nm)$ time. We assume without loss of generality that the given graph is connected and $n \leq m$, since otherwise we apply the algorithm to each connected component.

In **Step 1**, we use the linear-time algorithm of [22], which produces a linear extension of a transitive orientation \overline{F} of the complement \overline{G} if the given graph G is a cocomparability graph. As shown in Theorem 1, we can verify transitivity of \overline{F} in $O(nm)$ time, and hence **Step 1** can be performed in the same time bound.

In **Step 2**, we construct the auxiliary graph G^+ of G that is bipartite if and only if G is alternately orientable. Then, we make a 2CNF formula ϕ , an instance of the 2-satisfiability problem, from G^+ . The partial orientation F of G can be obtained from a truth assignment τ of ϕ . When G^+ is not bipartite or ϕ cannot be satisfied, the algorithm reports that G is not a simple-triangle graph.

The vertices of the auxiliary graph G^+ are all the ordered pairs (u, v) with $uv \in E(G)$. Each vertex (u, v) of G^+ is adjacent to the vertex (v, u) . Each vertex (u, v) of G^+ is also adjacent to every vertex (v, w) such that in G , the vertices (u, v, w) form a path of three vertices on a chordless cycle of length 4. We can see that G is alternately orientable if and only if G^+ is bipartite, and an alternating orientation of G can be obtained from one color class of G^+ . The number of vertices of G^+ is $2m$, and the number of edges of G^+ is at most $m + 2nm$ since the number of paths of three vertices in G is at most nm . Hence we can test the bipartiteness of G^+ in $O(nm)$ time.

Now, we show that the auxiliary graph G^+ can be constructed from G in $O(nm)$ time. The *neighborhood* of a vertex v of G is the set $N(v) = \{u \in V(G) : uv \in E(G)\}$. We define that the *upper set* of v is the set $U(v) = \{u \in V(G) : (v, u) \in \overline{F}\}$ and the *lower set* of v is the set $L(v) = \{u \in V(G) : (u, v) \in \overline{F}\}$. Let uv be an edge of G . If $N(u) \cap U(v) \neq \emptyset$ and $N(v) \cap U(u) \neq \emptyset$, then for any two vertices $w \in N(u) \cap U(v)$ and $z \in N(v) \cap U(u)$, we have $wz \in E(G)$, that is, (u, v, z, w) is a chordless cycle of length 4; otherwise either $(w, z) \in \overline{F}$ or $(z, w) \in \overline{F}$, but if $(w, z) \in \overline{F}$ then $(v, w) \in \overline{F}$ implies $(v, z) \in \overline{F}$, contradicting $z \in N(v)$, and if $(z, w) \in \overline{F}$ then $(u, z) \in \overline{F}$ implies $(u, w) \in \overline{F}$, contradicting $w \in N(u)$. Similarly, for any two vertices $w \in N(u) \cap L(v)$ and $z \in N(v) \cap L(u)$, we have that (u, v, z, w) is a chordless cycle of length 4. Conversely, each chordless cycle of length 4 has two edges uv and zw such that $N(u) \cap U(v) \neq \emptyset$, $N(v) \cap U(u) \neq \emptyset$, $N(z) \cap L(w) \neq \emptyset$, and $N(w) \cap L(z) \neq \emptyset$.

Therefore, we can construct the auxiliary graph G^+ by the following method. (1) For each edge $uv \in E(G)$, test whether $N(u) \cap U(v) \neq \emptyset$ and $N(v) \cap U(u) \neq \emptyset$ [resp. $N(u) \cap L(v) \neq \emptyset$ and $N(v) \cap L(u) \neq \emptyset$]. (2) If it is, then for any two vertices $w \in N(u) \cap U(v)$ and $z \in N(v) \cap U(u)$ [resp. $w \in N(u) \cap L(v)$ and $z \in N(v) \cap L(u)$], add to G^+ the edge joining (u, v) and (v, z) , the edge joining (z, v) and (v, u) , the edge joining (v, u) and (u, w) , and the edge joining (w, u) and (u, v) . (3) Finally, for each edge $uv \in E(G)$, add to G^+ the edge joining (u, v) and (v, u) . The first procedure takes $O(n)$ time for each edge, and the second procedure takes $O(nm)$ time in total since the number of paths of three vertices in G is at most nm . The third procedure takes $O(m)$ time in total, and hence the auxiliary graph G^+ can be constructed in $O(nm)$ time.

We next construct the 2CNF formula ϕ from G^+ . Recall that each vertex (u, v) of G^+ is adjacent to (v, u) . A connected component of G^+ consists of only two vertices (u, v) and (v, u) if and only if the edge uv of G is not on a chordless cycle of length 4. We remove such components from G^+ , and let c_1, c_2, \dots, c_k be the remaining components. To each component c_i , we assign the Boolean variable x_i . Since we assume that G^+ is bipartite, the vertices of each component can be partitioned into two color classes. We assign the literal x_i to the vertices of one color class of c_i , and we assign the literal $\overline{x_i}$ (the negation of x_i) to the vertices of the other color class of c_i .

Notice that a vertex (u, v) of G^+ has its literal if and only if the edge uv of G is on a cycle of length 4. Let $l_{(u,v)}$ be the literal assigned to a vertex (u, v) of G^+ . The 2CNF formula ϕ consists of all the clauses $(l_{(u,v)} \vee l_{(v,w)})$

such that the vertices (u, v, w) in G form a path of three vertices with $(w, u) \in \overline{F}$. When no literal is assigned to (u, v) or (v, w) , the 2CNF formula ϕ does not contain the clause.

Let τ be a truth assignment of the variables in ϕ . Notice that $l_{(u,v)} = 0$ in τ if and only if $l_{(v,u)} = 1$ since each vertex (u, v) of G^+ is adjacent to (v, u) . We obtain the partial orientation F of G from τ by orienting each edge $uv \in E(G)$ as $(u, v) \in F$ if $l_{(u,v)} = 0$ in τ . It is obvious from the construction of ϕ that a truth assignment τ satisfies ϕ if and only if $F \cup \overline{F}$ contains no Δ -obstructions. We can also see that the edges on every chordless cycle of length 4 is alternately oriented and the other edges remain undirected.

The 2CNF formula ϕ has at most $2m$ Boolean variables and at most nm clauses. We can also see that ϕ can be constructed in $O(nm)$ time, since all the paths of three vertices can be found in $O(nm)$ time. Since a satisfying truth assignment of a 2CNF formula ϕ can be computed in time linear to the size of ϕ (see [1] for example), **Step 2** can be performed in $O(nm)$ time.

It is easy to see that **Step 3** and **Step 4** can be performed in $O(nm)$ time and $O(n^2)$ time, respectively. Now, we have the following.

Theorem 5. *Simple-triangle graphs can be recognized in $O(nm)$ time, and an apex ordering of a simple-triangle graph can be computed in the same time bound.*

From **Step 1** and **Step 2** of the algorithm, we also have the following as a byproduct.

Theorem 6. *Alternately orientable cocomparability graphs can be recognized in $O(nm)$ time, and alternating orientation of a cocomparability graph can be computed in the same time bound.*

3. CORRECTNESS OF THE ALGORITHM

In this section, we prove correctness of our algorithm. In order to prove the correctness, it suffices to show that when **Step 3** is finished, $F'' \cup \overline{F}$ is acyclic, where F'' is the resultant orientation of **Step 3**.

Recall that the graph G is an alternately orientable graph and the complement \overline{G} of G is a comparability graph with a transitive orientation \overline{F} . Since G is a cocomparability graph, the length of every chordless cycle in G is at most 4. Recall also that G has a partial orientation F that satisfies the following three conditions: (1) $F \cup \overline{F}$ contains no Δ -obstructions, (2) the edges on every chordless cycle of length 4 are oriented so that the directions of the edges alternate, and (3) the other edges remain undirected. For a vertex v of G , let F_v be the set of all the edges $(w, u) \in F$ such that (u, v, w) is a directed cycle in F , and let F' be the partial orientation of G obtained from F by reversing the orientation of all the edges in F_v , that is, $F' = (F - F_v) \cup F_v^{-1}$.

The outline of the proof is as follows. An *alternating $2k$ -cycle* in $F \cup \overline{F}$ with $k \geq 2$ is a directed cycle $(a_0, b_0, a_1, b_1, \dots, a_{k-1}, b_{k-1})$ of length $2k$ with $(a_i, b_i) \in F$ and $(b_i, a_{i+1}) \in \overline{F}$ for any $i = 0, 1, \dots, k-1$ (indices are modulo k). See Figure 2 for example. We first show that

- F is acyclic if and only if $F \cup \overline{F}$ contains no alternating 6-cycles.

We next show the following:

- $F' \cup \overline{F}$ contains no alternating 6-cycles having the vertex v ,
- the reversing the direction of the edges in F_v generates no alternating 6-cycles,
- the directions of the edges still alternate in F' on every chordless cycle of length 4, and
- $F' \cup \overline{F}$ is still contains no Δ -obstructions.

Thus continuing in this way for each vertex, we obtain the acyclic orientation F'' such that $F'' \cup \overline{F}$ contains no Δ -obstructions, that is, $F'' \cup \overline{F}$ is acyclic.

3.1. Definitions and facts. We begin to state some definitions and facts.

Claim 7. *Let a_0, b_0, a_1, b_1 be four vertices of G . If $a_0b_0, a_1b_1 \in E(G)$ and $(a_0, b_1), (a_1, b_0) \in \overline{F}$, then $a_0a_1, b_0b_1 \in E(G)$, that is, (a_0, a_1, b_1, b_0) is a chordless cycle of length 4.*

Proof. Recall that \overline{F} is a transitive orientation. If $(a_0, a_1) \in \overline{F}$, then $(a_1, b_0) \in \overline{F}$ implies $(a_0, b_0) \in \overline{F}$, contradicting $a_0b_0 \in E(G)$. If $(a_1, a_0) \in \overline{F}$, then $(a_0, b_1) \in \overline{F}$ implies $(a_1, b_1) \in \overline{F}$, contradicting $a_1b_1 \in E(G)$. Thus




FIGURE 2. An alternating 6-cycles. An arrow $a \rightarrow b$ denotes edge $(a, b) \in F$, and a dashed arrow $a \dashrightarrow b$ denotes edge $(a, b) \in \overline{F}$.

$a_0a_1 \in E(G)$. Similarly, if $(b_0, b_1) \in \overline{F}$, then $(a_1, b_0) \in \overline{F}$ implies $(a_1, b_1) \in \overline{F}$, contradicting $a_1b_1 \in E(G)$. If $(b_1, b_0) \in \overline{F}$, then $(a_0, b_1) \in \overline{F}$ implies $(a_0, b_0) \in \overline{F}$, contradicting $a_0b_0 \in E(G)$. Thus $b_0b_1 \in E(G)$. \square

Suppose that $F \cup \overline{F}$ contains an alternating 4-cycle (a_0, b_0, a_1, b_1) with $(a_0, b_0), (a_1, b_1) \in F$ and $(b_0, a_1), (b_1, a_0) \in \overline{F}$. We have from Claim 7 that (a_0, a_1, b_1, b_0) is a chordless cycle of length 4. The directions of the edges on this cycle must alternate in F , but $(a_0, b_0), (a_1, b_1) \in F$, a contradiction. Thus $F \cup \overline{F}$ contains no alternating 4-cycles. An *alternating 4-anticycle* of $F \cup \overline{F}$ is a subgraph consisting of four vertices a_0, b_0, a_1, b_1 with $(a_0, b_0), (a_1, b_1) \in F$ and $(a_0, b_1), (a_1, b_0) \in \overline{F}$. We have from Claim 7 that (a_0, a_1, b_1, b_0) is a chordless cycle of length 4. The directions of the edges on this cycle must alternate in F , but $(a_0, b_0), (a_1, b_1) \in F$, a contradiction. Thus $F \cup \overline{F}$ contains no alternating 4-anticycles.

The following claim states the direction of chords of an alternating 6-cycle.

Claim 8. *If $F \cup \overline{F}$ contains an alternating 6-cycle $(a_0, b_0, a_1, b_1, a_2, b_2)$ with $(a_i, b_i) \in F$ and $(b_i, a_{i+1}) \in \overline{F}$ for any $i = 0, 1, 2$ (indices are modulo 3), then $(b_2, a_1), (b_2, b_0), (a_0, a_1), (b_0, a_2), (b_0, b_1), (a_1, a_2), (b_1, a_0), (b_1, b_2), (a_2, a_0) \in F$. See Figure 2.*

Proof. If $(b_2, a_1) \in \overline{F}$, then (a_1, b_1, a_2, b_2) is an alternating 4-cycle, a contradiction. If $(a_1, b_2) \in \overline{F}$, then $(b_0, a_1), (b_2, a_0) \in \overline{F}$ implies $(b_0, a_0) \in \overline{F}$, contradicting $(a_0, b_0) \in F$. Thus $b_2a_1 \in E(G)$. We have from Claim 7 that (a_0, b_0, b_2, a_1) is a chordless cycle of length 4. Since the directions of the edges on this cycle alternate in F , we have $(b_2, a_1), (b_2, b_0), (a_0, a_1) \in F$. By similar arguments, we also have $(b_0, a_2), (b_0, b_1), (a_1, a_2) \in F$ and $(b_1, a_0), (b_1, b_2), (a_2, a_0) \in F$. \square

Recall that for any directed edge $(u, v) \in F$, there are another two vertices w, z of G such that (u, v, w, z) is a chordless cycle of length 4, since an edge of G remains undirected if it is not on a chordless cycle of length 4.

Claim 9. *If there are four vertices a, b, c, d with $(a, b), (b, c), (c, d) \in F$ and $(d, a) \in \overline{F}$, then there are another two vertices e, f with $(b, e), (f, c), (f, e) \in F$ and $bf, ce \in E(\overline{G})$. In addition,*

- $(a, f), (d, b), (d, f), (c, a), (e, a), (e, d) \in F$ and
- *there is an alternating 6-cycle consisting of the vertices a, b, c, d, e, f .*

Proof. Since (b, c) is oriented in F , there are another two vertices e, f with $(b, e), (f, c), (f, e) \in F$ and $bf, ce \in E(\overline{G})$. If $(a, c) \in \overline{F}$, then $(d, a) \in \overline{F}$ implies $(d, c) \in \overline{F}$, contradicting $(c, d) \in F$. If $(c, a) \in \overline{F}$, then (a, b, c) is a Δ -obstruction, a contradiction. Thus $ac \in E(G)$. Similarly, if $(b, d) \in \overline{F}$, then $(d, a) \in \overline{F}$ implies $(b, a) \in \overline{F}$, contradicting $(a, b) \in F$. If $(d, b) \in \overline{F}$, then (b, c, d) is a Δ -obstruction, a contradiction. Thus $bd \in E(G)$.

If $(e, a) \in \overline{F}$, then (a, b, e) is a Δ -obstruction, a contradiction. Suppose $(a, e) \in \overline{F}$. If $af \in E(G)$, then (a, b, e, f) is a chordless cycle of length 4 with $(a, b), (b, e) \in F$, contradicting the definition of F . Thus

$af \in E(\overline{G})$. If $(f, a) \in \overline{F}$, then $(a, e) \in \overline{F}$ implies $(f, e) \in \overline{F}$, contradicting $(f, e) \in F$. If $(a, f) \in \overline{F}$, then $(d, a) \in \overline{F}$ implies $(d, f) \in \overline{F}$, but then (f, c, d) is a Δ -obstruction, a contradiction. Thus $ae \in E(G)$.

If $af \in E(\overline{G})$, then (a, e, f, c) is a chordless cycle of length 4. Since $(f, c), (f, e) \in F$, we have $(a, c), (a, e) \in F$, but then (a, c, d) is a Δ -obstruction, a contradiction. Thus $af \in E(G)$. If $(d, f) \in \overline{F}$, then (f, c, d) is a Δ -obstruction, a contradiction. If $(f, d) \in \overline{F}$, then $(d, a) \in \overline{F}$ implies $(f, a) \in \overline{F}$, contradicting $af \in E(G)$. Thus $df \in E(G)$. If $de \in E(\overline{G})$, then (d, b, e, f) is a chordless cycle of length 4. Since $(b, e), (f, e) \in F$, we have $(b, d), (f, d) \in F$, but then (a, b, d) is a Δ -obstruction, a contradiction. Thus $de \in E(G)$.

Since (a, b, d, f) is a chordless cycle of length 4 and $(a, b) \in F$, we have $(a, f), (d, b), (d, f) \in F$. Since (a, e, d, c) is a chordless cycle of length 4 and $(c, d) \in F$, we also have $(c, a), (e, a), (e, d) \in F$. We can verify that for any directions are assigned to the edges $bf, ce \in E(\overline{G})$, there is an alternating 6-cycle consisting of the vertices a, b, c, d, e, f . \square

Claim 10. *If there is a directed cycle (a, b, c) in F , then there is an alternating 6-cycle of $F \cup \overline{F}$ containing the vertices a, b, c .*

Proof. Since (a, b) is oriented in F , there are another two vertices d, e with $(a, d), (e, b), (e, d) \in F$ and $ae, bd \in E(\overline{G})$. We have $c \neq d$ from $bc \in E(G)$ and $bd \in E(\overline{G})$. We also have $c \neq e$ from $(b, c), (e, b) \in F$. If $(a, e) \in \overline{F}$, then we have the claim from Claim 9 since (e, b, c, a) is a directed cycle of length 4 in $F \cup \overline{F}$. If $(d, b) \in \overline{F}$, then we also have the claim from Claim 9 since (b, c, a, d) is a directed cycle of length 4 in $F \cup \overline{F}$. Therefore, we assume $(e, a), (b, d) \in \overline{F}$.

Since (b, c) is oriented in F , there are another two vertices f, g with $(b, f), (g, c), (g, f) \in F$ and $bg, cf \in E(\overline{G})$. We have $a \neq f$ from $(a, b), (b, f) \in F$. We also have $a \neq g$ from $ab \in E(G)$ and $bg \in E(\overline{G})$. If $(b, g) \in \overline{F}$, then we have the claim from Claim 9 since (g, c, a, b) is a directed cycle of length 4 in $F \cup \overline{F}$. If $(f, c) \in \overline{F}$, then we also have the claim from Claim 9 since (c, a, b, f) is a directed cycle of length 4 in $F \cup \overline{F}$. Therefore, we assume $(g, b), (c, f) \in \overline{F}$.

We have $d \neq f$ from $bd \in E(\overline{G})$ and $bf \in E(G)$. We also have $d \neq g$ from $(b, d), (g, b) \in \overline{F}$. Similarly, we have $e \neq f$ from $(e, b), (b, f) \in F$. We also have $e \neq g$ from $be \in E(G)$ and $bg \in E(\overline{G})$. Therefore, the seven vertices a, b, c, d, e, f, g are distinct.

If $(e, f) \in \overline{F}$, then four vertices b, f, e, d form an alternating 4-anticycle, a contradiction. If $(f, e) \in \overline{F}$, then (e, b, f) is a Δ -obstruction, a contradiction. Thus $ef \in E(G)$. Now, we have from Claim 7 that (a, c, e, f) is a chordless cycle of length 4. Since $(c, a) \in F$, we have $(c, e), (f, a), (f, e) \in F$.

If $(d, c) \in \overline{F}$, then $(b, d) \in \overline{F}$ implies $(b, c) \in \overline{F}$, contradicting $(b, c) \in F$. If $(c, d) \in \overline{F}$, then four vertices c, a, e, d form an alternating 4-anticycle, a contradiction. Thus $cd \in E(G)$. Now, we have from Claim 7 that (c, d, f, b) is a chordless cycle of length 4. Since $(b, c), (b, f) \in F$, we have $(d, c), (d, f) \in F$.

Now, we can see that (a, b, d, c, f, e) is an alternating 6-cycle in $F \cup \overline{F}$. \square

Claim 11. *If there is a directed cycle (a, b, c) in F , then there is another vertex d with $(c, d), (d, b) \in F$ and either $(a, d) \in \overline{F}$ or $(d, a) \in \overline{F}$.*

Proof. We have the claim from Claims 10 and 8. \square

3.2. Main part of the proof. Now, we show the following series of claims, which proves the correctness of the algorithm. It is immediate from Claim 8 that if $F \cup \overline{F}$ contains an alternating 6-cycle, then F contains a directed cycle of length 3. The following claim states that the converse is also true. (Recall that F contains undirected edges. Thus F may contain directed cycles of length greater than 3, and Claim 10 does not imply the converse.)

Claim 12. *If F is not acyclic, then $F \cup \overline{F}$ contains an alternating 6-cycle.*

Proof. Before proving the claim, we introduce the *middle edge* of a chordless cycle of length 4. Let $C_4 = (a, b, c, d)$ be a cycle of length 4 in G with $(a, b), (a, d), (c, b), (c, d) \in F$ and $(a, c), (b, d) \in \overline{F}$. We call the edge (c, b) the *middle edge* of C_4 in $F \cup \overline{F}$. We also say an edge is a middle edge in $F \cup \overline{F}$ if there is a chordless

cycle of length 4 such that the edge is the middle edge of the cycle. Notice that for any edge (u, v) on C_4 , there is the directed path from u to v in $F \cup \overline{F}$ consisting of the edges in \overline{F} and the middle edge. For example, there is the directed path (a, c, b, d) for the edge (a, d) .

Recall that each directed edge in F is on a cycle of length 4. Thus if F contains a directed cycle, then $F \cup \overline{F}$ contains a directed cycle $C = (v_0, v_1, \dots, v_{k-1})$ with $k \geq 3$ consisting of the edges in \overline{F} and the middle edges in $F \cup \overline{F}$.

We now show that if (v_i, v_{i+1}) and (v_{i+1}, v_{i+2}) are middle edges in $F \cup \overline{F}$ for some i (indices are modulo k), then either $(v_i, v_{i+2}) \in F$ or $(v_{i+2}, v_i) \in F$. In addition, we show that if $(v_i, v_{i+2}) \in F$, then (v_i, v_{i+2}) is a middle edge in $F \cup \overline{F}$. Since (v_i, v_{i+1}) is a middle edge, there are another two vertices a, b with $(v_i, a), (b, v_{i+1}), (b, a) \in F$ and $(b, v_i), (v_{i+1}, a) \in \overline{F}$. Similarly, since (v_{i+1}, v_{i+2}) is a middle edge, there are another two vertices c, d with $(v_{i+1}, c), (d, v_{i+2}), (d, c) \in F$ and $(v_{i+2}, c), (d, v_{i+1}) \in \overline{F}$. We have $v_i \neq c$ from $(v_i, v_{i+1}), (v_{i+1}, c) \in F$. We also have $v_i \neq d$ from $v_i v_{i+1} \in E(G)$ and $v_{i+1} d \in E(\overline{G})$. Similarly, we have $a \neq v_{i+2}$ from $av_{i+1} \in E(\overline{G})$ and $v_{i+1} v_{i+2} \in E(G)$. We also have $a \neq c$ from $av_{i+1} \in E(\overline{G})$ and $v_{i+1} c \in E(G)$. Moreover, we have $a \neq d$ from $(v_{i+1}, a), (d, v_{i+1}) \in \overline{F}$. By similar arguments, we have $b \neq v_{i+2}, c, d$. Therefore, the seven vertices $v_i, v_{i+1}, v_{i+2}, a, b, c, d$ are distinct.

If $(b, c) \in \overline{F}$, then four vertices v_{i+1}, c, b, a form an alternating 4-anticycle, a contradiction. If $(c, b) \in \overline{F}$, then (b, v_{i+1}, c) is a Δ -obstruction, a contradiction. Thus $bc \in E(G)$. If $(v_i, v_{i+2}) \in \overline{F}$, then $(b, v_i), (v_{i+2}, c) \in \overline{F}$ implies $(b, c) \in \overline{F}$, contradicting $bc \in E(G)$. If $(v_{i+2}, v_i) \in \overline{F}$, then (v_i, v_{i+1}, v_{i+2}) is a Δ -obstruction, a contradiction. Thus $v_i v_{i+2} \in E(G)$. Now, we have from Claim 7 that (v_i, v_{i+2}, b, c) is a chordless cycle of length 4, and hence the edge $v_i v_{i+2}$ is oriented in F . When $(v_i, v_{i+2}) \in F$, the edge (v_i, v_{i+2}) is the middle edge of the cycle (v_i, v_{i+2}, b, c) .

We have from Claim 10 that if $(v_i, v_{i+1}), (v_{i+1}, v_{i+2}), (v_{i+2}, v_i) \in F$, then $F \cup \overline{F}$ contains an alternating 6-cycle, and we have the claim. Thus we can assume that the directed cycle C contains no two middle edges that are consecutive on C . Obviously, if $(v_i, v_{i+1}), (v_{i+1}, v_{i+2}) \in \overline{F}$ for some i (indices are modulo k), then $(v_i, v_{i+2}) \in \overline{F}$. Therefore, we can assume that C contains no two edges in \overline{F} that are consecutive on C , that is, C is an alternating cycle.

It remains to show that if $F \cup \overline{F}$ contains an alternating $2k$ -cycle with $k \geq 4$, then $F \cup \overline{F}$ also contains an alternating 6-cycle. Suppose that $F \cup \overline{F}$ contains an alternating $2k$ -cycle with $k \geq 4$ but it contains no alternating 6-cycles. Let $(a_0, b_0, a_1, b_1, \dots, a_{k-1}, b_{k-1})$ be such an alternating $2k$ -cycle with $(a_i, b_i) \in F$ and $(b_i, a_{i+1}) \in \overline{F}$ for any $i = 0, 1, \dots, k-1$ (indices are modulo k). We assume without loss of generality that the length of this cycle is minimal, that is, $F \cup \overline{F}$ contains no alternating cycle of length smaller than $2k$. Since $k \geq 4$, the vertices $a_0, b_0, a_1, b_1, a_2, b_2, a_3, b_3$ are distinct. If $(b_2, a_1) \in \overline{F}$, then (a_1, b_1, a_2, b_2) is an alternating 4-cycle, a contradiction. If $(b_0, a_3) \in \overline{F}$, then $(a_0, b_0, a_3, b_3, \dots, a_{k-1}, b_{k-1})$ is an alternating $2(k-2)$ -cycle, a contradiction. If $(a_1, b_2) \in \overline{F}$, then $(b_0, a_1), (b_2, a_3) \in \overline{F}$ implies $(b_0, a_3) \in \overline{F}$, a contradiction. If $(a_3, b_0) \in \overline{F}$, then $(b_2, a_3), (b_0, a_1) \in \overline{F}$ implies $(b_2, a_1) \in \overline{F}$, a contradiction. Thus $a_1 b_2, b_0 a_3 \in E(G)$. Now, we have from Claim 7 that (a_1, b_2, b_0, a_3) is a chordless cycle of length 4. If $(a_1, b_2), (a_1, a_3), (b_0, b_2), (b_0, a_3) \in F$, then $(a_0, b_0, a_1, b_2, a_3, b_3, \dots, a_{k-1}, b_{k-1})$ is an alternating $2(k-1)$ -cycle, a contradiction. If $(b_2, a_1), (a_3, a_1), (b_2, b_0), (a_3, b_0) \in F$, then $(a_1, b_1, a_2, b_2, a_3, b_0)$ is an alternating 6-cycle, a contradiction. \square

Recall that F_v is the set of all the edges $(w, u) \in F$ such that (u, v, w) is a directed cycle in F . Recall also that F' is the partial orientation of G obtained from F by reversing the orientation of all the edges in F_v , that is, $F' = (F - F_v) + F_v^{-1}$. The following claim states that $F' \cup \overline{F}$ contains no alternating 6-cycles having the vertex v .

Claim 13. *There is no alternating 6-cycles of $F' \cup \overline{F}$ having the vertex v .*

Proof. We prove the claim by contradiction. Suppose that $F' \cup \overline{F}$ contains an alternating 6-cycle $(a_0, b_0, a_1, b_1, a_2, b_2)$ with $(a_i, b_i) \in F'$ and $(b_i, a_{i+1}) \in \overline{F}$ for any $i = 0, 1, 2$ (indices are modulo 3). Suppose $v = a_0$. It is obvious that $(a_0, b_0), (a_2, b_2) \notin F_v^{-1}$, that is, $(a_0, b_0), (a_2, b_2) \in F$. If $(a_1, b_1) \notin F_v^{-1}$, then we have from Claim 8 that $(a_0, a_1), (a_1, b_1), (b_1, a_0) \in F$, and then $(a_1, b_1) \in F_v$, a contradiction. If $(a_1, b_1) \in F_v^{-1}$, then $(a_0, b_1), (b_1, a_1), (a_1, a_0) \in F$, but then $F \cup \overline{F}$ contains an alternating 4-cycle (a_0, b_1, a_2, b_2) , a contradiction. Thus $v \neq a_0$. By similar arguments, we have $v \neq a_1, a_2$.

Suppose $v = b_0$. It is obvious that $(a_0, b_0), (a_1, b_1) \notin F_v^{-1}$, that is, $(a_0, b_0), (a_1, b_1) \in F$. If $(a_2, b_2) \notin F_v^{-1}$, then we have from Claim 8 that $(b_0, a_2), (a_2, b_2), (b_2, b_0) \in F$, and then $(a_2, b_2) \in F_v$, a contradiction. If $(a_2, b_2) \in F_v^{-1}$, then $(b_0, b_2), (b_2, a_2), (a_2, b_0) \in F$, but then $F \cup \overline{F}$ contains an alternating 4-cycle (a_2, b_0, a_1, b_1) , a contradiction. Thus $v \neq b_0$. By similar arguments, we have $v \neq b_1, b_2$. \square

The following claim states that the reversing the orientation of the edges in F_v generates no alternating 6-cycles.

Claim 14. *No edges in F_v^{-1} is an edge of any alternating 6-cycle of $F' \cup \overline{F}$.*

Proof. We prove the claim by contradiction. Suppose that $F' \cup \overline{F}$ contains an alternating 6-cycle $(a_0, b_0, a_1, b_1, a_2, b_2)$ with $(a_i, b_i) \in F'$ and $(b_i, a_{i+1}) \in \overline{F}$ for any $i = 0, 1, 2$ (indices are modulo 3). We have from Claim 13 that the vertex v is not on the alternating 6-cycle. If $(a_0, b_0), (a_1, b_1) \in F_v^{-1}$, then $(v, b_0), (b_0, a_0), (a_0, v) \in F$ and $(v, b_1), (b_1, a_1), (a_1, v) \in F$, but then (a_1, v, b_0) is a Δ -obstruction in $F \cup \overline{F}$, a contradiction. Thus we can see that at most one edge on the alternating 6-cycle is in F_v^{-1} .

We assume without loss of generality that $(a_0, b_0) \in F_v^{-1}$ and $(a_1, b_1), (a_2, b_2) \notin F_v^{-1}$. We have $(v, b_0), (b_0, a_0), (a_0, v) \in F$. It is obvious from $a_1 b_0, a_0 b_2 \in E(\overline{G})$ that $v \neq a_1, b_2$. We have $v \neq b_1$ since otherwise $(a_1, b_1 = v, b_0)$ is a Δ -obstruction, a contradiction. We also have $v \neq a_2$ since otherwise $(a_0, v = a_2, b_2)$ is a Δ -obstruction, a contradiction. Therefore, the seven vertices $v, a_0, b_0, a_1, b_1, a_2, b_2$ are distinct.

We have from Claim 11 that there is a vertex z such that $(a_0, z), (z, b_0) \in F$ and either $(v, z) \in \overline{F}$ or $(z, v) \in \overline{F}$. We first suppose $(z, v) \in \overline{F}$. It is obvious from $a_1 b_0, a_0 b_2 \in E(\overline{G})$ that $z \neq a_1, b_2$. We have $z \neq b_1$ since otherwise $(a_1, b_1 = z, b_0)$ is a Δ -obstruction, a contradiction. We also have $z \neq a_2$ since otherwise $(a_0, z = a_2, b_2)$ is a Δ -obstruction, a contradiction. Therefore, the eight vertices $v, z, a_0, b_0, a_1, b_1, a_2, b_2$ are distinct.

If $(b_1, v) \in \overline{F}$, then (v, b_0, a_1, b_1) is an alternating 4-cycle, a contradiction. If $(z, a_2) \in \overline{F}$, then (a_2, b_2, a_0, z) is an alternating 4-cycle, a contradiction. If $(v, b_1) \in \overline{F}$, then $(z, v), (b_1, a_2) \in \overline{F}$ implies $(z, a_2) \in \overline{F}$, a contradiction. If $(a_2, z) \in \overline{F}$, then $(b_1, a_2), (z, v) \in \overline{F}$ implies $(b_1, v) \in \overline{F}$, a contradiction. Thus $vb_1, za_2 \in E(G)$. Now, we have from Claim 7 that (v, b_1, z, a_2) is a chordless cycle of length 4.

If $(b_1, v), (b_1, z), (a_2, v), (a_2, z) \in F$, then $(v, b_0, a_1, b_1, a_2, z)$ is an alternating 6-cycle in $F \cup \overline{F}$, and hence we have from Claim 8 that $(a_1, b_1) \in F_v$, a contradiction. If $(v, b_1), (v, a_2), (z, b_1), (z, a_2) \in F$, then $(v, b_1, a_2, b_2, a_0, z)$ is an alternating 6-cycle in $F \cup \overline{F}$, and hence we have from Claim 8 that $(a_2, b_2) \in F_v$, a contradiction.

When we suppose $(v, z) \in \overline{F}$, we also have either $(a_1, b_1) \in F_v$ or $(a_2, b_2) \in F_v$, a contradiction. Thus we have the claim. \square

The following claim states that the directions of the edges still alternate in F' on every chordless cycle of length 4.

Claim 15. *If an edge on a chordless cycle of length 4 is in F_v , then all the edges on the cycle is in F_v .*

Proof. Suppose that there is a chordless cycle (a, b, c, d) of length 4 in G such that $(a, b) \in F_v$, that is, $(v, a), (a, b), (b, v), (a, d), (c, b), (c, d) \in F$ and $ac, bd \in E(\overline{G})$. We have from Claim 11 that there is a vertex z such that $(b, z), (z, a) \in F$ and either $(v, z) \in \overline{F}$ or $(z, v) \in \overline{F}$. In either case, we have from Claim 9 that $(v, c), (d, v) \in F$. Thus $(a, d), (c, b), (c, d) \in F_v$. \square

The following claim states that the reversing the orientation of the edges in F_v generates no Δ -obstructions, and thus $F' \cup \overline{F}$ still contains no Δ -obstructions.

Claim 16. *No edges in F_v^{-1} is an edge of any Δ -obstruction of $F' \cup \overline{F}$.*

Proof. We prove the claim by contradiction. Suppose that $F' \cup \overline{F}$ contains a Δ -obstruction (a, b, c) with $(a, b), (b, c) \in F'$ and $(c, a) \in \overline{F}$. If $(a, b), (b, c) \in F_v^{-1}$, then $(v, b), (b, a), (a, v) \in F$ and $(v, c), (c, b), (b, v) \in F$, a contradiction.

Suppose $(a, b) \in F_v^{-1}$ and $(b, c) \notin F_v^{-1}$. We have $(v, b), (b, a), (a, v), (b, c) \in F$ and $(c, a) \in \overline{F}$. We have from Claim 11 that there is a vertex z such that $(a, z), (z, b) \in F$ and either $(v, z) \in \overline{F}$ or $(z, v) \in \overline{F}$. We first suppose

$(v, z) \in \overline{F}$. If $(z, c) \in \overline{F}$, then $(c, a) \in \overline{F}$ implies $(z, a) \in \overline{F}$, contradicting $(a, z) \in F$. If $(c, z) \in \overline{F}$, then (z, b, c) is a Δ -obstruction, a contradiction. Thus $zc \in E(G)$. We now have from Claim 7 that (z, c, v, a) is a chordless cycle of length 4. Since $(a, v), (a, z) \in F$, we have $(c, v), (c, z) \in F$, but then $(v, b), (b, c), (c, v) \in F$ implies $(b, c) \in F_v$, a contradiction. We next suppose $(z, v) \in \overline{F}$. If $(v, c) \in \overline{F}$, then $(c, a) \in \overline{F}$ implies $(v, a) \in \overline{F}$, contradicting $(a, v) \in F$. If $(c, v) \in \overline{F}$, then (v, b, c) is a Δ -obstruction, a contradiction. Thus $vc \in E(G)$. We now have from Claim 7 that (v, c, z, a) is a chordless cycle of length 4. Since $(a, v), (a, z) \in F$, we have $(c, v), (c, z) \in F$, but then $(v, b), (b, c), (c, v) \in F$ implies $(b, c) \in F_v$, a contradiction.

Suppose $(b, c) \in F_v^{-1}$ and $(a, b) \notin F_v^{-1}$. We have $(v, c), (c, b), (b, v), (a, b) \in F$ and $(c, a) \in \overline{F}$. We have from Claim 11 that there is a vertex z such that $(b, z), (z, c) \in F$ and either $(v, z) \in \overline{F}$ or $(z, v) \in \overline{F}$. We first suppose $(v, z) \in \overline{F}$. If $(a, v) \in \overline{F}$, then $(c, a) \in \overline{F}$ implies $(c, v) \in \overline{F}$, contradicting $(v, c) \in F$. If $(v, a) \in \overline{F}$, then (a, b, v) is a Δ -obstruction, a contradiction. Thus $va \in E(G)$. We now have from Claim 7 that (v, c, z, a) is a chordless cycle of length 4. Since $(v, c), (z, c) \in F$, we have $(v, a), (z, a) \in F$, but then $(v, a), (a, b), (b, v) \in F$ implies $(a, b) \in F_v$, a contradiction. We next suppose $(z, v) \in \overline{F}$. If $(a, z) \in \overline{F}$, then $(c, a) \in \overline{F}$ implies $(c, z) \in \overline{F}$, contradicting $(z, c) \in F$. If $(z, a) \in \overline{F}$, then (a, b, z) is a Δ -obstruction, a contradiction. Thus $za \in E(G)$. We now have from Claim 7 that (z, c, v, a) is a chordless cycle of length 4. Since $(v, c), (z, c) \in F$, we have $(v, a), (z, a) \in F$, but then $(v, a), (a, b), (b, v) \in F$ implies $(a, b) \in F_v$, a contradiction. \square

4. CONCLUDING REMARKS

In this paper, we provided a new algorithm for the recognition of simple-triangle graphs to improve the time bound from $O(n^2\overline{m})$ to $O(nm)$. The algorithm uses the vertex ordering characterization in our previous paper [32] that a graph is a simple-triangle graph if and only if there is a linear ordering of the vertices containing both an alternating orientation of the graph and a transitive orientation of the complement of the graph. The algorithm finds such a vertex ordering or report that the given graph is not a simple-triangle graph. Correctness of the algorithm is due to the following structural characterization of simple-triangle graphs (Theorem 4): a graph G is a simple-triangle graph if and only if there is a pair of an alternating orientation F of G and a transitive orientation \overline{F} of the complement \overline{G} of G such that $F \cup \overline{F}$ contains no Δ -obstructions. We also showed, as a byproduct, that the recognition problem is NP-complete for acyclic alternately orientable graphs (Theorem 2), and alternately orientable cocomparability graphs can be recognized in $O(nm)$ time (Theorem 6).

Finally, we list some open problems related to the results of this paper:

- Is there a recognition algorithm for simple-triangle graphs with time complexity less than $O(nm)$?
- Is there a polynomial time algorithm for the isomorphism problem for simple-triangle graphs [30, 32, 33]?
- Is there a forbidden characterization for simple-triangle graphs?

REFERENCES

- [1] B. Aspvall, M. F. Plass, and R. E. Tarjan. A linear-time algorithm for testing the truth of certain quantified boolean formulas. *Inf. Process. Lett.*, 8(3):121–123, 1979.
- [2] K. P. Bogart, P. C. Fishburn, G. Isaak, and L. Langley. Proper and unit tolerance graphs. *Discrete Appl. Math.*, 60(1–3):99–117, 1995.
- [3] A. Brandstädt, V. B. Le, and J. P. Spinrad. *Graph Classes: A Survey*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1999.
- [4] M. R. Cerioli, F. de S. Oliveira, and J. L. Szwarcfiter. Linear-interval dimension and PI orders. *Electron. Notes in Discrete Math.*, 30:111–116, 2008.
- [5] F. Cheah and D. G. Corneil. On the structure of trapezoid graphs. *Discrete Appl. Math.*, 66(2):109–133, 1996.
- [6] O. Cogis. On the Ferrers dimension of a digraph. *Discrete Math.*, 38(1):47–52, 1982.
- [7] D. G. Corneil and P. A. Kamula. Extensions of Permutation and Interval Graphs. *Congr. Numer.*, 58:267–275, 1987.
- [8] I. Dagan, M. C. Golumbic, and R. Y. Pinter. Trapezoid graphs and their coloring. *Discrete Appl. Math.*, 21(1):35–46, 1988.
- [9] S. Felsner. Tolerance graphs, and orders. *J. Graph Theory*, 28(3):129–140, 1998.
- [10] S. Felsner, M. Habib, and R. H. Möhring. On the Interplay Between Interval Dimension and Dimension. *SIAM J. Discrete Math.*, 7(1):32–40, 1994.
- [11] T. Gallai. Transitiv orientierbare graphen. *Acta Math. Acad. Sci. Hungar.*, 18(1–2):25–66, 1967.

- [12] A. Ghouila-Houri. Caractérisation des graphes non orientés dont on peut orienter les arêtes de manière à obtenir le graphe d'une relation d'ordre. *C. R. Acad. Sci. Paris*, 254:1370–1371, 1962.
- [13] M. C. Golumbic. *Algorithmic Graph Theory and Perfect Graphs*, volume 57 of *Ann. Discrete Math.* Elsevier B.V., Amsterdam, The Netherlands, 2 edition, 2004.
- [14] M. C. Golumbic, C. L. Monma, and W. T. Trotter, Jr. Tolerance graphs. *Discrete Appl. Math.*, 9(2):157–170, 1984.
- [15] M. C. Golumbic and A. N. Trenk. *Tolerance Graphs*. Cambridge University Press, Cambridge, UK, 2004.
- [16] W. Guttmann and M. Maucher. Variations on an Ordering Theme with Constraints. In G. Navarro, L. Bertossi, and Y. Kohayakawa, editors, *Fourth IFIP International Conference on Theoretical Computer Science (TCS 2006)*, volume 209 of *IFIP*, pages 77–90. Springer, 2006.
- [17] P. Hell and J. Huang. Lexicographic orientation and representation algorithms for comparability graphs, proper circular arc graphs, and proper interval graphs. *J. Graph Theory*, 20(3):361–374, 1995.
- [18] C. T. Hoàng. Alternating orientation and alternating colouration of perfect graphs. *J. Combin. Theory Ser. B*, 42(3):264–273, 1987.
- [19] S. Hougardy. Counterexamples to three conjectures concerning perfect graphs. *Discrete Math.*, 117(1–3):245–251, 1993.
- [20] L. J. Langley. A recognition algorithm for orders of interval dimension two. *Discrete Appl. Math.*, 60(1–3):257–266, 1995.
- [21] T.-H. Ma and J. P. Spinrad. On the 2-Chain Subgraph Cover and Related Problems. *J. Algorithms*, 17(2):251–268, 1994.
- [22] R. M. McConnell and J. P. Spinrad. Modular decomposition and transitive orientation. *Discrete Math.*, 201(1–3):189–241, 1999.
- [23] T. A. McKee and F. R. McMorris. *Topics in Intersection Graph Theory*. Society for Industrial and Applied Mathematics, Philadelphia, PA, USA, 1999.
- [24] G. B. Mertzios. The recognition of triangle graphs. *Theor. Comput. Sci.*, 438:34–47, 2012.
- [25] G. B. Mertzios. The Recognition of Simple-Triangle Graphs and of Linear-Interval Orders is Polynomial. *SIAM J. Discrete Math.*, 29(3):1150–1185, 2015.
- [26] G. B. Mertzios and D. G. Corneil. Vertex splitting and the recognition of trapezoid graphs. *Discrete Appl. Math.*, 159(11):1131–1147, 2011.
- [27] G. B. Mertzios, I. Sau, and S. Zaks. The Recognition of Tolerance and Bounded Tolerance Graphs. *SIAM J. Comput.*, 40(5):1234–1257, 2011.
- [28] A. Pnueli, A. Lempel, and S. Even. Transitive orientation of graphs and identification of permutation graphs. *Canad. J. Math.*, 23(1):160–175, 1971.
- [29] J. P. Spinrad. *Efficient Graph Representations*. American Mathematical Society, Providence, RI, USA, 2003.
- [30] A. Takaoka. Graph Isomorphism Completeness for Trapezoid Graphs. *IEICE Trans. Fundamentals*, E98–A(8):1838–1840, 2015.
- [31] A. Takaoka. Recognizing Simple-Triangle Graphs by Restricted 2-Chain Subgraph Cover. In S.-H. Poon, M. Rahman, and H.-C. Yen, editors, *Proceedings of the 11th International Conference and Workshops on Algorithms and Computation (WALCOM 2017)*, volume 10167 of *Lecture Notes in Comput. Sci.*, pages 177–189. Springer, Cham, 2017.
- [32] A. Takaoka. A vertex ordering characterization of simple-triangle graphs. *Discrete Math.*, 341(12):3281–3287, 2018.
- [33] R. Uehara. The graph isomorphism problem on geometric graphs. *Discrete Math. Theor. Comput. Sci.*, 16(2):87–96, 2014.

DEPARTMENT OF INFORMATION SYSTEMS CREATION, KANAGAWA UNIVERSITY, ROKKAKUBASHI 3-27-1 KANAGAWA-KU, KANAGAWA, 221–8686, JAPAN

E-mail address: takaoka@kanagawa-u.ac.jp