

MUKAI'S PROGRAM (RECONSTRUCTING A K3 SURFACE FROM A CURVE) VIA WALL-CROSSING

SOHEYLA FEYZBAKHS

ABSTRACT. Let C be a curve of genus $g \geq 11$ such that $g - 1$ is a composite number. Suppose C is on a K3 surface whose Picard group is generated by the curve class $[C]$. We use wall-crossing with respect to Bridgeland stability conditions to generalise Mukai's program to this situation: we show how to reconstruct the K3 surface containing the curve C as a Fourier-Mukai transform of a Brill-Noether locus of vector bundles on C .

1. INTRODUCTION

In this paper, we consider the problem of reconstructing a K3 surface from a curve on that surface. The main result is the following which extends a program proposed by Mukai in [Muk01, Section 10].

Theorem 1.1. *Let (X, H) be a polarised K3 surface with $\text{Pic}(X) = \mathbb{Z}H$. Let C be any curve in the linear system $|H|$ of genus $g \geq 11$ such that $g - 1$ is a composite number¹. Then X is the unique K3 surface of Picard rank one and genus g containing C , and can be reconstructed as a Fourier-Mukai partner of a certain Brill-Noether locus of vector bundles on C .*

Any K3 surface of Picard rank one has a canonical primitive polarisation and therefore a well-defined genus. Note that the curve C in Theorem 1.1 is not necessarily smooth and can be singular. The missing cases where $g - 1$ is a prime number are considered in [Fey20].

Write $g = rs + 1$ for two integers $r \geq 2$ and $s \geq \max\{r, 5\}$. We consider the Brill-Noether locus $\mathcal{BN} := M_C(r, 2rs, r + s)$ of slope semistable rank r -vector bundles on the curve C having degree $2rs$ and possessing at least $r + s$ linearly independent global sections. Let $M_{X,H}(v)$ be the moduli space of H -Gieseker semistable sheaves with Mukai vector $v = (r, H, s)$ on X . We have chosen the Mukai vector v such that it is a primitive class with $v^2 = 0$, hence $M_{X,H}(v)$ is a K3 surface as well. Moreover, any H -Gieseker semistable sheaf $E \in M_{X,H}(v)$ is a slope stable locally free sheaf. The choice of the Brill-Noether locus \mathcal{BN} is justified by the following Theorem.

¹In the published version of this paper [Fey19], we also considered the case (B) where $g - 1$ is a prime number. But there is a mistake in the proof of [Fey19, Proposition 5.2 (a)] which affects the validity of [Fey19, Theorem 1.2] in case (B), see [Fey20, Remark 4.1] for details. That is why this case has been investigated in another paper [Fey20] using a new treatment so that eventually the main result (Theorem 1.1) in [Fey19] is proved valid.

Theorem 1.2. *Let (X, H) be a polarised K3 surface with $\text{Pic}(X) = \mathbb{Z}.H$ and let C be any curve in the linear system $|H|$. We have an isomorphism*

$$(1) \quad \psi: M_{X,H}(v) \rightarrow \mathcal{BN}$$

with \mathcal{BN} as defined above, which sends a bundle E on X to its restriction $E|_C$.

In other words, special vector bundles on the curve C , which have an unexpected number of global sections, are the restriction of vector bundles on the surface X . This is analogous to the case of line bundles, where a well-known theorem by Green and Lazarsfeld [GL87] says that the Clifford index of a non-Clifford general curve on a K3 surface can be computed by the restriction of a line bundle on the surface.

There exists a Brauer class $\alpha \in \text{Br}(\mathcal{BN})$ and a universal $(1 \times \alpha)$ -twisted sheaf \mathcal{E} on $C \times \mathcal{BN}$. Define $v' \in H^*(\mathcal{BN}, \mathbb{Z})$ to be the Mukai vector of $\mathcal{E}|_{p \times \mathcal{BN}}$ for a point p on the curve C (see [HS05] for definition in case $\alpha \neq 1$).

Theorem 1.3. *Let (X, H) be a polarised K3 surface with $\text{Pic}(X) = \mathbb{Z}.H$ of genus $g \geq 11$ such that $g-1$ is a composite number, and let C be any curve in the linear system $|H|$. Then any K3 surface of Picard rank one and genus g which contains the curve C is isomorphic to the moduli space $M_{\mathcal{BN}, H'}^\alpha(v')$ of α -twisted sheaves on \mathcal{BN} of Mukai vector v' which are semistable with respect to a generic polarisation H' on \mathcal{BN} .*

The embedding of the curve C into the K3 surface $M_{\mathcal{BN}, H'}^\alpha(v')$ is given by $p \mapsto \mathcal{E}|_{p \times \mathcal{BN}}$. Combining Theorems 1.2 and 1.3 gives Theorem 1.1.

1.1. Previous work. Let \mathcal{F}_g be the moduli space of polarised K3 surfaces (X, H) where H is a primitive ample line bundle on X and $H^2 = 2g - 2$. This space is a quasi-projective variety of dimension 19. Let \mathcal{P}_g be the moduli space of triples (X, H, C) such that $(X, H) \in \mathcal{F}_g$ and C is a smooth curve in the linear system $|H|$. Therefore, its dimension is $19 + g$. Finally, let \mathcal{M}_g be the moduli space of smooth curves of genus g . Its dimension is $3g - 3$. The space \mathcal{P}_g has natural projections to \mathcal{F}_g and \mathcal{M}_g which we denote by ϕ_g and m_g , respectively;

$$\begin{array}{ccc} & \mathcal{P}_g & \\ m_g \swarrow & & \searrow \phi_g \\ \mathcal{M}_g & & \mathcal{F}_g \end{array}$$

The map m_g is dominant for $g \leq 11$ and $g \neq 10$ [Muk88]. In [CLM93, Theorem 5], Ciliberto, Lopez and Miranda proved that for $g \geq 11$ and $g \neq 12$, the map m_g is birational onto its image. For the exceptional cases $g = 10$ or $g = 12$, the map m_g is neither dominant nor generically finite [Muk01].

In [Muk01], Mukai introduced a geometric program to find the rational inverse of m_g where $g = 2s + 1$ and $s \geq 5$ odd. His idea to reconstruct the K3 surface is as follows. Let C be a general curve in the image of m_g . Consider the Brill-Noether locus $M_C(2, K_C, s + 2)^{\text{st}}$ of stable rank 2-vector bundles on the curve C with canonical determinant and possessing at least $s + 2$ linearly independent global sections. Then $M_C(2, K_C, s + 2)^{\text{st}}$ is a K3 surface

and the K3 surface containing the curve C can be obtained uniquely as a Fourier-Mukai transform of the Brill-Noether locus.

This program was completely proved by him in [Muk96] for $g = 11$. The key idea is that all vector bundles in the Brill-Noether locus $M_C(2, K_C, 7)$ are the restriction of vector bundles on the surface. He first considers a point $(X', C') \in \mathcal{P}_g$ of a special type and shows that the Brill-Noether locus $M_{C'}(2, K_{C'}, 7)$ is isomorphic to X' . Indeed, he proves that both surfaces are isomorphic to the moduli space $M_{X', H'}(v)$ where $v = (2, H, 5)$. Given a general pair $(X, C) \in \mathcal{P}_g$, the Brill-Noether locus $M_C(2, K_C, 7)$ is a flat deformation of $M_{C'}(2, K_{C'}, 7)$ and has expected dimension. Thus, it is again a K3 surface and the original K3 surface can be obtained as an appropriate Fourier-Mukai transform of it.

Arbarello, Bruno and Sernesi [ABS14] generalised this strategy to higher genera. They proved that for a general pair $(X, C) \in \mathcal{P}_g$ where $g = 2s + 1 \geq 11$, there is a unique irreducible component V_C of $M_C(2, K_C, s + 2)$ such that $(V_C)_{\text{red}}$ is a K3 surface isomorphic to the moduli space $M_{X, H}(v)$ where $v = (2, H, s)$. Then they showed that the original K3 surface can be reconstructed using this component whenever $g \equiv 3 \pmod{4}$.

In this paper, without any deformation argument, we show that for a general pair $(X, C) \in \mathcal{P}_g$, when $g = rs + 1 \geq 11$, the Brill-Noether locus $M_C(r, K_C, s + r)$ is isomorphic to the moduli space $M_{X, H}(r, H, s)$. As a result, we prove the uniqueness of the K3 surface of Picard rank one which contains the curve C of genus $g \geq 11$ when $g - 1$ is a composite number.

1.2. The strategy of the proof. We prove Theorem 1.2 by wall-crossing for the push-forward of semistable vector bundles on the curve C , with respect to Bridgeland stability conditions on the bounded derived category $\mathcal{D}(X)$ of X . There exists a region in the space of stability conditions where the Brill-Noether behaviour of *stable* objects is completely controlled by the nearby *Brill-Noether wall*. This wall destabilises objects with non-zero global sections, and arguments similar to [Bay18] show that the Brill-Noether loci are mostly of expected dimension. Our first key result, Proposition 3.4, gives an extension to *unstable* objects: it gives a bound on the number of global sections in terms of their *mass*, i.e. the length of their Harder-Narasimhan polygon.

Consequently, we only need a polygon that circumscribes this Harder-Narasimhan polygon on the left, to bound the number of global sections. For any coherent sheaf, there exists a chamber which is called the Gieseker chamber, where the notion of Bridgeland stability coincides with the old notion of Gieseker stability. Unlike the case of push-forward of line bundles considered in [Bay18], the Brill-Noether wall is not adjacent to the Gieseker chamber for the push-forward of semistable vector bundles F of higher ranks on the curve C . However, the wall that bounds the Gieseker chamber provides an extremal polygon which contains the Harder-Narasimhan polygon, see e.g. Lemma 4.3. Combined with Proposition 3.4, this gives a bound on the number of global sections of vector bundles on the curve C ; the proof also shows that the bound is achieved if and only if the vector bundle F is the restriction of a vector bundle on the surface.

Plan of the paper. Section 2 reviews the definition of geometric stability conditions on K3 surfaces and describes a two-dimension family of stability conditions. Section 3 deals

with the Brill-Noether wall; we provide an upper bound for the number of global sections via the geometry of Harder-Narasimhan polygon. Section 4 concerns the proof of bijectivity of the morphism ψ in (1). The proof of the main result is contained in Section 5.

Acknowledgement. I would like to thank Arend Bayer for many useful discussions. I am grateful for comments by Benjamin Bakker, Gavril Farkas, Chunyi Li, Hsueh-Yung Lin, Richard Thomas, Yukinobu Toda and Bach Tran. I would also like to thank the referees for their careful reading of the paper, and for many useful suggestions. The author was supported by the ERC starting grant WallXBirGeom 337039.

2. BRIDGELAND STABILITY CONDITIONS ON K3 SURFACES

In this section, we give a brief review of a two-dimensional family of Bridgeland stability conditions on the bounded derived category of coherent sheaves on a K3 surface. The main references are [Bri07, Bri08].

2.1. Bridgeland stability conditions. Let (X, H) be a smooth polarised K3 surface with $\text{Pic}(X) = \mathbb{Z} \cdot H$. We denote by $\mathcal{D}(X)$ the bounded derived category of coherent sheaves on the surface X . The Mukai vector of an object $E \in \mathcal{D}(X)$ is an element of the lattice $\mathcal{N}(X) = \mathbb{Z} \oplus \text{NS}(X) \oplus \mathbb{Z} \cong \mathbb{Z}^3$ defined via

$$v(E) = (\text{rk}(E), c(E)H, s(E)) = \text{ch}(E)\sqrt{\text{td}(X)} \in H^*(X, \mathbb{Z}),$$

where $\text{ch}(E)$ is the Chern character of E . The Mukai bilinear form

$$\langle v(E), v(E') \rangle = c(E)c(E')H^2 - \text{rk}(E)s(E') - \text{rk}(E')s(E)$$

makes $\mathcal{N}(X)$ into a lattice of signature $(2, 1)$. The Riemann-Roch theorem implies that this form is the negative of the Euler form, defined as

$$\chi(E, E') = \sum_i (-1)^i \dim_{\mathbb{C}} \text{Hom}_X^i(E, E') = -\langle v(E), v(E') \rangle.$$

Note that the Euler form $\chi(-, -)$ defines a bilinear form on the Grothendieck group $K(X)$ which descends to a non-degenerate form on the lattice

$$\mathcal{N}(X) = K(X)/K(X)^{\perp},$$

where $K(X)^{\perp}$ is the left-radical. Recall that for a coherent sheaf E with positive rank $\text{rk}(E) > 0$, the slope is defined as

$$\mu_H(E) := \frac{c(E)}{\text{rk}(E)},$$

and if $\text{rk}(E) = 0$, define $\mu_H(E) := +\infty$.

Definition 2.1. We say that an object $E \in \text{Coh}(X)$ is μ_H -(semi)stable if for all proper non-trivial subsheaves $F \subset E$, we have $\mu_H(F) < (\leq) \mu_H(E)$.

A stability function on an abelian category \mathcal{A} is a group homomorphism $Z: K(\mathcal{A}) \rightarrow \mathbb{C}$ such that for any non zero object $E \in \mathcal{A}$,

$$Z(E) \in \mathbb{R}^{>0} \exp(i\pi\phi(E)) \text{ with } 0 < \phi(E) \leq 1.$$

By [Bri08, Proposition 3.5], to give a stability condition on a triangulated category \mathcal{D} is equivalent to giving a bounded t-structure on \mathcal{D} and a stability function on its heart which has the Harder-Narasimhan property. Given a real number $b \in \mathbb{R}$, denote by $\mathcal{T}^b \subset \text{Coh}(X)$ the subcategory of sheaves E whose quotients $E \twoheadrightarrow F$ satisfy $\mu_H(F) > b$ and by $\mathcal{F}^b \subset \text{Coh}(X)$ the subcategory of sheaves E' whose subsheaves $F' \hookrightarrow E'$ satisfy $\mu_H(F') \leq b$. Tilting with respect to the torsion pair $(\mathcal{T}^b, \mathcal{F}^b)$ on $\text{Coh}(X)$ gives a bounded t-structure on $\mathcal{D}(X)$ with heart

$$\mathcal{A}(b) := \{E \in \mathcal{D}(X) : E \cong [E^{-1} \xrightarrow{d} E^0], \ker d \in \mathcal{F}^b \text{ and } \text{cok } d \in \mathcal{T}^b\} \subset \mathcal{D}(X).$$

All the stability functions that we will consider in this paper factor through the surjection $K(X) \twoheadrightarrow \mathcal{N}(X)$. For a pair $(b, w) \in \mathbb{H} = \mathbb{R} \times \mathbb{R}^{>0}$, the stability function $Z_{(b,w)}: \mathcal{N}(X) \rightarrow \mathbb{C}$ is defined as

$$Z_{(b,w)}(r, cH, s) = \left\langle (r, cH, s), \left(1, bH, \frac{H^2}{2}(b^2 - w^2)\right) \right\rangle + i \left\langle (r, cH, s), \left(0, \frac{H}{H^2}, b\right) \right\rangle.$$

We denote the root system by $\Delta(X) := \{\delta \in \mathcal{N}(X) : \langle \delta, \delta \rangle = -2\}$.

Theorem 2.2 ([Bri08]). *Suppose (X, H) is a polarised K3 surface with $\text{Pic}(X) = \mathbb{Z}H$. Then the pair $\sigma_{(b,w)} = (\mathcal{A}(b), Z_{(b,w)})$ defines a Bridgeland stability condition on $\mathcal{D}(X)$ if for all $\delta \in \Delta(X)$ with $\text{rk}(\delta) > 0$ and $\text{Im}[Z_{(b,w)}(\delta)] = 0$ we have $\text{Re}[Z_{(b,w)}(\delta)] > 0$. The family of stability conditions $\sigma_{(b,w)}$ varies continuously as the pair (b, w) varies in \mathbb{H} .*

Note that the stability condition $\sigma_{(b,w)}$, up to the action of $\tilde{\text{GL}}^+(2, \mathbb{R})$, is the same as the stability condition defined in [Bri08, Section 6]. We expand upon the statements in Theorem 2.2 by explaining the notion of $\sigma_{(b,w)}$ -stability and the associated Harder-Narasimhan filtration. For a stability condition $\sigma_{(b,w)}$ and $E \in \mathcal{A}(b)$, we have $Z_{(b,w)}(v(E)) \in \mathbb{R}^{>0} \exp(i\pi\phi_{(b,w)}(v(E)))$ where

$$\phi_{(b,w)}(v(E)) = \frac{1}{\pi} \tan^{-1} \left(-\frac{\text{Re}[Z_{(b,w)}(v(E))]}{\text{Im}[Z_{(b,w)}(v(E))]} \right) + \frac{1}{2} \in (0, 1].$$

We will abuse notations and write $Z(E)$ and $\phi(E)$ instead of $Z(v(E))$ and $\phi(v(E))$.

Definition 2.3. We say that an object $E \in \mathcal{D}(X)$ is $\sigma_{(b,w)}$ -(semi)stable if some shift $E[k]$ is contained in the abelian category $\mathcal{A}(b)$ and for any non-trivial subobject $E' \subset E[k]$ in $\mathcal{A}(b)$, we have $\phi_{(b,w)}(E') < (\leq) \phi_{(b,w)}(E[k])$.

Any object $E \in \mathcal{A}(b)$ admits a Harder-Narasimhan (HN) filtration: a sequence

$$(2) \quad 0 = \tilde{E}_0 \subset \tilde{E}_1 \subset \tilde{E}_2 \subset \dots \subset \tilde{E}_n = E$$

of objects in $\mathcal{A}(b)$ where the factors $E_i := \tilde{E}_i / \tilde{E}_{i-1}$ are $\sigma_{(b,w)}$ -semistable and

$$\phi_{(b,w)}^+(E) := \phi_{(b,w)}(E_1) > \phi_{(b,w)}(E_2) > \dots > \phi_{(b,w)}(E_n) =: \phi_{(b,w)}^-(E).$$

In addition, any $\sigma_{(b,w)}$ -semistable object $E \in \mathcal{A}(b)$ has a Jordan-Hölder (JH) filtration into stable factors of the same phase, see [Bri08, Section 2] for more details.

Suppose $E_1 \hookrightarrow E_2 \twoheadrightarrow E_3$ is a short exact sequence in $\mathcal{A}(b)$. Since $H^i(E_j) = 0$ for $j = 1, 2, 3$ and $i \neq 0, -1$, taking cohomology gives a long exact sequence of coherent sheaves

$$0 \rightarrow H^{-1}(E_1) \rightarrow H^{-1}(E_2) \rightarrow H^{-1}(E_3) \rightarrow H^0(E_1) \rightarrow H^0(E_2) \rightarrow H^0(E_3) \rightarrow 0.$$

For any pair of objects E and E' of $\mathcal{D}(X)$, Serre duality gives isomorphisms

$$\mathrm{Hom}_X^i(E, E') \cong \mathrm{Hom}_X^{2-i}(E', E)^*.$$

If the objects E and E' lie in the heart $\mathcal{A}(b)$, then $\mathrm{Hom}_X^i(E, E') = 0$ if $i < 0$ or $i > 2$. Suppose the object $E \in \mathcal{A}(b)$ is $\sigma_{(b,w)}$ -stable, then E does not have any non-trivial subobject with the same phase, thus $\mathrm{Hom}_X(E, E) = \mathrm{Hom}_X^2(E, E)^* = \mathbb{C}$. This implies

$$(3) \quad v(E)^2 + 2 = \mathrm{Hom}_X^1(E, E) \geq 0.$$

To simplify drawing the figures, we always consider the following projection:

$$pr: \mathcal{N}(X) \setminus \{s = 0\} \rightarrow \mathbb{R}^2, \quad pr(r, cH, s) = \left(\frac{c}{s}, \frac{r}{s} \right).$$

Take a pair $(b, w) \in \mathbb{H}$, the kernel of $Z_{(b,w)}$ is a line inside the negative cone in $\mathcal{N}(X) \otimes \mathbb{R} \cong \mathbb{R}^3$ spanned by the vector $(2, 2bH, H^2(b^2 + w^2))$. Its projection is denoted by

$$k(b, w) := pr(\ker Z_{(b,w)}) = \left(\frac{2b}{H^2(b^2 + w^2)}, \frac{2}{H^2(b^2 + w^2)} \right).$$

Thus, for any stability condition $\sigma_{(b,w)}$, we associate a point $k(b, w) \in \mathbb{R}^2$. The two dimensional family of stability conditions of form $\sigma_{(b,w)}$, is parametrised by the space

$$V(X) := \{k(b, w): \text{the pair } (\mathcal{A}(b), Z_{(b,w)}) \text{ is a stability condition on } \mathcal{D}(X)\} \subset \mathbb{R}^2$$

with the standard topology on \mathbb{R}^2 .




FIGURE 1. The grey area is the 2-dimensional subspace of stability conditions $V(X)$.

Lemma 2.4. *We have*

$$V(X) = \left\{ (x, y) \in \mathbb{R}^2 : y > \frac{H^2 x^2}{2} \right\} \setminus \bigcup_{\delta \in \Delta(X)} I_\delta$$

where I_δ is the closed line segment that connects $p_\delta =: \text{pr}(\delta)$ to q_δ which is the intersection point of the parabola $y = \frac{H^2}{2}x^2$ with the line through the origin and p_δ , see Figure 1.

Proof. By definition, the point $k(b, w)$ is above the parabola and for every point (x, y) above the parabola, there exists a unique pair $(b, w) \in \mathbb{H}$ such that $k(b, w) = (x, y)$. For any root $\delta \in \Delta(X)$, $\text{Im}[Z_{(b,w)}(\delta)] = 0$ if and only if the point $k(b, w)$ is on the line passing through the origin and the point $\text{pr}(\delta) = p_\delta$. If $\text{rk}(\delta) > 0$, the line segment I_δ is precisely the segment of this line on which $\text{Re}[Z_{(b,w)}(\delta)] > 0$, thus the claim follows from Theorem 2.2. \square

Remark 2.5. The point $k(b, w)$ is on the line $x = by$. As w gets larger, the point $k(b, w)$ gets closer to the origin. Take two non-parallel vectors $u, v \in \mathcal{N}(X) \otimes \mathbb{R} \cong \mathbb{R}^3$, then $Z_{(b,w)}(v)$ and $Z_{(b,w)}(u)$ are aligned if and only if the kernel of $Z_{(b,w)}$ in $\mathcal{N}(X) \otimes \mathbb{R}$ lies on the plane spanned by v and u , i.e. the points corresponding to $\mathbb{R} \cdot u$, $\mathbb{R} \cdot v$ and $\ker Z_{(b,w)}$ in the projective space $\mathbb{P}_{\mathbb{R}}^2$ are collinear. This, in particular, implies that if three objects E_1, E_2 and E_3 in $\mathcal{D}(X)$ have the same phase with respect to a stability condition $\sigma_{(b,w)}$, there must be a linear dependence relation among the vectors $v(E_1), v(E_2)$ and $v(E_3)$ in $\mathcal{N}(X) \otimes \mathbb{R}$.

The 2-dimensional family of stability conditions parametrised by the space $V(X)$ admits a chamber decomposition for any object $E \in \mathcal{D}(X)$.

Proposition 2.6. *Given an object $E \in \mathcal{D}(X)$, there exists a locally finite set of walls (line segments) in $V(X)$ with the following properties:*

- (a) *The $\sigma_{(b,w)}$ -(semi)stability or instability of E is independent of the choice of the stability condition $\sigma_{(b,w)}$ in any chamber (which is a connected component of the complement of the union of walls).*
- (b) *When $\sigma_{(b_0, w_0)}$ is on a wall \mathcal{W}_E , i.e. the point $k(b_0, w_0) \in \mathcal{W}_E$, then E is strictly $\sigma_{(b_0, w_0)}$ -semistable.*
- (c) *If E is semistable in one of the adjacent chambers to a wall, then it is unstable in the other adjacent chamber.*
- (d) *Any wall \mathcal{W}_E is a connected component of $L \cap V(X)$, where L is a line that passes through the point $\text{pr}(v(E))$ if $s(E) \neq 0$, or that has a slope of $\text{rk}(E)/c(E)$ if $s(E) = 0$.*

Proof. The existence of a locally finite set of walls which satisfies properties (a), (b) and (c) is proved in [Bri08, section 9], see also [Mac14] for the description of the walls. Remark 2.5 implies that a numerical wall for the class $v(E)$ is a line L as claimed in part (d), thus [MS17, Proposition 6.22.(7)] completes the proof of (d). \square

Note that in Proposition 2.6, we do not assume $v(E)$ is primitive; in particular, E might be strictly semistable in the interior of a chamber.

Remark 2.7. Take an object $E \in \mathcal{D}(X)$, let L_1 be a connected component of $L \cap V(X)$ where L is a line as described in Proposition 2.6, part (d). Suppose E is $\sigma_{(b_0, w_0)}$ -(semi)stable for a stability condition $\sigma_{(b_0, w_0)}$ on L_1 . Then the structure of walls shows that E is (semi)stable with respect to all stability conditions on L_1 . Moreover, if E is in the heart $\mathcal{A}(b_0)$, by a straightforward computation, one can show that when we deform the stability condition $\sigma_{(b_0, w_0)}$ along the line segment L_1 , the phase of E is fixed so it remains in the heart.

The two-dimensional subspace of stability conditions. To describe the space $V(X)$, we need to find out the possible positions of the projection of roots. We denote by γ_n the point $(\frac{1}{n}, \frac{H^2}{2n^2})$ on the parabola for any $n \in \mathbb{Q}$, see Figure 2.

Lemma 2.8. For any positive number $n \in \frac{1}{2}\mathbb{N}$, define

$$U_n := \left\{ (x, y) \in \mathbb{R}^2 : 0 < |x| < \frac{1}{n} \text{ and } \frac{H^2}{2n} |x| < y \right\}.$$

If $n \leq \frac{H^2}{2}$, then there is no projection of roots $pr(\delta)$ in U_n .




FIGURE 2. No projection of roots in the grey area U_n

Proof. Assume for a contradiction that $pr(\delta = (\tilde{r}, \tilde{c}H, \tilde{s})) \in U_n$, then

$$(4) \quad 0 < \frac{H^2}{2n} \left| \frac{\tilde{c}}{\tilde{s}} \right| < \left| \frac{\tilde{r}}{\tilde{s}} \right|,$$

which implies $|\tilde{c}^2 H^2| < |2n\tilde{r}\tilde{c}|$. By assumption $\delta^2 = \tilde{c}^2 H^2 - 2\tilde{r}\tilde{s} = -2$, so

$$(5) \quad 0 < \left| \tilde{s} - \frac{1}{\tilde{r}} \right| < |n\tilde{c}|.$$

Moreover,

$$(6) \quad 0 < \left| \frac{\tilde{c}}{\tilde{s}} \right| < \frac{1}{n} \Rightarrow 0 < |n\tilde{c}| < |\tilde{s}|.$$

If $n \in \mathbb{N}$, there is no triple $(\tilde{r}, \tilde{c}, \tilde{s}) \in \mathbb{Z}^3$ that satisfies both inequalities (5) and (6) and if $n \in \frac{1}{2}\mathbb{N}$, the only possible case is $\tilde{r} = \pm 1$. But we assumed $2n \leq H^2$ and inequality (4) implies $0 < |\tilde{c}| < 1$, a contradiction. \square

Remark 2.9. Note that if the point $pr(\delta = (\tilde{r}, \tilde{c}H, \tilde{s})) = (\tilde{c}/\tilde{s}, \tilde{r}/\tilde{s})$ is on the y-axis, then $\tilde{c} = 0$. Since $\delta^2 = -2\tilde{r}\tilde{s} = -2$, we have $\tilde{r} = \tilde{s} = \pm 1$ and $pr(\delta) = (0, 1) = pr(v(\mathcal{O}_X))$. This point is denoted by o' in Figure 2.

Given three positive numbers $m, n, \epsilon \in \frac{1}{2}\mathbb{N}$ such that $m < n$, the point on the line segment $\overline{\gamma_m \gamma_n}$ with the x -coordinate $\frac{1}{m+\epsilon}$ is denoted by $q'_{m,n,\epsilon}$. Also, the point where the line segments $\overline{\gamma_m \gamma_n}$ and $\overline{o\gamma_{n-\epsilon}}$ intersect is denoted by $q_{m,n,\epsilon}$, see Figure 3. One can define similar points for the triple $(-m, -n, -\epsilon)$. For two points $q_1, q_2 \in \mathbb{R}^2$, we denote by $[\overline{q_1 q_2}]$ the closed line segment which contains both q_1 and q_2 . The open line segment which contains neither q_1 nor q_2 is denoted by $(\overline{q_1 q_2})$ and if it contains only q_1 not q_2 is denoted by $[\overline{q_1 q_2})$.

Lemma 2.10. Take $m, n, \epsilon \in \frac{1}{2}\mathbb{N}$ such that $\epsilon + \frac{1}{2} < n \leq \frac{H^2}{2}$ and

$$(7) \quad m < \frac{2\epsilon}{2\epsilon + 1}n - \epsilon.$$

Then there is no projection of roots in the grey area in Figure 3 and on the open line segments $(\overline{q_{m,n,\epsilon} q'_{m,n,\epsilon}})$ and $(\overline{q_{-m,-n,-\epsilon} q'_{-m,-n,-\epsilon}})$.




FIGURE 3. No projection of roots in the grey area

Proof. We show that the claimed region is contained in a suitable union of the U_k 's. Clearly, it is enough to prove that the open line segment $(\overline{q_{m,n,\epsilon} q'_{m,n,\epsilon}})$ is covered completely by a union of the U_k 's. Given a number $k \in \frac{1}{2}\mathbb{N}$ where $m < k < n$, the point where the line segments $\overline{\gamma_m \gamma_n}$ and $\overline{o \gamma'_k}$ intersect is denoted by γ'_k , see Figure 4.




FIGURE 4. Two consecutive points

The line segment $(\overline{q_{m,n,\epsilon} q'_{m,n,\epsilon}})$ is covered by a union of the U_k 's if the point γ'_k is inside the area $U_{k+\frac{1}{2}}$ for $m + \epsilon \leq k \leq n - \epsilon - \frac{1}{2}$. The x -coordinate of the point γ'_k is

$$x_k = \frac{1/mn}{1/m + 1/n - 1/k}.$$

One can easily show

$$f(k) := \frac{1}{mn} \left(k + \frac{1}{2} \right) + \frac{1}{k} \leq \max \{ f(n - \epsilon - 1/2), f(m + \epsilon) \} \stackrel{(*)}{<} \frac{1}{n} + \frac{1}{m},$$

where $(*)$ follows by the inequality (7). This gives $x_k < \frac{1}{k + \frac{1}{2}}$, so the point γ'_k is in $U_{k+\frac{1}{2}}$.

Therefore, the grey region in Figure 3 is contained in $\bigcup_{m+\epsilon \leq k \leq n} U_k$ where $k \in \frac{1}{2}\mathbb{N}$ and the claim follows from Lemma 2.8. \square

Wall and chamber decomposition of $V(X)$. For any coherent sheaf E on X , there is a chamber in the subspace of stability conditions $V(X)$ where the notion of Bridgeland stability coincides with the old notion of Gieseker stability which is defined using the Hilbert polynomial [Bri08, Proposition 14.2]. Recall that the Hilbert polynomial of a coherent sheaf E is defined as

$$P(E, m) := \frac{\text{rk}(E)H^2}{2}m^2 + c(E)H^2m + s(E) + \text{rk}(E).$$

The reduced Hilbert polynomial is $p(E, m) := P(E, m)/\alpha(E)$ where $\alpha(E)$ is the leading coefficient of $P(E, m)$.

Definition 2.11. A coherent sheaf E on X is called H -Gieseker (semi)stable if E is pure (has no subsheaf with lower dimensional support) and for all proper non-trivial subsheaves $F \subset E$, one has $p(F, m) < (\leq) p(E, m)$ for $m \gg 0$.

The notion of slope stability for coherent sheaves on a curve (i.e. an integral separated scheme of dimension one and of finite type over \mathbb{C}) is also defined as follows.

Definition 2.12. A torsion-free sheaf F on a curve C is slope (semi)stable if for all non-trivial subsheaves $F' \subset F$, we have

$$\frac{\chi(\mathcal{O}_C, F')}{\chi(\mathcal{O}_p, F')} < (\leq) \frac{\chi(\mathcal{O}_C, F)}{\chi(\mathcal{O}_p, F)},$$

where \mathcal{O}_p is the skyscraper sheaf at the generic point p on the curve C and \mathcal{O}_C is the structure sheaf of C .

Note that $\chi(\mathcal{O}_p, F)$ for the generic point p on the curve C is equal to the rank of the torsion-free sheaf F . If we have a closed embedding $i: C \hookrightarrow X$ of the curve C into the surface X , then the adjoint functors $Li^* \dashv Ri_*$ give $\chi(\mathcal{O}_C, F) = \chi(\mathcal{O}_X, i_*F) = \text{ch}_2(i_*F) = s(i_*F)$. Therefore, the torsion-free sheaf F on C is slope-(semi)stable if and only if i_*F is H -Gieseker (semi)stable. The following Lemma introduces the stability conditions that i_*F is stable.

Lemma 2.13 ([Mac14, Theorem 3.11]). *Let F be a vector bundle on the curve C .*

- (a) *If F is slope-(semi)stable, then there exists $w_0 > 0$ such that the push-forward i_*F is $\sigma_{(b,w)}$ -(semi)stable for any $b \in \mathbb{R}$ and $w > w_0$.*
- (b) *If i_*F is (semi)stable with respect to some stability condition $\sigma_{(b,w)}$, then F is slope-(semi)stable.*

Proof. Any coherent sheaf with rank zero is inside the heart $\mathcal{A}(b)$ for every $b \in \mathbb{R}$. Part (a) follows from [Mac14, Theorem 3.11] and the fact that $\lim_{w \rightarrow \infty} \phi_{(b,w)}(F') = 0$ for any object $F' \in \mathcal{A}(b)$ with $\text{rk}(F') > 0$. For part (b), suppose i_*F is $\sigma_{(b,w)}$ -(semi)stable. Let F' be a subsheaf of F , then i_*F' is a subobject of i_*F in the heart $\mathcal{A}(b)$, so

$$\phi_{(b,w)}(i_*F') < (\leq) \phi_{(b,w)}(i_*F) \quad \Rightarrow \quad \frac{s(i_*F')}{c(i_*F')} < (\leq) \frac{s(i_*F)}{c(i_*F)}.$$

This implies i_*F is H -Gieseker (semi)stable which gives slope (semi)stability of F . \square

For any slope semistable vector bundle F on the curve C , the chamber which contains the stability conditions $\sigma_{(b,w)}$ for $w \gg 0$, is called the *Gieseker chamber*. Note that the corresponding point $k(b, w)$ is close to the origin. We use the next lemma to describe regions in $V(X)$ with no walls for a given object.

Lemma 2.14. *Given a stability condition $\sigma_{(b,w)}$ and an object $E \in \mathcal{D}(X)$ such that*

$$0 < |\text{Im}[Z_{(b,w)}(E)]| = \min \left\{ |\text{Im}[Z_{(b,w)}(v')]| : v' \in \mathcal{N}(X) \text{ and } \text{Im}[Z_{(b,w)}(v')] \neq 0 \right\},$$

then the stability condition $\sigma_{(b,w)}$ cannot be on a wall for the object E . In particular, if $v(E) = (r, cH, s)$ and $b_0 = m/n$ for some $m, n \in \mathbb{Z}$ such that $nc - mr = \pm 1$, then the stability condition $\sigma_{(b_0,w)}$ cannot be on a wall for E .

Proof. If the stability condition $\sigma_{(b,w)}$ is on a wall \mathcal{W}_E , then the object E is strictly $\sigma_{(b,w)}$ -semistable. Up to shift, we may assume $E \in \mathcal{A}(b)$, so there are non-trivial objects $E_1, E_2 \in \mathcal{A}(b)$ of the same phase as E such that there is a short exact sequence $E_1 \hookrightarrow E \twoheadrightarrow E_2$. Since $\text{Im}[Z_{(b,w)}(E)] \neq 0$, we have $0 < \text{Im}[Z_{(b,w)}(E_i)]$ for $i = 1, 2$ and

$$\text{Im}[Z_{(b,w)}(E)] = \text{Im}[Z_{(b,w)}(E_1)] + \text{Im}[Z_{(b,w)}(E_2)].$$

This is a contradiction to our minimality assumption. If $b_0 = m/n$, then

$$\text{Im}[Z_{(b_0,w)}(E)] = c - \frac{m}{n}r = \frac{\pm 1}{n}$$

which clearly satisfies the minimality condition. \square

We finish this section by the following lemma which introduces stability conditions that a μ_H -stable vector bundle on X is stable with respect to them.

Lemma 2.15. *Let E be a μ_H -stable locally free sheaf with Mukai vector $v(E) = (r, cH, s)$. Then $E[1]$ is $\sigma_{(b_0,w)}$ -stable of phase one where $b_0 = c/r$.*

Proof. By definition, $E[1] \in \mathcal{A}(b_0)$ and has phase one which automatically implies it is $\sigma_{(b_0,w)}$ -semistable. Assume for a contradiction that $E[1]$ is strictly $\sigma_{(b_0,w)}$ -semistable. Let F_1 be the $\sigma_{(b_0,w)}$ -stable subobject of $E[1]$ and F_2 be the quotient

$$(8) \quad F_1 \hookrightarrow E[1] \twoheadrightarrow F_2.$$

Taking cohomology implies that $H^0(F_2) = 0$, so $F_2 = E'[1]$ for a torsion-free sheaf E' . Since F_1 is $\sigma_{(b_0,w)}$ -stable of phase one, [Bri08, Lemma 10.1] implies that F_1 is a skyscraper sheaf $k(x)$ or shift of a locally free sheaf. The sheaf E is locally free, so $\text{Hom}_X(k(x), E[1]) = 0$. Thus F_1 must be the shift of a locally free sheaf and we have the following exact sequence of coherent sheaves

$$0 \rightarrow H^{-1}(F_1) \rightarrow E \rightarrow E' \rightarrow 0.$$

Since $\text{Im}[Z_{(b,w)}(H^{-1}(F_1))] = \text{Im}[Z_{(b,w)}(E')] = 0$, the sheaves E' and $H^{-1}(F_1)$ have the same slope as E , which contradicts the μ_H -stability of E . \square

3. AN UPPER BOUND FOR THE NUMBER OF GLOBAL SECTIONS

In this section, we study the *Brill-Noether wall* and introduce an upper bound for the number of global sections of objects in $\mathcal{D}(X)$ depending only on the geometry of their Harder-Narasimhan polygons at a certain limit point, see Proposition 3.4.

We always assume X is a smooth K3 surface with $\text{Pic}(X) = \mathbb{Z}H$. Given an object $E \in \mathcal{D}(X)$, we denote its Mukai vector by $v(E) = (\text{rk}(E), c(E)H, s(E))$.

Lemma 3.1. *Let $E \in \mathcal{A}(0)$ be a $\sigma_{(0,w)}$ -semistable object with $\phi_{(0,w)}(E) < 1$. Then*

$$v(E)^2 \geq -2c(E)^2.$$

Proof. Let $0 = \tilde{E}_0 \subset \tilde{E}_1 \subset \dots \subset \tilde{E}_{n-1} \subset \tilde{E}_n = E$ be the Jordan-Hölder filtration of E with respect to the stability condition $\sigma_{(0,w)}$. Since the stable factors $E_i = \tilde{E}_i / \tilde{E}_{i-1}$ have the same phase as E , we have $\text{Im}[Z_{(0,w)}(E_i)] = c(E_i) > 0$. Therefore, the length of the filtration n is at most $c(E)$. Given two factors E_i and E_j , we know $\text{Hom}_X(E_i, E_j) = 0$ if $E_i \not\cong E_j$ and $\text{Hom}_X(E_i, E_i) = \mathbb{C}$. Thus, for any $0 < i, j \leq n$,

$$\langle v(E_i), v(E_j) \rangle = -\text{hom}_X(E_i, E_j) + \text{hom}_X^1(E_i, E_j) - \text{hom}_X(E_j, E_i) \geq -2,$$

which implies

$$v(E)^2 = \sum_{i=1}^n v(E_i)^2 + 2 \sum_{1 \leq i < j \leq n} \langle v(E_i), v(E_j) \rangle \geq -2n^2 \geq -2c(E)^2.$$

□

A generalization of the argument in [Bay18, Section 6] implies the following lemma.

Lemma 3.2. (Brill-Noether wall) *Let $\sigma_{(b_0, w_0)}$ be a stability condition with $b_0 < 0$ and $k(b_0, w_0)$ sufficiently close to the point $\text{pr}(v(\mathcal{O}_X)) = (0, 1) = o'$. Let $E \in \mathcal{D}(X)$ be a $\sigma_{(b_0, w_0)}$ -semistable object with the same phase as the structure sheaf \mathcal{O}_X . Then*

$$(9) \quad h^0(X, E) \leq \frac{\chi(E)}{2} + \frac{\sqrt{(rk(E) - s(E))^2 + c(E)^2(2H^2 + 4)}}{2},$$

where $h^0(X, E) = \dim_{\mathbb{C}} \text{Hom}_X(\mathcal{O}_X, E)$ and $\chi(E) = rk(E) + s(E)$ is the Euler characteristic of E .

Proof. We first claim that the structure sheaf \mathcal{O}_X is $\sigma_{(b,w)}$ -stable where $k(b, w)$ is sufficiently close to the point $\text{pr}(v(\mathcal{O}_X)) = o'$. By Lemma 2.15, \mathcal{O}_X is $\sigma_{(0,w)}$ -stable where $k(0, w)$ is on the line segment $(\overline{oo'})$, i.e. $w > \sqrt{2/H^2}$. Moreover, Lemma 2.14 implies that there is no wall for \mathcal{O}_X passing the line with equation $y = nx$ for any $n \in \mathbb{Z}$, because for any stability condition $\sigma_{(b,w)}$ on such a line, we have $b = \frac{1}{n}$ and

$$|\text{Im}[Z_{(b,w)}(\mathcal{O}_X)]| = \frac{1}{n} = \min \left\{ |\text{Im}[Z_{(b,w)}(r, cH, s)]| = \left| c - \frac{r}{n} \right| \neq 0 \right\}.$$

If the object E satisfies $c(E) = 0$, then the projection $\text{pr}(v(E))$ lies on the y -axis. Remark 2.5 implies that \mathcal{O}_X cannot have the same phase as E with respect to $\sigma_{(b_0, w_0)}$ with $b_0 < 0$ unless $\text{pr}(v(E)) = \text{pr}(v(\mathcal{O}_X))$, i.e. $v(E) = kv(\mathcal{O}_X)$. Thus the uniqueness of spherical sheaf with Mukai vector $(1, 0, 1)$ (see e.g. [Muk87, Corollary 3.5]) implies that E is the direct sum of k -copies of \mathcal{O}_X , hence the inequality (9) holds. Thus we may assume $c(E) \neq 0$. Let L_E be the line through o' which passes the point $\text{pr}(v(E))$ if $s(E) \neq 0$, or it has slope $rk(E)/c(E)$ if $s(E) = 0$. By assumption, the point $k(b_0, w_0)$ is on the line L_E . Since we assumed $b_0 < 0$, the structure sheaf \mathcal{O}_X is in the heart $\mathcal{A}(b_0)$. This implies $E \in \mathcal{A}(b_0)$ because E is $\sigma_{(b_0, w_0)}$ -semistable of the same phase as \mathcal{O}_X . When we deform the stability condition $\sigma_{(b_0, w_0)}$ along the line L_E towards the point o' , the object E remains in the corresponding heart, by Remark 2.7. Hence $\lim_{b \rightarrow 0^-} \text{Im}[Z_{(b,w)}(E)] \geq 0$ which implies $c(E) > 0$.

We may assume $\dim \operatorname{Hom}_X(\mathcal{O}_X, E) \neq 0$, otherwise there is nothing to prove. Consider the evaluation map $\operatorname{ev}: \operatorname{Hom}_X(\mathcal{O}_X, E) \otimes \mathcal{O}_X \rightarrow E$. As we saw above, the structure sheaf \mathcal{O}_X is $\sigma_{(b_0, w_0)}$ -stable, so it is a simple object in the abelian category of semistable objects with the same phase as \mathcal{O}_X . Therefore, the morphism ev is injective and the cokernel $\operatorname{cok}(\operatorname{ev})$ is also $\sigma_{(b_0, w_0)}$ -semistable. Let $\{E_i\}_{i=1}^{i=n}$ be the Jordan-Hölder factors of $\operatorname{cok}(\operatorname{ev})$ with respect to the stability condition $\sigma_{(b_0, w_0)}$. By Remark 2.5, the Mukai vector of any factor can be written as $v(E_i) = m_i v(\mathcal{O}_X) + t_i v(E)$ for some $m_i, t_i \in \mathbb{R}$. Note that $\sum_{i=1}^n v(E_i) = v(E) - h^0(E) v(\mathcal{O}_X)$.

If we deform the stability condition $\sigma_{(b_0, w_0)}$ along the line L_E towards the point o' , Remark 2.7 shows that the objects E_i remain stable and of the same phase as E and \mathcal{O}_X . Thus, in particular, they remain in the heart and

$$\lim_{k(b, w) \rightarrow (0^-, 1)} \operatorname{Im}[Z_{(b, w)}(E_i)] = \lim_{b \rightarrow 0^-} [t_i(c(E) - b \operatorname{rk}(E)) + m_i(-b)] \geq 0.$$

This gives $t_i \geq 0$. We have $\sum_{i=1}^n c(E_i) = \sum_{i=1}^n t_i c(E) = c(E)$, therefore

$$(10) \quad \sum_{i=1}^n t_i = 1$$

If $t_i = 0$, then since $v(E_i)^2 \geq -2$, we have $m_i = 1$ so the uniqueness of spherical sheaf again implies $E_i \cong \mathcal{O}_X$. We have $c(E_i) = t_i c(E) \in \mathbb{Z}$. Combining this with (10) proves that the maximum number of factors with $t_i \neq 0$ is equal to $c(E)$. By reordering of the factors, we can assume $E_i \cong \mathcal{O}_X$ for $1 \leq i \leq i_0$ and the other factors satisfy $t_i \neq 0$. Therefore,

$$v(E) - (h^0(X, E) + i_0)v(\mathcal{O}_X) = \sum_{i=i_0+1}^n w_i$$

where $0 \leq n - i_0 \leq k$. Since $\langle w_i, w_j \rangle \geq -2$ for $1 \leq i, j \leq n$, the same argument as in Lemma 3.1 implies that

$$(v(E) - (h^0(X, E) + i_0)v(\mathcal{O}_X))^2 = \left(\sum_{i=i_0+1}^n w_i \right)^2 \geq -2c(E)^2.$$

Now solving the quadratic equation

$$f(x) = (v(E) - xv(\mathcal{O}_X))^2 + 2k^2 = -2x^2 + 2x\chi(E) + v(E)^2 + 2c(E)^2 = 0$$

shows that

$$(11) \quad h^0(X, E) \leq h^0(X, E) + i_0 \leq \frac{\chi(E)}{2} + \frac{\sqrt{(\operatorname{rk}(E) - s(E))^2 + (2H^2 + 4)c(E)^2}}{2}.$$

□

Definition 3.3. Given a stability condition $\sigma_{(b, w)}$ and an object $E \in \mathcal{A}(b)$, the Harder-Narasimhan polygon of E is the convex hull of the points $Z_{(b, w)}(E')$ for all subobjects $E' \subset E$ of E .

If the Harder-Narasimhan filtration of E is the sequence

$$0 = \tilde{E}_0 \subset \tilde{E}_1 \subset \dots \subset \tilde{E}_{n-1} \subset \tilde{E}_n = E,$$

then the points $\{p_i = Z_{(b,w)}(\tilde{E}_i)\}_{i=0}^{i=n}$ are the extremal points of the Harder-Narasimhan polygon of E on the left side of the line segment $\overline{oZ_{(b,w)}(E)}$, see Figure 5.




FIGURE 5. The HN polygon is in the grey area.

We define the following non-standard norm on \mathbb{C} :

$$(12) \quad \|x + iy\| = \sqrt{x^2 + (2H^2 + 4)y^2}.$$

For two points p and q on the complex plane, the length of the line segment \overline{pq} induced by the above norm is denoted by $\|\overline{pq}\|$. The function $\overline{Z}: K(X) \rightarrow \mathbb{C}$ is defined as

$$\overline{Z}(E) = Z_{(0, \sqrt{2/H^2})}(E) = \text{rk}(E) - s(E) + i c(E).$$

The next proposition shows that we can bound the number of global sections of an object in $\mathcal{A}(0)$ via the length of the Harder-Narasimhan polygon at some limit point.

Proposition 3.4. *Consider an object $E \in \mathcal{A}(0)$ which has no subobject $F \subset E$ in $\mathcal{A}(0)$ with $ch_1(F) = 0$.*

- (a) *There exists $w^* > \sqrt{2/H^2}$ such that the Harder-Narasimhan filtration of E is a fixed sequence*

$$0 = \tilde{E}_0 \subset \tilde{E}_1 \subset \dots \subset \tilde{E}_{n-1} \subset \tilde{E}_n = E,$$

for all stability conditions $\sigma_{(0,w)}$ where $\sqrt{2/H^2} < w < w^$.*

- (b) *Let $p_i := \overline{Z}(\tilde{E}_i)$ for $0 \leq i \leq n$, then*

$$h^0(X, E) \leq \frac{\chi(E)}{2} + \frac{1}{2} \sum_{i=1}^n \|\overline{p_i p_{i-1}}\|.$$

Proof. We first show that there exists $w_1 > \sqrt{2/H^2}$ such that the semistable factor \tilde{E}_1 is fixed for the stability conditions of form $\sigma_{(0,w)}$ where $\sqrt{2/H^2} < w < w_1$. Let $\sigma_{(0,w)}$ be a stability condition such that $\sqrt{2/H^2} < w < \sqrt{4/H^2} := w_0$ and $v_1 = (r_1, c_1 H, s_1)$ be a possible class of the semistable factor \tilde{E}_1 . We have $0 < \text{Im}[Z_{(0,w)}(\tilde{E}_1)] = c_1 < \text{Im}[Z_{(0,w)}(E)] = c(E)$. Lemma 3.1 implies that

$$r_1 s_1 \leq c_1^2 \left(\frac{H^2}{2} + 1 \right) \leq c(E)^2 \left(\frac{H^2}{2} + 1 \right).$$

Hence, if $r_1 s_1 > 0$, there are only finitely many possibilities for the class v_1 . If $r_1 \geq 0$ and $s_1 \leq 0$, then since $\phi_{(0,w)}(v(E)) \leq \phi_{(0,w)}(v_1)$, we have

$$\max\{r_1, -s_1\} \leq \frac{r_1 H^2 w^2}{2} - s_1 = \text{Re}[Z_{(0,w)}(v_1)] \leq \max\{\text{Re}[Z_{(0,w)}(v(E))], 0\},$$

and if $r_1 \leq 0$ and $s_1 \geq 0$, the existence of HN filtration for E at $\sigma_{(0,w_0)}$ implies that there exists a real number M_0 such that

$$M_0 \leq \text{Re}[Z_{(0,w_0)}(v_1)] = \frac{r_1 H^2 w_0^2}{2} - s_1 \leq r_1 - s_1.$$

Thus in any case, there are only finitely many possibilities for v_1 . Note that the heart $\mathcal{A}(0)$ for the stability conditions $\sigma_{(0,w)}$ is fixed and does not depend on w . Moreover, the ordering of the phase function $\phi_{(0,w)}$ is the same as the ordering given by the linear function $-\frac{\text{Re}[Z_{(0,w)}(-)]}{\text{Im}[Z_{(0,w)}(-)]}$. Let \tilde{E}_1 be the semistable subobject of E of maximum phase in the HN filtration of E with respect to $\sigma_{(0,w)}$ where $w = w_0$. When we decrease w , the subobject \tilde{E}_1 in the HN filtration changes if another subobject of E gets bigger phase. Since there are only finitely many possibilities for the class of these subobjects which achieve the maximum phase, there is $w_1 > \sqrt{2/H^2}$ such that the subobject \tilde{E}_1 is fixed in the HN filtration of E with respect to $\sigma_{(0,w)}$ when $w \in (\sqrt{2/H^2}, w_1)$.

Continuing this argument by induction, one shows that there is a number w_i such that $\sqrt{2/H^2} < w_i < w_{i-1}$ and the semistable factor $E_i = \tilde{E}_i / \tilde{E}_{i-1}$ which is the semistable subobject of E / \tilde{E}_{i-1} with the maximum phase, is fixed for the stability conditions $\sigma_{(0,w)}$ where $\sqrt{2/H^2} < w < w_i$. Note that $0 < \text{Im}[Z_{(0,w)}(E_i)] < c(E)$, so the length of the HN filtration of E is at most $c(E)$. This completes the proof of (a).

Since $c(E_i) \neq 0$ for $1 \leq i \leq n$, the point $\overline{pr}(v(E_i))$ is not on the y -axis. Proposition 2.6, part (d) implies that the line segment $\overline{o o'}$ is not a wall for the semistable factor E_i . Moreover, since E_i is $\sigma_{(0,w)}$ -semistable for $\sqrt{2/H^2} < w < w^*$, there is no wall for E_i which passes the line segment $\overline{o' o^*}$ where $o^* = k(0, w^*)$. In other words, these stability conditions are all inside one chamber for E_i . If $s(E_i) \neq 0$, we define V_i as a cone in \mathbb{R}^2 with two rays $\overline{pr_i o'}$ and $\overline{pr_i o^*}$ where $pr_i := \overline{pr}(v(E_i))$ and if $s(E_i) = 0$, then V_i is defined as the area between two parallel lines of slope $\text{rk}(E_i)/c(E_i)$ which pass through the points o' and o^* , see Figure 6.

Lemma 2.8 implies that there is a small rectangle $a_1 a_2 a_3 a_4$ around the point $o' = (0, 1)$ such that there is no projection of roots other than $\overline{pr}(v(\mathcal{O}_X))$ inside it. Let V'_i be the

intersection of V_i and the rectangle $a_1 a_2 a_3 a_4$, see the dashed area in Figure 6. The structure of the wall and chamber decomposition implies that E_i is semistable with respect to the stability conditions in V'_i . In particular, it is $\sigma_i := \sigma_{(\tilde{b}_i, \tilde{w}_i)}$ -semistable where σ_i is on the top boundary of V'_i , i.e. the associated point $k(\tilde{b}_i, \tilde{w}_i)$ is on the top boundary of V'_i . In the figures, by abuse of notation, we denote the point $k(\tilde{b}_i, \tilde{w}_i)$ by the corresponding stability condition σ_i .




FIGURE 6. The object E_i remains semistable when we go to σ_i

We may assume $-1 \ll \tilde{b}_i < 0$. Since $E_i \in \mathcal{A}(0)$ is of phase less than one, it remains in the heart $\mathcal{A}(\tilde{b}_i)$. By Remark 2.5, the objects E_i and \mathcal{O}_X have the same phase with respect to σ_i . Hence Lemma 3.2 gives

$$h^0(X, E_i) \leq \frac{\text{rk}(E_i) + s(E_i) + \sqrt{(\text{rk}(E_i) - s(E_i))^2 + c(E_i)^2(2H^2 + 4)}}{2} = \frac{\text{rk}(E_i) + s(E_i)}{2} + \frac{\|\overline{p_i p_{i-1}}\|}{2}.$$

Thus,

$$h^0(X, E) \leq \sum_{i=1}^n h^0(X, E_i) \leq \frac{\chi(E)}{2} + \frac{1}{2} \sum_{i=1}^n \|\overline{p_i p_{i-1}}\|.$$

□

We denote by P_E the convex hull of the points $\{p_0, p_1, \dots, p_n\}$ as defined in Proposition 3.4, part (b). We think of P_E as the Harder-Narasimhan polygon of E on the left at the limit point. We finish this section by stating two useful inequalities which are the result of deformation of stability conditions.

Lemma 3.5. *Consider a stability condition $\sigma_{(b_0, w_0)}$ and a $\sigma_{(b_0, w_0)}$ -semistable object $E \in \mathcal{A}(b_0)$ with $c(E) \neq 0$. Let L be a line through the point $\text{pr}(v(E))$ if $s(E) \neq 0$ or it has a slope of $\text{rk}(E)/c(E)$ if $s(E) = 0$. Let $q_1 = (x_1, y_1)$ and $q_2 = (x_2, y_2)$ be two points on the line L where $y_1 y_2 \neq 0$ and $x_1/y_1 \leq x_2/y_2$. Suppose the point $k(b_0, w_0)$ is on the open line segment $(\overline{q_1 q_2})$. If every point on the open line segment $(\overline{q_1 q_2})$ is in correspondence to a stability condition, i.e., if $(\overline{q_1 q_2}) \subset V(X)$, then*

$$(13) \quad \mu_H^+(H^{-1}(E)) \leq \frac{x_1}{y_1} \quad \text{and} \quad \frac{x_2}{y_2} \leq \mu_H^-(H^0(E)).$$

Proof. Remark 2.7 implies that E is $\sigma_{(b,w)}$ -semistable and it is in the heart $\mathcal{A}(b)$ whenever the point $k(b, w)$ is on the open line segment $(\overline{q_1 q_2})$. Therefore,

$$\mu_H^+(H^{-1}(E)) \leq b < \mu_H^-(H^0(E)).$$

If $k(b_i, w_i) = q_i$, then $b_i = x_i/y_i$. Thus the stability conditions close to the points q_1 or q_2 give the inequalities (13). \square

4. THE BRILL-NOETHER LOCI

In this section, we first show that the morphism $\psi: M_{X,H}(v) \rightarrow \mathcal{BN}$ described in (1) is well-defined. Then we consider a slope semistable rank r -vector bundle F on the curve C of degree $2rs$ and describe the location of the wall that bounds the Gieseker chamber for the push-forward of F . Finally, in Proposition 4.4, we show that if the number of global sections of F is high enough, then it must be the restriction of a vector bundle on the surface.

We assume throughout Section 4 that X is a K3 surface with $\text{Pic}(X) = \mathbb{Z}H$ and $H^2 = 2rs$ for some $r \geq 2$ and $s \geq \max\{r, 5\}$. We also assume C is a curve in the linear system $|H|$ and $i: C \hookrightarrow X$ is the embedding of the curve C into the surface X . The push-forward of a rank r -vector bundle F on the curve C of degree $2rs$ has Mukai vector $v(i_*F) = (0, rH, 2rs - r^2s)$. Let $M_{X,H}(v)$ be the moduli space of H -Gieseker semistable sheaves on the surface X with Mukai vector $v = (r, H, s)$. Since $v^2 = 0$ and v is primitive, the moduli space $M_{X,H}(v)$ is a smooth projective K3 surface [Huy16, Proposition 10.2.5 and Corollary 10.3.5]. Any coherent sheaf $E \in M_{X,H}(v)$ is a μ_H -stable locally free sheaf [HL10, Remark 6.1.9]. Note that $E(-H)$ is also μ_H -stable. Let $u := v(i_*F) - v = v(E(-H)[1])$,

$$p_v := pr(v) = \left(\frac{1}{s}, \frac{r}{s}\right) \quad \text{and} \quad p_u := pr(u) = \left(\frac{-1}{s(r-1)}, \frac{r}{s(r-1)^2}\right).$$

We also denote by \tilde{o} the point at which the line segments $\overline{p_u p_v}$ and $\overline{o' o}$ intersect, where $o' = pr(v(\mathcal{O}_X))$. Define the object $K_E \in \mathcal{D}(X)$ as the cone of the evaluation map:

$$(14) \quad \mathcal{O}_X^{h^0(X,E)} \xrightarrow{\text{ev}_E} E \rightarrow K_E.$$

Denote the point $(-1/r, s/r)$ by q . Lemma 2.10 for $m = r$, $n = s(r-1)$ and $\epsilon = 1$ implies that there is no projection of roots in the grey area and on the open line segment (\overline{et}) in Figure 7, where $e = q_{-m, -n, -\epsilon}$ and $t = q'_{-m, -n, -\epsilon}$. As before, we denote by γ_n the point on the parabola $y = rsx^2$ with the x -coordinate $1/n$.

Proposition 4.1. *Let $E \in M_{X,H}(v)$ be a μ_H -stable vector bundle on the surface X . Then we have*

- (a) $\text{Hom}_X(E, E(-H)[1]) = 0$.
- (b) *The restriction $E|_C$ is a slope stable vector bundle on the curve C and $h^0(C, E|_C) = r + s$. In particular, the morphism ψ described in (1) is well-defined.*
- (c) *The object K_E is of the form $K_E = E'[1]$ where E' is a μ_H -stable locally free sheaf on X and $\text{Hom}_X(E', E(-H)[1]) = 0$.*




FIGURE 7. No projection of roots in the grey area

Proof. The objects E and $E(-H)[1]$ have the same phase with respect to the stability condition $\tilde{\sigma} := \sigma_{(0, \tilde{w})}$ where $k(0, \tilde{w}) = \tilde{o}$, see Figure 7. There are no homomorphisms between non-isomorphic stable objects of the same phase. Hence to prove claim (a), we only need to show both E and $E(-H)[1]$ are $\tilde{\sigma}$ -stable.

By [Bri08, Proposition 14.2], the μ_H -stable sheaf E is $\sigma_{(0, w)}$ -stable where $w \gg 0$. Lemma 2.14 implies that there is no wall for E intersecting the line segment $(\overline{oo'})$. Thus E is $\tilde{\sigma}$ -stable. Lemma 2.15 implies that $E(-H)[1]$ with Mukai vector $(-r, (r-1)H, -s(r-1)^2)$ is $\sigma_1 := \sigma_{(b_1, w_1)}$ -stable where $b_1 = -(r-1)/r$ and w_1 is arbitrary. Let e' be the point that the line segment $\overline{qp_u}$ intersects the line $x = b_2 y$ where

$$b_2 = -\frac{r-2}{r-1} \quad \text{if } r > 2 \quad \text{or} \quad b_2 = -\frac{1}{3} \quad \text{if } r = 2.$$

If $s \geq \max\{r, 5\}$, then

$$-\frac{s(r-1)-1}{rs} < b_2 < -\frac{r+1}{rs}.$$

Thus the line segment $\overline{oe'}$ is located between two lines $\overline{o\gamma_{-r-1}}$ and $\overline{o\gamma_{-s(r-1)+1}}$ and it is on the grey area with no projection of roots. We claim that there is no wall for $E(-H)[1]$ intersecting the line segment $(\overline{oe'})$. Consider a stability condition of form $\sigma_{(b_2, w)}$ where the point $k(b_2, w)$ is on $(\overline{oe'})$. If $r > 2$, then

$$|\operatorname{Im}[Z_{(b_2, w)}(E(-H))]| = \frac{1}{r-1} = \min \left\{ |\operatorname{Im}[Z_{(b_2, w)}(r', c'H, s')]| = \left| c' + \frac{r-2}{r-1} r' \right| \neq 0 \right\},$$

and if $r = 2$, then $|\operatorname{Im}[Z_{(b_2, w)}(E(-H))]| = \frac{1}{3}$. Thus in any case, the minimality condition of Lemma 2.14 is satisfied. Hence the stability condition $\sigma_{(b_2, w)}$ cannot be on a wall for $E(-H)[1]$. This in particular implies $E(-H)[1]$ is $\tilde{\sigma}$ -stable which finishes the proof of claim (a).

Consider the short exact sequence

$$(15) \quad E \hookrightarrow i_* E|_C \twoheadrightarrow E(-H)[1].$$

Since E and $E(-H)[1]$ are $\tilde{\sigma}$ -stable of the same phase, their extension $i_* E|_C$ is $\tilde{\sigma}$ -semistable and the objects E and $E(-H)[1]$ are its JH factors. We have $\phi_{(0, w)}(E) < \phi_{(0, w)}(i_* E|_C)$ for $w > \tilde{w}$, thus the object $i_* E|_C$ is $\sigma_{(0, w)}$ -stable and by Lemma 2.13, $E|_C$ is slope stable.

The next step is to show $h^0(X, E) = r + s$. Consider a stability condition of form $\sigma_{(b_3, w_3)} =: \sigma_3$ such that the point $k(b_3, w_3)$ is on the line segment $(\overline{qo'})$ and it is sufficiently close to the point o' . Lemma 2.14 implies that there is no wall for E which intersects the open line segment $(\overline{oo'})$. Therefore $\sigma_{(0, w)}$ -stability of E for $w \gg 0$ implies that it is σ_3 -semistable. Moreover, E has the same phase as the structure sheaf \mathcal{O}_X with respect to σ_3 , thus Lemma 3.2 implies that

$$h^0(X, E) \leq \left\lfloor \frac{r+s}{2} + \frac{\sqrt{(r+s)^2 + 4}}{2} \right\rfloor = r + s.$$

The coherent sheaf E is μ_H -stable and has positive slope, so $\operatorname{Hom}_X(E, \mathcal{O}_X) = 0$ and

$$\chi(E) = r + s = h^0(X, E) - h^1(X, E).$$

Therefore $h^0(X, E) = r + s$ and the object K_E has Mukai vector $v(K_E) = (-s, H, -r)$. On the other hand, since there is no wall for $E(-H)[1]$ which passes the line segment $(\overline{oe'})$, it is stable with respect to the stability conditions on the line segment $(\overline{p_u o'})$, where \mathcal{O}_X is also stable and has the same phase as $E(-H)[1]$. Thus

$$\operatorname{Hom}_X(\mathcal{O}_X, E(-H)[1]) = 0.$$

Thus the short exact sequence (15) gives $h^0(C, E|_C) = h^0(X, E) = r + s$, which completes the proof of (b).

The sheaves E and \mathcal{O}_X are σ_3 -semistable of the same phase. Since \mathcal{O}_X is σ_3 -stable, the evaluation map ev_E defined in (14) is injective in the abelian category of semistable objects with the same phase as \mathcal{O}_X , hence the cokernel K_E is σ_3 -semistable. We claim that K_E is σ_3 -stable. Assume otherwise. Let E_1 be a σ_3 -stable factor of K_E . Remark 2.5 implies that $v(E_1) = t_1 v(E) + s_1 v(\mathcal{O}_X)$. Similar to the argument in Lemma 3.2, we deform the stability condition σ_3 towards the point o' , then

$$0 \leq \lim_{b \rightarrow 0} \operatorname{Im}[Z_{(b, w)}(E_1)] = t_1 c(E) \leq \lim_{b \rightarrow 0} \operatorname{Im}[Z_{(b, w)}(E)] = c(E) = 1.$$

Since $t_1 c(E) = t_1 \in \mathbb{Z}$, we have $t_1 = 0$ or 1 . Therefore, \mathcal{O}_X is a subobject or a quotient of K_E . But $\operatorname{Hom}_X(\mathcal{O}_X, K_E) = 0$ and since $\operatorname{Hom}_X(E, \mathcal{O}_X) = 0$, we have $\operatorname{Hom}_X(K_E, \mathcal{O}_X) = 0$, a contradiction.

Note that K_E has Mukai vector $v(K_E) = v(E) - (r + s)v(\mathcal{O}_X) = (-s, H, -r)$ with the projection $pr(v(K_E)) = q$. Lemma 2.14 shows that there is no wall for K_E intersecting

the open line segment $(\overline{o\sigma'})$. Therefore, it is $\sigma_{(0,w)}$ -stable where $w \gg 0$. By [MS17, Lemma 6.18], $H^0(K_E)$ is zero or a skyscraper sheaf and $H^{-1}(K_E)$ is a μ_H -stable sheaf. If $H^0(K_E) \neq 0$, then for some $k > 0$, we have

$$v(H^{-1}(K_E))^2 = (s, -H, r+k)^2 = -2sk < -2,$$

a contradiction. Therefore $K_E = E'[1]$ for a μ_H -stable coherent sheaf E' on X . Since $E'^{\vee\vee}$ is also μ_H -stable, we have

$$-2 \leq v(E'^{\vee\vee})^2 = v(E)^2 - 2\mathrm{rk}(E')l(E'^{\vee\vee}/E) = -2\mathrm{rk}(E')l(E'^{\vee\vee}/E').$$

Since $\mathrm{rk}(E') = s \geq 5$, we must have $E' = E'^{\vee\vee}$, so E' is a locally free sheaf.

To prove the final claim of part (c), we find a stability condition such that E' and $E(-H)[1]$ are stable of the same phase. Lemma 2.15 implies that $E'[1]$ is $\sigma_4 := \sigma_{(b_4, w_4)}$ -stable where $b_4 = -1/s$ and w_4 is arbitrary. We claim that E' is stable with respect to the stability condition at the point e' . Let t' be the point that the line segment $\overline{qp_u}$ intersects the line given by the equation $y = x(-s+1)$. Then the x -coordinate of the point t' is equal to $-1/(2r-1)$ which is bigger than $-1/(r+1)$ if $r > 2$ and $t' = t$ if $r = 2$. We claim that for $r = 2$ the point $t = (-1/3, (s-1)/3)$ cannot be the projection of a root. Indeed, if there exists a root $\delta = (\tilde{r}, \tilde{c}H, \tilde{s})$ with $pr(\delta) = t$, then

$$\frac{\tilde{c}}{\tilde{s}} = \frac{-1}{3} \quad \text{and} \quad \frac{\tilde{r}}{\tilde{s}} = \frac{s-1}{3}.$$

This implies $|\tilde{s}| \geq 3$. Since $\delta^2 = -2$, we have $\tilde{s}^2(s-3) = 9$ which is impossible for $s \geq 5$. Therefore there is a stability condition corresponding to any point on the line segment $(\overline{ot'})$. By Lemma 2.14, there is no wall for E' intersecting the line segment $(\overline{ot'})$. Thus σ_4 -stability of E' implies that it is stable with respect to the stability condition at the point e' and it has the same phase as $E(-H)[1]$. On the other hand, as we have seen there is no wall for $E(-H)[1]$ intersecting the line segment $(\overline{oe'})$. Thus $E(-H)[1]$ is also stable with respect to the stability condition at the point e' , so there is no non-trivial homomorphism between E' and $E(-H)[1]$. This finishes the proof of (c). \square

4.1. The first wall. Let $M_C(r, 2rs)$ be the moduli space of slope semistable rank r -vector bundles on the curve C of degree $2rs$. Note that the vector bundle F in $M_C(r, 2rs)$ can be strictly semistable. The push-forward of the vector bundle F to the surface X has Mukai vector $v(i_*F) = (0, rH, 2rs - r^2s)$. Lemma 2.13 implies that i_*F is semistable with respect to the stability conditions $\sigma_{(b,w)}$ in the Gieseker chamber which means w is large enough and b is arbitrary. By Proposition 2.6, part (d), any wall for i_*F is part of a line which goes through the point $p := pr(v(i_*F))$ if $r > 2$ or it is a horizontal line segment if $r = 2$. The next proposition describes the location of the wall that bounds the Gieseker chamber for i_*F . Recall that $p_v = pr(v)$ and $p_u = pr(u)$ where $v = (r, H, s)$ and $u = v(i_*F) - v$.

Proposition 4.2. *Given a vector bundle $F \in M_C(r, 2rs)$, the wall that bounds the Gieseker chamber for i_*F is not below the line segment $\overline{p_u p_v}$ and it coincides with the line segment $\overline{p_u p_v}$ if and only if F is the restriction of a vector bundle $E \in M_{X,H}(v)$ to the curve C .*

Proof. Assume that the wall \mathcal{W}_{i_*F} that bounds the Gieseker chamber for i_*F , is below or on the line segment $\overline{p_u p_v}$, see Figure 8.




FIGURE 8. The first wall \mathcal{W}_{i_*F}

Suppose the stability condition $\sigma_{(0,w')}$ is on the wall \mathcal{W}_{i_*F} . Then there is a destabilising sequence $F_1 \hookrightarrow i_*F \twoheadrightarrow F_2$ of objects in $\mathcal{A}(0)$ such that F_1 and F_2 are $\sigma_{(0,w')}$ -semistable of the same phase as i_*F and

$$(16) \quad \phi_{(0,w)}(F_1) > \phi_{(0,w)}(i_*F) \quad \text{for } w < w'.$$

Taking cohomology gives a long exact sequence of sheaves

$$(17) \quad 0 \rightarrow H^{-1}(F_1) \rightarrow 0 \rightarrow H^{-1}(F_2) \rightarrow H^0(F_1) \xrightarrow{d_0} i_*F \xrightarrow{d_1} H^0(F_2) \rightarrow 0.$$

Thus $H^{-1}(F_1) = 0$ and $H^0(F_1) \cong F_1$. Let $v(F_1) = (r', c'H, s')$. If $r' = 0$, then the projection $pr(v(F_1))$ lies on the x -axis. By Remark 2.5, F_1 and i_*F cannot have the same phase with respect to $\sigma_{(0,w')}$ unless $pr(v(F_1)) = pr(v(i_*F))$, i.e. $v(F_1) = kv(i_*F)$ for some $k \in \mathbb{R}$, which is in contradiction to (16). Hence $r' > 0$.

Let $T(F_1)$ be the maximal torsion subsheaf of F_1 and $F_1/T(F_1)$ be its torsion-free part. Let $v(T(F_1)) = (0, \tilde{r}H, \tilde{s})$. Right-exactness of the underived pull-back i^* applied to the short exact sequence $T(F_1) \hookrightarrow F_1 \twoheadrightarrow F_1/T(F_1)$ implies that

$$(18) \quad \text{rank}(i^*F_1) \leq \text{rank}(i^*T(F_1)) + \text{rank}(i^*(F_1/T(F_1))).$$

Since F is a vector bundle on C and $T(F_1)$ is a subsheaf of i_*F , we have $\text{rank}(i^*T(F_1)) = \tilde{r}$. Thus inequality (18) implies that $\text{rank}(i^*F_1) \leq \tilde{r} + r'$. Let $v(H^0(F_2)) = (0, c''H, s'')$. Then the right-exactness of i^* implies

$$(19) \quad \text{rank}(i^* \ker d_1) \leq \text{rank}(i^*F_1) \quad \Rightarrow \quad r - c'' \leq r' + \tilde{r}.$$

Let q_1 and q_2 be the intersection points of the wall \mathcal{W}_{i_*F} with the line segments $\overline{op_u}$ and $\overline{op_v}$, respectively. Then Lemma 3.5 implies that

$$\frac{1}{r} \leq \mu_H^-(F_1) = \mu_H^-(F_1/T(F_1)) \quad \text{and} \quad \mu_H^+(H^{-1}(F_2)) \leq \frac{1-r}{r}.$$

Therefore,

$$\begin{aligned} \frac{r - c'' - \tilde{r}}{r'} &= \mu_H(F_1/T(F_1)) - \mu_H(H^{-1}(F_2)) \geq \\ \mu_H^-(F_1/T(F_1)) - \mu_H^+(H^{-1}(F_2)) &\geq \frac{1}{r} - \frac{1-r}{r} = 1. \end{aligned}$$

Combined with the inequalities (19), this is only possible if all these inequalities are equalities, i.e. $r' = r - c'' - \tilde{r}$,

$$(20) \quad \mu_H^+(H^{-1}(F_2)) = \mu_H(H^{-1}(F_2)) = \frac{1-r}{r},$$

and

$$(21) \quad \frac{c' - \tilde{r}}{r - c'' - \tilde{r}} = \mu_H(F_1/T(F_1)) = \mu_H^-(F_1/T(F_1)) = \frac{1}{r}.$$

Since $c'' \geq 0$ and $\tilde{r} \geq 0$, the denominators of the first and last sentences in (21) imply that $c'' = \tilde{r} = 0$, hence $c' = 1$, $v(F_1) = (r, H, s')$ and $v(T(F_1)) = (0, 0, \tilde{s})$. However $T(F_1)$ cannot be a skyscraper sheaf because $T(F_1)$ is a subsheaf of i_*F . Thus $T(F_1) = 0$ and F_1 is torsion-free. Moreover, the equations (20) and (21) imply that F_1 and $H^{-1}(F_2)$ are μ_H -semistable sheaves. These sheaves are indeed μ_H -stable because their rank and degree $\frac{\text{ch}_1(-) \cdot H}{H^2}$ are co-prime.

The point $pr(v(F_1))$ lies on the line through the points q_1 and q_2 . Moreover, $\frac{\text{rk}(F_1)}{c(F_1)} = r$, so it also lies on the line through the origin and p_v . Thus $pr(v(F_1)) = q_2$, see Figure 8. On the other hand, μ_H -stability of F_1 implies that $v(F_1)^2 = 2r(s - s') \geq -2$, i.e. $s' \leq s$. Hence q_2 cannot be on the open line segment $(\overline{op_v})$. Therefore, the wall \mathcal{W}_{i_*F} is above or on the line segment $\overline{p_u p_v}$.

If the wall \mathcal{W}_{i_*F} coincides with the line segment $\overline{p_u p_v}$, then $v(F_1) = (r, H, s)$. The non-zero morphism d_0 in the long exact sequence (17) factors via the morphism $d'_0: i_*F_1|_C \rightarrow i_*F$. The objects $i_*F_1|_C$ and i_*F have the same Mukai vector and so have the same phase. Proposition 4.1 implies that $i_*F_1|_C$ is H -Gieseker stable. Hence the morphism d'_0 is injective. Since $i_*F_1|_C$ and i_*F have the same Mukai vector, we must have isomorphism $F_1|_C \cong F$. \square

Now instead of checking the possible walls above the line segment $\overline{p_u p_v}$, we consider the stability conditions of form $\sigma_{(0,w)}$ which are close to the point $(0, 1)$ and examine the Harder-Narasimhan filtrations. Given a semistable vector bundle $F \in M_C(r, 2rs)$, the $\sigma_{(0,w)}$ -semistability of i_*F for $w \gg 0$ implies that it does not have any subobject F' in $\mathcal{A}(0)$ with $\text{ch}_1(F') = 0$ because $\phi_{(0,w)}(F') = 1 > \phi_{(0,w)}(i_*F)$. Proposition 3.4 for i_*F implies that there exists $w^* > 0$ such that for every stability condition $\sigma_{(0,w)}$ where $\sqrt{1/(rs)} < w < w^*$, the HN filtration of i_*F is a fixed sequence $0 = \tilde{E}_0 \subset \tilde{E}_1 \subset \dots \subset \tilde{E}_{n-1} \subset \tilde{E}_n = i_*F$ with the semistable factors $E_i = \tilde{E}_i/\tilde{E}_{i-1}$ for $1 \leq i \leq n$. Recall that the stability function $\overline{Z}: K(X) \rightarrow \mathbb{C}$ is defined as $\overline{Z}(E) = Z_{(0, \sqrt{1/rs})}(E) = \text{rk}(E) - s(E) + ic(E)$ and the

polygon P_{i_*F} has vertices $\{p_i\}_{i=0}^{i=n}$ where $p_i = \overline{Z}(\tilde{E}_i)$. Let T be a triangle with the vertices $z_1 := \overline{Z}(v) = r - s + i$, $z_2 := \overline{Z}(i_*F) = r^2s - 2rs + i$ and the origin.

Lemma 4.3. *The polygon P_{i_*F} for any vector bundle $F \in M_C(r, 2rs)$ is contained in the triangle $T = \triangle oz_1z_2$ and they coincide if and only if the bundle F is the restriction of a vector bundle $E \in M_{X,H}(v)$ to the curve C , see Figure 9.*




FIGURE 9. The polygon P_{i_*F} is inside the triangle T

Proof. The polygon P_{i_*F} is convex. Thus for the first statement we only need to show the following two conditions are satisfied: firstly, the line through $\overline{op_1}$ is between the lines through $\overline{oz_1}$ and $\overline{oz_2}$, and secondly the line through $\overline{z_2p_{n-1}}$ is between the lines through $\overline{z_2z_1}$ and $\overline{z_2o}$. By definition, the points $\{p_i\}_{i=0}^{i=n}$ are on the left hand-side of the line segment $\overline{oz_2}$. Therefore it suffices to show that

$$(22) \quad -\frac{\operatorname{Re}[\overline{Z}(E_1)]}{\operatorname{Im}[\overline{Z}(E_1)]} \leq -\frac{\operatorname{Re}[z_1]}{\operatorname{Im}[z_1]} \quad \text{and} \quad -\frac{\operatorname{Re}[z_2 - z_1]}{\operatorname{Im}[z_2 - z_1]} \leq -\frac{\operatorname{Re}[\overline{Z}(E_n)]}{\operatorname{Im}[\overline{Z}(E_n)]}.$$

Let $v(E_1) = (r_1, c_1H, s_1)$. We have $0 < \operatorname{Im}[Z_{(0,w)}(E_1)] = c_1 \leq \operatorname{Im}[Z_{(0,w)}(i_*F)] = r$. Assume for a contradiction that the first inequality in (22) does not hold, then

$$(23) \quad \frac{s_1}{c_1} - \frac{r_1}{c_1} > s - r.$$

Therefore, the point $q_1 := (r_1/c_1, s_1/c_1)$ is above the line L_1 given by the equation $y - x = s - r$. Proposition 4.2 implies that i_*F is $\sigma_{(0,\tilde{w})}$ -semistable where $k(0, \tilde{w}) = \tilde{o}$ is on the line segment $\overline{p_u p_v}$, see Figure 8. Therefore,

$$(24) \quad \phi_{(0,\tilde{w})}(E_1) \leq \phi_{(0,\tilde{w})}(v) \quad \Rightarrow \quad \frac{s_1}{c_1} - \frac{r_1}{c_1}(rs\tilde{w}^2) \leq s - r(rs\tilde{w}^2).$$

This shows q_1 is below or on the line L_2 given by the equation $y - x(rs\tilde{w}^2) = s - r^2s\tilde{w}^2$, see Figure 10. Since the point of intersection of the lines L_1 and L_2 is (r, s) , we must have

$$r < \frac{r_1}{c_1} \quad \Rightarrow \quad r \leq \frac{r_1}{c_1} - \frac{1}{c_1} \leq \frac{r_1}{c_1} - \frac{1}{r}.$$

Therefore, the point q_1 is in the dashed area in Figure 10. The point on the line L_1 with the first coordinate $r + 1/r$, which is denoted by q' , has the second coordinate $s + 1/r$. On the other hand, E_1 is $\sigma_{(0,w)}$ -semistable where $\sqrt{1/(rs)} < w < w^*$, so Lemma 3.1 implies that

$$-2c_1^2 \leq c_1^2(H^2) - 2r_1s_1 = c_1^2(2rs) - 2r_1s_1 \Rightarrow \frac{r_1s_1}{c_1^2} \leq rs + 1.$$

Therefore, the point q_1 is below or on the hyperbola with equation $xy = rs + 1$. But the point q' and so whole of the dashed area is above the hyperbola, a contradiction.




FIGURE 10. The point q_1 is in the dashed area

Similarly, if the semistable factor E_n with Mukai vector $v(E_n) = (r_n, c_nH, s_n)$ does not satisfy the second inequality in (22), then the point $q_n := (r_n/c_n, s_n/c_n)$ is below the line L'_1 by the equation $y = x - s(r - 1) + r/(r - 1)$ and is above or on the line L'_2 with the equation $y = x(rs\tilde{w}^2) - s(r - 1) + r^2s\tilde{w}^2/(r - 1)$. Since the point of intersection of these two lines is $(-r/(r - 1), -s(r - 1))$, we have

$$\frac{r_n}{c_n} < \frac{-r}{r - 1} \Rightarrow \frac{r_n}{c_n} \leq \frac{-r}{r - 1} - \frac{1}{c_n(r - 1)} \leq \frac{-r}{r - 1} - \frac{1}{(r - 1)^2}.$$

Then the same argument as above leads to a contradiction for $s \geq r$.

If the vector bundle F is the restriction of a vector bundle $E \in M_{X,H}(v)$, then Proposition 4.1 implies that the HN factors of $i_*E|_C$ with respect to the stability conditions close to the point $(0, 1)$, are E and $E(-H)[1]$. Therefore, the polygon P_{i_*F} coincides with the triangle T . Conversely, assume for a vector bundle $F \in M_C(r, 2rs)$, we have $P_{i_*F} = T$. Then $z_1 = p_1$ so $v(E_1) = (r + k, H, s + k)$ for some $k \in \mathbb{Z}$. The point $q_1 = (r + k, s + k)$ is on the line L_1 . Proposition 4.2 and inequality (24) imply that the point q_1 is below on the line L_2 , thus $k \geq 0$ (see Figure 10). Since $c(E_1) = 1$ is minimal, lemma 2.14 implies

that E_1 is $\sigma_{(0,w)}$ -stable. Therefore $v(E_1)^2 = -2k(r+k+s) \geq -2$ which gives $k = 0$ and Proposition 4.2 implies that F is the restriction of the vector bundle $E_1 \in M_{X,H}(v)$. \square

4.2. The maximum number of global sections. The next proposition shows that any vector bundle $F \in M_C(r, 2rs)$ with high enough number of global sections is the restriction of a vector bundle on the surface.

Proposition 4.4. *Let F be a slope-semistable rank r -vector bundle on the curve C of degree $2rs$, where $r \geq 2$ and $s \geq \max\{5, r\}$. If $h^0(C, F) \geq r + s$, then F is the restriction of a unique vector bundle $E \in M_{X,H}(v)$ to the curve C . In other words, the morphism $\psi: M_{X,H}(v) \rightarrow \mathcal{BN} = M_C(r, 2rs, r+s)$, which sends a vector bundle to its restriction, is bijective.*

Proof. If the vector bundle $F \in \mathcal{BN}$ is the restriction of a vector bundle $E \in M_{X,H}(v)$, then E is a Harder-Narasimhan factor of i_*F with respect to $\sigma_{(0,w)}$ where $\sqrt{1/rs} < w < w^*$. Thus the uniqueness of the Harder-Narasimhan filtration implies that ψ is injective.

For the surjectivity part, by Lemma 4.3 we only need to show for any $F \in \mathcal{BN}$ the polygon P_{i_*F} coincides with the triangle $T = \triangle oz_1z_2$. Assume for a contradiction that P_{i_*F} is strictly inside T . Since the vertices of P_{i_*F} are Gaussian integers, P_{i_*F} must be contained in the polygon $oz'_1z'_2z_2$, where $z'_1 = r-s+1+i$ and $z'_2 = s(r-2)+r-r/(r-1)+2i$, see Figure 11.




FIGURE 11. The polygon P_{i_*F} is inside the polygon $oz'_1z'_2z_2$

The convexity of the polygon P_{i_*F} and the polygon $oz'_1z'_2z_2$ gives

$$\sum_{i=1}^n \|\overline{p_i p_{i-1}}\| \leq \|\overline{oz'_1}\| + \|\overline{z'_1 z'_2}\| + \|\overline{z'_2 z_2}\| =: l_{in}.$$

Note that $\|\cdot\|$ is the non-standard norm defined in (12). In our case, we have $H^2 = 2rs$, so $\|x + iy\| = \sqrt{x^2 + (4rs + 4)y^2}$. Let $l := \|\overline{oz_1}\| + \|\overline{z_1 z_2}\|$, then

$$l - l_{in} = \|\overline{oz_1}\| - \|\overline{oz'_1}\| + \|\overline{z_1 z'_2}\| - \|\overline{z'_1 z'_2}\|.$$

Define $\epsilon := \frac{2rs-r^2s}{2} + \frac{l}{2} - (r+s)$. Since $r+s \leq h^0(F)$, Proposition 3.4 implies

$$\frac{2rs-r^2s}{2} + \frac{l}{2} - \epsilon = r+s \leq h^0(F) \leq \frac{\chi(F)}{2} + \frac{1}{2} \sum_{i=1}^n \|\overline{p_i p_{i-1}}\| \leq \frac{2rs-r^2s}{2} + \frac{l_{\text{in}}}{2}.$$

Thus $l - l_{\text{in}} \leq 2\epsilon$. We have $l = \sqrt{(s+r)^2+4} + \sqrt{(s(r-1)^2+r)^2+4(r-1)^2}$, so

$$2\epsilon = \frac{4}{\sqrt{(r+s)^2+4} + (r+s)} + \frac{4(r-1)^2}{\sqrt{(s(r-1)^2+r)^2+4(r-1)^2} + (s(r-1)^2+r)}.$$

Therefore, we have $2\epsilon \leq \frac{4}{2(r+s)} + \frac{4(r-1)^2}{2(s(r-1)^2+r)} \leq \frac{2}{r+s} + \frac{2}{s} \leq \frac{24}{35}$, because we assumed $r \geq 2$ and $s \geq \max\{r, 5\}$. We will show $l - l_{\text{in}} \geq 0.6868$, thus $l - l_{\text{in}} > 2\epsilon$ which gives a contradiction.

We have $\|\overline{oz_1}\| - \|\overline{oz'_1}\| = \sqrt{4rs+4+(r-s)^2} - \sqrt{4rs+4+(r-s+1)^2}$, hence

$$f_1(r, s) := \frac{2s-2r-1}{2\sqrt{(r+s)^2+4}} \leq \frac{2s-2r-1}{\sqrt{(r+s)^2+4} + \sqrt{(r+s)^2+5+2(r-s)}} = \|\overline{oz_1}\| - \|\overline{oz'_1}\|.$$

Also, $\|\overline{z_1 z'_2}\| - \|\overline{z'_1 z'_2}\| = \sqrt{4rs+4+\left(\frac{s(r-1)^2-r}{r-1}\right)^2} - \sqrt{4rs+4+\left(\frac{s(r-1)^2-r}{r-1}-1\right)^2}$, thus

$$f_2(r, s) := \frac{\frac{s(r-1)^2+r}{r-1} - \frac{2}{r-1} - \frac{5}{2}}{\sqrt{4+\left(\frac{s(r-1)^2+r}{r-1}\right)^2}} = \frac{\frac{s(r-1)^2-r}{r-1} - \frac{1}{2}}{\sqrt{4rs+4+\left(\frac{s(r-1)^2-r}{r-1}\right)^2}} \leq \|\overline{z_1 z'_2}\| - \|\overline{z'_1 z'_2}\|.$$

The function $f_1(r, s)$ is positive unless $s = r \geq 5$, and $f_2(r, s) = \sqrt{\frac{m}{4+m}} + \frac{-\frac{2}{r-1}-\frac{5}{2}}{\sqrt{4+m}}$ where $m = \left(\frac{s(r-1)^2+r}{r-1}\right)^2$, so f_2 is an increasing function with respect to both r and s . Consider the following three cases:

- If $r \geq 4$, then $f_1(r, s) \geq f_1(s, s) \geq f_1(5, 5) \geq -0.05$ and $f_2(r, s) \geq f_2(4, 5) \geq 0.78$.
- If $r = 3$, then $f_1(3, s) = \frac{s-3.5}{\sqrt{(s+3)^2+4}} \geq f_1(3, 5) \geq 0.18$ and $f_2(3, s) \geq f_2(3, 5) \geq 0.685$.
- If $r = 2$, then $f_1(2, s) + f_2(2, s) = 2 \cdot \frac{s-2.5}{\sqrt{(s+2)^2+4}} \geq \frac{5}{\sqrt{53}} \geq 0.6868$.

Therefore, $l - l_{\text{in}} \geq 0.6868$ as claimed, which is a contradiction. Thus the polygon P_{i_*F} coincides with the triangle T and the morphism $\psi: M_{X,H}(v) \rightarrow \mathcal{BN}$ is surjective in case (A). \square

Corollary 4.5. *Let F be a slope-semistable rank r -vector bundle on the curve C of degree $2rs$ such that $r \geq 2$ and $s \geq \max\{5, r\}$. Then $h^0(F) \leq r+s$ and if $h^0(F) = r+s$, then F is slope-stable.*

Proof. Using the same notations as in the proof of Proposition 4.4, we have

$$h^0(F) \leq \frac{2rs - r^2s}{2} + \frac{1}{2} \sum_{i=1}^n \|\overline{p_i p_{i-1}}\| \leq \frac{2rs - r^2s}{2} + \frac{l}{2} = (r+s) + \epsilon \stackrel{(*)}{<} r+s+1,$$

where $(*)$ is the result of the above computation which shows $\epsilon \leq \frac{12}{35}$. If $h^0(F) = r+s$, then by Proposition 4.4, F is the restriction of a stable vector bundle on the surface, so it is stable. \square

5. THE FINAL RESULTS

In this section we prove the main results.

Proof of Theorem 1.2. By Proposition 4.4, the morphism $\psi: M_{X,H}(v) \rightarrow \mathcal{BN}$ is bijective. Therefore Proposition 4.1 implies that any vector bundle F in the Brill-Noether locus $\mathcal{BN} = M_C(r, 2rs, r+s)$ is slope-stable and $h^0(F) = r+s$. The moduli space $M_{X,H}(v)$ is a smooth projective variety of dimension 2. Hence we only need to show derivative of the restriction map $d\psi$ is surjective. The Zariski tangent space to the Brill-Noether locus \mathcal{BN} at the point $[F]$ is the kernel of the map

$$k_1: \text{Ext}^1(F, F) \rightarrow \text{Hom}(H^0(C, F), H^1(C, F)),$$

where any $f: F \rightarrow F[1] \in \text{Ext}^1(F, F) = \text{Hom}_C(F, F[1])$ goes to

$$k_1(f) = H^0(f): \text{Hom}_C(\mathcal{O}_C, F) \rightarrow \text{Hom}_C(\mathcal{O}_C, F[1]),$$

see [BS13, Proposition 4.3] for details. Note that the proof in [BS13] is valid for any family of simple sheaves on a variety. In addition, for any vector bundle E in the moduli space $M_{X,H}(v)$,

$$T_{[E]}(M_{X,H}(v)) = \text{Hom}_X(E, E[1]).$$

Let $i: C \hookrightarrow X$ be the closed embedding of the curve C into the surface X , then $Ri_*(-) = i_*(-)$ and for a vector bundle E on X , we have $Li^*(E) = i^*(E)$. The derivative of the restriction map

$$d\psi: T_{[E]}M_{X,H}(v) \rightarrow T_{[E|_C]}\mathcal{BN},$$

sends any $f: E \rightarrow E[1] \in \text{Hom}_X(E, E[1])$ to its restriction $i^*f: i^*E \rightarrow i^*E[1] \in \ker(k_1)$.

Define $h: \text{id}_{\mathcal{D}(X)} \rightarrow Ri_*Li^*$ as the natural transformation for the pair of adjoint functors $Li^* \dashv Ri_*$. Given a vector bundle E in the moduli space $M_{X,H}(v)$ and a morphism $\varphi \in \text{Hom}_X(E, E[1])$, we have the commutative diagram

$$(25) \quad \begin{array}{ccc} E & \xrightarrow{h_E} & i_*i^*E \\ \varphi \downarrow & & \downarrow i_*i^*\varphi \\ E[1] & \xrightarrow{h_{E[1]}} & i_*i^*E[1]. \end{array}$$

Therefore the following diagram is also commutative

$$\begin{array}{ccccc}
 \mathrm{Hom}_X(E, E[1]) & \xrightarrow{d\psi} & \mathrm{Hom}_C(i^*E, i^*E[1]) & \xrightarrow{k_1} & \mathrm{Hom}(H^0(C, i^*E), H^1(C, i^*E)). \\
 & \searrow k_2 := h_{E[1]} \circ (-) & \downarrow \sim i_*(-) \circ h_E =: k_3 & & \\
 & & \mathrm{Hom}_X(E, i_*i^*E[1]) & &
 \end{array}$$

Consider the following distinguished triangle

$$(26) \quad E[1] \xrightarrow{h_{E[1]}} i_*i^*E[1] \xrightarrow{g} E(-H)[2].$$

Given a morphism $\xi \in \ker(k_1) \subseteq \mathrm{Hom}_C(i^*E, i^*E[1])$, we first claim that the composition $g \circ i_*\xi \circ h_E = 0$ vanishes, so there exist morphisms ξ' and ξ'' such that the following diagram commutes.

$$(27) \quad \begin{array}{ccccc}
 E & \xrightarrow{h_E} & i_*i^*E & \longrightarrow & E(-H)[1] \\
 \downarrow \exists \xi' & & \downarrow i_*\xi & & \downarrow \exists \xi'' \\
 E[1] & \xrightarrow{h_{E[1]}} & i_*i^*E[1] & \xrightarrow{g} & E(-H)[2].
 \end{array}$$

Since $\xi \in \ker(k_1)$, the composition

$$\xi \circ i^*ev_E: i^*\mathcal{O}_X^{h^0(E)} \rightarrow i^*E \rightarrow i^*E[1] \in \mathrm{Hom}_C(i^*\mathcal{O}_X^{h^0(E)}, i^*E[1])$$

vanishes. Thus the adjunction $Li^* \dashv Ri_*$ gives $i_*(\xi \circ i^*ev_E) \circ h_{\mathcal{O}_X^{h^0(E)}} = 0$. We know $i_*i^*ev_E \circ h_{\mathcal{O}_X^{h^0(E)}} = h_E \circ ev_E$, hence the morphism

$$i_*\xi \circ (h_E \circ ev_E): \mathcal{O}_X^{h^0(E)} \rightarrow i_*i^*E \rightarrow i_*i^*E[1]$$

vanishes. Moreover, Propositions 4.1 implies that we have the short exact sequence

$$0 \rightarrow E' \rightarrow \mathcal{O}_X^{h^0(E)} \xrightarrow{ev_E} E \rightarrow 0$$

in $\mathrm{Coh}(X)$. Since $\mathrm{Hom}_X(E', E(-H)[1]) = 0$, applying the functor $\mathrm{Hom}_X(-, E(-H)[2])$ gives the exact sequence

$$0 \rightarrow \mathrm{Hom}_X(E, E(-H)[2]) \xrightarrow{\Phi} \mathrm{Hom}_X(\mathcal{O}_X^{h^0(E)}, E(-H)[2]) \rightarrow \mathrm{Hom}_X(E', E(-H)[2]) \rightarrow 0.$$

Since $\Phi(g \circ i_*\xi \circ h_E) = g \circ (i_*\xi \circ h_E \circ ev_E) = 0$, the injectivity of Φ implies $g \circ i_*\xi \circ h_E = 0$ as claimed. Therefore there exists morphism $\xi': E \rightarrow E[1]$ in the diagram (27) such that $i_*\xi \circ h_E = h_{E[1]} \circ \xi'$. The commutative diagram (25) gives $h_{E[1]} \circ \xi' = i_*i^*\xi' \circ h_E$. Therefore $i_*\xi \circ h_E = i_*i^*\xi' \circ h_E \in \mathrm{Hom}_X(E, i_*i^*E[1])$ and the isomorphism k_3 implies $\xi = i^*\xi'$ which shows $d\psi$ is surjective.

□

Proof of Theorem 1.3. Let (X, H) be a polarised K3 surface with $\text{Pic}(X) = \mathbb{Z} \cdot H$, and let C be any curve in the linear system $|H|$. The moduli space $N = M_{X,H}(v)$ is a smooth projective K3 surface. There exists a Brauer class $\alpha \in \text{Br}(N)$ and a universal $(1 \times \alpha)$ -twisted sheaf $\tilde{\mathcal{E}}$ on $X \times N$.

Theorem 1.2 implies that the moduli space N is isomorphic to the Brill-Noether locus \mathcal{BN} and the restriction of the universal twisted sheaf $\tilde{\mathcal{E}}|_{C \times \mathcal{BN}}$ is a universal $(1 \times \alpha)$ -twisted sheaf on $C \times \mathcal{BN}$, so $v' = v(\tilde{\mathcal{E}}|_{p \times \mathcal{BN}})$ for a point p on the curve C .

Let H' be a generic polarisation on N . Then the moduli space $M_{N,H'}^\alpha(v')$ of α -twisted semistable sheaves on N with respect to H' , is isomorphic to the original K3 surface X (see e.g. [Yos15, Theorem 2.7.1]). Therefore, $M_{\mathcal{BN},H'}^\alpha(v') \cong X$ which completes the proof of Theorem 1.3. \square

REFERENCES

- [ABS14] E. Arbarello, A. Bruno, and E. Sernesi. Mukai’s program for curves on a K3 surface. *Algebr. Geom.*, 1(5):532–557, 2014.
- [Bay18] A. Bayer. Wall-crossing implies Brill-Noether: applications of stability conditions on surfaces. In *Algebraic geometry: Salt Lake City 2015*, volume 97 of *Proc. Sympos. Pure Math.*, pages 3–27. Amer. Math. Soc., Providence, RI, 2018.
- [Bri07] T. Bridgeland. Stability conditions on triangulated categories. *Ann. of Math. (2)*, 166(2):317–345, 2007.
- [Bri08] T. Bridgeland. Stability conditions on K3 surfaces. *Duke Math. J.*, 141(2):241–291, 2008.
- [BS13] U. N. Bhosle and S. K. Singh. Brill-Noether loci and generated torsion free sheaves over nodal and cuspidal curves. *Manuscripta Math.*, 141(1-2):241–271, 2013.
- [CLM93] C. Ciliberto, A. Lopez, and R. Miranda. Projective degenerations of K3 surfaces, Gaussian maps, and Fano threefolds. *Invent. Math.*, 114(3):641–667, 1993.
- [Fey19] S. Feyzbakhsh. Mukai’s program (reconstructing a K3 surface from a curve) via wall-crossing, arXiv:1710.06692, to appear in J. Reine Angew. Math. DOI: 10.1515/crelle-2019-0025, 2019.
- [Fey20] S. Feyzbakhsh. Mukai’s program (reconstructing a K3 surface from a curve) via wall-crossing, II, arXiv:2006.08410, 2020.
- [GL87] M. Green and R. Lazarsfeld. Special divisors on curves on a K3 surface. *Invent. Math.*, 89(2):357–370, 1987.
- [HL10] D. Huybrechts and M. Lehn. *The geometry of moduli spaces of sheaves*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010.
- [HS05] D. Huybrechts and P. Stellari. Equivalences of twisted K3 surfaces. *Math. Ann.*, 332(4):901–936, 2005.
- [Huy16] D. Huybrechts. *Lectures on K3 surfaces*, volume 158 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2016.
- [Mac14] A. Maciocia. Computing the walls associated to Bridgeland stability conditions on projective surfaces. *Asian J. Math.*, 18(2):263–279, 2014.
- [MS17] E. Macri and B. Schmidt. Lectures on Bridgeland stability. In *Moduli of curves*, volume 21 of *Lect. Notes Unione Mat. Ital.*, pages 139–211. Springer, Cham, 2017.
- [Muk87] S. Mukai. On the moduli space of bundles on K3 surfaces. I. In *Vector bundles on algebraic varieties (Bombay, 1984)*, volume 11 of *Tata Inst. Fund. Res. Stud. Math.*, pages 341–413. Tata Inst. Fund. Res., Bombay, 1987.
- [Muk88] S. Mukai. Curves, K3 surfaces and Fano 3-folds of genus ≤ 10 . In *Algebraic geometry and commutative algebra, Vol. I*, pages 357–377. Kinokuniya, Tokyo, 1988.

- [Muk96] S. Mukai. Curves and $K3$ surfaces of genus eleven. In *Moduli of vector bundles (Sanda, 1994; Kyoto, 1994)*, volume 179 of *Lecture Notes in Pure and Appl. Math.*, pages 189–197. Dekker, New York, 1996.
- [Muk01] S. Mukai. Non-abelian Brill-Noether theory and Fano 3-folds [translation of Sugaku **49** (1997), no. 1, 1–24; MR1478148 (99b:14012)]. *Sugaku Expositions*, 14(2):125–153, 2001. Sugaku Expositions.
- [Yos15] K. Yoshioka. Perverse coherent sheaves and Fourier-Mukai transforms on surfaces, II. *Kyoto J. Math.*, 55(2):365–459, 2015.

SCHOOL OF MATHEMATICS, IMPERIAL COLLEGE LONDON, HUXLEY BUILDING, SOUTH KENSINGTON
CAMPUS, SW7 2AZ, LONDON, UNITED KINGDOM
E-mail address: `s.feyzbakhsh@imperial.ac.uk`