

On vertex coloring without monochromatic triangles

Michał Karpiński*, Krzysztof Piecuch†

Institute of Computer Science
University of Wrocław
Joliot-Curie 15, 50-383 Wrocław, Poland

February 9, 2022

Abstract

We study a certain relaxation of the classic vertex coloring problem, namely, a coloring of vertices of undirected, simple graphs, such that there are no monochromatic triangles. We give the first classification of the problem in terms of classic and parametrized algorithms. Several computational complexity results are also presented, which improve on the previous results found in the literature. We propose the new structural parameter for undirected, simple graphs – the triangle-free chromatic number χ_3 . We bound χ_3 by other known structural parameters. We also present two classes of graphs with interesting coloring properties, that play pivotal role in proving useful observation about our problem. We give/ask several conjectures/questions throughout this paper to encourage new research in the area of graph coloring.

1 Introduction

Graph coloring is probably the most popular subject in graph theory. It is an interesting topic from both algorithmic and combinatoric points of view. The coloring problems have many practical applications in areas such as operations research, scheduling and computational biology. For a recent survey one can turn to [9]. In this paper we study a variation of the classic coloring – we call it the *triangle-free coloring* problem, in which we ask for an assignment of colors to the vertices of a given graph, such that the number of colors used is minimum and that each cycle of length 3 has at least two vertices colored differently. We show that our problem has interesting graph-theoretical properties and we also present some evidence that this problem might be easier than the classic vertex coloring. This suggests that studying our variation, new results can be achieved in the field of classic vertex coloring,

*karp@cs.uni.wroc.pl

†kpiecuch@cs.uni.wroc.pl

which is known to be one of the hardest known optimization problems. Apart from theoretical motivation, there is also a practical one – vertex coloring without monochromatic cycles can be used in the study of consumption behavior [7].

1.1 Related work

Some researchers have already considered coloring problems that are similar to our variation. The class of planar graphs has been of particular interest, for example, Angelini and Frati [2] study planar graphs that admit an acyclic 3-coloring – a proper coloring in which every 2-chromatic subgraph is acyclic. Algorithms for acyclic coloring can be used to solve/approximate a triangle-free coloring, although we do not explore this possibility in this paper. Another result is of Kaiser and Škrekovski [11], where they prove that every planar graph has a 2-coloring such that no cycle of length 3 or 4 is monochromatic. Thomassen [21], on the other hand, considers list-coloring of planar graphs without monochromatic triangles. Few hardness results for our problem are known – Karpiński [12] showed that verifying whether a graph admits a 2-coloring without monochromatic cycles of fixed length is \mathcal{NP} -complete. His proof was then simplified by Shitov [19], who also proposed and proved the hardness of an extension of our problem, where additional restriction is imposed on the coloring in the form of the set of *polar* edges – edges that must not be monochromatic in the resulting coloring.

1.2 Our contribution

Several novel results are presented in this paper. First, we explore the graph-theoretical side of our problem. We propose the new structural parameter $\chi_3(G)$ which is the minimum number of colors needed to label the vertices of an undirected, simple graph G , such that there are no monochromatic triangles. We then bound this new parameter by $\omega(G)$, $\chi(G)$ and $\Delta(G)$, which are clique number, chromatic number and the largest vertex degree of G , respectively. We conjecture that $\chi_3(G) \leq \lceil \omega(G)/2 \rceil + 1$ and we construct an infinite class of graphs for which this upper bound is met. In our construction we use what we call *cycle-cliques* as building blocks. Those gadgets have interesting coloring properties and are also used in proving hardness results later in the paper.

For the positive side, several known graph classes are presented, for which our problem can be solved efficiently, for example, we can find χ_3 on planar graphs in polynomial time, whereas finding χ is \mathcal{NP} -complete, even on planar graphs with maximum degree 4 [6]. We use the fact that when χ is small (less than 5), then we can reduce our problem to the problem of deciding if a given graph is triangle-free. In general, the time needed for listing all triangles of a graph is $O(m\sqrt{m})$ [1], but in the presented graphs, we can find if a graph is triangle-free in time $O(n)$. We also prove that our problem is fixed-parameter tractable, when the parameter is the vertex cover number.

We present several hardness results, which improve on the work of Karpiński [12] and Shitov [19]. We show that given any fixed number $q \geq 2$, determining if graph is triangle-free q -colorable is \mathcal{NP} -hard. This improves on the result given in [12], where the author shows

hardness only for $q = 2$. Another improvement to Karpiński’s result [12] is \mathcal{NP} -hardness proof of triangle-free 2-coloring problem for the graphs that does not contain clique of size 4 as a subgraph. In [19], author formulates and proves the \mathcal{NP} -hardness of triangle-free 2-coloring problem where additional set of polar edges – edges that must not be monochromatic – are given on input. We show that this variation of our problem remains \mathcal{NP} -hard, even on graphs with maximum degree 3 – the sub-cubic graphs.

Throughout the paper we ask many questions and propose conjectures, which we hope will spark an interest in this new variant of vertex coloring.

1.3 Structure of the paper

In Section 2 we give definitions and notations used throughout the paper, as well as formulation of all considered problems. We give several bounds on χ_3 in Section 3 and we construct an infinite class of graphs for which one of the conjectured bounds is tight. In Section 4 we present efficient algorithms for the triangle-free coloring problem for certain classes of graphs. A single FPT result are also shown there. Hardness results are presented in Section 5. We finish the paper with some concluding remarks in Section 6, where we also present the reader several open problems.

2 Preliminaries

We recall several properties and notions from graph theory that are of interest to us. Let $G = (V, E)$ be a finite, undirected, unweighted, simple graph with the vertex set V and the edge set E . Set of vertices adjacent to some vertex $v \in V$ is called the neighborhood of v and is denoted by $N_G(v)$. The degree of a vertex v is defined as $d_G(v) = |N_G(v)|$ (we omit index G in the notation if it’s clear from the context which graph is considered). The degree of G is $\Delta(G) = \max_{v \in V} d(v)$. We write that H is a subgraph of G as $H \subseteq G$, and H is isomorphic to G as $H \sim G$. We say that graph G is H -free if there is no $H' \subseteq G$ such that $H' \sim H$.

An elementary contraction of graph G is obtained by *identification* of (ordered) pair of vertices $\langle u, v \rangle$. This process is performed by removing v and adding an edge uw , for every $w \in N(v)$. After that, multi-edges and loops are removed.

Let K_n ($n \in \mathbb{N}$) be the clique of size n . The greatest integer r such that $K_r \subseteq G$ is the *clique number* $\omega(G)$ of G . A subset I of vertices V is an independent set, if none of its vertices are adjacent. A subset W of vertices V is called a vertex cover, if every edge in the graph is incident to at least one vertex of W . The smallest integer r such that $|W| = r$ and W is a vertex cover of G , is called the *vertex cover number*.

In this paper we study coloring of vertices, and to avoid confusion, we distinguish two types of coloring. A *classic k -coloring* of a graph is a function $c : V \rightarrow \{1, \dots, k\}$, such that there are no two adjacent vertices u and v , for which $c(u) = c(v)$. Given G , the smallest k for which there exists a classic k -coloring for G is called the *chromatic number* and is denoted as $\chi(G)$. A *triangle-free k -coloring* of a graph is a function $c : V \rightarrow \{1, \dots, k\}$, such that there are no three mutually adjacent vertices u, v and w , for which $c(u) = c(v) = c(w)$. If such

vertices exist, then the induced subgraph (K_3) is called a *monochromatic triangle*. Given G , the smallest k for which there exists a triangle-free k -coloring for G we call the *triangle-free chromatic number* and we denote it as $\chi_3(G)$.

We now formulate the decision problems investigated in this paper, for any fixed integer $q > 0$:

TRIANGLEFREE-q-COLORING

Input: A finite, undirected, simple graph G .

Question: Is there a triangle-free q -coloring of G ?

TRIANGLEFREEPOLAR-q-COLORING

Input: A finite, undirected, simple graph $G = (V, E)$ and a subset $S \subseteq E$.

Question: Is there a triangle-free q -coloring of G , such that no edge in S is monochromatic?

Fixed-parameter tractable (FPT) algorithm (w.r.t. parameter k) is an algorithm for whose running time is polynomial for any fixed value of k , i.e., running time is $O(f(k) \cdot n^c)$, where n is a size of the input, c is a constant and f is a computable function.

We denote $\lceil x \rceil$ to be the smallest integer not less than x . Let $r \in \mathbb{N}$, we define a binary operator $+_r$ on the set $\mathbb{Z}_r = \{0, 1, 2, \dots, r-1\}$, which is addition modulo r , i.e., for every $n, m \in \mathbb{Z}_r$, $n +_r m = n + m \pmod{r}$. The operator $-_r$ can be defined in a similar way.

3 Bounds on the triangle-free chromatic number

We first give simple bounds on $\chi_3(G)$ in terms of $\omega(G)$ and $\chi(G)$:

Theorem 1. *For any graph G : $\lceil \frac{\omega(G)}{2} \rceil \leq \chi_3(G) \leq \lceil \frac{\chi(G)}{2} \rceil$.*

Proof. To see that the lower bound holds, take any clique in G of maximum cardinality. We can use one color for at most two vertices of that clique, otherwise we would create a monochromatic triangle. Therefore we need at least $\lceil \omega(G)/2 \rceil$ colors in order to make the coloring triangle-free in this clique, and therefore at least $\lceil \omega(G)/2 \rceil$ colors are needed to triangle-free color the entire graph.

The upper bound can be justified by the following argument. Let $k = \chi(G)$ and take any classic k -coloring of G . Let V_i be the set of vertices colored i , where $1 \leq i \leq k$. For each $0 \leq j \leq \lceil k/2 \rceil - 1$ recolor sets $V_{2j+1} \cup V_{2j+2}$ with j (we may need to add empty set V_{k+1} , if k is odd). Since every V_i is an independent set, then after recoloring, any monochromatic cycle is of even length. Therefore the resulting coloring is triangle-free. \square

Remark 1. *The recoloring procedure from the above proof we call the standard recoloring strategy. It will be used in the construction of algorithms in the next section.*

The fundamental question we ask is how tight are the bounds in Theorem 1 and what kind of algorithmic consequences are implied by these observations. The following theorem shows that the upper bound can be arbitrarily large, which means that we should not expect to find an algorithm for general graphs that would acceptably approximate $\chi_3(G)$ based on a chromatic number alone.

Theorem 2. For any $k \geq 1$, there exists a graph G for which $\chi_3(G) = 1$ and $\chi(G) = k$.

Proof. The class of graphs called Mycielski graphs [14] meet this property, as they are triangle-free and can have arbitrarily large chromatic number. \square

On the other hand, we conjecture that the lower bound of Theorem 1 is almost tight.

Conjecture 1. For any graph G : $\chi_3(G) \leq \left\lceil \frac{\omega(G)}{2} \right\rceil + 1$.

To show that it is possible to reach the conjectured bound we construct an infinite class of graphs for which $\chi_3(G) = \lceil \omega(G)/2 \rceil + 1$. First, we present an auxiliary class of graphs that we call *cycle-cliques*, which will serve as building blocks of our construction.

Definition 1 (cycle-clique). Let $k \in \mathbb{N}$. The k -cycle-clique is a graph that have exactly $5 \cdot k$ vertices $v_{i,j}$ for $i \in \{0, \dots, 4\}$ and $j \in \{0, \dots, k-1\}$ and there is an edge between $v_{i,j}$ and $v_{i',j'}$ if and only if $|i - i'| \leq 1$. We will define a set $J_i = \{v_{i,j} : j \in \{0, \dots, k-1\}\}$. Each J_i is called a joint.

Observation 1. For any k -cycle-clique it holds that $J_i \sim K_k$ and $J_i \cup J_{i+5} \sim K_{2k}$, for $i \in \{0, \dots, 4\}$.

Example 1. In Figures 1b and 1c we present k -cycle-cliques for $k \in \{1, 2\}$. In both graphs joints are marked with dashed lines. The 1-cycle-clique is simply a cycle of length 5. Notice how $J_i \cup J_{i+5}$ forms a clique of size $2k$, for $i \in \{0, \dots, 4\}$, whereas each J_i is a clique of size k . We also note that the graph in Figure 1c is the same as the gadget presented in [12], where the author uses it to encode what he calls super-edges, in his \mathcal{NP} -completeness proof of the TRIANGLEFREE-2-COLORING problem. The cycle-clique is a generalization of this gadget.

Observation 2. Let $k \geq 1$. In every triangle-free k -coloring of graph K_{2k} every color is used exactly twice.

Lemma 3. Let $k \geq 1$. In every triangle-free k -coloring of k -cycle-clique each vertex of J_i has a unique color (w.r.t. other vertices of J_i), for $i \in \{0, \dots, 4\}$.

Proof. Proof by contradiction. Let's say that there is a triangle-free k -coloring of k -cycle-clique and $i \in \{0, \dots, 4\}$ such that there exist two vertices in J_i that have the same color. Using Observation 2 we know that this color cannot appear in J_{i+5} or J_{i-5} . Using Observation 2 again, this color need to be used twice in each of J_{i+5} and J_{i-5} . Since $J_{i+5} \cup J_{i-5} \sim K_{2k}$ from the definition and that four vertices of $J_{i+5} \cup J_{i-5}$ are colored with the same color, we reach a contradiction with the assumption that the initial coloring was triangle-free. \square

Now we present the contraction scheme of cliques, which is a procedure used in our construction.

Clique contraction scheme: Let $U = \langle u_0, \dots, u_{k-1} \rangle$, $V = \langle v_0, \dots, v_{k-1} \rangle$, $W = \langle w_0, \dots, w_{k-1} \rangle$ be the cliques of size k . Perform vertex identification on the following pairs of vertices (in that order):




Figure 1: Examples of new classes of graphs.

1. (v_i, u_i) for $i \in \{0, \dots, k-2\}$,
2. (v_i, w_i) for $i \in \{0, \dots, k-3\}$,
3. (v_{k-1}, w_{k-2}) and (u_{k-1}, w_{k-1}) .

Example 2. In Figure 2 we present a contraction of three cliques: $U = \langle u_0, u_1, u_2 \rangle$, $V = \langle v_0, v_1, v_2 \rangle$ and $W = \langle w_0, w_1, w_2 \rangle$, using the above procedure. Figure 2a show the initial state. Dashed lines connect vertices that will be identified in the first step of the scheme, namely, $\langle v_0, u_0 \rangle$ and $\langle v_1, u_1 \rangle$. The result of the first step is presented in Figure 2b. Next, vertices $\langle v_0, w_0 \rangle$ are identified (Figure 2c) and lastly, we identify vertices $\langle v_2, w_1 \rangle$ and $\langle u_2, w_2 \rangle$. The final result is presented in Figure 2d. Notice that the clique contraction scheme, given three cliques of size k , outputs one clique of size $k+1$.

We are ready to present the desired construction.

Definition 2 (k -clover graph). Let $G_1 = (\cup_{i=0}^4 V_i, E_1)$, $G_2 = (\cup_{i=0}^4 U_i, E_2)$ and $G_3 = (\cup_{i=0}^4 W_i, E_3)$ be k -cycle-cliques, where $k \geq 2$. A graph created by running the clique contraction scheme on sets V_0 , U_0 and W_0 is called a k -clover graph.

Example 3. The example of k -clover graph is presented in Figure 1a, for $k = 2$. Notice how contracting three joints of size 2 results in a creation of a triangle (K_3) in the middle.

Lemma 4. There is no k -clover graph G such that $K_{2k+1} \subseteq G$, for $k \geq 2$.




Figure 2: Sample run of the clique contraction scheme.

Proof. Only vertices that can have degree at least $2k$ are those that participated in vertex identification. In graph G there is exactly $k + 1$ such vertices. \square

Lemma 5. *There is no triangle-free k -coloring of k -clover graph.*

Proof. Proof by contradiction. By Lemma 3 we know that in each set $V_0 = \langle v_0, \dots, v_{k-1} \rangle$, $U_0 = \langle u_0, \dots, u_{k-1} \rangle$ and $W_0 = \langle w_0, \dots, w_{k-1} \rangle$ vertices need to have different colors. Because we identify vertices (v_i, u_i) for $i \in \{0, \dots, k-2\}$, it means that v_{k-1} and u_{k-1} need to have the same color. But because we identify (v_{k-1}, w_{k-2}) and (u_{k-1}, w_{k-1}) that mean w_{k-2} and w_{k-1} need to have the same color, which contradicts Lemma 3. \square

Theorem 6. *For any $k \geq 2$, there exists a graph G , such that $\chi_3(G) = \lceil \omega(G)/2 \rceil + 1$.*

Proof. Let G be a k -clover graph. From the definition of a cycle-clique, $\omega(G) \geq 2k$ and from Lemma 4 and Theorem 1, $\omega(G) \leq 2k$, therefore $\omega(G) = 2k$. We have $\chi_3(G) \geq k + 1$ by Lemma 5. We show that there exist a triangle-free coloring of graph G that uses $k + 1$ colors. Let U, V, W be the cliques used in the contraction scheme, when constructing G . The result of the contraction is a clique of size $k + 1$. Assign a unique color to each of its vertices. Notice that U, V and W are cliques of size k where each vertex has a unique color. By Lemma 3 we can extend the coloring of U, V and W to their respective k -cycle-cliques. This proves that $\chi_3(G) = k + 1 = \frac{2k}{2} + 1 = \lceil \omega(G)/2 \rceil + 1$. \square

Remark 2. *Notice that we have used 5 joints in the construction of a single k -cycle-clique. We comment on why constant 5 is important here. Notice that if we have used even number of joints, then the property of Lemma 3 would not hold. Also, if we were to use 3 joints, then the k -cycle-clique would be isomorphic to K_{3k} , and therefore would not be triangle-free k -colorable – we leave the verification of those facts to the reader, as an easy exercise. In conclusion, 5 is the smallest possible number of joints in the k -cycle-clique for it to hold the desired properties. But in fact, any odd number of joints greater or equal to 5 can be used.*

Another issue worth discussing is that the proof of Theorem 6 shows the existence of graphs for which $\chi_3(G) = \lceil \omega(G)/2 \rceil + 1$, but only when $\omega(G)$ is even. We now show that there exists a construction for $r = 3$, and we leave the existence of such graphs for other values of r as an open problem.




Figure 3: The K_4 -free polar gadget. It consists of 12 vertices, 30 edges and 20 triangles.

Definition 3. *The K_4 -free polar gadget is the graph $G = (V, E)$ with $V = \{u, v, w_1, w_2, w_3, w_4, z_1, z_2, z_3, y_1, y_2, y_3\}$, and edge set defined as in Figure 3.*

We say that edge xy is *polar*, if in every triangle-free 2-coloring c , $c(x) \neq c(y)$.

Lemma 7. *Let G be a K_4 -free polar gadget. Then the following are true: (i) G is triangle-free 2-colorable, (ii) uv is a polar edge, (iii) G is K_4 -free.*

Proof. *Ad. (i):* Let $S = \{u, z_1, y_1\}$. Set $c(S) = 1$ and $c(V \setminus S) = 2$.

Ad. (ii): Assume by contradiction, that $c(u) = c(v) = 1$ and G is triangle-free 2-colorable. Then $c(\{w_1, w_2, w_3, w_4\}) = 2$, which is forced. Since $c(v) = 1$, then either $c(z_1) = 2$, or $c(y_1) = 2$. Assume the former, without loss of generality. Then $c(z_2) = 1$, otherwise $\{w_1, z_1, z_2\}$ would be monochromatic. But this means that we cannot color z_3 , because if $c(z_3) = 1$, then $\{u, z_2, z_3\}$ is monochromatic, and if $c(z_3) = 2$, then $\{w_2, z_1, z_3\}$ is monochromatic – a contradiction.

Ad. (iii): Let $V_1 = \{u, v\}$, $V_2 = \{w_1, w_2, w_3, w_4\}$, $V_3 = \{z_1, z_2, z_3, y_1, y_2, y_3\}$. If there exists $H \subset G$ that is isomorphic to K_4 , then we can observe that:

- H contains at most one vertex from V_2 , since V_2 is an independent set, and
- H contains at most one vertex from V_1 , because otherwise H would contain at least one vertex from V_3 , and there is no vertex (or pair of vertices) in V_3 that is connected to both vertices in V_1 .

Therefore we are left with the following four cases:

- H uses only vertices from V_3 . But in $G(V_3)$ only z_1 and y_1 have degree at most 3.

- H uses 3 vertices from V_3 and one vertex from V_2 . But each vertex from V_2 is adjacent to exactly two vertices from V_3 .
- H uses 3 vertices from V_3 and one vertex from V_1 . Then $v \notin H$ because it is adjacent to only z_1 and y_1 . Thus $u \in H$, but it is easy to verify that $N(u)$ does not contain a triangle.
- H uses 2 vertices from V_3 , one vertex from V_2 and one vertex from V_1 . But for each $\{s, t\} \subset V_3$, there exists exactly one triangle $\{s, t, r\}$, such that $r \in V_1 \cup V_2$.

We have reached contradictions in all possible cases, thus we conclude that G is K_4 -free. \square

Theorem 8. *There exists a graph G , such that $\chi_3(G) = \lceil \omega(G)/2 \rceil + 1$, and $\omega(G) = 3$.*

Proof. To construct G , take three K_4 -free polar gadgets and identify their polar edges so that they form a triangle. Properties (i) and (ii) of Lemma 7 forces the use of the third color in G and Property (iii) makes the construction K_4 -free. \square

We believe that the gadgets that we presented in this section are interesting from the graph-theoretic point of view. It would be instructive to find answers to the following questions.

Question 1. *For any $k \geq 2$, what is the smallest graph (in terms of vertices and/or edges) for which the following are true: (i) $\chi_3(G) = k$, and (ii) for any triangle-free k -coloring of G , there exists a polar clique of size k in G , i.e., a clique of size k where each vertex has a unique color?*

Question 2. *Is K_4 -free polar gadget the smallest is terms of vertices/edges/triangles graph that satisfy the properties of Lemma 7?*

To finish this section we give another bound on $\chi_3(G)$. We recall the well-known Brook's Theorem [4] which states that for any graph G , we have $\chi(G) \leq \Delta(G)$, unless G is a complete graph or an odd cycle. Using this theorem we can prove the following.

Theorem 9. *Let $G = (V, E)$ be any graph where $|V| > 3$, where G is not a complete graph of odd number of vertices. Then $\chi_3(G) \leq \left\lceil \frac{\Delta(G)}{2} \right\rceil$.*

Proof. If G is an odd cycle of length at least 5, then $\chi_3(G) = 1$ and $\left\lceil \frac{\Delta}{2} \right\rceil = 1$, so the inequality holds. If G is a complete graph (a clique) of n vertices, then $\chi_3(G) = \left\lceil \frac{n}{2} \right\rceil$ and $\Delta = n - 1$. The inequality $\left\lceil \frac{n}{2} \right\rceil \leq \left\lceil \frac{n-1}{2} \right\rceil$ holds iff n is even. If the above cases does not occur, then the application of the Brook's Theorem combined with the upper bound of Theorem 1 completes the proof. \square

4 Tractable classes of graphs

Here we present several classes of graphs for which efficient algorithms for triangle-free coloring problem exist. We use the bounds derived in the previous section to show that for small values of structural parameters polynomial time complexity can be achieved in many cases.

Given graph $G = (V, E)$, we distinguish between two optimization variants of the triangle-free coloring problem: (i) finding the number $\chi_3(G)$, and (ii) finding the triangle-free coloring that uses exactly $\chi_3(G)$ colors, i.e., the mapping from V to $\{1, \dots, \chi_3(G)\}$ is the required output. As we will soon see, complexities for these two variants can be different. To this end we propose the following notion of time complexity for our problem. For convenience, we say that n is the number of vertices of graph G and m is the number of edges of G , unless stated otherwise.

Definition 4. *Let \mathcal{G} be a class of graphs. We say that the triangle-free coloring problem is solvable in time $O(f(n+m), g(n+m))$ on \mathcal{G} , iff there exist algorithms \mathcal{A} and \mathcal{B} , such that for every input $G \in \mathcal{G}$, (i) algorithm \mathcal{A} outputs $\chi_3(G)$ in $O(f(n+m))$ time, and (ii) algorithm \mathcal{B} outputs the triangle-free coloring that uses exactly $\chi_3(G)$ colors, in $O(g(n+m))$ time.*

4.1 Graphs with bounded chromatic number

Here we present a single theorem that presents the complexity of triangle-free coloring problem on several popular classes of graphs. We exploit the fact that if we know in advance that $\chi(G) \leq 4$ – and therefore $\chi_3(G) \leq 2$, by Theorem 1 – then $\chi_3(G)$ can be found easily and is one of the following: (a) $\chi_3(G) = 0$ iff G is empty, (b) $\chi_3(G) = 1$ iff G is triangle-free (K_3 -free), (c) $\chi_3(G) = 2$ iff the above two cases does not hold. Checking the first case is trivial. Therefore the challenge of finding $\chi_3(G)$ lies in checking whether G contains a triangle.

For finding the actual coloring, we use the fact that in the given graphs we can find “good-enough” classic coloring, and then recolor vertices so that there are no monochromatic triangles using the *standard recoloring strategy*.

Theorem 10. *The triangle-free coloring problem is solvable:*

- in time $O(n, n^2)$ on planar graphs,
- in time $O(n, n)$ on: outerplanar graphs, chordal graphs, Δ -regular graphs, with $\Delta \leq 4$.

We first give the required definitions, then, the proof.

We say that the graph is *planar*, if we can embed it in the plane, i.e., it can be drawn on the plane in such a way that its edges intersect only at their endpoints. Graph G is *outerplanar* if it is planar and there exists a plane embedding of G in which all vertices lie on the exterior (unbounded) face. Graph is *chordal* if every cycle of length greater than 3 has a chord – an edge connecting two non-adjacent vertices on a cycle. Graph is *Δ -regular*, if its every vertex has degree Δ .

Proof of Theorem 10. We begin with chordal graphs. Chordal graphs are the part of the larger class called perfect graphs. A perfect graph G has a property $\chi(G) = \omega(G)$, therefore $\chi_3(G) = \left\lceil \frac{\chi(G)}{2} \right\rceil = \left\lceil \frac{\omega(G)}{2} \right\rceil$, by Theorem 1. This means that determining $\chi_3(G)$ is equivalent to determining either $\omega(G)$ or $\chi(G)$. Fortunately, this can be done in linear-time in chordal graphs [10]. Moreover, not only can we find $\chi(G)$ in linear-time, but also the (classic) coloring that uses $\chi(G)$ colors. Using standard recoloring strategy, we can produce a triangle free coloring that uses $\left\lceil \frac{\chi(G)}{2} \right\rceil = \chi_3(G)$ colors.

We now turn our attention to other classes of graphs. First, we show that we can find whether given graph G is triangle-free in $O(n)$ time, for the following cases:

- *planar graphs* - using linear-time algorithm by Papadimitriou and Yannakakis [15],
- *outerplanar graphs* - same as above,
- Δ -*regular graphs* - we loop through all the vertices, and for every vertex v , check if any pair of vertices from $N(v)$ are adjacent to each other. This takes at most $O(n \cdot \Delta^2) = O(n)$ time.

Hence, we can check if $\chi_3(G) > 1$ in linear-time (checking if $\chi_3(G) = 0$ is trivial). Furthermore, from Theorems 1 and 9 we know that $\chi_3(G) \leq 2$ in all three cases, therefore determining $\chi_3(G)$ can be done in time $O(n)$.

From now on we assume that the input graph contains at least one triangle. Otherwise, we can output a trivial coloring $c(V) = 1$.

Finding the actual coloring of G has different time complexity depending on the class of G . First, we find a “good-enough” classic coloring of G :

- *planar graphs* - The well-known Four Color Theorem states that every planar graph is classically 4-colorable, that is, for each planar G , $\chi(G) \leq 4$. The theorem was proved by Appel and Haken in [3], where the authors propose a quadratic-time algorithm based on their proof. The algorithm was later improved by Robertson et al. [17], but the asymptotic running time is still $O(n^2)$.
- *outerplanar graphs* - Proskurowski and Sysłó [16] give an algorithm for finding $\chi(G)$ on outerplanar graphs which runs in $O(n)$ time.
- Δ -*regular graphs (where $\Delta \leq 4$)* - Skulrattanakulchai [20] shows how to find classic Δ -coloring in $O(n)$ time.

Given the classic coloring of G , we use the standard recoloring strategy to obtain the final coloring. Since we know that $\chi(G) \leq 4$ and $\chi(G) \geq 3$ (by the assumption that it contains at least one triangle), then $\chi_3(G) = 2$ and the resulting coloring is optimal. \square

We do not know if there exists a sub-quadratic algorithm for finding a triangle-free coloring in planar graphs, so we leave it as an open problem. It is worth noting that the situation would not improve even if we knew that the the input graph is classically 3-colorable, as the best know algorithm for classically 4-coloring planar graphs with this property still runs in $O(n^2)$ time [13].

Question 3. *Is there an algorithm that finds a triangle-free coloring of a planar graph in time $o(n^2)$?*

Remark 3. *Theorem 10 serves as an evidence, that the triangle-free coloring problem is easier than the classic coloring problem. We have efficient algorithms for finding χ_3 on planar graphs and graphs with $\Delta = 4$, but it is \mathcal{NP} -hard to find the χ even on 4-regular planar graphs [6]. This suggests that the above question might have a positive answer.*

4.2 FPT result

In this subsection we look at our problem from the parametrized complexity point of view. The main goal here is not to give efficient algorithms, but rather provide the initial classification of the problem in the context of \mathcal{FPT} theory. We prove that TRIANGLEFREE- q -COLORING problem is fixed-parameter tractable when parametrized by vertex cover number.

In the article by Fiala, Golovach and Kratochvíl [8], authors explore differences in the complexity of several coloring problems when parametrized by either vertex cover number or treewidth. We use techniques similar to theirs in the proof of the following theorem.

Theorem 11. *The TRIANGLEFREE- q -COLORING problem is \mathcal{FPT} when parametrized by the vertex cover number.*

Proof. Let W be a minimum vertex cover of G , and let $|W| = k$. Then $I = V \setminus W$ is an independent set. The goal is to find a triangle-free coloring $c : V \rightarrow \{1, \dots, q\}$.

The algorithm works in two steps: first, we find the triangle-free q -coloring of W by exhaustive search. Then, since I is an independent set, we can use greedy strategy to color its vertices, once we know the coloring of W . By greedy strategy we mean taking each $v \in I$, checking its colored neighborhood $N(v)$ (notice that $N(v) \subseteq W$), and coloring v in a way which does not create a monochromatic triangle in G . Clearly this procedure will find a triangle-free q -coloring iff such coloring exists. The analysis of running time now follows.

Since $\chi_3(W) \leq \lceil \chi(W)/2 \rceil \leq \lceil k/2 \rceil$, it is natural to consider the following two cases:

(i) If $q \leq \lceil k/2 \rceil$, then we consider all q -colorings of W and their extensions to I by the greedy algorithm. Since W has at most $k^{\lceil k/2 \rceil}$ colorings, the running time is $O(k^{\lceil k/2 \rceil + 1}n)$.

(ii) If $q > \lceil k/2 \rceil$, then assuming that $W = \{w_0, \dots, w_{k-1}\}$, we can set $c(w_{2i}) = c(w_{2i+1}) = i + 1$, for each $0 \leq i < \lceil k/2 \rceil$, which clearly uses $\lceil k/2 \rceil$ colors, then set $c(I) = \lceil k/2 \rceil + 1$. Thus, in this case, the triangle-free q -coloring always exists, and we can find it in $O(n)$ time.

We conclude that the total running time of the algorithm is $O(k^{\lceil k/2 \rceil + 1}n)$, which proves the claim. \square

5 Hardness results

Here we improve the results of Karpiński [12] and Shitov [19]. In their papers they show that the generalized version of TRIANGLEFREE-2-COLORING problem is \mathcal{NP} -complete – they not only consider monochromatic triangles, but monochromatic cycles of arbitrary (fixed) length. Karpiński first presented the proof, which was then simplified by Shitov, by introducing an auxiliary problem, which we call the TRIANGLEFREEPOLAR-2-COLORING problem. In Shitov’s proof, this new problem serves as a connection between NOTALLEQUAL 3-SAT and TRIANGLEFREE-2-COLORING.

Our contribution is as follows: we first prove that the triangle-free coloring remains \mathcal{NP} -hard even if we can use any fixed number of colors. Then we show that the TRIANGLEFREE-2-COLORING problem is \mathcal{NP} -hard in the restricted class of graphs, which do not contain K_4 as a subgraph. These improve the results of Karpiński [12]. Lastly, we show that the TRIANGLEFREEPOLAR-2-COLORING remains \mathcal{NP} -hard even in graphs of maximum degree 3, which improves the result of Shitov [19].

Theorem 12. *For any fixed $q \geq 2$, the TRIANGLEFREE- q -COLORING problem is \mathcal{NP} -hard.*

Proof. We prove the theorem by induction on q . For $q = 2$ the hardness has already been proved by Karpiński [12]. What remains is to show the polynomial reduction from the TRIANGLEFREE- q -COLORING problem to TRIANGLEFREE- $(q+1)$ -COLORING problem, for any fixed $q \geq 2$. Take any graph $G = (V, E)$, where $|V| = n$ and $|E| = m$, and assume that the vertices are arbitrarily ordered, i.e., $V = \{v_1, \dots, v_n\}$. We construct the graph $G' = (V', E')$ in the following way:

1. Let $V' = V \cup (\bigcup_{i=1}^n V_i)$ and $E' = E \cup (\bigcup_{i=1}^n E_i)$, where $V_i = V_i^0 \cup V_i^1 \cup V_i^2 \cup V_i^3 \cup V_i^4$ and (V_i, E_i) is a $(q + 1)$ -cycle-clique with the set $\{V_i^0, V_i^1, V_i^2, V_i^3, V_i^4\}$ being its joints.
2. For every $1 \leq i \leq n$, pick arbitrary vertex from V_i^0 and call it u_i . Identify the pairs of vertices in order: $\langle u_1, u_2 \rangle, \dots, \langle u_1, u_n \rangle$. Rename u_1 to u .
3. For every $1 \leq i \leq n$, take any $s \in V_i^0$ such that $s \neq u$ and identify vertices $\langle v_i, s \rangle$.

Observe that $v_i u$ is an edge in V_i^0 (for $1 \leq i \leq n$) and therefore in any triangle-free $(q + 1)$ -coloring of G' , v_i has a different color than u , by Lemma 3. It is now easy to verify that G is triangle-free q -colorable iff G' is triangle-free $(q + 1)$ -colorable. Also, since q is part of the problem (and therefore a constant), the reduction is polynomial w.r.t. n and m . \square

Theorem 13. *The TRIANGLEFREE-2-COLORING problem is \mathcal{NP} -hard on K_4 -free graphs.*

Proof. We create a polynomial reduction from the following problem, which is known to be \mathcal{NP} -complete [18]:

NOTALLEQUAL 3-SAT

Input: Boolean formula ϕ in conjunctive normal form, where each clause consists of exactly three literals.

Question: Does ϕ have a nae-satisfying assignment, i.e., in each clause at least one literal is true and at least one literal is false?

Let ϕ be a Boolean formula as described above, with n variables X and m clauses C_1, \dots, C_m . We assume, without loss of generality, that each clause consists of at least two unique literals, otherwise the instance is trivially false. We show a construction of K_4 -free graph $G = (V, E)$, such that ϕ is nae-satisfiable iff there exist a triangle-free 2-coloring of G .

1. For every clause $C_i \equiv (l_1^i \vee l_2^i \vee l_3^i)$, $1 \leq i \leq m$, add vertices l_1^i, l_2^i , and l_3^i to V , and add edges $\{l_1^i, l_2^i\}$, $\{l_2^i, l_3^i\}$ and $\{l_3^i, l_1^i\}$ to E .
2. For every variable $x \in X$, add vertices t_x and f_x to V , and edge $\{t_x, f_x\}$ to E .
3. For every variable x and every occurrence of literal $l \equiv \neg x$ in ϕ , add edge $\{t_x, l\}$ to E . For every variable x and every occurrence of literal $l' \equiv x$ in ϕ , add edge $\{f_x, l'\}$ to E .
4. For every edge $e \in E$, except for edges added in step 1, create K_4 -free polar gadget (Definition 3), with e begin the polar edge.

Observe that introducing polar gadgets in step 4 forces every vertex corresponding to the literal l to have to same color, and also every vertex corresponding to literal $\neg l$ to have the same color, but different than the color of l 's. Therefore the nae-satisfying assignment of each clause can be obtained from the coloring of each triangle introduced in step 1. Note that K_4 -free polar gadget is 2-colorable, by Lemma 7. From the above observations, we can conclude that ϕ has a nae-satisfying assignment iff G is triangle-free 2-colorable. Also, the reduction is polynomial w.r.t. n and m , and G is K_4 -free, since every polar gadget used is K_4 -free, by Lemma 7. \square

Theorem 14. *The TRIANGLEFREEPOLAR-2-COLORING problem is \mathcal{NP} -hard on graphs with maximum degree at most 3, but it is linear-time solvable on graphs with maximum degree at most 2.*

Proof. For hardness part, we create a polynomial reduction from the following problem, which we have proved to be \mathcal{NP} -complete (see Appendix A):

NOTALLEQUAL 3-SAT-4

Input: Boolean formula ϕ in conjunctive normal form, where each clause consists of exactly three literals and every variable appears at most four times in the formula.

Question: Does ϕ have a nae-satisfying assignment, i.e., in each clause at least one literal is true and at least one literal is false?

Let ϕ be a Boolean formula as described above, with n variables X and m clauses C_1, \dots, C_m . We show a construction of graph $G = (V, E)$ and $S \subseteq E$, such that the maximum vertex degree in G is at most 3, and ϕ is nae-satisfiable iff there exist a triangle-free 2-coloring of G with the set S of polar edges.

1. For every clause $C_i \equiv (l_1^i \vee l_2^i \vee l_3^i)$, $1 \leq i \leq m$, add vertices l_1^i , l_2^i , and l_3^i to V , and add edges $\{l_1^i, l_2^i\}$, $\{l_2^i, l_3^i\}$ and $\{l_3^i, l_1^i\}$ to E .
2. For every variable $x \in X$ and $\alpha \in \{t, f\}$, let $B_x^\alpha = (V_x^\alpha, E_x^\alpha)$, where $V_x^\alpha = \{\alpha_x^1, \dots, \alpha_x^7\}$ and $E_x^\alpha = \{\{\alpha_x^i, \alpha_x^{2i}\}, \{\alpha_x^i, \alpha_x^{2i+1}\} \mid i \in \{1, 2, 3\}\}$, i.e., B_x^α is a complete binary tree of height 2, with α_x^1 being the root, and $\{\alpha_x^4, \alpha_x^5, \alpha_x^6, \alpha_x^7\}$ the set of leaves. For each $x \in X$ add $V_x^t \cup V_x^f$ to V and $S_1 = E_x^t \cup E_x^f \cup \{t_x^1, f_x^1\}$ to E .
3. For every $x \in X$, do the following: take every occurrence of literal x (let's say it occurs $p \leq 4$ times) and order their corresponding vertices in V in an arbitrary way: x_1^t, \dots, x_p^t . Do similar with literals $\neg x$, and let the resulting sequence be x_1^f, \dots, x_q^f , for some $q \leq 4$. Add edges $S_2 = \{\{x_i^t, t_x^{i+3}\} \mid 1 \leq i \leq p\} \cup \{\{x_i^f, f_x^{i+3}\} \mid 1 \leq i \leq q\}$ to E . Here, we have connected all occurrences of positive literals x to the distinct leaves of B_x^t , and all occurrences of negative literals $\neg x$ to the distinct leaves of B_x^f .
4. Let $S = S_1 \cup S_2$.

It is easy to verify that every vertex in G has degree at most 3, and that we can construct G in time polynomial w.r.t. n and m . The correctness of the reduction now follows.

Assume that ϕ has a satisfying assignment $A : X \rightarrow \{true, false\}$, such that in each clause at least one literal is true and at least one literal is false. For every variable x and every vertex x_i^t and x_i^f , set $c(x_i^t) = 1$ and $c(x_i^f) = 2$, if $A(x) = true$, and $c(x_i^t) = 2$ and $c(x_i^f) = 1$, if $A(x) = false$. With this partial coloring we eliminated any possibility of there being any monochromatic triangle in G . We extend this coloring to the rest of the graph. Notice that by the choice of S , we have ensured that in every coloring of B_x^α , $c(\alpha_x^1) \neq c(\{\alpha_x^2, \alpha_x^3\}) \neq c(\{\alpha_x^4, \alpha_x^5, \alpha_x^6, \alpha_x^7\})$. Furthermore, since we have introduced the edge $\{t_x^1, f_x^1\}$, the coloring of B_x^t is simply a reflection of the coloring of B_x^f , in particular, $c(\{t_x^4, t_x^5, t_x^6, t_x^7\}) \neq c(\{f_x^4, f_x^5, f_x^6, f_x^7\})$. Therefore to properly color the vertices of G (to satisfy the choice of S), we set $c(t_x^1) = 2$ if $A(x) = true$ and $c(t_x^1) = 1$ if $A(x) = false$. The coloring can then be extended to the rest of B_x^t and to B_x^f in a straightforward way.

Now assume that there exists a triangle-free 2-coloring of G . Then by the coloring property of B_x^t and B_x^f mentioned earlier, every occurrence of literal x will have the same color, let's say 1, and every occurrence of $\neg x$ will be colored 2. Since each triangle corresponding to the clause $C_i \equiv (l_1^i \vee l_2^i \vee l_3^i)$ uses two colors, we can build the satisfying assignment A of ϕ , such that in each clause at least one literal is true and at least one literal is false, by setting $A(l) = true$ if $c(l) = 1$ and $A(l) = false$ if $c(l) = 2$.

When $\Delta(G) \leq 2$, then G is a union of isolated vertices, paths and/or cycles. Therefore G is not triangle-free 2-colorable iff G contains an odd cycle in which every edge is polar. The existence of such cycle can be checked in linear time. □

6 Conclusions and Future Work

In this paper we have shown several results regarding vertex coloring without monochromatic triangles. Many new interesting problems can be derived from this new coloring variant. We have already asked a few questions throughout this paper, in particular, it would be instructive to see the answer to Conjecture 1. Apart from that, we propose a handful of ways to extend our research:

- For positive results one can look for an algorithm that finds $\chi_3(G)$ in graphs with $\Delta(G) \geq 5$. Using an algorithm from [20] when $\Delta(G) = 5$ can only get us classic 5-coloring of G , and therefore using standard recoloring strategy may not produce an optimal solution. This issue requires more complicated algorithmic approach.
- For negative side, one can look for the smallest $\Delta(G)$, for which the TRIANGLEFREE- q -COLORING problem is \mathcal{NP} -hard.
- In the context of parametrized complexity we have only shown two results. Constructing algorithms for other choices of parameters is a fine research direction. It would also be good to know where the problems proved to be \mathcal{NP} -hard in Section 5 reside in W -hierarchy. From the proofs of the theorems in Section 5 it seems that one will need to first consider parametrized complexity of the NOTALLEQUAL SAT problem (and its variants) to achieve meaningful results.

Finally, we note that it is possible to further generalize the notion of χ_3 to get the parameter χ_r , for any $r \geq 3$ (notice that $\chi_2 = \chi$). This new parameter can restrict the coloring in two ways: either not allow monochromatic K_r , or not allow monochromatic $C_{r'}$ (cycle of length r'), for any $3 \leq r' \leq r$. Both extensions are interesting.

References

- [1] Alon, N., Yuster, R., Zwick, U. (1997), *Finding and counting given length cycles*, Algorithmica, 17(3), pp. 209–223.
- [2] Angelini, P., Frati, F. (2012), *Acyclically 3-colorable planar graphs*, Journal of Combinatorial Optimization, Vol. 24, Issue 2, pp. 116–130.
- [3] Appel, K., Haken, W. (1989), *Every planar map is four colorable*, Contemp. Math., 98, pp. 1–741.
- [4] Brooks, R. (1941), *On colouring the nodes of a network*, Mathematical Proceedings of the Cambridge Philosophical Society, 37(2), pp. 194–197.
- [5] Courcelle, B. (1990), *The monadic second-order logic for graphs I: Recognizable of finite graphs*, Information and Computation, Volume 85, pp. 12–75.

- [6] Dailey, D., P. (1980), *Uniqueness of colorability and colorability of planar 4-regular graphs are NP-complete*, Discrete Mathematics, 30 (3), pp. 289–293.
- [7] Deb R. (2008), *An efficient nonparametric test of the collective household model*, Available at SSRN: <http://ssrn.com/abstract=1107246>.
- [8] Fiala, J., Golovach, P., Kratochvíl, J. (2011) *Parametrized complexity of coloring problems: Treewidth versus vertex cover*, Theoretical Computer Science, Vol. 412, pp. 2513–2523.
- [9] Formanowicz, P., Tanaś, K. (2012), *A survey of graph coloring - its types, methods and applications*, Foundations of Computing and Decision Sciences 37(3), pp. 223–238.
- [10] Golumbic, M., C., Rheinboldt, W. (1980), *Algorithmic Graph Theory and Perfect Graphs*, Academic Press, New York.
- [11] Kaiser, T., Škrekovski, R. (2004), *Planar graph colorings without short monochromatic cycles*, Journal of Graph Theory, Vol. 46, Issue 1, pp. 25–38.
- [12] Karpiński, M. (2017), *Vertex 2-coloring without monochromatic cycles of fixed size is NP-complete*, Theoretical Computer Science, Vol. 659, pp. 88–94.
- [13] Kawarabayashi, K., Ozeki, K. (2010), *A simple algorithm for 4-coloring 3-colorable planar graphs*, Theoretical Computer Science, Vol. 411, Issues 26–28, pp. 2619–2622.
- [14] Mycielski, J. (1955), *Sur le coloriage des graphs*, Colloquium Mathematicae, Vol. 3, Issue 2, pp. 161–162.
- [15] Papadimitriou, C., Yannakakis, M. (1981), *The clique problem for planar graphs*, Inform. Proc. Letters 13, pp. 131–133.
- [16] Proskurowski, A., Sysło, M., M. (1986), *Efficient vertex-and edge-coloring of outerplanar graphs*, SIAM Journal on Algebraic and Discrete Methods, 7: 131–136.
- [17] Robertson, N., Sanders, D., P., Seymour, P., Thomas, R. (1996), *Efficiently four-coloring planar graphs*, STOC '96, pp. 571–575.
- [18] Schaefer, T., J. (1978), *The complexity of satisfiability problems*, STOC '78, pp. 216–226.
- [19] Shitov, Y. (2017), *A tractable NP-completeness proof for the two-coloring without monochromatic cycles of fixed length*, Theoretical Computer Science, Volume 674, pp. 116–118.
- [20] Skulrattanakulchai, S. (2006), *Delta-list vertex coloring in linear time*, Inf. Process. Lett. 98(3), pp. 101–106.
- [21] Thomassen, C. (2008), *2-List-coloring planar graphs without monochromatic triangles*, Journal of Combinatorial Theory, Series B 98, pp. 1337–1348.

- [22] Tovey, C. (1984), *A simplified NP-complete satisfiability problem*, Discrete Applied Mathematics, 8(1):85–89, 1984.

A NotAllEqual 3-SAT-4 is NP-complete

For the sake of the proof below we define the following: a clause C is NAE-satisfied by assignment A , if there exists $l_1 \in C$, such that $A(l_1) = true$ and there exists $l_2 \in C$, such that $A(l_2) = false$. A CNF formula ψ is NAE-satisfied by A , if all its clauses are NAE-satisfied by A .

Proof. We construct a polynomial reduction from the following \mathcal{NP} -complete problem:

3-SAT-4

Input: Boolean formula ϕ in conjunctive normal form, where each clause consists of exactly three literals and every variable appears exactly four times in the formula.

Question: Does ϕ have a satisfying assignment?

The 3-SAT-4 problem was proved to be \mathcal{NP} -complete by Tovey [22].

Let ϕ be the Boolean formula in CNF, as described above. Let X be the set of variables in ϕ and let $|X| = n$. Let $C = \{C_1, \dots, C_m\}$ be the set of clauses of ϕ . We now construct ϕ' , the instance of NOTALLEQUAL 3-SAT-4.

The variable set of ϕ' is $X \cup \{c_1, \dots, c_m\} \cup \{f_1, \dots, f_m\}$. We add the following clauses to ϕ' :

1. $(x \vee y \vee c_i) \wedge (z \vee \neg c_i \vee f_i)$, for every clause $C_i \equiv (x \vee y \vee z)$, where $1 \leq i \leq m$,
2. $(\neg f_m \vee \neg f_m \vee f_1) \wedge \bigwedge_{i=1}^{m-1} (\neg f_i \vee \neg f_i \vee f_{i+1})$

First, observe that in every assignment A' of ϕ' and every $1 \leq i, j \leq m$, $A'(f_i) = A'(f_j)$. This property is maintained by clauses from 2. This also implies that clauses from 2. are always NAE-satisfied.

Assume that A is an assignment satisfying ϕ . The assignment that NAE-satisfies ϕ' is then defined as follows:

- For every $x \in X$, set $A'(x) = A(x)$,
- For every $C_i \equiv (x \vee y \vee z)$, where $1 \leq i \leq m$, if $A(x) = A(y) = false$, then set $A'(c_i) = true$, otherwise $A'(c_i) = false$.
- For every $1 \leq i \leq m$, set $A'(f_i) = false$.

Clearly, assignment A' NAE-satisfies ϕ' .

Now assume that A' is some NAE-satisfying assignment of ϕ' . If in this assignment all f_i 's are true, then we can negate the assignment of all the variables of ϕ' , which would still result in a NEA-satisfying assignment. Therefore we can assume without loss of generality, that all f_i 's are false in A' . Take any $C_i \equiv (x \vee y \vee z)$, for $1 \leq i \leq m$. Because of clauses introduced in 1., if $A'(c_i) = true$, then $A'(z) = true$, and if $A'(c_i) = false$, then $A'(x) = true$

or $A'(y) = true$. Therefore at least one of the variables from $\{x, y, z\}$ will be true in A' . Hence, the assignment A , where $A(x) = A'(x)$ for all $x \in X$, is a satisfying assignment for ϕ .

To complete the proof we note that in ϕ' , every variable occurs at most 4 times and each clause has exactly 3 literals. Moreover, the ϕ' can be constructed in polynomial time w.r.t. n and m . We also note that given an arbitrary assignment, we can check whether it NAE-satisfies a CNF formula, in polynomial time. \square