

**SPECTRUM, TRACE AND OSCILLATION OF A
STURM-LIOUVILLE TYPE RETARDED DIFFERENTIAL
OPERATOR WITH INTERFACE CONDITIONS**

ERDOĞAN ŞEN

ABSTRACT. In this study, a formula for regularized sums of eigenvalues of a Sturm-Liouville problem with retarded argument at the point of discontinuity is obtained. Moreover, oscillation properties of the related problem is investigated.

2010 MATHEMATICS SUBJECT CLASSIFICATION. 34B24, 47A10, 47A55.

KEYWORDS AND PHRASES. Differential equation with retarded argument; interface conditions; spectrum; regularized trace; nodal points; oscillation.

1. INTRODUCTION

In this paper, we consider the boundary value problem for the differential equation with retarded argument

$$(1) \quad p(x)y''(x) + q(x)y(x - \Delta(x)) + \lambda^2 y(x) = 0,$$

on $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$, with boundary conditions

$$(2) \quad a_1 y(0) + a_2 y'(0) = 0,$$

$$(3) \quad y'(\pi) + d y(\pi) = 0$$

and interface conditions

$$(4) \quad \gamma_1 y\left(\frac{\pi}{2} - 0\right) - \delta_1 y\left(\frac{\pi}{2} + 0\right) = 0,$$

$$(5) \quad \gamma_2 y'\left(\frac{\pi}{2} - 0\right) - \delta_2 y'\left(\frac{\pi}{2} + 0\right) = 0,$$

where $p(x) = p_1^2$ for $x \in [0, \frac{\pi}{2})$ and $p(x) = p_2^2$ for $x \in (\frac{\pi}{2}, \pi]$; the real-valued function $q(x)$ is continuous in $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$ and has finite limits $q(\frac{\pi}{2} \pm 0) = \lim_{x \rightarrow \frac{\pi}{2} \pm 0} q(x)$, the real-valued function $\Delta(x) \geq 0$ is continuous in $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$ has finite limits $\Delta(\frac{\pi}{2} \pm 0) = \lim_{x \rightarrow \frac{\pi}{2} \pm 0} \Delta(x)$, if $x \in [0, \frac{\pi}{2})$ then $x - \Delta(x) \geq 0$; if $x \in (\frac{\pi}{2}, \pi]$ then $x - \Delta(x) \geq \frac{\pi}{2}$; λ is a spectral parameter; $p_i, a_i, d, \gamma_i, \delta_i$ ($i = 1, 2$) are arbitrary real numbers such that $p_i a_i d \neq 0$ ($i = 1, 2$).

Differential equations with retarded argument appeared as far back as the eighteenth century in connection with the solution of the problem of Euler on the investigation of the general form of curves similar to their own evolutes. Differential equations with retarded argument arise in many application of mathematical modelling: for example, combustion in the chamber of a liquid propellant rocket engine [1,2] and vibrations of the hammer in an electromagnetic circuit breaker [3,4].

Boundary value problems with interface conditions arise in varied assortment of physical transfer problems (see [5]). Also, some problems with interface conditions arise in thermal conduction problems for a thin laminated plate (i.e., a plate composed by materials with different characteristics piled in the thickness). In this class of problems, interface conditions across the interfaces should be added since the plate is laminated. The study of the structure of the solution in the matching region of the layer with the basis solution in the plate leads to consideration of an eigenvalue problem for a second-order differential operator with piecewise continuous coefficients and interface conditions [6].

The asymptotic formulas for eigenvalues of boundary value problems with retarded argument obtained in [7-17]. In [17], the principal term of asymptotic distribution of eigenvalues of the problem (1)-(5) was obtained up to $O(\frac{1}{n})$. But we need sharper asymptotic formulas. Therefore, we improve this formula up to $O(\frac{1}{n^3})$.

The theory of regularized traces of Sturm-Liouville operators stems from the paper [18] of Gelfand and Levitan. Trace formulas for the Sturm-Liouville equation with a complex valued potential and with two point boundary conditions obtained in [19]. Regularized trace of the Sturm-Liouville problem with irregular boundary conditions investigated in [20]. A regularized trace formula for the matrix Sturm-Liouville operator found in [21]. The first regularized traces of boundary value problems with unbounded operator coefficient in a finite interval obtained in [22-24]. The second regularized trace of a differential operator with bounded operator coefficient and with the mixed boundary conditions can be found in [25]. For a comprehensive review of traces of Schrödinger operators, the interested reader is referred to [26].

As mentioned above partly, the literature is about the regularized traces of classic type differential operators is so rich and diverse. For a more comprehensive list, one can refer to the survey paper [27] and the paper [28]. However, there are only a few works on regularized traces and oscillation properties for differential operators with retarded argument. M. Pikula in [8] obtained trace formula of first order:

if $\tau \geq \pi$

$$\sum_{n=1}^{\infty} (\lambda_n(\tau) - n^2) = q(0) \varphi(-\tau) - \frac{h^2 + H^2}{2}$$

and if $\tau \leq \pi$

$$\begin{aligned} & \sum_{n=1}^{\infty} \left[\lambda_n(\tau) - n^2 - \frac{2}{\pi} \left(h + H + \frac{\cos n\tau}{2} \int_{\tau}^{\pi} q(t) dt \right) \right] \\ &= q(0) \varphi(-\tau) - \frac{h^2 + H^2}{2} + \frac{h + H}{2\pi} \int_{\tau}^{\pi} q(t) dt + \left[\frac{q(\tau) + q(\pi)}{4} - q(\tau) \right] \frac{\pi - \tau}{\pi} \\ &+ \left[\frac{b}{4} - \left(\frac{1}{\sqrt{8}} \int_{\tau}^{\pi} q(t) dt \right)^2 \right] \frac{\pi - 2\tau}{\pi} \end{aligned}$$

for the boundary-value problem of second order with retarded argument:

$$\begin{aligned} -y'' + q(x)y(x - \tau) &= \lambda y, \\ y'(0) - hy(0) &= y'(\pi) + Hy(\pi) = 0, \\ y(x - \tau) &= y(0)\varphi(x - \tau), \quad x \leq \tau, \quad \varphi(0) = 1. \end{aligned}$$

C-F. Yang in [11] obtained formula of the first regularized trace, oscillations of the eigenfunctions and the solutions of inverse nodal problem for discontinuous boundary value problems with retarded argument and with interface conditions at the one point of discontinuity. F. Hira in [14] obtained a formula for regularized sums of eigenvalues for a Sturm-Liouville problem with retarded argument at the one point of discontinuity which contains a spectral parameter in the boundary conditions and as a most recent study in this topic, Şen studied regularized trace formula and oscillation of eigenfunctions of a Sturm-Liouville operator with retarded argument at two points of discontinuity [16].

The goals of this article are to calculate the regularized trace and to find the nodal points of eigenfunctions for the problem (1)-(5). We point out that our results are extension and/or generalization to those in [7-11, 17-18, 28, 30, 31]. For example, if the retardation function $\Delta \equiv 0$ in (1) and $p(x) \equiv 1, \delta = 1, \gamma = 1$ we have the formula of the first regularized trace for the classical Sturm-Liouville operator which is called Gelfand-Levitan formula (see [29]).

2. THE SPECTRUM

Let $\omega_1(x, \lambda)$ be a solution of Eq. (1) on $[0, \frac{\pi}{2}]$, satisfying the initial conditions

$$(6) \quad \omega_1(0, \lambda) = a_2 \text{ and } \omega_1'(0, \lambda) = -a_1.$$

The conditions (6) define a unique solution of Eq. (1) on $[0, \frac{\pi}{2}]$ (see [7,17]).

After defining the above solution, then we will define the solution $\omega_2(x, \lambda)$ of Eq. (1) on $[\frac{\pi}{2}, \pi]$ by means of the solution $\omega_1(x, \lambda)$ using the initial conditions

$$(7) \quad \omega_2\left(\frac{\pi}{2}, \lambda\right) = \gamma_1 \delta_1^{-1} \omega_1\left(\frac{\pi}{2}, \lambda\right) \text{ and } \omega_2'\left(\frac{\pi}{2}, \lambda\right) = \gamma_2 \delta_2^{-1} \omega_1'\left(\frac{\pi}{2}, \lambda\right).$$

The conditions (7) define a unique solution of Eq. (1) on $[\frac{\pi}{2}, \pi]$ (see [17]).

Consequently, the function $\omega(x, \lambda)$ is defined on $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$ by the equality

$$\omega(x, \lambda) = \begin{cases} \omega_1(x, \lambda), & x \in [0, \frac{\pi}{2}), \\ \omega_2(x, \lambda), & x \in (\frac{\pi}{2}, \pi] \end{cases}$$

is a solution of (1) on $[0, \frac{\pi}{2}) \cup (\frac{\pi}{2}, \pi]$, which satisfies one of the boundary conditions and the interface conditions (4)-(5) Then the following integral equations hold:

$$(8) \quad \begin{aligned} \omega_1(x, \lambda) &= a_2 \cos \frac{\lambda}{p_1} x - \frac{a_1 p_1}{\lambda} \sin \frac{\lambda}{p_1} x \\ &- \frac{1}{\lambda p_1} \int_0^x q(\tau) \sin \frac{\lambda}{p_1} (x - \tau) \omega_1(\tau - \Delta(\tau), \lambda) d\tau, \end{aligned}$$

$$(9) \quad \begin{aligned} \omega_2(x, \lambda) &= \frac{\gamma_1}{\delta_1} \omega_1\left(\frac{\pi}{2}, \lambda\right) \cos \frac{\lambda}{p_2} \left(x - \frac{\pi}{2}\right) + \frac{\gamma_2 p_2 \omega_1'\left(\frac{\pi}{2}, \lambda\right)}{\lambda \delta_2} \sin \frac{\lambda}{p_2} \left(x - \frac{\pi}{2}\right) \\ &\quad - \frac{1}{\lambda p_2} \int_{\frac{\pi}{2}}^x q(\tau) \sin \frac{\lambda}{p_2} (x - \tau) \omega_2(\tau - \Delta(\tau), \lambda) d\tau. \end{aligned}$$

Solving the equations (8)-(9) by the method of successive approximation, we obtain the following asymptotic equalities for $|\lambda| \rightarrow \infty$:

$$(10) \quad \begin{aligned} \omega_1(x, \lambda) &= a_2 \cos \frac{\lambda}{p_1} x - \frac{a_1 p_1}{\lambda} \sin \frac{\lambda}{p_1} x - \frac{a_2}{2\lambda p_1} \int_0^x q(\tau) \sin \frac{\lambda}{p_1} (x - \Delta(\tau)) d\tau \\ &\quad - \frac{a_2}{2\lambda p_1} \int_0^x q(\tau) \sin \frac{\lambda}{p_1} (x - (2\tau - \Delta(\tau))) d\tau + O\left(\frac{1}{\lambda^2}\right). \end{aligned}$$

Differentiating (10) with respect to x , we get

$$(11) \quad \begin{aligned} \omega_1'(x, \lambda) &= -\frac{a_2 \lambda}{p_1} \sin \frac{\lambda}{p_1} x - a_1 \cos \frac{\lambda}{p_1} x - \frac{a_2}{2p_1^2} \int_0^x q(\tau) \cos \frac{\lambda}{p_1} (x - \Delta(\tau)) d\tau \\ &\quad - \frac{a_2}{2p_1^2} \int_0^x q(\tau) \cos \frac{\lambda}{p_1} (x - (2\tau - \Delta(\tau))) d\tau + O\left(\frac{1}{\lambda}\right). \end{aligned}$$

Using the fact that,

$$O\left(\frac{1}{\lambda}\right) = \begin{cases} \int_0^x q(\tau) \sin \frac{\lambda}{p_1} (2\tau - \Delta(\tau)) d\tau, & x \in [0, \frac{\pi}{2}]; \\ \int_0^x q(\tau) \cos \frac{\lambda}{p_1} (2\tau - \Delta(\tau)) d\tau, & x \in [0, \frac{\pi}{2}]; \\ \int_{\frac{\pi}{2}}^x q(\tau) \sin \frac{\lambda}{p_2} (2\tau - \Delta(\tau)) d\tau, & x \in [\frac{\pi}{2}, \pi]; \\ \int_{\frac{\pi}{2}}^x q(\tau) \cos \frac{\lambda}{p_2} (2\tau - \Delta(\tau)) d\tau, & x \in [\frac{\pi}{2}, \pi] \end{cases}$$

(see [9]), (10) and (11) we have

$$(12) \quad \begin{aligned} \omega_2(x, \lambda) &= \frac{\gamma_1 a_2}{\delta_1} \cos \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x\right) - \frac{\gamma_1 a_1 p_1}{\lambda \delta_1} \sin \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x\right) \\ &\quad - \frac{\gamma_1 a_2 (p_1 + p_2)}{2\lambda \delta_1 p_1 p_2} \left\{ \left[B\left(\frac{\pi}{2}, \lambda\right) + D(x, \lambda) \right] \sin \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x\right) \right. \\ &\quad \left. + \left[A\left(\frac{\pi}{2}, \lambda\right) + C(x, \lambda) \right] \cos \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x\right) \right\} + O\left(\frac{1}{\lambda^2}\right). \end{aligned}$$

Here,

$$\begin{aligned} A(x, \lambda) &= \int_0^x q(\tau) \sin \frac{\lambda \Delta(\tau)}{p_1} d\tau \quad x \in \left[0, \frac{\pi}{2}\right]; \\ B(x, \lambda) &= \int_0^x q(\tau) \cos \frac{\lambda \Delta(\tau)}{p_1} d\tau \quad x \in \left[0, \frac{\pi}{2}\right]; \\ C(x, \lambda) &= \int_{\frac{\pi}{2}}^x q(\tau) \sin \frac{\lambda \Delta(\tau)}{p_2} d\tau, \quad x \in \left[\frac{\pi}{2}, \pi\right]; \\ D(x, \lambda) &= \int_{\frac{\pi}{2}}^x q(\tau) \cos \frac{\lambda \Delta(\tau)}{p_2} d\tau, \quad x \in \left[\frac{\pi}{2}, \pi\right]. \end{aligned}$$

Differentiating (12) with respect to x , we get

$$\begin{aligned} \omega'_2(x, \lambda) &= -\frac{\gamma_1 a_2 \lambda}{\delta_1 p_2} \sin \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) - \frac{\gamma_1 a_1 p_1}{\delta_1 p_2} \cos \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \\ &\quad - \frac{\gamma_1 a_2 (p_1 + p_2)}{2\delta_1 p_1 p_2^2} \left\{ \left[B\left(\frac{\pi}{2}, \lambda\right) + D(x, \lambda) \right] \cos \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \right. \\ (13) \quad &\quad \left. - \left[A\left(\frac{\pi}{2}, \lambda\right) + C(x, \lambda) \right] \sin \frac{\lambda}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \right\} + O\left(\frac{1}{\lambda}\right). \end{aligned}$$

The solution $\omega(x, \lambda)$ defined above is a nontrivial solution of (1) satisfying conditions (2) and (4)-(5). Putting $\omega(x, \lambda)$ into (3), we get the characteristic equation

$$(14) \quad \Theta(\lambda) \equiv \omega'(\pi, \lambda) + d\omega(\pi, \lambda).$$

The set of eigenvalues of boundary value problem (1)-(5) coincides with the set of the squares of roots of (14), and eigenvalues are simple (see [17]). From (12), (13) and (14), we obtain

$$\begin{aligned} \Theta(\lambda) &\equiv -\frac{\gamma_1 a_2 \lambda}{\delta_1 p_2} \sin \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) - \frac{\gamma_1 a_1 p_1}{\delta_1 p_2} \cos \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \\ &\quad - \frac{\gamma_1 a_2 (p_1 + p_2)}{2\delta_1 p_1 p_2^2} \left\{ \left[B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right] \cos \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \right. \\ &\quad \left. - \left[A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right] \sin \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \right\} \\ &\quad + \frac{d\gamma_1 a_2}{\delta_1} \cos \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) + O\left(\frac{1}{\lambda}\right). \end{aligned}$$

Define

$$\Theta_0(\lambda) \equiv -\frac{\gamma_1 a_2 \lambda}{\delta_1 p_2} \sin \frac{\lambda \pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right).$$

Denote by $\lambda_n^0 = \frac{2p_1 p_2 n}{p_1 + p_2}$, $n \in \mathbb{Z}$, zeros of the function $\Theta_0(\lambda)$. It is simple algebraically except for $\lambda_{\pm 0}^0$ and we have

$$(15) \quad \lambda_n \sim \lambda_n^0 + O\left(\frac{1}{n}\right).$$

Denote by C_n the circle of radius, $0 < \varepsilon < \frac{1}{2}$, centered at the origin λ_n^0 and by Γ_{N_0} the counterclockwise square contours with four vertices

$$\begin{aligned} K &= N_0 + \varepsilon + N_0i, & L &= -N_0 - \varepsilon + N_0i, \\ M &= -N_0 - \varepsilon - N_0i, & N &= N_0 + \varepsilon - N_0i, \end{aligned}$$

where $i = \sqrt{-1}$ and N_0 is a natural number. Obviously, if $\lambda \in C_n$ or $\lambda \in \Gamma_{N_0}$, then $|\Theta_0(\lambda)| \geq M|\lambda|e^{|\operatorname{Im}\lambda|\pi}$ ($M > 0$) by using a similar method in [11, 32]. Thus, on $\lambda \in C_n$ or $\lambda \in \Gamma_{N_0}$, we have

$$\begin{aligned} \frac{\Theta(\lambda)}{\Theta_0(\lambda)} &= 1 + \frac{1}{\lambda} \left\{ \frac{a_1 p_1}{a_2} \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \right. \\ &+ \frac{p_1 + p_2}{2p_1 p_2} \left[\left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) - \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \right] \\ &\left. - dp_2 \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \right\} + O\left(\frac{1}{\lambda^2}\right). \end{aligned}$$

Expanding $\ln \frac{\Theta(\lambda)}{\Theta_0(\lambda)}$ by the Maclaurin formula, we find that

$$\begin{aligned} \ln \frac{\Theta(\lambda)}{\Theta_0(\lambda)} &= \frac{1}{\lambda} \left\{ \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \right. \\ &\times \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) - \frac{p_1 + p_2}{2p_1 p_2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \left. \right\} \\ &- \frac{1}{2\lambda^2} \left\{ \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right]^2 \right. \\ &\times \cot^2 \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) + \frac{(p_1 + p_2)^2}{4p_1^2 p_2^2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right)^2 \\ &+ \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \\ &\left. \times \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) \right\} + O\left(\frac{1}{\lambda^3}\right). \end{aligned}$$

Using the well-known Rouché Theorem, we get that $\Theta(\lambda)$ has the same number of zeros inside Γ_{N_0} as $\Theta_0(\lambda)$ (see [11]). Using the residue theorem, we have

$$\begin{aligned} \lambda_n - \lambda_n^0 &= -\frac{1}{2\pi i} \oint_{C_n} \ln \frac{\Theta(\lambda)}{\Theta_0(\lambda)} d\lambda \\ &= -\frac{1}{2\pi i} \oint_{C_n} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \frac{\cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{\lambda} d\lambda \\ &\quad + \frac{1}{2\pi i} \oint_{C_n} \frac{p_1 + p_2}{2p_1 p_2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \frac{d\lambda}{\lambda} \end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\pi i} \oint_{C_n} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right]^2 \frac{\cot^2 \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{2\lambda^2} d\lambda \\
& \quad + \frac{1}{2\pi i} \oint_{C_n} \frac{(p_1 + p_2)^2}{4p_1^2 p_2^2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right)^2 \frac{d\lambda}{2\lambda^2} \\
& \quad + \frac{1}{2\pi i} \oint_{C_n} \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \\
& \quad \times \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \frac{\cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{2\lambda^2} d\lambda + O\left(\frac{1}{n^3}\right).
\end{aligned}$$

Thus, using (15) and residue calculation we have proven the following theorem.

Theorem 2.1. *The spectrum of the problem (1)-(5) has the*

$$\begin{aligned}
\lambda_n &= \lambda_n^0 - \frac{1}{\lambda_n^0 \pi} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \\
& \quad + \frac{1}{(\lambda_n^0)^2 \pi} \left\{ \frac{p_1 + p_2}{p_1 p_2} \left(A\left(\frac{\pi}{2}, \lambda_n^0\right) + C(\pi, \lambda_n^0) \right) \right. \\
& \quad \times \left. \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \right\} \\
& \quad - \frac{1}{(\lambda_n^0)^3} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right]^2 + O\left(\frac{1}{n^3}\right)
\end{aligned}$$

asymptotic distribution for sufficiently large $|n|$.

3. THE REGULARIZED TRACE FORMULA

In this section, we will get regularized trace formula for the problem (1)-(5).

The asymptotic formula (15) for the eigenvalues implies that for all sufficiently large N_0 , the numbers λ_n with $|n| \leq N_0$ are inside Γ_{N_0} , and the numbers λ_n with $|n| > N_0$ are outside Γ_{N_0} . It follows that

$$\begin{aligned}
\lambda_{-0}^2 + \lambda_0^2 + \sum_{0 \neq n = -N_0}^{N_0} \left(\lambda_n^2 - (\lambda_n^0)^2 \right) &= -\frac{1}{2\pi i} \oint_{\Gamma_n} 2\lambda \ln \frac{\Delta(\lambda)}{\Delta_0(\lambda)} d\lambda \\
&= -\frac{1}{2\pi i} \oint_{\Gamma_n} 2 \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \\
& \times \cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right) d\lambda + \frac{1}{2\pi i} \oint_{\Gamma_n} 2 \left[\frac{p_1 + p_2}{2p_1 p_2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \right] d\lambda \\
& + \frac{1}{2\pi i} \oint_{\Gamma_n} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right]^2 \frac{\cot^2 \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{\lambda} d\lambda
\end{aligned}$$

$$\begin{aligned}
& + \frac{1}{2\pi i} \oint_{\Gamma_n} \frac{(p_1 + p_2)^2}{4p_1^2 p_2^2} \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right)^2 \frac{d\lambda}{\lambda} \\
& + \frac{1}{2\pi i} \oint_{\Gamma_n} \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \\
& \times \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \frac{\cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{\lambda} d\lambda + O\left(\frac{1}{N_0}\right),
\end{aligned}$$

by calculations, which implies that

$$\begin{aligned}
& \lambda_{-0}^2 + \lambda_0^2 + \sum_{0 \neq n = -N_0}^{N_0} \lambda_n^2 - (\lambda_n^0)^2 \\
& = -\frac{2}{\pi} \sum_{0 \neq n = -N_0}^{N_0} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \\
& \quad - \frac{2}{\pi} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, 0\right) + D(\pi, 0) \right) - dp_2 \right] \\
& + \sum_{0 \neq n = -N_0}^{N_0} \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \\
& \quad \times \left(A\left(\frac{\pi}{2}, \lambda_n^0\right) + C(\pi, \lambda_n^0) \right) \frac{1}{\lambda_n^0 \pi} + R \\
& \quad - \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, 0\right) + D(\pi, 0) \right) - dp_2 \right]^2 \\
(16) \quad & + \frac{(p_1 + p_2)^2}{4p_1^2 p_2^2} \left(A\left(\frac{\pi}{2}, 0\right) + C(\pi, 0) \right)^2 + O\left(\frac{1}{N_0}\right),
\end{aligned}$$

where,

$$\begin{aligned}
R = Res_{\lambda=0} & \left\{ \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda\right) + D(\pi, \lambda) \right) - dp_2 \right] \right. \\
& \left. \times \left(A\left(\frac{\pi}{2}, \lambda\right) + C(\pi, \lambda) \right) \frac{\cot \frac{\lambda\pi}{p_2} \left(\frac{p_2 - p_1}{2p_1} + 1 \right)}{\lambda} d\lambda \right\}.
\end{aligned}$$

Passing to the limit as $N_0 \rightarrow \infty$ in (16), we have

$$\begin{aligned}
& \lambda_{-0}^2 + \lambda_0^2 + \sum_{0 \neq n = -\infty}^{\infty} \left\{ \lambda_n^2 - (\lambda_n^0)^2 \right. \\
& \quad + \frac{2}{\pi} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \\
& \quad - \frac{p_1 + p_2}{p_1 p_2} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right] \\
& \quad \left. \times \left(A\left(\frac{\pi}{2}, \lambda_n^0\right) + C(\pi, \lambda_n^0) \right) \frac{1}{\lambda_n^0 \pi} \right\}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{2}{\pi} \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, 0\right) + D(\pi, 0) \right) - dp_2 \right] + R \\
&\quad - \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, 0\right) + D(\pi, 0) \right) - dp_2 \right]^2 \\
(17) \quad &\quad + \frac{(p_1 + p_2)^2}{4p_1^2 p_2^2} \left(A\left(\frac{\pi}{2}, 0\right) + C(\pi, 0) \right)^2.
\end{aligned}$$

The series on the left side of (17) is called the regularized trace of the problem (1)-(5).

4. THE OSCILLATION

In this chapter, we will find nodal points of eigenfunctions of the problem (1)-(5).

Let us rewrite the equation (10) and replace λ by λ_n

$$\begin{aligned}
\omega_1(x, \lambda_n) &= a_2 \cos \frac{\lambda_n x}{p_1} - \frac{a_1 p_1}{\lambda_n} \sin \frac{\lambda_n x}{p_1} - \frac{a_2 \sin \frac{\lambda_n x}{p_1}}{2\lambda_n p_1} \int_0^x q(\tau) \cos \left(\frac{\lambda_n \Delta(\tau)}{p_1} \right) d\tau \\
&\quad + \frac{a_2 \cos \frac{\lambda_n x}{p_1}}{2\lambda_n p_1} \int_0^x q(\tau) \sin \left(\frac{\lambda_n \Delta(\tau)}{p_1} \right) d\tau + O\left(\frac{1}{\lambda_n^2}\right).
\end{aligned}$$

Let us assume that x_n^j are the nodal points of the eigenfunction $\omega_1(x, \lambda_n)$.

Taking $\sin\left(\frac{\lambda_n x}{p_1}\right) \neq 0$ into account for sufficiently large n , we get

$$\cot \left(\frac{\lambda_n x}{p_1} \right) \left[1 + \frac{A(x, \lambda_n)}{2\lambda_n p_1} \right] = \frac{a_1 p_1}{a_2 \lambda_n} + \frac{B(x, \lambda_n)}{2\lambda_n p_1} + O\left(\frac{1}{\lambda_n^2}\right).$$

It follows easily that

$$(18) \quad \tan \left(\frac{\lambda_n x}{p_1} + \frac{\pi}{2} \right) = -\frac{a_1 p_1}{a_2 \lambda_n} - \frac{B(x, \lambda_n)}{2\lambda_n p_1} + O\left(\frac{1}{\lambda_n^2}\right).$$

Thus, solving the equation (18), one obtains

$$(19) \quad x_n^j = \frac{(j - \frac{1}{2}) \pi p_1}{\lambda_n} - \frac{a_1 p_1^2}{a_2 \lambda_n^2} - \frac{B(x_n^j, \lambda_n)}{2\lambda_n^2} + O\left(\frac{1}{\lambda_n^3}\right).$$

Note that

$$(20) \quad \lambda_n^{-1} = \frac{1}{\lambda_n^0} + \frac{\left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right]}{(\lambda_n^0)^3 \pi} + O\left(\frac{1}{n^4}\right)$$

and

$$(21) \quad \lambda_n^{-2} = \frac{1}{(\lambda_n^0)^2} + O\left(\frac{1}{n^4}\right).$$

Substituting (20) and (21) into (19) we have

$$x_n^j = \frac{(j - \frac{1}{2}) \pi p_1}{\lambda_n^0} - \frac{(j - \frac{1}{2}) p_1 \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} \left(B\left(\frac{\pi}{2}, \lambda_n^0\right) + D(\pi, \lambda_n^0) \right) - dp_2 \right]}{(\lambda_n^0)^3}$$

$$(22) \quad -\frac{a_1 p_1^2}{a_2 (\lambda_n^0)^2} - \frac{B\left(\frac{(j-\frac{1}{2})\pi p_1}{\lambda_n^0}, \lambda_n^0\right)}{2(\lambda_n^0)^2} + O\left(\frac{1}{n^3}\right), \quad j = 1, \overline{\left[\frac{n}{2}\right]}.$$

Similarly, from (12), we get

$$\begin{aligned} \frac{\delta_1 \omega_2(x, \lambda_n)}{\gamma_1} &= a_2 \cos \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) - \frac{a_1 p_1}{\lambda_n} \sin \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \\ &\quad - \frac{a_2(p_1 + p_2)}{2\lambda_n p_1 p_2} \left\{ \left[B\left(\frac{\pi}{2}, \lambda_n\right) + D(x, \lambda_n) \right] \sin \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \right. \\ &\quad \left. + \left[A\left(\frac{\pi}{2}, \lambda_n\right) + C(x, \lambda_n) \right] \cos \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \right\} + O\left(\frac{1}{\lambda_n^2}\right). \end{aligned}$$

For nodal points of $\omega_2(x, \lambda_n)$, again, taking $\sin \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \neq 0$ into account for sufficiently large n , we get

$$\begin{aligned} &\cot \frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) \left[1 - \frac{(A(\frac{\pi}{2}, \lambda_n) + C(x, \lambda_n))(p_1 + p_2)}{2\lambda_n p_1 p_2} \right] \\ &= \frac{a_1 p_1}{a_2 \lambda_n} + \frac{(p_1 + p_2)(B(\frac{\pi}{2}, \lambda_n) + D(x, \lambda_n))}{2\lambda_n p_1 p_2} + O\left(\frac{1}{\lambda_n^2}\right). \end{aligned}$$

and thus

$$\begin{aligned} &\tan \left(\frac{\lambda_n}{p_2} \left(\frac{\pi(p_2 - p_1)}{2p_1} + x \right) + \frac{\pi}{2} \right) \\ &= -\frac{a_1 p_1}{a_2 \lambda_n} - \frac{(p_1 + p_2)(B(\frac{\pi}{2}, \lambda_n) + D(x, \lambda_n))}{2\lambda_n p_1 p_2} + O\left(\frac{1}{\lambda_n^2}\right). \end{aligned} \quad (23)$$

Thus, solving the equation (23), one obtains

$$\begin{aligned} x_n^j &= -\frac{\pi(p_2 - p_1)}{2p_1} + \frac{(j - \frac{1}{2})\pi p_2}{\lambda_n} - \frac{\alpha_1 p_1 p_2}{a_2 \lambda_n^2} \\ (24) \quad &- \frac{(p_1 + p_2)(B(\frac{\pi}{2}, \lambda_n) + D(x_n^j, \lambda_n))}{2\lambda_n^2 p_1} + O\left(\frac{1}{\lambda_n^3}\right). \end{aligned}$$

Substituting (20) and (21) into (24) we have

$$\begin{aligned} x_n^j &= -\frac{\pi(p_2 - p_1)}{2p_1} + \frac{(j - \frac{1}{2})\pi p_2}{\lambda_n^0} \\ &\quad - \frac{(j - \frac{1}{2})p_2 \left[\frac{a_1 p_1}{a_2} + \frac{p_1 + p_2}{2p_1 p_2} (B(\frac{\pi}{2}, \lambda_n^0) + D(\pi, \lambda_n^0)) - d p_2 \right]}{(\lambda_n^0)^3} - \frac{\alpha_1 p_1 p_2}{a_2 (\lambda_n^0)^2} \\ (25) \quad &- \frac{(p_1 + p_2) \left[B(\frac{\pi}{2}, \lambda_n^0) + D\left(\frac{(j-\frac{1}{2})\pi p_2}{\lambda_n^0}, \lambda_n^0\right) \right]}{2(\lambda_n^0)^2 p_1} + O\left(\frac{1}{n^3}\right), \quad j = \overline{\left[\frac{n}{2}\right]} + 1, n. \end{aligned}$$

Thus we have proven the following theorem:

Theorem 4.1. *For sufficiently large n , we have the formulas (22) and (25) of the nodal points for the problem (1)-(5).*

REFERENCES

- [1] L. Crocco and S. Chang, *Theory of combustion instability in liquid propellant rocket motors*, Butterworths, London, 1956.
- [2] J. Kolesovas, D. Svitra, *Mathematical modelling of the combustion process in the chamber of a liquid propellant rocket engine*, Lithuanian Mathematical Journal, 15 (4) (1975) 632-642.
- [3] K.F. Teodorcik, *Self-oscillatory systems*, 3rd ed., Gostehizdat, Moscow, 1952. (Russian).
- [4] A.A. Harkevic, *Auto-oscillations*, Gostehizdat, Moscow, 1954. (Russian).
- [5] A. V. Likov, Y. A. Mikhaililov, *The Theory of Heat and Mass Transfer*, Qosenergaizdat, 1963 (In Russian).
- [6] I. Titeux, Y. Yakubov, *Completeness of root functions for thermal conduction in a strip with piecewise continuous coefficients*, Math. Models Methods Appl. Sci. 7 (7) (1997) 1035-1050.
- [7] S.B. Norkin, *Differential equations of the second order with retarded argument*, Translations of Mathematical Monographs, AMS, Providence, RI, 1972.
- [8] M. Pikula, *Regularized traces of differential operator of Sturm-Liouville type with retarded argument*, Differentsialnye Uravneniya, 26 (1) (1990) 103-109 (Russian); English translation: *Differential Equations*, 26 (1) (1990) 92-96.
- [9] A. Bayramov, S. Çalışkan and S. Uslu, *Computation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument*, Appl. Math. Comput., 191 (2007) 592-600.
- [10] E. Şen, A. Bayramov, *Calculation of eigenvalues and eigenfunctions of a discontinuous boundary value problem with retarded argument which contains a spectral parameter in the boundary condition*, Math. Comput. Model. 54 (11-12) (2011) 3090-3097.
- [11] C-F. Yang, *Trace and inverse problem of a discontinuous Sturm-Liouville operator with retarded argument*, J. Math. Anal. Appl., 395 (2012) 30-41.
- [12] F.A. Cetinkaya, K.R. Mamedov, *A boundary value problem with retarded argument and discontinuous coefficient in the differential equation*, Azerbaijan Journal of Mathematics, 7 (2017) 135-145.
- [13] E. Şen, M. Acikgoz, S. Araci, *Spectral problem for Sturm-Liouville operator with retarded argument which contains a spectral parameter in the boundary condition*, Ukrainian Mathematical Journal, 68 (8) (2017) 1263-1277.
- [14] F. Hira, *A trace formula for the Sturm-Liouville type equation with retarded argument*, Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat., 66 (1) (2017) 124-132.
- [15] M. Bayramoglu, A. Bayramov, E. Şen, *A regularized trace formula for a discontinuous Sturm-Liouville operator with delayed argument*, Electronic Journal of Differential Equations (EJDE), 2017 (104) (2017) 1-12.
- [16] E. Şen, *A regularized trace formula and oscillation of eigenfunctions of a Sturm-Liouville operator with retarded argument at two points of discontinuity*, Mathematical Methods in the Applied Sciences, doi: 10.1002/mma.4512.
- [17] E. Şen, A. Bayramov, *Spectral analysis of boundary value problems with retarded argument*, Commun. Fac. Sci. Univ. Ank. Sér. A1 Math. Stat., 66 (2) (2017) 175-194.
- [18] I.M. Gelfand, B.M. Levitan, *On a formula for eigenvalues of a differential operator of second order*, Doklady Akademii Nauk SSSR, 88 (1953) 593-596 (Russian).
- [19] A. Makin, *Regularized trace of the Sturm-Liouville operator with irregular boundary conditions*, Electronic Journal of Differential Equations (EJDE), 2009 (27) (2009) 1-8.
- [20] A. Makin, *Trace formulas for the Sturm-Liouville operator with regular boundary conditions*, Doklady Mathematics, 76 (2) (2007) 702-707.
- [21] C-F. Yang, *Trace formula for the matrix Sturm-Liouville operator*, Anal. Math. Phys. 6(2016) 31-41.
- [22] F.G. Maksudov, M. Bayramoglu, E.E. Adiguzelov, *On a regularized traces of the Sturm-Liouville operator on a finite interval with the unbounded operator coefficient*, Doklady Akademii Nauk SSSR, 277 (4) (1984); English translation: *Soviet Math. Dokl.*, 30 (1984) 169-173.

- [23] E. Adiguzelov, Y. Sezer, *The regularized trace of a self adjoint differential operator of higher order with unbounded operator coefficient*, Appl. Math. Comput., 218 (2011) 2113-2121.
- [24] E. Şen, A. Bayramov, K. Oruçoğlu; *Regularized trace formula for higher order differential operators with unbounded coefficients*, Electronic Journal of Differential Equations (EJDE), 2016 (31) (2016), 1-12.
- [25] E. Adiguzelov, Y. Sezer, *The second regularized trace of a self adjoint differential operator given in a finite interval with bounded operator coefficient*, Math. Comput. Model., 53 (2011) 553-565.
- [26] F. Gesztesy F., H. Holden, *On Trace Formulas for Schrödinger-Type Operators*. In: Truhlar D.G., Simon B. (eds) *Multiparticle Quantum Scattering With Applications to Nuclear, Atomic and Molecular Physics*. The IMA Volumes in Mathematics and its Applications, vol 89. Springer, New York, 1997.
- [27] V.A. Sadovnichii, V.E. Podolskii, *Traces of operators*, Uspekhi Mat. Nauk, 61 (2006) 89-156; English translation: Russian Math. Surveys 61 (2006) 885-953.
- [28] C.T. Fulton and S.A. Pruess, *Eigenvalue and eigenfunction asymptotics for regular Sturm-Liouville problems*, J. Math. Anal. Appl. 188 (1994) 297-340.
- [29] L.A. Dikii, *Trace formulas for Sturm-Liouville differential equations*, Uspekhi Mat. Nauk, 12 (3) (1958) 111-143.
- [30] C. T. Shieh, V.A. Yurko, *Inverse nodal and inverse spectral problems for discontinuous boundary value problem*, J. Math. Anal. Appl. 347 (2008) 266–272.
- [31] Y.H. Cheng, C.K. Law, *On the quasi-nodal map for the Sturm-Liouville problem*, Proc. Soc. Edinburgh, 136A (2006) 71-86.
- [32] V.A. Yurko, *Inverse Spectral Problems for Differential Operators and Their Applications*, Gordon and Breach, Amsterdam, 2000.

DEPARTMENT OF MATHEMATICS, NAMIK KEMAL UNIVERSITY, 59030, TEKİRDAĞ,
TURKEY

E-mail address: erdogan.math@gmail.com