

FUNDAMENTAL GROUPS AS LIMITS OF DISCRETE FUNDAMENTAL GROUPS

FEDERICO VIGOLO

ABSTRACT. In this note we investigate to what extent the fundamental group of a metric space can be described as the inverse limit of its discrete fundamental groups. We show that some mild conditions suffice to imply the existence of an isomorphism and we provide a list of counterexamples to possible weakenings of these hypotheses.

1. INTRODUCTION

The discrete fundamental group (at scale 1) of a simplicial complex was first introduced by Barcelo, Kramer, Laubenbacher and Weaver [BKLW01] as a special case of A-theory—a general theory of connectivity of simplicial complexes. Barcelo, Capraro and White [BCW14] later extended this definition to the more general context of metric spaces and included in the definition a parameter $\theta > 0$ encoding the ‘scale’ of discretisation.

The discrete fundamental group $\pi_{1,\theta}(X, x_0)$ consists of the group of discrete closed θ -paths (finite sequences of points, $x_0 = z_0, z_1, \dots, z_n = x_0$, with $d(z_{i-1}, z_i) \leq \theta$ for every $i = 1, \dots, n$) up to θ -homotopy; see Section 2 for the precise definition. For a (locally path connected) metric space X , $\pi_{1,\theta}(X, x_0)$ can roughly be thought of as a quotient of the fundamental group $\pi_1(X, x_0)$ where all the loops of diameter at most θ are ignored.

The study of the groups $\pi_{1,\theta}(X, x_0)$ already proved to be very fruitful (see for example the survey [BL05]). When the parameter θ is large it can provide rather strong coarse invariants [DK17, FNvL17, Vig17]. On the other hand, the behaviour of $\pi_{1,\theta}(X, x_0)$ as θ tends to zero has not yet been investigated thoroughly. The following has been asked in [BCW14, Section 7]:

Question 1.1. *Can the fundamental group $\pi_1(X, x_0)$ be recovered as some sort of limit of $\pi_{1,\theta}(X, x_0)$ as θ tends to 0?*

In the same paper, the authors remarked that a natural choice of limit would be the inverse limit $\varprojlim \pi_{1,\theta}(X, x_0)$, since for every $\theta' < \theta$ there is a natural homomorphism $\pi_{1,\theta'}(X, x_0) \rightarrow \pi_{1,\theta}(X, x_0)$. Still, they also warned that $\varprojlim \pi_{1,\theta}(X, x_0)$ is not isomorphic to $\pi_1(X, x_0)$ in general, as can be easily seen by considering $X = \mathbb{R}^2 \setminus \{0\}$.

This note was born as a spin-off of [Vig17]. In that paper, we utilize the relation between the fundamental group and $\pi_{1,\theta}(X, x_0)$ for large θ to produce coarse invariants of warped systems. Since the formalism developed for that purpose could be applied for small parameters as well and proved

to be useful in tackling problems such as Question 1.1, we thought it was worthwhile to reorganise it in this self-contained note where we try to answer the following modification of Question 1.1:

Question 1.2. *For which metric spaces is the natural ‘discretisation homomorphism’ (see Section 3) from the fundamental group $\pi_1(X, x_0)$ to the projective limit $\varprojlim \pi_{1,\theta}(X, x_0)$ an isomorphism?*

The example of $\mathbb{R}^2 \setminus \{0\}$ suggests that completeness is a reasonable requirement to ask for X . Moreover, the example of manifolds with cusps illustrates the fact that compactness could be essential as well. It is fairly easy to realise that local path connectedness is also required, and further analysis suggests that semi-locally simply connectedness should be included too (definitions are given in Section 2). Our positive result is that these three conditions are actually enough:

Theorem 1.3. *If a metric space X is compact, locally path connected and semi-locally simply connected then $\pi_1(X, x_0) \cong \varprojlim \pi_{1,\theta}(X, x_0)$.*

Perhaps more interestingly, we also provide a series of counterexamples to possible extensions of Theorem 1.3. Indeed, we show that any combination of two of the three conditions above does not imply neither injectivity nor surjectivity of the natural homomorphism $\pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$. Extended arguments for the examples that we construct can be found in [Vig18].

Remark 1.4. Our result is actually more general than what we state in the introduction (see Theorem 3.2). This suggests that there might be room for improvement, but we doubt it is possible to find meaningful ‘if and only if’ conditions: one can always obtain ‘monstrous’ spaces X for which the isomorphism holds trivially, for example by letting X be the contractible space obtained by coning-off a ‘monstrous’ space Y .

Structure of the paper. In Section 2 we introduce the discrete fundamental group of a metric space and recall some notions of general topology. In Section 3 we prove Theorem 1.3, and in Section 4 and 5 we show that we can construct counterexamples (to surjectivity and injectivity, respectively) if the hypotheses are weakened.

Acknowledgements. I wish to thank my advisor Cornelia Druţu for her support and Gareth Wilkes for his helpful comments.

The author was funded by the EPSRC Grant 1502483 and the J.T.Hamilton Scholarship. We also thank the Isaac Newton Institute for Mathematical Sciences, Cambridge (EPSRC grant no EP/K032208/1), for support and hospitality during the programme “Non-positive curvature, group actions and cohomology”.

2. NOTATION AND PRELIMINARIES

2.1. General topology. Throughout the paper X will always be a metric space, $B(x, r)$ will denote the open ball of radius r centred at $x \in X$, and $\overline{B}(x, r)$ will be the closed ball. A *homotopy* between two paths $\gamma, \gamma': [0, 1] \rightarrow X$ must keep the endpoints fixed, while a *free homotopy* is allowed to move

them. A *closed path* or *loop* is a path $\gamma: [0, 1] \rightarrow X$ with $\gamma(0) = \gamma(1)$. We say that a loop is *null-homotopic* if it is homotopic to a constant path or, equivalently, if it is freely homotopic to a constant path (a free homotopy of loops is not allowed to break the loops into open paths).

Recall that X is *locally path connected* (shortened as l.p.c.) if for every point $x \in X$ and neighbourhood U of x there exists a neighbourhood $x \in V \subseteq U$ such that every two points in V are joined by a path in U (this is equivalent to the fact that every point of X admits a neighbourhood basis of path-connected open neighbourhoods [Sak13, Lemma 5.14.4]). We say that X is *uniformly locally path connected* (u.l.p.c.) if for every $\epsilon > 0$ there exists a $\delta > 0$ such that for every $x \in X$ any two points in $B(x, \delta)$ are connected by a path γ with image completely contained in $B(x, \epsilon)$.

The space X is *semi-locally simply connected* (s.l.s.c.) if for every $x \in X$ there exists a $\epsilon(x) > 0$ small enough so that every loop with image contained in $B(x, \epsilon(x))$ is null-homotopic in X (but is not necessarily null-homotopic in the ball). We say that X is *uniformly semi-locally simply connected* (u.s.l.s.c.) if it is semi-locally simply connected and one can choose $\epsilon(x)$ to be constant (*i.e.* uniform over $x \in X$).

Remark 2.1. It is easy to show that if X is compact then both local path connectedness and semi-local simple connectedness imply their uniform analogues [Sak13, Proposition 5.14.8].

2.2. Discrete fundamental group. Let X be a metric space and fix a parameter $\theta > 0$. A *discrete path* at *scale* θ (or θ -*path*) is a θ -Lipschitz map $Z: [n] \rightarrow X$ where $[n]$ is the set $\{0, 1, 2, \dots, n\}$ seen as a subset of \mathbb{R} . Equivalently, Z can be seen as an ordered sequence of points (z_0, \dots, z_n) in X with $d(z_i, z_{i+1}) \leq \theta$; we will use both notations throughout the paper.

We say that a θ -path $Z': [m] \rightarrow X$ is a *lazy version* (or *lazification*) of Z if it is obtained from it by repeating some values, *i.e.* if $m > n$ and $Z' = Z \circ f$ where $f: [m] \rightarrow [n]$ is a surjective monotone map.

Given two θ -paths Z_1 and Z_2 of the same length n , a *free θ -grid homotopy* between them is a θ -Lipschitz map $H: [n] \times [m] \rightarrow X$ such that $H(\cdot, 0) = Z_1$ and $H(\cdot, m) = Z_2$ (here the product $[n] \times [m]$ is equipped the ℓ^1 metric). A *θ -grid homotopy* is a free θ -grid homotopy so that $H(0, t) = Z_1(0) = Z_2(0)$ and $H(n, t) = Z_1(n) = Z_2(n)$ for every $t \in [m]$.

Definition 2.2. Two θ -paths Z_1 and Z_2 are *θ -homotopic* (denoted by $Z_1 \sim_\theta Z_2$) if they are equivalent under the equivalence relation induced by lazifications and θ -grid homotopies. Equivalently, $Z_1 \sim_\theta Z_2$ if and only if there exist lazy versions Z'_1 and Z'_2 of Z_1 and Z_2 which are homotopic via a θ -grid homotopy.

If the endpoint of a θ -path coincides with the starting point of a second θ -path, the two θ -paths can be concatenated. Since the operation of concatenation is clearly compatible with θ -homotopies, we can make the following definition:

Definition 2.3 ([BCW14]). Let x_0 be a base point in a metric space X . The *discrete fundamental group* at *scale* θ (or *θ -fundamental group*) is the group $\pi_{1,\theta}(X, x_0)$ consisting of θ -homotopy classes of closed θ -paths with endpoints x_0 , equipped with the operation of concatenation.

Note that, given $\theta' \leq \theta$, every θ' -path is also a θ -path and every θ' -homotopy is also a θ -homotopy. We therefore have a natural homomorphism $\pi_{1,\theta'}(X, x_0) \rightarrow \pi_{1,\theta}(X, x_0)$ and we can hence consider the inverse limit $\varprojlim \pi_{1,\theta}(X, x_0)$ of this direct system. We will show in Section 3 that continuous paths can be discretised—this will imply that there is a well-defined homomorphism $\hat{\cdot}: \pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$.

Remark 2.4. If the image of a continuous path has diameter smaller than θ then its θ -discretisation will be trivially θ -homotopic to a constant θ -path. Moreover, every closed θ -path of length at most 4θ is θ -homotopic to a constant path as follows: $(z_0, z_1, z_2, z_3, z_4 = z_0) \sim_{\theta} (z_0, z_1, z_1, z_0, z_0) \sim_{\theta} (z_0, z_0, z_0, z_0, z_0)$.

In some sense, the above is the only way in which loops become null-homotopic in $\pi_{1,\theta}$. For instance, it is proved in [BKLW01] that the 1-fundamental group of a graph is isomorphic to the fundamental group of the graph quotiented by the normal group generated by all the loops of length at most 4 in the graph. Under the assumption of some ‘niceness’ properties, this fact can be broadly generalised. For example, in [Vig17] we show that if X is a geodesic metric space then $\pi_{1,\theta}(X)$ is isomorphic to the quotient of $\pi_1(X)$ where all the loops of length at most 4θ (up to free homotopy) are killed. In this sense we can say that the θ -fundamental group is the quotient of the fundamental group where short cycles are ignored.

3. THE POSITIVE RESULTS

Given a continuous path $\gamma: [0, 1] \rightarrow X$ and a sequence of times $0 = t_0 \leq t_1 \leq \dots \leq t_n = 1$, we say that the discrete path $(\gamma(t_0), \dots, \gamma(t_n))$ is a θ -discretisation of γ if $\gamma(t_{i-1}, t_i) \subseteq \overline{B}(\gamma(t_{i-1}), \theta)$ for every $i = 1, \dots, n$. We will denote this θ -path by $\hat{\gamma}_{(t_0, \dots, t_n)}^{\theta}$. We will systematically use the following simple lemma:

Lemma 3.1. *Every continuous path admits a θ -discretisation. If two paths α and β are (freely) homotopic then any two discretisations $\hat{\alpha}_{t_0, \dots, t_n}^{\theta}$ and $\hat{\beta}_{t'_0, \dots, t'_m}^{\theta}$ are (freely) θ -homotopic.*

Proof. Fix the parameter $\theta > 0$. For any continuous path $\gamma: [0, 1] \rightarrow X$ consider the open cover of $[0, 1]$ given by the preimages $\gamma^{-1}(B(x, \frac{\theta}{2}))$ with $x \in X$. By the Lebesgue’s Number Lemma, there exists $N \in \mathbb{N}$ large enough so that for every $t \in [0, 1]$ the segment $[t - 1/N, t + 1/N]$ is fully contained in one of those open set. It follows that letting $t_i := i/N$ for $i = 0, \dots, N$ yields a θ -discretisation of γ .

It remains to prove that continuous homotopies induce discrete homotopies. Let $H: [0, 1]^2 \rightarrow X$ be an homotopy between two paths α and β . As above, we can use Lebesgue’s Number Lemma to deduce that there exists $N \in \mathbb{N}$ large enough so that for every $(t, s) \in [0, 1]^2$ the image under H of the ball $B((t, s), 1/N)$ has diameter smaller than θ . Consider the map $\hat{H}^{\theta}: [N]^2 \rightarrow X$ defined by

$$\hat{H}^{\theta}(i, j) := H\left(\frac{i}{N}, \frac{j}{N}\right).$$

Then the maps $\widehat{H}^\theta(\cdot, 0): [N] \rightarrow X$ and $\widehat{H}^\theta(\cdot, N): [N] \rightarrow X$ are θ -discretisations of α and β respectively, and \widehat{H}^θ is a free θ -grid homotopy between them.

To conclude the proof of the lemma it is hence enough to show that any two θ -discretisations of the same path are θ -homotopic. This is readily done. Let $\widehat{\gamma}_{t_0, \dots, t_n}^\theta$ and $\widehat{\gamma}_{t'_0, \dots, t'_m}^\theta$ be two θ -discretisations of a path γ . We can assume that the inequalities $t_0 < \dots < t_n$ are strict and—up to choosing a common refinement for the discretisations—we can also assume that $n \leq m$ and that for every t_i there exists a $j \geq i$ so that $t_i = t'_j$. We now define a surjective function $f: [m] \rightarrow [n]$ letting $f(j) := \max\{i \mid t_i \leq t'_j\}$. Then the θ -path $(\gamma(t_{f(0)}), \dots, \gamma(t_{f(m)}))$ is a lazification of $(\gamma(t_0), \dots, \gamma(t_n))$ and it is θ -homotopic to $(\gamma(t'_0), \dots, \gamma(t'_m))$ via a θ -grid homotopy consisting of a single step.

The same proof clearly implies the statement for free homotopies as well. \square

From Lemma 3.1 it follows that there is a well-defined θ -discretisation map $\widehat{\cdot}^\theta: \pi_1(X, x_0) \rightarrow \pi_{1,\theta}(X, x_0)$. As it is clear that the θ -discretisation of a concatenation of paths is θ -homotopic to the concatenation of their θ -discretisations, the map $\widehat{\cdot}^\theta$ is a homomorphism. To simplify the notation, we will generally drop the specific times t_i from the notation for the discretisation of a path and simply write $\widehat{\gamma}^\theta$.

Note that, given $\theta' < \theta$, any θ' -discretisation $\widehat{\gamma}_{t_0, \dots, t_n}^{\theta'}$ of a path γ is also a θ -discretisation. Therefore Lemma 3.1 also implies that the following diagram commutes:

$$\begin{array}{ccc} & \pi_1(X, x_0) & \\ \widehat{\cdot}^{\theta'} \swarrow & & \searrow \widehat{\cdot}^\theta \\ \pi_{1,\theta'}(X, x_0) & \longrightarrow & \pi_{1,\theta}(X, x_0). \end{array}$$

and hence the discretisation maps induce a homomorphism to the inverse limit $\widehat{\cdot}: \pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$. We will denote the image of (the homotopy class of) a continuous path γ by $\widehat{\gamma}$.

We can now prove the main result:

Theorem 3.2. *If X is u.l.p.c. and u.s.l.s.c. then the discretisation map $\widehat{\cdot}: \pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$ is an isomorphism.*

Proof. Injectivity: since X is u.s.l.s.c., there exists $\epsilon > 0$ so that every loop contained in a ball of radius at most ϵ is null-homotopic in X . Moreover, since X is u.l.p.c. there exists a $0 < \delta \leq \epsilon/2$ so that any two points in $B(x, \delta)$ are joined by a path in $B(x, \epsilon/2)$. Fix a parameter $0 < \theta < \delta$ and let γ be a continuous closed path whose θ -discretisation $\widehat{\gamma}^\theta$ is trivial in $\pi_{1,\theta}(X, x_0)$; we claim that γ is null-homotopic in X .

We can assume that $\widehat{\gamma}^\theta: [n] \rightarrow X$ is homotopic to the constant path via a θ -grid homotopy $H: [n] \times [m] \rightarrow X$. By construction, for every $0 \leq i \leq n-1$ and $0 \leq j \leq m$ there exist continuous paths $\alpha_{i,j}$ joining the point $H(i, j)$ with $H(i+1, j)$ and having its image completely contained in the ball of radius $\epsilon/2$ centred at $H(i, j)$. Similarly, for every $0 \leq i \leq n$ and $0 \leq j \leq m-1$ there exist analogous paths $\beta_{i,j}$ going from $H(i, j)$ to $H(i, j+1)$. We can




FIGURE 1. Constructing a continuous homotopy

concatenate these paths to obtain a path $\xi_{i,j}$ joining x_0 to $H(i,j)$ letting $\xi_{i,j} := [(\beta_{0,0}) \cdots (\beta_{0,j-1})][(\alpha_{0,j}) \cdots (\alpha_{i-1,j})]$. We now define some closed loops as follows (see Figure 1):

$$\eta_{i,j} := (\xi_{i,j})(\alpha_{i,j})(\beta_{i+1,j})(\alpha_{i,j+1}^{-1})(\beta_{i,j}^{-1})(\xi_{i,j}^{-1}).$$

By construction, the loops $\eta_{i,j}$ are null-homotopic and it is easy to see that the path γ is homotopic to the product of the $\eta_{i,j}$ (concatenating them in the appropriate order), and therefore $[\gamma]$ is the trivial element in $\pi_1(X, x_0)$. It follows that the map $\hat{\cdot}^\theta$ is injective and, a fortiori, $\hat{\cdot}$ is injective as well.

Surjectivity: a generic element of $\varprojlim \pi_{1,\theta}(X, x_0)$ can be represented by a family $(Z_\theta)_{\theta>0}$ —where Z_θ is a closed θ -path based at x_0 —so that for every $\theta' < \theta$ we have $Z_{\theta'} \sim_\theta Z_\theta$. Again, since X is u.l.p.c., for every $\theta > 0$ there exists a $0 < \delta(\theta) \leq \theta$ such that any two points in $B(x, \delta(\theta))$ are joined by a path in $B(x, \theta)$ (in what follows we assume $\delta(\theta)$ to be a decreasing function of θ). It follows that the points of the $\delta(\theta)$ -path $Z_{\delta(\theta)}$ can be joined with some small continuous paths and, concatenating these paths, we obtain a continuous loop γ_θ and we see that $Z_{\delta(\theta)}$ is a θ -discretisation of γ_θ .

From the proof of injectivity it follows that there exists a $\bar{\theta} > 0$ small enough so that $\hat{\cdot}^{\bar{\theta}}$ is injective. Note that for every $0 < \theta < \bar{\theta}$ we have

$$\hat{\gamma}_\theta^\theta \sim_\theta Z_{\delta(\theta)} \sim_{\delta(\bar{\theta})} Z_{\delta(\bar{\theta})} \sim_{\bar{\theta}} \hat{\gamma}_{\bar{\theta}}^{\bar{\theta}}.$$

It follows that $\hat{\gamma}_\theta^\theta$ and $\hat{\gamma}_{\bar{\theta}}^{\bar{\theta}}$ are $\bar{\theta}$ -homotopic and hence γ_θ and $\gamma_{\bar{\theta}}$ are homotopic. Therefore

$$\hat{\gamma}_{\bar{\theta}}^\theta \sim_\theta \hat{\gamma}_\theta^\theta \sim_\theta Z_{\delta(\theta)} \sim_\theta Z_\theta$$

for every $0 < \theta < \bar{\theta}$, and hence $\hat{\gamma}_{\bar{\theta}} = ([Z_\theta])_{\theta>0} \in \varprojlim \pi_{1,\theta}(X, x_0)$. \square

By Remark 2.1 we obtain the following:

Corollary 3.3 (Theorem 1.3). *If the space X is l.p.c., s.l.s.c. and compact then $\hat{\cdot}: \pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$ is an isomorphism.*

Note that from the proof of Theorem 3.2 it follows that if X is u.l.p.c. and u.s.l.s.c., then the projections $\varprojlim \pi_{1,\theta}(X, x_0) \rightarrow \pi_{1,\theta}(X, x_0)$ are injective for every θ small enough. However, such projections need not be surjective as shown by the following example:




FIGURE 2. Example of space with non-surjective projections

Example 3.4. Consider in \mathbb{R}^3 the cylindrical shell whose bases are disks of radius 1 centred at $(0, 0, 0)$ and $(1, 0, 0)$. Let $X_0 \subset \mathbb{R}^3$ be the space obtained by adding to the cylinder the segments joining $(\frac{1}{2}, e^{\frac{k\pi}{4}i})$ with $(\frac{1}{2}, \frac{1}{2}e^{\frac{k\pi}{4}i})$ for $k = 1, \dots, 8$. Let $X_1 \subset \mathbb{R}^3$ be a copy of X_0 , but translated by 1 on the x coordinate and with the y and z coordinates rescaled by a half. Similarly, X_2 is a translated and rescaled copy of X_1 and so on. Finally, let $X := \cup_{n \in \mathbb{N}} X_n$ be the ‘telescope space’ obtained taking the union all such cylinders (with the ambient metric of \mathbb{R}^3). See Figure 2.

Now, X is u.l.p.c. and u.s.l.s.c. (in fact, it is simply connected). We can hence use Theorem 3.2 to deduce that $\varprojlim \pi_{1,\theta}(X) = \{0\}$. Still, for every $n \in \mathbb{N}$ the discrete path in $X_n \subset X$ given by the points $(n + \frac{1}{2}, \frac{1}{2^{n+1}}e^{\frac{k\pi}{4}i})$ represents a non-trivial θ -path in $\pi_{1,\theta}(X)$ for $\theta = \|\frac{1}{2^{n+1}}e^0 - \frac{1}{2^{n+1}}e^{\frac{\pi}{4}i}\|$.


One may think that it should be true in general that if $\hat{\cdot}: \pi_1(X, x_0) \rightarrow \varprojlim \pi_{1,\theta}(X, x_0)$ is an isomorphism then the maps $\hat{\cdot}$ should be injective for θ small enough. The next example shows that this is not the case.

Example 3.5. Let $\frac{1}{2^n} \cdot \mathbb{S}^1$ be the circle of radius $1/2^n$ in \mathbb{R}^2 equipped with the metric d_n induced from \mathbb{R}^2 and let $X := \prod_{n \in \mathbb{N}} \frac{1}{2^n} \cdot \mathbb{S}^1$ equipped with the ℓ_1 metric, i.e. $d((x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}}) = \sum_{n \in \mathbb{N}} d_n(x_n, y_n)$. One can then check that the fundamental group of X is isomorphic to the direct product $\prod_{n \in \mathbb{N}} \mathbb{Z}$ and the discretisation $\hat{\cdot}$ is an isomorphism. Still, it is easy to see that $\pi_{1,\theta}(X, x_0)$ is isomorphic to a product of only finitely many copies of \mathbb{Z} because the small loops do not contribute.


4. THE COUNTEREXAMPLES: SURJECTIVITY

In this section we show how the surjectivity of $\hat{\cdot}$ may fail if one hypothesis among local path connectedness, semi-local simple connectedness and compactness is dropped.


Example 4.1 (not compact). In \mathbb{R}^3 , for every $n \in \mathbb{N}$ let S_n be the unit circle centred at $(0, 0, n)$ lying on the plane orthogonal to the axis $(0, 0, 1)$, and let $Y_n \subset S_n$ be the subset of points at angle $2\pi \frac{k+1/2}{2^n}$ for some $k \in \mathbb{N}$. Consider now the space $X \subset \mathbb{R}^3$ obtained as the union of the segments joining a point in Y_n with the two closest points in Y_{n+1} (see Figure 3a). The space X is contractible, as it is homeomorphic to a tree, still it is easy to see that $\varprojlim \pi_{1,\theta}(X)$ contains \mathbb{Z} as a subgroup.




(a) Non compact example



(b) Non locally path connected example



(c) Hawaiian earring




(d) Hawaiian window


FIGURE 3. Counterexamples to surjectivity

Example 4.2 (not l.p.c.). In \mathbb{R}^2 , let X be the union of the graph of the function $\sin(1/x)$ for $x \in (0, 1]$ with the segment I joining $(0, -1)$ to $(0, 1)$ and a path joining one end of the graph to I (see Figure 3b). This space is compact and simply connected, but $\varprojlim \pi_{1,\theta}(X) = \mathbb{Z}$.

Example 4.3 (not s.l.s.c.). Let X be the Hawaiian earring, *i.e.* the union of the circles of radius $1/n$ centred at $(1/n, 0)$ in \mathbb{R}^2 (Figure 3c). This space is the most common example of a non locally simply connected space and—despite still being a rather mysterious object—its fundamental group has been quite thoroughly studied. It is known that $\pi_1(X)$ injects in the projective limit of the fundamental group of the largest cycles $\pi_1(X) \hookrightarrow \varprojlim_n (\mathbb{Z}^{*n})$ but its image (which can be completely described) is not the whole group. Specifically, let $F_n = \mathbb{Z}^{*n}$ be freely generated by the set $\{a_1, \dots, a_n\}$, labelled so that the projection $F_{n+1} \rightarrow F_n$ is the map sending a_{n+1} to the identity. Let w_n be a reduced word in F_n and let $(w_n)_{n \in \mathbb{N}}$ represent an element of $\varprojlim_n (F_n)$. Then $(w_n)_{n \in \mathbb{N}}$ is in the image of $\pi_1(X)$ if and only if for every $i \in \mathbb{N}$ the number of times that the letter a_i appears in w_n is bounded uniformly on n (and hence eventually constant)[MM86, Eda92].



(a) Non locally path connected example



(b) Non semi-locally simply path connected

FIGURE 4. Counterexamples to injectivity

Now one can show that for every $n \in \mathbb{N}$ there is an appropriate value of θ so that $\pi_{1,\theta}(X) \cong F_n$, and that the discretisation $\hat{\cdot}$ coincides with the injection $\pi_1(X) \hookrightarrow \varprojlim_n (\mathbb{Z}^{*n})$ mentioned above. In particular, the discretisation homomorphism is not surjective.

Judging from the above example, one might hope to be able to characterise the image of $\pi_1(X)$ into $\varprojlim \pi_{1,\theta}(X)$ as the ‘sequences of words (in some generating set of $\pi_{1,\theta}(X)$) whose projections in $\pi_{1,\theta}(X)$ are eventually constant for every fixed θ ’. The next example shows that this is unlikely.

Example 4.4 (not s.l.s.c. bis). Let $X_0 \subset \mathbb{R}^2$ be the (empty) square with corners $(1, 1), (-1, 1), (-1, -1), (1, -1)$ and let X_1 be obtained from the square X_0 by adding the central cross (*i.e.* the two segments joining $(-1, 0)$ to $(1, 0)$ and $(0, -1)$ to $(0, 1)$). Now, let X_{n+1} be obtained from X_n by adding the central crosses to all the left-most squares of X_n and let $X = \bigcup_n X_n$ be a ‘Hawaiian window’.

Consider now the infinite path $\gamma: [0, \infty) \rightarrow X$ starting from the far right and zig-zagging from the top to the bottom while moving to the left—as the graph of $\sin(1/x)$ would do—(see Figure 3d). For every θ , the discretisation $\hat{\gamma}^\theta$ is θ -homotopic to a finite θ -path, and hence (by closing up the loop) γ can be used to define an element in $\varprojlim \pi_{1,\theta}(X)$. Still, γ is not homotopic to any continuous path $\gamma': [0, 1] \rightarrow X$ and hence $\hat{\gamma}^\theta$ cannot be in the image of $\pi_1(X)$, despite having ‘constant projections onto $\pi_{1,\theta}(X)$ ’.

5. THE COUNTEREXAMPLES: INJECTIVITY

As in Section 4, we show how injectivity of $\hat{\cdot}$ may fail if one hypothesis is dropped.




FIGURE 5. Choosing appropriate paths and points in the subset $C \subset X$

FIGURE 6. Counterexamples to injectivity

Example 5.1 (not compact). It is sufficient to consider $\mathbb{R}^2 \setminus \{0\}$, or a manifold with cusps if one still desires completeness.

Example 5.2 (not l.p.c.). In \mathbb{R}^3 , consider the graph of $z = \sin(\pi/x)$ as (x, y) range in $[0, 1] \times [-1, 1]$. Let $X \subset \mathbb{R}^3$ be this graph together with the limit square A and two squares B_1, B_2 running sideways the graph as in Figure 4a. The path γ cutting through these three squares and closing up straight through the graph of $\sin(1/x)$ is not homotopic to a constant path.

Indeed, if there was a continuous map $F: \mathbb{D} \rightarrow X$ with $F|_{\partial\mathbb{D}} = \gamma$ (here \mathbb{D} is the disk) then there would be a path α in $\mathbb{D} \setminus F^{-1}(B_1 \cup B_2)$ joining a point of $F^{-1}(A)$ to a point $X \setminus F^{-1}(A \cup B_1 \cup B_2)$. But this is impossible, as $F \circ \alpha$ would be a path reaching the limit square A by running along the graph of $\sin(1/x)$.

Example 5.3 (not s.l.s.c.). The basic idea is similar to that of Example 5.2, but it is a little more technical. Instead of adding to the graph of $\sin(\pi/x)$ three squares, the space X is obtained adding a whole triangular prism built over the triangle with vertices $(0, -1), (0, 1), (1, 1)$. Moreover, we also add on the vertical plane passing through $(0, -1), (1, -1)$ the subset of \mathbb{R}^2 obtained by filling in the space contained between the graph of $\sin(\pi/x)$ and the x -axis as x ranges in $(0, 1]$ (see Figure 4b). Let B denote the prisms, C the subset of the plane before mentioned and A the segment joining $(0, -1, -1)$ and $(0, -1, 1)$. We claim that the same path γ as in Example 5.2 is not homotopically trivial.

Assume by contradiction that γ is null-homotopic. Then there is a map $F: \mathbb{D} \rightarrow X$ that coincides with γ on $\partial\mathbb{D}$. Let $x_n \in \partial\mathbb{D}$ be the point sent to the midpoint between $(\frac{1}{n}, -1, 0)$ and $(\frac{1}{n+1}, -1, 0)$, and let $z_n \in C \subset X$ be the point on the vertical line through $F(x_n)$ lying at distance $1/2$ away from $F(x_n)$ (Figure 5). Let also $c_n \subset C_n$ denote the graph of $\sin(\pi/x)$ as x ranges in $[\frac{1}{n+1}, \frac{1}{n}]$. Note that c_n disconnects X .

It is simple to show that the image of F must contain the whole of $X \setminus B$. Moreover, using some extra care one can also show that we can choose a point

$y_n \in F^{-1}(z_n)$ and a path β_n connecting x_n to y_n such that the image $F(\beta_n)$ does not intersect curve c_n (Figure 5). In particular, any path $F(\beta_n)$ with n odd stays at distance at least $1/2$ from all the points z_m with m even (and vice versa). Since \mathbb{D} is compact, the map F must be uniformly continuous, and hence there exists an $\epsilon > 0$ such that the path β_n with n odd stays at distance at least ϵ from all the points y_m with m even (and vice versa).

Since \mathbb{D} is compact, both the sequence $(y_{2n})_{n \in \mathbb{N}}$ and $(y_{2n+1})_{n \in \mathbb{N}}$ will admit converging subsequences. It follows that there exist intertwined indices $2n < 2m + 1 < 2n' < 2m' + 1$ such that both $d_{\mathbb{D}}(y_{2n}, y_{2n'}) < \frac{\epsilon}{2}$ and $d_{\mathbb{D}}(y_{2m+1}, y_{2m'+1}) < \frac{\epsilon}{2}$. We can hence join up those pairs of points with segments of length at most $\epsilon/2$. Concatenating these small paths with the paths $\beta_{2n}, \beta_{2n'}^{-1}$ and $\beta_{2m+1}, \beta_{2m'+1}^{-1}$ we would obtain two disjoint continuous paths in \mathbb{D} linking intertwined pairs of points of $\partial\mathbb{D}$, which yields the desired contradiction.

REFERENCES

- [BCW14] Hélène Barcelo, Valerio Capraro, and Jacob A. White, *Discrete homology theory for metric spaces*, Bulletin of the London Mathematical Society **46** (2014), no. 5, 889–905.
- [BKLW01] Helene Barcelo, Xenia Kramer, Reinhard Laubenbacher, and Christopher Weaver, *Foundations of a connectivity theory for simplicial complexes*, Advances in Applied Mathematics **26** (2001), no. 2, 97–128.
- [BL05] Hélène Barcelo and Reinhard Laubenbacher, *Perspectives on a-homotopy theory and its applications*, Discrete mathematics **298** (2005), no. 1, 39–61.
- [DK17] Thiebaut Delabie and Ana Khukhro, *Coarse fundamental groups and box spaces*, arXiv preprint arXiv:1701.02919 (2017).
- [Eda92] Katsuya Eda, *Free σ -products and noncommutatively slender groups*, Journal of algebra **148** (1992), 243–263.
- [FNvL17] David Fisher, Thang Nguyen, and Wouter van Limbeek, *Coarse geometry of expanders from rigidity of warped cones*, arXiv preprint arXiv:1710.03085 (2017).
- [MM86] John W. Morgan and Ian Morrison, *A van kampen theorem for weak joins*, Proceedings of the London Mathematical Society **3** (1986), no. 3, 562–576.
- [Sak13] Katsuro Sakai, *Geometric aspects of general topology*, Springer, 2013.
- [Vig17] Federico Vigolo, *Discrete fundamental groups of warped cones and expanders*, arXiv preprint arXiv:1710.04955 (2017).
- [Vig18] Federico Vigolo, *Geometry of actions, expanders and warped cones*, Ph.D. thesis, University of Oxford, Trinity 2018.

Current address: Mathematical Institute, University of Oxford, Woodstock Road, Oxford, OX2 6GG, United Kingdom