arXiv:1710.07250v1 [math.NT] 19 Oct 2017

On k-normal elements over finite fields

Lucas Reis[1]

School of Mathematics and Statistics, Carleton University, 1125 Colonel By Drive, Ottawa ON (Canada), K1S 5B6

Abstract

The so called k-normal elements appear in the literature as a generalization of normal elements over finite fields. Recently, questions concerning the construction of k-normal elements and the existence of k-normal elements that are also primitive have attracted attention from many authors. In this paper we give alternative constructions of k-normal elements and, in particular, we obtain a sieve inequality for the existence of primitive, k-normal elements. As an application, we show the existence of primitive k-normal elements for a significant proportion of k's in many field extensions. In particular, we prove that there exist primitive k-normals in $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ in the case when k lies in the interval $[1, n/4]$, n has a special property and $q, n \geq 420$.

Keywords: primitive elements, normal bases, k-normal elements
2010 MSC: 12E20, 11T30, 12E20

1. Introduction

Let $\mathbb{F}_{q^n}$ be the finite field with q^n elements, where q is a prime power and n is a positive integer. We have two special notions of generators in the theory of finite fields. The multiplicative group $\mathbb{F}_{q^n}^*$ is cyclic, with $q^n - 1$ elements, and any generator is called *primitive*. Also, $\mathbb{F}_{q^n}$ can be regarded as an $\mathbb{F}_q$-vector space over $\mathbb{F}_q$: its dimension is n and, in particular, $\mathbb{F}_{q^n}$ is isomorphic to $\mathbb{F}_q^n$. An element $\alpha \in \mathbb{F}_{q^n}$ is said to be *normal* over $\mathbb{F}_q$ if $A = \{\alpha, \alpha^q, \cdots, \alpha^{q^{n-1}}\}$ is a basis of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$: A is frequently called a *normal basis*. Due to their high efficiency, normal bases are frequently used in cryptography and computer algebra systems; sometimes it is also interesting to use normal bases composed by primitive elements. The *Primitive Normal Basis Theorem* states that for any extension field $\mathbb{F}_{q^n}$ of $\mathbb{F}_q$, there exists a normal basis composed by primitive elements; this result was first proved by Lenstra and Schoof [8] and a proof without the use of a computer was later given in [3].

[1]Permanent address: Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte MG (Brazil), 30123-970.

Recently, Huczynska *et al* [7] introduce k-normal elements, extending the notion of normal elements. There are many equivalent definitions and here we present the most natural in the sense of vector spaces.

Definition 1.1. *For $\alpha \in \mathbb{F}_{q^n}$, consider the set $S_\alpha = \{\alpha, \alpha^q, \cdots, \alpha^{q^{n-1}}\}$ comprising the conjugates of α by the action of the Galois Group of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$. The element α is said to be k-normal over $\mathbb{F}_q$ if the vector space V_α generated by S_α has dimension $n-k$, i.e., $V_\alpha \subseteq \mathbb{F}_{q^n}$ has co-dimension k.*

From definition, 0-normal elements correspond to normal elements in the usual sense. Also, the concept of k-normal depends strongly on the base field that we are working. For this reason, unless otherwise stated, $\alpha \in \mathbb{F}_{q^n}$ is k-normal if it is k-normal over $\mathbb{F}_q$.

In [7], the authors find a formula for the number of k-normals and, consequently, obtain some results on the density of these elements. Motivated by the Primitive Normal Basis Theorem, they obtain an existence result on primitive, 1-normal elements.

Theorem 1.2 ([7], Theorem 5.10)**.** *Let $q = p^e$ be a prime power and n a positive integer not divisible by p. Assume that $n \geq 6$ if $q \geq 11$ and that $n \geq 3$ if $3 \leq q \leq 9$. Then there exists a primitive 1-normal element of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.*

In [12], we explore some ideas of [7] and, in particular, we obtain a characterization of k-normals in the case when n is not divisible by p. In the same paper, we find some asymptotic results on the existence of k-normal elements that are primitive or have a reasonable high multiplicative order. In [1], the author obtains alternative characterizations of k-normals via some recursive constructions and, in particular, he presents a method for constructing 1-normal elements in even characteristic. Recently, Theorem 1.2 was extended to arbitrary $n \geq 3$ (see [13]), where the authors use many techniques for completing the case $\gcd(n, p) = 1$ and extending this existence result to the case when n is divisible by p.

In this paper we deal with the general question concerning the existence of primitive, k-normal elements. We introduce a particular class of k-normal elements, obtained from normal elements, that removes the pertinent obstruction $n \equiv 0 \pmod{p}$. In particular, we obtain a character sum formula, derived from the Lenstra-Schoof method, for the number of certain primitive k-normals. As an application, we obtain some existence results on primitive k-normals in special extensions of $\mathbb{F}_q$. We also extend some results of [13] to k-normal elements.

2. Preliminaries

Here we introduce some basic definitions related to the k-normal elements, adding some recent results. We start with some arithmetic functions and their polynomial version.

Definition 2.1. (a) *Let $f(x)$ be a monic polynomial with coefficients in $\mathbb{F}_q$. The Euler Phi Function for polynomials over $\mathbb{F}_q$ is given by*

$$\Phi(f) = \left| \left(\frac{\mathbb{F}_q[x]}{\langle f \rangle} \right)^* \right|,$$

where $\langle f \rangle$ is the ideal generated by $f(x)$ in $\mathbb{F}_q[x]$.

(b) *If t is a positive integer (or a monic polynomial over $\mathbb{F}_q$), $W(t)$ denotes the number of square-free (monic) divisors of t.*

(c) *If $f(x)$ is a monic polynomial with coefficients in $\mathbb{F}_q$, the Polynomial Mobius Function μ_q is given by $\mu_q(f) = 0$ is f is not square-free and $\mu_q(f) = (-1)^r$ if f writes as a product of r distinct irreducible factors over $\mathbb{F}_q$.*

2.1. q-polynomials and k-normals

For $f \in \mathbb{F}_q[x]$, $f = \sum_{i=0}^{s} a_i x^i$, we set $L_f(x) = \sum_{i=0}^{s} a_i x^{q^i}$ as the q-associate of f. Also, for $\alpha \in \mathbb{F}_{q^n}$, we set $f \circ \alpha = L_f(\alpha) = \sum_{i=0}^{s} a_i \alpha^{q^i}$.

As follows, we have some basic properties of the q-associates:

Lemma 2.2. *([9], Theorem 3.62) Let $f, g \in \mathbb{F}_q[x]$. The following hold:*

(i) $L_f(L_g(x)) = L_{fg}(x)$,
(ii) $L_f(x) + L_g(x) = L_{f+g}(x)$.

For $\alpha \in \mathbb{F}_{q^n}$, we define $\mathcal{I}_\alpha$ as the subset of $\mathbb{F}_q[x]$ comprising the polynomials $f(x)$ for which $f(x) \circ \alpha = 0$, i.e., $L_f(\alpha) = 0$. Notice that $x^n - 1 \in \mathcal{I}_\alpha$ and, from the previous Lemma, it can be verified that $\mathcal{I}_\alpha$ is an ideal of $\mathbb{F}_q[x]$, hence $\mathcal{I}_\alpha$ is generated by a polynomial $m_\alpha(x)$. We can suppose $m_\alpha(x)$ monic. The polynomial $m_\alpha(x)$ is defined as the $\mathbb{F}_q$-order of α. This is a dual definition of multiplicative order in $\mathbb{F}_{q^n}^*$.

Clearly $m_\alpha(x)$ is always a divisor of $x^n - 1$, hence its degree is j for some $0 \leq j \leq n$. Notice that $j = 0$ if and only if $m_\alpha(x) = 1$, i.e., $\alpha = 0$. The following result shows a connection between k-normal elements and their $\mathbb{F}_q$-order.

Proposition 2.3. *([7], Theorem 3.2) Let $\alpha \in \mathbb{F}_{q^n}$. Then α is k-normal if and only if $m_\alpha(x)$ has degree $n - k$.*

In particular, an element α is normal if and only if $m_\alpha(x) = x^n - 1$. We see that the existence of k-normals depends on the existence of a polynomial of degree $n - k$ dividing $x^n - 1$ over $\mathbb{F}_q$; as follows, we have a formula for the number of k-normal elements.

Lemma 2.4. *([7], Theorem 3.5) The number N_k of $k-$normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$ is given by*

$$N_k = \sum_{\substack{h | x^n - 1 \\ \deg(h) = n-k}} \Phi(h), \tag{1}$$

where the divisors are monic and polynomial division is over $\mathbb{F}_q$.

One of the main steps in the proof of the result above is the fact that, for each monic divisor $f(x)$ of $x^n - 1$, there exists $\Phi_q(f(x))$ elements $\alpha \in \mathbb{F}_{q^n}$ for which $m_\alpha(x) = f(x)$. This fact will be further used in this paper.

Since $\frac{x^n-1}{x-1}$ divides $x^n - 1$, there are 1-normal elements over any finite field extension. However, the sum in equality (1) can be empty: for instance, if $q = 5$ and $n = 7$:

$$x^7 - 1 = (x - 1)(x^6 + x^5 + x^4 + x^3 + x^2 + x + 1)$$

is the factorization of $x^7 - 1$ into irreducible factors over $\mathbb{F}_5$. In particular, there are no 2, 3, 4 or 5-normal elements of $\mathbb{F}_{5^7}$ over $\mathbb{F}_5$. More generally, if n is a prime and q is primitive $\mod n$, $x^n - 1$ factors as $(x - 1)(x^{n-1} + \cdots + x + 1)$ and we do not have k-normal elements for any $1 < k < n - 1$. The existence of k-normals is not guaranteed for generic values of k.

As follows, we may construct k-normal elements from a given normal element.

Lemma 2.5. *Let $\beta \in \mathbb{F}_{q^n}$ be a normal element over $\mathbb{F}_q$ and $f(x)$ be a polynomial of degree k such that $f(x)$ divides $x^n - 1$. Then $\alpha = f(x) \circ \beta$ is k-normal.*

Proof. We prove that $m_\alpha(x) = \frac{x^n-1}{f(x)}$ and this implies the desired result. Notice that $\frac{x^n-1}{f(x)} \circ \alpha = \frac{x^n-1}{f(x)} \circ (f(x) \circ \beta) = (x^n - 1) \circ \beta = 0$, hence $m_\alpha(x)$ divides $\frac{x^n-1}{f(x)}$. It cannot divide strictly because $m_\beta(x) = x^n - 1$ and this completes the proof. □

In particular, we have a method for constructing k-normal elements when they exist: if we find a divisor $f(x)$ of $x^n - 1$ of degree k and a normal element $\beta \in \mathbb{F}_{q^n}$, the element $\alpha = f(x) \circ \beta$ is k-normal. There are many ways of finding normal elements in finite field extensions, including constructive and random methods; this is a classical topic in the theory of finite fields and the reader can easily find a wide variety of papers regarding those methods. For instance, see [6].

2.2. A characteristic equation for elements with prescribed $\mathbb{F}_q$-order

We have noticed that we may construct k-normal elements from the normal elements. However, it is not guaranteed that this method describes every k-normal in $\mathbb{F}_{q^n}$. For a polynomial f dividing $x^n - 1$, set

$$\Psi_f(x) = \prod_{m_\alpha = f(x)} (x - \alpha),$$

the polynomial of least degree that vanishes in every element $\alpha \in \mathbb{F}_{q^n}$ with $m_\alpha = f(x)$. Clearly $m_\alpha = f$ if and only if $\Psi_f(\alpha) = 0$. Also, for $f(x) \in \mathbb{F}_q[x]$ and $\alpha \in \mathbb{F}_{q^n}$, we have $f(x) \circ \alpha = L_f(\alpha) = 0$ if and only if m_α divides $f(x)$. In particular, this shows that $L_f(x) = \prod_{g|f} \Psi_g(x)$. This identity is similar to the one describing cyclotomic polynomials

$$x^n - 1 = \prod_{n|d} \Phi_d(x).$$

From the Mobius Inversion formula, we may deduce $\Phi_d(x) = \prod_{r|d}(x^r - 1)^{\mu(d/r)}$. This last equality describes the elements of multiplicative order d in finite fields. Motivated by this characterization, we obtain the following:

Proposition 2.6. *Let $f(x)$ be any divisor of $x^n - 1$ over $\mathbb{F}_q$. The following holds:*

$$\Psi_f(x) = \prod_{g|f} L_g(x)^{\mu_q(f/g)}, \tag{2}$$

where g is monic and polynomial division is over $\mathbb{F}_q$.

Proof. Notice that

$$\prod_{g|f} L_g(x)^{\mu_q(f/g)} = \prod_{g|f} L_{f/g}(x)^{\mu_q(g)},$$

and, from $L_{f/g}(x) = \prod_{h|f/g} \Psi_h(x)$, we obtain

$$\prod_{g|f} L_{f/g}(x)^{\mu_q(g)} = \prod_{h|f} \Psi_h(x)^{\sum_{g|f/h} \mu_q(g)}.$$

Writing f/h as product of irreducibles over $\mathbb{F}_q$, we can easily see that

$$\sum_{g|f/h} \mu_q(g) = \begin{cases} 1 & \text{if } f/h = 1, \\ 0 & \text{otherwise.} \end{cases}.$$

This shows that

$$\prod_{g|f} L_g(x)^{\mu_q(f/g)} = \Psi_f(x).$$

□

In particular, if we set

$$\Lambda_k(x) = \prod_{\substack{f|x^n-1 \\ \deg f = n-k}} \Psi_f(x),$$

an element $\alpha \in \mathbb{F}_{q^n}$ is k-normal if and only if $\Lambda_k(\alpha) = 0$. For instance, we may characterize the normal elements in $\mathbb{F}_{q^n}$: an element $\alpha \in \mathbb{F}_{q^n}$ is normal if and only if $\Lambda_0(\alpha) = 0$, where

$$\Lambda_0(x) = \Psi_{x^n-1}(x) = \prod_{g|x^n-1} L_g(x)^{\mu_q(x^n-1/g)}.$$

Example 2.7. *Suppose that $q \equiv 3 \pmod 4$ and $n = 4$. Then $\Lambda_0(x) = \frac{x^{q^4-1}-1}{(x^{q^2-1}+1)(x^{q^2-1}-1)} = \sum_{i=0}^{\frac{q^2-1}{2}} x^{2i(q^2-1)}$. Hence, $\alpha \in \mathbb{F}_{q^4}$ is normal if and only if*

$$\sum_{i=0}^{\frac{q^2-1}{2}} \alpha^{2i(q^2-1)} = 0.$$

In general, we have shown that the k-normals can be described as the zeroes of a univariate polynomial over $\mathbb{F}_q$; this polynomial can be computed from the factorization of $x^n - 1$ over $\mathbb{F}_q$.

2.3. Some recent results

The proof of Theorem 1.2 is based in an application of the *Lenstra-Schoof* method, introduced in [8]; this method has been used frequently in the characterization of elements in finite fields with particular properties like being primitive, normal and of zero-trace. In particular, from Corollary 5.8 of [7], we can easily deduce the following:

Lemma 2.8. *Suppose that q is a power of a prime p, $n \geq 2$ is a positive integer not divisible by p and $T(x) = \frac{x^n-1}{x-1}$. If*

$$W(T) \cdot W(q^n - 1) < q^{n/2-1}, \tag{3}$$

there exist primitive, 1-normal elements of $\mathbb{F}_{q^n}$ over $\mathbb{F}_q$.

Inequality (3) is an essential step in the proof of Theorem 1.2 and it was studied in [4]; if $n \geq 6$ for $q \geq 11$ and $n \geq 3$ for $3 \leq q \leq 9$, this inequality holds for all but a finite number of pairs (q, n).

In [7], the authors propose an extension of the Theorem 1.2 for all pairs (q, n) with $n \geq 3$ as a problem ([7], Problem 6.2); they conjectured that such elements always exist. This was recently proved in [13], where the authors use many different techniques to complete Theorem 1.2 in the case $\gcd(n, p) = 1$ and add the case when p divides n.

In particular, when n is divisible by p^2, they obtain the following:

Lemma 2.9. *([13], Lemma 5.2) Suppose that $\mathbb{F}_q$ has characteristic p and let $n = p^2 s$ for any $s \geq 1$. Then $\alpha \in \mathbb{F}_{q^n}$ is such that $m_\alpha(x) = \frac{x^n-1}{x-1}$ if and only if $\beta = \mathrm{Tr}_{q^n/q^{ps}}(\alpha) = \sum_{i=0}^{p} \alpha^{q^{psi}}$ satisfies $m_\beta(x) = \frac{x^{ps}-1}{x-1}$.*

Using the well-known result on the existence of primitive elements with prescribed trace due to Cohen (see [5]), they prove the existence of primitive 1-normals in the case when p^2 divides n. The case $n = ps$ with $\gcd(p, s) = 1$ is dealt in a similar way of [7].

As follows, we have a generalization of Lemma 2.9.

Lemma 2.10. *([13], Lemma 5.2) Suppose that $\mathbb{F}_q$ has characteristic p and let $n = p^2 s$ for any $s \geq 1$. Also, let $f(x)$ be a polynomial dividing $x^s - 1$. Then $\alpha \in \mathbb{F}_{q^n}$ is such that $m_\alpha(x) = \frac{x^n-1}{f(x)}$ if and only if $\beta = \mathrm{Tr}_{q^n/q^{ps}}(\alpha) = \sum_{i=0}^{p-1} \alpha^{q^{psi}}$ satisfies $m_\beta(x) = \frac{x^{ps}-1}{f(x)}$.*

The proof is entirely similar to the proof of Lemma 2.9 (see [13]) so we will omit. Motivated by Theorem 5.3 of [13], we have the following.

Theorem 2.11. *Suppose that $n = p^2 \cdot s$ and $x^s - 1$ is divisible by a polynomial of degree k. Then there exists a primitive, k-normal element over $\mathbb{F}_q$.*

Proof. Let $f(x)$ be a polynomial of degree k such that $f(x)$ divides $x^s - 1$. In particular, $f(x)$ divides $x^{ps} - 1$, i.e., $g(x) = \frac{x^{ps}-1}{f(x)}$ is a polynomial. Since $g(x)$ divides $x^{ps} - 1$, we know that there exists an element $\beta \in \mathbb{F}_{q^{ps}}$ such that the $\mathbb{F}_q$-order $m_\beta(x)$ of β is $g(x)$. Clearly $g(x) \neq 1$, hence $\beta \neq 0$. According to [5], there exists a primitive element $\alpha \in \mathbb{F}_{q^n}$ such that $\mathrm{Tr}_{q^n/q^{ps}}(\alpha) = \beta$ and then, from Lemma 2.10, it follows that such an α satisfies $m_\alpha = \frac{x^n-1}{f(x)}$, i.e., α is a primitive, k-normal element. □

3. Characteristic function for a class of primitive k-normals

In this section, we use the method of Lenstra and Schoof in the characterization of primitive and normal elements. This method has been used by many different authors in a wide variety of existence problems. For this reason, we skip some details, which can be found in [3]. We recall the notion of *freeness*.

Definition 3.1. 1. *If m divides $q^n - 1$, an element $\alpha \in \mathbb{F}_{q^n}^*$ is said to be m-free if $\alpha = \beta^d$ for any divisor d of m implies $d = 1$.*
2. *If $m(x)$ divides $x^n - 1$, an element $\alpha \in \mathbb{F}_{q^n}$ is $m(x)$-free if $\alpha = h \circ \beta$ for any divisor $h(x)$ of $m(x)$ implies $h = 1$.*

It follows from definition that primitive elements correspond to the $(q^n - 1)$-free elements. Also, $\alpha \in \mathbb{F}_{q^n}$ is normal if and only if is $(x^n - 1)$-free. The concept of *freeness* derives some characteristic functions for primitive and normal elements. We pick the notation of [7].

Multiplicative Part: $\int\limits_{d|m} \eta_d$ denotes $\sum_{d|m} \frac{\mu(d)}{\varphi(d)} \sum_{(d)} \eta_d$, where μ and φ are the Mobius and Euler functions for integers, respectively, η_d is a typical multiplicative character of $\mathbb{F}_{q^n}$ of order d, and the sum $\sum_{(d)} \eta_d$ runs through all the multiplicative characters of order d.

Additive Part: χ denotes the canonical additive character on $\mathbb{F}_{q^n}$, i.e.

$$\chi(\omega) = \lambda\left(\sum_{i=0}^{n-1} \omega^{q^i}\right), \omega \in \mathbb{F}_{q^n},$$

where λ is the canonical additive character of $\mathbb{F}_q$ to $\mathbb{F}_p$. If D is a monic polynomial dividing $x^n - 1$ over $\mathbb{F}_q$, a typical character χ_* of $\mathbb{F}_{q^n}$ of $\mathbb{F}_q$-order D is one such that $\chi_*(D \circ^q \cdot)$ is the trivial additive character in $\mathbb{F}_{q^n}$ and D is minimal (in terms of degree) with this property. Let Δ_D be the set of all $\delta \in \mathbb{F}_{q^n}$ such that χ_δ has $\mathbb{F}_q$-order D, where $\chi_\delta(\omega) = \chi(\delta\omega)$ for any $\omega \in \mathbb{F}_{q^n}$. For instance, $\Delta_1 = \{0\}$ and $\Delta_{x-1} = \mathbb{F}_q^*$.

In the same way of the multiplicative part, $\int\limits_{D|T} \chi_{\delta_D}$ denotes the sum

$$\sum_{D|T} \frac{\mu_q(D)}{\Phi(D)} \sum_{(\delta_D)} \chi_{\delta_D},$$

where μ_q and Φ are the Mobius and Euler functions for polynomials over $\mathbb{F}_q$, respectively, χ_{δ_D} denotes a typical additive character of $\mathbb{F}_{q^n}$ of $\mathbb{F}_q$-Order D and the sum $\sum_{(\delta_D)} \chi_{\delta_D}$ runs through all the additive characters whose $\mathbb{F}_q$-order equals D, i.e., $\delta_D \in \Delta_D$.

For t dividing q^n-1 and D dividing x^n-1, set $\theta(t) = \frac{\varphi(t)}{t}$ and $\Theta(D) = \frac{\Phi(D)}{q^{\deg D}}$.

Theorem 3.2. [7, Section 5.2]

1. *For $w \in \mathbb{F}_{q^n}^*$ and t be a positive divisor of q^n-1,*

$$\omega_t(w) = \theta(t) \int_{d|t} \eta_d(w) = \begin{cases} 1 & \text{if } w \text{ is } t\text{-free,} \\ 0 & \text{otherwise.} \end{cases}$$

2. *For $w \in \mathbb{F}_{q^n}$ and D be a monic divisor of x^n-1,*

$$\Omega_D(w) = \Theta(D) \int_{E|D} \chi_{\delta_E}(w) = \begin{cases} 1 & \text{if } w \text{ is } D\text{-free,} \\ 0 & \text{otherwise.} \end{cases}$$

In particular, for $t = q^n-1$ and $D = x^n-1$, we obtain characteristic functions for primitive and normal elements, respectively. We write $\omega_{q^n-1} = \omega$ and $\Omega_{x^n-1} = \Omega$. As usual, we may extend the multiplicative characters to 0 by setting $\eta_1(0) = 1$, where η_1 is the trivial multiplicative character and $\eta(0) = 0$ if η is not trivial.

We have seen that some k-normals arise from a normal element β via the composition $f(x) \circ \beta$, where $f(x)$ is a divisor of x^n-1, $\deg f = k$; more than that, $\alpha = f(x) \circ \beta$ satisfies $m_\alpha = \frac{x^n-1}{f(x)}$. In particular, we may obtain the characteristic function for the primitive elements arising from this construction: note that $\Omega(w) \cdot \omega(L_f(w)) = 1$ if and only if w is normal and $L_f(w) = f(x) \circ w$ is primitive. This characteristic function describes a particular class of primitive k-normal elements. The following is straightforward.

Proposition 3.3. *Let $f(x)$ be a divisor of x^n-1 of degree k and n_f be the number of primitive elements of the form $f(x) \circ \alpha$, where α is a normal element. The following holds:*

$$\frac{n_f}{\theta(q^n-1)\Theta(x^n-1)} = \sum_{w \in \mathbb{F}_{q^n}} \int_{d|q^n-1} \int_{D|x^n-1} \eta_d(L_f(w)) \chi_{\delta_D}(w). \tag{4}$$

In particular, the number of primitive, k-normal elements in $\mathbb{F}_{q^n}$ is at least n_f.

3.1. Character sums and a sieve inequality

Here we make some estimates for the character sums that appear naturally from Eq. (4): note that η_d is the trivial multiplicative character if and only if $d = 1$. Also, χ_{δ_D} is the trivial additive character if and only if $\delta_D = 0$, i.e., $D = 1$.

As usual, we split the sum in Eq. (4) as Gauss sums types, according to the trivial and non-trivial characters. For each d dividing q^n-1 and D dividing x^n-1, set $G_f(\eta_d, \chi_{\delta_D}) = \sum_{w\in\mathbb{F}_{q^n}} \eta_d(L_f(w))\chi_{\delta_D}(w)$.

Note that, from Proposition 3.3,

$$\frac{n_f}{\theta(q^n-1)\Theta(x^n-1)} = s_0 + S_1 + S_2 + S_3,$$

where $s_0 = G_f(\eta_1, \chi_0)$, $S_1 = \sum_{\substack{D|x^n-1 \\ D\neq 1}} G_f(\eta_1, \chi_{\delta_D})$, $S_2 = \sum_{\substack{d|q^n-1 \\ d\neq 1}} G_f(\eta_d, \chi_0)$ and $S_3 = \sum_{\substack{d|q^n-1 \\ d\neq 1}} \sum_{\substack{D|x^n-1 \\ D\neq 1}} G_f(\eta_d, \chi_{\delta_D})$.

From definition, $s_0 = q^n$. Also, note that

$$G_f(\eta_1, \chi_{\delta_D}) = \sum_{w\in\mathbb{F}_{q^n}} \eta_1(L_f(w))\chi_{\delta_D}(w) = \sum_{w\in\mathbb{F}_{q^n}} \chi_{\delta_D}(w) = 0,$$

for any divisor D of x^n-1 with $D\neq 1$. In particular, $S_1 = 0$. For the quantities S_2 and S_3, we use some general bounds on character sums.

Lemma 3.4. *([9], Theorem 5.41) Let η be a multiplicative character of $\mathbb{F}_{q^n}$ of order $r>1$ and $f\in\mathbb{F}_{q^n}[x]$ be a monic polynomial of positive degree such that f is not of the form $g(x)^r$ for some $g\in\mathbb{F}_{q^n}[x]$ with degree at least 1. Suppose that e is the number of distinct roots of f in its splitting field over $\mathbb{F}_{q^n}$. For every $a\in\mathbb{F}_{q^n}$,*

$$\left|\sum_{c\in\mathbb{F}_{q^n}} \eta(af(c))\right| \leq (e-1)q^{n/2}.$$

Lemma 3.5. *([2]) Let η be a non trivial multiplicative character of order r and χ be a nontrivial additive character of $\mathbb{F}_{q^n}$. Let f and g be rational functions in $\mathbb{F}_{q^n}(x)$ such that f is not of the form $y\cdot h^r$ for any $y\in\mathbb{F}_{q^n}$ and $h\in\mathbb{F}_{q^n}(x)$, and g is not of the form h^p-h+y for any $y\in\mathbb{F}_{q^n}$ and $h\geq\mathbb{F}_{q^n}(x)$. Then,*

$$\left|\sum_{w\in\mathbb{F}_{q^n}\setminus S} \eta(f(w))\cdot\chi(g(w))\right| \leq (\deg(g)_\infty + m + m' - m'' - 2)q^{n/2},$$

where S is the set of poles of f and g, $(g)_\infty$ is the pole divisor of g, m is the number of distinct zeros and finite poles of f in the algebraic closure $\overline{\mathbb{F}}_q$ of $\mathbb{F}_q$, m' is the number of distinct poles of g (including ∞) and m'' is the number of finite poles of f that are poles or zeros of g.

Write $f(x) = \sum_{i=0}^k a_i x^i$. Note that, since L_f is a q-polynomial, its formal derivative equals a_0. In particular, L_f does not have repeated roots, hence cannot be of the form $y\cdot g(x)^r$ for some $r>1$.

Also, if $f(x)$ divides x^n-1 and has degree k, we know that $L_f = 0$ has exactly q^k roots over $\mathbb{F}_{q^n}$; these roots describe a k-dimensional $\mathbb{F}_q$-vector subspace of $\mathbb{F}_{q^n}$. Finally, notice that $g(x) = x$ cannot be written as $h^p - h - y$ for any rational function $h \in \mathbb{F}_{q^n}(x)$ and $y \in \mathbb{F}_{q^n}$. From Lemma 3.4 we conclude that, for $d > 1$ a divisor of $q^n - 1$,

$$|G_f(\eta_d, \chi_0)| = \left| \sum_{c \in \mathbb{F}_{q^n}} \eta_d(L_f(c)) \right| \leq (q^k - 1)q^{n/2}.$$

From Lemma 3.5, it follows that, for any D divisor of $x^n - 1$ and d a divisor of $q^n - 1$, with $D, d \neq 1$,

$$|G_f(\eta_d, \chi_{\delta_D})| = \left| \sum_{w \in \mathbb{F}_{q^n}} \eta_d(L_f(w))\chi_{\delta_D}(w) \right| \leq (1 + q^k + 1 - 0 - 2)q^{n/2} = q^{n/2+k}.$$

Combining all the previous bounds, we obtain the following:

Theorem 3.6. *Let $f(x)$ be a divisor of $x^n - 1$ of degree k and let n_f be the number of primitive elements of the form $f(x) \circ \alpha$, where α is a normal element. The following holds:*

$$\frac{n_f}{\theta(q^n - 1)\Theta(x^n - 1)} > q^n - q^{n/2+k} W(q^n - 1)W(x^n - 1).$$

In particular, if

$$q^{n/2-k} \geq W(q^n - 1)W(x^n - 1), \tag{5}$$

then there exist primitive k-normal elements in $\mathbb{F}_{q^n}$.

Proof. We have seen that $\frac{n_f}{\theta(q^n-1)\Theta(x^n-1)} = s_0 + S_1 + S_2 + S_3 = q^n + S_2 + S_3$. In particular,

$$\frac{n_f}{\theta(q^n - 1)\Theta(x^n - 1)} \geq q^n - |S_2| - |S_3|.$$

Applying estimates to the sums S_2, S_3, we obtain

$$|S_2| \leq (W(q^n - 1) - 1)(q^k - 1)q^{n/2}$$

and $|S_3| \leq (W(q^n - 1) - 1)(W(x^n - 1) - 1)q^{n/2+k}$, hence

$$|S_1| + |S_2| < q^{n/2+k} W(q^n - 1)W(x^n - 1).$$

In particular, if Eq. (5) holds, $n_f > 0$ and we know that the number of primitive k-normals is at least n_f. This completes the proof. □

Notice that Eq. (5) generalizes the sieve inequality in Lemma 2.8.

4. Existence of primitive k-normals

In this section, we discuss the existence of primitive k-normals in some special extensions of $\mathbb{F}_q$. Essentially, we explore the sieve inequality present in Theorem 3.6 and the result contained in Theorem 2.11.

We have seen that the existence of k-normals is not always ensured for generic values of $1 < k < n-1$: from Lemma 2.4, the number of k-normals is strongly related to the factorization of $x^n - 1$ over $\mathbb{F}_q$. This motivates us to introduce the concept of practical numbers.

Definition 4.1. *(Practical numbers) A positive integer n is said to be $\mathbb{F}_q$-practical if, for any $1 \leq k \leq n-1$, $x^n - 1$ is divisible by a polynomial of degree k over $\mathbb{F}_q$.*

Notice that, if n is $\mathbb{F}_q$-practical, there exist k-normals in $\mathbb{F}_{q^n}$ for any $1 < k < n-1$. This definition arises from the so called *φ-practical numbers*: they are the positive integers n for which $x^n - 1 \in \mathbb{Z}[x]$ is divisible by a polynomial of degree k for any $1 \leq k \leq n-1$. These φ-practical numbers have been extensively studied in many aspects, such as their density over $\mathbb{N}$ and their asymptotic number. In particular, if $s(t)$ denotes the number of φ-practical numbers up to t, according to [11], there exists a constant $C > 0$ such that $\lim_{t\to\infty} \frac{s(t)\cdot \log t}{t} = C$. This shows that the φ-practical numbers behaves like the primes on integers and, in particular, their density in $\mathbb{N}$ is zero.

Notice that the factorization of $x^n - 1$ over $\mathbb{Z}$ also holds over any finite field: we take the coefficients $\pmod p$ and recall that $\mathbb{F}_p \subseteq \mathbb{F}_q$. This shows that any φ-practical number is also $\mathbb{F}_q$-practical. In particular, the number of $\mathbb{F}_q$-practicals up to t has growth at least $\frac{Ct}{\log t}$. The exact growth of the number of $\mathbb{F}_q$-practicals is still an open problem. However, we can find infinite families of such numbers.

Proposition 4.2. *([12], Theorem 4.4) Let q be a power of a prime p and let n be a positive integer such that every prime divisor of n divides $p(q-1)$. Then n is $\mathbb{F}_q$-practical.*

From Theorem 2.11, we obtain the following.

Corollary 4.3. *Let $n = p^2 s$, where s is an $\mathbb{F}_q$-practical number. Then, for any $1 \leq k \leq s$, there exists a primitive, k-normal element of $\mathbb{F}_{q^n}$.*

The previous corollary ensures the existence of primitive k-normals for k in the interval $[1, \frac{n}{p^2}]$. This corresponds to a proportion close to $1/p^2$ of the possible values of k.

In the rest of this paper, we explore Eq. (5) in extensions of degree n, where n is an $\mathbb{F}_q$-practical number. Essentially, we obtain effective bounds on the functions $W(x^n - 1)$ and $W(q^n - 1)$.

4.1. Some estimates for the square-free divisors counting

Here we obtain some estimates on the functions $W(q^n-1)$ and $W(x^n-1)$. We start with some general bounds. The bound $W(x^n-1) \leq 2^n = q^{n\log_q 2}$ is trivial. Also, we have a general bound for $W(q^n-1)$: if $d(q^n-1)$ denotes the number of divisors of q^n-1, $W(q^n-1) \leq d(q^n-1)$. For the number of divisors function, we have the following.

Lemma 4.4. *If $d(m)$ denotes the number of divisors of m, then for all $m \geq 3$,*

$$d(m) \leq m^{\frac{1.5379\log 2}{\log\log m}} < m^{\frac{1.06}{\log\log m}}.$$

Proof. This inequality is a direct consequence of the result in [10]. □

In particular, we can easily obtain $W(q^n-1) < q^{\frac{1.06n}{\log\log(q^n-1)}}$. We now study the special case when n is a power of two.

Lemma 4.5. *Suppose that q is odd. For any $i \geq 1$, if r_0 is an odd prime that divides $q^{2^i}+1$, then $r_0 \equiv 1 \pmod{2^{i+1}}$.*

Proof. Let l be the least positive integer such that $q^l \equiv 1 \pmod{r_0}$. Clearly l divides $\varphi(r_0) = r_0 - 1$. Also, r_0 divides $q^{2^{i+1}}-1$ but not $q^{2^i}-1$, it follows that l divides 2^{i+1} but not 2^i. This shows that $2^{i+1} = l$, hence 2^{i+1} divides r_0-1. □

In particular, we obtain the following.

Proposition 4.6. *For any $q \geq 3$ odd and $t \geq 2$,*

$$W(q^{2^t}-1) < 2q^{\frac{2^t}{t-1}}. \tag{6}$$

Proof. For $1 \leq i < t-1$, set $d_i = q^{2^i}+1$. Note that $W(q^{2^t}-1) \leq W(q^2-1)\cdot\prod_{i=2}^{t-1} W(d_i/2)$. For a fixed $2 \leq i \leq t$, let $s_1 < \cdots < s_{d(i)}$ be the distinct odd primes that divide d_i. Clearly $W(d_i/2) = 2^{d(i)}$. As we have seen, $s_j \equiv 1 \pmod{2^{i+1}}$, hence $s_j > 2^{i+1}$ and then

$$d_i > s_1 \cdots s_{d(i)} > 2^{(i+1)\cdot d(i)},$$

hence $2^{d(i)} < q^{\frac{2^i}{i+1}}$. It follows by induction that $\sum_{i=1}^{t-1} \frac{2^i}{i+1} \leq \frac{2^t}{t-1} - 1$ for $t \geq 2$. Moreover, we have the trivial bound $W(q^2-1) < 2q$. This completes the proof. □

4.2. Applications of Theorem 3.6

The following is straightforward.

Proposition 4.7. *Let n be an $\mathbb{F}_q$-practical number. If k is a positive integer such that*

$$k \leq n\cdot\left(\frac{1}{2} - \frac{1.06}{\log\log(q^n-1)} - \frac{\log 2}{\log q}\right),$$

there exist primitive k-normals in $\mathbb{F}_{q^n}$.

For $h(n,q) = \frac{1}{2} - \frac{1.06}{\log\log(q^n-1)} - \frac{\log 2}{\log q}$, we have $\lim_{q\to\infty} h(n,q) = 1/2$ uniformly on n. In particular, given $\varepsilon > 0$, for q sufficiently large, we can guarantee the existence of k-normals in the interval $[1, (\frac{1}{2} - \varepsilon)n]$ in the case when n is $\mathbb{F}_q$-practical. In general, the bounds on character sums over $\mathbb{F}_{q^n}$ yield the factor $q^{n/2}$. In particular, this result is somehow sharp based on our character sums estimates. However, $h(n,q)$ goes to $1/2$ slowly and we do not have much control on n, since we are assuming that n is $\mathbb{F}_q$-practical. Far from the extreme $n/2$, we have effective results:

Corollary 4.8. *Let $q \geq 420$ be a power of a prime and let $n \geq 420$ be an $\mathbb{F}_q$-practical number. For any $k \in [1, n/4]$, there exist primitive k-normals in $\mathbb{F}_{q^n}$.*

Proof. Note that $h(n,q)$ is an increasing function on q and n. Also, $h(420, 420) > 1/4$ and the result follows. □

The previous corollary gives a wide class of extensions having primitive k-normals in the interval $[1, n/4]$; for instance, we can consider $n = r^t, t \geq 1$, where $r \geq 420$ is a prime dividing $q - 1$. In the special case when n is a power of two, we obtain the following.

Corollary 4.9. *Set $n = 2^t, t \geq 2$ and let q be a power of a prime. Additionally, suppose that $t \geq 9$ and $q \geq 259$ if q is odd. Then, for $k \in [1, n/4]$, there exist primitive k-normal elements in $\mathbb{F}_{q^n}$.*

Proof. Since 2 always divides $p(q-1)$, from Proposition 4.2, 2^t is $\mathbb{F}_q$-practical. For q even, the result follows from Corollary 4.3. Suppose that q is odd. According to Proposition 4.6, $W(q^{2^t} - 1) < 2q^{\frac{2^t}{t-1}}$ and we have the trivial bound $W(x^{2^t} - 1) \leq 2^{2^t}$. We can verify that, for $t \geq 9$ and $q \geq 259$, $2^{2^t+1} q^{\frac{2^t}{t-1}} \leq q^{2^t/4}$ and the result follows from Theorem 3.6. □

5. Conclusions and an additional remark

In this paper we have discussed the existence of primitive k-normal elements over finite fields. We recall some recent results on 1-normal elements and partially extend them to more general k-normals. In particular, we obtain a sieve inequality for the existence of primitive k-normal elements in extension fields that contains k-normals. As an application, we give some families of pairs (q, n) for which we can guarantee the existence of k-normals in $\mathbb{F}_{q^n}$ for $k \in [1, n/4]$: this corresponds to the first quarter of the possible values of k. For q large enough, we can extend the range of k to $[1, (1/2 - \varepsilon)n]$, where ε is close to 0: in other words, we can asymptotically reach the first half of the interval $[1, n]$. As we have pointed out, the typical bounds for our character sums always have the component $q^{n/2}$ and, for this reason, in order to obtain results on primitive k-normals for $k \in [n/2, n]$, we need methods beyond the approach of Lenstra and Schoof or even better character sums estimates.

Here we make a brief discussion on the existence of primitive k-normals for k in the other extreme, i.e., k close to n. We recall that $0 \in \mathbb{F}_{q^n}$ is the only n-normal element. Also, as pointed out in [7], there do no exist primitive $(n-1)$-normals in $\mathbb{F}_{q^n}$: in particular, a primitive element and its conjugates cannot all lie in a "line".

For the special case $k = n - 2$ and $n > 2$, we consider the following: set $q = p$ a prime, $n = p$ and let a be a primitive element of $\mathbb{F}_p$. It can be verified that $f(x) = x^p - x - a$ is irreducible over $\mathbb{F}_p$ and any root $\alpha \in \mathbb{F}_{p^p}$ of $f(x)$ satisfies $(x-1)^2 \circ \alpha = 0$, i.e., α is $(n-2)$-normal. It is conjectured that such an α is always a primitive element of $\mathbb{F}_{p^p}$. This conjecture is verified for every $p \leq 100$ and some few primes $p > 100$. If this conjecture is true, we obtain an interesting example of a primitive element α of "low normalcy" (in the sense that α and their conjugates generates an $\mathbb{F}_q$-vector space of low dimension).

Acknowledgments

This work was conducted during a scholarship supported by the Program CAPES-PDSE (process - 88881.134747/2016-01) at Carleton University.

References

[1] M. Alizadeh, *Some notes on the k-normal elements and k-normal polynomials over finite fields*, Journal of Algebra and its Applications, 16 (2017), 11 pages.

[2] F.N. Castro, C.J. Moreno, *Mixed exponential sums over finite fields*, Proc. Am. Math. Soc. 128 (2000) 2529-2537.

[3] S.D. Cohen, S. Huczynska. *The primitive normal basis theorem - without a computer*, Journal of the London Mathematical Society 67 (2003) 41-56.

[4] S. D. Cohen, D. Hachenberger, *Primitive normal bases with prescribed trace*, Applicable Algebra in Engineering, Communication and Computing 9 (1999) 383-403.

[5] S. D. Cohen, *Primitive elements and polynomials with arbitrary trace.* Discrete Mathematics, 83 (1990), 1-7.

[6] J. von zur Gathen, M. Giesbrecht, *Constructing normal bases in finite fields*, Journal of Symbolic Computation 10 (1990) 547-570.

[7] S. Huczynska, G.L. Mullen, D. Panario, and D. Thomson, *Existence and properties of k-normal elements over finite fields*, Finite Fields Appl. 24 (2013) 170-183.

[8] H.W. Lenstra, R. Schoof, *Primitive normal bases for finite fields*, Mathematics of Computation 48 (1987) 217-231.

[9] R. Lidl, H. Niederreiter, *Finite Fields: Encyclopedia of Mathematics and Its Applications*, vol. 20, 2nd ed. Cambridge University Pres, Cambridge, 1997.

[10] J.L. Nicolas, G. Robin, *Majorations explicites pour le nombre de diviseurs de N*, Canad. Math. Bull. 26 (1983) 485-492.

[11] C. Pomerance, L. Thompson, A. Weingartner, *On integers n for which $x^n - 1$ has a divisor of every degree*, Acta Arith. 175 (2016) 225-243.

[12] L. Reis, *Existence results on K-normal elements over finite fields*, Arxiv (2017) `https://arxiv.org/pdf/1612.05931.pdf`.

[13] L. Reis, D. Thomson, *Existence of primitive 1-normal elements in Finite Fields*, Arxiv (2017) `https://arxiv.org/pdf/1710.06131.pdf`.