

# Nesterov's Acceleration For Approximate Newton

**Haishan Ye**

*yhs12354123@gmail.com*

*Department of Computer Science and Engineering  
Shanghai Jiao Tong University  
800 Dong Chuan Road, Shanghai, China 200240*

**Zhihua Zhang**

*zhzhang@math.pku.edu.cn*

*School of Mathematical Sciences  
Peking University  
Beijing, China 100871*

February 4, 2022

## Abstract

Optimization plays a key role in machine learning. Recently, stochastic second-order methods have attracted much attention due to their low computational cost in each iteration. However, these algorithms might perform poorly especially if it is hard to approximate the Hessian well and efficiently. As far as we know, there is no effective way to handle this problem. In this paper, we resort to Nesterov's acceleration technique to improve the convergence performance of a class of second-order methods called approximate Newton. We give a theoretical analysis that Nesterov's acceleration technique can improve the convergence performance for approximate Newton just like for first-order methods. We accordingly propose an accelerated regularized sub-sampled Newton. Our accelerated algorithm performs much better than the original regularized sub-sampled Newton in experiments, which validates our theory empirically. Besides, the accelerated regularized sub-sampled Newton has good performance comparable to or even better than classical algorithms.

## 1. Introduction

Optimization has become an increasingly popular issue in machine learning. Many machine learning models can be reformulated as the following optimization problems:

$$\min_{x \in \mathbb{R}^d} F(x) = \frac{1}{n} \sum_{i=1}^n f_i(x). \quad (1)$$

where each  $f_i$  is the loss with respect to (w.r.t.) the  $i$ -th training sample. There are many examples such as logistic regressions, smoothed support vector machines, neural networks, and graphical models.

In the era of big data, large-scale optimization algorithms have become an important challenge. The stochastic gradient descent algorithm (SGD) has been widely employed to reduce the computational cost [5, 14, 20]. However, SGD has poor convergence property. Hence, many variants have been proposed to improve the convergence rate of SGD [11, 22, 23, 27].

For the first-order methods which only make use of the gradient information, Nesterov's acceleration technique is a very useful tool [17]. It greatly improves the convergence of gradient descent [17], proximal gradient descent [2, 18], and stochastic gradient with variance reduction [1, 13], etc.

Recently, second-order methods have also received great attention due to their high convergence rate. However, conventional second-order methods are very costly because they take heavy computational cost to obtain the Hessian matrices. To conquer this weakness, one proposed a sub-sampled Newton which only samples a subset of functions  $f_i$  randomly to construct a sub-sampled Hessian [21, 3, 25]. Pilanci and Wainwright [19] applied the sketching technique to alleviate the computational burden of computing Hessian and brought up *sketch Newton*. Regularized sub-sampled Newton methods were also devised to deal with the ill-condition problem [7, 21].

In the latest work, Ye et al. [26] cast these stochastic second-order procedures into a so-called *approximate Newton* framework. They showed that if approximate Hessian  $H^{(t)}$  satisfies

$$(1 - \pi)\nabla^2 F(x^{(t)}) \preceq H^{(t)} \preceq (1 + \pi)\nabla^2 F(x^{(t)}), \quad (2)$$

where  $0 < \pi < 1$ , then approximate Newton converges with rate  $\pi$ . If  $H^{(t)}$  is a poor approximation like  $\pi = 1 - 1/\kappa$ , where  $\kappa$  is the condition number of object function  $F(x)$ , then approximate Newton has the same convergence rate with gradient descent.

Since approximate Newton converges with a linear rate, it is natural to ask whether approximate Newton can be accelerated just like gradient descent. If it can be accelerated, can the convergence rate be promoted to  $1 - \sqrt{1 - \pi}$  compared to original  $\pi$ ? In this paper, we aim to introduce Nesterov's acceleration technique to improve the performance of second-order methods, specifically approximate Newton.

We summarize our work and contribution as follows:

- First, we introduce Nesterov's acceleration technique to improve the convergence rate of the stochastic second-order methods (approximate Newton). This acceleration is very important especially when  $n$  and  $d$  are close to each other and object function in question is ill-conditioned. In these cases, it is very hard to construct a good approximate Hessian with low cost.
- Our theoretical analysis shows that by Nesterov's acceleration, the convergence rate of approximate Newton can be improved to  $1 - \sqrt{1 - \pi}$  from original rate  $\pi$  where  $0 < \pi < 1$  when the object function is quadratic. For general smooth convex functions, we also show that the similar acceleration also holds when the initial point is close to the optimal point.
- We propose Accelerated Regularized Sub-sampled Newton. Compared with classical stochastic first-order methods, our algorithm shows competitive or even better performance. This demonstrates the efficiency of the accelerated second-order method. Our experimental study shows that Nesterov's acceleration technique can improve approximate Newton methods effectively. Our experiments also reveal a fact that adding curvature information properly can always improve the algorithm's convergence performance.

## 2. Notation and Preliminaries

We first introduce notation that will be used in this paper. Then, we give some properties of object function that will be used.

### 2.1 Notation

Given a matrix  $A = [a_{ij}] \in \mathbb{R}^{m \times n}$  of rank  $\ell$  and a positive integer  $k \leq \ell$ , its SVD is given as  $A = U\Sigma V^T = U_k \Sigma_k V_k^T + U_{\setminus k} \Sigma_{\setminus k} V_{\setminus k}^T$ , where  $U_k$  and  $U_{\setminus k}$  contain the left singular vectors of  $A$ ,  $V_k$  and  $V_{\setminus k}$  contain the right singular vectors of  $A$ , and  $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_\ell)$  with  $\sigma_1 \geq \sigma_2 \geq \dots \geq \sigma_\ell > 0$  are the nonzero singular values of  $A$ . Additionally,  $\|A\| \triangleq \sigma_1$  is the spectral norm. If  $A$  is positive semidefinite, then  $U = V$  and the eigenvalue decomposition of  $A$  is the same to singular value

decomposition. It also holds that  $\lambda_i(A) = \sigma_i(A)$ , where  $\lambda_i(A)$  is the  $i$ -th largest eigenvalue of  $A$ . Let  $\lambda_{\max}(A)$  and  $\lambda_{\min}(A)$  denote the largest and smallest eigenvalue of  $A$ , respectively.

## 2.2 Assumptions

In this paper, we focus on the problem described in Eqn. (1). Moreover, we will make the following two assumptions.

**Assumption 1** The objective function  $F$  is  $\mu$ -strongly convex, that is,

$$F(y) \geq F(x) + [\nabla F(x)]^T(y - x) + \frac{\mu}{2}\|y - x\|^2, \text{ for } \mu > 0.$$

**Assumption 2**  $\nabla F(x)$  is  $L$ -Lipschitz continuous, that is,

$$\|\nabla F(x) - \nabla F(y)\| \leq L\|y - x\|, \text{ for } L > 0.$$

By Assumptions 1 and 2, we define the condition number of function  $F(x)$  as:  $\kappa \triangleq \frac{L}{\mu}$ .

Besides, we will also use the notion of Lipschitz continuity of  $\nabla^2 F(x)$  in this paper. We say  $\nabla^2 F(x)$  is  $\hat{L}$ -Lipschitz continuous if

$$\|\nabla^2 F(x) - \nabla^2 F(y)\| \leq \hat{L}\|y - x\|, \text{ for } \hat{L} > 0.$$

## 2.3 Row Norm Squares Sampling

The row norm squares sampling matrix  $S = D\Omega \in \mathbb{R}^{s \times n}$  w.r.t.  $A \in \mathbb{R}^{n \times d}$  is determined by sampling probability  $p_i$ , a sampling matrix  $\Omega \in \mathbb{R}^{s \times n}$  and a diagonal rescaling matrix  $D \in \mathbb{R}^{s \times s}$ . The sampling probability is defined as

$$p_i = \frac{\|A_{i,:}\|^2}{\|A\|_F^2},$$

where  $A_{i,:}$  means the  $i$ -th row of  $A$ . We construct  $S$  as follows. For every  $j = 1, \dots, s$ , independently and with replacement, pick an index  $i$  from the set  $\{1, 2, \dots, m\}$  with probability  $p_i$  and set  $\Omega_{ji} = 1$  and  $\Omega_{jk} = 0$  for  $k \neq i$  as well as  $D_{jj} = 1/\sqrt{p_i s}$ .

Row norm squares sampling matrix has the following important property.

**Theorem 1** Let  $S \in \mathbb{R}^{s \times n}$  be a row norm squares sampling matrix w.r.t.  $A \in \mathbb{R}^{n \times d}$ , then it holds that

$$\mathbb{E}\|A^T S^T S A - A^T A\| \leq \left( \sqrt{\frac{4 \cdot \text{sr}(A) \cdot \log(2d)}{s}} + \frac{2 \cdot \text{sr}(A) \cdot \log(2d)}{3s} \right) \cdot \|A\|^2$$

## 2.4 Randomized sketching matrices

We first give the definition of the  $\epsilon$ -subspace embedding property. Then we list some useful types of randomized sketching matrices including Gaussian sketching [10, 12], random sampling [6], count sketch [4, 16, 15].

**Definition 2**  $S \in \mathbb{R}^{s \times m}$  is said to be an  $\epsilon$ -subspace embedding matrix for any fixed matrix  $A \in \mathbb{R}^{m \times d}$ , if  $\|SAx\|^2 = (1 \pm \epsilon)\|Ax\|^2$  (i.e.,  $(1 - \epsilon)\|Ax\|^2 \leq \|SAx\|^2 \leq (1 + \epsilon)\|Ax\|^2$ ) for all  $x \in \mathbb{R}^d$ .

**Gaussian sketching matrix:** The most classical sketching matrix is the Gaussian sketching matrix  $S \in \mathbb{R}^{s \times m}$  with i.i.d normal random entries with mean 0 and variance  $1/s$ . Because of well-known concentration properties of Gaussian random matrices [24], they are very attractive. Besides,  $s = O(d/\epsilon^2)$  is enough to guarantee the  $\epsilon$ -subspace embedding property for any fixed matrix  $A \in \mathbb{R}^{m \times d}$ . And  $s = O(d/\epsilon^2)$  is the tightest bound in known types of sketching matrices. However, Gaussian random matrices are dense, so it is costly to compute  $SA$ .

**Count sketch matrix:** Count sketch matrix  $S \in \mathbb{R}^{s \times m}$  is of the form that there is only one non-zero entry uniformly sampling from  $\{1, -1\}$  in each column [4]. Hence it is very efficient to compute  $SA$ , especially when  $A$  is a sparse matrix. To achieve an  $\epsilon$ -subspace embedding property for  $A \in \mathbb{R}^{m \times d}$ ,  $s = O(d^2/\epsilon^2)$  is sufficient [15, 24].

Other types of sketching matrices like Sub-sampled Randomized Hadamard Transformation and detailed properties of sketching matrices can be found in the survey [24].

### 3. Accelerated Approximate Newton

In practice, it is common that the problem is ill-condition and the data size  $n$  and data dimension  $d$  are close to each other. Therefore, conventional Sketch Newton, sub-sampled Newton and regularized sub-sampled Newton can not construct a good approximate Hessian efficiently. However, Ye et al. [26] showed that a poor approximate Hessian will lead to a slow converge rate. In the extreme case, these approximate Newton methods have the same convergence rate with gradient descent. To improve the convergence property of approximate Newton when the Hessian can only be approximate poorly, we resort to Nesterov's acceleration technique and propose *accelerated approximate Newton*. We summarize the algorithmic procedure of accelerated approximate Newton as follows.

First, we construct an approximate Hessian  $[H^{(t)}]$  satisfies

$$(1 - \pi)[\nabla^2 F(y^{(t)})]^{-1} \preceq \mathbb{E}([H^{(t)}]^{-1}) \preceq [\nabla^2 F(y^{(t)})]^{-1}, \quad (3)$$

where  $0 < \pi < 1$ . This condition is a little stronger than the one of approximate Newton which satisfies Eqn. (2). We will see that Condition (3) can be easily satisfied in practice in next section.

And we update sequence  $x^{(t)}$  as follows,

$$\begin{cases} y^{(t+1)} = (1 + \theta)x^{(t)} - \theta x^{(t-1)}, \\ x^{(t+1)} = y^{(t+1)} - [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}), \end{cases} \quad (4)$$

where  $\theta$  is chosen in terms of the value of  $\pi$ . We can see that the iteration (4) is much like the update procedure of Nesterov's accelerated gradient descent but replacing step size with  $H^{-1}$ . Besides, when  $\theta = 0$ , the above update procedure reduces to the update step of approximate Newton [26]. Thus, we refer a class of methods satisfying Eqn. (3) and algorithm procedure (4) as *accelerated approximate Newton*.

Similar to approximate Newton method, we can update  $x^{(t+1)}$  with a direction vector  $p^{(t)}$  which is an inexact solution of following problem

$$\frac{1}{2} p^T H^{(t)} p - p^T \nabla F(y^{(t)}). \quad (5)$$

There are different ways to solve problem (5) like conjugate gradient.

#### 3.1 Theoretical Analysis

Because a general smooth convex object function can be approximated by quadratic functions in a region close to optimal point, we show the convergence properties of Algorithm 1 when applied to a quadratic object function. The following lemma (Lemma 3) gives the detailed reason why we can only analyze convergence properties of Algorithm 1 applied to a quadratic object function.

Furthermore, we will first give the convergence analysis of accelerated approximate Newton which satisfies Eqn. (3) and of algorithm procedure (4) in Theorem 4.

**Lemma 3** *Let Assumptions 1 and 2 hold. Suppose that  $\nabla^2 F(x)$  exists and is continuous in a neighborhood of a minimizer  $x^*$ . Let  $H^{(t)}$  be an approximation of  $\nabla^2 F(y^{(t)})$ . Consider the iteration (4). If  $x^{(t)}$  is sufficient close to  $x^*$  then we have the following result*

$$\begin{aligned}
& [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t+1)}) \\
&= \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) \\
&\quad \times \left( (1 + \theta) [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t)}) - \theta [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t-1)}) \right) \\
&\quad + o(\nabla F(x^{(t)})) + o(\nabla F(x^{(t-1)})).
\end{aligned}$$

Furthermore, if  $\nabla^2 F(x)$  is  $\hat{L}$ -Lipschitz continuous, then the above result holds whenever  $x^{(t)}$  satisfies

$$\|x^{(t)} - x^*\| \leq \frac{o(1)}{\hat{L}}.$$

Lemma 3 shows that the convergence property of iteration (4) is mainly determined by  $I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}}$  and  $\theta$ . Lemma 3 also describes such a fact that if  $x^{(t-1)}$  and  $x^{(t)}$  are sufficient close to  $x^*$ , that is,  $o(F(x^{(t)}))$  and  $o(F(x^{(t-1)}))$  are very small, then the convex function can be well approximated by a quadratic function. Therefore, we will demonstrate the convergence analysis of accelerated approximate Newton on the strongly convex quadratic functions. Because the Hessian of a quadratic function  $F(x)$  is a constant matrix, that is the Lipschitz constant of  $\nabla^2 F(x)$  is zero. Hence, the result of Lemma 3 degenerates to

$$\begin{aligned}
& [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t+1)}) \\
&= \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) \\
&\quad \times \left( (1 + \theta) [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t)}) - \theta [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t-1)}) \right).
\end{aligned}$$

This equation describes a linear dynamic system which contains the convergence property of iteration (4).

**Theorem 4** Let  $F(x)$  be a quadratic function with Assumption 1 and 2 holding. Let  $H^{(t)}$  be an approximation of  $\nabla^2 F(y^{(t)})$  satisfying Eqn. (3) and  $\mathbb{E}([H^{(t)}]^{-1})$  is a constant matrix for all  $t$ . We set  $\theta = \frac{1-\sqrt{1-\pi}}{1+\sqrt{1-\pi}} - \epsilon_0$  with  $0 < \epsilon_0 < 1$ .  $T_1$  is a matrix of the form

$$T_i = \begin{bmatrix} (1 + \theta)\pi & -\theta\pi \\ 1 & 0 \end{bmatrix}.$$

Let  $S_1$  and  $q$  be the eigenmatrix and larger eigenvalue of  $T_1$  respectively. Then  $q$  is of the value  $q = 1 - \sqrt{1 - \pi} + O(\sqrt{\epsilon_0})$ . And  $c$  is a constant defined by  $c_1 = \|S_1\| \cdot \|S_1^{-1}\|$ . Let direction vector  $p^{(t)}$  satisfy

$$\left\| H^{(t)} p^{(t)} - \nabla F(y^{(t)}) \right\| \leq \frac{\epsilon_1}{12c_1\sqrt{\kappa}} \left\| \nabla F(y^{(t)}) \right\|, \quad (6)$$

where  $\kappa$  is the condition number of the Hessian matrix  $\nabla^2 F(x^*)$ . Then after  $t$  iterations, Algorithm 1 has

$$\left\| \mathbb{E} \begin{bmatrix} M_* \nabla F(x^{(t+1)}) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} \right\| \leq 2c_1 p^t \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\|,$$

with  $p = q + \epsilon_1$  and  $M_* = [\nabla^2 F(x^*)]^{-1/2}$ .

From Theorem 4, we can see that if we choose  $\theta = \frac{1-\sqrt{1-\pi}}{1+\sqrt{1-\pi}} - \epsilon_0$  and  $\epsilon_1 = 0$ , then the convergence rate of Algorithm 1 is  $1 - \sqrt{1 - \pi} + O(\sqrt{\epsilon_0})$  in contrast to  $\pi$  of the conventional regularized sub-sampled Newton method. If we choose  $H = (1/L)I$  and  $\theta = \frac{\sqrt{L}-\sqrt{\mu}}{\sqrt{L}+\sqrt{\mu}}$ , where  $L = \lambda_{\max}(\nabla^2 F(x^*))$  and  $\mu = \lambda_{\min}(\nabla^2 F(x^*))$ , then Theorem 4 shows that the convergence rate of Nesterov's acceleration for the least square regression problem is  $1 - \sqrt{\mu/L} + O(\sqrt{\epsilon_0})$ .

---

**Algorithm 1** Accelerated Regularized Sub-sample Newton (ARSSN).

---

1: **Input:**  $x^{(0)}$  and  $x^{(1)}$  are initial points sufficient close to  $x^*$ .  $\theta$  is the acceleration parameter.  
2: **for**  $t = 1, \dots$  until termination **do**  
3:   Select a sample set  $\mathcal{S}$  of size  $|\mathcal{S}|$  by random sampling and construct  $H^{(t)}$  of the form (7) satisfying Eqn. (3);  
4:    $y^{(t+1)} = (1 + \theta)x^{(t)} - \theta x^{(t-1)}$ ;  
5:   Obtain the direction vector  $p^{(t)}$  by solving problem .  
6:    $x^{(t+1)} = y^{(t+1)} - p^{(t+1)}$ .  
7: **end for**

---

#### 4. Accelerated Regularized Sub-sampled Newton

In practice, it is common that data size  $n$  and data dimension  $d$  are close to each other. Conventional Sketch Newton method is not suitable because sketching size  $|\mathcal{S}|$  will be less than  $d$ , and the sketched Hessian is not invertible. Hence, adding a proper regularizer is a good approach. On the other hand, sub-sampled Newton method needs lots of samples when the problem is ill-conditioned. Hence, regularized sub-sample Newton methods are proposed [21]. However, Ye et al. [26] showed that regularized sub-sample Newton will converge slowly as the sample size decreases. In the extreme case, regularized sub-sample Newton has the same convergence rate with gradient descent.

To conquer the weakness of slow convergence rate of regularized sub-sample Newton, we resort to Nesterov's acceleration technique and propose accelerated regularized sub-sample Newton method in Algorithm 1. Just like conventional second-order methods, we assume initial point  $x^{(0)}$  and  $x^{(1)}$  are sufficient close to optimal point  $x^*$  in our algorithms.

In Algorithm 1,  $H^{(t)}$  is an approximation of  $\nabla^2 F(y^{(t)})$  constructed by random sampling.  $H^{(t)}$  is of the following structure

$$H^{(t)} = \hat{H}^{(t)} + \alpha^{(t)} I. \quad (7)$$

$\hat{H}^{(t)}$  is the sub-sampled Hessian. And  $\alpha^{(t)} I$  is the regularizer.  $H^{(t)}$  also satisfies Algorithm 1 specifies the way to construct the approximate Hessian by random sampling and is a kind of accelerated approximate Newton. Therefore, the convergence property of Algorithm 1 can be analyzed by Theorem 4.

First, we consider the following case that the Hessian of  $\nabla^2 F(x)$  has the following structure

$$\nabla^2 F(x) = B(x)^T B(x), \quad B(x) \in \mathbb{R}^{n \times d}. \quad (8)$$

For this case, we construct approximate Hessian's as

$$H^{(t)} = [S^{(t)} B^{(t)}]^T S^{(t)} B^{(t)} + \alpha^{(t)} I. \quad (9)$$

$S^{(t)}$  is a random sampling matrix or other sketching matrix. And  $\alpha^{(t)}$  is a regularizer scaler.

**Theorem 5** Let  $F(x)$  be a quadratic function with Assumption 1 and 2 holding.  $\nabla^2 F(x)$  satisfies Eqn. (8). Given sample size parameter  $0 < c < 1$ ,  $S^{(t)} \in \mathbb{R}^{s \times n}$  is a row norm squares sampling matrix w.r.t.  $B^{(t)}$  with  $s = O(c^{-2} \cdot \text{sr}(B^{(t)}) \log d)$ . And set regularizer  $\alpha = c \|B^{(t)}\|^2$ . Construct the approximate Hessian  $H^{(t)} = [S^{(t)} B^{(t)}]^T S^{(t)} B^{(t)} + \alpha^{(t)} I$ . Let  $\pi = \frac{2c\kappa}{1+2c\kappa}$ , and set parameters as Theorem 4, Algorithm 1 converges as

$$\left\| \mathbb{E} \begin{bmatrix} M_* \nabla F(x^{(t+1)}) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} \right\| \leq 2c_1 \left( 1 - \frac{1}{\sqrt{1+2c\kappa}} + O(\sqrt{\epsilon_0}) + \epsilon_1 \right)^t \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\|,$$

with  $M_* = [\nabla^2 F(x^*)]^{-1/2}$  and  $\epsilon_0, \epsilon_1$  defined in Theorem 4.

**Proof** For notation convenience, we will omit superscript and just use  $S$ ,  $B$  and  $\alpha$  instead of  $S^{(t)}$ ,  $B^{(t)}$  and  $\alpha^{(t)}$ . By Theorem 1 and omit the high order term, we have

$$\mathbb{E}\|B^T B - B^T S^T S B\| \leq c\|B\|^2.$$

Hence, in expectation, we have

$$\begin{aligned} & |y^T (B^T B - B^T S^T S B) y| \leq c\|B\|^2 \cdot \|y\|^2 \\ \Rightarrow & y^T B^T S^T S B y - c\|B\|^2 \cdot \|y\|^2 \leq y^T B^T B y \leq y^T B^T S^T S B y + c\|B\|^2 \cdot \|y\|^2 \\ \Rightarrow & y^T B^T B y \leq y^T H^{(t)} y \leq y^T B^T B y + 2c\|B\|^2 \cdot \|y\|^2 \\ \Rightarrow & y^T [B^T B]^{-1} y \geq y^T [H^{(t)}]^{-1} y \geq y^T (B^T B + 2c\|B\|^2 \cdot I)^{-1} y \end{aligned}$$

Furthermore, we have

$$\begin{aligned} & y^T B^T B y + 2c\|B\|^2 \cdot \|y\|^2 \leq y^T B^T B y + \frac{2c\|B\|^2}{\sigma_{\min}(B)^2} y^T B^T B y \\ \Rightarrow & B^T B + 2c\|B\|^2 \cdot I \preceq (1 + 2c\kappa) B^T B \\ \Rightarrow & \left(1 - \frac{2c\kappa}{1 + 2c\kappa}\right) B^T B \preceq (B^T B + 2c\|B\|^2 \cdot I)^{-1} \end{aligned}$$

Thus, we have

$$\left(1 - \frac{2c\kappa}{1 + 2c\kappa}\right) \cdot [\nabla^2 F(x^*)]^{-1} \preceq \mathbb{E}([H^{(t)}]^{-1}) \preceq [\nabla^2 F(x^*)]^{-1}.$$

Then final convergence property can be obtained by Theorem 4 using  $\pi = \frac{2c\kappa}{1+2c\kappa}$ . ■

Then, we consider the case that each  $f_i(x)$  and  $F(x)$  in (1) have the following properties:

$$\max_{1 \leq i \leq n} \|\nabla^2 f_i(x)\| \leq K < \infty, \quad (10)$$

$$\lambda_{\min}(\nabla^2 F(x)) \geq \sigma > 0. \quad (11)$$

In this case, we do not need the Hessian of the special form (9) but need that each individual Hessian is upper bounded. We construct the approximate Hessian  $H^{(t)}$  just by uniformly sampling as follows

$$H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)}) + \alpha^{(t)} I \quad (12)$$

**Theorem 6** Let  $F(x)$  be a quadratic function with Assumption 1 and 2 holding. And Eqns (10) and (11) also hold. We construct the approximate Hessian  $H^{(t)}$  as Eqn. (12) by uniformly sampling with sample size  $|\mathcal{S}| = O(c^{-2} K^2 \log d)$ . And set regularizer  $\alpha^{(t)} = c\|\nabla^2 F(x^{(t)})\|^2$ . Set parameters as Theorem 4, Algorithm 1 converges as

$$\left\| \mathbb{E} \begin{bmatrix} M_* \nabla F(x^{(t+1)}) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} \right\| \leq 2c_1 \left( 1 - \frac{1}{\sqrt{1 + 2c\kappa}} + O(\sqrt{\epsilon_0}) + \epsilon_1 \right)^t \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\|,$$

with  $M_* = [\nabla^2 F(x^*)]^{-1/2}$  and  $\epsilon_0, \epsilon_1$  defined in Theorem 4.

**Proof** Consider  $|\mathcal{S}|$  i.i.d random matrices  $H_j^{(t)}, j = 1, \dots, |\mathcal{S}|$  sampled uniformly. Then, we have  $\mathbb{E}(H_j^{(t)}) = \nabla^2 F(x^{(t)})$  for all  $j = 1, \dots, |\mathcal{S}|$ . By (10) and the positive semi-definite property of  $H_j^{(t)}$ , we have  $\lambda_{\max}(H_j^{(t)}) \leq K$  and  $\lambda_{\min}(H_j^{(t)}) \geq 0$ .

We define random matrices  $X_j = H_j^{(t)} - \nabla^2 F(x^{(t)})$  for all  $j = 1, \dots, |\mathcal{S}|$ . We have  $\mathbb{E}[X_j] = 0$ ,  $\|X_j\| \leq 2K$  and  $\|X_j\|^2 \leq 4K^2$ . By the matrix Bernstein inequality, we have

$$\mathbb{E}(\|H^{(t)} - \nabla^2 F(x^{(t)})\|) \leq \sqrt{\frac{2 \cdot 4K^2 \log 2d}{|\mathcal{S}|}} + \frac{2K \log 2d}{3|\mathcal{S}|}.$$

When the sample size  $|\mathcal{S}| = O(c^{-2}K^2 \log d)$ ,  $\|H^{(t)} - \nabla^2 F(x^{(t)})\| \leq cK$  holds in expectation, that is

$$\mathbb{E}\|H^{(t)} - \nabla^2 F(x^{(t)})\| \leq c\nabla^2 F(x^{(t)}).$$

For notation convenience, we will omit superscript. Hence, in expectation, we have

$$\begin{aligned} |y^T (H - \nabla^2 F(x))y| &\leq c\|\nabla^2 F(x)\| \cdot \|y\|^2 \\ \Rightarrow y^T H Y y - c\|\nabla^2 F(x)\| \cdot \|y\|^2 &\leq y^T \nabla^2 F(x)y \leq y^T H y + c\|\nabla^2 F(x)\| \cdot \|y\|^2 \\ \Rightarrow y^T \nabla^2 F(x)y &\leq y^T H y \leq y^T \nabla^2 F(x)y + 2c\|\nabla^2 F(x)\| \cdot \|y\|^2 \\ \Rightarrow y^T [\nabla^2 F(x)]^{-1}y &\geq y^T [H]^{-1}y \geq y^T (\nabla^2 F(x) + 2c\|\nabla^2 F(x)\| \cdot I)^{-1}y \end{aligned}$$

Furthermore, we have

$$\begin{aligned} y^T \nabla^2 F(x)y + 2c\|\nabla^2 F(x)\| \cdot \|y\|^2 &\leq y^T \nabla^2 F(x)y + \frac{2c\|\nabla^2 F(x)\|}{\sigma_{\min}(\nabla^2 F(x))} y^T \nabla^2 F(x)y \\ \Rightarrow \nabla^2 F(x) + 2c\|\nabla^2 F(x)\| \cdot I &\preceq (1 + 2c\kappa) \nabla^2 F(x) \\ \Rightarrow \left(1 - \frac{2c\kappa}{1 + 2c\kappa}\right) \nabla^2 F(x) &\preceq (\nabla^2 F(x) + 2c\|\nabla^2 F(x)\| \cdot I)^{-1} \end{aligned}$$

Thus, we have

$$\left(1 - \frac{2c\kappa}{1 + 2c\kappa}\right) \cdot [\nabla^2 F(x^*)]^{-1} \preceq \mathbb{E}([H^{(t)}]^{-1}) \preceq [\nabla^2 F(x^*)]^{-1}.$$

Then final convergence property can be obtained by Theorem 4 using  $\pi = \frac{2c\kappa}{1+2c\kappa}$ . ■

#### 4.1 Fast Sub-problem Solver

In practice, we will sub-sample a small subset of samples, that is sample size  $s$  is much smaller than data dimension  $d$ . And Eqn. (9) becomes

$$H^{(t)} = \tilde{B}^T \tilde{B} + \alpha^{(t)} I \quad (13)$$

where  $\tilde{B} = S^{(t)} B^{(t)} \in \mathbb{R}^{s \times d}$  with  $s \ll d$ . Therefore, by Sherman-Woodbury identity formula, we can get the exact solution of Eqn. (5) by

$$p^{(t)} = [H^{(t)}]^{-1} \nabla F(y^{(t)}) = (\alpha^{(t)})^{-1} \nabla F(y^{(t)}) - \tilde{B}^T (\tilde{B} \tilde{B}^T + \alpha^{(t)} I)^{-1} \tilde{B} \nabla F(y^{(t)}) / \alpha^{(t)} \quad (14)$$

And  $p^{(t)}$  can be obtain in  $O(ds^2 + s^3)$  time. Since Theorem 5 shows that the direction vector can be an approximate solution of Eqn. (5), we can get an inexact solution in a more efficient manner.

The main computational burden of Eqn. (14) is the matrix product of  $\tilde{B}$  and  $\tilde{B}^T$ . Rather than Sherman-Woodbury identity formula, we can approximate the matrix inversion and get rid of the matrix multiplication by conjugate gradient. The main computational cost of conjugate gradient to compute  $p^{(t)}$  is the matrix vector multiplication. This is very suitable for sparse dataset. However, its computational complexity depends on  $\sqrt{\kappa(H^{(t)})}$  linearly. This may be expensive when the condition number of  $H^{(t)}$  is large.

Therefore, instead of using conjugate gradient method directly, we resort to preconditioned conjugate gradient method (Algorithm 5) to obtain an approximation of  $(\tilde{B}\tilde{B}^T + \alpha I)^{-1}g$  in Eqn. (14) with  $g = \tilde{B} \cdot \alpha^{-1} \nabla F(y^{(t)})$ . Because of  $\tilde{B} \in \mathbb{R}^{s \times d}$  with  $s \ll d$ , we can use sketching tools to construct a good preconditioner very efficiently. We describe the detailed algorithm in Algorithm 2. And we have the following result.

**Theorem 7** *Let the approximate Hessian  $H^{(t)}$  be of form Eqn. (13) with  $\tilde{B} \in \mathbb{R}^{s \times d}$  and  $s \ll d$ . Set iteration number  $T$  as*

$$T = \log \left( \frac{\alpha^{-\frac{3}{2}} \|H^{(t)}\|^{\frac{3}{2}} c_1 \sqrt{\kappa}}{\epsilon_1} \right).$$

where  $c_1$  and  $\kappa$  are defined in Theorem 4.  $p^{(t)}$  is returned from Algorithm 2, then we have

$$\|H^{(t)}p^{(t)} - \nabla F(y^{(t)})\| \leq \frac{\epsilon_1}{12c_1\sqrt{\kappa}} \|\nabla F(y^{(t)})\|.$$

And the computation complexity of computing  $p^{(t)}$  is  $O(Tds + s^3)$ .

**Proof** First, by the property of subspace embedding property (Definition 2) and  $G$  is a  $1/3$ -subspace embedding matrix w.r.t  $\tilde{B}$ , we have

$$\left(1 - \frac{1}{3}\right) \tilde{B}\tilde{B}^T \preceq \tilde{B}GG^T\tilde{B}^T \preceq \left(1 + \frac{1}{3}\right) \tilde{B}\tilde{B}^T.$$

With a simple transformation, we obtain

$$\left(1 - \frac{1}{2}\right) \tilde{B}GG^T\tilde{B}^T \preceq \tilde{B}\tilde{B}^T \preceq \left(1 + \frac{1}{2}\right) \tilde{B}GG^T\tilde{B}^T.$$

Thus sketched matrix  $P$  is a good preconditioner. And by the convergence property of preconditioned conjugate gradient method (Lemma 8), after Algorithm 5 running  $T$  iterations, we have

$$\begin{aligned} \|A^{-1}g - q\|_A &\leq \left(\frac{1}{2}\right)^T \|A^{-1}g\|_A \\ \Rightarrow \sqrt{\lambda_{\min}(A)} \cdot \|A^{-1}g - q\| &\leq \left(\frac{1}{2}\right)^T \sqrt{g^T A^{-1}g} \end{aligned}$$

where  $A = \tilde{B}\tilde{B}^T + \alpha I$  and  $g = \tilde{B} \cdot \alpha^{-1} \nabla F(y^{(t)})$ . We also have

$$g^T A^{-1}g = \alpha^{-2} \cdot \nabla F(y^{(t)})^T \tilde{B}^T (\tilde{B}\tilde{B}^T + \alpha I)^{-1} \tilde{B} \nabla F(y^{(t)}) \leq \alpha^{-2} \|\nabla F(y^{(t)})\|^2.$$

The second inequality is because  $\tilde{B}^T (\tilde{B}\tilde{B}^T + \alpha I)^{-1} \tilde{B}$  is positive definite and its largest eigenvalue is smaller than 1. Thus, we have

$$\|A^{-1}g - q\| \leq \left(\frac{1}{2}\right)^T \alpha^{-\frac{3}{2}} \|\nabla F(y^{(t)})\|.$$

---

**Algorithm 2** Fast sub-problem solver.

---

- 1: **Input:** Matrix  $\tilde{B}$ , regularizer  $\alpha$ , gradient  $\nabla F(y^{(t)})$ , and iteration number  $T$ ;
  - 2: Compute  $g = \tilde{B} \cdot \alpha^{-1} \nabla F(y^{(t)})$ . Construct a 1/3-subspace embedding matrix  $G$  w.r.t  $\tilde{B}$  and compute preconditioner matrix  $P = \tilde{B}GG^T\tilde{B}$ .
  - 3: Using  $A = \tilde{B}\tilde{B}^T + \alpha I$ ,  $T$ , and  $P$  as input of Algorithm 5, we get a approximate solution vector  $q$  by Algorithm 5.
  - 4: **Output:**  $p^{(t)} = \alpha^{-1} \nabla F(y^{(t)}) - \tilde{B}^T q$ .
- 

Furthermore, we have

$$\begin{aligned}
\|H^{(t)}p^{(t)} - \nabla F(y^{(t)})\| &\leq \|H^{(t)}\| \cdot \|p^{(t)} - [H^{(t)}]^{-1} \nabla F(y^{(t)})\| \\
&= \|H^{(t)}\| \cdot \|\alpha^{-1} \nabla F(y^{(t)}) - \tilde{B}^T q - (\alpha^{-1} \nabla F(y^{(t)}) - \tilde{B}^T A^{-1} g)\| \\
&\leq \left(\frac{1}{2}\right)^T \alpha^{-\frac{3}{2}} \|H^{(t)}\| \cdot \|\tilde{B}\| \cdot \|\nabla F(y^{(t)})\| \\
&\leq \left(\frac{1}{2}\right)^T \alpha^{-\frac{3}{2}} \|H^{(t)}\|^{\frac{3}{2}} \cdot \|\nabla F(y^{(t)})\|
\end{aligned}$$

Since we set iteration number  $T$  to be

$$T = \log \left( \frac{\alpha^{-\frac{3}{2}} \|H^{(t)}\|^{\frac{3}{2}} c_1 \sqrt{\kappa}}{\epsilon_1} \right),$$

we obtain the result.

Because the main operation in preconditioned conjugate gradient is matrix-vector production, the total cost is  $O(Tds + s^3)$ . ■

From Theorem 7, we can see that iteration number  $T$  is only of order  $O(\log \kappa)$ . Therefore the time of computing a  $p^{(t)}$  by Algorithm 2 is cheap than by Eqn. (14) when  $O(\log \kappa) < s$ . And this is common in practice.

## 5. Experiments

In Section 3.1, we have shown that Nesterov's acceleration technique can improve the performance of approximate Newton method theoretically. In this section, we will validate our theory empirically. In particular, we first compare accelerated regularized sub-sampled Newton (Algorithm 1 referred as ARSSN) with regularized sub-sampled Newton (Algorithm 3 referred as RSSN) on the ridge regression whose objective function is a quadratic function to validate the theoretical analysis in Section 3.1. Then we conduct more experiments on a popular machine learning problem called Ridge Logistic Regression, and compare accelerated regularized sub-sampled Newton with other classical algorithms.

### 5.1 Experiments on the Ridge Regression

The ridge regression is defined as follows

$$F(x) = \|Ax - b\|^2 + \lambda \|x\|^2, \quad (15)$$

where  $\lambda$  is a regularizer controls the condition number of the Hessian. In our experiment In our experiments, we choose dataset 'gisette' just as depicted in Table 1. And we set the regularizer  $\lambda = 1$ .




Figure 1: Experiment on the least square regression with  $|\mathcal{S}|$  being different sample size

Table 1: Datasets summary (sparsity =  $\frac{\text{\#Non-Zero Entries}}{n \times d}$ )

Dataset	$n$	$d$	sparsity	source
gisette	5,000	6,000	dense	libsvm dataset
sido0	12,678	4,932	dense	[9]
svhn	19,082	3,072	dense	libsvm dataset
rcv1	20,242	47,236	0.16%	libsvm dataset
real-sim	72,309	20,958	0.24%	libsvm dataset
avazu	2,085,163	999,975	0.0015%	libsvm dataset

In experiments, we set the sample size  $|\mathcal{S}|$  to be  $1\%n$ ,  $5\%n$  and  $10\%n$ . The regularizer  $\alpha$  of Algorithm 1 is properly chosen according to  $|\mathcal{S}|$ . ARSSN and RSSN share the same  $|\mathcal{S}|$ . And the acceleration parameter  $\theta$  of is fixed and appropriately selected. We report the experiments result in Figure 1.

From Figure 1, we can see that ARSSN and RSSN have significant difference in convergence rate and ARSSN is much faster. This validates the analysis in Section 3.1. Besides, we can also observe that ARSSN runs faster as sample size  $|\mathcal{S}|$  increases. When  $|\mathcal{S}| = 10\%n$ , ARSSN takes only about 3000 iterations to achieve an  $10^{-14}$  error while it needs about 6000 iterations to achieve the same precision when  $|\mathcal{S}| = 5\%n$ .

## 5.2 Experiments on the Ridge Logistic Regression

We conduct experiments on the Ridge Logistic Regression problem whose objective is

$$F(x) = \frac{1}{n} \sum_{i=1}^n \log[1 + \exp(-b_i \langle a_i, x \rangle)] + \frac{\lambda}{2} \|x\|^2, \quad (16)$$

where  $a_i \in \mathbb{R}^d$  is the  $i$ -th input vector, and  $b_i \in \{-1, 1\}$  is the corresponding label.

We conduct our experiments on six datasets: ‘gisette’, ‘protein’, ‘svhn’, ‘rcv1’, ‘sido0’, and ‘real-sim’. The first three datasets are dense and the last three ones are sparse. We give the detailed description of the datasets in Table 1. Notice that the size and dimension of dataset are close to each other, so the sketch Newton method [19, 25] can not be used. In our experiments, we try different settings of the regularizer  $\lambda$ , as  $1/n$ ,  $10^{-1}/n$ , and  $10^{-2}/n$  to represent different levels of regularization.

We compare Algorithm 1 (ARSSN) with RSSN (Algorithm 3), AGD ([17]) and SVRG ([11]) which are classical and popular optimization methods in machine learning.

In our experiments, the sample size  $|\mathcal{S}|$  and regularizer  $\alpha$  of ARSSN are chosen according to Theorem 5 while those of RSSN are chosen by Theorem 9. For a fixed  $|\mathcal{S}|$ , a proper  $\alpha$  can be found after several tries.




Figure 2: Experiment on ‘gisette’

In our experiment, the sub-sampled Hessian  $H^{(t)}$  constructed in Algorithm 1 can be written as

$$H^{(t)} = \tilde{A}^T \tilde{A} + (\alpha + \lambda)I,$$

where  $\tilde{A} \in \mathbb{R}^{\ell \times d}$ , where  $\ell < n$ . We can resort to Woodbury’s identity to compute the inverse of  $H^{(t)}$ . If  $\tilde{A}$  is sparse, we use conjugate gradient (Algorithm 4 in Appendix) to obtain an approximation of  $[H^{(t)}]^{-1} \nabla F(x^{(t)})$  which exploits the sparsity of  $\tilde{A}$ . In our experiments on sparse datasets, we set  $tol = 0.1 \|\nabla F(x^{(t)})\|$  for conjugate gradient (Algorithm 4).

For the acceleration parameter  $\theta$ , it is hard to get the best value for ARSSN just like AGD. However, our theoretical analysis implies that for large sample size  $|\mathcal{S}|$ , a small  $\theta$  should be chosen. In our experiments, we set  $\theta^{(t)} = \frac{t}{t+16}$  in Algorithm 1 for the dense datasets and  $\theta^{(t)} = \frac{t}{t+30}$  for the sparse datasets. We set  $x^{(0)} = 0$  for all the datasets and all the algorithms.

We report our result in Figure 2, 3, 4, 5, 6 and 7. We can see that ARSSN converges much faster than RSSN when these two algorithms have the same sample size. This shows Nesterov’s acceleration technique can promote the performance of regularized sub-sampled Newton effectively. We can also observe that ARSSN outperforms AGD significantly even when the sample size  $\mathcal{S}$  is  $1\%n$  or even less. This validates the fact that adding curvature information is an effective way to improve the ability of accelerated gradient descent.

Compared with SVRG, we can see that ARSSN also has better performance. Specifically, ARSSN performs much better than SVRG when  $\lambda$  is small like  $\lambda = 10^{-2}/n$ . When  $\lambda = 10^{-1}$ , ARSSN has comparable performance with SVRG on ‘sido0’, ‘rcv1’, ‘real-sim’, and ‘avazu’ while ARSSN performs much better on ‘gisette’ and ‘svhn’. This means that ARSSN is an efficient algorithm compared with classical stochastic first order method.

The experiments also reveal the fact that ARSSN has great advantages over other algorithms when the problem is ill-conditioned. On ‘gisette’, ‘sido0’, ‘svhn’, other algorithms have very poor performance on the case that  $\lambda = 10^{-2}/n$ . But ARSSN shows good convergence property.

### 5.3 Conclusion of Empirical Study

The above experiments show that Nesterov’s acceleration is an effective way to promote the convergence rate of approximate Newton methods. The experiments also show that adding some curvature information always help AGD to obtain a faster convergence rate.

## 6. Conclusion

In this paper, we have exploited Nesterov’s acceleration technique to promote the performance of second-order methods and propose Accelerated Regularized Sub-sample Newton. We have presented the theoretical analysis on the convergence properties of accelerated second-order methods, showing that accelerated approximate Newton has higher convergence rate, especially when the approximate Hessian is not a good approximation. Based on our theory, we have developed ARSSN. Our




Figure 3: Experiment on ‘sido0’




Figure 4: Experiment on ‘svhn’




Figure 5: Experiment on ‘rcv1’




Figure 6: Experiment on ‘real-sim’




Figure 7: Experiment on ‘avazu’

experiments have shown that our ARSSN performs much better than the conventional RSSN, which meets our theory well. ARSSN also has several advantages over other classical algorithms, demonstrating the efficiency of accelerated second-order methods.

## References

- [1] Zeyuan Allen-Zhu. Katyusha: The first direct acceleration of stochastic gradient methods. *arXiv preprint arXiv:1603.05953*, 2016.
- [2] Amir Beck and Marc Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM journal on imaging sciences*, 2(1):183–202, 2009.
- [3] Richard H Byrd, Gillian M Chin, Will Neveitt, and Jorge Nocedal. On the use of stochastic hessian information in optimization methods for machine learning. *SIAM Journal on Optimization*, 21(3):977–995, 2011.
- [4] Kenneth L Clarkson and David P Woodruff. Low rank approximation and regression in input sparsity time. In *Proceedings of the forty-fifth annual ACM symposium on Theory of computing*, pages 81–90. ACM, 2013.
- [5] Andrew Cotter, Ohad Shamir, Nati Srebro, and Karthik Sridharan. Better mini-batch algorithms via accelerated gradient methods. In *Advances in neural information processing systems*, pages 1647–1655, 2011.
- [6] Petros Drineas, Michael W Mahoney, and S Muthukrishnan. Sampling algorithms for l 2 regression and applications. In *Proceedings of the seventeenth annual ACM-SIAM symposium on Discrete algorithm*, pages 1127–1136. Society for Industrial and Applied Mathematics, 2006.
- [7] Murat A Erdogdu and Andrea Montanari. Convergence rates of sub-sampled newton methods. In *Advances in Neural Information Processing Systems*, pages 3034–3042, 2015.
- [8] Gene H Golub and Charles F Van Loan. *Matrix computations*, volume 3. JHU Press, 2012.
- [9] I. Guyon. Sido: A phamacology dataset. URL <http://www.causality.inf.ethz.ch/data/SIDO.html>.
- [10] N Halko, P G Martinsson, and J A Tropp. Finding Structure with Randomness : Probabilistic Algorithms for Matrix Decompositions. *SIAM Review*, 53(2):217–288, 2011.
- [11] Rie Johnson and Tong Zhang. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in Neural Information Processing Systems*, pages 315–323, 2013.

- [12] William B. Johnson and Joram Lindenstrauss. Extensions of Lipschitz mappings into a Hilbert space. *Contemporary mathematics*, 26(189-206), 1984.
- [13] Guanghui Lan and Yi Zhou. An optimal randomized incremental gradient method. *arXiv preprint arXiv:1507.02000*, 2015.
- [14] Mu Li, Tong Zhang, Yuqiang Chen, and Alexander J Smola. Efficient mini-batch training for stochastic optimization. In *Proceedings of the 20th ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 661–670. ACM, 2014.
- [15] Xiangrui Meng and Michael W Mahoney. Low-distortion subspace embeddings in input-sparsity time and applications to robust linear regression. In *Proceedings of the forty-fifth annual ACM symposium on Theory of computing*, pages 91–100. ACM, 2013.
- [16] Jelani Nelson and Huy L Nguyễn. Osnap: Faster numerical linear algebra algorithms via sparser subspace embeddings. In *Foundations of Computer Science (FOCS), 2013 IEEE 54th Annual Symposium on*, pages 117–126. IEEE, 2013.
- [17] Yurii Nesterov. A method of solving a convex programming problem with convergence rate  $o(1/k^2)$ . In *Soviet Mathematics Doklady*, volume 27, pages 372–376, 1983.
- [18] Yurii Nesterov et al. Gradient methods for minimizing composite objective function, 2007.
- [19] Mert Pilanci and Martin J. Wainwright. Newton sketch: A near linear-time optimization algorithm with linear-quadratic convergence. *SIAM Journal on Optimization*, 27(1):205–245, 2017.
- [20] Herbert Robbins and Sutton Monro. A stochastic approximation method. *The annals of mathematical statistics*, pages 400–407, 1951.
- [21] Farbod Roosta-Khorasani and Michael W Mahoney. Sub-sampled newton methods ii: Local convergence rates. *arXiv preprint arXiv:1601.04738*, 2016.
- [22] Nicolas L Roux, Mark Schmidt, and Francis R Bach. A stochastic gradient method with an exponential convergence rate for finite training sets. In *Advances in Neural Information Processing Systems*, pages 2663–2671, 2012.
- [23] Mark Schmidt, Nicolas Le Roux, and Francis Bach. Minimizing finite sums with the stochastic average gradient. *arXiv preprint arXiv:1309.2388*, 2013.
- [24] David P Woodruff. Sketching as a tool for numerical linear algebra. *Foundations and Trends® in Theoretical Computer Science*, 10(1–2):1–157, 2014.
- [25] Peng Xu, Jiyang Yang, Farbod Roosta-Khorasani, Christopher Ré, and Michael W Mahoney. Sub-sampled newton methods with non-uniform sampling. In *Advances in Neural Information Processing Systems*, pages 3000–3008, 2016.
- [26] Haishan Ye, Luo Luo, and Zhihua Zhang. Approximate newton methods and their local convergence. In *International Conference on Machine Learning*, pages 3931–3939, 2017.
- [27] Lijun Zhang, Mehrdad Mahdavi, and Rong Jin. Linear convergence with condition number independent access of full gradients. In *Advance in Neural Information Processing Systems 26 (NIPS)*, pages 980–988, 2013.

## Appendix A. Conjugate Gradient Descent

Conjugate Gradient is a classical method to solve the linear system of equation

$$Ax = b,$$

where  $A$  is a  $d \times d$  symmetric positive definite matrix. Algorithm 4 gives the detailed implementation to solve above linear system. Conjugate gradient has the following convergence properties [8]:

$$\|x_k - x_*\|_A \leq 2 \left( \frac{\sqrt{\kappa(A)} - 1}{\sqrt{\kappa(A)} + 1} \right)^k \|x_0 - x_*\|_A.$$

As we can see, the convergence behavior of conjugate gradient method depends on the condition of  $A$ . Hence, it suffers from poor convergence rate When  $A$  is ill-conditioned. Precondition method is an important way to improve the convergence properties. In the following Lemma, we give the sufficient condition of a preconditioner.

**Lemma 8** *If  $A$  and  $B$  are  $d \times d$  symmetric positive matrices, and  $(1 - \varepsilon)B \preceq A \preceq (1 + \varepsilon)B$ , where  $0 < \varepsilon_0 < 1$ , then for the optimization problem  $\min_x \|Ax - b\|$ , we have*

$$\|x_1 - x_*\|_A \leq \varepsilon \|x_*\|_A,$$

where  $x_* = A^{-1}b$  and  $x_1 = B^{-1}b$ . Besides, if we set  $r_k = Ax_k - b$ ,  $x_{kr} = B^{-1}r_k$  and  $x_{k+1} = x_k + x_{kr}$ , then  $\|x_{k+1} - x_*\|_A \leq \varepsilon_0 \|x_k - x_*\|_A$ .

**Proof** Because  $A \preceq (1 + \varepsilon)B$ , we have

$$\begin{aligned} \lambda_{\max}(B^{-1}A) &= \lambda_{\max}(B^{-1/2}AB^{-1/2}) \\ &= \max_{x \neq 0} \frac{x^T B^{-1/2}AB^{-1/2}x}{x^T x} \\ &= \max_{y \neq 0} \frac{y^T Ay}{y^T By} \\ &\leq 1 + \varepsilon, \end{aligned}$$

where the last equality is by setting  $y = B^{-1/2}x$ . Similarly, we have  $\lambda_{\min}(B^{-1}A) \geq 1 - \varepsilon$ .

Since  $B^{-1}A$  and  $A^{1/2}B^{-1}A^{1/2}$  are similar, the eigenvalues of  $A^{1/2}B^{-1}A^{1/2}$  are all between  $1 - \varepsilon$  and  $1 + \varepsilon$ . Therefore, we have

$$\|A^{1/2}B^{-1}A^{1/2} - I\| \leq \varepsilon,$$

where  $I$  is the  $d \times d$  identity matrix.

Hence, we have

$$\begin{aligned} \|B^{-1}b - x_*\|_A &= \|A^{1/2}(B^{-1}Ax_* - x_*)\| \\ &= \|A^{1/2}B^{-1}A^{1/2}(A^{1/2}x_*) - A^{1/2}x_*\| \\ &\leq \|A^{1/2}B^{-1}A^{1/2} - I\| \cdot \|A^{1/2}x_*\| \\ &= \varepsilon \|x_*\|_A. \end{aligned}$$

For the convergence property, we have

$$\begin{aligned} \|x_{k+1} - x_*\|_A &= \|B^{-1}(Ax_k - b) + x_k - A^{-1}b\|_A \\ &= \|B^{-1}(Ax_k - b) - A^{-1}(Ax_k - b)\|_A \\ &\leq \varepsilon_0 \|x_k - A^{-1}b\|_A \\ &= \varepsilon \|x_k - x_*\|_A. \end{aligned}$$

---

**Algorithm 3** Regularized Sub-sample Newton (RSSN).

---

- 1: **Input:**  $x^{(0)}$ ,  $0 < \delta < 1$ , regularizer parameter  $\alpha$ , sample size  $|\mathcal{S}|$  ;
  - 2: **for**  $t = 0, 1, \dots$  until termination **do**
  - 3:   Select a sample set  $\mathcal{S}$ , of size  $|\mathcal{S}|$  and  $H^{(t)} = \frac{1}{|\mathcal{S}|} \sum_{j \in \mathcal{S}} \nabla^2 f_j(x^{(t)}) + \alpha I$ ;
  - 4:   Update  $x^{(t+1)} = x^{(t)} - [H^{(t)}]^{-1} \nabla F(x^{(t)})$ ;
  - 5: **end for**
- 

The inequality is due to the property  $\|B^{-1}b - x_*\|_A \leq \|x_*\|_A$  and  $b$  replaced with  $Ax_k - b$  and  $x_*$  replaced with  $A^{-1}(Ax_k - b)$ .  $\blacksquare$

Let  $A$  and  $b$  be replaced by  $B^{-1/2}AB^{-1/2}$  and  $B^{-1/2}b$  respectively. And with some transformation, the iterations described in Lemma 8 can be transformed into preconditioned conjugate gradient method described in Algorithm 5. From Lemma 8, we can see that preconditioned conjugate gradient converges linearly with a constant rate using a good preconditioner.

## Appendix B. Regularized Sub-sampled Newton

The regularized Sub-sampled Newton method is depicted in Algorithm 3, and we now give its local convergence properties in the following theorem [26].

**Theorem 9** *Let  $F(x)$  satisfy Assumption 1 and 2. Assume Eqns. (10) and (11) hold, and let  $0 < \delta < 1$ ,  $0 \leq \epsilon_1 < 1$  and  $0 < \alpha$  be given. Assume  $\beta$  is a constant such that  $0 < \beta < \alpha + \frac{\sigma}{2}$ , the subsampled size  $|\mathcal{S}|$  satisfies  $|\mathcal{S}| \geq \frac{16K^2 \log(2d/\delta)}{\beta^2}$ , and  $H^{(t)}$  is constructed as in Algorithm 3. Define*

$$\epsilon_0 = \max \left( \frac{\beta - \alpha}{\sigma + \alpha - \beta}, \frac{\alpha + \beta}{\sigma + \alpha + \beta} \right),$$

*which implies that  $0 < \epsilon_0 < 1$ . And we define  $\|x\|_{M^*} = \|[M^*]^{-\frac{1}{2}}x\|$ . Then Algorithm 3 has the following convergence properties:*

1. *There exists a sufficient small value  $\gamma$ ,  $0 < \nu(t) < 1$ , and  $0 < \eta(t) < 1$  such that when  $\|x^{(t)} - x^*\| \leq \gamma$ , each iteration satisfies*

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \left( \epsilon_0 + \frac{2\eta(t)}{1 - \epsilon_0} \right) \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}.$$

*Besides,  $\nu(t)$  and  $\eta(t)$  will go to 0 as  $x^{(t)}$  goes to  $x^*$ .*

2. *If  $\nabla^2 F(x^{(t)})$  is also Lipschitz continuous with parameter  $\hat{L}$  and  $x^{(t)}$  satisfies*

$$\|x^{(t)} - x^*\| \leq \frac{\mu}{\hat{L}\kappa} \nu(t),$$

*where  $0 < \nu(t) < 1$ , then it holds that*

$$\|\nabla F(x^{(t+1)})\|_{M^*} \leq \epsilon_0 \frac{1 + \nu(t)}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*} + \frac{2}{(1 - \epsilon_0)^2} \frac{\hat{L}\kappa}{\mu\sqrt{\mu}} \frac{(1 + \nu(t))^2}{1 - \nu(t)} \|\nabla F(x^{(t)})\|_{M^*}^2.$$

## Appendix C. Proof of Lemma 3

**Proof** By Taylor's theorem, we have

$$\begin{aligned}
& \nabla F(x^{(t+1)}) \\
&= \nabla F(y^{(t+1)}) + \nabla^2 F(y^{(t+1)})(-p^{(t+1)}) + o(p^{(t+1)}) \\
&= \nabla F(y^{(t+1)}) - \nabla^2 F(y^{(t+1)})[H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) + o(p^{(t+1)}) \\
&= \nabla F(y^{(t+1)}) - \nabla^2 F(x^*)[H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) - (\nabla^2 F(x^*) - \nabla^2 F(y^{(t+1)}))[H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) + o(p^{(t+1)}) \\
&= [\nabla^2 F(x^*)]^{\frac{1}{2}} \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(y^{(t+1)}) \\
&\quad + (\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)) [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) + o(p^{(t+1)}).
\end{aligned}$$

For  $\nabla F(y^{(t+1)})$ , we have

$$\begin{aligned}
\nabla F(y^{(t+1)}) &= \nabla F(x^{(t)} + \theta s^{(t)}) \\
&= \nabla F(x^{(t)}) + \theta \nabla^2 F(x^{(t)})(s^{(t)}) + o(s^{(t)}) \\
&= \nabla F(x^{(t)}) + \theta (\nabla F(x^{(t)}) - \nabla F(x^{(t-1)})) + o(s^{(t)}) \\
&= (1 + \theta) \nabla F(x^{(t)}) - \theta \nabla F(x^{(t-1)}) + o(s^{(t)}).
\end{aligned}$$

Besides, we have

$$\begin{aligned}
o(s^{(t)}) &= o(x^{(t)} - x^{(t-1)}) = o(x^{(t)} - x^* - (x^{(t-1)} - x^*)) \\
&= o(\nabla F(x^{(t)})) + o(\nabla F(x^{(t-1)})),
\end{aligned}$$

where the last equality is because  $\nabla F(x)$  is  $L$ -Lipschitz continuous. Hence, it holds that

$$\nabla F(y^{(t+1)}) = (1 + \theta) \nabla F(x^{(t)}) - \theta \nabla F(x^{(t-1)}) + o(\nabla F(x^{(t)}) + \nabla F(x^{(t-1)})) \quad (17)$$

For  $(\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)) [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)})$ , we show that it is of order  $o(\nabla F(y^{(t+1)}))$  as follows. First, if  $\nabla^2 F(x)$  is not Lipschitz continuous, then there exists a  $\gamma$  such that when  $\|y - x^*\| \leq \gamma$ , it holds that

$$\|\nabla^2 F(y) - \nabla^2 F(x^*)\| = o(1).$$

Such  $\gamma$  exists because  $\nabla^2 F(x)$  is continuous near optimal point  $x^*$ . Hence,  $(\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)) [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)})$  is of order  $o(\nabla F(y^{(t+1)}))$  when  $y^{(t+1)}$  is sufficient close to  $x^*$ .

If  $\nabla^2 F(x)$  is  $\hat{L}$ -Lipschitz continuous and  $F(x)$  is  $\mu$ -strongly convex, then we have

$$\|\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)\| \leq \hat{L} \|y^{(t+1)} - x^*\| \leq \frac{\hat{L}}{\mu} \|\nabla F(y^{(t+1)})\|.$$

Then, it holds that

$$\|(\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)) [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)})\| = O(\|\nabla F(y^{(t+1)})\|^2).$$

Besides, because of  $p^{(t+1)} = [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)})$ , we have

$$o(p^{(t+1)}) = o(\nabla F(y^{(t+1)})).$$

Combining Eqn. (17), we have

$$(\nabla^2 F(y^{(t+1)}) - \nabla^2 F(x^*)) [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) = o(\nabla F(x^{(t)})) + o(\nabla F(x^{(t-1)}))$$

---

**Algorithm 4** Conjugate Gradient Descent Method.

---

```
1: Input:  $A, b, x_0$ , and  $tol$ ;  
2: Set  $r_0 = Ax_0 - b$ ,  $p_0 = -r_0$ ,  $k = 0$ ;  
3: while  $\|r_k\| > tol$  do  
4:   Calculate  $\alpha_k = \frac{r_k^T r_k}{p_k^T A p_k}$ ;  
5:   Calculate  $x_{k+1} = x_k + \alpha_k p_k$  and  $r_{k+1} = r_k + \alpha_k A p_k$ ;  
6:   Calculate  $\beta_{k+1} = \frac{r_{k+1}^T r_{k+1}}{r_k^T r_k}$  and  $p_{k+1} = -r_{k+1} + \beta_{k+1} p_k$ ;  
7:    $k = k + 1$ ;  
8: end while  
9: Output:  $x_k$ .
```

---

---

**Algorithm 5** Preconditioned Conjugate Gradient Descent Method.

---

```
1: Input:  $x_0$ , iteration number  $T$ , and preconditioner  $P$ ;  
2: Set  $r_0 = Ax_0 - b$ , solve  $Py_0 = r_0$ , set  $p_0 = -y_0$ ,  $k = 0$ ;  
3: while  $k < T$  do  
4:   Calculate  $\alpha_k = \frac{r_k^T r_k}{p_k^T A p_k}$ ;  
5:   Calculate  $x_{k+1} = x_k + \alpha_k p_k$  and  $r_{k+1} = r_k + \alpha_k A p_k$ ;  
6:   Solve  $Py_{k+1} = r_{k+1}$ ;  
7:   Calculate  $\beta_{k+1} = \frac{r_{k+1}^T y_{k+1}}{r_k^T y_k}$  and  $p_{k+1} = -y_{k+1} + \beta_{k+1} p_k$ ;  
8:    $k = k + 1$ ;  
9: end while  
10: Output:  $x_k$ .
```

---

and

$$o(p^{(t+1)}) = o(\nabla F(x^{(t)})) + o(\nabla F(x^{(t-1)})).$$

Hence, we have the following result

$$\begin{aligned} & [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t+1)}) \\ = & \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) \left( (1 + \theta) [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t)}) - \theta [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(x^{(t-1)}) \right) \\ & + o(\nabla F(x^{(t)})) + o(\nabla F(x^{(t-1)})). \end{aligned}$$

■

## Appendix D. Proof of Theorem 4

**Proof of Theorem 4** By Taylor's theorem and the property of quadratic function, we have

$$\begin{aligned} & \nabla F(x^{(t+1)}) \\ = & \nabla F(y^{(t+1)}) + \nabla^2 F(y^{(t+1)})(-p^{(t+1)}) \\ = & \nabla F(y^{(t+1)}) - \nabla^2 F(y^{(t+1)})[H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) \\ = & \nabla F(y^{(t+1)}) - \nabla^2 F(x^*)[H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) + \nabla^2 F(x^*) \left( [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) - p^{(t)} \right) \\ = & [\nabla^2 F(x^*)]^{\frac{1}{2}} \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} [H^{(t+1)}]^{-1} [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) [\nabla^2 F(x^*)]^{-\frac{1}{2}} \nabla F(y^{(t+1)}) \\ & + \nabla^2 F(x^*) \left( [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) - p^{(t)} \right). \end{aligned}$$

For  $\nabla F(y^{(t+1)})$ , we have

$$\begin{aligned}
\nabla F(y^{(t+1)}) &= \nabla F(x^{(t)} + \theta s^{(t)}) \\
&= \nabla F(x^{(t)}) + \theta \nabla^2 F(x^{(t)})(s^{(t)}) + o(s^{(t)}) \\
&= \nabla F(x^{(t)}) + \theta(\nabla F(x^{(t)}) - \nabla F(x^{(t-1)})) \\
&= (1 + \theta)\nabla F(x^{(t)}) - \theta\nabla F(x^{(t-1)}).
\end{aligned}$$

For notational convinience, we use  $M_*$  to denote  $[\nabla^2 F(x^*)]^{-\frac{1}{2}}$ , we have

$$\begin{aligned}
&\mathbb{E} \left( M_* \nabla F(x^{(t+1)}) \right) \\
&= \left( I - [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(t+1)}]^{-1} \right) [\nabla^2 F(x^*)]^{\frac{1}{2}} \right) \left( (1 + \theta) M_* \nabla F(x^{(t)}) - \theta M_* \nabla F(x^{(t-1)}) \right) \\
&\quad + [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) - p^{(t)} \right). \tag{18}
\end{aligned}$$

Because sketching matrices share the same distribution,  $\mathbb{E}([H^{(t+1)}]^{-1})$  is a constant matrix. we use the following notation for convenience

$$K = I - [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(t+1)}]^{-1} \right) [\nabla^2 F(x^*)]^{\frac{1}{2}}.$$

And  $K$  has the following spectral decomposition

$$K = U \Lambda U^T,$$

with  $\lambda_1 \geq \lambda_2 \dots \lambda_d$  We can reformulate the Eqn. (18) as

$$\begin{aligned}
\begin{bmatrix} \mathbb{E} (M_* \nabla F(x^{(t+1)})) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} &= \begin{bmatrix} (1 + \theta)K, & -\theta K \\ I, & 0 \end{bmatrix} \cdot \begin{bmatrix} M_* \nabla F(x^{(t)}) \\ M_* \nabla F(x^{(t-1)}) \end{bmatrix} \\
&\quad + [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(t+1)}]^{-1} \nabla F(y^{(t+1)}) - p^{(t)} \right).
\end{aligned}$$

Define the matrix

$$T = \begin{bmatrix} (1 + \theta)K, & -\theta K \\ I, & 0 \end{bmatrix}.$$

Thus, we obtain

$$\begin{aligned}
&\begin{bmatrix} \mathbb{E} (M_* \nabla F(x^{(t+1)})) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} \\
&= T^t \cdot \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} + \sum_{i=0}^t T^{t-i} [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \nabla F(y^{(i+1)}) - p^{(i)} \right).
\end{aligned}$$

Let  $\Pi$  be the  $2d \times 2d$  matrix with entries

$$\Pi_{i,j} = \begin{cases} 1 & i \text{ odd, } j = i \\ 1 & i \text{ even, } j = 2n + i \\ 0 & \text{otherwise} \end{cases}$$

Then, by conjugation, we have

$$\begin{aligned}
& \Pi \begin{bmatrix} U & 0 \\ 0 & U \end{bmatrix}^T \begin{bmatrix} (1+\theta)K & -\theta K \\ I & 0 \end{bmatrix} \begin{bmatrix} U & 0 \\ 0 & U \end{bmatrix} \Pi^T \\
&= \Pi \begin{bmatrix} (1+\theta)\Lambda & -\theta\Lambda \\ I & 0 \end{bmatrix} \Pi^T \\
&= \begin{bmatrix} T_1 & 0 & \cdots & 0 \\ 0 & T_2 & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & 0 & \cdots & T_d \end{bmatrix}
\end{aligned}$$

with

$$T_i = \begin{bmatrix} (1+\theta)\lambda_i & -\theta\lambda_i \\ 1 & 0 \end{bmatrix}.$$

That is  $T$  is similar to the block diagonal matrix with  $2 \times 2$  diagonal blocks  $T_i$ . And the eigenvalues of  $T_i$  are the roots of

$$a^2 - a(1+\theta)\lambda_i + \theta\lambda_i = 0.$$

Because of

$$(1-\pi)\nabla F(x^*) \preceq \mathbb{E}[H^{-1}] \preceq \nabla F(x^*),$$

we have

$$0 \leq \lambda_i \leq \pi.$$

Setting  $\theta = \frac{1-\sqrt{1-\pi}}{1+\sqrt{1-\pi}} - \epsilon_0$ , the larger eigenvalue of  $T_1$  is  $q = 1 - \sqrt{1-\pi} + O(\sqrt{\epsilon_0})$ . And Let  $S_1$  be the matrix consisting of eigenvectors of  $T_1$ . We denote

$$c_1 = \|S_1\| \cdot \|S_1^{-1}\|.$$

In fact,  $T_1$  dominates the convergence rate of  $T^t$ . Then we have

$$\|T^t\| \leq c_1 \cdot (1 - \sqrt{1-\pi})^t = c_1 q^t.$$

Thus, we have

$$\begin{aligned}
& \left\| \mathbb{E} \begin{bmatrix} M_* \nabla F(x^{(t+1)}) \\ M_* \nabla F(x^{(t)}) \end{bmatrix} \right\| \\
& \leq c_1 q^t \cdot \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\| + c_1 \sum_{i=0}^t q^{t-i} \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \nabla F(y^{(i+1)}) - p^{(i)} \right) \right\| \\
& = c_1 q^t \cdot A_0 + c_1 \sum_{i=0}^t q^{t-i} \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \nabla F(y^{(i+1)}) - p^{(i)} \right) \right\|,
\end{aligned}$$

where  $A_0 = \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\|.$

Due to Eqn. (3), (6) and using notation

$$\tilde{\epsilon}_1 = \frac{\epsilon_1}{12c_1\sqrt{\kappa}},$$

we also have

$$\begin{aligned}
& \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \nabla F(y^{(i+1)}) - p^{(i)} \right) \right\| \\
&= \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \left( \nabla F(y^{(i+1)}) - H^{(i+1)} p^{(i)} \right) \right) \right\| \\
&\leq \tilde{\epsilon}_1 \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \right) \right\| \cdot \left\| \mathbb{E} \left( \nabla F(y^{(i+1)}) \right) \right\| \\
&\leq \tilde{\epsilon}_1 \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \right) [\nabla^2 F(x^*)]^{\frac{1}{2}} M_\star \right\| \cdot \left\| M_\star^{-1} \mathbb{E} \left( M_\star \nabla F(y^{(i+1)}) \right) \right\| \\
&\leq \tilde{\epsilon}_1 (1 + \pi) \sqrt{\kappa} \left\| \mathbb{E} \left( M_\star \nabla F(y^{(i+1)}) \right) \right\| \\
&= \tilde{\epsilon}_1 (1 + \pi) \sqrt{\kappa} \left\| [1 + \theta, -\theta] \mathbb{E} \left[ \begin{array}{c} \nabla F(x^{(i)}) \\ \nabla F(x^{(i-1)}) \end{array} \right] \right\| \\
&\leq 6\tilde{\epsilon}_1 \sqrt{\kappa} \left\| \mathbb{E} \left[ \begin{array}{c} \nabla F(x^{(i)}) \\ \nabla F(x^{(i-1)}) \end{array} \right] \right\|
\end{aligned}$$

We use induction to prove our convergence rate, it is trivial for  $t = 1$  because

$$\begin{aligned}
\left\| \mathbb{E} \left[ \begin{array}{c} M_\star \nabla F(x^{(2)}) \\ M_\star \nabla F(x^{(1)}) \end{array} \right] \right\| &\leq c_1 q A_0 + \frac{\epsilon_1}{2c_1} A_0 \\
&= c_1 A_0 \left( q + \frac{\epsilon_1}{2c_1^2} \right) \\
&\leq c_1 A_0 (q + \epsilon_1) \\
&< 2c_1 p A_0.
\end{aligned}$$

The second inequality is because  $c_1$  is no smaller than 1. We assume that

$$\left\| \mathbb{E} \left[ \begin{array}{c} \nabla F(x^{(t+1)}) \\ \nabla F(x^{(t)}) \end{array} \right] \right\| \leq C p^t, \text{ with } p = q + \epsilon_1 \text{ and } C = 2c_1 A_0.$$

Then, we have

$$\begin{aligned}
& \left\| \mathbb{E} \left[ \begin{array}{c} M_\star \nabla F(x^{(t+1)}) \\ M_\star \nabla F(x^{(t)}) \end{array} \right] \right\| \\
&\leq c_1 q^t \cdot A_0 + c_1 \sum_{i=0}^t q^{t-i} \left\| [\nabla^2 F(x^*)]^{\frac{1}{2}} \mathbb{E} \left( [H^{(i+1)}]^{-1} \nabla F(y^{(i+1)}) - p^{(i)} \right) \right\| \\
&\leq c_1 q^t \cdot A_0 + 6\tilde{\epsilon}_1 c_1 \sqrt{\kappa} \sum_{i=1}^t q^{t-i} \left\| \mathbb{E} \left[ \begin{array}{c} \nabla F(x^{(i)}) \\ \nabla F(x^{(i-1)}) \end{array} \right] \right\| \\
&\leq c_1 q^t \cdot A_0 + 6C\tilde{\epsilon}_1 c_1 \sqrt{\kappa} \sum_{i=1}^t q^{t-i} p^{i-1} \\
&= c_1 q^t \cdot A_0 + 6C\tilde{\epsilon}_1 c_1 \sqrt{\kappa} \sum_{i=1}^t q^i p^{i-i-1} \\
&\leq c_1 q^t \cdot A_0 + 6C\tilde{\epsilon}_1 c_1 \sqrt{\kappa} \frac{p^t}{p - q} \\
&\leq \left( c_1 A_0 \left( \frac{q}{p} \right)^t + \frac{C}{2} \right) \cdot p^t \\
&\leq (c_1 A_0 + c_1 A_0) \cdot p^t = C p^t
\end{aligned}$$

Therefore, for all  $t = 1, \dots$ , we have

$$\left\| \mathbb{E} \begin{bmatrix} \nabla F(x^{(t+1)}) \\ \nabla F(x^{(t)}) \end{bmatrix} \right\| \leq Cp^t = 2c_1p^t \left\| \begin{bmatrix} M_* \nabla F(x^{(1)}) \\ M_* \nabla F(x^{(0)}) \end{bmatrix} \right\|.$$

■