

Approximation Algorithms for ℓ_0 -Low Rank Approximation

Karl Bringmann¹, Pavel Kolev^{*1}, and David P. Woodruff²

¹Max Planck Institute for Informatics, Saarland Informatics Campus, Saarbrücken,
Germany

{kbringma,pkolev}@mpi-inf.mpg.de

²Department of Computer Science, Carnegie Mellon University
dwoodruf@cs.cmu.edu

Abstract

We study the ℓ_0 -Low Rank Approximation Problem, where the goal is, given an $m \times n$ matrix A , to output a rank- k matrix A' for which $\|A' - A\|_0$ is minimized. Here, for a matrix B , $\|B\|_0$ denotes the number of its non-zero entries. This NP-hard variant of low rank approximation is natural for problems with no underlying metric, and its goal is to minimize the number of disagreeing data positions.

We provide approximation algorithms which significantly improve the running time and approximation factor of previous work. For $k > 1$, we show how to find, in $\text{poly}(mn)$ time for every k , a rank $O(k \log(n/k))$ matrix A' for which $\|A' - A\|_0 \leq O(k^2 \log(n/k)) \text{OPT}$. To the best of our knowledge, this is the first algorithm with provable guarantees for the ℓ_0 -Low Rank Approximation Problem for $k > 1$, even for bicriteria algorithms.

For the well-studied case when $k = 1$, we give a $(2+\epsilon)$ -approximation in *sublinear time*, which is impossible for other variants of low rank approximation such as for the Frobenius norm. We strengthen this for the well-studied case of binary matrices to obtain a $(1+O(\psi))$ -approximation in sublinear time, where $\psi = \text{OPT} / \|A\|_0$. For small ψ , our approximation factor is $1 + o(1)$.

^{*}This work has been funded by the Cluster of Excellence “Multimodal Computing and Interaction” within the Excellence Initiative of the German Federal Government.

Contents

1	Introduction	3
1.1	ℓ_0 -Low Rank Approximation	3
1.2	Our Results	4
2	Preliminaries	5
3	Algorithms for Reals ℓ_0-Rank-k	6
3.1	Structural Results	6
3.2	Basic Algorithm	8
3.3	Bicriteria Algorithm	9
4	Algorithm for Reals ℓ_0-Rank-1	12
4.1	Correctness	14
4.2	Implementing Step 2.2	16
4.3	Implementing Step 3	17
4.4	Analyzing the Runtime of Algorithm 2	19
5	Algorithms for Boolean ℓ_0-Rank-1	19
5.1	Preparations	20
5.2	Approximating α and β	21
5.3	The Algorithm	22
5.4	The Exact Algorithm	26
6	Sample Complexity Lower Bound for Boolean ℓ_0-Rank-1	27
6.1	Core Probabilistic Result	27
6.2	Hard Instance	29

1 Introduction

Low rank approximation of an $m \times n$ matrix A is an extremely well-studied problem, where the goal is to replace the matrix A with a rank- k matrix A' which well-approximates A , in the sense that $\|A - A'\|$ is small under some measure $\|\cdot\|$. Since any rank- k matrix A' can be written as $U \cdot V$, where U is $m \times k$ and V is $k \times n$, this allows for a significant parameter reduction. Namely, instead of storing A , which has mn entries, one can store U and V , which have only $(m + n)k$ entries in total. Moreover, when computing Ax , one can first compute Vx and then $U(Vx)$, which takes $(m + n)k$ instead of mn time. We refer the reader to several surveys [KV09, Mah11, Woo14] for references to the many results on low rank approximation.

We focus on approximation algorithms for the low-rank approximation problem, i.e. we seek to output a rank- k matrix A' for which $\|A - A'\| \leq \alpha \|A - A_k\|$, where $A_k = \operatorname{argmin}_{\operatorname{rank}(B)=k} \|A - B\|$ is the best rank- k approximation to A , and the approximation ratio α is as small as possible. One of the most widely studied error measures is the Frobenius norm $\|A\|_F = (\sum_{i=1}^m \sum_{j=1}^n A_{i,j}^2)^{1/2}$, for which the optimal rank- k approximation can be obtained via the singular value decomposition (SVD). Using randomization and approximation, one can compute an $\alpha = 1 + \epsilon$ -approximation, for any $\epsilon > 0$, in time much faster than the $\min(mn^2, mn^2)$ time required for computing the SVD, namely, in $O(\|A\|_0 + n \cdot \operatorname{poly}(k/\epsilon))$ time [CW13, MM13, NN13], where $\|A\|_0$ denotes the number of non-zero entries of A . For the Frobenius norm $\|A\|_0$ time is also a lower bound, as any algorithm that does not read nearly all entries of A might not read a very large entry, and therefore cannot achieve a relative error approximation.

The rank- k matrix A_k obtained by computing the SVD is also optimal with respect to any rotationally invariant norm, such as the operator and Schatten- p norms. Thus, such norms can also be solved exactly in polynomial time. Recently, however, there has been considerable interest [CW15, AGKM16, RSW16] in obtaining low rank approximations for NP-hard error measures such as the *entrywise* ℓ_p -norm $\|A\|_p = (\sum_{i,j} |A_{i,j}|^p)^{1/p}$, where $p \geq 1$ is a real number. Note that for $p < 1$, this is not a norm, though it is still a well-defined quantity. For $p = \infty$, this corresponds to the max-norm or Chebyshev norm. It is known that one can achieve a $\operatorname{poly}(k \log(mn))$ -approximation in $\operatorname{poly}(mn)$ time for the low-rank approximation problem with entrywise ℓ_p -norm for every $p \geq 1$ [SWZ16, CGK⁺17].

1.1 ℓ_0 -Low Rank Approximation

A natural variant of low rank approximation which the results above do not cover is that of *ℓ_0 -low rank approximation*, where the measure $\|A\|_0$ is the number of non-zero entries. In other words, we seek a rank- k matrix A' for which the number of entries (i, j) with $A'_{i,j} \neq A_{i,j}$ is as small as possible. Letting $\operatorname{OPT} = \min_{\operatorname{rank}(B)=k} \sum_{i,j} \delta(A_{i,j} \neq B_{i,j})$, where $\delta(A_{i,j} \neq A'_{i,j}) = 1$ if $A_{i,j} \neq A'_{i,j}$ and 0 otherwise, we would like to output a rank- k matrix A' for which there are at most $\alpha \operatorname{OPT}$ entries (i, j) with $A'_{i,j} \neq A_{i,j}$. Approximation algorithms for this problem are essential since solving the problem exactly is NP-hard [DAJ⁺15, GV15], even when $k = 1$ and A is a binary matrix.

The ℓ_0 -low rank approximation problem is quite natural for problems with no underlying metric, and its goal is to minimize the number of disagreeing data positions with a low rank matrix. Indeed, this error measure directly answers the following question: if we are allowed to ignore some data - outliers or anomalies - what is the best low-rank model we can get? One well-studied case is when A is binary, but A' and its factors U and V need not necessarily be binary. This is called unconstrained Binary Matrix Factorization in [JPHY14], which has applications to association rule

mining [KG03], biclustering structure identification [ZLD⁺10, ZLDZ07], pattern discovery for gene expression [SJY09], digits reconstruction [MGNR06], mining high-dimensional discrete-attribute data [KGR05, KGR06], market based clustering [Li05], and document clustering [ZLDZ07]. There is also a body of work on Boolean Matrix Factorization which restricts the factors to also be binary, which is referred to as constrained Binary Matrix Factorization in [JPHY14]. This is motivated in applications such as classifying text documents and there is a large body of work on this, see, e.g. [MV14, RPG15].

The ℓ_0 -low rank approximation problem coincides with a number of problems in different areas. It exactly coincides with the famous matrix rigidity problem over the reals, which asks for the minimal number OPT of entries of A that need to be changed in order to obtain a matrix of rank at most k . The matrix rigidity problem is well-studied in complexity theory [Gri76, Gri80, Val77] and parameterized complexity [FLM⁺17]. These works are not directly relevant here as they do not provide approximation algorithms. There are also other variants of ℓ_0 -low rank approximation, corresponding to cases such as when A is binary, $A' = UV$ is required to have binary factors U and V , and multiplication is either performed over a binary field [Yer11, GGYT12, DAJ⁺15, PRF15], or corresponds to an OR of ANDs. The latter is known as the Boolean model [BV10, DAJ⁺15, MMG⁺08, SBM03, SH06, VAG07]. These different notions of inner products lead to very different algorithms and results for the ℓ_0 -low rank approximation problem. However, all these models coincide in the special and important case in which A is binary and $k = 1$. This case was studied in [KG03, SJY09, JPHY14], as their algorithm for $k = 1$ forms the basis for their successful heuristic for general k , e.g. the PROXIMUS technique [KG03].

Another related problem is robust PCA [CLMW11a], in which there is an underlying matrix A that can be written as a low rank matrix L plus a sparse matrix S [CLMW11b]. Candès et al. [CLMW11b] argue that both components are of arbitrary magnitude, and we do not know the locations of the non-zeros in S nor how many there are. Moreover, grossly corrupted observations are common in image processing, web data analysis, and bioinformatics where some measurements are *arbitrarily* corrupted due to occlusions, malicious tampering, or sensor failures. Specific scenarios include video surveillance, face recognition, latent semantic indexing, and ranking of movies, books, etc. [CLMW11b]. These problems have the common theme of being an arbitrary magnitude sparse perturbation to a low rank matrix with no natural underlying metric, and so the ℓ_0 -distance measure (which is just the Hamming distance, or number of disagreements) is appropriate. In order to solve robust PCA in practice, Candès et al. [CLMW11b] relaxed the ℓ_0 -distance measure to the ℓ_1 -norm. Understanding theoretical guarantees for solving the original ℓ_0 -problem is of fundamental importance, and we study this problem in this paper.

Finally, interpreting 0^0 as 0, the ℓ_0 -low rank approximation problem coincides with the aforementioned notion of entrywise ℓ_p -approximation when $p = 0$. It is not hard to see that previous work [CGK⁺17] for general $p \geq 1$ fails to give any approximation factor for $p = 0$. Indeed, critical to their analysis is the scale-invariance property of a norm, which does not hold for $p = 0$ since ℓ_0 is not a norm.

1.2 Our Results

We provide approximation algorithms for the ℓ_0 -low rank approximation problem which significantly improve the running time or approximation factor of previous work. In some cases our algorithms even run in *sublinear time*, i.e., faster than reading all non-zero entries of the matrix. This is provably impossible for other measures such as the Frobenius norm and more generally, any ℓ_p -norm for $p > 0$. For $k > 1$, our approximation algorithms are, to the best of our knowledge, the

first with provable guarantees for this problem.

First, for $k = 1$, we significantly improve the polynomial running time of previous $(2 + \epsilon)$ -approximations for this problem. The best previous algorithm due to Jiang et al. [JPHY14] was based on the observation that there exists a column u of A spanning a 2-approximation. Therefore, solving the problem $\min_v \|A - uv\|_0$ for each column u of A yields a 2-approximation, where for a matrix B the measure $\|B\|_0$ counts the number of non-zero entries. The problem $\min_v \|A - uv\|_0$ decomposes into $\sum_i \min_{v_i} \|A_{:,i} - v_i u\|_0$, where $A_{:,i}$ is the i -th column of A , and v_i the i -th entry of vector v . The optimal v_i is the mode of the ratios $A_{i,j}/u_j$, where j ranges over indices in $\{1, 2, \dots, m\}$ with $u_j \neq 0$. As a result, one can find a rank-1 matrix uv^T providing a 2-approximation in $O(\|A\|_0 n)$ time, which was the best known running time. Somewhat surprisingly, we show that one can achieve *sublinear time* for solving this problem. Namely, we obtain a $(2 + \epsilon)$ -approximation in $(m + n) \text{poly}(\epsilon^{-1} \psi^{-1} \log(mn))$ time, for any $\epsilon > 0$, where $\psi = \text{OPT} / \|A\|_0$. This significantly improves upon the earlier $O(\|A\|_0 n)$ time for not too small ϵ and ψ . Our result should be contrasted to Frobenius norm low rank approximation, for which $\Omega(\|A\|_0)$ time is required even for $k = 1$, as otherwise one might miss a very large entry in A . Since ℓ_0 -low rank approximation is insensitive to the magnitude of entries of A , we bypass this general impossibility result.

Next, still considering the case of $k = 1$, we show that if the matrix A is binary, a well-studied case coinciding with the abovementioned $GF(2)$ and Boolean models, we obtain an approximation algorithm parameterized in terms of the ratio $\psi = \text{OPT} / \|A\|_0$, showing it is possible in time $(m + n) \psi^{-1} \text{poly}(\log(mn))$ to obtain a $(1 + O(\psi))$ -approximation. Note that our algorithm is again sublinear, unlike all algorithms in previous work. Moreover, when A is itself very well approximated by a low rank matrix, then ψ may actually be sub-constant, and we obtain a significantly better $(1 + o(1))$ -approximation than the previous best known 2-approximations. Thus, we simultaneously improve the running time and approximation factor. We also show that the running time of our algorithm is optimal up to $\text{poly}(\log(mn))$ factors by proving that any $(1 + O(\psi))$ -approximation succeeding with constant probability must read $\Omega((m + n) \psi^{-1})$ entries of A in the worst case.

Finally, for arbitrary $k > 1$, we first give an impractical algorithm, running in $n^{O(k)}$ time and achieving an $\alpha = \text{poly}(k)$ -approximation. To the best of our knowledge this is the first approximation algorithm for the ℓ_0 -low rank approximation problem with any non-trivial approximation factor. To make our algorithm practical, we reduce the running time to $\text{poly}(mn)$, with an exponent independent of k , if we allow for a bicriteria solution. In particular, we allow the algorithm to output a matrix A' of somewhat larger rank $O(k \log(n/k))$, for which $\|A - A'\|_0 \leq O(k^2 \log(n/k)) \cdot \min_{\text{rank}(B)=k} \|A - B\|_0$. Although we do not obtain rank exactly k , many of the motivations for finding a low rank approximation, such as reducing the number of parameters and fast matrix-vector product, still hold if the output rank is $O(k \log(n/k))$. We are not aware of any alternative algorithms which achieve $\text{poly}(mn)$ time and any provable approximation factor, even for bicriteria solutions.

2 Preliminaries

For an matrix $A \in \mathbb{A}^{m \times n}$ with entries $A_{i,j}$, we write $A_{i,:}$ for its i -th row and $A_{:,j}$ for its j -th column.

Input Formats We always assume that we have random access to the entries of the given matrix A , i.e. we can read any entry $A_{i,j}$ in constant time. For our sublinear time algorithms we need more efficient access to the matrix, specifically the following two variants:

- (1) We say that we are given A *with column adjacency arrays* if we are given arrays B_1, \dots, B_n

and lengths ℓ_1, \dots, ℓ_n such that for any $1 \leq k \leq \ell_j$ the pair $B_j[k] = (i, A_{i,j})$ stores the row i containing the k -th nonzero entry in column j as well as that entry $A_{i,j}$. This is a standard representation of matrices used in many applications. Note that given only these adjacency arrays B_1, \dots, B_n , in order to access any entry $A_{i,j}$ we can perform a binary search over B_j , and hence random access to any matrix entry is in time $O(\log n)$. Moreover, we assume to have random access to matrix entries in constant time, and note that this is optimistic by at most a factor $O(\log n)$.

(2) We say that we are given matrix A *with row and column sums* if we can access the numbers $\sum_j A_{i,j}$ for $i \in [m]$ and $\sum_i A_{i,j}$ for $j \in [n]$ in constant time (and, as always, access any entry $A_{i,j}$ in constant time). Notice that storing the row and column sums takes $O(m + n)$ space, and thus while this might not be standard information it is very cheap to store.

We show that the first access type even allows to sample from the set of nonzero entries uniformly in constant time.

Lemma 1. *Given a matrix $A \in \mathbb{R}^{m \times n}$ with column adjacency arrays, after $O(n)$ time preprocessing we can sample a uniformly random nonzero entry (i, j) from A in time $O(1)$.*

Proof. Note that we are in particular given the number of nonzero entries $\ell_j = \|A_{:,j}\|_0$ for each column. We want to first sample a column $X \in [n]$ such that $\Pr[X = j] = \ell_j / \sum_{k \in [n]} \ell_k$, then sample $Y \in [\ell_j]$ uniformly, read $B_X[Y] = (i, A_{i,X})$, and return $A_{i,X}$. Observe that this process indeed samples each nonzero entry of A with the same probability, since the probability of sampling a particular nonzero entry (i, j) is $(\ell_j / \sum_{k \in [n]} \ell_k) \cdot (1/\ell_j) = 1 / \sum_{k \in [n]} \ell_k$. Sampling $Y \in [\ell_j]$ uniformly can be done in constant time by assumption. For sampling X , we use the classic Alias Method by Walker [Wal74], which is given the probabilities $\Pr[X = 1], \dots, \Pr[X = n]$ as input and computes, in $O(n)$ time, a data structure that allows to sample from X in time $O(1)$. This finishes the construction. \square

3 Algorithms for Reals ℓ_0 -Rank- k

Given a matrix $A \in \mathbb{R}^{m \times n}$, the Reals ℓ_0 -Rank- k problem asks to find a matrix $A' \in \mathbb{R}^{m \times n}$ with rank k such that the difference between A and A' measured in ℓ_0 -distance is minimized. We denote the optimum value by

$$\text{OPT}^{(k)} \stackrel{\text{def}}{=} \min_{\text{rank}(A')=k} \|A - A'\|_0 = \min_{U \in \mathbb{R}^{m \times k}, V \in \mathbb{R}^{k \times n}} \|A - UV\|_0. \quad (1)$$

In this section, we establish several new results on the Reals ℓ_0 -Rank- k problem. In Subsection 3.1, we prove a structural lemma that shows the existence of k columns which provide a $(k + 1)$ -approximation to $\text{OPT}^{(k)}$, and we also give an $\Omega(k)$ -approximation lower bound for any algorithm that selects k columns from the input matrix A . In Subsection 3.2, we give an approximation algorithm that runs in $\text{poly}(n^k, m)$ time and achieves an $O(k^2)$ -approximation. To the best of our knowledge, this is the first algorithm with provable non-trivial approximation guarantees. In Subsection 3.3, we design a practical algorithm that runs in $\text{poly}(n, m)$ time with an exponent independent of k , if we allow for a bicriteria solution.

3.1 Structural Results

We give a new structural result for ℓ_0 -distance showing that any matrix A contains k columns which provide a $(k + 1)$ -approximation for the Reals ℓ_0 -Rank- k problem.

Lemma 2. Let $A \in \mathbb{R}^{m \times n}$ be a matrix and $k \in [n]$. There is a subset $J^{(k)} \subset [n]$ of size k and a matrix $Z \in \mathbb{R}^{k \times n}$ such that $\|A - A_{:,J^{(k)}}Z\|_0 \leq (k+1)\text{OPT}^{(k)}$.

Proof. Let $Q^{(0)}$ be the set of columns j with $UV_{:,j} = 0$, and let $R^{(0)} \stackrel{\text{def}}{=} [n] \setminus Q^{(0)}$. Let $S^{(0)} \stackrel{\text{def}}{=} [n]$, $T^{(0)} \stackrel{\text{def}}{=} \emptyset$. We split the value $\text{OPT}^{(k)}$ into $\text{OPT}(S^{(0)}, R^{(0)}) \stackrel{\text{def}}{=} \|A_{S^{(0)}, R^{(0)}} - UV_{S^{(0)}, R^{(0)}}\|_0$ and

$$\text{OPT}(S^{(0)}, Q^{(0)}) \stackrel{\text{def}}{=} \|A_{S^{(0)}, Q^{(0)}} - UV_{S^{(0)}, Q^{(0)}}\|_0 = \|A_{S^{(0)}, Q^{(0)}}\|_0.$$

Suppose $\text{OPT}(S^{(0)}, R^{(0)}) \geq |S^{(0)}||R^{(0)}|/(k+1)$. Then, for any subset $J^{(k)}$ it follows that

$$\min_Z \|A - A_{S^{(0)}, J^{(k)}}Z\|_0 \leq |S^{(0)}||R^{(0)}| + \|A_{S^{(0)}, Q^{(0)}}\|_0 \leq (k+1)\text{OPT}^{(k)}.$$

Otherwise, there is a column $i^{(1)}$ such that

$$\|A_{S^{(0)}, i^{(1)}} - (UV)_{S^{(0)}, i^{(1)}}\|_0 \leq \text{OPT}(S^{(0)}, R^{(0)})/|R^{(0)}| \leq \text{OPT}^{(k)}/|R^{(0)}|.$$

Let $T^{(1)}$ be the set of indices on which $(UV)_{S^{(0)}, i^{(1)}}$ and $A_{S^{(0)}, i^{(1)}}$ disagree, and similarly $S^{(1)} \stackrel{\text{def}}{=} S^{(0)} \setminus T^{(1)}$ on which they agree. Then we have $|T^{(1)}| \leq \text{OPT}^{(k)}/|R^{(0)}|$. Hence, in the submatrix $T^{(1)} \times R^{(0)}$ the total error is at most $|T^{(1)}| \cdot |R^{(0)}| \leq \text{OPT}^{(k)}$. Let $R^{(1)}, D^{(1)}$ be a partitioning of $R^{(0)}$ such that $A_{S^{(1)}, j}$ is linearly dependent on $A_{S^{(1)}, i^{(1)}}$ iff $j \in D^{(1)}$. Then by selecting column $A_{:, i^{(1)}}$ the incurred cost on matrix $S^{(1)} \times D^{(1)}$ is zero. For the remaining submatrix $S^{(1)} \times R^{(1)}$, we perform a recursive call of the algorithm.

We make at most k recursive calls, on instances $S^{(\ell)} \times R^{(\ell)}$ for $\ell \in \{0, \dots, k-1\}$. In the ℓ^{th} iteration, either $\text{OPT}(S^{(\ell)}, R^{(\ell)}) \geq |S^{(\ell)}||R^{(\ell)}|/(k+1-\ell)$ and we are done, or there is a column $i^{(\ell+1)}$ which partitions $S^{(\ell)}$ into $S^{(\ell+1)}, T^{(\ell+1)}$ and $R^{(\ell)}$ into $R^{(\ell+1)}, D^{(\ell+1)}$ such that

$$|S^{(\ell+1)}| \geq m \cdot \prod_{i=0}^{\ell} \left(1 - \frac{1}{k+1-i}\right) = \frac{k-\ell}{k+1} \cdot m$$

and for every $j \in D^{(\ell)}$ the column $A_{S^{(\ell+1)}, j}$ belongs to the span of $\{A_{S^{(\ell+1)}, i^{(t)}}\}_{t=1}^{\ell+1}$.

Suppose we performed k recursive calls. We show now that the incurred cost in submatrix $S^{(k)} \times R^{(k)}$ is at most $\text{OPT}(S^{(k)}, R^{(k)}) \leq \text{OPT}^{(k)}$. By construction, $|S^{(k)}| \geq m/(k+1)$ and the subcolumns $\{A_{S^{(k)}, i}\}_{i \in I^{(k)}}$ are linearly independent, where $I^{(k)} = \{i^{(1)}, \dots, i^{(k)}\}$ is the set of the selected columns, and $A_{S^{(k)}, I^{(k)}} = (UV)_{S^{(k)}, I^{(k)}}$. Since $\text{rank}(A_{S^{(k)}, I^{(k)}}) = k$, it follows that $\text{rank}(U_{S^{(k)}, :}) = k$, $\text{rank}(V_{:, I^{(k)}}) = k$ and the matrix $V_{:, I^{(k)}} \in \mathbb{R}^{k \times k}$ is invertible. Hence, for matrix $Z = (V_{:, I^{(k)}})^{-1} V_{:, R^k}$ we have

$$\text{OPT}(S^{(k)}, R^{(k)}) = \|A_{S^k, R^k} - A_{S^k, I^k}Z\|_0.$$

The statement follows by noting that the recursive calls accumulate a total cost of at most $k \cdot \text{OPT}^{(k)}$ in the submatrices $T^{(\ell+1)} \times R^{(\ell)}$ for $\ell \in \{0, 1, \dots, k-1\}$, as well as cost at most $\text{OPT}^{(k)}$ in submatrix $S^{(k)} \times R^{(k)}$. \square

We also show that any algorithm that selects k columns of a matrix A incurs at least an $\Omega(k)$ -approximation for the Reals ℓ_0 -Rank- k problem.

Lemma 3. Let $k \leq n/2$. Suppose $A = (G_{k \times n}; I_{n \times n}) \in \mathbb{R}^{(n+k) \times n}$ is a matrix composed of a Gaussian random matrix $G \in \mathbb{R}^{k \times n}$ with $G_{i,j} \sim N(0, 1)$ and identity matrix $I_{n \times n}$. Then for any subset $J^{(k)} \subset [n]$ of size k , we have $\min_{Z \in \mathbb{R}^{k \times n}} \|A - A_{:, J^{(k)}}Z\|_0 = \Omega(k) \cdot \text{OPT}^{(k)}$.

Proof. Notice that the optimum cost is at most n , achieved by selecting $U = (I_{k \times k}; 0_{n \times k})$ and $V = G_{k \times n}$. It is well known that Gaussian matrices are invertible with probability 1, see e.g. [SST06, Thm 3.3]. Hence, $G_{:,J^{(k)}}$ is a nonsingular matrix for every subset $J^{(k)} \subset [n]$ of size k .

We will show next that for any subset $J^{(k)}$ of k columns the incurred cost is at least $(n-k)k \geq nk/2$. Without loss of generality, the chosen columns $J^{(k)} = [k]$ are the first k columns of A . Let $R = [2k]$ be the first $2k$ rows and $C = [n] \setminus J$ be the last $n-k$ columns. We bound

$$\begin{aligned} \min_Z \|A - A_{:, [k]} Z\|_0 &\geq \min_Z \|A_{R,C} - A_{R, [k]} Z\|_0 \\ &= \sum_{j \in C} \min_{z^{(j)}} \|A_{R,j} - A_{R, [k]} z^{(j)}\|_0, \end{aligned}$$

i.e. we ignore all rows and columns except R and C . Consider any column $j \in C$. Since $A_{R,j} = (G_{:,j}, 0_k)$ and $A_{R, [k]} = (G_{:, [k]}, I_{k \times k})$, for any vector $z \in \mathbb{R}^k$ we have

$$\begin{aligned} \|A_{R,j} - A_{R, J^{(k)}} z\|_0 &= \|G_{:,j} - G_{:, [k]} z\|_0 + \|I_{k \times k} z\|_0 \\ &= \|G_{:,j} - G_{:, [k]} z\|_0 + \|z\|_0. \end{aligned}$$

Let $\ell \stackrel{\text{def}}{=} \|z\|_0$. By symmetry, without loss of generality we can assume that the first ℓ entries of z are non-zero and the remaining entries are 0. Let $x \in \mathbb{R}^\ell$ be the vector containing the first ℓ entries of z . Then we have

$$\|A_{R,j} - A_{R, J^{(k)}} z\|_0 = \|G_{:,j} - G_{:, [\ell]} x\|_0 + \ell.$$

We consider w.l.o.g. the first k columns of A , and we construct the optimum matrix Z that minimizes $\|A_{:, 1:k} Z - A\|_0$. Observe that it is optimal to set the first k columns of Z to $I_{k \times k}$, and since $A_{2k+1:n, 1:k} = 0$ we can focus only on the submatrix $A_{1:2k, k+1:n} = (G_{1:k, k+1:n}; 0_{k \times n-k})$.

Consider a column $A_{1:2k, j}$ for $j \in [k+1, n]$. Our goal is to find a vector $v \in \mathbb{R}^k$ minimizing the objective function $\Psi = \min_v \{\|v\|_0 + \|G^{(k)} v - g\|_0\}$, where $G^{(k)} \stackrel{\text{def}}{=} \{G_{1:k, 1:k}\}$ and $g \stackrel{\text{def}}{=} G_{1:k, j}$. It holds with probability 1 that $G^{(k)}$ and g do not have an entry equal to zero. Moreover, since $G^{(k)}$ is invertible every row in $[G^{(k)}]^{-1}$ is non-zero, and thus with probability 1 a vector $v = [G^{(k)}]^{-1} g$ has entry equal to zero.

Let $v = (x; 0)$ be an arbitrary vector with $\|x\|_1 = \ell$. Let $G^{(\ell)}$ be a submatrix of $G^{(k)}$ induced by the first ℓ columns. For every subset $S \subset [m]$ of ℓ rows the corresponding submatrix $G_{S, :}^{(\ell)}$ has a full rank. Suppose there is a subset S such that for $G_{S, :}^{(\ell)}$ and g_S there is a vector $x \in \mathbb{R}^k$ satisfying $G_{S, :}^{(\ell)} x = g_S$. Since $G_{S, :}^{(\ell)}$ is invertible, the existence of x implies its uniqueness. On the other hand, for any row $i \in [m] \setminus S$ the probability of the event $G_{i, :}^{(\ell)} x = g_i$ is equals to 0. Since $G^{(k)} v = G^{(\ell)} x$ and there are finitely many possible subsets S as above, i.e. $\binom{m}{\ell} \leq m^\ell$, by union bound it follows that $\|G^{(k)} v - g\|_0 \geq k - \ell$. Therefore, it holds that $\phi \geq k$.

The statement follows by noting that the total cost incurred by $A_{:, 1:k}$ and any Z is lower bounded by $(n-k)k + (n-k) = (1-k/n)(k+1)n$. \square

3.2 Basic Algorithm

We give an impractical algorithm that runs in $\text{poly}(n^k, m)$ time and achieves an $O(k^2)$ -approximation. To the best of our knowledge this is the first approximation algorithm for the Reals ℓ_0 -Rank- k problem with non-trivial approximation guarantees.

Theorem 4. *Given $A \in \mathbb{R}^{m \times n}$ and $k \in [n]$ we can compute in $O(n^{k+1}m^2k^{\omega+1})$ time a set of k indices $J^{(k)} \subset [n]$ and a matrix $Z \in \mathbb{R}^{k \times n}$ such that $\|A - A_{:,J^{(k)}}Z\|_0 \leq O(k^2) \cdot \text{OPT}^{(k)}$.*

We use as a subroutine the algorithm of Berman and Karpinski [BK02] (attributed also to Kannan in that paper) which given a matrix U and a vector b approximates $\min_x \|Ux - b\|_0$ in polynomial time. Specifically, we invoke in our algorithm the following variant of this result established by Alon, Panigrahy, and Yekhanin [APY09].

Theorem 5. [APY09] *There is an algorithm that given a matrix $A \in \mathbb{R}^{m \times k}$ and a vector $b \in \mathbb{R}^m$, outputs in time $O(m^2k^{\omega+1})$ a vector $z \in \mathbb{R}^k$ such that w.h.p. $\|Az - b\|_0 \leq k \cdot \min_x \|Ax - b\|_0$.*

Proof of Theorem 4. The existence of a subset J^* of k columns of A and matrix $Z^* \in \mathbb{R}^{k \times n}$ with $\|A - A_{:,J^*}Z^*\|_0 \leq (k+1)\text{OPT}^{(k)}$ follows by Lemma 2. We enumerate all $\binom{n}{k}$ subsets $J^{(k)}$ of k columns. For each $J^{(k)}$, we split $\min_Z \|A_{:,J^{(k)}}Z - A\|_0 = \sum_{i=1}^n \min_{z^{(i)}} \|A_{:,J^{(k)}}z^{(i)} - A_{:,i}\|_0$, and we run the algorithm from Theorem 5 for each column $A_{:,i}$, obtaining approximate solutions $\tilde{z}^{(1)}, \dots, \tilde{z}^{(n)}$ that form a matrix \tilde{Z} . Then, we return the best solution $(A_{:,J^{(k)}}, \tilde{Z})$. To verify that this yields a $k(k+1)$ -approximation, note that for $J^{(k)} = J^*$ we have

$$\begin{aligned} \|A_{:,J^*}\tilde{Z} - A\|_0 &= \sum_{i=1}^n \|A_{:,J^*}\tilde{z}^{(i)} - A_{:,i}\|_0 \leq k \sum_{i=1}^n \min_{z^{(i)}} \|A_{:,J^*}z^{(i)} - A_{:,i}\|_0 \\ &= k \cdot \min_Z \|A_{:,J^*}Z - A\|_0 \leq k(k+1) \cdot \text{OPT}^{(k)}. \end{aligned}$$

The time bound $O(n^{k+1}m^2k^{\omega+1})$ is immediate from Theorem 5. This proves the statement. \square

3.3 Bicriteria Algorithm

Our main contribution in this section is to design a practical algorithm that runs in $\text{poly}(n, m)$ time with an exponent independent of k , if we allow for a bicriteria solution.

Theorem 6. *Given $A \in \mathbb{R}^{m \times n}$ with $m \geq n$ and $k \in \{1, \dots, n\}$, there is an algorithm that in expected $\text{poly}(m, n)$ time outputs a subset of indices $J \subset [n]$ with $|J| = O(k \log(n/k))$ and a matrix $Z \in \mathbb{R}^{|J| \times n}$ such that $\|A - A_{:,J}Z\|_0 \leq O(k^2 \log(n/k)) \cdot \text{OPT}^{(k)}$.*

The structure of the proof follows a recent approximation algorithm [CGK⁺17, Algorithm 3] for the ℓ_p -low rank approximation problem, for any $p \geq 1$. We note that the analysis of [CGK⁺17, Theorem 7] is missing an $O(\log^{1/p} n)$ approximation factor, and naïvely provides an $O(k \log^{1/p} n)$ -approximation rather than the stated $O(k)$ -approximation. Further, it might be possible to obtain an efficient algorithm yielding an $O(k^2 \log k)$ -approximation for Theorem 6 using unpublished techniques in [SWZ18]; we leave the study of obtaining the optimal approximation factor to future work.

There are two critical differences with the proof of [CGK⁺17, Theorem 7]. We cannot use the earlier [CGK⁺17, Theorem 3] which shows that any matrix A contains k columns which provide an $O(k)$ -approximation for the ℓ_p -low rank approximation problem, since that proof requires $p \geq 1$ and critically uses scale-invariance, which does not hold for $p = 0$. Our combinatorial argument in Lemma 2 seems fundamentally different than the maximum volume submatrix argument in [CGK⁺17] for $p \geq 1$.

Second, unlike for ℓ_p -regression for $p \geq 1$, the ℓ_0 -regression problem $\min_x \|Ux - b\|_0$ given a matrix U and vector b is not efficiently solvable since it corresponds to a nearest codeword problem,

Algorithm 1 Bicriteria Algorithm: Selecting $O(k \log(n/k))$ Columns

ApproximatelySelectColumns(A, k)

ensure A has at least $k \log(n/k)$ columns

1. **If** the number of columns of matrix A is less than or equal to $2k$
 2. **Return** all the columns of A
 3. **Else**
 4. **Repeat**
 5. Let R be a set of $2k$ uniformly random columns of A
 6. **Until** at least $1/10$ fraction of columns of A are nearly approximately covered
 7. Let $A_{\overline{R}}$ be the columns of A not nearly approximately covered by R
 8. **Return** $R \cup \text{ApproximatelySelectColumns}(A_{\overline{R}}, k)$
-

which is NP-hard [Ale11]. Thus, we resort to an approximation algorithm for ℓ_0 -regression, based on ideas for solving the nearest codeword problem in [APY09, BK02].

Note that $\text{OPT}^{(k)} \leq \|A\|_0$. Since there are only $mn + 1$ possibilities of $\text{OPT}^{(k)}$, we can assume we know $\text{OPT}^{(k)}$ and we can run the Algorithm 1 below for each such possibility, obtaining a rank- $O(k \log n)$ solution, and then outputting the solution found with the smallest cost.

This can be further optimized by forming instead $O(\log(mn))$ guesses of $\text{OPT}^{(k)}$. One of these guesses is within a factor of 2 from the true value of $\text{OPT}^{(k)}$, and we note that the following argument only needs to know $\text{OPT}^{(k)}$ up to a factor of 2.

We start by defining the notion of approximate coverage, which is different than the corresponding notion in [CGK⁺17] for $p \geq 1$, due to the fact that ℓ_0 -regression cannot be efficiently solved. Consequently, approximate coverage for $p = 0$ cannot be efficiently tested. Let $Q \subseteq [n]$ and $M = A_{:,Q}$ be an $m \times |Q|$ submatrix of A . We say that a column $M_{:,i}$ is (S, Q) -approximately covered by a submatrix $M_{:,S}$ of M , if $|S| = 2k$ and

$$\min_x \|M_{:,S} \cdot x - M_{:,i}\|_0 \leq \frac{100(k+1)\text{OPT}^{(k)}}{|Q|}. \quad (2)$$

Lemma 7. (Similar to [CGK⁺17, Lemma 6], but using Lemma 2) Let $Q \subseteq [n]$ and $M = A_{:,Q}$ be a submatrix of A . Suppose we select a subset R of $2k$ uniformly random columns of M . Then with probability at least $1/3$, at least a $1/10$ fraction of the columns of M are (R, Q) -approximately covered.

Proof. To show this, as in [CGK⁺17], consider a uniformly random column index i not in the set R . Let $T \stackrel{\text{def}}{=} R \cup \{i\}$, $\eta \stackrel{\text{def}}{=} \min_{\text{rank}(B)=k} \|M_{:,T} - B\|_0$, and $B^* \stackrel{\text{def}}{=} \arg \min_{\text{rank}(B)=k} \|M - B\|_0$. Since T is a uniformly random subset of $2k + 1$ columns of M , we have

$$\begin{aligned} \mathbb{E}_T \eta &\leq \mathbb{E}_T \|M_{:,T} - B^*_{:,T}\|_0 = \sum_{T \in \binom{[Q]}{2k+1}} \sum_{i \in T} \|M_{:,i} - B^*_{:,i}\|_0 \Pr[T] \\ &= \sum_{i \in Q} \frac{\binom{|Q|-1}{2k}}{\binom{|Q|}{2k+1}} \|M_{:,i} - B^*_{:,i}\|_0 = \frac{(2k+1)\text{OPT}_M^{(k)}}{|Q|} \leq \frac{(2k+1)\text{OPT}^{(k)}}{|Q|}. \end{aligned}$$

Then, by a Markov bound, we have $\Pr[\eta \leq \frac{10(2k+1)\text{OPT}^{(k)}}{|Q|}] \geq 9/10$. Let \mathcal{E}_1 denotes this event.

Fix a configuration $T = R \cup \{i\}$ and let $L(T) \subset T$ be the subset guaranteed by Lemma 2 such that $|L(T)| = k$ and

$$\min_X \|M_{:,L(T)}X - M_{:,T}\|_0 \leq (k+1) \min_{\text{rank}(B)=k} \|M_{:,T} - B\|_0.$$

Notice that

$$\mathbb{E}_i \left[\min_x \|M_{:,L(T)}x - M_{:,i}\|_0 \mid T \right] = \frac{1}{2k+1} \min_X \|M_{:,L(T)}X - M_{:,T}\|_0,$$

and thus by the law of total probability we have

$$\mathbb{E}_T \left[\min_x \|M_{:,L(T)}x - M_{:,i}\|_0 \right] \leq \frac{(k+1)\eta}{2k+1}.$$

Let \mathcal{E}_2 denote the event that $\min_x \|M_{:,L}x - M_{:,i}\|_0 \leq \frac{10(k+1)\eta}{2k+1}$. By a Markov bound, $\Pr[\mathcal{E}_2] \geq 9/10$.

Further, as in [CGK⁺17], let \mathcal{E}_3 be the event that $i \notin L$. Observe that there are $\binom{k+1}{k}$ ways to choose a subset $R' \subset T$ such that $|R'| = 2k$ and $L \subset R'$. Since there are $\binom{2k+1}{2k}$ ways to choose R' , it follows that $\Pr[L \subset R \mid T] = \frac{k+1}{2k+1} > 1/2$. Hence, by the law of total probability, we have $\Pr[\mathcal{E}_3] > 1/2$.

As in [CGK⁺17], $\Pr[\mathcal{E}_1 \wedge \mathcal{E}_2 \wedge \mathcal{E}_3] > 2/5$, and conditioned on $\mathcal{E}_1 \wedge \mathcal{E}_2 \wedge \mathcal{E}_3$,

$$\min_x \|M_{:,R}x - M_{:,i}\|_0 \leq \min_x \|M_{:,L}x - M_{:,i}\|_0 \leq \frac{10(k+1)\eta}{2k+1} \leq \frac{100(k+1)\text{OPT}^{(k)}}{|Q|}, \quad (3)$$

where the first inequality uses that L is a subset of R given \mathcal{E}_3 , and so the regression cost cannot decrease, while the second inequality uses the occurrence of \mathcal{E}_2 and the final inequality uses the occurrence of \mathcal{E}_1 .

As in [CGK⁺17], if Z_i is an indicator random variable indicating whether i is approximately covered by R , and $\tilde{Z} = \sum_{i \in Q} Z_i$, then $\mathbb{E}_R[\tilde{Z}] \geq \frac{2|Q|}{5}$ and $\mathbb{E}_R[|Q| - \tilde{Z}] \leq \frac{3|Q|}{5}$. By a Markov bound, it follows that $\Pr[|Q| - \tilde{Z} \geq \frac{9|Q|}{10}] \leq \frac{2}{3}$. Thus, probability at least $1/3$, at least a $1/10$ fraction of the columns of M are (R, Q) -approximately covered. \square

Given Lemma 7, we are ready to prove Theorem 6. As noted above, a key difference with the corresponding [CGK⁺17, Algorithm 3] for ℓ_p and $p \geq 1$, is that we cannot efficiently test if the i -th column is approximately covered by the set R . We will instead again make use of Theorem 5.

Proof of Theorem 6. The computation of matrix Z force us to relax the notion of (R, Q) -approximately covered to the notion of (R, Q) -nearly-approximately covered as follows: we say that a column $M_{:,i}$ is (R, Q) -nearly-approximately covered if, the algorithm in Theorem 5 returns a vector z such that

$$\|M_{:,R}z - M_{:,i}\|_0 \leq \frac{100(k+1)^2\text{OPT}^{(k)}}{|Q|}. \quad (4)$$

By the guarantee of Theorem 5, if $M_{:,i}$ is (R, Q) -approximately covered then it is also with probability at least $1 - 1/\text{poly}(mn)$ (R, Q) -nearly-approximately covered.

Suppose Algorithm 1 makes t iterations and let $A_{:, \cup_{i=1}^t R_i}$ and Z be the resulting solution. We bound now its cost. Let $B_0 = [n]$, and consider the i -th iteration of Algorithm 1. We denote

by R_i a set of $2k$ uniformly random columns of B_{i-1} , by G_i a set of columns that is (R_i, B_{i-1}) -nearly-approximately covered, and by $B_i = B_{i-1} \setminus \{G_i \cup R_i\}$ a set of the remaining columns. By construction, $|G_i| \geq |B_{i-1}|/10$ and

$$|B_i| \leq \frac{9}{10}|B_{i-1}| - 2k < \frac{9}{10}|B_{i-1}|.$$

Since Algorithm 1 terminates when $B_{t+1} \leq 2k$, we have

$$2k < |B_t| < \left(1 - \frac{1}{10}\right)^t |B_0| = \left(1 - \frac{1}{10}\right)^t n,$$

and thus the number of iterations $t \leq 10 \log(n/2k)$. By construction, $|G_i| = (1 - \alpha_i)|B_{i-1}|$ for some $\alpha_i \leq 9/10$, and hence

$$\sum_{i=1}^t \frac{|G_i|}{|B_{i-1}|} \leq t \leq 10 \log \frac{n}{2k}. \quad (5)$$

Therefore, the solution cost is bounded by

$$\begin{aligned} \|A_{:, \cup_{i=1}^t R_i} Z - A\|_0 &= \sum_{i=1}^t \sum_{j \in G_i} \|A_{:, R_i} z^{(j)} - A_{:, j}\|_0 \\ &\stackrel{\text{Lem. 7}}{\leq} \sum_{i=1}^t \sum_{j \in G_i} k \cdot \min_{x^{(j)}} \|A_{:, R_i} x^{(j)} - A_{:, j}\|_0 \stackrel{(4)}{\leq} \sum_{i=1}^t \sum_{j \in G_i} \frac{100(k+1)^2 \text{OPT}^{(k)}}{|B_{i-1}|} \\ &= 100(k+1)^2 \text{OPT}^{(k)} \cdot \sum_{i=1}^t \frac{|G_i|}{|B_{i-1}|} \stackrel{(5)}{\leq} O\left(k^2 \cdot \log \frac{n}{2k}\right) \cdot \text{OPT}^{(k)}. \end{aligned}$$

By Lemma 7, the expected number of iterations of selecting a set R_i such that $|G_i| \geq 1/10|B_{i-1}|$ is $O(1)$. Since the number of recursive calls t is bounded by $O(\log(n/k))$, it follows by a Markov bound that Algorithm 1 chooses $O(k \log(n/k))$ columns in total. Since the approximation algorithm of Theorem 5 runs in polynomial time, our entire algorithm has expected polynomial time. \square

4 Algorithm for Reals ℓ_0 -Rank-1

Given a matrix $A \in \mathbb{R}^{m \times n}$, the Reals ℓ_0 -Rank-1 problem asks to find a matrix $A' \in \mathbb{R}^{m \times n}$ with rank 1, i.e. $A' = uv^T$ for some vectors $u \in \mathbb{R}^m$ and $v \in \mathbb{R}^n$, such that the difference between A and A' measured in ℓ_0 -distance is minimized. We denote the optimum value by

$$\text{OPT}^{(1)} \stackrel{\text{def}}{=} \min_{\text{rank}(A')=1} \|A - A'\|_0 = \min_{u \in \mathbb{R}^m, v \in \mathbb{R}^n} \|A - uv^T\|_0. \quad (6)$$

In the trivial case when $\text{OPT}^{(1)} = 0$, there is an optimal algorithm that runs in time $O(\|A\|_0)$ and finds the exact rank-1 decomposition uv^T of a matrix A . In this work, we focus on the case when $\text{OPT}^{(1)} \geq 1$. We show that Algorithm 2 yields a $(2 + \epsilon)$ -approximation factor and runs in nearly linear time in $\|A\|_0$, for any constant $\epsilon > 0$. Furthermore, a variant of our algorithm even runs in sublinear time, if $\|A\|_0$ is large and

$$\psi \stackrel{\text{def}}{=} \text{OPT}^{(1)} / \|A\|_0 \quad (7)$$

is not too small. In particular, we obtain time $o(\|A\|_0)$ when $\text{OPT}^{(1)} \geq (\epsilon^{-1} \log(mn))^4$ and $\|A\|_0 \geq n(\epsilon^{-1} \log(mn))^4$.

Theorem 8. *There is an algorithm that, given $A \in \mathbb{R}^{m \times n}$ with column adjacency arrays and $\text{OPT}^{(1)} \geq 1$, and given $\epsilon \in (0, 0.1]$, runs w.h.p. in time*

$$O\left(\left(\frac{n \log m}{\epsilon^2} + \min\left\{\|A\|_0, n + \psi^{-1} \frac{\log n}{\epsilon^2}\right\}\right) \frac{\log^2 n}{\epsilon^2}\right)$$

and outputs a column $A_{:,j}$ and a vector $z \in \mathbb{R}^n$ such that w.h.p. $\|A - A_{:,j} \cdot z^T\|_0 \leq (2 + \epsilon)\text{OPT}^{(1)}$. The algorithm also computes an estimate Y satisfying w.h.p. $(1 - \epsilon)\text{OPT}^{(1)} \leq Y \leq (2 + 2\epsilon)\text{OPT}^{(1)}$.

In fact, our analysis of Theorem 8 directly applies to the Boolean ℓ_0 -Rank-1 problem, for the definition see Section 5, and yields as a special case the following result (which we use to prove Theorem 18 in Section 5).

Theorem 9. *Let $\text{OPT} = \min_{u \in \{0,1\}^m, v \in \{0,1\}^n} \|A - u \cdot v^T\|_0$. Given a binary matrix $A \in \{0,1\}^{m \times n}$ with column adjacency arrays and $\text{OPT} \geq 1$, and given $\epsilon \in (0, 0.1]$, we can compute w.h.p. in time*

$$O\left(\left(\frac{n \log m}{\epsilon^2} + \min\left\{\|A\|_0, n + \psi^{-1} \frac{\log n}{\epsilon^2}\right\}\right) \frac{\log^2 n}{\epsilon^2}\right)$$

a column $A_{:,j}$ and a binary vector $z \in \{0,1\}^n$ such that w.h.p. $\|A - A_{:,j} \cdot z^T\|_0 \leq (2 + \epsilon)\text{OPT}$. Further, we can compute an estimate Y such that w.h.p. $(1 - \epsilon)\text{OPT} \leq Y \leq (2 + 2\epsilon)\text{OPT}$.

The rest of this section is devoted to proving Theorem 8. We present the pseudocode of this result in Algorithm 2.

Algorithm 2 Reals ℓ_0 -Rank-1: Approximation Scheme

Input: $A \in \mathbb{R}^{m \times n}$ and $\epsilon \in (0, 0.1)$.

1. Partition the columns of A into weight-classes $\mathcal{S} = \{S^{(0)}, \dots, S^{(1+\log n)}\}$ such that
 - i) $S^{(0)}$ contains all columns j with $\|A_{:,j}\|_0 = 0$, and
 - ii) $S^{(i)}$ contains all columns j with $2^{i-1} \leq \|A_{:,j}\|_0 < 2^i$.
2. **For each** weight-class $S^{(i)}$ **do**:
 - 2.1 Sample a set $C^{(i)}$ of $\Theta(\epsilon^{-2} \log n)$ elements uniformly at random from $S^{(i)}$.
 - 2.2 Find a $(1 + \frac{\epsilon}{15})$ -approximate solution $z^{(j)} \in \mathbb{R}^n$ for each column $A_{:,j} \in C^{(i)}$, i.e.

$$\|A - A_{:,j} \cdot [z^{(j)}]^T\|_0 \leq \left(1 + \frac{\epsilon}{15}\right) \min_v \|A - A_{:,j} \cdot v^T\|_0. \quad (8)$$

3. Compute a $(1 + \frac{\epsilon}{15})$ -approximation Y_j of $\|A - A_{:,j} \cdot [z^{(j)}]^T\|_0$ for every $j \in \bigcup_{i \in [|\mathcal{S}|]} C^{(i)}$.
 4. **Return** the pair $(A_{:,j}, z^{(j)})$ corresponding to the minimal value Y_j .
-

The only steps for which the implementation details are not immediate are Steps 2.2 and 3. We will discuss them in Sections 4.2 and 4.3, respectively.

Note that the algorithm from Theorem 8 selects a column $A_{:,j}$ and then finds a good vector z such that the product $A_{:,j} z^T$ approximates A . We show that the approximation guarantee $2 + \epsilon$ is essentially tight for algorithms following this pattern.

Lemma 10. *There exist a matrix $A \in \mathbb{R}^{n \times n}$ such that $\min_{z \in \mathbb{R}^n} \|A - A_{:,j} \cdot z^T\|_0 \geq 2(1 - 1/n)\text{OPT}^{(1)}$, for every column $A_{:,j}$.*

Proof of Lemma 10. Let $A = I + J \in \mathbb{R}^{n \times n}$, where I is an identity matrix and $J = \mathbf{1}\mathbf{1}^T$ is an all-ones matrix. Note that $\text{OPT}^{(1)} \leq n$ is achieved by approximating A with the rank-1 matrix J . On the other hand, when we choose $u = A_{:,i}$ for any $i \in [n]$, the incurred cost on any column $A_{:,j}$, $j \neq i$, is $\min_x \|A_{:,j} - xA_{:,i}\|_0 = 2$, since there are two entries where $A_{:,i}$ and $A_{:,j}$ disagree. Hence, the total cost is at least $2n - 2 \geq (2 - 2/n)\text{OPT}^{(1)}$. \square

4.1 Correctness

We first prove the following structural result, capturing Steps 1-2.2 of Algorithm 2.

Lemma 11. *Let $C^{(0)}, \dots, C^{(\log n+1)}$ be the sets constructed in Step 2.1 of Algorithm 2, and let $C = C^{(0)} \cup \dots \cup C^{(\log n+1)}$. Then, w.h.p. C contains an index $j \in [n]$ such that*

$$\min_{z \in \mathbb{R}^n} \|A - A_{:,j} \cdot z^T\|_0 \leq (2 + \epsilon/2) \cdot \text{OPT}^{(1)}.$$

Proof. Let u, v be an optimum solution of (6). For the weight class $S^{(0)}$ containing all columns without nonzero entries, setting $z_c = 0$ for any $c \in S^{(0)}$ gives zero cost on these columns, no matter what column $A_{:,j}$ we picked. Hence, without loss of generality in the following we assume that $S^{(0)} = \emptyset$.

For any $i \geq 1$, we partition the weight class $S^{(i)}$ into $N^{(i)}, Z^{(i)}$ such that $v_i = 0$ for every $i \in Z^{(i)}$ and $v_i \neq 0$ for $i \in N^{(i)}$. We denote by \mathcal{S}^+ the set of weight-classes $S^{(i)}$ with $|N^{(i)}| \geq \frac{1}{3}|S^{(i)}|$. Let $\mathcal{R} = \bigcup_{i \in \mathcal{S}^+} S^{(i)}$ and $\mathcal{W} = [n] \setminus \mathcal{R}$. We partition $\mathcal{R} = \mathcal{N} \cup \mathcal{Z}$ such that $v_i = 0$ for every $i \in \mathcal{Z}$ and $v_i \neq 0$ for $i \in \mathcal{N}$. Further, using the three sets \mathcal{N}, \mathcal{Z} and \mathcal{W} we decompose $\text{OPT}^{(1)}$ into

$$\begin{aligned} \text{OPT}^{(1)} &= \text{OPT}_{\mathcal{N}} + \text{OPT}_{\mathcal{Z}} + \text{OPT}_{\mathcal{W}} \\ &= \|A_{:, \mathcal{N}} - u \cdot v_{\mathcal{N}}^T\|_0 + \|A_{:, \mathcal{Z}}\|_0 + \|A_{:, \mathcal{W}} - u \cdot v_{\mathcal{W}}^T\|_0. \end{aligned}$$

The proof proceeds by case distinction:

The set \mathcal{Z} : For any column $A_{:,j}$ of A , we have

$$\min_{z_{\mathcal{Z}}} \|A_{:, \mathcal{Z}} - A_{:,j} \cdot z_{\mathcal{Z}}^T\|_0 \leq \|A_{:, \mathcal{Z}} - A_{:,j} \cdot 0\|_0 = \|A_{:, \mathcal{Z}}\|_0 = \text{OPT}_{\mathcal{Z}}. \quad (9)$$

The set \mathcal{W} : Note that \mathcal{W} consists of all weight classes $S^{(i)}$ with $|Z^{(i)}| > \frac{2}{3}|S^{(i)}|$. For any such weight class $S^{(i)}$, the optimum cost satisfies

$$\|A_{:, S^{(i)}} - uv_{S^{(i)}}^T\|_0 \geq \|A_{:, Z^{(i)}}\|_0 \geq \frac{2}{3}|S^{(i)}|2^{i-1} = \frac{1}{3}|S^{(i)}|2^i.$$

Further, for any column $A_{:,j}$ of A , we have

$$\begin{aligned} \min_z \|A_{:, S^{(i)}} - A_{:,j} \cdot z^T\|_0 &\leq \|A_{:, S^{(i)}}\|_0 \leq \|A_{:, Z^{(i)}}\|_0 + \frac{1}{3}|S^{(i)}|2^i \\ &\leq 2\|A_{:, Z^{(i)}}\|_0 \leq 2\|A_{:, S^{(i)}} - uv_{S^{(i)}}^T\|_0, \end{aligned}$$

and thus the total cost in \mathcal{W} is bounded by

$$\min_{z_{\mathcal{W}}} \|A_{:, \mathcal{W}} - A_{:,j} \cdot z_{\mathcal{W}}^T\|_0 = \sum_{i \in \mathcal{W}} \min_{z_{S^{(i)}}} \|A_{:, S^{(i)}} - A_{:,j} \cdot z_{S^{(i)}}^T\|_0 \leq 2\|A_{:, \mathcal{W}} - uv_{\mathcal{W}}^T\|_0 = 2\text{OPT}_{\mathcal{W}}. \quad (10)$$

The set \mathcal{N} : By an averaging argument there is a subset $G \subseteq \mathcal{N}$ of size $|G| \geq \frac{\epsilon}{3}|\mathcal{N}|$ such that for every $j \in G$ we have

$$\|A_{:,j} - v_j \cdot u\|_0 \leq \frac{1}{1 - \epsilon/3} \cdot \frac{\text{OPT}_{\mathcal{N}}}{|\mathcal{N}|} \leq \left(1 + \frac{\epsilon}{2}\right) \cdot \frac{\text{OPT}_{\mathcal{N}}}{|\mathcal{N}|}.$$

Let $j \in G$ be arbitrary. Furthermore, let $P^{(j)}$ be the set of all rows i with $A_{i,j} = v_j \cdot u_i$, and let $Q^{(j)} = [m] \setminus P^{(j)}$. By construction, we have $|Q^{(j)}| \leq (1 + \frac{\epsilon}{2})\text{OPT}_{\mathcal{N}}/|\mathcal{N}|$. Moreover, since $j \in \mathcal{N}$ we have $v_j \neq 0$, and thus we may choose $z_{\mathcal{N}} = \frac{1}{v_j}v_{\mathcal{N}}$. This yields

$$\begin{aligned} \min_{z_{\mathcal{N}}} \|A_{:, \mathcal{N}} - A_{:,j} \cdot z_{\mathcal{N}}^T\|_0 &\leq \|A_{:, \mathcal{N}} - A_{:,j} \cdot \frac{1}{v_j}v_{\mathcal{N}}^T\|_0 \\ &= \|A_{P^{(j)}, \mathcal{N}} - u_{P^{(j)}} \cdot v_{\mathcal{N}}^T\|_0 + \|A_{Q^{(j)}, \mathcal{N}} - A_{:,j} \cdot \frac{1}{v_j}v_{\mathcal{N}}^T\|_0 \\ &\leq \text{OPT}_{\mathcal{N}} + \left(1 + \frac{\epsilon}{3}\right) \text{OPT}_{\mathcal{N}} = \left(2 + \frac{\epsilon}{3}\right) \text{OPT}_{\mathcal{N}}. \end{aligned} \quad (11)$$

Hence, by combining (9), (10) and (11), it follows for any index $j \in G$ that

$$\begin{aligned} &\min_z \|A - A_{:,j} \cdot z^T\|_0 \\ &= \min_{z_{\mathcal{W}}} \|A_{:, \mathcal{W}} - A_{:,j} \cdot z_{\mathcal{W}}^T\|_0 + \min_{z_{\mathcal{Z}}} \|A_{:, \mathcal{Z}} - A_{:,j} \cdot z_{\mathcal{Z}}^T\|_0 + \min_{z_{\mathcal{N}}} \|A_{:, \mathcal{N}} - A_{:,j} \cdot z_{\mathcal{N}}^T\|_0 \\ &\leq 2\text{OPT}_{\mathcal{W}} + \text{OPT}_{\mathcal{Z}} + \left(2 + \frac{\epsilon}{2}\right) \text{OPT}_{\mathcal{N}} \leq \left(2 + \frac{\epsilon}{2}\right) \text{OPT}^{(1)}. \end{aligned}$$

This yields the desired approximation guarantee, provided that we sampled a column $A_{:,j}$ from G . We show next that whenever $\mathcal{N} \neq \emptyset$, our algorithm samples with high probability at least one column from G .

Before that let us consider the case when $\mathcal{N} = \emptyset$. Then, since the bounds (9) and (10) hold for any column $A_{:,j}$ of A , the set C contains only good columns. Thus, we may assume that $\mathcal{N} \neq \emptyset$.

We now analyze the probability of sampling a column $A_{:,j}$ from G . By construction, the set \mathcal{N} is the union of all $N^{(i)}$ such that $|N^{(i)}| \geq \frac{1}{3}|S^{(i)}|$. As shown above, we have $|G| \geq \frac{\epsilon}{3}|\mathcal{N}|$, and thus there is an index i satisfying $|G \cap S^{(i)}| \geq \frac{\epsilon}{3}|N^{(i)}| \geq \frac{\epsilon}{9}|S^{(i)}|$. Hence, when sampling a uniformly random element from $S^{(i)}$ we hit G with probability at least $\frac{\epsilon}{9}$. Since we sample $\Theta(\epsilon^{-2} \log n)$ elements from $S^{(i)}$, we hit G with high probability. This finishes the proof. \square

Correctness Proof of Algorithm 2: It remains to show that the pair $(A_{:,j}, z^j)$ with minimum estimate Y_j yields a $(2 + \epsilon)$ -approximation to $\text{OPT}^{(1)}$. By Step 3, for every column j we have

$$\left(1 + \frac{\epsilon}{15}\right)^{-1} \cdot \|A - A_{:,j}[z^{(j)}]^T\|_0 \leq Y_j \leq \left(1 + \frac{\epsilon}{15}\right) \cdot \|A - A_{:,j}[z^{(j)}]^T\|_0. \quad (12)$$

Since $Y_j \leq Y_{j'}$ for any other column j' , (12) and the approximation guarantee of Steps 2.2 yield

$$\left(1 + \frac{\epsilon}{15}\right)^{-1} \|A - A_{:,j}[z^{(j)}]^T\|_0 \leq \left(1 + \frac{\epsilon}{15}\right) \|A - A_{:,j'}[z^{(j')}]^T\|_0 \leq \left(1 + \frac{\epsilon}{15}\right)^2 \min_z \|A - A_{:,j'}z^T\|_0.$$

By Lemma 11, w.h.p. there exists a column $j' \in C$ with $\min_z \|A - A_{:,j'}z^T\|_0 \leq (2 + \frac{\epsilon}{2})\text{OPT}^{(1)}$. We obtain a total approximation ratio of $(1 + \frac{\epsilon}{15})^3(2 + \frac{\epsilon}{2}) \leq 2 + \epsilon$ for any error $0 < \epsilon \leq 0.1$, i.e. we

have $\|A - A_{:,j}[z^{(j)}]^T\|_0 \leq (2 + \epsilon)\text{OPT}^{(1)}$. Therefore, it holds that

$$(1 - \epsilon)\text{OPT}^{(1)} \leq \left(1 + \frac{\epsilon}{15}\right)^{-1} \text{OPT}^{(1)} \leq Y_j \leq \left(1 + \frac{\epsilon}{15}\right) (2 + \epsilon)\text{OPT}^{(1)} \leq (2 + 2\epsilon)\text{OPT}^{(1)}.$$

This finishes the correctness proof.

4.2 Implementing Step 2.2

Step 2.2 of Algorithm 2 uses the following sublinear procedure.

Algorithm 3 Reals ℓ_0 -Rank-1: Objective Value Estimation

Input: $A \in \mathbb{R}^{m \times n}$, $u \in \mathbb{R}^m$ and $\epsilon \in (0, 1)$.

let $t \stackrel{\text{def}}{=} \Theta(\epsilon^{-2} \log m)$, $N \stackrel{\text{def}}{=} \text{supp}(u)$, and $p \stackrel{\text{def}}{=} t/|N|$.

1. Select each index $i \in N$ with probability p and let S be the resulting set.
 2. Compute a vector $z \in \mathbb{R}^n$ such that $z_j = \arg \min_{r \in \mathbb{R}} \|A_{S,j} - r \cdot u_S\|_0$ for all $j \in [n]$.
 3. **Return** the vector z .
-

We prove now the correctness of Algorithm 3 and we analyze its runtime.

Lemma 12. *Given $A \in \mathbb{R}^{m \times n}$ with $m \geq n$, $u \in \mathbb{R}^m$ and $\epsilon \in (0, 1)$ we can compute in time $O(\epsilon^{-2} n \log m)$ a vector $z \in \mathbb{R}^n$ such that w.h.p. for every index $i \in [n]$ it holds that*

$$\|A_{:,i} - z_i \cdot u\|_0 \leq (1 + \epsilon) \min_{v_i \in \mathbb{R}} \|A_{:,i} - v_i \cdot u\|_0.$$

Proof. Let N, Z be a partitioning of $[m]$ such that $u_i = 0$ for $i \in Z$ and $u_i \neq 0$ for $i \in N$. Since $\|A - u \cdot z^T\|_0 = \|A_{N,:} - u_N \cdot z^T\|_0 + \|A_{Z,:}\|_0$, it suffices to find a vector z such that for every $j \in [n]$ we have

$$\|A_{N,j} - z_j \cdot u_N\|_0 \leq (1 + \epsilon) \cdot \min_{v_j} \|A_{N,j} - v_j \cdot u_N\|_0. \quad (13)$$

Let $j \in [n]$ be arbitrary. For $r \in \mathbb{R}$ let $G(r) \stackrel{\text{def}}{=} \{i \in N : A_{i,j}/u_i = r\}$ be the set of entries of $A_{N,j}$ that we correctly recover by setting $z_j = r$. Note that $\|A_{N,j} - z_j \cdot u_N\|_0 = |N| - |G(z_j)|$ holds for any $z_j \in \mathbb{R}$. Hence, the optimal solution sets $z_j = r^* \stackrel{\text{def}}{=} \arg \max_{r \in \mathbb{R}} |G(r)|$.

Let $X_{G(r)}$ be a random variable indicating the number of elements selected from group $G(r)$ in Step 1 of Algorithm 3. Notice that $\mathbb{E}[X_{G(r)}] = t \cdot |G(r)|/|N|$, and by Chernoff bound w.h.p. we have

$$|X_{G(r)} - \mathbb{E}[X_{G(r)}]| \leq (\epsilon/8) \cdot t. \quad (14)$$

Let $S \subseteq N$ be the set of selected indices. Further, since $|S| = \sum_{\ell} X_{G(r)}$ and $\mathbb{E}[|S|] = t$, by Chernoff bound we have w.h.p. $|S| \leq (1 + \epsilon)t$. Observe that Step 2 of Algorithm 3 selects $z_j = \arg \max_{r \in \mathbb{R}} X_{G(r)}$, since $\|A_{S,j} - r \cdot u_S\|_0 = |S| - X_{G(r)}$. We now relate z_j to r^* . The proof proceeds by case distinction on $\delta^* \stackrel{\text{def}}{=} |G(r^*)|/|N|$.

Case 1: Suppose $\delta^* \leq \epsilon/4$. Then $\|A_{N,j} - r \cdot u_N\|_0 \geq (1 - \epsilon/4)|N|$ for every $r \in \mathbb{R}$, and thus no matter which z_j is selected we obtain a $(1 + \epsilon)$ -approximation, since

$$\|A_{N,j} - z_j \cdot u_N\|_0 \leq |N| \leq (1 + \epsilon) \min_r \|A_{N,j} - r \cdot u_N\|_0.$$

Case 2: Suppose $\delta^* \geq 1/2 + \epsilon$. Then, by (14) w.h.p. we have

$$X_{G(r^*)} \geq \mathbb{E}[X_{G(r^*)}] - (\epsilon/8)t = (\delta^* - \epsilon/8)t > (1 + \epsilon)t/2 \geq |S|/2,$$

and thus X_{r^*} is maximal among all X_r . Hence, we select the optimal $z_j = r^*$.

Case 3: Suppose $\epsilon/4 < \delta^* < 1/2 + \epsilon$. Let $z_j = r$ be the value chosen by Algorithm 3. By (14), the event of making a mistake, given by $X_{G(r)} \geq X_{G(r^*)}$, happens when

$$\mathbb{E}[X_{G(r)}] + (\epsilon/8)t \geq \mathbb{E}[X_{G(r^*)}] - (\epsilon/8)t. \quad (15)$$

Let $\delta \stackrel{\text{def}}{=} |G(r)|/|N|$ and note that (15) implies $\delta \geq \delta^* - \epsilon/4$. Hence, for the selected $r \neq r^*$ we have

$$\begin{aligned} \|A_{N,j} - r \cdot u_N\|_0 &= (1 - \delta)|N| \leq (1 - \delta^* + \epsilon/4)|N| \\ &\leq (1 + \epsilon)(1 - \delta^*)|N| = (1 + \epsilon)\|A_{N,j} - r^* \cdot u_N\|_0. \end{aligned}$$

Therefore, in each of the preceding three cases, we obtain w.h.p. a $(1 + \epsilon)$ -approximate solution. The statement follows by the union bound. \square

4.3 Implementing Step 3

In Step 3 of Algorithm 2, our goal is to compute a $(1 + \frac{\epsilon}{15})$ -approximation Y_j of $\|A - A_{:,j}[z^{(j)}]^T\|_0$, for every $j \in \bigcup_{i \in [S]} C^{(i)}$.

In this subsection, our main algorithmic result is the following.

Theorem 13. *There is an algorithm that, given $A \in \mathbb{R}^{m \times n}$ with column adjacency arrays and $\text{OPT}^{(1)} \geq 1$, and given $j \in [n]$, $v \in \mathbb{R}^m$ and $\epsilon \in (0, 1)$, outputs an estimator Y that satisfies w.h.p.*

$$(1 - \epsilon) \|A - A_{:,j} \cdot v^T\|_0 \leq Y \leq (1 + \epsilon) \|A - A_{:,j} \cdot v^T\|_0.$$

The algorithm runs w.h.p. in time $O(\min\{\|A\|_0, n + \epsilon^{-2}\psi^{-1} \log n\})$, where $\psi = \text{OPT}^{(1)} / \|A\|_0$.

We prove Theorem 13, by designing: i) an exact deterministic algorithm, see Lemma 14; and ii) a randomized approximation algorithm running in sublinear-time, see Lemma 16.

Lemma 14. *Suppose $A, B \in \mathbb{R}^{m \times n}$ are represented by column adjacency arrays. Then, we can compute in $O(\|A\|_0 + n)$ time the measure $\|A - B\|_0$.*

Proof. We partition the entries of A into five sets:

$$\begin{aligned} T_1 &= \{(i, j) : A_{ij} = 0 \text{ and } B_{ij} \neq 0\}, & T_4 &= \{(i, j) : 0 \neq A_{ij} = B_{ij} \neq 0\}, \\ T_2 &= \{(i, j) : A_{ij} \neq 0 \text{ and } B_{ij} = 0\}, & T_5 &= \{(i, j) : A_{ij} = B_{ij} = 0\}, \\ T_3 &= \{(i, j) : 0 \neq A_{ij} \neq B_{ij} \neq 0\}. \end{aligned}$$

Observe that $\|A - B\|_0 = |T_1| + |T_2| + |T_3|$ and $\|B\|_0 = |T_1| + |T_3| + |T_4|$. Since $\|A - B\|_0 = \|B\|_0 + |T_2| - |T_4|$, it suffices to compute the numbers $\|B\|_0$, $|T_2|$ and $|T_4|$. We compute $|T_2|$ and $|T_4|$ in $O(\|A\|_0)$ time, by enumerating all non-zero entries of A and performing $O(1)$ checks for each. For $\|B\|_0$, we sum the column lengths of B in $O(n)$ time. \square

For our second, sampling-based implementation of Step 3, we make use of an algorithm by Dagum et al. [DKLR00] for estimating the expected value of a random variable. We note that the runtime of their algorithm is a random variable, the magnitude of which is bounded w.h.p. within a certain range.

Theorem 15. [DKLR00] *Let X be a random variable taking values in $[0, 1]$ with $\mu \stackrel{\text{def}}{=} \mathbb{E}[X] > 0$. Let $0 < \epsilon, \delta < 1$ and $\rho_X = \max\{\text{Var}[X], \epsilon\mu\}$. There is an algorithm with sample access to X that computes an estimator $\tilde{\mu}$ in time t such that for a universal constant c we have*

$$i) \Pr[(1 - \epsilon)\mu \leq \tilde{\mu} \leq (1 + \epsilon)\mu] \geq 1 - \delta \quad \text{and} \quad ii) \Pr[t \geq c \cdot \epsilon^{-2} \mu^{-2} \rho_X \log(1/\delta)] \leq \delta.$$

We state now our key technical insight, on which we build upon our sublinear algorithm.

Lemma 16. *There is an algorithm that, given $A, B \in \mathbb{R}^{m \times n}$ with column adjacency arrays and $\|A - B\|_0 \geq 1$, and given $\epsilon > 0$, computes an estimator Z that satisfies w.h.p.*

$$(1 - \epsilon)\|A - B\|_0 \leq Z \leq (1 + \epsilon)\|A - B\|_0.$$

The algorithm runs w.h.p. in time $O(n + \epsilon^{-2} \frac{\|A\|_0 + \|B\|_0}{\|A - B\|_0} \log n)$.

Proof. By Lemma 1, after $O(n)$ preprocessing time we can sample a uniformly random non-zero entry from A or B in $O(1)$ time.

We consider the following random process:

1. Sample $C \in \{A, B\}$ such that $\Pr[C = A] = \frac{\|A\|_0}{\|A\|_0 + \|B\|_0}$ and $\Pr[C = B] = \frac{\|B\|_0}{\|A\|_0 + \|B\|_0}$.
2. Sample (i, j) uniformly at random from the non-zero entries of C
3. Return:

$$X = \begin{cases} 0, & \text{if } A_{ij} = B_{ij}; \\ 1/2, & \text{if } 0 \neq A_{ij} \neq B_{ij} \neq 0; \\ 1, & \text{if } A_{ij} \neq B_{ij} \text{ and either } A_{ij} \text{ or } B_{ij} \text{ equals } 0. \end{cases}$$

Observe that

$$\begin{aligned} \mathbb{E}[X] &= \sum_{(i,j): A_{ij} \neq B_{ij} = 0} \frac{\|A\|_0}{\|A\|_0 + \|B\|_0} \cdot \frac{1}{\|A\|_0} + \sum_{(i,j): 0 = A_{ij} \neq B_{ij}} \frac{\|B\|_0}{\|A\|_0 + \|B\|_0} \cdot \frac{1}{\|B\|_0} \\ &\quad + \sum_{(i,j): 0 \neq A_{ij} \neq B_{ij} \neq 0} \left(\frac{1}{2} \cdot \frac{\|A\|_0}{\|A\|_0 + \|B\|_0} \cdot \frac{1}{\|A\|_0} + \frac{1}{2} \cdot \frac{\|B\|_0}{\|A\|_0 + \|B\|_0} \cdot \frac{1}{\|B\|_0} \right) \\ &= \frac{\|A - B\|_0}{\|A\|_0 + \|B\|_0}. \end{aligned}$$

Straightforward checking shows that $X \in [0, 1]$ implies $\text{Var}[X] \leq \mathbb{E}[X]$, and thus

$$\rho_X = \max\{\text{Var}[X], \epsilon \cdot \mathbb{E}[X]\} \leq \mathbb{E}[X].$$

Setting $\delta = 1/\text{poly}(n)$ in Theorem 15, we can compute w.h.p. in time $O(\epsilon^{-2} \mathbb{E}[X]^{-1} \log n) = O(\epsilon^{-2} \frac{\|A\|_0 + \|B\|_0}{\|A - B\|_0} \log n)$ an estimator $(1 - \epsilon)\mathbb{E}[X] \leq \tilde{\mu} \leq (1 + \epsilon)\mathbb{E}[X]$. Then, w.h.p. the estimator $Z \stackrel{\text{def}}{=} (\|A\|_0 + \|B\|_0)\tilde{\mu}$ satisfies the statement. \square

We are now ready to prove Theorem 13.

Proof of Theorem 13. Let $B \stackrel{\text{def}}{=} A_{:,j} v^T$ and observe that $\|A - B\|_0 \geq \text{OPT}^{(1)} \geq 1$. Note that we implicitly have column adjacency arrays for B , since for any column c with $v_c = 0$ there are no non-zero entries in $B_{:,c}$, and for any column c with $v_c = 1$ the non-zero entries of $B_{:,c}$ are the same as for $A_{:,j}$. Hence, Lemma 14 and Lemma 16 are applicable.

We analyze the running time of Lemma 16. Note that if $\|B\|_0 \leq (1 + \psi)\|A\|_0$ then $\frac{\|A\|_0 + \|B\|_0}{\|A - B\|_0} \leq (2 + \psi)/\psi$, and otherwise, i.e. $(1 + \psi)\|A\|_0 \leq \|B\|_0$, we have

$$\|A - B\|_0 \geq \|B\|_0 - \|A\|_0 \geq \frac{\psi}{1 + \psi} \|B\|_0$$

and thus $\frac{\|A\|_0 + \|B\|_0}{\|A - B\|_0} \leq 2(1 + \psi)/\psi$. Hence, $\frac{\|A\|_0 + \|B\|_0}{\|A - B\|_0} < 4/\psi$, which yields with high probability time $O(n + \epsilon^{-2}\psi^{-1} \log n)$.

We execute in parallel the algorithms from Lemma 14 and Lemma 16. Once the faster algorithm outputs a solution, we terminate the execution of the slower one. Note that this procedure runs w.h.p in time $O(\min\{\|A\|_0, n + \epsilon^{-2}\psi^{-1} \log n\})$, and returns w.h.p. the desired estimator Y . \square

To implement Step 3 of Algorithm 2, we simply apply Theorem 13 with A , ϵ and $v = z^{(j)}$ to each sampled column $j \in \bigcup_{0 \leq i \leq \log n + 1} C^{(i)}$.

4.4 Analyzing the Runtime of Algorithm 2

Consider again Algorithm 2. In Steps 1, 2 and 2.1, from each of the $O(\log n)$ weight classes we sample $O(\epsilon^{-2} \log n)$ columns. In Step 2.2, for each sampled column we use Lemma 12, which takes time $O(\epsilon^{-2} n \log m)$ per column, or $O(\epsilon^{-4} n \log m \log^2 n)$ in total. Finally in Step 3, for each sampled column we use Theorem 13, which w.h.p. takes time $O(\min\{\|A\|_0, n + \epsilon^{-2}\psi^{-1} \log n\})$ per column, or $O(\min\{\|A\|_0, n + \epsilon^{-2}\psi^{-1} \log n\} \cdot \epsilon^{-2} \log^2 n)$ in total. Then, the total runtime is bounded by $O(\epsilon^{-4} n \log m \log^2 n + \min\{\|A\|_0 \epsilon^{-2} \log^2 n, \epsilon^{-4} \psi^{-1} \log^3 n\})$.

5 Algorithms for Boolean ℓ_0 -Rank-1

Given a matrix $A \in \{0, 1\}^{m \times n}$, the Boolean ℓ_0 -Rank-1 problem asks to find a matrix $A' \in \{0, 1\}^{m \times n}$ with rank 1, i.e. $A' = uv^T$ for some vectors $u \in \{0, 1\}^m$ and $v \in \{0, 1\}^n$, such that the difference between A and A' measured in ℓ_0 -distance is minimized. We denote the optimum value by

$$\text{OPT} = \text{OPT}_A \stackrel{\text{def}}{=} \min_{u \in \{0, 1\}^m, v \in \{0, 1\}^n} \|A - uv^T\|_0. \quad (16)$$

In practice, approximating a matrix A by a rank-1 matrix uv^T makes most sense if A is close to being rank-1. Hence, the above optimization problem is most relevant in the case $\text{OPT} \ll \|A\|_0$. For this reason, in this section we focus on the case $\text{OPT}/\|A\|_0 \leq \phi$ for sufficiently small $\phi > 0$. We prove the following.

Theorem 17. *Given $A \in \{0, 1\}^{m \times n}$ with row and column sums, and given $\phi \in (0, \frac{1}{80}]$ with $\text{OPT}/\|A\|_0 \leq \phi$, we can compute in time $O(\min\{\|A\|_0 + m + n, \phi^{-1}(m + n) \log(mn)\})$ vectors $\tilde{u} \in \{0, 1\}^m$ and $\tilde{v} \in \{0, 1\}^n$ such that w.h.p. $\|A - \tilde{u} \cdot \tilde{v}^T\|_0 \leq (1 + 5\phi)\text{OPT} + 37\phi^2 \|A\|_0$.*

Using Theorem 9, we can compute a $(2 + \epsilon)$ -approximation of OPT , and thus a $(2 + \epsilon)$ -approximation of the ratio $\text{OPT}/\|A\|_0$. Hence, combining Theorem 17 and Theorem 9, yields an algorithm that does not need ϕ as an input and computes a $(1 + 500\psi)$ -approximate solution of the Boolean ℓ_0 -Rank-1 problem.

Theorem 18. Given $A \in \{0, 1\}^{m \times n}$ with column adjacency arrays and with row and column sums, for $\psi = \text{OPT} / \|A\|_0$ we can compute w.h.p. in time $O(\min\{\|A\|_0 + m + n, \psi^{-1}(m + n)\} \cdot \log^3(mn))$ vectors $\tilde{u} \in \{0, 1\}^m$ and $\tilde{v} \in \{0, 1\}^n$ such that w.h.p. $\|A - \tilde{u} \cdot \tilde{v}^T\|_0 \leq (1 + 500\psi)\text{OPT}$.

Proof of Theorem 18. We compute a 3-approximation of OPT by applying Theorem 9 with $\epsilon = 0.1$. This yields a value ϕ satisfying $\psi \leq \phi \leq 3\psi$. If $\phi > 1/80$, then the 3-approximation is already good enough, since $\psi > 1/240$ and $1 + 500\psi > 3$. Otherwise, we run Theorem 17 with ϕ . Further, using $\phi^2 \|A\|_0 \leq 9\psi^2 \|A\|_0 = 9\psi\text{OPT}$, the total error is at most

$$(1 + 5\phi)\text{OPT} + 37\phi^2 \|A\|_0 \leq (1 + 15\psi)\text{OPT} + 37 \cdot 9\psi\text{OPT} \leq (1 + 500\psi)\text{OPT}.$$

A rough upper bound on the running time is $O(\min\{\|A\|_0 + m + n, \psi^{-1}(m + n)\} \cdot \log^3(mn))$. \square

A variant of the algorithm from Theorem 17 can also be used to solve the Boolean ℓ_0 -Rank-1 problem exactly. This yields the following theorem, which in particular shows that the problem is in polynomial time whenever $\text{OPT} \leq O(\sqrt{\|A\|_0} \log(mn))$.

Theorem 19. Given a matrix $A \in \{0, 1\}^{m \times n}$, if $\text{OPT}_A / \|A\|_0 \leq 1/240$ then we can exactly solve the Boolean ℓ_0 -Rank-1 problem in time $2^{O(\text{OPT}/\sqrt{\|A\|_0})} \cdot \text{poly}(mn)$.

The remainder of this section is devoted to the proof of Theorem 17 (and Theorem 19).

5.1 Preparations

Let $A \in \{0, 1\}^{m \times n}$ and fix an optimal solution u, v to the problem, realizing $\text{OPT} = \|A - uv^T\|_0$. Moreover, set $\alpha \stackrel{\text{def}}{=} \|u\|_0$ and $\beta \stackrel{\text{def}}{=} \|v\|_0$. We start with the following technical preparations.

Lemma 20. For any row $i \in [m]$ let x_i be the number of 0's in columns selected by v , i.e., $x_i \stackrel{\text{def}}{=} |\{j \in [n] \mid A_{i,j} = 0, v_j = 1\}|$, and let y_i be the number of 1's in columns not selected by v , i.e., $y_i \stackrel{\text{def}}{=} |\{j \in [n] \mid A_{i,j} = 1, v_j = 0\}|$. Let $R = \{i \in [m] \mid u_i = 1\}$ be the rows selected by u , and let $\bar{R} \stackrel{\text{def}}{=} [m] \setminus R$. Symmetrically, let C be the columns selected by v . Then we have

1. $\|A_{i,:}\|_0 = \beta - x_i + y_i$ for any $i \in [m]$,
2. $\text{OPT} = \sum_{i \in R} (x_i + y_i) + \sum_{i \in \bar{R}} (\beta - x_i + y_i)$,
3. $\text{OPT} \geq \sum_{i \in R} |x_i - y_i|$,
4. $x_i \leq \beta/2$ for any $i \in R$, and $x_i \geq \beta/2$ for any $i \in \bar{R}$,
5. $\text{OPT} \geq \sum_{i=1}^m \min\{\|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta|\}$,
6. $|\|A\|_0 - \alpha\beta| \leq \text{OPT}$,
7. If $\text{OPT} \leq \phi \|A\|_0$ then $(1 - \phi) \|A\|_0 \leq \alpha\beta \leq (1 + \phi) \|A\|_0$.

Proof. For (1), note that in the β columns C selected by v , row i has $\beta - x_i$ 1's, and in the remaining $n - |C|$ columns row i has y_i 1's. Hence, the total number of 1's in row i is $\|A_{i,:}\|_0 = \beta - x_i + y_i$.

(2) We split $\text{OPT} = \|A - uv^T\|_0$ into a sum over all rows, so that $\text{OPT} = \sum_{i=1}^m \|A_{i,:} - u_i v^T\|_0$. For $i \in \bar{R}$, the i -th term of this sum is simply $\|A_{i,:}\|_0 = \beta - x_i + y_i$. For $i \in R$, the i -th term is $\|A_{i,:} - v^T\|_0 = x_i + y_i$.

(3) follows immediately from (2).

(4) follows from (2), since for $x_i > \beta/2$ and $i \in R$ we can change u_i from 1 to 0, reducing the contribution of row i from $x_i + y_i$ to $\beta - x_i + y_i$, which contradicts optimality of OPT.

For (5), we use that $x_i + y_i \geq |x_i - y_i| = |\|A_{i,:}\|_0 - \beta|$ by (1), and

$$\text{OPT} = \sum_{i \in R} (x_i + y_i) + \sum_{i \in \bar{R}} (\beta - x_i + y_i) = \sum_{i \in R} (x_i + y_i) + \sum_{i \in \bar{R}} \|A_{i,:}\|_0.$$

(6) is shown similarly to (5) by noting that

$$\begin{aligned} \text{OPT} &= \sum_{i \in R} (x_i + y_i) + \sum_{i \in \bar{R}} (\beta - x_i + y_i) \geq \sum_{i \in R} |\|A_{i,:}\|_0 - \beta| + \sum_{i \in \bar{R}} \|A_{i,:}\|_0 \\ &\geq \sum_{i \in R} (\|A_{i,:}\|_0 - \beta) + \sum_{i \in \bar{R}} \|A_{i,:}\|_0 = \|A\|_0 - \alpha\beta, \end{aligned}$$

and similarly

$$\text{OPT} \geq \sum_{i \in R} |\|A_{i,:}\|_0 - \beta| + \sum_{i \in \bar{R}} \|A_{i,:}\|_0 \geq \sum_{i \in R} (\beta - \|A_{i,:}\|_0) - \sum_{i \in \bar{R}} \|A_{i,:}\|_0 = \alpha\beta - \|A\|_0.$$

Finally, (7) follows immediately from (6) by plugging in the upper bound $\text{OPT} \leq \phi \|A\|_0$. \square

5.2 Approximating α and β

We now show how to approximate $\alpha = \|u\|_0$ and $\beta = \|v\|_0$, where u, v is an optimal solution.

Lemma 21. *Given $A \in \{0, 1\}^{m \times n}$ and $\phi \in (0, 1/30]$ with $\text{OPT}/\|A\|_0 \leq \phi$, we can compute in time $O(\|A\|_0 + m + n)$ an integer $\tilde{\beta} \in [m]$ with*

$$\frac{1 - 3\phi}{1 - \phi} \beta \leq \tilde{\beta} \leq \frac{1 + \phi}{1 - \phi} \beta.$$

Symmetrically, we can approximate α by $\tilde{\alpha}$. If we are additionally given the number of 1's in each row and column, then the running time becomes $O(m + n)$.

Proof. Let

$$\Lambda \stackrel{\text{def}}{=} \min_{\beta' \in [m]} \sum_{i=1}^m \min \left\{ \|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta'| \right\},$$

and let $\tilde{\beta}$ be the value of β' realizing Λ .

We first verify the approximation guarantee. Consider the set of rows R selected by u . Let x_i, y_i for $i \in R$ be as in Lemma 20. Then we have

$$\Lambda \geq \sum_{i \in R} \min \left\{ \|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \tilde{\beta}| \right\} = \sum_{i \in R} \min \left\{ \beta + y_i - x_i, |\beta - \tilde{\beta} + y_i - x_i| \right\},$$

where we used Lemma 20.1. Assume for the sake of contradiction that $|\beta - \tilde{\beta}| > \frac{2\phi}{1-\phi} \beta$. Since $|x - y| \geq |x| - |y|$ for any numbers x, y , we obtain

$$|\beta - \tilde{\beta} + y_i - x_i| \geq |\beta - \tilde{\beta}| - |x_i - y_i| > \frac{2\phi}{1-\phi} \beta - |x_i - y_i|.$$

Similarly, we have $\beta + y_i - x_i > \frac{2\phi}{1-\phi}\beta - |x_i - y_i|$. Hence,

$$\Lambda > \sum_{i \in R} \left(\frac{2\phi\beta}{1-\phi} - |x_i - y_i| \right) \geq \frac{2\phi\beta}{1-\phi}|R| - \text{OPT},$$

where we used Lemma 20.3. Since R is the set of rows selected by u , we have $|R| = \alpha$. By Lemma 20.7, we have $\text{OPT} \leq \phi \|A\|_0 \leq \frac{\phi}{1-\phi}\alpha\beta$. Together, this yields $\Lambda > \text{OPT}$, contradicting

$$\Lambda \leq \sum_{i=1}^m \min\{\|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta|\} \leq \text{OPT}$$

by Lemma 20.5. Hence, $|\beta - \tilde{\beta}| \leq \frac{2\phi}{1-\phi}\beta$.

It remains to design a fast algorithm. We first compute all numbers $\|A_{i,:}\|_0$ in time $O(\|A\|_0)$ (this step can be skipped if we are given these numbers as input). We sort these numbers, obtaining a sorted order $\|A_{\pi(1),:}\|_0 \leq \dots \leq \|A_{\pi(m),:}\|_0$. Using counting sort, this takes time $O(m + n)$. We precompute prefix sums $P(k) \stackrel{\text{def}}{=} \sum_{\ell=1}^k \|A_{\pi(\ell),:}\|_0$, which allows us to evaluate in constant time any sum

$$\sum_{\ell=x}^y \|A_{\pi(\ell),:}\|_0 = P(y) - P(x-1).$$

Finally, we precompute the inverse

$$\ell(\beta') \stackrel{\text{def}}{=} \max\{\ell \mid \|A_{\pi(\ell),:}\|_0 \leq \beta'\},$$

or $\ell(\beta') = 0$ if there is no ℓ with $\|A_{\pi(\ell),:}\|_0 \leq \beta'$. By a simple sweep, all values $\ell(\beta')$ can be computed in total time $O(m + n)$.

Note that for any fixed β' and row i , the term realizing $\min\{\|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta'|\}$ is equal to: (a) $\|A_{i,:}\|_0$ if $\|A_{i,:}\|_0 \leq \beta'/2$; (b) $\beta' - \|A_{i,:}\|_0$, if $\beta'/2 < \|A_{i,:}\|_0 \leq \beta'$; and (c) $\|A_{i,:}\|_0 - \beta'$, if $\|A_{i,:}\|_0 > \beta'$. Hence, we obtain

$$\begin{aligned} & \sum_{i=1}^n \min\left\{\|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta'|\right\} \\ &= \left(\sum_{i=1}^{\ell(\beta'/2)} \|A_{\pi(i),:}\|_0 \right) + \left(\sum_{i=\ell(\beta'/2)+1}^{\ell(\beta')} \beta' - \|A_{\pi(i),:}\|_0 \right) + \left(\sum_{i=\ell(\beta')+1}^n \|A_{\pi(i),:}\|_0 - \beta' \right) \\ &= P(n) - 2[P(\ell(\beta')) - P(\ell(\beta'/2))] - [n + \ell(\beta'/2) - 2\ell(\beta')] \beta'. \end{aligned}$$

This shows that after the above precomputation the sum $\sum_{j=1}^n \min\{\|A_{i,:}\|_0, |\|A_{i,:}\|_0 - \beta'|\}$ can be evaluated in time $O(1)$ for any β' . Minimizing over all $\beta' \in [m]$ yields $\tilde{\beta}$. This finishes our algorithm, which runs in total time $O(\|A\|_0 + m + n)$, or $O(m + n)$ if we are given the number of 1's in each row and column. \square

5.3 The Algorithm

We now design an approximation algorithm for the Boolean ℓ_0 -Rank-1 problem that will yield Theorem 17. We present the pseudocode of this Algorithm 4 below.

Algorithm 4 Boolean ℓ_0 -Rank-1 With Small Optimal Value

Input: $A \in \{0, 1\}^{m \times n}$ and $\phi \in (0, 1/80]$ such that $\text{OPT}/\|A\|_0 \leq \phi$.

Output: Vectors $\tilde{u} \in \{0, 1\}^m$, $\tilde{v} \in \{0, 1\}^n$ such that $\|A - \tilde{u} \cdot \tilde{v}^T\|_0 \leq (1 + \epsilon)\text{OPT}$.

1. Compute approximations $\frac{1-3\phi}{1-\phi}\alpha \leq \tilde{\alpha} \leq \frac{1+\phi}{1-\phi}\alpha$ and $\frac{1-3\phi}{1-\phi}\beta \leq \tilde{\beta} \leq \frac{1+\phi}{1-\phi}\beta$ using Lemma 21.
Initialize $R^R \stackrel{\text{def}}{=} [m]$, $C^R \stackrel{\text{def}}{=} [n]$, $R^S \stackrel{\text{def}}{=} \emptyset$, $C^S \stackrel{\text{def}}{=} \emptyset$.
 2. For any row i , if $\|A_{i,:}\|_0 < \frac{1-\phi}{1+\phi} \cdot \frac{\tilde{\beta}}{2}$ then set $\tilde{u}_i = 0$ and remove i from R^R .
For any column j , if $\|A_{:,j}\|_0 < \frac{1-\phi}{1+\phi} \cdot \frac{\tilde{\alpha}}{2}$ then set $\tilde{v}_j = 0$ and remove j from C^R .
 3. For any $i \in R^R$ compute an estimate X_i with $|X_i - \|A_{i,C^R}\|_0| \leq \frac{1}{9}|C^R|$.
For any $j \in C^R$ compute an estimate Y_j with $|Y_j - \|A_{R^R,j}\|_0| \leq \frac{1}{9}|R^R|$.
 4. For any $i \in R^R$, if $X_i > \frac{2}{3}\tilde{\beta}$ then set $\tilde{u}_i = 1$ and add i to R^S .
For any $j \in C^R$, if $Y_j > \frac{2}{3}\tilde{\alpha}$ then set $\tilde{v}_j = 1$ and add j to C^S .
 5. For any $i \in R^R \setminus R^S$, compute an estimate X'_i with $|X'_i - \|A_{i,C^S}\|_0| \leq \phi|C^S|$,
For any $j \in C^R \setminus C^S$, compute an estimate Y'_j with $|Y'_j - \|A_{R^S,j}\|_0| \leq \phi|R^S|$.
 6. For any $i \in R^R \setminus R^S$, set $\tilde{u}_i = 1$ if $X'_i \geq |C^S|/2$ and 0 otherwise,
For any $j \in C^R \setminus C^S$, set $\tilde{v}_j = 1$ if $Y'_j \geq |R^S|/2$ and 0 otherwise.
 7. **Return** (\tilde{u}, \tilde{v}) .
-

Running Time By Lemma 21, Step 1 runs in time $O(\|A\|_0 + m + n)$, or in time $O(m + n)$ if we are given the number of 1's in each row and column. Steps 2, 4, and 6 clearly run in time $O(m + n)$. For steps 3 and 5, there are two ways to implement them.

Variant (1) is an exact algorithm. We enumerate all nonzero entries of A and count how many contribute to the required numbers $\|A_{i,C^R}\|_0, \|A_{R^R,j}\|_0$ etc. This takes total time $O(\|A\|_0)$, and hence the total running time of the algorithm is $O(\|A\|_0 + m + n)$.

Variant (2) uses random sampling. In order to estimate $\|A_{i,C^R}\|_0$, consider a random variable Z that draws a uniformly random column $j \in C^R$ and returns 1 if $A_{i,j} \neq 0$ and 0 otherwise. Then $\mathbb{E}[Z] = \|A_{i,C^R}\|_0/|C^R|$. Taking independent copies Z_1, \dots, Z_ℓ of Z , where $\ell = \Theta(\log(mn)/\delta^2)$ with sufficiently large hidden constant, a standard Chernoff bound argument shows that w.h.p.

$$\left| (Z_1 + \dots + Z_\ell) \cdot \frac{|C^R|}{\ell} - \|A_{i,C^R}\|_0 \right| \leq \delta \cdot |C^R|,$$

which yields the required approximation. For Step 3 we use this procedure with $\delta = \frac{1}{9}$ and obtain running time $O(\log(mn))$ per row and column, or $O((m + n) \log(mn))$ in total. For Step 5 we use $\delta = \phi$, resulting in time $O(\phi^{-2} \log(mn))$ for computing one estimate X'_i or Y'_j . By Claim 25 below there are only $O(\phi(m + n))$ rows and columns in $R^R \setminus R^S$ and $C^R \setminus C^S$, and hence the total running time for Step 5 is $O(\phi^{-1}(m + n) \log(mn))$. This dominates the total running time.

Combining both variants, we obtain the claimed running time of

$$O(\min\{\|A\|_0 + m + n, \phi^{-1}(m + n) \log(mn)\}).$$

Correctness In the following we analyze the correctness of the above algorithm.

Claim 22. *For any row i deleted in Step 2 we have $\tilde{u}_i = u_i$. Symmetrically, for any column j deleted in Step 2 we have $\tilde{v}_j = v_j$.*

Proof. If row i is deleted, then by the approximation guarantee of $\tilde{\beta}$ we have

$$\|A_{i,:}\|_0 < \frac{1-\phi}{1+\phi} \cdot \frac{\tilde{\beta}}{2} \leq \frac{\beta}{2}$$

Note that for x_i (the number of 0's in row i in columns selected by v) we have $x_i \geq \beta - \|A_{i,:}\|_0$. Together, we obtain $x_i > \beta/2$, and thus row i cannot be selected by u , by Lemma 20.4. Hence, we have $u_i = 0 = \tilde{u}_i$. The statement for the columns is symmetric. \square

Claim 23. *After Step 2, it holds for the remaining rows R^R and columns C^R that*

$$|R^R| \leq \left(1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi}\right) \alpha \quad \text{and} \quad |C^R| \leq \left(1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi}\right) \beta.$$

Proof. By Claim 22 the α rows R selected by u remain. We split the rows R^R remaining after Step 2 into $R \cup R'$, and bound $|R'|$ from above. Since any $i \in R'$ is not selected by u , it contributes $\|A_{i,:}\|_0$ to OPT. Note that

$$\|A_{i,:}\|_0 \geq \frac{1-\phi}{1+\phi} \cdot \frac{\tilde{\beta}}{2} \geq \frac{1-\phi}{1+\phi} \cdot \frac{1-3\phi}{1-\phi} \cdot \frac{\beta}{2} = \frac{1-3\phi}{1+\phi} \cdot \frac{\beta}{2},$$

and thus $|R'| \leq \text{OPT} \cdot \frac{1+\phi}{1-3\phi} \cdot \frac{2}{\beta}$. Since

$$\text{OPT} \leq \phi \|A\|_0 \leq \frac{\phi}{1-\phi} \cdot \alpha \beta$$

by Lemma 20.7, we obtain $|R'| \leq \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi} \cdot \alpha$. Thus, we have in total

$$|R^R| = |R| + |R'| \leq \left(1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi}\right) \alpha.$$

The statement for the columns is symmetric. \square

Claim 24. *The rows and columns selected in Step 4 are also selected by the optimal solution u, v , i.e., for any $i \in R^S$ we have $u_i = 1$ and for any $j \in C^S$ we have $v_j = 1$.*

Proof. If row i is selected in Step 4, then we have by the approximation guarantee of X_i , definition of Step 4, Claim 23, and Lemma 21

$$\begin{aligned} \|A_{i,C^R}\|_0 &\geq X_i - \frac{1}{9}|C^R| > \frac{2}{3}\tilde{\beta} - \frac{1}{9}\left(1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi}\right) \beta \\ &\geq \frac{2}{3} \cdot \frac{1-3\phi}{1-\phi} \beta - \frac{1}{9}\left(1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi}\right) \beta. \end{aligned}$$

It is easy to see that for sufficiently small $\phi \geq 0$ this yields

$$\|A_{i,C^R}\|_0 > \frac{\beta}{2} + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi} \beta.$$

One can check that $0 \leq \phi \leq 1/80$ is sufficient. Since there are $|C^R| \leq (1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi})\beta$ columns remaining, in particular the β columns $C \subseteq C^R$ which are selected by v , we obtain

$$\|A_{i,C}\|_0 \geq \|A_{i,C^R}\|_0 - (|C^R| - \beta) > \beta/2.$$

By Lemma 20.4, we thus obtain that row i is selected by the optimal u . The statement for the columns is symmetric. \square

Claim 25. *After Step 4 there are $|R^R \setminus R^S| \leq 6\phi\alpha$ remaining rows and $|C^R \setminus C^S| \leq 6\phi\beta$ remaining columns.*

Proof. After Step 4, every remaining row i , for any $0 \leq \phi \leq 1/80$, satisfies

$$\|A_{i,:}\|_0 \geq \frac{1-\phi}{1+\phi} \cdot \frac{\tilde{\beta}}{2} \geq \frac{1-\phi}{1+\phi} \cdot \frac{1-3\phi}{1-\phi} \cdot \frac{\beta}{2} \geq \frac{2}{5}\beta,$$

Moreover, each such row satisfies

$$\|A_{i,C^R}\|_0 \leq X_i + \frac{1}{9}|C^R| \leq \frac{2}{3}\tilde{\beta} + \frac{1}{9}|C^R|,$$

which together with $\tilde{\beta} \leq \frac{1+\phi}{1-\phi}\beta$ (Lemma 21) and $|C^R| \leq (1 + \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi})\beta$ (Claim 23) yields

$$\|A_{i,C^R}\|_0 \leq \left(\frac{2}{3} \cdot \frac{1+\phi}{1-\phi} + \frac{1}{9} + \frac{1}{9} \cdot \frac{1+\phi}{1-3\phi} \cdot \frac{2\phi}{1-\phi} \right) \beta.$$

It is easy to see that for sufficiently small $\phi \geq 0$ we have $\|A_{i,C^R}\|_0 \leq \frac{4}{5}\beta$, and it can be checked that $0 \leq \phi \leq 1/80$ is sufficient.

If i is not selected by u , then its contribution to OPT is $\|A_{i,:}\|_0 \geq \frac{2}{5}\beta$. If i is selected by u , then since $C \subseteq C^R$ its contribution to OPT is at least

$$\beta - \|A_{i,C}\|_0 \geq \beta - \|A_{i,C^R}\|_0 \geq \beta - \frac{4}{5}\beta = \frac{1}{5}\beta.$$

Thus, the number of remaining rows is at most

$$\frac{\text{OPT}}{\beta/5} \leq \frac{5\phi\alpha\beta}{(1-\phi)\beta} \leq 6\phi\alpha,$$

where we used Lemma 20.7. The statement for the columns is symmetric. \square

We are now ready to prove correctness of Algorithm 4.

Proof of Theorem 17. The rows and columns removed in Step 2 are also not picked by the optimal solution, by Claim 22. Hence, in the region $([m] \setminus R^R) \times [n]$ and $[m] \times ([n] \setminus C^R)$ we incur the same error as the optimal solution. The rows and columns chosen in Step 4 are also picked by the optimal solution, by Claim 24. Hence, in the region $R^S \times C^S$ we incur the same error as the optimal solution. We split the remaining matrix into three regions: $(R^R \setminus R^S) \times C^S$, $R^S \times (C^R \setminus C^S)$, and $(R^R \setminus R^S) \times (C^R \setminus C^S)$.

In the region $(R^R \setminus R^S) \times C^S$ we compute for any row $i \in R^R \setminus R^S$ an additive $\phi|C|$ -approximation X'_i of $\|A_{i,C}\|_0$, and we pick row i iff $X'_i \geq |C|/2$. In case $|\|A_{i,C}\|_0 - |C|/2| > \phi|C|$, we have $X'_i \geq |C|/2$ if and only if $\|A_{i,C}\|_0 \geq |C|/2$, and thus our choice for row i is optimal, restricted to

region $(R^R \setminus R^S) \times C^S$. Otherwise, if $||A_{i,C}||_0 - |C|/2| \leq \phi|C|$, then no matter whether we choose row i or not, we obtain approximation ratio

$$\frac{|C|/2 + \phi|C|}{|C|/2 - \phi|C|} = \frac{1 + 2\phi}{1 - 2\phi} \leq 1 + 5\phi,$$

restricted to region $(R^R \setminus R^S) \times C^S$. The region $R^S \times (C^R \setminus C^S)$ is symmetric.

Finally, in region $(R^R \setminus R^S) \times (C^R \setminus C^S)$ we pessimistically assume that every entry is an error. By Claim 25 and Lemma 20.7, this submatrix has size at most

$$6\phi\alpha \cdot 6\phi\beta \leq 36\phi^2(1 + \phi) \|A\|_0 \leq 37\phi^2 \|A\|_0.$$

In total, over all regions, we computed vectors \tilde{u}, \tilde{v} such that

$$\|A - \tilde{u}\tilde{v}^T\|_0 \leq (1 + 5\phi)\text{OPT} + 37\phi^2 \|A\|_0.$$

This completes the correctness prove of Algorithm 4. \square

5.4 The Exact Algorithm

We now prove Theorem 19, i.e., given a matrix $A \in \{0, 1\}^{m \times n}$ we exactly solve the Boolean ℓ_0 -Rank-1 problem in time $2^{O(\text{OPT}/\sqrt{\|A\|_0})} \cdot \text{poly}(mn)$ if we have $\psi \stackrel{\text{def}}{=} \text{OPT}/\|A\|_0 \leq 1/240$. This algorithm builds upon the algorithmic results established in Theorem 18 and Theorem 8, and it consists of the following three phases:

1. Run the algorithm in Theorem 8 to compute a 3-approximation of $\psi = \text{OPT}/\|A\|_0$, i.e. a number $\phi \in [\psi, 3\psi]$.
2. Run Steps 1-4 of Algorithm 4, resulting in selected rows R^S and columns C^S , and undecided rows $R' = R^R \setminus R^S$ and columns $C' = C^R \setminus C^S$. As shown above, the choices made by these steps are optimal.
3. For the remaining rows R' and columns C' we use brute force to find the optimum solution. Specifically, assume without loss of generality that $|R'| \leq |C'|$. Enumerate all Boolean vectors $u' \in \{0, 1\}^{R'}$. For each u' , set $\tilde{u}_i = u'_i$ for all $i \in R'$ to complete the specification of a vector $\tilde{u} \in \{0, 1\}^m$. We can now find the optimal choice of vector \tilde{v} in polynomial time, since the optimal choice is to set $\tilde{v}_j = 1$ iff column $A_{:,j}$ has more 1's than 0's in the support of \tilde{u} . Since some u' gives rise to the optimal vector $\tilde{u} = u$, we solve the Boolean ℓ_0 -Rank-1 problem exactly.

To analyze the running time, note that by Claim 25 we have

$$\min\{|R'|, |C'|\} \leq 6\phi \min\{\alpha, \beta\} \leq 6\phi\sqrt{\alpha\beta}.$$

By Lemma 20.7 and $\phi \leq 3\psi$, we obtain $\min\{|R'|, |C'|\} = O(\psi\sqrt{\|A\|_0})$. Hence, we enumerate $2^{O(\psi\sqrt{\|A\|_0})} = 2^{O(\text{OPT}/\sqrt{\|A\|_0})}$ vectors u' , and the total running time is $2^{O(\text{OPT}/\sqrt{\|A\|_0})} \cdot \text{poly}(mn)$. This completes the proof of Theorem 19.

6 Sample Complexity Lower Bound for Boolean ℓ_0 -Rank-1

We now give a lower bound of $\Omega(n/\phi)$ on the number of samples of any $(1 + O(\phi))$ -approximation algorithm for the Boolean ℓ_0 -Rank-1 problem, where $\phi \geq \text{OPT}/\|A\|_0$ as before.

Theorem 6.1. *Let $C \geq 1$. Given an $n \times n$ binary matrix A with column adjacency arrays and with row and column sums, and given $\sqrt{\log(n)/n} \ll \phi \leq 1/100C$ such that $\text{OPT}/\|A\|_0 \leq \phi$, computing a $(1 + C\phi)$ -approximation of the Boolean ℓ_0 -Rank-1 problem requires to read $\Omega(n/\phi)$ entries of A (in the worst case over A).*

6.1 Core Probabilistic Result

The technical core of our argument is the following lemma.

Lemma 26. *Let $\phi \in (0, 1/2)$. Let X_1, \dots, X_k be binary random variables with expectations p_1, \dots, p_k , where $p_i \in \{1/2 - \phi, 1/2 + \phi\}$ for each i . Let \mathcal{A} be an algorithm which can adaptively obtain any number of samples of each random variable, and which outputs bits b_i for every $i \in [1 : k]$. Suppose that with probability at least 0.95 over the joint probability space of \mathcal{A} and the random samples, \mathcal{A} outputs for at least a 0.95 fraction of all i that $b_i = 1$ if $p_i = 1/2 + \phi$ and $b_i = 0$ otherwise. Then, with probability at least 0.05, \mathcal{A} makes $\Omega(k/\phi^2)$ samples in total, asymptotically in k .*

Proof. Consider the following problem P : let X be a binary random variable with expectation p drawn uniformly in $\{1/2 - \phi, 1/2 + \phi\}$. It is well-known that any algorithm which, with probability at least 0.6, obtains samples from X and outputs 0 if $p = 1/2 - \phi$ and outputs 1 if $p = 1/2 + \phi$, requires $\Omega(1/\phi^2)$ samples; see, e.g., Theorem 4.32 of [BY02]. Let $c > 0$ be such that c/ϕ^2 is a lower bound on the number of samples for this problem P .

Let \mathcal{A} be an algorithm solving the problem in the lemma statement. Since \mathcal{A} succeeds with probability at least 0.95 in obtaining the guarantees of the lemma for given sequence p_1, \dots, p_k , it also succeeds with this probability when (p_1, \dots, p_k) is drawn from the uniform distribution on $\{1/2 - \phi, 1/2 + \phi\}^k$.

Suppose, towards a contradiction, that \mathcal{A} takes less than $0.05 \cdot ck/\phi^2$ samples with probability at least 0.95. By stopping \mathcal{A} before taking $0.05 \cdot ck/\phi^2$ samples, we obtain an algorithm A' that always takes less than $0.05 \cdot ck/\phi^2$ samples. By the union bound, A' obtains the guarantees of the lemma for the output bits b_i with probability at least 0.9, over the joint probability space of A' and the random samples.

Note that the expected number of samples A' takes from a given X_i is less than $0.05 \cdot c/\phi^2$. By Markov's inequality, for a 0.95 fraction of indices i , A' takes less than c/ϕ^2 samples from X_i . We say that i is *good* if A' takes less than c/ϕ^2 samples from X_i and the output bit b_i is correct. By union bound, at least a $1 - (1 - 0.9) - (1 - 0.95) = 0.85$ fraction of indices i is good.

Since (p_1, \dots, p_k) is drawn from the uniform distribution on $\{1/2 - \phi, 1/2 + \phi\}^k$, with probability at least 0.95 the number $k_+ = |\{i : p_i = 1/2 + \phi\}|$ satisfies $0.45k \leq k_+ \leq 0.55k$ (for sufficiently large k). This implies that a 0.65 fraction of indices $\{i : p_i = 1/2 + \phi\}$ is good, as otherwise the number of bad i 's is at least $(1 - 0.65) \cdot 0.45k > 0.15k$. Similarly, a 0.65 fraction of indices $\{i : p_i = 1/2 - \phi\}$ is good.

Given an instance of problem P with random variable X and expectation p , we choose a uniformly random $i \in [k]$, and set $X_i = X$. For $j \neq i$, we independently and uniformly at random choose $p_j \in \{1/2 - \phi, 1/2 + \phi\}$. We then run algorithm A' . Whenever A' samples from X_i , we

sample a new value of X as in problem P . Whenever A' samples from X_j for $j \neq i$, we flip a coin with probability p_j and report the output to A' . If A' takes c/ϕ^2 samples from X_i , then we abort, thus ensuring that A' always takes less than c/ϕ^2 samples from $X_i = X$. Observe that the input to A' is a sequence of random variables X_1, \dots, X_k with expectations p_1, \dots, p_k which are independent and uniformly distributed in $\{1/2 - \phi, 1/2 + \phi\}$. In particular, except for their expectation these random variables are indistinguishable.

We now condition on $0.45k \leq k_+ \leq 0.55k$, which has success probability at least 0.95 for sufficiently large k . Then no matter whether $p_i = 1/2 + \phi$ or $p_i = 1/2 - \phi$, at least a 0.65 fraction of indices j with $p_j = p_i$ is good. Since i was chosen to be a uniformly random position independently of the randomness of the sampling and the algorithm A' , and the X_j with $p_j = p_i$ are indistinguishable, with probability at least 0.65 index i is good. In this case, A' takes less than c/ϕ^2 samples from $X_i = X$ and correctly determines the output bit b_i , i.e., whether $p_i = 1/2 + \phi$. As by union bound the total success probability is $1 - (1 - 0.65) - (1 - 0.95) = 0.6$, this contradicts the requirement of c/ϕ^2 samples mentioned above for solving P . Hence, the assumption was wrong, and \mathcal{A} takes $\Omega(k/\phi^2)$ samples with probability at least 0.05. \square

We start with a simplified version of our result, where we only have random access to the matrix entries. Below we extend this lower bound to the situation where we even have random access to the adjacency lists of all rows and columns.

Theorem 27. *Let $C \geq 1$. Given an $n \times n$ binary matrix A by random access to its entries, and given $\sqrt{\log(n)/n} \ll \phi \leq 1/100C$ such that $\text{OPT}/\|A\|_0 \leq \phi$, computing a $(1 + C\phi)$ -approximation of OPT requires to read $\Omega(n/\phi)$ entries of A (in the worst case over A).*

Proof. Set $\phi' \stackrel{\text{def}}{=} 25C\phi$ and $k \stackrel{\text{def}}{=} \phi n/2$. As in Lemma 26, consider binary random variables X_1, \dots, X_k with expectations p_1, \dots, p_k , where $p_i \in \{1/2 - \phi', 1/2 + \phi'\}$ for each i . We (implicitly) construct an $n \times n$ matrix A as follows. For ever $k < i \leq n$, $1 \leq j \leq n$ we set $A_{i,j} \stackrel{\text{def}}{=} 1$. For any $1 \leq i \leq k$, $1 \leq j \leq n$ we sample a bit $b_{i,j}$ from X_i and set $A_{i,j} \stackrel{\text{def}}{=} b_{i,j}$. Note that we can run any Boolean ℓ_0 -Rank-1 algorithm implicitly on A : whenever the algorithm reads an entry $A_{i,j}$ we sample a bit from X_i to determine the entry (and we remember the entry for possible further accesses).

Let us determine the optimal solution for A . Note that for each $i > k$, since the row $A_{i,:}$ is all-ones, it is always better to pick this row than not to pick it, and thus without loss of generality any solution u, v has $u_i = 1$. Similarly, for any j , since the column $A_{:,j}$ has $n - k > n/2$ 1's in rows picked by u , it is always better to pick the column than not to pick it, and thus $v_j = 1$, i.e., v is the all-ones vector. Hence, the only choice is for any $1 \leq i \leq k$ to pick or not to pick row i . Note that no matter whether we pick these rows or not, the total error is at most $\phi n^2/2$, since these rows in total have $kn = \phi n^2/2$ entries, and all remaining entries of A are correctly recovered by the product uv^T by the already chosen entries of u and v . Hence, $\text{OPT} \leq \phi n^2/2$, and since $\|A\|_0 \geq (n - k)n \geq n^2/2$, we obtain, as required, $\text{OPT}/\|A\|_0 \leq \phi$.

Now consider the rows $1 \leq i \leq k$ more closely. Since v is the all-ones vector, not picking row i incurs cost for each 1 in the row, which is cost $\|A_{i,:}\|_0$, while picking row i incurs cost for each 0 in the row, which is cost $n - \|A_{i,:}\|_0$. Note that the expected number of 1's in row $1 \leq i \leq k$ is $p_i n$. The Chernoff bound yields concentration: We have w.h.p. $|\|A_{i,:}\|_0 - p_i n| \leq 0.01 \cdot \phi' n$, where we used $\phi' \gg \sqrt{\log(n)/n}$. In the following we condition on this event and thus drop “w.h.p.” from our statements. In particular, for any i with $p_i = 1/2 + \phi'$ we have $\|A_{i,:}\|_0 \geq (1/2 + 0.99\phi')n$, and for any i with $p_i = 1/2 - \phi'$ we have $\|A_{i,:}\|_0 \leq (1/2 - 0.99\phi')n$.

By picking all rows $i \leq k$ with $p_i = 1/2 + \phi'$ and not pick the rows with $p_i = 1/2 - \phi'$, we see that $\text{OPT} \leq (1/2 - 0.99\phi')kn$. Now consider a solution u that among the rows $1 \leq i \leq k$ with $p_i = 1/2 + \phi'$ picks g_+ many and does not pick b_+ many. Similarly, among the rows with $p_i = 1/2 - \phi'$ it picks g_- and does not pick b_- . Note that each of the g_+ “good” rows incurs cost

$$n - \|A_{i:}\|_0 \geq n - (1/2 + 1.01\phi')n = (1/2 - 1.01\phi')n.$$

Each of the b_+ “bad” rows incurs a cost of $\|A_{i:}\|_0 \geq (1/2 + 0.99\phi')n$. Similar statements hold for g_- and b_- , and thus for $g \stackrel{\text{def}}{=} g_+ + g_-$ and $b \stackrel{\text{def}}{=} b_+ + b_-$, with $g + b = k$, we obtain a total cost of

$$\begin{aligned} \|A - uv^T\|_0 &\geq g \cdot (1/2 - 1.01\phi')n + b \cdot (1/2 + 0.99\phi')n \\ &= k(1/2 - 0.99\phi')n + 2b\phi'n - 0.02k\phi'n \\ &\geq \text{OPT} + 2b\phi'n - 0.02\phi'kn. \end{aligned}$$

If $b \geq 0.02k$, then

$$\|A - uv^T\|_0 \geq \text{OPT} + 0.02\phi'kn \geq (1 + 0.04\phi')\text{OPT}.$$

By contraposition, if we compute a $(1 + 0.04\phi' = 1 + C\phi)$ -approximation on A , then $b \leq 0.02k$, and thus the vector u correctly identifies for at least a 0.98 fraction of the random variables X_i whether $p_i = 1/2 + \phi'$ or $p_i = 1/2 - \phi'$. Since this holds w.h.p., by Lemma 26 we need $\Omega(k/\phi'^2) = \Omega(n/(\phi C^2))$ samples from the variables X_i , and thus $\Omega(n/(\phi C^2))$ reads in A . Since $C \geq 1$ is constant, we obtain a lower bound of $\Omega(n/\phi)$. This lower bound holds in expectation over the constructed distribution of A -matrices, and thus also in the worst case over A . \square

6.2 Hard Instance

The construction in Theorem 27 does not work in the case when we have random access to the adjacency lists of the rows, since this allows us to quickly determine the numbers of 1's per row, which is all we need to know in order to decide whether we pick a particular row in the matrix constructed above. We overcome this issue, using the adapted construction in Theorem 6.1.

We now present the correctness of Theorem 6.1.

Proof of Theorem 6.1. We assume that n is even. Let $\phi', k, X_1, \dots, X_k, p_1, \dots, p_k$ be as in the proof of Theorem 27. We adapt the construction of the matrix A as follows. For any $2k < i \leq n$, $1 \leq j \leq n/2$ we set $A_{i,2j} \stackrel{\text{def}}{=} 1$ and $A_{i,2j-1} \stackrel{\text{def}}{=} 0$. For any $1 \leq i \leq k$, $1 \leq j \leq n/2$ we sample a bit $b_{i,j}$ from X_i and set $A_{2i,2j} \stackrel{\text{def}}{=} A_{2i-1,2j-1} \stackrel{\text{def}}{=} b_{i,j}$ and $A_{2i-1,2j} \stackrel{\text{def}}{=} A_{2i,2j-1} \stackrel{\text{def}}{=} 1 - b_{i,j}$.

As before, when running any Boolean ℓ_0 -Rank-1 algorithm on A we can easily support random accesses to entries $A_{i,j}$, by sampling from $X_{\lceil i/2 \rceil}$ to determine the entry (and remembering the sampled bit for possible further accesses). Furthermore, we can now allow random accesses to the *adjacency arrays* of rows and columns. Specifically, if we want to determine the ℓ -th 1 in row $i \leq 2k$, we know that among the entries $A_{i,1}, \dots, A_{i,2\ell}$ there are exactly ℓ 1's, since by construction $A_{i,2j-1} + A_{i,2j} = 1$. Hence, the ℓ -th 1 in row i is at position $A_{i,2\ell-1}$ or $A_{i,2\ell}$, depending only on the sample $b_{\lceil i/2 \rceil, \ell}$ from $X_{\lceil i/2 \rceil}$. For rows $i > 2k$, the ℓ -th 1 is simply at position $A_{i,2\ell}$. Thus, accessing the ℓ -th 1 in any row takes at most one sample, so we can simulate any algorithm on A with random access to the adjacency lists of rows. The situation for columns is essentially symmetric. Similarly, we can allow constant time access to the row and column sums.

In the remainder we show that the constructed matrix A has essentially the same properties as the construction in Theorem 27. We first argue that any 2-approximation u, v for the Boolean

ℓ_0 -Rank-1 problem on A picks all rows $i > 2k$ and picks all even columns and does not pick any odd column. Thus, the only remaining choice is which rows $i \leq 2k$ to pick. To prove this claim, first note that any solution following this pattern has error at most $2kn = \phi n^2$, since the $2k$ undecided rows have $2kn$ entries, and all other entries are correctly recovered by the already chosen parts of uv^T . Hence, we have $\text{OPT} \leq \phi n^2$. Now consider any 2-approximation u, v , which must have cost at most $2\phi n^2$. Note that u picks at least $(1 - 5\phi)n$ of the rows $\{2k + 1, \dots, n\}$, since each such row contains $n/2$ 1's that can only be recovered if we pick the row, so we can afford to ignore at most $8k = 4\phi n$ of these $n - 2k = (1 - \phi)n$ rows. Now, each even column contains at least $(1 - 5\phi)n > n/2$ 1's in picked rows, and thus it is always better to pick the even columns. Similarly, each odd column contains at least $n/2$ 0's in picked rows, and thus it is always better not to pick the odd columns. Hence, we obtain without loss of generality $v_{2j} = 1$ and $v_{2j-1} = 0$. Finally, each row $i > 2k$ contains $n/2$ 1's in columns picked by v and $n/2$ 0's in columns not picked by v , and thus it is always better to pick row i . Hence, we obtain without loss of generality $u_i = 1$ for $i > 2k$.

Our goal now is to lower bound $\|A - uv^T\|_0$ in terms of OPT and the error term $b\phi'n$, similarly to the proof in Theorem 27. Notice that we may ignore the odd columns, as they are not picked by v . Restricted to the even columns, row $2i$ is exactly as row i in the construction in Theorem 27, while row $2i - 1$ is row $2i$ negated. Thus, analogously as in the proof of Theorem 27, we obtain w.h.p. $\text{OPT} \leq (1/2 - 0.99\phi')2kn$ and

$$\|A - uv^T\|_0 \geq \text{OPT} + 2b\phi'n - 0.04k\phi'n \geq (1 + 0.04\phi')\text{OPT},$$

where $b \geq 0.04k$ is the number of “bad” rows $i \leq 2k$. Again analogously, if we compute a $(1 + 0.04\phi') = 1 + C\phi$ -approximation on A , then $b \leq 0.04k$, and thus w.h.p. we correctly identify for at least a 0.96 fraction of the random variables X_i whether $p_i = 1/2 + \phi'$ or $p_i = 1/2 - \phi'$. As before, this yields a lower bound of $\Omega(n/\phi)$ samples. \square

References

- [AGKM16] Sanjeev Arora, Rong Ge, Ravi Kannan, and Ankur Moitra. Computing a nonnegative matrix factorization - provably. *SIAM J. Comput.*, 45(4):1582–1611, 2016.
- [Ale11] Michael Alekhovich. More on average case vs approximation complexity. *Computational Complexity*, 20(4):755–786, 2011.
- [APY09] Noga Alon, Rina Panigrahy, and Sergey Yekhanin. Deterministic approximation algorithms for the nearest codeword problem. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques, 12th International Workshop, APPROX 2009, and 13th International Workshop, RANDOM 2009, Berkeley, CA, USA, August 21-23, 2009. Proceedings*, pages 339–351, 2009.
- [BK02] Piotr Berman and Marek Karpinski. Approximating minimum unsatisfiability of linear equations. In *Proceedings of the Thirteenth Annual ACM-SIAM Symposium on Discrete Algorithms, January 6-8, 2002, San Francisco, CA, USA.*, pages 514–516, 2002.
- [BV10] Radim Belohlávek and Vilém Vychodil. Discovery of optimal factors in binary data via a novel method of matrix decomposition. *J. Comput. Syst. Sci.*, 76(1):3–20, 2010.

- [BY02] Ziv Bar-Yossef. *The complexity of massive data set computations*. PhD thesis, University of California, Berkeley, 2002.
- [CGK⁺17] Flavio Chierichetti, Sreenivas Gollapudi, Ravi Kumar, Silvio Lattanzi, Rina Panigrahy, and David P. Woodruff. Algorithms for ℓ_p low-rank approximation. In *Proceedings of the 34th International Conference on Machine Learning, ICML 2017, Sydney, NSW, Australia, 6-11 August 2017*, pages 806–814, 2017.
- [CLMW11a] Emmanuel J Candès, Xiaodong Li, Yi Ma, and John Wright. Robust principal component analysis? *Journal of the ACM (JACM)*, 58(3):11, 2011.
- [CLMW11b] Emmanuel J. Candès, Xiaodong Li, Yi Ma, and John Wright. Robust principal component analysis? *J. ACM*, 58(3):11:1–11:37, June 2011.
- [CW13] Kenneth L. Clarkson and David P. Woodruff. Low rank approximation and regression in input sparsity time. In *Symposium on Theory of Computing Conference, STOC’13, Palo Alto, CA, USA, June 1-4, 2013*, pages 81–90, 2013.
- [CW15] Kenneth L. Clarkson and David P. Woodruff. Input sparsity and hardness for robust subspace approximation. In *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, pages 310–329, 2015.
- [DAJ⁺15] C. Dan, K. Arnsfelt Hansen, H. Jiang, L. Wang, and Y. Zhou. Low Rank Approximation of Binary Matrices: Column Subset Selection and Generalizations. *ArXiv e-prints*, 2015.
- [DKLR00] Paul Dagum, Richard M. Karp, Michael Luby, and Sheldon M. Ross. An optimal algorithm for monte carlo estimation. *SIAM J. Comput.*, 29(5):1484–1496, 2000.
- [FLM⁺17] Fedor V. Fomin, Daniel Lokshtanov, S. M. Meesum, Saket Saurabh, and Meirav Zehavi. Matrix rigidity: Matrix theory from the viewpoint of parameterized complexity. In *STACS*. Springer, March 2017.
- [GGYT12] Harold W. Gutch, Peter Gruber, Arie Yeredor, and Fabian J. Theis. ICA over finite fields - separability and algorithms. *Signal Processing*, 92(8):1796–1808, 2012.
- [Gri76] D. Grigoriev. Using the notions of separability and independence for proving the lower bounds on the circuit complexity (in russian). *Notes of the Leningrad branch of the Steklov Mathematical Institute, Nauka*, 1976.
- [Gri80] D. Grigoriev. Using the notions of separability and independence for proving the lower bounds on the circuit complexity. *Journal of Soviet Math.*, 14(5):1450–1456, 1980.
- [GV15] Nicolas Gillis and Stephen A. Vavasis. On the complexity of robust PCA and ℓ_1 -norm low-rank matrix approximation. *CoRR*, abs/1509.09236, 2015.
- [JPHY14] Peng Jiang, Jiming Peng, Michael Heath, and Rui Yang. A clustering approach to constrained binary matrix factorization. In *Data Mining and Knowledge Discovery for Big Data*, pages 281–303. Springer, 2014.

- [KG03] Mehmet Koyutürk and Ananth Grama. Proximus: a framework for analyzing very high dimensional discrete-attributed datasets. In *Proceedings of the ninth ACM SIGKDD international conference on Knowledge discovery and data mining*, pages 147–156. ACM, 2003.
- [KGR05] Mehmet Koyutürk, Ananth Grama, and Naren Ramakrishnan. Compression, clustering, and pattern discovery in very high-dimensional discrete-attribute data sets. *IEEE Trans. Knowl. Data Eng.*, 17(4):447–461, 2005.
- [KGR06] Mehmet Koyutürk, Ananth Grama, and Naren Ramakrishnan. Nonorthogonal decomposition of binary matrices for bounded-error data compression and analysis. *ACM Transactions on Mathematical Software (TOMS)*, 32(1):33–69, 2006.
- [KV09] Ravi Kannan and Santosh Vempala. Spectral algorithms. *Foundations and Trends in Theoretical Computer Science*, 4(3-4):157–288, 2009.
- [Li05] Tao Li. A general model for clustering binary data. In *Proceedings of the eleventh ACM SIGKDD international conference on Knowledge discovery in data mining*, pages 188–197. ACM, 2005.
- [Mah11] Michael W. Mahoney. Randomized algorithms for matrices and data. *Foundations and Trends in Machine Learning*, 3(2):123–224, 2011.
- [MGNR06] Edward Meeds, Zoubin Ghahramani, Radford M. Neal, and Sam T. Roweis. Modeling dyadic data with binary latent factors. In *Advances in Neural Information Processing Systems 19, Proceedings of the Twentieth Annual Conference on Neural Information Processing Systems, Vancouver, British Columbia, Canada, December 4-7, 2006*, pages 977–984, 2006.
- [MM13] Xiangrui Meng and Michael W. Mahoney. Low-distortion subspace embeddings in input-sparsity time and applications to robust linear regression. In *Symposium on Theory of Computing Conference, STOC’13, Palo Alto, CA, USA, June 1-4, 2013*, pages 91–100, 2013.
- [MMG⁺08] Pauli Miettinen, Taneli Mielikäinen, Aristides Gionis, Gautam Das, and Heikki Mannila. The discrete basis problem. *IEEE Trans. Knowl. Data Eng.*, 20(10):1348–1362, 2008.
- [MV14] Pauli Miettinen and Jilles Vreeken. MDL4BMF: minimum description length for boolean matrix factorization. *TKDD*, 8(4):18:1–18:31, 2014.
- [NN13] Jelani Nelson and Huy L. Nguyen. OSNAP: faster numerical linear algebra algorithms via sparser subspace embeddings. In *54th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2013, 26-29 October, 2013, Berkeley, CA, USA*, pages 117–126, 2013.
- [PRF15] A. Painsky, S. Rosset, and M. Feder. Generalized Independent Component Analysis Over Finite Alphabets. *ArXiv e-prints*, 2015.
- [RPG15] S. Ravanbakhsh, B. Poczos, and R. Greiner. Boolean Matrix Factorization and Noisy Completion via Message Passing. *ArXiv e-prints*, 2015.

- [RSW16] Ilya P. Razenshteyn, Zhao Song, and David P. Woodruff. Weighted low rank approximations with provable guarantees. In *Proceedings of the 48th Annual ACM SIGACT Symposium on Theory of Computing, STOC 2016, Cambridge, MA, USA, June 18-21, 2016*, pages 250–263, 2016.
- [SBM03] Jouni K. Seppänen, Ella Bingham, and Heikki Mannila. A simple algorithm for topic identification in 0-1 data. In *Knowledge Discovery in Databases: PKDD 2003, 7th European Conference on Principles and Practice of Knowledge Discovery in Databases, Cavtat-Dubrovnik, Croatia, September 22-26, 2003, Proceedings*, pages 423–434, 2003.
- [SH06] Tomás Singliar and Milos Hauskrecht. Noisy-or component analysis and its application to link analysis. *Journal of Machine Learning Research*, 7:2189–2213, 2006.
- [SJY09] Bao-Hong Shen, Shuiwang Ji, and Jieping Ye. Mining discrete patterns via binary matrix factorization. In *Proceedings of the 15th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, Paris, France, June 28 - July 1, 2009*, pages 757–766, 2009.
- [SST06] Arvind Sankar, Daniel A. Spielman, and Shang-Hua Teng. Smoothed analysis of the condition numbers and growth factors of matrices. *SIAM J. Matrix Analysis Applications*, 28(2):446–476, 2006. Available at: <http://www.cs.yale.edu/homes/spielman/Research/nopivotdas.pdf>.
- [SWZ16] Zhao Song, David P. Woodruff, and Peilin Zhong. Low rank approximation with entrywise ℓ_1 -norm error. *CoRR*, abs/1611.00898, 2016.
- [SWZ18] Zhao Song, David P. Woodruff, and Peilin Zhong. Entrywise low rank approximation of general functions, 2018. Manuscript.
- [VAG07] Jaideep Vaidya, Vijayalakshmi Atluri, and Qi Guo. The role mining problem: finding a minimal descriptive set of roles. In *12th ACM Symposium on Access Control Models and Technologies, SACMAT 2007, Sophia Antipolis, France, June 20-22, 2007, Proceedings*, pages 175–184, 2007.
- [Val77] Leslie G. Valiant. Graph-theoretic arguments in low-level complexity. In *Mathematical Foundations of Computer Science 1977, 6th Symposium, Tatranska Lomnica, Czechoslovakia, September 5-9, 1977, Proceedings*, pages 162–176, 1977.
- [Wal74] A. J. Walker. New fast method for generating discrete random numbers with arbitrary frequency distributions. *Electronics Letters*, 10(8):127–128, April 1974.
- [Woo14] David P. Woodruff. Sketching as a tool for numerical linear algebra. *Foundations and Trends in Theoretical Computer Science*, 10(1-2):1–157, 2014.
- [Yer11] Arie Yeredor. Independent component analysis over galois fields of prime order. *IEEE Trans. Information Theory*, 57(8):5342–5359, 2011.
- [ZLD⁺10] Zhong-Yuan Zhang, Tao Li, Chris Ding, Xian-Wen Ren, and Xiang-Sun Zhang. Binary matrix factorization for analyzing gene expression data. *Data Mining and Knowledge Discovery*, 20(1):28–52, 2010.

- [ZLDZ07] Zhongyuan Zhang, Tao Li, Chris Ding, and Xiangsun Zhang. Binary matrix factorization with applications. In *Data Mining, 2007. ICDM 2007. Seventh IEEE International Conference on*, pages 391–400. IEEE, 2007.