

Effective Tensor Sketching via Sparsification*

Dong Xia and Ming Yuan[†]

Columbia University

(March 8, 2024)

Abstract

In this paper, we investigate effective sketching schemes via sparsification for high dimensional multilinear arrays or tensors. More specifically, we propose a novel tensor sparsification algorithm that retains a subset of the entries of a tensor in a judicious way, and prove that it can attain a given level of approximation accuracy in terms of tensor spectral norm with a much smaller sample complexity when compared with existing approaches. In particular, we show that for a k th order $d \times \cdots \times d$ cubic tensor of *stable rank* r_s , the sample size requirement for achieving a relative error ε is, up to a logarithmic factor, of the order $r_s^{1/2} d^{k/2} / \varepsilon$ when ε is relatively large, and $r_s d / \varepsilon^2$ and essentially optimal when ε is sufficiently small. It is especially noteworthy that the sample size requirement for achieving a high accuracy is of an order independent of k . To further demonstrate the utility of our techniques, we also study how higher order singular value decomposition (HOSVD) of large tensors can be efficiently approximated via sparsification.

*This research was supported by NSF Grant DMS-1721584, and NIH Grant 1U54AI117924-01.

[†]Address for Correspondence: Department of Statistics, Columbia University, 1255 Amsterdam Avenue, New York, NY 10027.

1 Introduction

Massive datasets are being generated everyday across diverse fields and can often be formatted into matrices or higher order tensors. For example, in biomedical research, huge data matrices and tensors arise in gene expression analysis (see, e.g., [Kluger et al., 2003](#)), protein-to-protein interaction (see, e.g., [Stelzl et al., 2005](#)), and MRI image analysis (see, e.g., [Smith et al., 2004](#)). They also occur frequently in statistical physics (see, e.g., [Orús, 2014](#); [Cichocki et al., 2015](#)), video processing (see, e.g., [Li and Li, 2010](#); [Liu et al., 2013](#)), and analyzing large graphs and social networks (see, e.g., [Clauset et al., 2004](#); [Abadi et al., 2016](#); [Scott, 2017](#)), to name a few. As the size of these data matrices or tensors grows, it becomes costly and sometimes prohibitively expensive to store, communicate or manipulate them. This naturally brings about the task of “sketching”: approximate the original data matrices or tensors with a more manageable amount of sketches.

In the case of data matrices, numerous sketching approaches have been proposed in recent years. See [Woodruff et al. \(2014\)](#) for a recent review. A popular idea behind many of these approaches is *sparsification* – creating a sparse matrix by zeroing out some entries of the original data matrix. Sparse sketching of a large data matrix not only reduces space complexity but also allows for efficient computations. See, e.g., [Frieze et al. \(2004\)](#); [Arora et al. \(2006\)](#); [Achlioptas and McSherry \(2007\)](#); [Drineas and Zouzias \(2011\)](#); [Achlioptas et al. \(2013\)](#); [Krishnamurthy and Singh \(2013\)](#), among others. The main purpose of this article is to investigate to what extent sparsification can be used to effectively sketch higher order tensors. There have been some recent attempts along this direction. In particular, our work is inspired by [Nguyen et al. \(2015\)](#) who showed that for a k th order cubic tensor $\mathbf{A} \in \mathbb{R}^{d \times \dots \times d}$, there is a randomized sparsification scheme that yields another tensor $\tilde{\mathbf{A}}$ of same dimension but with

$$\text{nnz}(\tilde{\mathbf{A}}) = \tilde{O}_p\left(\frac{d^{k/2}\text{sr}(\mathbf{A})}{\varepsilon^2}\right), \quad \text{as } d \rightarrow \infty, \quad (1)$$

such that

$$\|\tilde{\mathbf{A}} - \mathbf{A}\| \leq \varepsilon \|\mathbf{A}\|.$$

Here, $\text{nnz}(\cdot)$ stands for the number of nonzero entries of a tensor, $\text{sr}(\mathbf{A}) = \|\mathbf{A}\|_{\text{F}}^2 / \|\mathbf{A}\|^2$ is

the so-called stable rank (see, e.g., [Achlioptas et al., 2013](#); [Nguyen et al., 2015](#)) of a tensor \mathbf{A} , $\|\cdot\|$ is the usual tensor spectral norm, and $\tilde{O}(\cdot)$ means $O(\cdot)$, up to a certain polynomial of logarithmic factor. Similar results have also been obtained by [Bhojanapalli and Sanghavi \(2015\)](#) in the case when $k = 3$. On the one hand, the sample size requirement given by (1) is satisfying because it is essentially optimal in the matrix case, that is $k = 2$. See, e.g., [Achlioptas et al. \(2013\)](#). On the other hand, the exponential dependence on k suggests a large amount of entries still need to be retained to yield a good approximation. Our goal is to investigate if this aspect could be improved.

In particular, we propose a novel tensor sparsification algorithm that randomly retain entries from \mathbf{A} in a judicious way to yield a tensor $\widehat{\mathbf{A}}^{\text{SPA}}$ such that

$$\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\| \leq \varepsilon \|\mathbf{A}\|,$$

and

$$\text{nnz}(\widehat{\mathbf{A}}^{\text{SPA}}) = \tilde{O}_p \left(\max \left\{ \frac{d \cdot \text{sr}(\mathbf{A})}{\varepsilon^2}, \frac{d^{k/2} \cdot \text{sr}(\mathbf{A})^{1/2}}{\varepsilon} \right\} \right). \quad (2)$$

Here, to fix ideas, we focus on the case of cubic tensors although our results deal with more general rectangular tensors as well. This sample size requirement significantly improves those earlier ones. Especially if a high accuracy approximation is sought, that is $\varepsilon \leq \text{sr}(\mathbf{A}) \cdot d^{-k/2+1}$, then our sparsification algorithm can achieve relative approximation error ε in terms of tensor spectral norm by retaining as few as $\tilde{O}_p(d \cdot \text{sr}(\mathbf{A}) \cdot \varepsilon^{-2})$ entries of \mathbf{A} , regardless of the order of the tensor. Furthermore, for larger ε , the number of nonzero entries we keep is smaller than $\tilde{\mathbf{A}}$ by a factor of $\text{sr}(\mathbf{A})^{1/2} \varepsilon^{-1}$.

Similar to many other sparsification algorithms, we treat different entries according to their magnitude: large entries are always kept, and moderate ones are sampled proportion to their square values. The key difference between our approach and the existing ones is in the treatment of small entries. Instead of zeroing them out as, for example, [Nguyen et al. \(2015\)](#), we sample them in a uniform fashion, which proves to be essential for obtaining good approximation with tighter number of nonzero entries. This modification is motivated by the concentration behavior of randomly sampled tensors recently observed by [Yuan and Zhang \(2016, 2017\)](#); [Xia and Yuan \(2017\)](#).

To demonstrate the effectiveness of our tensor sketching schemes, we show how they can be used for efficient approximation of the leading singular spaces from higher order singular value decomposition (HOSVD). Let $\mathbf{U}_j \in \mathbb{R}^{d \times r}$ be the top r left singular vectors of the flattening of \mathbf{A} along its j th mode. We show that it is possible to construct an approximation $\widehat{\mathbf{U}}_j$ obeying

$$\|\widehat{\mathbf{U}}_j \widehat{\mathbf{U}}_j^\top - \mathbf{U}_j \mathbf{U}_j^\top\| \leq \varepsilon,$$

if we retain

$$\tilde{O}_p \left(\max \left\{ \frac{rd}{\varepsilon^2}, \frac{rd^{k/2}}{\varepsilon} \right\} \right)$$

carefully chosen entries. As before, we note that for high accuracy approximations, the sample complexity is essentially independent of the order of the tensor. Although our primary focus is on higher order tensors, as a byproduct, our results indicate that our sparsification scheme improves the sample complexity of earlier approaches for approximating the singular vectors of highly rectangular matrix.

The rest of the paper is organized as follows. We first discuss the new tensor sparsification algorithm in Section 2. In Section 3 we consider the application to HOSVD. All proofs are relegated to Section 4.

2 Tensor Sparsification

Sketches of a tensor $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ are its approximations. We consider measuring the quality in terms of relative error in terms of tensor spectral norm. Recall that the spectral norm of a tensor $\mathbf{B} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ is defined as

$$\|\mathbf{B}\| = \sup_{\mathbf{u}_j \in \mathbb{R}^{d_j}, \|\mathbf{u}_j\|_{\ell_2} \leq 1} \langle \mathbf{B}, \mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k \rangle.$$

We seek an approximation $\widehat{\mathbf{A}}$ of \mathbf{A} such that

$$\|\widehat{\mathbf{A}} - \mathbf{A}\| \leq \varepsilon \|\mathbf{A}\|,$$

for some $\varepsilon \in (0, 1)$.

We first consider sketching a tensor by sparsification. The idea is to systematically zero out entries of \mathbf{A} and scale the remaining entries to yield a good approximation of \mathbf{A} . We focus here on sparsification strategies that are carried out in an entry-by-entry fashion. Our approach can be characterized as *keeping* large entries, *sampling proportionally* moderate entries, and *sampling uniformly* small entries. The key is determining how to classify entries into these categories, and how to sample the moderate entries, so that the number of nonzero entries retained are as small as possible. Details are presented in Algorithm 1.

Algorithm 1 Tensor Sparsification

Input: $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$, sampling budget $1 \leq n \leq d_1 \cdots d_k$.

2: Output: $\widehat{\mathbf{A}}^{\text{SPA}} \in \mathbb{R}^{d_1 \times \dots \times d_k}$.

for $i_1 \in [d_1], i_2 \in [d_2], \dots, i_k \in [d_k]$ **do**

4: **if** $|A(i_1, \dots, i_k)| \geq \|\mathbf{A}\|_{\text{F}}/n^{1/2}$, **then** } Large Entries

$\widehat{A}(i_1, \dots, i_k) = A(i_1, \dots, i_k)$.

6: **end if**

if $|A(i_1, \dots, i_k)|/\|\mathbf{A}\|_{\text{F}} \in \left(\frac{1}{(d_1 \cdots d_k)^{1/2}}, \frac{1}{n^{1/2}}\right)$, **then** } Moderate Entries

$$\widehat{A}(i_1, \dots, i_k) = \begin{cases} \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}, & \text{with probability } P(i_1, \dots, i_k) := \frac{nA^2(i_1, \dots, i_k)}{\|\mathbf{A}\|_{\text{F}}^2} \\ 0, & \text{with probability } 1 - P(i_1, \dots, i_k). \end{cases}$$

8: **end if** }

if $|A(i_1, \dots, i_k)| \leq \|\mathbf{A}\|_{\text{F}}/(d_1 \cdots d_k)^{1/2}$, **then** } Small Entries

10:
$$\widehat{A}(i_1, \dots, i_k) = \begin{cases} \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}, & \text{with probability } P(i_1, \dots, i_k) := \frac{n}{d_1 d_2 \cdots d_k} \\ 0, & \text{with probability } 1 - P(i_1, \dots, i_k) \end{cases}$$

end if

12: **end for**

 Output: $\widehat{\mathbf{A}}^{\text{SPA}} = \widehat{\mathbf{A}}$.

In particular, we keep all entries whose absolute value is greater than $n^{-1/2}\|\mathbf{A}\|_{\text{F}}$, sample

uniformly all entries whose absolute value is smaller than $(d_1 \cdots d_k)^{-1/2} \|\mathbf{A}\|_F$, and sample proportional to their squared values entries whose absolute value is in-between. Here n is a sampling parameter. Note that $\mathbb{E}[\text{nnz}(\widehat{\mathbf{A}}^{\text{SPA}})] \leq 2n$. And it is not hard to see, by Chernoff bound, that $\text{nnz}(\widehat{\mathbf{A}}^{\text{SPA}}) = O_p(n)$. In other words, n represents essentially the targeted sampling budget.

We note that our sparsification algorithm is similar to the one proposed earlier by [Nguyen et al. \(2015\)](#). But the two schemes also have several key differences. The main difference between the two algorithms is their treatment of “small” entries. [Nguyen et al. \(2015\)](#) suggests to zero them out, while ours sample them in a uniform fashion. This is largely motivated by the concentration behavior of randomly sampled tensors observed earlier. In particular, it can be shown that a uniformly sampled tensor concentrates much sharply around its mean if its entries are sufficiently small (see, e.g., [Yuan and Zhang, 2016](#)). Therefore, instead of discarding small entries, we could derive a good estimate of them by sampling uniformly. Another subtle difference between the two algorithm is in the criteria for “small” entries. Our criterion for “small” entries is that their absolute values are smaller than $(d_1 \cdots d_k)^{-1/2} \|\mathbf{A}\|_F$, whereas [Nguyen et al. \(2015\)](#) treats only cubic tensors, that is $d_1 = d_2 = \cdots = d_k =: d$, and small entries of their scheme are those smaller than $n^{-1/2} d^{-k/4} \|\mathbf{A}\|_F \log^{k/2} d$.

We now present the performance bounds for our sparsification algorithm.

Theorem 1. *Let $\mathbf{A} \in \mathbb{R}^{d_1 \times \cdots \times d_k}$ and $\widehat{\mathbf{A}}^{\text{SPA}}$ be the output from Algorithm 1 with sampling budget n . There exists a constant $C > 0$ depending on k only such that if for any $\alpha \geq 4 \log(k \log d_{\max})$ and $\varepsilon \in (0, 1)$, if*

$$n \geq C \max \left\{ \alpha^4 \frac{d_{\max} \cdot \text{sr}(\mathbf{A})}{\varepsilon^2} \log^2 d_{\max}, \alpha^2 \frac{(d_1 \cdots d_k \cdot \text{sr}(\mathbf{A}))^{1/2}}{\varepsilon} \log^{k+4} d_{\max} \right\},$$

then, with probability at least $1 - d_{\max}^{-\alpha}$,

$$\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\| \leq \varepsilon \|\mathbf{A}\|,$$

where $d_{\max} = \max\{d_1, \dots, d_k\}$.

In the light of Theorem 1, we can achieve relative error ε in terms of tensor spectral

norm with a sparse tensor such that

$$\text{nnz}(\widehat{\mathbf{A}}^{\text{SPA}}) = \begin{cases} \tilde{O}\left(\varepsilon^{-2} \cdot d_{\max} \cdot \text{sr}(\mathbf{A})\right), & \text{if } \varepsilon \leq d_{\max} \cdot \text{sr}(\mathbf{A})^{1/2} \cdot (d_1 \dots d_k)^{-1/2}; \\ \tilde{O}\left(\varepsilon^{-1} \cdot (d_1 \dots d_k \cdot \text{sr}(\mathbf{A}))^{1/2}\right), & \text{otherwise.} \end{cases}$$

This significantly improves earlier work by [Bhojanapalli and Sanghavi \(2015\)](#) and [Nguyen et al. \(2015\)](#). It is worth noting that for small ε , or high accuracy approximation, the number of nonzero entries of $\widehat{\mathbf{A}}^{\text{SPA}}$ is of the order $\varepsilon^{-2} d_{\max} \cdot \text{sr}(\mathbf{A})$, regardless of k . This, in particular, is known to be optimal in the matrix ($k = 2$) case (see, e.g., [Achlioptas et al., 2013](#)).

The main technical tool for proving Theorem 1 is the following concentration inequality for random tensors which might be of independent interest.

Theorem 2. *Let $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ and $\mathbf{P} \in [0, 1]^{d_1 \times \dots \times d_k}$ be two fixed tensors, $\Delta \in \{0, 1\}^{d_1 \times \dots \times d_k}$ be a random tensor such that $\mathbb{E}\Delta(i_1, \dots, i_k) = P(i_1, \dots, i_k)$. Define a random tensor $\widehat{\mathbf{A}} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ by*

$$\widehat{A}(i_1, \dots, i_k) = A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k) / P(i_1, \dots, i_k).$$

Then, there exist absolute constants $C_1, C_2, C_3 > 0$ such that for any $\alpha > 0$, with probability at least $1 - 3d_{\max}^{-\alpha}$,

$$\|\widehat{\mathbf{A}} - \mathbf{A}\| \leq C_1 \left(\left(\sum_{j=1}^k d_j \right)^{1/2} + \alpha k \log d_{\max} \right) \alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) + C_2 \alpha k^3 \log^{k+2}(d_{\max}) \sqrt{\nu} \alpha_{\infty}(\mathbf{A}, \mathbf{P}),$$

where

$$\nu = C_3 \alpha \max \left\{ \beta(\mathbf{P}), k \log d_{\max} \right\},$$

$$\beta(\mathbf{P}) = \max_{j=1, \dots, k} \max_{i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_k} \sum_{i_j=1}^{d_j} P(i_1, \dots, i_k),$$

$$\alpha_{\infty}(\mathbf{A}, \mathbf{P}) = \max_{i_j \in [d_j], j=1, \dots, k} \frac{|A(i_1, \dots, i_k)|}{P(i_1, \dots, i_k)},$$

and

$$\alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) = \max_{i_j \in [d_j], j=1, \dots, k} \left(\frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right)^{1/2}.$$

Here we follow the convention that $0/0 = 0$.

3 HOSVD via Tensor Sketching

To further illustrate the merits of the sketching schemes introduced earlier, we now consider a specific application to HOSVD, a popular technique for analyzing high dimensional tensor data. See, e.g., [Kolda and Bader \(2009\)](#); [Sidiropoulos et al. \(2017\)](#) and references therein.

For a k -th order tensor $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$, let $\mathbf{M}_j = \mathcal{M}_j(\mathbf{A}) \in \mathbb{R}^{d_j \times d_{-j}}$ be its j -th matricization where $1 \leq j \leq k$, that is,

$$\mathcal{M}_j(\mathbf{A}) \left(i_j, \sum_{s=1, s \neq j}^k (i_s - 1) \left(\prod_{s'=s+1, s' \neq j}^k d_{s'} \right) + 1 \right) = A(i_1, \dots, i_k), \quad \forall i_j \in [d_j], 1 \leq j \leq k.$$

Here $d_{-j} = (d_1 \cdots d_k)/d_j$. Denote by $\mathbf{U}_j^{(r_j)}$ the collection of the top r_j left singular vectors of \mathbf{M}_j . Clearly, $\mathbf{U}_j^{(r_j)}$ is computable via the standard matrix singular value decomposition on \mathbf{M}_j whose computation complexity is $O(d_j d_1 d_2 \dots d_k)$, see [Golub and Van Loan \(2012\)](#). Efficient computation of singular value decomposition for large matrices is an actively researched topic in numerical algebra and computational science. See [Berry \(1992\)](#); [Kobayashi et al. \(2001\)](#); [Achlioptas and McSherry \(2007\)](#); [Holmes et al. \(2007\)](#); [Drineas and Zouzias \(2011\)](#); [Menon and Elkan \(2011\)](#), among numerous others.

A general idea is to first obtain an approximation of \mathbf{M}_j , say $\widehat{\mathbf{M}}_j \in \mathbb{R}^{d_j \times d_{-j}}$, that is amenable for fast computation of singular value decomposition; and then approximate $\mathbf{U}_j^{(r_j)}$ by the top left singular vectors of $\widehat{\mathbf{M}}_j$. In particular, sparsification is commonly used to yield $\widehat{\mathbf{M}}_j$. Denote by $\Delta_j = \widehat{\mathbf{M}}_j - \mathbf{M}_j$ and by $\widehat{\mathbf{U}}_j^{(r_j)}$ the leading r_j left singular vectors of $\widehat{\mathbf{M}}_j$. By Davis-Kahan Theorem ([Davis and Kahan, 1970](#)), we get

$$\left\| \widehat{\mathbf{U}}_j^{(r_j)} \left(\widehat{\mathbf{U}}_j^{(r_j)} \right)^\top - \mathbf{U}_j^{(r_j)} \left(\mathbf{U}_j^{(r_j)} \right)^\top \right\| \leq \frac{2\|\Delta_j\|}{\bar{g}_{r_j}(\mathbf{M}_j)} \quad (3)$$

where $\sigma_k(\cdot)$ denotes the k -th singular value, and

$$\bar{g}_{r_j}(\mathbf{M}_j) = \sigma_{r_j}(\mathbf{M}_j) - \sigma_{r_j+1}(\mathbf{M}_j),$$

is the r_j -th eigengap. In particular, we can consider applying this strategy by taking $\widehat{\mathbf{M}}_j = \mathcal{M}_j(\widehat{\mathbf{A}}^{\text{SPA}})$. The following result characterizes its performance.

Theorem 3. Let $\mathbf{U}_j^{(r_j)}$ and $\widehat{\mathbf{U}}_j^{(r_j)}$ be the top r_j left singular vectors of $\mathcal{M}_j(\mathbf{A})$ and $\mathcal{M}_j(\widehat{\mathbf{A}}^{\text{SPA}})$ respectively. Then there exists a constant $C > 0$ depending on k only such that for any $t > 0$,

$$\begin{aligned} & \left\| \widehat{\mathbf{U}}_j^{(r_j)} \left(\widehat{\mathbf{U}}_j^{(r_j)} \right)^\top - \mathbf{U}_j^{(r_j)} \left(\mathbf{U}_j^{(r_j)} \right)^\top \right\| \\ & \leq C \frac{\|\mathbf{M}_j\|_F}{\bar{g}_{r_j}(\mathbf{M}_j)} \left(\sqrt{\frac{d_1 \dots d_k (t + \log d_{\max})}{n d_j}} + \frac{(d_1 \dots d_k)^{1/2} (t + \log d_{\max})}{n} \right), \end{aligned}$$

with probability at least $1 - e^{-t}$.

By Theorem 3, in the case when $\frac{\|\mathbf{M}_j\|_F}{\bar{g}_{r_j}(\mathbf{M}_j)} = O(\sqrt{r_j})$, we can ensure

$$\left\| \widehat{\mathbf{U}}_j^{(r_j)} \left(\widehat{\mathbf{U}}_j^{(r_j)} \right)^\top - \mathbf{U}_j^{(r_j)} \left(\mathbf{U}_j^{(r_j)} \right)^\top \right\| \leq \varepsilon$$

by taking

$$n \geq C \cdot \max \left\{ \frac{r_j d_1 \dots d_k}{d_j \varepsilon^2}, \frac{(r_j d_1 \dots d_k)^{1/2}}{\varepsilon} \right\} \log d_{\max}. \quad (4)$$

A critical fact that is neglected by this approach is that we are interested in approximating the left singular vectors of a potentially very “fat” matrix because d_{-j} is generally much larger than d_j . As such, this type of approach turns out to be suboptimal for our purpose.

Alternatively, we adopt a new spectral method similar in spirit to a recent proposal from [Xia and Yuan \(2017\)](#). More specifically, we shall approximate $\mathbf{U}_j^{(r_j)}$ by the leading eigenvectors of an approximation of $\mathbf{M}_j \mathbf{M}_j^\top$ instead. In particular, we can run Algorithm 1 twice to obtain two *independent* sparsifications of \mathbf{A} , denoted by $\widehat{\mathbf{A}}_1^{\text{SPA}}$ and $\widehat{\mathbf{A}}_2^{\text{SPA}}$, and then proceed to approximate $\mathbf{M}_j \mathbf{M}_j^\top$ by $\mathcal{M}_j(\widehat{\mathbf{A}}_1^{\text{SPA}}) \mathcal{M}_j(\widehat{\mathbf{A}}_2^{\text{SPA}})^\top$. Details are presented in Algorithm 2.

Algorithm 2 Computing HOSVD via Tensor Sparsification

- Input: $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$, sampling budget $n \geq 1$.
- 2: Output: the r_j leading left singular vectors $\widehat{\mathbf{U}}_j^{(r_j)}$ as an estimate of HOSVD of $\mathcal{M}_j(\mathbf{A})$.
Run Algorithm 1 on \mathbf{A} with sampling budget n . Denote the output by $\widehat{\mathbf{A}}_1^{\text{SPA}}$.
 - 4: Run Algorithm 1 on \mathbf{A} with sampling budget n . Denote the output by $\widehat{\mathbf{A}}_2^{\text{SPA}}$.
Compute $\widehat{\mathbf{U}}_j^{(r_j)}$ as the r_j leading left singular vectors of $\mathcal{M}_j(\widehat{\mathbf{A}}_1^{\text{SPA}}) \mathcal{M}_j(\widehat{\mathbf{A}}_2^{\text{SPA}})^\top$.
 - 6: Output $\widehat{\mathbf{U}}_j^{(r_j)}$.
-

The following theorem provides the performance bound for approximate the singular space $\mathbf{U}_j^{(r_j)}$ s.

Theorem 4. Denote by $\mathbf{U}_j^{(r_j)}$ the r_j leading left singular vectors of $\mathcal{M}_j(\mathbf{A})$. Let $\widehat{\mathbf{U}}_j^{(r_j)}$ be the output from Algorithm 2. There exists a constant $C > 0$ such that for any $\alpha \geq 1$ and $\varepsilon \in (0, 1)$, if

$$n \geq C\alpha \left(\frac{d_j \log d_{\max}}{\varepsilon^2} \frac{\|\mathbf{A}\|_{\text{F}}^2 \sigma_{\max}^2(\mathbf{M}_j)}{\bar{g}_{r_j}^2(\mathbf{M}_j \mathbf{M}_j^\top)} + \frac{(d_1 \dots d_k)^{1/2} \log d_{\max}}{\varepsilon} \frac{\|\mathbf{A}\|_{\text{F}}^2}{\bar{g}_{r_j}(\mathbf{M}_j \mathbf{M}_j^\top)} \right),$$

then

$$\left\| \widehat{\mathbf{U}}_j^{(r_j)} \left(\widehat{\mathbf{U}}_j^{(r_j)} \right)^\top - \mathbf{U}_j^{(r_j)} \left(\mathbf{U}_j^{(r_j)} \right)^\top \right\| \leq \varepsilon,$$

with probability at least $1 - d_{\max}^{-\alpha}$.

From Theorem 4, if $\frac{\|\mathbf{A}\|_{\text{F}}^2}{\bar{g}_{r_j}(\mathbf{M}_j \mathbf{M}_j^\top)} = O(r_j)$ and $\frac{\|\mathbf{A}\|_{\text{F}}^2 \sigma_{\max}^2(\mathbf{M}_j)}{\bar{g}_{r_j}^2(\mathbf{M}_j \mathbf{M}_j^\top)} = O(r_j)$, then the required sample complexity for sparsification is

$$\tilde{O}_p \left(\frac{r_j d_j \log d_{\max}}{\varepsilon^2} + \frac{r_j (d_1 \dots d_k)^{1/2} \log d_{\max}}{\varepsilon} \right).$$

It is worth noting that, even though our main focus is on higher order tensors, in the case of matrices ($k = 2$) this sample complexity compares favorable with other sparsification techniques that have been developed for computing singular vectors. For example, consider computing the top r left singular vectors of a $d_1 \times d_2$ ($d_1 \leq d_2$) matrix. The approach from Achlioptas and McSherry (2007) needs to sample

$$\tilde{O}_p \left(\frac{r d_1 d_2^2}{\varepsilon^2} \cdot \frac{\max_{i,j} |A(i, j)|^2}{\|\mathbf{A}\|_{\text{F}}^2} \right)$$

entries; the technique of Drineas et al. (2006) requires

$$\tilde{O}_p \left(\frac{r d_2}{\varepsilon^2} \right)$$

entries. These are to be compared with Algorithm 2 which needs

$$\tilde{O}_p \left(\frac{r d_1}{\varepsilon^2} + \frac{r (d_1 d_2)^{1/2}}{\varepsilon} \right)$$

sampled entries, which could be much smaller than the previous two when $d_1 \ll d_2$.

4 Proofs

We now present the proofs to our main results.

4.1 Proof of Theorem 1

Theorem 1 follows immediately from the concentration bound for $\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\|$ below.

Lemma 1. *Let $\mathbf{A} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ and $\widehat{\mathbf{A}}^{\text{SPA}}$ be the output from Algorithm 1 with sampling budget n . Then there exist absolute constants $C_1, C_2 > 0$ such that, for any $\alpha \geq 4 \log(k \log d_{\max})$, the following bound holds with probability at least $1 - d_{\max}^{-\alpha}$:*

$$\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\| \leq C_1 \alpha^2 k^4 \log(d_{\max}) \sqrt{\frac{\|\mathbf{A}\|_{\text{F}}^2 d_{\max}}{n}} + C_2 \alpha^2 k^5 \log^{k+4}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_{\text{F}}}{n}.$$

Proof of Lemma 1. Given \mathbf{A} , we define the disjoint subsets of $[d_1] \times \dots \times [d_k]$

$$\Omega_1 = \{(i_1, \dots, i_k) : |A(i_1, \dots, i_k)| \leq \|\mathbf{A}\|_{\text{F}} / (d_1 \dots d_k)^{1/2}\},$$

$$\Omega_2 = \left\{ (i_1, \dots, i_k) : |A(i_1, \dots, i_k)| / \|\mathbf{A}\|_{\text{F}} \in \left(\frac{1}{(d_1 \dots d_k)^{1/2}}, \frac{1}{n^{1/2}} \right) \right\},$$

and

$$\Omega_3 = \{(i_1, \dots, i_k) : |A(i_1, \dots, i_k)| \geq \|\mathbf{A}\|_{\text{F}} / n^{1/2}\}.$$

Note that $\Omega_1, \Omega_2, \Omega_3$ are non-random subsets for given \mathbf{A} . Then,

$$\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\| \leq \|\widehat{\mathbf{A}}_{\Omega_1}^{\text{SPA}} - \mathbf{A}_{\Omega_1}\| + \|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\| + \|\widehat{\mathbf{A}}_{\Omega_3}^{\text{SPA}} - \mathbf{A}_{\Omega_3}\|.$$

By definition of $\widehat{\mathbf{A}}^{\text{SPA}}$ in Algorithm 1, we have $\|\widehat{\mathbf{A}}_{\Omega_3}^{\text{SPA}} - \mathbf{A}_{\Omega_3}\| = 0$ so that it suffices to bound $\|\widehat{\mathbf{A}}_{\Omega_1}^{\text{SPA}} - \mathbf{A}_{\Omega_1}\|$ and $\|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\|$.

Upper bound of $\|\widehat{\mathbf{A}}_{\Omega_1}^{\text{SPA}} - \mathbf{A}_{\Omega_1}\|$. In order to apply Theorem 2, we introduce auxiliary tensors \mathbf{B} and $\tilde{\mathbf{B}}$ such that $\mathbf{B}_{\Omega_1} = \mathbf{A}_{\Omega_1}$ and $\mathbf{B}_{\Omega_1^\dagger} = \mathbf{0}$, where Ω_1^\dagger denotes the complement of Ω_1 . Define a tensor $\mathbf{P} \in [0, 1]^{d_1 \times \dots \times d_k}$ such that

$$P(i_1, \dots, i_k) = \begin{cases} \frac{n}{d_1 \dots d_k}, & \text{if } (i_1, \dots, i_k) \in \Omega_1 \\ 0, & \text{otherwise.} \end{cases}$$

Then, random tensor $\tilde{\mathbf{B}}$ is defined as

$$\tilde{B}(i_1, \dots, i_k) = \begin{cases} \frac{B(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}, & \text{with probability } P(i_1, \dots, i_k) \\ 0, & \text{with probability } 1 - P(i_1, \dots, i_k), \end{cases}$$

where we followed the convention $0/0 = 0$. Clearly, $\tilde{\mathbf{B}} - \mathbf{B}$ has the same distribution as $\widehat{\mathbf{A}}_{\Omega_1}^{\text{SPA}} - \mathbf{A}_{\Omega_1}$. To apply Theorem 2, we observe that

$$\nu = C_3 t \max \left\{ \frac{nd_{\max}}{d_1 \dots d_k}, k \log d_{\max} \right\}$$

and

$$\alpha_{\infty}(\mathbf{B}, \mathbf{P}) = \max_{i_1, \dots, i_k} \frac{|B(i_1, \dots, i_k)|}{P(i_1, \dots, i_k)} = \max_{i_1, \dots, i_k} \frac{d_1 \dots d_k}{n} |B(i_1, \dots, i_k)| \leq \frac{(d_1 \dots d_k)^{1/2}}{n} \|\mathbf{A}\|_{\text{F}}$$

and

$$\alpha_{2,\infty}(\mathbf{B}, \mathbf{P}) = \max_{i_1, \dots, i_k} \frac{|B(i_1, \dots, i_k)|}{\sqrt{P(i_1, \dots, i_k)}} = \max_{i_1, \dots, i_k} \frac{(d_1 \dots d_k)^{1/2} |B(i_1, \dots, i_k)|}{n^{1/2}} \leq \frac{\|\mathbf{A}\|_{\text{F}}}{n^{1/2}}.$$

By Theorem 2, with probability at least $1 - d_{\max}^{-t}$,

$$\|\widehat{\mathbf{A}}_{\Omega_1}^{\text{SPA}} - \mathbf{A}_{\Omega_1}\| = \|\tilde{\mathbf{B}} - \mathbf{B}\| \leq C_1 t k^3 \sqrt{\frac{d_{\max}}{n}} \|\mathbf{A}\|_{\text{F}} + C_2 t k^4 \log^{k+3}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_{\text{F}}}{n}.$$

Upper bound of $\|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\|$. Bounding $\|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\|$ is more involved. For $s = 1, 2, \dots, \lceil \log(d_1 \dots d_k/n) \rceil$, define

$$\Omega_{2,s} = \left\{ (i_1, \dots, i_k) : |A(i_1, \dots, i_k)|^2 \in \left[\frac{\|\mathbf{A}\|_{\text{F}}^2}{n} 2^{-s}, \frac{\|\mathbf{A}\|_{\text{F}}^2}{n} 2^{-s+1} \right) \right\}.$$

Clearly,

$$\Omega_2 = \bigcup_{s=1}^{\lceil \log(d_1 \dots d_k/n) \rceil} \Omega_{2,s},$$

so that

$$\|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\| \leq \sum_{s=1}^{\lceil \log(d_1 \dots d_k/n) \rceil} \|\widehat{\mathbf{A}}_{\Omega_{2,s}}^{\text{SPA}} - \mathbf{A}_{\Omega_{2,s}}\|.$$

We now apply Theorem 2 to bound each term on the righthand side. We follow the same strategy as before and define auxiliary tensors $\tilde{\mathbf{B}}_s$ and \mathbf{B}_s such that $(\mathbf{B}_s)_{\Omega_{2,s}} = \mathbf{A}_{\Omega_{2,s}}$ and

$(\mathbf{B}_s)_{\Omega_{2,s}^\dagger} = \mathbf{0}$. The probability tensor \mathbf{P}_s is defined as

$$P_s(i_1, \dots, i_k) = \begin{cases} \frac{nA^2(i_1, \dots, i_k)}{\|\mathbf{A}\|_F^2}, & \text{if } (i_1, \dots, i_k) \in \Omega_{2,s} \\ 0, & \text{otherwise.} \end{cases}$$

The random tensor $\tilde{\mathbf{B}}_s$ is defined as

$$\tilde{B}_s(i_1, \dots, i_k) = \begin{cases} \frac{B_s(i_1, \dots, i_k)}{P_s(i_1, \dots, i_k)}, & \text{with probability } P_s(i_1, \dots, i_k) \\ 0, & \text{with probability } 1 - P_s(i_1, \dots, i_k). \end{cases}$$

Clearly, $\widehat{\mathbf{A}}_{\Omega_{2,s}}^{\text{SPA}} - \mathbf{A}_{\Omega_{2,s}}$ has the same distribution as $\tilde{\mathbf{B}}_s - \mathbf{B}_s$. To apply Theorem 2, observe that

$$\alpha_{2,\infty}(\mathbf{B}_s, \mathbf{P}_s) = \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \sqrt{\frac{B_s^2(i_1, \dots, i_k)}{P_s(i_1, \dots, i_k)}} = \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \sqrt{\frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}} = \sqrt{\frac{\|\mathbf{A}\|_F^2}{n}}.$$

Since

$$\nu = C_1 t \max \left\{ \max_{j \in [k]} \max_{i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_k} \sum_{i_j: (i_1, \dots, i_k) \in \Omega_{2,s}} P(i_1, \dots, i_k), k \log d_{\max} \right\},$$

we obtain

$$\begin{aligned} \sqrt{\nu} \alpha_{\infty}(\mathbf{B}_s, \mathbf{P}_s) &\leq C_1 t^{1/2} k^{1/2} \log^{1/2}(d_{\max}) \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \frac{|A(i_1, \dots, i_k)|}{P(i_1, \dots, i_k)} \\ &+ C_1 t^{1/2} \left(\max_{j \in [k]} \max_{i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_k} \sqrt{\sum_{i_j: (i_1, \dots, i_k) \in \Omega_{2,s}} P(i_1, \dots, i_k)} \right) \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \frac{|A(i_1, \dots, i_k)|}{P(i_1, \dots, i_k)}. \end{aligned}$$

By definition of $\Omega_{2,s}$, we have

$$\frac{\max_{(i_1, \dots, i_k) \in \Omega_{2,s}} P(i_1, \dots, i_k)}{\min_{(i_1, \dots, i_k) \in \Omega_{2,s}} P(i_1, \dots, i_k)} \leq 2.$$

Therefore,

$$\begin{aligned} &\left(\max_{j \in [k]} \max_{i_1, \dots, i_{j-1}, i_{j+1}, \dots, i_k} \sqrt{\sum_{i_j: (i_1, \dots, i_k) \in \Omega_{2,s}} P(i_1, \dots, i_k)} \right) \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \frac{|A(i_1, \dots, i_k)|}{P(i_1, \dots, i_k)} \\ &\leq \sqrt{2d_{\max}} \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \frac{|A(i_1, \dots, i_k)|}{\sqrt{P(i_1, \dots, i_k)}}. \end{aligned}$$

By the fact $P(i_1, \dots, i_k) = \frac{nA^2(i_1, \dots, i_k)}{\|\mathbf{A}\|_F^2}$ and $|A(i_1, \dots, i_k)| \geq \|\mathbf{A}\|_F / (d_1 \dots d_k)^{1/2}$, we get

$$\begin{aligned} \sqrt{\nu} \alpha_{\infty}(\mathbf{B}_s, \mathbf{P}_s) &\leq C_1 k^{1/2} t^{1/2} \log^{1/2}(d_{\max}) \max_{(i_1, \dots, i_k) \in \Omega_{2,s}} \frac{\|\mathbf{A}\|_F^2}{n |A(i_1, \dots, i_k)|} + C_2 t^{1/2} d_{\max}^{1/2} \sqrt{\frac{\|\mathbf{A}\|_F^2}{n}} \\ &\leq C_1 k^{1/2} t^{1/2} \log^{1/2}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_F}{n} + C_2 t^{1/2} \sqrt{\frac{\|\mathbf{A}\|_F^2 d_{\max}}{n}}. \end{aligned}$$

By Theorem 2, with probability at least $1 - d_{\max}^{-t}$,

$$\|\widehat{\mathbf{A}}_{\Omega_{2,s}}^{\text{SPA}} - \mathbf{A}_{\Omega_{2,s}}\| \leq C_1 t^2 k^3 \sqrt{\frac{\|\mathbf{A}\|_{\text{F}}^2 d_{\max}}{n}} + C_2 t^2 k^4 \log^{k+3}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_{\text{F}}}{n}.$$

By taking a uniform bound for all $s = 1, 2, \dots, \lceil \log(d_1 \dots d_k / n) \rceil$, we conclude that with probability at least $1 - k \log(d_{\max}) d_{\max}^{-t}$,

$$\|\widehat{\mathbf{A}}_{\Omega_2}^{\text{SPA}} - \mathbf{A}_{\Omega_2}\| \leq C_1 t^2 k^4 \log(d_{\max}) \sqrt{\frac{\|\mathbf{A}\|_{\text{F}}^2 d_{\max}}{n}} + C_2 t^2 k^5 \log^{k+4}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_{\text{F}}}{n}.$$

Finalize the proof of Lemma 1. Put the above bounds together, we end up with, for any $t > 1$,

$$\|\widehat{\mathbf{A}}^{\text{SPA}} - \mathbf{A}\| \leq C_1 t^2 k^4 \log(d_{\max}) \sqrt{\frac{\|\mathbf{A}\|_{\text{F}}^2 d_{\max}}{n}} + C_2 t^2 k^5 \log^{k+4}(d_{\max}) \frac{(d_1 \dots d_k)^{1/2} \|\mathbf{A}\|_{\text{F}}}{n}$$

which holds with probability at least $1 - (1 + k \log d_{\max}) d_{\max}^{-t} = 1 - d_{\max}^{-t + \log(k \log d_{\max})}$. \square

4.2 Proof of Theorem 2

We begin with symmetrization (see, e.g., [Yuan and Zhang, 2016](#)) and obtain for any $t > 0$,

$$\mathbb{P}\left(\|\widehat{\mathbf{A}} - \mathbf{A}\| \geq t\right) \leq 4\mathbb{P}\left(\|\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}\| \geq 2t\right) + 4 \exp\left(\frac{-t^2/2}{\alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P}) + t\alpha_{\infty}(\mathbf{A}, \mathbf{P})/3}\right)$$

where $\boldsymbol{\varepsilon} \in \mathbb{R}^{d_1 \times \dots \times d_k}$ is a random tensor with i.i.d. Rademacher entries, and

$$\alpha_{\infty}(\mathbf{A}, \mathbf{P}) = \max_{i_j \in [d_j], j=1, \dots, k} \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}$$

and

$$\alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) = \max_{i_j \in [d_j], j \in [k]} \left(\frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right)^{1/2}.$$

The \odot operator stands for entrywise multiplication, that is

$$(\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}})(i_1, \dots, i_k) = \boldsymbol{\varepsilon}(i_1, \dots, i_k) \widehat{\mathbf{A}}(i_1, \dots, i_k).$$

By definition, the operator norm $\|\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}\|$ is given by

$$\|\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}\| = \sup_{\mathbf{u}_j \in \mathbb{R}^{d_j}, \|\mathbf{u}_j\|_{\ell_2} \leq 1, 1 \leq j \leq k} \langle \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}, \mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k \rangle.$$

We begin with the discretization of ℓ_2 -norm balls. For each $j = 1, \dots, k$, define

$$\mathfrak{B}_{m_j, d_j} = \left\{0, \pm 1, \pm 2^{-1/2}, \dots, \pm 2^{-m_j/2}\right\}^{d_j} \cap \left\{\mathbf{u} \in \mathbb{R}^{d_j} : \|\mathbf{u}\|_{\ell_2} \leq 1\right\}$$

where $m_j = 2(\lceil \log_2 d_j \rceil + 3)$. Define the “digitalization” operator \mathbf{D}_s which zeros out the entries of \mathbf{A} whose absolute value is not $2^{-s/2}$. Then,

$$\mathbf{D}_s(\mathbf{A}) = \sum_{i_1, \dots, i_k} \mathbf{1}\left\{\left|\langle \mathbf{A}, \mathbf{e}_{i_1} \otimes \dots \otimes \mathbf{e}_{i_k} \rangle\right| = 2^{-s/2}\right\} A(i_1, \dots, i_k) \mathbf{e}_{i_1} \otimes \dots \otimes \mathbf{e}_{i_k}$$

where we denote by \mathbf{e}_{i_j} the canonical basis vectors in \mathbb{R}^{d_j} . Clearly, for all $\mathbf{u}_j \in \mathfrak{B}_{m_j, d_j}$,

$$\left\langle \mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \right\rangle = \sum_{s=1}^{m_1 + \dots + m_k} \left\langle \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \right\rangle.$$

For a subset $\mathcal{T} \subset [d_1] \times \dots \times [d_k]$, the aspect ratio $\mu_{\mathcal{T}}$ is defined by

$$\mu_{\mathcal{T}} := \max_{\ell=1, \dots, k} \max_{i_j: j \in [\ell]} \text{Card}\left(\left\{i_\ell : (i_1, \dots, i_k) \in \mathcal{T}\right\}\right).$$

Define the sampling locations

$$\Omega = \left\{(i_1, \dots, i_k) : \Delta(i_1, \dots, i_k) = 1\right\}$$

and the associated sampling operator

$$\mathcal{P}_{\Omega}(\mathbf{A}) = \sum_{i_1, \dots, i_k} \mathbf{1}\left((i_1, \dots, i_k) \in \Omega\right) A(i_1, \dots, i_k) \mathbf{e}_{i_1} \otimes \dots \otimes \mathbf{e}_{i_k}.$$

We shall now make use of the following version of the Chernoff bound:

Lemma 2. *Let X_1, \dots, X_n be independent binary random variables such that $\mathbb{P}(X_j = 1) = p_j \in [0, 1], j = 1, \dots, n$. Then, for any $t \geq 0$,*

$$\mathbb{P}\left(\sum_{j=1}^n (X_j - p_j) \geq 2t \sqrt{\sum_{j=1}^n p_j(1 - p_j)}\right) \leq e^{-t^2}.$$

Lemma 2 is fairly standard and we include its proof in the Appendix for completeness.

By Lemma 2, there exists an absolute constant $C > 0$ such that for all $\alpha \geq 1$,

$$\mathbb{P}\left(\mu_{\Omega} \geq C\alpha \max\left\{\beta(\mathbf{P}), k \log d_{\max}\right\}\right) \leq d_{\max}^{-\alpha}$$

where

$$\beta(\mathbf{P}) = \max_{j=1,\dots,k} \max_{i_1,\dots,i_{j-1},i_{j+1},\dots,i_k} \sum_{i_j=1}^{d_j} P(i_1, \dots, i_k)$$

and $d_{\max} := \max_{1 \leq j \leq k} d_j$. Denote the above event by \mathcal{E}_1 with $\mathbb{P}(\mathcal{E}_1) \geq 1 - d_{\max}^{-\alpha}$. The rest of our analysis is conditioned on event \mathcal{E}_1 . Observe that

$$\langle \mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle = \sum_{s=1}^{m_1+\dots+m_k} \langle \mathcal{P}_\Omega(\mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k)), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle.$$

For $\mathbf{u}_j \in \mathfrak{B}_{m_j, d_j}$, let $\mathcal{A}_{b_j} = \{i_j : |u_j(i_j)| = 2^{-b_j/2}\}$ for $j = 1, \dots, k$. Then, we write

$$\mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k) = \sum_{(b_1, \dots, b_k) : b_1 + \dots + b_k = s} \mathcal{P}_{\mathcal{A}_{b_1} \times \dots \times \mathcal{A}_{b_k}} \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k).$$

By definition of μ_Ω , on event \mathcal{E}_1 , there exist $\tilde{\mathcal{A}}_{b_1} \subset \mathcal{A}_{b_1}, \dots, \tilde{\mathcal{A}}_{b_k} \subset \mathcal{A}_{b_k}$ such that

$$(\mathcal{A}_{b_1} \times \dots \times \mathcal{A}_{b_k}) \cap \Omega = (\tilde{\mathcal{A}}_{b_1} \times \dots \times \tilde{\mathcal{A}}_{b_k}) \cap \Omega$$

and

$$\text{Card}^2(\tilde{\mathcal{A}}_{b_j}) \leq \mu_\Omega \prod_{j=1}^k \text{Card}(\tilde{\mathcal{A}}_{b_j}), \quad j = 1, 2, \dots, k.$$

We conclude with

$$\langle \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle = \sum_{s=1}^{m_1+\dots+m_k} \sum_{b_1+\dots+b_k=s} \langle \mathcal{P}_{\tilde{\mathcal{A}}_{b_1} \times \dots \times \tilde{\mathcal{A}}_{b_k}} \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle.$$

Given Ω , we define the balanced version of digitalization operator

$$\widetilde{\mathbf{D}}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k) = \sum_{(b_1, \dots, b_k) : b_1 + \dots + b_k = s} \mathcal{P}_{\tilde{\mathcal{A}}_{b_1} \times \dots \times \tilde{\mathcal{A}}_{b_k}} \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k)$$

where $\tilde{\mathcal{A}}_j$ are defined as above. Then, $\mathcal{P}_\Omega \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k) = \mathcal{P}_\Omega \widetilde{\mathbf{D}}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k)$. Given Ω , define

$$\mathfrak{B}_{\Omega, m_\star} := \left\{ \sum_{0 \leq s \leq m_\star} \widetilde{\mathbf{D}}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k) + \sum_{m_\star < s \leq m^\star} \mathbf{D}_s(\mathbf{u}_1 \otimes \dots \otimes \mathbf{u}_k) : \mathbf{u}_j \in \mathfrak{B}_{m_j, d_j}, j = 1, \dots, k \right\}$$

for any $0 < m_\star \leq m^\star \leq \sum_{j=1}^k m_j$. Conditioned on \mathcal{E}_1 , we shall focus on $\{\Omega : \mu_\Omega \leq \nu\}$ where $\nu = C\alpha \max\{\beta(\mathbf{P}), k \log d_{\max}\}$. Denote $\mathfrak{B}_{\nu, m_\star}^\star = \bigcup_{\mu_\Omega \leq \nu} \mathfrak{B}_{\Omega, m_\star}^\star$. Following an identical argument as that in [Yuan and Zhang \(2016\)](#), we get

$$\|\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}\| \leq 2^k \max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^\star} \langle \mathbf{Y}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle.$$

The entropy number of $\mathfrak{B}_{\nu, m_\star}^\star$ plays an essential role in bounding $\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^\star} \langle \mathbf{Y}, \mathbf{X} \rangle$. Observe that $\mathfrak{B}_{\nu, m_\star}^\star \subset \mathfrak{B}_{m_1, d_1} \times \dots \times \mathfrak{B}_{d_k, m_k}$ and

$$\begin{aligned} \text{Card}(\mathfrak{B}_{m_j, d_j}) &\leq \prod_{k=0}^{m_j} \binom{d_j}{2^k \wedge d_j} 2^{2^k \wedge d_j} \\ &\leq \prod_{k=0}^{m_j} \exp \left((2^k \wedge d_j) (\log 2 + 1 + (\log d_j / 2^k)_+) \right) \\ &\leq \exp \left(d_j \sum_{\ell=1}^{\infty} 2^{-\ell} (\log 2 + 1 + \log(2^\ell)) \right) \\ &\leq \exp(21d_j/4), \end{aligned}$$

which implies that

$$\log \text{Card}(\mathfrak{B}_{\nu, m_\star}^\star) \leq \frac{21}{4} (d_1 + \dots + d_k).$$

See [Yuan and Zhang \(2016\)](#) for more details. More precise characterizations of $\text{Card}(\mathfrak{B}_{\nu, m_\star}^\star)$ can also be derived. For any $0 \leq q \leq s \leq m_\star$, define

$$\mathfrak{D}_{\nu, s, q} = \left\{ \mathbf{D}_s(\mathbf{Y}) : \mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^\star, \|\mathbf{D}_s(\mathbf{Y})\|_{\ell_2}^2 \leq 2^{q-s} \right\}.$$

Lemma 3. *Let $\nu \geq 1$. For all $0 \leq q \leq s \leq m_\star$, the following bound holds*

$$\log \text{Card}(\mathfrak{D}_{\nu, s, q}) \leq qs^k \log 2 + 2k^2 s^k \sqrt{\nu 2^q} L(\sqrt{\nu 2^q}, d_{\max} s^{k/2})$$

where $L(x, y) = \max \{1, \log(ey/x)\}$.

We write

$$\begin{aligned} \|\varepsilon \odot \widehat{\mathbf{A}}\| &\leq 2^k \max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^\star} \langle \mathbf{Y}, \varepsilon \odot \widehat{\mathbf{A}} \rangle \\ &= 2^k \max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^\star} \left(\sum_{0 \leq s \leq m_\star} \langle \mathbf{D}_s(\mathbf{Y}), \varepsilon \odot \widehat{\mathbf{A}} \rangle + \langle \mathbf{S}_\star(\mathbf{Y}), \varepsilon \odot \widehat{\mathbf{A}} \rangle \right) \end{aligned}$$

where $\mathbf{S}_\star(\mathbf{Y}) = \sum_{s > m_\star} \mathbf{D}_s(\mathbf{Y})$. The actual value of m_\star is to be determined later.

Upper bound of $\left| \langle \mathbf{D}_s(\mathbf{Y}), \varepsilon \odot \widehat{\mathbf{A}} \rangle \right|$. Recall the definition of $\mathfrak{D}_{\nu, s, q}$ and that

$$2^{-s} \leq \|\mathbf{D}_s(\mathbf{Y})\|_{\ell_2}^2 \leq 1,$$

we can write

$$\mathbf{D}_s(\mathbf{Y}) \in \bigcup_{q=1}^s \left(\mathfrak{D}_{\nu,s,q} \setminus \mathfrak{D}_{\nu,s,q-1} \right).$$

Then

$$\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^*} \langle \mathbf{D}_s(\mathbf{Y}), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle = \max_{1 \leq q \leq s} \max_{\mathbf{Y}_{s,q} \in \mathfrak{D}_{\nu,s,q} \setminus \mathfrak{D}_{\nu,s,q-1}} \langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle.$$

Observe that

$$\langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle = \sum_{i_j \in [d_j], j=1, \dots, k} \frac{\Delta(i_1, \dots, i_k)}{P(i_1 \dots i_k)} \varepsilon(i_1, \dots, i_k) A(i_1, \dots, i_k) Y_{s,q}(i_1, \dots, i_k),$$

where Δ is a binary random tensor and $\boldsymbol{\varepsilon}$ is a Rademacher random tensor. Both of them have i.i.d. entries. By definition of $\mathbf{Y}_{s,q}$ and $\mathfrak{D}_{\nu,s,q}$, we have $\max_{i_1, \dots, i_k} |Y_{s,q}(i_1, \dots, i_k)| \leq 2^{-s/2}$.

Moreover,

$$\text{Var}(\langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle) = \sum_{i_j \in [d_j], j=1, \dots, k} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} Y_{s,q}^2(i_1, \dots, i_k).$$

Since $\|\mathbf{Y}_{s,q}\|_{\text{F}}^2 \leq 2^{q-s}$, we obtain

$$\text{Var}(\langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle) \leq \max_{i_j \in [d_j], j \in [k]} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \|\mathbf{Y}_{s,q}\|_{\text{F}}^2 \leq 2^{q-s} \max_{i_j \in [d_j], j \in [k]} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}.$$

Recall the definition of $\alpha_\infty(\mathbf{A}, \mathbf{P})$ and $\alpha_{2,\infty}(\mathbf{A}, \mathbf{P})$. By Bernstein inequality for sum of bounded random variables, there exist absolute constants $C_0, C_1, C_2 > 0$ such that

$$\mathbb{P}\left(\left|\langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle\right| \geq t\right) \leq \exp\left(-\frac{C_0 t^2}{C_1 2^{q-s} \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P}) + C_2 2^{-s/2} t \alpha_\infty(\mathbf{A}, \mathbf{P})}\right)$$

for any $t > 0$. By the union bound and Lemma 3, we get

$$\begin{aligned} & \mathbb{P}\left(\max_{\mathbf{Y}_{s,q} \in \mathfrak{D}_{\nu,s,q}} \left|\langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle\right| \geq t\right) \\ & \leq \text{Card}(\mathfrak{D}_{\nu,s,q}) \exp\left(-\frac{C_0 t^2}{C_1 2^{q-s} \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P}) + C_2 2^{-s/2} t \alpha_\infty(\mathbf{A}, \mathbf{P})}\right) \\ & \leq \exp\left(21\left(\sum_{j=1}^k d_j\right)/4 - \frac{C_0 t^2}{C_1 2^{q-s} \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})}\right) \\ & \quad + \exp\left(qs^k \log 2 + 2k^2 s^k \sqrt{\nu 2^q} L\left(\sqrt{\nu 2^q}, d_{\max} s^{k/2}\right) - \frac{C_0 2^{s/2} t^2}{C_2 t \alpha_\infty(\mathbf{A}, \mathbf{P})}\right). \end{aligned}$$

Recall that

$$0 \leq q \leq s \leq m_\star \lesssim k \log d_{\max}$$

and

$$L\left(\sqrt{\nu 2^q}, d_{\max} s^{k/2}\right) \lesssim \frac{k}{2} \log d_{\max}.$$

For large enough constants $C_3, C_4 > 0$, by choosing $t > 0$ such that

$$t \geq C_3 2^{(q-s)/2} \left(\sum_{j=1}^k d_j \right)^{1/2} \alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) + C_4 k^3 \log^{k+1} d_{\max} \sqrt{\nu} 2^{q-s} \alpha_{\infty}(\mathbf{A}, \mathbf{P}),$$

we get for any $0 \leq q \leq s \leq m_*$,

$$\mathbb{P}\left(\max_{\mathbf{Y}_{s,q} \in \mathfrak{D}_{\nu,s,q}} \left| \langle \mathbf{Y}_{s,q}, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle \right| \geq t\right) \leq \exp\left(-\frac{C_0 t^2}{C_1 2^{q-s} \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})}\right) + \exp\left(-\frac{C_0 2^{s/2} t}{C_2 \alpha_{\infty}(\mathbf{A}, \mathbf{P})}\right).$$

By making the above bound uniform over all pairs $0 \leq q \leq s \leq m_*$, we obtain

$$\begin{aligned} \mathbb{P}\left(\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_*}^*} \left| \sum_{0 \leq s \leq m_*} \langle \mathbf{D}_s(\mathbf{Y}), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle \right| \geq (m_* + 1)t\right) &\leq \binom{m_* + 1}{2} \exp\left(-\frac{C_0 t^2}{C_1 \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})}\right) \\ &\quad + \binom{m_* + 1}{2} \exp\left(-\frac{C_0 t}{C_2 \alpha_{\infty}(\mathbf{A}, \mathbf{P})}\right). \end{aligned}$$

Upper bound of $\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_*}^*} |\langle \mathbf{S}_*(\mathbf{Y}), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle|$. For notation simplicity, we write \mathbf{S}_* in short for $\mathbf{S}_*(\mathbf{Y})$. We apply Bernstein inequality to

$$\langle \mathbf{S}_*, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle = \sum_{i_j \in [d_j], j=1, \dots, k} \frac{\Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \varepsilon(i_1, \dots, i_k) A(i_1, \dots, i_k) S_*(i_1, \dots, i_k).$$

Clearly, $|S_*(i_1, \dots, i_k)| \leq 2^{-m_*/2}$. Meanwhile,

$$\text{Var}\left(\langle \mathbf{S}_*, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle\right) = \sum_{i_j \in [d_j], j=1, \dots, k} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} S_*^2(i_1, \dots, i_k).$$

Following an identical approach as previously, we show that

$$\text{Var}\left(\langle \mathbf{S}_*, \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle\right) \leq \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P}).$$

By Bernstein inequality and the union bound

$$\begin{aligned} &\mathbb{P}\left(\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_*}^*} \left| \langle \mathbf{S}_*(\mathbf{Y}), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle \right| \geq t\right) \\ &\leq \text{Card}(\mathfrak{B}_{\nu, m_*}^*) \exp\left(-\frac{C_0 t^2}{C_1 \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P}) + C_2 2^{-m_*/2} t \alpha_{\infty}(\mathbf{A}, \mathbf{P})}\right) \\ &\leq \exp\left(21 \sum_{j=1}^k d_j / 4 - \frac{C_0 t^2}{C_1 \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})}\right) + \exp\left(21 \sum_{j=1}^k d_j / 4 - \frac{C_0 2^{m_*/2} t}{C_2 \alpha_{\infty}(\mathbf{A}, \mathbf{P})}\right) \end{aligned}$$

for some absolute constants $C_0, C_1, C_2 > 0$. For large enough constants $C_3, C_4 > 0$, by choosing t such that

$$t \geq C_3 \left(\sum_{j=1}^k d_j \right)^{1/2} \alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) + C_4 \left(\sum_{j=1}^k d_j \right) 2^{-m_\star/2} \alpha_\infty(\mathbf{A}, \mathbf{P}),$$

we obtain

$$\mathbb{P} \left(\max_{\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^*} \left| \langle \mathbf{S}_\star(\mathbf{Y}), \boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}} \rangle \right| \geq t \right) \leq \exp \left(- \frac{C_0 t^2}{C_1 \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})} \right) + \exp \left(- \frac{C_0 2^{m_\star/2} t}{C_2 \alpha_\infty(\mathbf{A}, \mathbf{P})} \right).$$

Finalize the proof of Theorem 2. Combining above bounds, we conclude that if for large enough constants $C_3, C_4, C_5 > 0$ such that

$$t \geq C_3 \left(\sum_{j=1}^k d_j \right)^{1/2} \alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) + C_4 k^3 \log^{k+1}(d_{\max}) \sqrt{\nu} \alpha_\infty(\mathbf{A}, \mathbf{P}) + C_5 \left(\sum_{j=1}^k d_j \right) 2^{-m_\star/2} \alpha_\infty(\mathbf{A}, \mathbf{P}).$$

Thus

$$\begin{aligned} \mathbb{P} \left(\|\boldsymbol{\varepsilon} \odot \widehat{\mathbf{A}}\| \geq (m_\star + 2)t \right) &\leq \left(\binom{m_\star + 1}{2} + 1 \right) \exp \left(- \frac{C_0 t^2}{C_1 \alpha_{2,\infty}^2(\mathbf{A}, \mathbf{P})} \right) \\ &\quad + \left(\binom{m_\star + 1}{2} + 1 \right) \exp \left(- \frac{C_0 t}{C_2 \alpha_\infty(\mathbf{A}, \mathbf{P})} \right). \end{aligned}$$

Recall that $\nu = C_1 \alpha \max \{ \beta(\mathbf{P}), k \log d_{\max} \}$ and $m_\star \leq \sum_{j=1}^k 2 \left(\lceil \log_2 d_j \rceil + 3 \right)$. By choosing m_\star large enough such that $2^{-m_\star/2} \left(\sum_{j=1}^k d_j \right) \leq \sqrt{\nu}$, we conclude that for any $\gamma > 0$ such that

$$t \geq C_3 \left(\left(\sum_{j=1}^k d_j \right)^{1/2} + \gamma k \log d_{\max} \right) \alpha_{2,\infty}(\mathbf{A}, \mathbf{P}) + C_4 \gamma k^3 \log^{k+2}(d_{\max}) \sqrt{\nu} \alpha_\infty(\mathbf{A}, \mathbf{P}).$$

It follows immediately, by adjusting the constant C_3 , that

$$\mathbb{P} \left(\|\widehat{\mathbf{A}} - \mathbf{A}\| \geq t \right) \leq 2d_{\max}^{-\gamma}.$$

4.3 Proof of Theorem 3

It suffices to prove the upper bound of $\|\widehat{\mathbf{M}}_j - \mathbf{M}_j\|$ where $\mathbf{M}_j = \mathcal{M}_j(\mathbf{A})$ and $\widehat{\mathbf{M}}_j = \mathcal{M}_j(\widehat{\mathbf{A}}^{\text{SPA}})$.

Without loss of generality, let $j = 1$. Recall the notation $d_{-1} = d_2 \dots d_k$. By denoting

$\mathbf{E}_{i_1(i_2 \dots i_k)} \in \mathbb{R}^{d_1 \times d_{-1}}$ the canonical basis matrices of $\mathbb{R}^{d_1 \times d_{-1}}$ that is $\mathbf{E}_{i_1(i_2 \dots i_k)}$ has exactly value 1 on the $(i_1, i_2 \dots i_k)$ position and all 0's elsewhere. Then,

$$\widehat{\mathbf{M}}_j - \mathbf{M}_j = \sum_{i_j \in [d_j], 1 \leq j \leq k} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{E}_{i_1(i_2 \dots i_k)}$$

where $\mathbb{P}(\Delta(i_1, \dots, i_k) = 1) = P(i_1, \dots, i_k)$. We shall apply the matrix Bernstein inequality to bound the sum of random matrices for $\widehat{\mathbf{M}}_j - \mathbf{M}_j$. Denote the locations of small entries by

$$\Omega_1 := \left\{ (i_1, \dots, i_k) : \|A(i_1, \dots, i_k)\| \leq \|\mathbf{A}\|_{\text{F}} / (d_1 \dots d_k)^{1/2} \right\} \subset [d_1] \times \dots \times [d_k]$$

moderate entries by

$$\Omega_2 := \left\{ (i_1, \dots, i_k) : \|A(i_1, \dots, i_k)\| / \|\mathbf{A}\|_{\text{F}} \in \left(1/(d_1 \dots d_k)^{1/2}, 1/n^{1/2} \right) \right\} \subset [d_1] \times \dots \times [d_k]$$

and large entries by

$$\Omega_3 := \left\{ (i_1, \dots, i_k) : \|A(i_1, \dots, i_k)\| \geq \|\mathbf{A}\|_{\text{F}} / n^{1/2} \right\} \subset [d_1] \times \dots \times [d_k].$$

Recall that $P(i_1, \dots, i_k) = 1$ for $(i_1, \dots, i_k) \in \Omega_3$. Then, for any $(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2$, we have

$$\left\| \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{E}_{i_1(i_2 \dots i_k)} \right\| \leq \max_{i_j \in [d_j], 1 \leq j \leq k} \left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right|.$$

Moreover,

$$\begin{aligned} & \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{E}_{i_1(i_2 \dots i_k)} \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \right\| \\ & \leq \max_{1 \leq i_1 \leq d_1} \sum_{i_j \in [d_j], 2 \leq j \leq k} \frac{A^2(i_1, \dots, i_k) (1 - P(i_1, \dots, i_k))}{P(i_1, \dots, i_k)} \\ & \leq \max_{1 \leq i_1 \leq d_1} \sum_{i_j \in [d_j], j \geq 2, (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}. \end{aligned}$$

Similarly,

$$\begin{aligned} & \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \mathbf{E}_{i_1(i_2 \dots i_k)} \right\| \\ & \leq \max_{i_j \in [d_j], 2 \leq j \leq k} \sum_{i_1=1}^{d_1} \frac{A^2(i_1, \dots, i_k) (1 - P(i_1, \dots, i_k))}{P(i_1, \dots, i_k)} \\ & \leq \max_{i_j \in [d_j], 2 \leq j \leq k} \sum_{i_1 \in [d_1], (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)}. \end{aligned}$$

Observe that if $(i_1, \dots, i_k) \in \Omega_1$, then

$$\left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right| = \frac{(d_1 \dots d_k)}{n} |A(i_1, \dots, i_k)| \leq \frac{(d_1 \dots d_k)^{1/2}}{n} \|\mathbf{A}\|_{\text{F}}$$

and

$$\frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} = \frac{(d_1 \dots d_k) A^2(i_1, \dots, i_k)}{n} \leq \frac{\|\mathbf{A}\|_{\text{F}}^2}{n}.$$

Similarly, if $(i_1, \dots, i_k) \in \Omega_2$, then

$$\left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right| = \frac{\|\mathbf{A}\|_{\text{F}}^2}{n |A(i_1, \dots, i_k)|} \leq \frac{(d_1 \dots d_k)^{1/2}}{n} \|\mathbf{A}\|_{\text{F}}$$

and

$$\frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} = \frac{\|\mathbf{A}\|_{\text{F}}^2}{n}.$$

By matrix Bernstein inequality ([Tropp, 2012](#)), for any $t \geq 0$, with probability at least $1 - e^{-t}$ that

$$\|\widehat{\mathbf{M}}_j - \mathbf{M}_j\| \leq 2\|\mathbf{A}\|_{\text{F}} \left(\sqrt{\frac{d_2 d_3 \dots d_k (t + k \log d_{\max})}{n}} + \frac{(d_1 \dots d_k)^{1/2} (t + k \log d_{\max})}{n} \right).$$

Since $\widehat{\mathbf{M}}_j = \mathbf{M}_j + (\widehat{\mathbf{M}}_j - \mathbf{M}_j)$, the claim follows directly from Davis-Kahan Theorem as in [\(3\)](#).

4.4 Proof of Theorem [4](#)

Theorem [4](#) is an immediate consequence of the following concentration bound.

Lemma 4. *Let $\mathbf{U}_j^{(r_j)}$ be the r_j leading left singular vectors of $\mathcal{M}_j(\mathbf{A})$, and $\widehat{\mathbf{U}}_j^{(r_j)}$ be the output from Algorithm [2](#). There exist constants $C_1, C_2 > 0$ depending on k only such that if*

$$n \geq C_1 (d_1 \dots d_k)^{1/2} (t + \log d_{\max}),$$

then for any $t \geq 0$, the following bound holds with probability at least $1 - e^{-t}$:

$$\begin{aligned} & \left\| \widehat{\mathbf{U}}_j^{(r_j)} (\widehat{\mathbf{U}}_j^{(r_j)})^\top - \mathbf{U}_j^{(r_j)} (\mathbf{U}_j^{(r_j)})^\top \right\| \\ & \leq C_2 \frac{\|\mathbf{A}\|_{\text{F}}}{\bar{g}_{r_j}(\mathbf{M}_j \mathbf{M}_j^\top)} \left(\sigma_{\max}(\mathbf{M}_j) \sqrt{\frac{d_j (t + \log d_{\max})}{n}} + \|\mathbf{A}\|_{\text{F}} \frac{(d_1 \dots d_k)^{1/2} (t + \log d_{\max})}{n} \right). \end{aligned}$$

Proof of Lemma 4. With out loss of generality, we assume $j = 1$ without loss of generality.

In this case, $\widehat{\mathbf{M}}_j^{(1)} = \mathcal{M}_j(\widehat{\mathbf{A}}_1^{\text{SPA}})$, $\widehat{\mathbf{M}}_j^{(2)} = \mathcal{M}_j(\widehat{\mathbf{A}}_2^{\text{SPA}}) \in \mathbb{R}^{d_1 \times (d_2 \dots d_k)}$. Observe that

$$\begin{aligned} \widehat{\mathbf{M}}_j^{(1)} \left(\widehat{\mathbf{M}}_j^{(2)} \right)^\top &= \mathbf{M}_j \mathbf{M}_j^\top + \left(\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j \right) \mathbf{M}_j^\top + \mathbf{M}_j \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top \\ &\quad + \left(\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j \right) \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top. \end{aligned}$$

Upper bound of $\left\| \left(\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j \right) \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top \right\|$. Denote by $\mathbf{Z}_1 = \widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j$. By Theorem 3, there exists an event \mathcal{E}_1 with $\mathbb{P}(\mathcal{E}_1) \geq 1 - e^{-t}$ such that on event \mathcal{E}_1 ,

$$\|\mathbf{Z}_1\| \leq C \|\mathbf{A}\|_F \left(\sqrt{\frac{d_2 d_3 \dots d_k (t + \log d_{\max})}{n}} + \frac{(d_1 \dots d_k)^{1/2} (t + \log d_{\max})}{n} \right).$$

Denote by $\|\mathbf{Z}_1\|_{2,\infty}$ the maximal column ℓ_2 norm., i.e., $\|\mathbf{Z}_1\|_{2,\infty} = \max_{j \in [d_2 \dots d_k]} \|\mathbf{Z}_1 \mathbf{e}_j\|_{\ell_2}$.

Clearly, there exists a constant C_1 depending on k only such that

$$\begin{aligned} \|\mathbf{Z}_1\|_{2,\infty} &\leq C_1 \left(\max_{i_j \in [d_j], 2 \leq j \leq k} \sqrt{\sum_{i_1 \in [d_1]: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} (t + \log d_{\max})} \right. \\ &\quad \left. + \max_{(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right| (t + \log d_{\max}) \right) \\ &\leq C_1 \|\mathbf{A}\|_F \left(\sqrt{\frac{d_1 (t + \log d_{\max})}{n}} + \frac{(d_1 \dots d_k)^{1/2} (t + \log d_{\max})}{n} \right), \end{aligned}$$

which holds with probability at least $1 - e^{-t}$. Denote the above event by \mathcal{E}_3 . We shall proceed conditional on $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3$. Write

$$\mathbf{Z}_1 \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top = \sum_{i_j \in [d_j], 1 \leq j \leq k} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{Z}_1 \mathbf{E}_{i_1(i_2 \dots i_k)}^\top$$

which is again a sum of random matrices. Clear, for any $(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2$,

$$\begin{aligned} &\left\| \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{Z}_1 \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \right\| \\ &\leq \max_{(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right| \|\mathbf{Z}_1\|_{2,\infty} \\ &\leq \frac{(d_1 \dots d_k)^{1/2}}{n} \|\mathbf{A}\|_F \|\mathbf{Z}_1\|_{2,\infty}. \end{aligned}$$

Moreover,

$$\begin{aligned}
& \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{Z}_1 \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \mathbf{E}_{i_1(i_2 \dots i_k)} \mathbf{Z}_1^\top \right\| \\
& \leq \max_{i_j \in [d_j], 2 \leq j \leq k} \|\mathbf{Z}_1\|^2 \sum_{i_1 \in [d_1]: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \\
& \leq \frac{d_1 \|\mathbf{A}\|_{\text{F}}^2}{n} \|\mathbf{Z}_1\|^2.
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{E}_{i_1(i_2 \dots i_k)} \mathbf{Z}_1^\top \mathbf{Z}_1 \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \right\| \\
& \leq \max_{(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \|\mathbf{Z}_1\|_{\text{F}}^2 \\
& \leq \frac{d_1 \|\mathbf{A}\|_{\text{F}}^2}{n} \|\mathbf{Z}_1\|^2.
\end{aligned}$$

By matrix Bernstein inequality, the following bound holds with probability at least $1 - e^{-t}$,

$$\begin{aligned}
& \left\| \left(\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j \right) \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top \right\| \\
& \leq C \|\mathbf{A}\|_{\text{F}} \left(\sqrt{\frac{d_1(t + \log d_{\max})}{n}} \|\mathbf{Z}_1\| + \frac{(d_1 \dots d_k)^{1/2}(t + \log d_{\max})}{n} \|\mathbf{Z}_1\|_{2, \infty} \right).
\end{aligned}$$

Denote the above event by \mathcal{E}_4 . On event $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \cap \mathcal{E}_4$, if

$$n \geq C_1 (d_1 d_2 \dots d_k)^{1/2} (t + \log d_{\max}),$$

then

$$\begin{aligned}
& \left\| \left(\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j \right) \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top \right\| \\
& \leq C_2 \|\mathbf{A}\|_{\text{F}}^2 \left(\frac{(d_1 \dots d_k)^{1/2}(t + \log d_{\max})}{n} + \frac{d_1^{1/2} (d_1 \dots d_k)^{1/2} (t + \log d_{\max})^{3/2}}{n^{3/2}} \right) \\
& \leq C_2 \|\mathbf{A}\|_{\text{F}}^2 \frac{(d_1 \dots d_k)^{1/2}(t + \log d_{\max})}{n}.
\end{aligned}$$

Upper bound of $\left\| \mathbf{M}_j \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top \right\|$. We write

$$\mathbf{M}_j \left(\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j \right)^\top = \sum_{i_j \in [d_j], 1 \leq j \leq k} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{M}_j \mathbf{E}_{i_1(i_2 \dots i_k)}^\top.$$

The proof follows identically as above. Indeed, for any $(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2$,

$$\begin{aligned}
& \left\| \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right) \mathbf{M}_j \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \right\| \\
& \leq \max_{(i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \left| \frac{A(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right| \max_{i_j \in [d_j], 2 \leq j \leq k} \sqrt{\sum_{i_1: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} A^2(i_1, \dots, i_k)} \\
& \leq \frac{(d_1 \dots d_k)^{1/2}}{n} \left(\frac{d_1}{n} \right)^{1/2} \|\mathbf{A}\|_{\text{F}}^2.
\end{aligned}$$

Moreover,

$$\begin{aligned}
& \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{M}_j \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \mathbf{E}_{i_1(i_2 \dots i_k)} \mathbf{M}_j^\top \right\| \\
& \leq \max_{i_j \in [d_j], 2 \leq j \leq k} \sum_{i_1: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \|\mathbf{M}_j\|^2 \\
& \leq \frac{d_1}{n} \|\mathbf{A}\|_{\text{F}}^2 \sigma_{\max}^2(\mathbf{M}_j).
\end{aligned}$$

Similarly,

$$\begin{aligned}
& \left\| \sum_{i_j \in [d_j], 1 \leq j \leq k} \mathbb{E} \left(\frac{A(i_1, \dots, i_k) \Delta(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} - A(i_1, \dots, i_k) \right)^2 \mathbf{E}_{i_1(i_2 \dots i_k)} \mathbf{M}_j^\top \mathbf{M}_j \mathbf{E}_{i_1(i_2 \dots i_k)}^\top \right\| \\
& \leq \left(\max_{i_j \in [d_j], 2 \leq j \leq k} \sum_{i_1: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} A^2(i_1, \dots, i_k) \right) \left(\max_{i_1 \in [d_1]} \sum_{i_j \in [d_j], 2 \leq j \leq k: (i_1, \dots, i_k) \in \Omega_1 \cup \Omega_2} \frac{A^2(i_1, \dots, i_k)}{P(i_1, \dots, i_k)} \right) \\
& \leq \frac{d_1 d_2 \dots d_k}{n^2} \|\mathbf{A}\|_{\text{F}}^4.
\end{aligned}$$

By matrix Bernstein inequality ([Tropp, 2012](#)), if $n \geq C_1(d_1 \dots d_k)^{1/2}(t + \log d_{\max})$, then with probability at least $1 - e^{-t}$ such that

$$\begin{aligned}
& \left\| \mathbf{M}_j (\widehat{\mathbf{M}}_j^{(2)} - \mathbf{M}_j)^\top \right\| \\
& \leq C_2 \|\mathbf{A}\|_{\text{F}} \left(\sigma_{\max}(\mathbf{M}_j) \sqrt{\frac{d_1(t + \log d_{\max})}{n}} + \|\mathbf{A}\|_{\text{F}} \frac{(d_1 \dots d_k)^{1/2}(t + \log d_{\max})}{n} \right).
\end{aligned}$$

Denote this event by \mathcal{E}_5 . Clearly, an identical bound holds for $\left\| (\widehat{\mathbf{M}}_j^{(1)} - \mathbf{M}_j) \mathbf{M}_j^\top \right\|$ with the same probability. Denote this event by \mathcal{E}_6 .

Finalize the proof of Theorem 4. On event $\mathcal{E}_1 \cap \mathcal{E}_2 \cap \mathcal{E}_3 \cap \mathcal{E}_4 \cap \mathcal{E}_5 \cap \mathcal{E}_6$, if $n \geq C_1(d_1 \dots d_k)^{1/2}(t + \log d_{\max})$, there exists a constant C_2 depending on k only such that

$$\begin{aligned} & \left\| \widehat{\mathbf{M}}_j^{(1)} \left(\widehat{\mathbf{M}}_j^{(2)} \right)^\top - \mathbf{M}_j \mathbf{M}_j^\top \right\| \\ & \leq C_2 \|\mathbf{A}\|_F \left(\sigma_{\max}(\mathbf{M}_j) \sqrt{\frac{d_1(t + \log d_{\max})}{n}} + \|\mathbf{A}\|_F \frac{(d_1 \dots d_k)^{1/2}(t + \log d_{\max})}{n} \right), \end{aligned}$$

which concludes the proof by adjusting the constant C_2 and applying Davis-Kahan Theorem. \square

References

- Martín Abadi, Ashish Agarwal, Paul Barham, Eugene Brevdo, Zhifeng Chen, Craig Citro, Greg S Corrado, Andy Davis, Jeffrey Dean, and Matthieu Devin. Tensorflow: Large-scale machine learning on heterogeneous distributed systems. *arXiv preprint arXiv:1603.04467*, 2016.
- Dimitris Achlioptas and Frank McSherry. Fast computation of low-rank matrix approximations. *Journal of the ACM (JACM)*, 54(2):9, 2007.
- Dimitris Achlioptas, Zohar S Karnin, and Edo Liberty. Near-optimal entrywise sampling for data matrices. In *Advances in Neural Information Processing Systems*, pages 1565–1573, 2013.
- Sanjeev Arora, Elad Hazan, and Satyen Kale. A fast random sampling algorithm for sparsifying matrices. In *APPROX-RANDOM*, volume 6, pages 272–279. Springer, 2006.
- Michael W Berry. Large-scale sparse singular value computations. *The International Journal of Supercomputing Applications*, 6(1):13–49, 1992.
- Srinadh Bhojanapalli and Sujay Sanghavi. A new sampling technique for tensors. *arXiv preprint arXiv:1502.05023*, 2015.
- Andrzej Cichocki, Danilo Mandic, Lieven De Lathauwer, Guoxu Zhou, Qibin Zhao, Cesar Caiafa, and Huy Anh Phan. Tensor decompositions for signal processing applications:

- From two-way to multiway component analysis. *IEEE Signal Processing Magazine*, 32(2):145–163, 2015.
- Aaron Clauset, Mark EJ Newman, and Cristopher Moore. Finding community structure in very large networks. *Physical review E*, 70(6):066111, 2004.
- Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM Journal on Numerical Analysis*, 7(1):1–46, 1970.
- Petros Drineas and Anastasios Zouzias. A note on element-wise matrix sparsification via a matrix-valued bernstein inequality. *Information Processing Letters*, 111(8):385–389, 2011.
- Petros Drineas, Michael W Mahoney, and S Muthukrishnan. Subspace sampling and relative-error matrix approximation: Column-based methods. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 316–326. Springer, 2006.
- Alan Frieze, Ravi Kannan, and Santosh Vempala. Fast monte-carlo algorithms for finding low-rank approximations. *Journal of the ACM (JACM)*, 51(6):1025–1041, 2004.
- Gene H Golub and Charles F Van Loan. *Matrix computations*, volume 3. JHU Press, 2012.
- Michael Holmes, Alexander Gray, and Charles Isbell. Fast SVD for large-scale matrices. In *Workshop on Efficient Machine Learning at NIPS*, volume 58, pages 249–252, 2007.
- Yuval Kluger, Ronen Basri, Joseph T Chang, and Mark Gerstein. Spectral biclustering of microarray data: coclustering genes and conditions. *Genome research*, 13(4):703–716, 2003.
- Mei Kobayashi, Georges Dupret, Oliver King, and Hikaru Samukawa. Estimation of singular values of very large matrices using random sampling. *Computers & Mathematics with Applications*, 42(10-11):1331–1352, 2001.
- Tamara G Kolda and Brett W Bader. Tensor decompositions and applications. *SIAM review*, 51(3):455–500, 2009.

- Akshay Krishnamurthy and Aarti Singh. Low-rank matrix and tensor completion via adaptive sampling. In *Advances in Neural Information Processing Systems*, pages 836–844, 2013.
- Nan Li and Baoxin Li. Tensor completion for on-board compression of hyperspectral images. In *Image Processing (ICIP), 2010 17th IEEE International Conference on*, pages 517–520. IEEE, 2010.
- Ji Liu, Przemyslaw Musialski, Peter Wonka, and Jieping Ye. Tensor completion for estimating missing values in visual data. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 35(1):208–220, 2013.
- Aditya Krishna Menon and Charles Elkan. Fast algorithms for approximating the singular value decomposition. *ACM Transactions on Knowledge Discovery from Data (TKDD)*, 5(2):13, 2011.
- Nam H Nguyen, Petros Drineas, and Trac D Tran. Tensor sparsification via a bound on the spectral norm of random tensors. *Information and Inference: A Journal of the IMA*, 4(3):195–229, 2015.
- Román Orús. A practical introduction to tensor networks: Matrix product states and projected entangled pair states. *Annals of Physics*, 349:117–158, 2014.
- John Scott. *Social network analysis*. Sage, 2017.
- Nicholas D Sidiropoulos, Lieven De Lathauwer, Xiao Fu, Kejun Huang, Evangelos E Papalexakis, and Christos Faloutsos. Tensor decomposition for signal processing and machine learning. *IEEE Transactions on Signal Processing*, 65(13):3551–3582, 2017.
- Stephen M Smith, Mark Jenkinson, Mark W Woolrich, Christian F Beckmann, Timothy EJ Behrens, Heidi Johansen-Berg, Peter R Bannister, Marilena De Luca, Ivana Drobnjak, and David E Flitney. Advances in functional and structural mr image analysis and implementation as FSL. *Neuroimage*, 23:S208–S219, 2004.

Ulrich Stelzl, Uwe Worm, Maciej Lalowski, Christian Haenig, Felix H Brembeck, Heike Goehler, Martin Stroedicke, Martina Zenkner, Anke Schoenherr, and Susanne Koeppen. A human protein-protein interaction network: a resource for annotating the proteome. *Cell*, 122(6):957–968, 2005.

Joel A Tropp. User-friendly tail bounds for sums of random matrices. *Foundations of computational mathematics*, 12(4):389–434, 2012.

David P Woodruff et al. Sketching as a tool for numerical linear algebra. *Foundations and Trends® in Theoretical Computer Science*, 10(1–2):1–157, 2014.

Dong Xia and Ming Yuan. On polynomial time methods for exact low rank tensor completion. *arXiv preprint arXiv:1702.06980*, 2017.

Ming Yuan and Cun-Hui Zhang. On tensor completion via nuclear norm minimization. *Foundations of Computational Mathematics*, 16(4):1031–1068, 2016.

Ming Yuan and Cun-Hui Zhang. Incoherent tensor norms and their applications in higher order tensor completion. *IEEE Transactions on Information Theory*, 63(10):6753–6766, 2017.

A Technical Lemmas

A.1 Proof of Lemma 2

Clearly, for any t and $\lambda > 0$,

$$\begin{aligned}
\mathbb{P}\left(\sum_{j=1}^n (X_j - p_j) \geq t\right) &= \mathbb{P}\left(\exp\left\{\lambda \sum_{j=1}^n (X_j - p_j)\right\} \geq \exp\{\lambda t\}\right) \\
&\leq e^{-\lambda t} \mathbb{E} \exp\left\{\lambda \sum_{j=1}^n (X_j - p_j)\right\} \\
&\leq e^{-\lambda t} \prod_{j=1}^n \mathbb{E} e^{\lambda(X_j - p_j)} \\
&\leq e^{-\lambda t} \prod_{j=1}^n (p_j e^{\lambda(1-p_j)} + (1-p_j) e^{-\lambda p_j}).
\end{aligned}$$

Note that $e^x \leq 1 + x + x^2$ for any $x \in [-1, 1]$. Then,

$$p_j e^{\lambda(1-p_j)} + (1-p_j)e^{-\lambda p_j} \leq 1 + \lambda^2 p_j(1-p_j) \leq e^{\lambda^2 p_j(1-p_j)}.$$

Therefore, we obtain

$$\mathbb{P}\left(\sum_{j=1}^n (X_j - p_j) \geq t\right) \leq e^{-\lambda t} \prod_{j=1}^n e^{\lambda^2 p_j(1-p_j)} = \exp\left\{-\lambda t + \lambda^2 \sum_{j=1}^n p_j(1-p_j)\right\}.$$

By choosing $\lambda = t/2 \sum_{j=1}^n p_j(1-p_j)$, we end up with

$$\mathbb{P}\left(\sum_{j=1}^n (X_j - p_j) \geq t\right) \leq \exp\left\{-t^2/4 \sum_{j=1}^n p_j(1-p_j)\right\}.$$

The proof is closed after choosing $t = 2s\sqrt{\sum_{j=1}^n p_j(1-p_j)}$ for $s \geq 0$.

A.2 Proof of Lemma 3

The proof follows from the same argument as that for Lemma 12 of [Yuan and Zhang \(2016\)](#).

More specifically, denote the aspect ratio for a block $A_1 \times \dots \times A_k \subset [d_1] \times \dots \times [d_k]$,

$$h(A_1 \times \dots \times A_k) = \min \left\{ \nu : |A_j|^2 \leq \nu \prod_{j=1}^k |A_j|, j = 1, 2, \dots, k \right\}.$$

We bound the entropy of a single block. Let

$$\begin{aligned} \mathfrak{D}_{\nu, \ell}^{(\text{block})} &= \left\{ \text{sgn}(u_1(a_1)) \dots \text{sgn}(u_k(a_k)) \mathbf{1}\{(a_1, \dots, a_k) \in A_1 \times \dots \times A_k\} : \right. \\ &\quad \left. h(A_1 \times \dots \times A_k) \leq \nu, \prod_{j=1}^k |A_j| = \ell \right\}. \end{aligned}$$

By definition, we obtain

$$\max(|A_1|^2, \dots, |A_k|^2) \leq \nu |A_1| |A_2| \dots |A_k| \leq \nu \ell.$$

By dividing $\mathfrak{D}_{\nu, \ell}^{(\text{block})}$ into subsets according to $(\ell_1, \dots, \ell_k) = (|A_1|, \dots, |A_k|)$, we find

$$\left| \mathfrak{D}_{\nu, \ell}^{(\text{block})} \right| \leq \sum_{\ell_1 \dots \ell_k = \ell, \max_j \ell_j \leq \sqrt{\nu \ell}} 2^{\ell_1 + \dots + \ell_k} \binom{d_1}{\ell_1} \dots \binom{d_k}{\ell_k}.$$

By the Stirling formula, for $j = 1, 2, \dots, k$,

$$\binom{d_j}{\ell_j} \leq \frac{d_j^{\ell_j}}{(\ell_j!)} \leq \left(\frac{d_j}{\ell_j}\right)^{\ell_j} e^{\ell_j} \frac{1}{\sqrt{2\pi \ell_j}},$$

then

$$\log \left[\sqrt{2\pi\ell_j} 2^{\ell_j} \binom{d_j}{\ell_j} \right] \leq \ell_j L(\ell_j, 2d_{\max}) \leq \sqrt{\nu\ell} L(\sqrt{\nu\ell}, 2d_{\max})$$

where $L(x, y) := \max\{1, \log(ey/x)\}$. Let $\ell = \prod_{j=1}^m p_j^{v_j}$ with distinct prime factors p_j . Since $(v_j + 1)v_j/(2p_j^{v_j/2})$ is upper bounded by 2.66 for $p_j = 2$, by 1.16 for $p_j = 3$ and by 1 for $p_j \geq 5$, we get

$$\begin{aligned} \left| \{(\ell_1, \dots, \ell_k) : \ell_1 \dots \ell_k = \ell\} \right| &= \prod_{j=1}^m \binom{v_j + 1}{k - 1} \\ &\leq \prod_{j=1}^m \binom{v_j + 1}{2}^{k/2} \\ &\leq (2.66 \times 1.16)^{k/2} (\sqrt{\ell})^{k/2} \\ &\leq \prod_{j=1}^k \left(2\sqrt{2\pi\ell_j} \right)^{k/2}, \quad \forall \prod_{j=1}^k \ell_j = \ell. \end{aligned}$$

Therefore,

$$\begin{aligned} \left| \mathfrak{D}_{\nu, \ell}^{(\text{block})} \right| &\leq \frac{\exp \left(k\sqrt{\nu\ell} L(\sqrt{\nu\ell}, 2d_{\max}) \right)}{\prod_{j=1}^k \sqrt{2\pi\ell_j}} \prod_{j=1}^k \left(2\sqrt{2\pi\ell_j} \right)^{k/2}, \quad \forall (\ell_1 \dots \ell_k) = \ell \\ &\leq 2^{k^2/2} (2\pi)^{k(k-2)/4} \ell^{(k-2)/4} \exp \left(k\sqrt{\nu\ell} L(\sqrt{\nu\ell}, 2d_{\max}) \right) \\ &\leq 2^{k^2/2} (2\pi)^{k(k-2)/4} \exp \left(2k\sqrt{\nu\ell} L(\sqrt{\nu\ell}, 2d_{\max}) \right). \end{aligned}$$

Due to the constraint $b_1 + b_2 + \dots + b_k = s$ in defining $\mathfrak{B}_{\nu, m_\star}^*$, for any $\mathbf{Y} \in \mathfrak{B}_{\nu, m_\star}^*$, $\mathbf{D}_s(\mathbf{Y})$ is composed of at most $i^\star := \binom{s+k-1}{k-1}$ blocks. Since the sum of the sizes of the blocks is bounded by 2^q , we obtain

$$\begin{aligned} \left| \mathfrak{D}_{\nu, s, q} \right| &\leq \sum_{\ell_1 + \dots + \ell_{i^\star} \leq 2^q} \prod_{i=1}^{i^\star} \left| \mathfrak{D}_{\nu, \ell_i}^{(\text{block})} \right| \\ &\leq \sum_{\ell_1 + \dots + \ell_{i^\star} \leq 2^q} (2\pi)^{i^\star k(k-2)/4} 2^{i^\star k^2/2} \exp \left(2k \sum_{i=1}^{i^\star} \sqrt{\nu\ell_i} L(\sqrt{\nu\ell_i}, 2d_{\max}) \right) \\ &\leq 2^{i^\star k^2/2} (2^q)^{i^\star} (2\pi)^{i^\star k(k-2)/4} \max_{\ell_1 + \dots + \ell_{i^\star} \leq 2^q} \exp \left(2k \sum_{i=1}^{i^\star} \sqrt{\nu\ell_i} L(\sqrt{\nu\ell_i}, 2d_{\max}) \right). \end{aligned}$$

As shown in [Yuan and Zhang \(2016\)](#), $\sum_{i=1}^{i^\star} \sqrt{\ell_i} L(\sqrt{\nu\ell_i}, 2d_{\max}) \leq \sqrt{i^\star 2^q} \left(L(\sqrt{\nu 2^q}, 2d_{\max}) + \log(\sqrt{i^\star}) \right)$, we obtain

$$\log \left| \mathfrak{D}_{\nu, s, q} \right| \leq i^\star \log(2^q) + i^\star k(k-2)/2 + i^\star k^2/2 + 2k\sqrt{i^\star \nu 2^q} L(\sqrt{\nu 2^q}, 2d_{\max} \sqrt{i^\star}).$$

Since $i^\star = \binom{s+k-1}{k-1} \leq s^k$, it follows that

$$\log \left| \mathfrak{D}_{\nu, s, q} \right| \leq q s^k \log 2 + 2k^2 s^k \sqrt{\nu 2^q} L \left(\sqrt{\nu 2^q}, d_{\max} s^{k/2} \right).$$