

# Empirical likelihood inference for partial functional linear regression models based on B-spline

Mingao Yuan and Yue Zhang

*Department of mathematical science,  
Indiana University-Purdue University Indianapolis,  
Indianapolis, IN 46202, USA*  
E-mail: mingyuan@iupui.edu, yz65@umail.iu.edu

November 2, 2017

## Abstract

In this paper, we apply empirical likelihood method to inference for the regression parameters in the partial functional linear regression models based on B-spline. We prove that the empirical log-likelihood ratio for the regression parameters converges in law to a weighted sum of independent chi-square distributions. Our simulation shows that the proposed empirical likelihood method produces more accurate confidence regions in terms of coverage probability than the asymptotic normality method.

**Keywords:** empirical likelihood, partial functional linear regression, B-spline.

## 1. Introduction

With the rapid development of measurement apparatus and computers, it is possible that the data are collected over an entire time period. In the literature, this kind of data are called *functional data*. One of the goal of functional data analysis is to explain the variations of a dependent variable by using an independent functional variable. Functional linear model is one of the most popular models to realize this. There are mainly two approaches for estimation and inference in functional linear analysis. One of them is the functional principle component analysis(FPCA), which has the advantages of interpretability and availability of a good estimate of the slope function(Cai

and Hall(2006), Hall and Horowitz (2007), Shin (2009), Yuan and Cai (2010), Cai and Yuan (2012)). Another approach is the polynomial spline method. As a commonly used method in nonparametric or semiparametric analysis, the polynomial spline method has been introduced into functional data analysis by different authors(Ramsay and Silverman(1997), Cardot et al. (2003, 2005)). Ramsay and Silverman (1997,2005) and Cardot et al.(2003) employed spline to estimate the functional slope and Cardot et al.(2005) proposed a spline estimator for the functional coefficient in quantile regression.

To improve the accuracy of prediction and make the functional linear model more interpretable, sometimes other predictor variables should be incorporated into the model. Especially, Zhang et al. (2007) introduced the *partial functional linear model* and applied it to analyze the effect of women's hormone on the total hip bone mineral density. Later, using functional principle component analysis, Shin (2009) proposed a new estimator for the parameters. For estimators based on B-spline in partial functional linear model, Zhou et al. (2016) established the asymptotic normality for the regression parameters and the global convergence rate for the slope function.

The empirical likelihood(EL) was introduced by Owen(1990, 2001) to construct confidence region in a nonparametric setting. As an analog of the parametric likelihood method, it has been extensively applied to different fields due to some of the nice properties. As nonparametric method, it doesn't require a prespecified distribution for the data. The confidence region respects the range of the data and usually performs better than that based on asymptotic normality. Recently, the EL method has been used for inferences under different models(Zhao (2010), Cheng et al.(2012)). In this paper, we propose the empirical likelihood based confidence region for the regression parameters in partial functional linear model and compare it with the ones based on asymptotic normality proposed in Zhou et al. (2016).

The paper is organized as follows. Section 2 presents the partial functional linear model and the asymptotic normality based confidence region. In section 3, we develop the empirical likelihood confidence regions for the regression parameters. Section 4 includes simulation studies to illustrate the advantage of the EL based confidence region over asymptotic normality based confidence region in terms of coverage probability. The proof is presented in Section 5.

## 2. Asymptotic normality method

The partial functional linear model is

$$Y = \mathbf{Z}^T \beta + \int_0^1 X(t) \alpha(t) dt + \epsilon,$$

where  $Y$  is scalar response variable,  $\mathbf{Z}$  is  $p$ -dimensional predictor variable and  $X(t)$  is a random process in  $L^2([0, 1])$ . We assume that the random error  $\epsilon$  is independent of  $X(t)$  and  $\mathbf{Z}$ ,  $E(\epsilon) = 0$  and  $Var(\epsilon) = \sigma^2$ . The  $\beta$  is unknown  $p$ -dimensional parameter vector and  $\alpha(t)$  is an unknown slope function in  $L^2([0, 1])$ . Denote  $\|\phi_1\| = (\int_0^1 \phi_1^2 dt)^{\frac{1}{2}}$  and  $\langle \phi_1, \phi_2 \rangle = \int_0^1 \phi_1(t) \phi_2(t) dt$ , for  $\phi_1, \phi_2 \in L^2([0, 1])$ .

Let  $0 = t_0 < t_1 < \dots < t_{N_n} < t_{N_n+1} = 1$  be a knot sequence in  $[0, 1]$ . A polynomial spline of degree  $k$  ( $k \geq 0$ ) is a function such that it is a polynomial of degree  $k$  on each interval  $[t_j, t_{j+1}]$  ( $j = 0, 1, \dots, N_n$ ) and is  $k - 1$  times continuously differentiable in  $[0, 1]$ . Let  $S_{k, N_n}$  be the linear space spanned by splines with degree  $k$  and number of knots  $N_n$ . It is well known that  $S_{k, N_n}$  is of dimension  $k_n = N_n + k + 1$ . Let  $B_1, \dots, B_{k_n}$  be the B-spline basis. For  $\alpha(t) \in C^{k+1}([0, 1])$ , we can approximate it by a unique linear combination of splines, that is,

$$\alpha(t) \approx \sum_{i=1}^{k_n} b_i B_i.$$

Let  $(X_i, \mathbf{Z}_i, Y_i), i = 1, \dots, n$  be the data from the model. Then the model can be written approximately as

$$Y_i = \mathbf{Z}_i^T \beta + \sum_{s=1}^{k_n} b_s \langle X_i, B_s \rangle + \epsilon_i, \quad i = 1, \dots, n.$$

Then the least square estimators of  $\mathbf{b} = (b_1, \dots, b_{k_n})^T$  and  $\beta$  are the minimizers of the following loss function

$$\sum_{i=1}^n \left( Y_i - \mathbf{Z}_i^T \beta - \sum_{s=1}^{k_n} b_s \langle X_i, B_s \rangle \right)^2.$$

The profile least square estimators are given by

$$\hat{\beta} = (\mathbf{Z}^T (I - A) \mathbf{Z})^{-1} \mathbf{Z}^T (I - A) \mathbf{Y}, \quad \hat{\mathbf{b}} = (B^T B)^{-1} B^T (\mathbf{Y} - \mathbf{Z} \hat{\beta}),$$

where  $\mathbf{Y} = (Y_1, \dots, Y_n)^T$ ,  $\mathbf{Z} = (\mathbf{Z}_1, \dots, \mathbf{Z}_n)^T$ ,  $B = [\langle X_i, B_j \rangle]_{\substack{1 \leq j \leq k_n \\ 1 \leq i \leq n}}$ ,  $A = B(B^T B)^{-1} B^T$ . The estimators of  $\alpha(t)$  and  $\sigma^2$  are

$$\hat{\alpha}(t) = \sum_{s=1}^{k_n} \hat{b}_s B_s, \quad \hat{\sigma}^2 = \frac{1}{n} \sum (Y_i - \langle X_i, \hat{\alpha} \rangle - \mathbf{Z}_i^T \hat{\beta}).$$

For two positive sequences  $\{a_n\}_{n=1}^{+\infty}$  and  $\{b_n\}_{n=1}^{+\infty}$ ,  $a_n \asymp b_n$  means  $a_n/b_n$  is uniformly bounded away from infinity and zero. To get the asymptotic normality, the following assumptions are required in Zhou et al.(2016).

(A1) There are positive constants  $M$  and  $\frac{1}{4(k+1)} < r < \frac{1}{2}$  such that

$$h = \max_{0 \leq j \leq N_n} (t_{j+1} - t_j) \asymp n^{-r}, \quad k_n \asymp n^{-r}, \quad h / \min_{0 \leq j \leq N_n} (t_{j+1} - t_j) \leq M.$$

(A2)  $E\|X\|^4 < +\infty$  and the eigenvalues of the covariance operator  $\Gamma$  of  $X$  are strictly positive.

(A3)  $E|Z_{11}|^4 + \dots + E|Z_{1p}|^4 + E|\epsilon_1|^4 < +\infty$ .

(A4)  $E(Z_{1j}|X_1) = \langle X_1, g_j \rangle$  for some function  $g_j \in L^2([0, 1])$ , for  $j = 1, \dots, p$ . Besides, we assume  $g_j, \alpha \in C^{k+1}([0, 1])$ .

(A5) Let  $\eta_1 = (\eta_{11}, \dots, \eta_{1p})^T$  with  $\eta_{1j} = Z_{1j} - E(Z_{1j}|X_1)$ . Assume  $\Sigma = \text{Var}(\eta_1)$  is positive definite.

Under conditions (A1)-(A5), Zhou et al. (2016) established the asymptotic normality for the regression parameters and global convergence rate for the slope function as follows.

**Lemma 1.** *Under condition (A1)-(A5), we have*

$$\sqrt{n}(\hat{\beta} - \beta_0) \Rightarrow N(0, \sigma^2 \Sigma^{-1}),$$

$$\|\hat{\alpha} - \alpha\|^2 = O_p\left(\frac{k_n}{n} + k_n^{-2(k+1)}\right),$$

$$\frac{n}{\hat{\sigma}^2}(\hat{\beta} - \beta_0)^T \hat{\Sigma}^{-1}(\hat{\beta} - \beta_0) \Rightarrow \chi_p^2,$$

where  $\Sigma = \text{Var}(\mathbf{Z}_1 - E(\mathbf{Z}_1|X_1))$ , "  $\Rightarrow$  " represents convergence in law and

$$\hat{\Sigma} = \frac{1}{n} \mathbf{Z}^T (I - A) \mathbf{Z}.$$

Then the asymptotic  $(1 - \gamma)\%$  confidence region is

$$\left\{ \beta : \frac{n}{\hat{\sigma}^2} (\hat{\beta} - \beta)^T \hat{\Sigma}^{-1} (\hat{\beta} - \beta) \leq \chi_p^2(1 - \gamma) \right\}.$$

### 3. The empirical likelihood method

Motivated by the estimation equation, we define

$$\mathbf{W}_i(\beta) = \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - B_i^{*T} (B^T B)^{-1} B^T (Y - \mathbf{Z} \beta) \right),$$

where  $B_i^{*T}$  is the  $i$ -th row of the matrix  $B$ . The empirical likelihood at  $\beta$  is given by

$$\mathcal{R}_n(\beta) = \sup \left\{ \prod_{i=1}^n n \pi_i \mid \sum_{i=1}^n \pi_i \mathbf{W}_i = 0, \sum_{i=1}^n \pi_i = 1, \pi_i \geq 0 \right\}.$$

By Lagrange method as in Owen(2001), the solution is

$$\pi_i = \frac{1}{n} \frac{1}{1 + \lambda^T \mathbf{W}_i},$$

where  $\lambda$  satisfies

$$\frac{1}{n} \sum_{i=1}^n \frac{\mathbf{W}_i}{1 + \lambda^T \mathbf{W}_i} = 0. \quad (1)$$

**Theorem 1.** *Suppose (A1)-(A5) hold. Then at the true  $\beta_0$ , we have*

$$-2 \log \mathcal{R}_n(\beta_0) \Rightarrow \mathbf{U}^T \Sigma_0 \mathbf{U},$$

where " $\Rightarrow$ " means "convergence in distribution",  $\mathbf{U} \sim \mathcal{N}(0, I_p)$ ,  $\Sigma_0 = \Sigma^{\frac{1}{2}} \Sigma_1^{-1} \Sigma^{\frac{1}{2}}$ ,  $\Sigma = \text{Var}(\mathbf{Z}_1 - E(\mathbf{Z}_1 | X_1))$  and  $\Sigma_1 = E(\mathbf{Z}_1 \mathbf{Z}_1^T)$ .

The limiting distribution of the empirical likelihood ratio is not the usual chi-square distribution. Actually, it is a weighted sum of independent chi-square distributions with degree of freedom 1. However, the confidence regions based on the this empirical likelihood ratio still have the advantages of having natural shape and respecting the range of  $\beta$ . In application,  $\Sigma$  and  $\Sigma_1$  are unknown and we propose the following consistent estimators

$$\hat{\Sigma} = \frac{1}{n} \sum_{i=1}^n \mathbf{Z}_i \mathbf{Z}_i^T, \quad \hat{\Sigma}_1 = \frac{1}{n} \mathbf{Z}^T (I - A) \mathbf{Z}. \quad (2)$$

To construct confidence regions, we use the Monte Carlo method to simulate the limiting distribution and compute the desired quantile.

#### 4. Simulation study

In this section, we present simulation results to compare the finite sample behaviors of empirical likelihood(EL) method and the asymptotic normality(NA) method in terms of coverage probability of the confidence regions. As in Lian (2011), we generate  $X_i$  by the following method,

$$X_i = \sum_{j=1}^{50} \xi_{ij} j^{-1} \phi_j(t),$$

where  $\phi_1(t) = 1$ ,  $\phi_j(t) = \sqrt{2} \cos((j-1)\pi t)$  for  $j \geq 2$  and  $\xi_{ij}$  are iid with uniform distribution  $U[-\sqrt{3}, \sqrt{3}]$ . We simulate data based on the following three models, which are similar to the models in Zhou et al.(2016).

**Model 1:**  $Y_i = Z_{i1} + Z_{i2} + \int_0^1 X_i(t)\alpha(t)dt + \epsilon_i$ ,  $\alpha(t) = \frac{\sqrt{2}}{2} + \sum_{j=2}^{50} 4j^{-2}\phi_j(t)$ .  $(Z_{i1}, Z_{i2})$  are from bivariate normal with zero mean vector,  $Var(Z_{i1}) = 0.9$ ,  $Var(Z_{i2}) = 0.5$  and  $Cov(Z_{i1}, Z_{i2}) = 0.2$ . The error follows  $\mathcal{N}(0, 0.36)$  and skewed normal distribution with mean 0, standard deviation 1 and skewness parameter 5.

**Model 2:**  $Y_i = 5Z_{i1} - 1.7Z_{i2} + \int_0^1 X_i(t)\alpha(t)dt + \epsilon_i$ . We generate  $(Z_{i1}, Z_{i2})$  from standard bivariate normal distribution and use the following functional coefficient,

$$\alpha(t) = 2 \sin(0.5\pi t) + 4 \sin(1.5\pi t) + 5 \sin(2.5\pi t).$$

The error follows  $\mathcal{N}(0, 1)$  and skewed normal distribution with mean 0, standard deviation 1 and skewness parameter 5.

**Model 3:**  $Y_i = 2Z_{i1} - Z_{i2} + \int_0^1 X_i(t)\alpha(t) + \epsilon_i$ , where

$$Z_{i1} = \int_0^1 X_i(t)\alpha_1(t)dt + \epsilon_{i1}, \quad Z_{i2} = \int_0^1 X_i(t)\alpha_2(t)dt + \epsilon_{i2},$$

$\alpha_1(t) = \sum_{j=1}^{50} b_{1j}\phi_j(t)$ ,  $\alpha_2(t) = \sum_{j=1}^{50} b_{2j}\phi_j(t)$ ,  $b_{11} = 1$ ,  $b_{21} = -0.5$ ,  $b_{1j} = 2j^{-2}$ ,  $b_{2j} = 3j^{-2}$  for  $j \geq 2$ . The random error  $\epsilon_{i1} \sim \mathcal{N}(0, 0.25)$  and  $\epsilon_{i2} \sim \mathcal{N}(0, 0.64)$ . The random model error  $\epsilon_i \sim \mathcal{N}(0, 0.25)$  and skewed normal distribution with mean 0, standard deviation 0.5 and skewness parameter 5.

We use B-spline basis with equally spaced knots and degree 2. The number of knots are selected by the "leave-one-out" cross-validation(Rice and Silverman 1991). We compute the confidence regions for  $\beta$  based on the EL

Table 1: Coverage Probabilities Under Model 1

		Normal Error		Skewed Normal Error	
Sample size $n$	$1 - \gamma$	EL	NA	EL	NA
30	0.90	0.793	0.760	0.779	0.763
	0.95	0.867	0.841	0.854	0.825
50	0.90	0.840	0.830	0.831	0.822
	0.95	0.909	0.897	0.895	0.890
80	0.90	0.843	0.843	0.862	0.853
	0.95	0.914	0.906	0.922	0.915
150	0.90	0.895	0.889	0.902	0.909
	0.95	0.945	0.934	0.952	0.962

method and the NA method. To simulate the limiting distribution in Theorem 1, we firstly estimate the unknown variance and second moment by (2) and then use Monte Carlo method to simulate the  $(1 - \gamma)$  quantiles. The sample sizes are 30, 50, 80 and 150, representing small, moderate and large sample sizes. The coverage probability is computed by 1000 simulation runs, with nominal confidence level 0.90 and 0.95 respectively.

The simulation results under Model 1 are summarized in Table 1, with results under Model 2 and Model 3 in Table 2 and Table 3 respectively. There is a similar pattern in all three tables. As the sample size increases, the coverage probabilities increase towards the nominal level, regardless of the error type. However, the EL method outperforms the NA method, since the coverage probabilities based on EL method is larger and closer to the nominal level than that of the NA method. Hence, the EL method yields more accurate confidence regions than the NA method.

## 5. Proof

In this section, we present the proof of Theorem 1.

**Lemma 2.** *Under conditions (A1)-(A5), we have*

$$\frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{W}_i(\beta_0) \Rightarrow \mathcal{N}(0, \sigma^2 \Sigma^{-1}).$$

Table 2: Coverage Probabilities Under Model 2

		Normal Error		Skewed Normal Error	
Sample size $n$	$1 - \gamma$	EL	NA	EL	NA
30	0.90	0.776	0.752	0.770	0.746
	0.95	0.850	0.819	0.846	0.820
50	0.90	0.831	0.818	0.821	0.802
	0.95	0.907	0.897	0.888	0.876
80	0.90	0.875	0.875	0.860	0.858
	0.95	0.934	0.929	0.914	0.909
150	0.90	0.889	0.883	0.874	0.867
	0.95	0.939	0.937	0.925	0.917

Table 3: Coverage Probabilities Under Model 3

		Normal Error		Skewed Normal Error	
Sample size $n$	$1 - \gamma$	EL	NA	EL	NA
30	0.90	0.770	0.744	0.786	0.760
	0.95	0.839	0.826	0.832	0.816
50	0.90	0.841	0.829	0.837	0.834
	0.95	0.906	0.893	0.879	0.877
80	0.90	0.873	0.862	0.859	0.856
	0.95	0.926	0.921	0.918	0.915
150	0.90	0.889	0.888	0.879	0.874
	0.95	0.941	0.927	0.933	0.919



**Proof:** Note that

$$\begin{aligned}
\frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{W}_i &= \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - B_i^{*T} (B^T B)^{-1} B^T (Y - \mathbf{Z} \beta) \right) \\
&= \frac{1}{\sqrt{n}} \mathbf{Z} \left( Y - \mathbf{Z} \beta - B^T (B^T B)^{-1} B^T (Y - \mathbf{Z} \beta) \right) \\
&= \frac{1}{\sqrt{n}} \mathbf{Z} \left( (I - A) Y - (I - A) \mathbf{Z} \beta \right) \\
&= \frac{1}{\sqrt{n}} \mathbf{Z} (I - A) \mathbf{Z}^T \left( (\mathbf{Z} (I - A) \mathbf{Z}^T)^{-1} \mathbf{Z} (I - A) Y - \beta \right) \\
&= \frac{1}{n} \mathbf{Z} (I - A) \mathbf{Z}^T \sqrt{n} (\hat{\beta} - \beta).
\end{aligned}$$

By the Lemma 2 and Theorem 1 in Zhou et al. (2016), the last term converges to  $\mathcal{N}(0, \sigma^2 \Sigma^{-1})$ . □

**Lemma 3.** *Under conditions (A1)-(A5), we have*

$$\frac{1}{n} \sum_{i=1}^n \mathbf{W}_i(\beta_0) \mathbf{W}_i(\beta_0)^T \xrightarrow{P} \sigma^2 \Sigma_1.$$

**Proof:** Firstly, we rewrite the  $W_i$  below

$$\begin{aligned}
\mathbf{W}_i(\beta) &= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - B_i^{*T} (B^T B)^{-1} B^T (Y - \mathbf{Z} \beta) \right) \\
&= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - B_i^{*T} (B^T B)^{-1} B^T (Y - \mathbf{Z} \hat{\beta} + \mathbf{Z} (\hat{\beta} - \beta)) \right) \\
&= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - B_i^{*T} \hat{\mathbf{b}} - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) \right) \\
&= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - \langle X_i, \hat{\alpha} \rangle - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) \right) \\
&= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - \langle X_i, \hat{\alpha} \rangle - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) \right) \\
&= \mathbf{Z}_i \left( Y_i - \mathbf{Z}_i^T \beta - \langle X_i, \alpha \rangle - \langle X_i, \hat{\alpha} - \alpha \rangle - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) \right) \\
&= \mathbf{Z}_i \left( \epsilon_i - \langle X_i, \hat{\alpha} - \alpha \rangle - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) \right).
\end{aligned}$$

Then

$$\begin{aligned}
\frac{1}{n} \sum_{i=1}^n \mathbf{W}_i \mathbf{W}_i^T &= \frac{1}{n} \sum_{i=1}^n \mathbf{Z}_i \mathbf{Z}_i^T \left\{ \epsilon_i^2 + \langle X_i, \hat{\alpha} - \alpha \rangle^2 - 2 \langle X_i, \hat{\alpha} - \alpha \rangle \epsilon_i \right. \\
&\quad + B_i^{*T} (B^T B)^{-1} B^T Z (\hat{\beta} - \beta) (\hat{\beta} - \beta)^T Z^T (B^T B)^{-1} B_i^* \\
&\quad - 2 B_i^{*T} (B^T B)^{-1} B^T Z (\hat{\beta} - \beta) \epsilon_i \\
&\quad \left. + 2 B_i^{*T} (B^T B)^{-1} B^T Z (\hat{\beta} - \beta) \langle X_i, \hat{\alpha} - \alpha \rangle \right\} \\
&= R_{n1} + R_{n2} + R_{n3} + R_{n4} + R_{n5} + R_{n6}.
\end{aligned}$$

By the law of large number and the fact  $\epsilon_i$  and  $\mathbf{Z}_i$  are independent, we have

$$R_{n1} = \frac{1}{n} \sum_{i=1}^n \mathbf{Z}_i \mathbf{Z}_i^T \epsilon_i^2 \xrightarrow{P} E(\epsilon_1^2 \mathbf{Z}_1 \mathbf{Z}_1^T) = \sigma^2 \Sigma_1.$$

Next, we show the rest terms converge to zero in probability.

$$E \langle X_i, \hat{\alpha} - \alpha \rangle^2 \leq E \|\hat{\alpha} - \alpha\|^2 \|X_i\|^2 \leq \sqrt{E \|\hat{\alpha} - \alpha\|^4 E \|X_i\|^4}$$

By Lemma 1 and (A2), we have

$$\langle X_i, \hat{\alpha} - \alpha \rangle^2 = o_p(1),$$

hence  $R_{n2} = o_p(1)$  and  $R_{n3} = o_p(1)$ .

Notice that by Theorem 1 and Lemma 1 in Zhang et al. (2016),

$$\begin{aligned}
B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z} (\hat{\beta} - \beta) &= \frac{1}{n\sqrt{n}} B_i^{*T} \left( \frac{B^T B}{n} \right)^{-1} B^T \mathbf{Z} \sqrt{n} (\hat{\beta} - \beta) \\
&\asymp \frac{1}{n\sqrt{n}} B_i^{*T} B^T \mathbf{Z}.
\end{aligned}$$

By (A2) and (A4), one has

$$\begin{aligned}
E\|B_i^{*T} B^T \mathbf{Z}\|_2 &= E\left\|\sum_{l=1}^n \sum_{j=1}^{k_n} \langle B_j, X_l \rangle \langle B_j, X_i \rangle \mathbf{Z}_l^T\right\|_2 \\
&\leq \sum_{l=1}^n \sum_{j=1}^{k_n} E\left\|\langle B_j, X_l \rangle \langle B_j, X_i \rangle \mathbf{Z}_l^T\right\|_2 \\
&\leq \sum_{l=1}^n \sum_{j=1}^{k_n} E\|B_j\|^2 \|X_l\| \|X_i\| \|\mathbf{Z}_l^T\|_2 \\
&\lesssim nk_n,
\end{aligned}$$

where  $\|\cdot\|_2$  is the Euclidean norm.

Hence,

$$B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z}(\hat{\beta} - \beta) = O_p\left(\frac{k_n}{\sqrt{n}}\right) = o_p(1).$$

Then  $R_{n4} = R_{n5} = R_{n6} = o_p(1)$ . □

**Lemma 4.** *Under the assumption (A1)-(A5), one has*

$$\max_{1 \leq i \leq n} \|\mathbf{W}_i\|_2 = o_p(\sqrt{n}), \quad \|\lambda\|_2 = O_p(n^{-\frac{1}{2}}).$$

**Proof:** By the proof of Lemma 3, we have

$$\epsilon_i - \langle X_i, \hat{\alpha} - \alpha \rangle - B_i^{*T} (B^T B)^{-1} B^T \mathbf{Z}(\hat{\beta} - \beta) = O_p(1).$$

And since  $0 < \text{Var}(\mathbf{Z}_1) < +\infty$ , then  $\max_{1 \leq i \leq n} \|\mathbf{Z}_i\|_2 = o_p(\sqrt{n})$  by Owen (2001), which completes the proof. □

**Proof of Theorem 1:** By (1), Lemma 2 and Lemma 4, we have

$$\lambda(\beta_0) = \left(\frac{1}{n} \sum_{i=1}^n \mathbf{W}_i \mathbf{W}_i^T\right)^{-1} \frac{1}{n} \sum_{i=1}^n \mathbf{W}_i + O_p\left(\frac{1}{\sqrt{n}}\right).$$

Then by Lemma 3, it yields the following Taylor expansion

$$\begin{aligned}
-2 \log \mathcal{R}_n(\beta_0) &= \sum_{i=1}^n \lambda(\beta_0) \mathbf{W}_i(\beta_0) + o_p(1) \\
&= \left( \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{W}_i \right)^T \left( \frac{1}{n} \sum_{i=1}^n \mathbf{W}_i \mathbf{W}_i^T \right)^{-1} \left( \frac{1}{\sqrt{n}} \sum_{i=1}^n \mathbf{W}_i \right) + o_p(1) \\
&\Rightarrow \mathbf{G}^T \Sigma_1^{-1} \mathbf{G},
\end{aligned}$$

where  $\mathbf{G} \sim \mathcal{N}(0, \Sigma^{-1})$ . The proof of Theorem 1 is complete.

## References

### References

- [1] Cai T. T. and Hall P.(2006). Prediction in functional linear regression. *Annals of Statistics*,**34**: 2159-2179.
- [2] Cai, T.T. and Yuan, M.(2016). Minimax and adaptive prediction for functional linear regression. *Journal of American Statistical Association*.**107**: 1201-1216.
- [3] Cardot,H., Crambes,C., Sarda, P.(2005). Quantile regression when the covariates are functions. *Journal of Nonparametric Statistics*,**17**:841-856.
- [4] Cardort, H., Ferraty F. and Sarda, P.(2003). Spline estimators for the functional linear model. *Statistica Sinica*,**13**: 571-591.
- [5] Cheng,G., Zhao,Y. and Li,B.(2012). Empirical likelihood inferences for the semiparametric additive isotonic regression. *Journal of Multivariate Analysis*.**112** :172-182.
- [6] Hall,P. and Horowitz, J.(2007). Methodology and convergence rates for functional linear regression. *Annals of Statistics*,**35**: 70-91.
- [7] Lian, H.(2011). Functional partial linear model. *Journal of Nonparametric Statistics*,**23**:115-128.
- [8] Owen, A.B.(1990). Empirical likelihood confidence region. *Annals of Statistics*,**18**: 90-120.

- [9] Owen, A.B.(2001). *Empirical likelihood*, Chapman Hall/CRC, London.
- [10] Ramsay, J. and Silverman, B.(1997). *Functional data analysis*. Springer, New York.
- [11] Rice, J.A. and Silverman, B.W.(1991). Estimating the mean and covariance structure nonparametrically when the data are curves. *Journal of Royal Statistical Society Series B*,**53**:233-243.
- [12] Shin, H.(2009). Partial functional linear regression. *Journal of Statistical Planning and Inference*,**139**:3405-3418.
- [13] Yuan, M. and Cai, T.T.(2010). A reproducing kernel Hilbert space approach to functional linear regression. *Annals of Statistics*,**38**:3412-3444.
- [14] Zhang, D., Lin, X. and Sowers, M.(2007). Two-stage functional mixed models for evaluating the effect of longitudinal covariate profiles on a scalar outcome. *Biometrics*.**63**: 351-362.
- [15] Zhao, Y.(2010). Semiparametric inference for transformation models via empirical likelihood. *Journal of Multivariate Analysis*.**101**:1846-1858.
- [16] Zhou, J.J., Chen, Z. and Peng, Q.Y.(2016). Polynomial spline estimation for partial functional linear regression models. *Comput Stat.* **31**: 1107-1129.