

# DYNAMICAL HARTREE-FOCK-BOGOLIUBOV APPROXIMATION OF INTERACTING BOSONS

JACKY J. CHONG AND ZEHUA ZHAO

**ABSTRACT.** We consider a many-body Bosonic system with pairwise particle interaction given by  $N^{3\beta-1}v(N^\beta x)$  where  $0 < \beta < 1$  and  $v$  a non-negative spherically-symmetric function. Our main result is the extension of the local-in-time Fock space approximation of the exact dynamics of squeezed states proved in [GM17] for  $0 < \beta < \frac{2}{3}$  to a global-in-time approximation for  $0 < \beta < 1$ . Our work can also be viewed as a generalization of the results in [BCS17] to a more general set of initial data that includes coherent states along with an improved error estimate. The key ingredients in establishing the Fock space approximation are the work of Grillakis and Machedon on the the local well-posedness theory [GM19], some recent established global estimates in [CGMZ20], and our quantitative results on the uniform in  $N$  global well-posedness of the time-dependent Hartree-Fock-Bogoliubov (TBHFB) system.

## 1. INTRODUCTION

We consider a system of  $N$  non-relativistic interacting spinless Bosons in three dimensional space whose dynamics is governed by the  $N$ -body linear Schrödinger equation<sup>1</sup>

$$\left( \frac{1}{i} \frac{\partial}{\partial t} + \sum_{j=1}^N -\Delta_{x_j} + \frac{1}{N} \sum_{i>j} v_N(x_i - x_j) \right) \Psi_N(t, x_1, \dots, x_N) = 0 \quad (1.1)$$

where  $x_i \in \mathbb{R}^3$  and  $v_N(x) = N^{3\beta}v(N^\beta x)$ . In particular, we work exclusively in the setting of repulsive interaction.

The two main goals of this paper are to study of the global-in-time behavior of quantum fluctuations about the mean-field dynamics of (1.1) and provide a quantitative method for effectively tracking the evolution of the many-body quantum system in state space. Unfortunately, the problem of tracking the exact dynamics of Bosonic systems in state space with arbitrary initial data is still not tractable with the current available tools. See [LGS15] for a in-depth overview of the physics of ultracold Bose gas. Nevertheless, if we restrict ourselves to a special class of initial data, such as coherent states or, more generally, squeezed states, we are able to obtain

---

<sup>1</sup>We work under the assumption that  $\hbar = 1$  and  $2m = 1$ , where  $m$  is the mass. However, it would be interesting to incorporate  $\hbar$  in the calculation to see the explicit dependence of  $\hbar$  in our results. Moreover, as written, (1.1) models a system of interacting particles in the mean-field scaling. Cf. section 1.8 of [Gol16].

some positive results in the direction of understanding the exact evolution of the many-body quantum system in state space by means of studying its effective dynamics.

Another interesting question that one could study is the range of the short-range scaling parameter  $\beta$  associated with the interaction potential  $v_N$ , which was first introduced in [ESY07] as a way to model the different length scales in the problem. More precisely, we adopt the convention of setting the scale of the system to order one so that the density  $\rho$  of the system is of order  $N$ . Then the interaction potential  $v_N$  has two length scales: the range of  $v_N$  and the scattering length of  $v_N$ , denoted by  $r_N$  and  $a_N$ , respectively. Here, we see that  $r_N \sim \mathcal{O}(N^{-\beta})$  and  $a_N \sim \mathcal{O}(N^{-1})$ . When  $\beta \in (0, 1)$ , we are in the situation where  $a_N \ll r_N \ll 1$ . Then the effective interaction potential of  $v_N$  is given by  $(\int v)\rho$  where  $\int v$  is the Born approximation of the two-body scattering process. However, when  $\beta = 1$ , also known as the Gross-Pitaevskii regime, we have  $r_N \sim a_N$ . Then the effective two-body scattering is just the full two-body scattering process that means the effective potential is given by  $8\pi\rho a_N \sim \mathcal{O}(1)$ . See appendices of [ESY07, LSSY05] for more detail.

Physically speaking, for  $\beta \in (0, \frac{1}{3})$ , we are in the situation where  $a_N \ll \rho^{-\frac{1}{3}} \ll r_N$  ( $\rho^{-\frac{1}{3}}$  is called the mean inter-particle distance), which model a weakly interacting dense gas. But, once  $\beta \in [\frac{1}{3}, 1)$ , we enter the self-interacting regime or sometimes called the weakly interacting diluted gas regime, that is,  $a_N \ll r_N \ll \rho^{-\frac{1}{3}}$ . A way to see that (1.1) models a weakly interacting gas when  $\beta \in (0, 1)$  is by moving to the microscopic coordinates, i.e.  $(t, x) \mapsto (\tau, q) := (N^{2\beta}t, N^\beta x)$ . Thence, we see that the particle system is weakly coupled with the coupling constant being proportional to  $N^{\beta-1}$ . Let us remark that in three dimensional space the case  $\beta = 1$  is the endpoint case in the heuristic scaling analysis. In fact, when  $\beta = 1$  the system can model a dilute gas with rare but strong collisions, which suggests a more refined analysis is needed since, heuristically, we can no longer treat the interactions as a mere perturbation of the non-interaction case, or, at the very least, not a small perturbation.

In recent years, many have contributed to the studies of effective dynamics for many particle systems. In the case of  $\beta = 0$  with repulsive Coulomb interactions, Erdős and Yau in [EY01] proved the qualitative result, via the method of BBGKY hierarchy, that the one-particle marginal density  $\gamma_{N,t}^{(1)}$  associated with the wave function  $\Psi_{N,t}$  with asymptotically factorized initial state, i.e.  $\Psi_{N,0} \rightarrow \phi^{\otimes N}$  as  $N \rightarrow \infty$ , converges to  $|\phi_t\rangle\langle\phi_t|$  in trace norm in the mean-field limit of  $N \rightarrow \infty$  where  $\phi_t$  satisfies the Hartree equation

$$\frac{1}{i} \frac{\partial}{\partial t} \phi(t, x) - \Delta_x \phi(t, x) + \left( \frac{1}{|\cdot|} * |\phi_t|^2 \right) \phi(t, x) = 0. \quad (1.2)$$

Using the Fock space method introduced by Hepp in [Hep74] and subsequently extended by Ginibre and Velo in [GV79a, GV79b], Rodnianski and

Schlein in [RS09] provided a rate of convergence of the one-particle marginal associated with the many-body quantum system towards the Hartree dynamics in trace norm, that is<sup>2</sup>,

$$\mathrm{Tr} \left| \gamma_{N,t}^{(1)} - |\phi_t\rangle\langle\phi_t| \right| \lesssim \frac{e^{Kt}}{\sqrt{N}}. \quad (1.3)$$

The estimate was later improved to  $e^{Kt}N^{-1}$  in [ES09, CLS11] with the optimal exponent in  $N$ . Using a second-order correction Fock space method introduced by Grillakis, Machedon and Margetis in [GMM10, GMM11], Kuz in [Kuz15] provides a rate of convergence of the many-body quantum system to the Hartree dynamics in the sense of Fock space marginal density<sup>3</sup>. Consequently, Kuz shown that

$$\mathrm{Tr} \left| \gamma_{N,t}^{(1)} - |\phi_t\rangle\langle\phi_t| \right| \lesssim \frac{\sqrt{1+t}}{N^{\frac{1}{4}}} \quad (1.4)$$

which in turn establishes the validity of the approximation for time  $t$  of the order  $\sqrt{N}$ . Similar results were derived in [FKS09, KP10] but the approaches are completely different from the above methods.

For the case  $0 < \beta \leq 1$ , Erdős, Schlein and Yau in a series of papers [ESY06, ESY07, ESY10, ESY09] showed qualitatively that the many-body dynamics with asymptotically factorized initial data converges to the cubic nonlinear Schrödinger dynamics when  $0 < \beta < 1$  or the Gross-Pitaevskii dynamics when  $\beta = 1$ . More precisely, they proved that  $\gamma_{N,t}^{(1)}$  converges to  $|\phi_t\rangle\langle\phi_t|$  in trace norm where  $\phi_t$  satisfies

$$i\partial_t\phi(t, x) + \Delta_x\phi(t, x) = \begin{cases} (\int v) |\phi_t|^2\phi_t & \text{if } 0 < \beta < 1 \\ 8\pi a |\phi_t|^2\phi_t & \text{if } \beta = 1 \end{cases} \quad (1.5)$$

with  $a$  being the scattering length corresponding to the potential  $v$ . Results on the rate of convergence of Fock space marginals can be found in [KP10, BdOS15, Kuz15].

After identifying the mean-field dynamics, it is natural to study the quantum fluctuations about it. A natural setting to account for the fluctuations is in the Bosonic Fock space

$$\mathcal{F}_s(\mathfrak{h}) = \mathbb{C} \oplus \bigoplus_{n \geq 1} \mathrm{Sym}(\mathfrak{h}^{\otimes n})$$

where  $\mathfrak{h} = L^2(\mathbb{R}^3)$ . Introducing  $\mathcal{F}_s$  allows us to deal with states with varying number of particles, which in our model are the excitation and condensate

---

<sup>2</sup>We adopt the standard notation  $A \lesssim B$  to mean there exists a constant, depending on some parameters, such that  $A \leq CB$ .

<sup>3</sup>One should note that the main result in Rodnianski and Schlein's paper is their result on the rate of convergence of the one-particle Fock marginal towards the Hartree dynamics. Whereas, the significance of Kuz's paper is that she was able to show that the mean-field estimate is actually valid for a much longer period of time than most proceeding results had indicated.

elements. Recent works on the evolution of coherent states in Fock space with quantum fluctuation can be found in [RS09, GMM10, GMM11, Che12, GM13a, GM13b, Kuz15, Kuz17, BCS17, NN17, Cho20]. Hence, by accounting for some quantum fluctuations, one is able to track the evolution of the coherent states in Fock space norm, which in effect allows one to obtain  $L^2$ -norm approximation of the evolution of many-body quantum system with factorized initial data. We refer the reader to [LSSY05, Gol16, GMM17] for a complete survey of the subject.

## 2. BACKGROUND AND MAIN RESULT

**2.1. Notations and Earlier Results.** In this section, we provide a brief summary of the results obtained in [GM13a, GM17, GM19] along with relevant notations and rudimentary background materials on second quantization necessary to capture the effects of quantum fluctuations about quasifree states<sup>4</sup>. The presentation is standard in the literature. For reference, the reader may consult [Ber12, Sol14].

Let us denote the one-particle base space by  $\mathfrak{h} := L^2(\mathbb{R}^3, dx)$  endowed with the inner product  $\langle \cdot, \cdot \rangle_{\mathfrak{h}}$  that is linear in the second variable and conjugate linear in the first variable. We define the symmetric (Bosonic) Fock space over  $\mathfrak{h}$  to be the closure of

$$\mathcal{F}_s(\mathfrak{h}) = \mathcal{F}_s := \mathbb{C} \oplus \bigoplus_{n=1}^{\infty} \text{Sym}(\mathfrak{h}^{\otimes n}) \quad (2.1)$$

with respect to the norm induced by the Fock inner product

$$\langle \varphi, \psi \rangle = \bar{\varphi}_0 \psi_0 + \sum_{n=1}^{\infty} \langle \varphi_n, \psi_n \rangle_{\mathfrak{h}^{\otimes n}} \quad (2.2)$$

for every  $\varphi = (\varphi_0, \varphi_1, \dots), \psi = (\psi_0, \psi_1, \dots) \in \mathcal{F}_s(\mathfrak{h})$ . The vacuum, denoted by  $\Omega$ , is defined to be the Fock vector  $(1, 0, 0, \dots) \in \mathcal{F}_s$ .

For every field  $\phi \in \mathfrak{h}$ , we can define the associated creation and annihilation operators on  $\mathcal{F}_s$ , denoted respectively by  $a^*(\phi)$  and  $a(\bar{\phi})$ , as follow

$$(a^*(\phi)\psi)_n(x_1, \dots, x_n) := \frac{1}{\sqrt{n}} \sum_{j=1}^n \phi(x_j) \psi_{n-1}(x_1, \dots, \hat{x}_j, \dots, x_n) \quad (2.3a)$$

$$(a(\bar{\phi})\psi)_n(x_1, \dots, x_n) := \sqrt{n+1} \int dx \bar{\phi}(x) \psi_{n+1}(x, x_1, \dots, x_n) \quad (2.3b)$$

---

<sup>4</sup>Cf. Chapter 10 in [Sol14] for the definition of quasifree.

on each sector with the property that  $a(\phi)\Omega = 0$ . In particular, we could define the corresponding creation and annihilation distribution-valued operators denote by  $a_x^*$  and  $a_x$  as follow

$$(a_x^*\psi)_n := \frac{1}{\sqrt{n}} \sum_{j=1}^n \delta(x - x_j) \psi_{n-1}(x_1, \dots, \hat{x}_j, \dots, x_n), \quad (2.4a)$$

$$(a_x\psi)_n := \sqrt{n+1} \psi_{n+1}(x, x_1, \dots, x_n). \quad (2.4b)$$

Hence, we have the relations

$$a^*(\phi) = \int dx \{ \phi(x) a_x^* \} \quad \text{and} \quad a(\bar{\phi}) = \int dx \{ \bar{\phi}(x) a_x \}.$$

Let us note that the creation and annihilation operators  $a(\bar{\phi})$  and  $a^*(\phi)$  associated with the field  $\phi$  are unbounded, densely defined, closed operators, which are dual to one another. Moreover, one could formally verify the pair  $(a_x^*, a_x)$  satisfies the canonical commutation relation (CCR):  $[a_x, a_y^*] = \delta(x - y)$ ,  $[a_x, a_y] = [a_x^*, a_y^*] = 0$ , and the number operator defined by

$$\mathcal{N} := \int dx a_x^* a_x \quad (2.5)$$

is a diagonal operator on  $\mathcal{F}_s$  that counts the number of particles in each sector; for instance,  $(\mathcal{N}\psi)_n = n\psi_n$ .

For each  $\phi \in \mathfrak{h}$ , we associate a skew-Hermitian operator

$$\mathcal{A}(\phi) := a(\bar{\phi}) - a^*(\phi) \quad (2.6)$$

and define the corresponding Weyl operator to be the unitary operator  $e^{-\sqrt{N}\mathcal{A}(\phi)}$ . Then the coherent state associated with  $\phi$  is given by

$$\psi(\phi) := e^{-\sqrt{N}\mathcal{A}(\phi)}\Omega. \quad (2.7)$$

Using the Baker-Campbell Hausdorff formula, one can show

$$\psi(\phi) = (\dots, c_n \phi^{\otimes n}, \dots) \quad \text{with} \quad c_n = \left( e^{-N\|\phi\|_{\mathfrak{h}}^2} N^n / n! \right).$$

In this context,  $\phi$  is called the condensate wave function and the Fock space marginal associated with the approximation scheme  $e^{-\sqrt{N}\mathcal{A}(\phi_t)}\Omega$  where  $\phi_t = \phi(t)$  satisfies the cubic NLS offers a first-order (mean-field) approximation to exact evolution of the coherent state in trace norm; see [RS09]. However, to track the exact dynamics in Fock space, a first-order approximation is not sufficient. We need to introduce the pair excitation function  $k(x, y) = k(y, x)$  and its corresponding quadratic operator  $\mathcal{B}(k)$ , called the squeezed operator, given by

$$\mathcal{B}(k) := \int dxdy \{ \bar{k}(x, y) a_x a_y - k(x, y) a_x^* a_y^* \}. \quad (2.8)$$

From the pair excitation function, we concoct a new approximation scheme, a second-order correction<sup>5</sup> to the mean-field approximation, given by

$$\psi_{\text{approx}} = e^{i\chi(t)} e^{-\sqrt{N}\mathcal{A}(\phi_t)} e^{-\mathcal{B}(k_t)} \Omega \quad (2.9)$$

where  $\chi(t)$  is some phase factor to be determined. With an appropriate choice of evolution equations for  $\phi$  and  $k$ , we will see that (2.9) will indeed allow us to track the exact dynamics of the evolution of quasifree states in Fock space. Hence we also refer to (2.9) as the quasifree approximation.

For a fixed  $N \in \mathbb{N}$ , we are interested in the time evolution generated by the Fock Hamiltonian operator<sup>6</sup> associated with  $N$  on the Fock space, which is a diagonal operator denoted by  $\mathcal{H}_N$ , defined by its action on Fock vectors of the form  $\psi = (0, \dots, 0, \Psi_n(x_1, \dots, x_n), 0, \dots)$

$$(\mathcal{H}_N \psi)_n = \left( \sum_{j=1}^n \Delta_{x_j} - \frac{1}{N} \sum_{i < j}^n v_N(x_i - x_j) \right) \Psi_n =: H_{N,n} \Psi_n. \quad (2.10)$$

For convenience, we drop the  $N$  subscript in  $\mathcal{H}_N$  since it should be clear from the context. Rewrite  $\mathcal{H}$  using creation and annihilation operators yields

$$\mathcal{H} := \mathcal{H}_0 - N^{-1} \mathcal{V} \quad \text{where} \quad (2.11a)$$

$$\mathcal{H}_0 := \int dx dy \{ \Delta_x \delta(x - y) a_x^* a_y \} \quad \text{and} \quad (2.11b)$$

$$\mathcal{V} := \frac{1}{2} \int dx dy \{ v_N(x - y) a_x^* a_y^* a_x a_y \}. \quad (2.11c)$$

In light of the Fock Hamiltonian, we are interested in studying the solution to the following Cauchy problem in Fock space

$$\begin{aligned} \mathbf{S}_D \psi &:= \left( \frac{1}{i} \frac{\partial}{\partial t} - \mathcal{H} \right) \psi = 0 \quad \text{with} \\ \psi(0) &= \psi_0 = e^{-\sqrt{N}\mathcal{A}(\phi_0)} e^{-\mathcal{B}(k_0)} \Omega \end{aligned} \quad (2.12)$$

which we shall formally write as

$$\psi_{\text{exact}}(t) = e^{it\mathcal{H}} e^{-\sqrt{N}\mathcal{A}(\phi_0)} e^{-\mathcal{B}(k_0)} \Omega. \quad (2.13)$$

Here  $\psi_0$  is usually referred to as a pure squeezed state, or more generally, a quasifree state.

Let  $\mathcal{M} = e^{-\sqrt{N}\mathcal{A}} e^{-\mathcal{B}}$ . Following [GM13a, GM17], we work with the reduced dynamics, which is also called the fluctuation dynamics. More

---

<sup>5</sup>In the mathematical physics literature,  $e^{\mathcal{B}}$  is called the infinite-dimensional Segal-Shale-Weil representation of the double cover of the group of symplectic matrices of integral operators. The elements of the corresponding  $C^*$ -algebra are called Bogoliubov transformations (cf. Chapter 4 of [Fol89] and Chapter 11 of [DG13]).

<sup>6</sup>We adopted the convention that  $\mathcal{H} \leq 0$ , a non-positive operator, which could make some of the signs in the paper not conventional when compare to the mathematical physics literature.

specifically, since  $\mathcal{M}$  is a unitary operator then the error of the quasifree approximation can be rewritten in terms of the reduced dynamics, that is,

$$\|E\|_{\mathcal{F}} := \|\psi_{\text{exact}}(t) - \psi_{\text{approx}}(t)\|_{\mathcal{F}} = \|\psi_{\text{red}}(t) - \Omega\|_{\mathcal{F}} \quad (2.14)$$

where

$$\psi_{\text{red}}(t) = e^{\mathcal{B}(t)} e^{\sqrt{N}\mathcal{A}(t)} e^{it\mathcal{H}} e^{-\sqrt{N}\mathcal{A}(\phi_0)} e^{-\mathcal{B}(k_0)} \Omega \quad (2.15)$$

is called the reduced Fock vector. Therefore, to study the error, it suffices to study the evolution of the reduced dynamics.

By direct computation, one can show that the evolution of  $\psi_{\text{red}}$  is determined by the equation

$$\frac{1}{i} \frac{\partial}{\partial t} \psi_{\text{red}} = \mathcal{H}_{\text{red}} \psi_{\text{red}} \quad \text{where} \quad \mathcal{H}_{\text{red}} = \frac{1}{i} (\partial_t \mathcal{M}^*) \mathcal{M} + \mathcal{M}^* \mathcal{H} \mathcal{M}. \quad (2.16)$$

$\mathcal{H}_{\text{red}}$  is called the reduced Hamiltonian. The reader should note that  $\mathcal{H}_{\text{red}}$  is a fourth degree polynomial in  $a$  and  $a^*$ . Then we see that the error  $E$  satisfies the inhomogeneous equation<sup>7</sup>

$$\left( \frac{1}{i} \frac{\partial}{\partial t} - \mathcal{H}_{\text{red}} \right) E = \mathcal{H}_{\text{red}} \Omega \quad (2.17)$$

where

$$\mathcal{H}_{\text{red}} \Omega = (X_0, X_1, X_2, X_3, X_4, 0, 0, \dots). \quad (2.18)$$

Thus, to estimate the error and show that it is small, for large  $N$ , we need to be able to control the forcing term  $\mathcal{H}_{\text{red}} \Omega$  uniformly in  $N$ . However, it has been shown in [GM13a, GM13b] that  $X_0, X_1$  and  $X_2$  are heuristically large since they are proportional to  $N, N^{\frac{1}{2}}$  and constant, respectively. On the other hand,  $X_3$  and  $X_4$  are proportional to  $N^{-\frac{1}{2}}$  and  $N^{-1}$ , respectively, making them heuristically small when  $N$  is sufficiently large.

Fortunately,  $X_0$  does not pose any serious problem. In fact, by simply modifying our quasifree approximation by a phase factor is sufficient to get rid of  $X_0$  in the forcing term (2.18), that is, if we consider the modified approximation

$$\tilde{\psi}_{\text{red}}(t) = e^{-i \int_0^t X_0(s) ds} \psi_{\text{red}}(t) \quad (2.19)$$

then we see that the modified error  $\tilde{E}$  satisfies the equation

$$\mathbf{S}_F \tilde{E} := \left( \frac{1}{i} \frac{\partial}{\partial t} - \mathcal{H}_{\text{red}} + X_0 \right) (\tilde{\psi}_{\text{red}} - \Omega) = \mathcal{H}_{\text{red}} \Omega - X_0 \Omega. \quad (2.20)$$

The explicit form of  $X_0$ , also denoted by  $X_0 := -N\mu$ , is given in Proposition 6.1 of [GM13a]. Next, since we have the two degrees of freedom  $\phi$  and  $k$  from our approximation scheme, then we can handle  $X_1$  and  $X_2$  by simply

---

<sup>7</sup>Here we made a slight abused of notation by identifying  $E$  with both  $\psi_{\text{exact}} - \psi_{\text{approx}}$  and  $\psi_{\text{red}} - \Omega$ . However, since our analysis only involves studying  $\psi_{\text{red}} - \Omega$ , the reader can safely assume  $E = \psi_{\text{red}} - \Omega$  for the remainder of the paper.

imposing the natural conditions that  $\phi_t$  and  $k_t$  must satisfy the abstract equations

$$X_1 = 0 \quad \text{and} \quad X_2 = 0. \quad (2.21)$$

These equations were first explicitly computed in Theorem 7.1 of [GM13a] (cf. appendix C of [BBCFS16]).

To compactly write down the evolution equations for  $\phi_t$  and  $k_t$ , we need to introduce some notations.

It turns out that the equations for  $k$  is most appropriately stated in terms of the following set of fields

$$\text{sh}(k) := k + \frac{1}{3!}k \circ \bar{k} \circ k + \frac{1}{5!}k \circ \bar{k} \circ k \circ \bar{k} \circ k + \dots \quad (2.22a)$$

$$\text{ch}(k) := \delta + \frac{1}{2!}\bar{k} \circ k + \frac{1}{4!}\bar{k} \circ k \circ \bar{k} \circ k + \dots \quad (2.22b)$$

where  $\circ$  indicates composition, that is, the kernel of  $k \circ l$  is given by

$$(k \circ l)(x, y) = \int dz \, k(x, z)l(z, y). \quad (2.23)$$

Adopting the convention of [GM13a], we also use the notation  $u = \text{sh}(k)$  and  $c = \text{ch}(k) = \delta + p$ .

Moreover, we define the operator kernels

$$\Lambda(t, x, y) := \phi(t, x)\phi(t, y) + \frac{1}{2N} \text{sh}(2k)(t, x, y) \quad (2.24a)$$

$$\Gamma(t, x, y) := \bar{\phi}(t, x)\phi(t, y) + \frac{1}{N} \left( \overline{\text{sh}(k)} \circ \text{sh}(k) \right) (t, x, y) \quad (2.24b)$$

which will play paramount roles in our subsequent discussion. To reconcile with some of the notations in the literature, in particular the notations of [BBCFS16], we also adopt the notation  $\sigma := u \circ c = \frac{1}{2} \text{sh}(2k)$  and  $\gamma := u \circ \bar{u} = \frac{1}{2}(\overline{\text{ch}(2k)} - \delta)$ . It is convenient to consider the trace density<sup>8</sup> of  $\Gamma$

$$\text{diag } \Gamma(t, x) = \frac{1}{N} [n_c(t, x) + \tilde{n}(t, x)] := |\phi(t, x)|^2 + \frac{1}{N} (\bar{u} \circ u)(t, x, x) \quad (2.25)$$

and define the operator kernel

$$\alpha(t, x, y) := (v_N * \text{diag } \Gamma)(t, x)\delta(x - y) + v_N(x - y)\Gamma(t, x, y) \quad (2.26a)$$

$$\begin{aligned} &= (v_N * |\phi|^2)(t, x)\delta(x - y) + v_N(x - y)\bar{\phi}(t, x)\phi(t, y) \\ &\quad + \frac{1}{N} \left\{ (v_N * \tilde{n})(t, x)\delta(x - y) + v_N(x - y)(\bar{u} \circ u)(t, x, y) \right\} \\ &=: \alpha_c(t, x, y) + N^{-1}\tilde{\alpha}(t, x, y) \end{aligned} \quad (2.26b)$$

---

<sup>8</sup>When an integral operator  $F$  is positive-definite, such as  $\Gamma$ ,  $\text{diag } F$ , denoted the restriction of  $F$  to the diagonal, is the trace density of  $F$ , that is  $\text{tr } F = \int F(x, x) dx$ . In the physics literature,  $N \text{diag } \Gamma(x)$  is called the total-number density and is often denoted by  $n(x)$ . Here we adopt the standard notation,  $n_c$  and  $\tilde{n}$  denote the condensate density and the non-condensate density, respectively, i.e.  $n(x) = n_c(x) + \tilde{n}(x)$ .



and the multiplication

$$(v_N F)(t, x, y) := v_N(x - y)F(t, x, y). \quad (2.27)$$

Let us also define the following Schrödinger-type operators

$$\mathbf{S}_\pm = \frac{1}{i} \frac{\partial}{\partial t} - \Delta_x - \Delta_y \quad \text{and} \quad \mathbf{S} = \frac{1}{i} \frac{\partial}{\partial t} - \Delta \quad (2.28)$$

where the latter is defined for either 6+1 or 3+1 dimensions, which will be clear from the context, and their corresponding operators with “potentials”: for any function  $s$  symmetric in  $(x, y)$ , i.e.  $s(x, y) = s(y, x)$ , and any function  $p$  conjugate symmetric in  $(x, y)$ , i.e.  $\bar{p}(y, x) = p(x, y)$  or  $\bar{p}^T = p$  where  $p^T(x, y) = p(y, x)$  denotes the transpose operator, we define

$$\tilde{\mathbf{S}}(s) := \mathbf{S}(s) + [\alpha^T, s]_+ \quad (2.29a)$$

$$\tilde{\mathbf{W}}(p) := \mathbf{S}_\pm(p) + [\alpha^T, p] \quad (2.29b)$$

where  $[A, B] := A \circ B - B^* \circ A^*$  is the (skew-Hermitian) commutator and  $[A, B]_+ := A \circ B + B^T \circ A^T$  is the symmetrization of the two operators  $A$  and  $B$ .

Then we have the following theorem.

**Theorem 2.1** (Theorem 7.1 of [GM13a]). *The equation  $X_1 = 0$  is equivalent to*

$$\mathbf{S}\phi_t(x_1) = - \int dy \left\{ (v_N \Lambda_t)(x_1, y) \bar{\phi}_t(y) + \frac{1}{N} \tilde{\alpha}^T(t, x_1, y) \phi_t(y) \right\}. \quad (2.30a)$$

*The equation  $X_2 = 0$  is equivalent to the pair of equations*

$$\tilde{\mathbf{S}}(\text{sh}(2k_t)) + [v_N \Lambda_t, \text{ch}(2k_t)]_+ = 0, \quad (2.30b)$$

$$\tilde{\mathbf{W}}(\overline{\text{ch}(2k_t)}) + [v_N \Lambda_t, \overline{\text{sh}(2k_t)}] = 0. \quad (2.30c)$$

By imposing the system of equations (2.30), we have reduced (2.20) to

$$\mathbf{S}_F \tilde{E} = (0, 0, 0, X_3, X_4, 0, \dots) \quad (2.31)$$

which we could apply energy method (or Strichartz) to estimate  $\tilde{E}$  in Fock space. If we write out  $X_3$  and  $X_4$  explicitly, up to symmetrization, normalization and some lower-order terms, we see that

$$X_3(t, x_1, x_2, x_3) \quad (2.32a)$$

$$\sim \int \frac{dy_1 dy_2}{\sqrt{N}} \overline{\text{ch}(k_t)}(x_1, y_1) \text{ch}(k_t)(y_2, x_2) v_N(y_1 - y_2) \phi_t(y_2) \text{sh}(k_t)(x_3, y_1)$$

and

$$X_4(t, x_1, x_2, x_3, x_4) \quad (2.32b)$$

$$\sim \int \frac{dy_1 dy_2}{N} \overline{\text{ch}(k_t)}(x_1, y_1) \text{ch}(k_t)(y_2, x_2) v_N(y_1 - y_2) \phi_t(y_2) \text{sh}(k_t)(x_3, y_1).$$

The complete list of the error terms can be found in Section 5 of [GM13b]. Hence we would like to obtain uniform in  $N$  estimates for  $\text{sh}(k)$  and  $\text{ch}(k)$ ,

which will in term show that the Fock space error tends to zero in the particle limit  $N \rightarrow \infty$ . However, due to the singular nature of the forcing term  $v_N \Lambda$  in (2.30b) and (2.30c), this poses some inconvenience when trying to estimate  $\text{sh}(2k)$  and  $\text{ch}(2k)$  uniformly in  $N$  for  $\beta \in (0, 1)$ . It was shown in [GM17] that  $k$  is best expressed in terms of auxiliary fields  $\Gamma$  and  $\Lambda$ , which are functions defined by (2.24) in terms of  $k$  and  $\phi$ . In fact, it was shown that  $\Gamma$  and  $\Lambda$  can be viewed as the “generalized” (Fock space) marginal density matrices for the pure quasifree state  $\psi_0$  in (2.12).

**2.2. Hartree-Fock-Bogoliubov System.** To write down the alternative system of nonlinear equations to (2.30), we define the kernel of the marginal density matrices  $\mathcal{L}_{m,n}$  associated with the state  $\psi_{\text{approx}}$ , which is sometimes called the  $(m+n)$ -point correlation function, as follows

$$\begin{aligned} \mathcal{L}_{m,n}(t, y_1, \dots, y_m; x_1, \dots, x_n) \\ &:= \frac{1}{N^{(m+n)/2}} \langle a_{y_1} \cdots a_{y_m} \psi_{\text{approx}}, a_{x_1} \cdots a_{x_n} \psi_{\text{approx}} \rangle \\ &= \frac{1}{N^{(m+n)/2}} \langle \mathcal{M} \Omega, \mathcal{P}_{m,n} \mathcal{M} \Omega \rangle \end{aligned} \quad (2.33)$$

where  $\mathcal{P}_{m,n} = a_{y_1}^* \cdots a_{y_m}^* a_{x_1} \cdots a_{x_n}$ . It is clear from the definition that  $\mathcal{L}_{m,n}^* = \mathcal{L}_{n,m}$ . It is also straightforward to write down the evolution equations for  $\mathcal{L}_{m,n}$  using (2.16), that is,

$$\frac{1}{i} \frac{\partial}{\partial t} (\mathcal{M}^* \mathcal{P} \mathcal{M}) = [\mathcal{H}_{\text{red}}, \mathcal{M}^* \mathcal{P} \mathcal{M}] + \mathcal{M}^* [\mathcal{P}, \mathcal{H}] \mathcal{M} \quad (2.34a)$$

which means

$$\frac{1}{i} \frac{\partial}{\partial t} \mathcal{L} = \frac{1}{N^{(m+n)/2}} (\langle \Omega, [\mathcal{H}_{\text{red}}, \mathcal{M}^* \mathcal{P} \mathcal{M}] \Omega \rangle + \langle \Omega, \mathcal{M}^* [\mathcal{P}, \mathcal{H}] \mathcal{M} \Omega \rangle). \quad (2.34b)$$

Notice the set of equations (2.34b) for different pairs  $(m, n)$  forms an infinite linear hierarchy of coupled equations where the evolution of each point correlation function depends on other higher-order point correlation functions. In the literature, (2.34b) is refer to as the von Neuman-Landau equation.

Since  $\psi_{\text{approx}}$  is a quasifree state, then it can be shown that any  $(n+m)$ -point correlation functions for  $n+m \geq 3$  can be expressed in terms of the one-particle or two-particle Fock marginal densities,  $\mathcal{L}_{0,1}, \mathcal{L}_{1,1}$  and  $\mathcal{L}_{0,2}$ . Hence it suffices for us to study the evolution equations for  $\mathcal{L}_{0,1}, \mathcal{L}_{1,1}$  and  $\mathcal{L}_{0,2}$ . In fact, it was shown in [GM17] that  $\mathcal{L}_{0,1} = \phi, \mathcal{L}_{1,1} = \Gamma, \mathcal{L}_{0,2} = \Lambda$  and that their evolution equations form a closed system of nonlinear coupled PDEs given by

$$\begin{aligned} \left\{ \frac{1}{i} \frac{\partial}{\partial t} - \Delta_{x_1} \right\} \phi(x_1) &= - \int dy \{ v_N(x_1 - y) \Gamma(y, y) \} \phi(x_1) \\ &\quad - \int dy \{ v_N(x_1 - y) (\Gamma(y, x_1) - \bar{\phi}(y) \phi(x_1)) \phi(y) \} \\ &\quad - \int dy \{ v_N(x_1 - y) (\Lambda(x_1, y) - \phi(y) \phi(x_1)) \bar{\phi}(y) \} \end{aligned} \quad (2.35a)$$

$$\left\{ \frac{1}{i} \frac{\partial}{\partial t} - \Delta_{x_1} + \Delta_{x_2} \right\} \bar{\Gamma}(x_1, x_2) \quad (2.35b)$$

$$\begin{aligned} &= - \int dy \{ (v_N(x_1 - y) - v_N(x_2 - y)) \Lambda(x_1, y) \bar{\Lambda}(y, x_2) \} \\ &\quad - \int dy \{ (v_N(x_1 - y) - v_N(x_2 - y)) (\bar{\Gamma}(x_1, y) \bar{\Gamma}(y, x_2) + \bar{\Gamma}(y, y) \bar{\Gamma}(x_1, x_2)) \} \\ &\quad + 2 \int dy \{ (v_N(x_1 - y) - v_N(x_2 - y)) |\phi(y)|^2 \phi(x_1) \bar{\phi}(x_2) \} \end{aligned}$$

$$\left\{ \frac{1}{i} \frac{\partial}{\partial t} - \Delta_{x_1} - \Delta_{x_2} + \frac{1}{N} v_N(x_1 - x_2) \right\} \Lambda(x_1, x_2) \quad (2.35c)$$

$$\begin{aligned} &= - \int dy \{ (v_N(x_1 - y) + v_N(x_2 - y)) \Gamma(y, y) \Lambda(x_1, x_2) \} \\ &\quad - \int dy \{ (v_N(x_1 - y) + v_N(x_2 - y)) (\Lambda(x_1, y) \Gamma(y, x_2) + \bar{\Gamma}(x_1, y) \Lambda(y, x_2)) \} \\ &\quad + 2 \int dy \{ (v_N(x_1 - y) + v_N(x_2 - y)) |\phi(y)|^2 \phi(x_1) \phi(x_2) \}. \end{aligned}$$

Notice we have suppressed the time dependence to compactify the notation. We refer to (2.35) as the time-dependent Hartree-Fock-Bogoliubov (TDHFB) system.

*Remark 2.2.* In the literature, the name TDHFB equations typically refers to the coupled system (2.30) where the pair interaction is given by the contact interaction potential  $v(x) = g_0 \delta(x)$  for some coupling constant  $g_0 \geq 0$  (In the context of diluted gases, one has  $g_0 = 4\pi a$  where  $a$  is some scattering length). Furthermore, it is standard to express the equation for  $\phi$  in terms of  $\Phi := \sqrt{N} \phi$  then (2.30a) comes

$$\mathbf{S}\Phi(x_1) = -g_0 [n_c(x_1) + 2\tilde{n}(x_1)] \Phi(x_1) - g_0 \tilde{m}(x_1) \bar{\Phi}(x_1) \quad (2.36a)$$

where  $\tilde{m}(x_1) := \sigma(x_1, x_1) = N\Lambda(x_1, x_1) - |\Phi(x_1)|^2$ . The other two equations (2.30b) and (2.30c) can be neatly rewritten in terms of a matrix equation of  $\gamma$  and  $\sigma$  as follows

$$\frac{1}{i} \frac{\partial}{\partial t} R = [R, \mathfrak{S}] \quad (2.36b)$$

where  $R$  is the  $2 \times 2$  matrix of operators with matrix kernel

$$R(x, y) = \begin{pmatrix} \gamma(x, y) & \sigma(x, y) \\ \bar{\sigma}(x, y) & \delta(x - y) + \bar{\gamma}(x, y) \end{pmatrix} \quad (2.37a)$$

and  $\mathfrak{S}$  has the matrix kernel

$$\mathfrak{S}(x, y) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} G(x, y) & M(x, y) \\ \bar{M}(x, y) & \bar{G}(x, y) \end{pmatrix} \quad (2.37b)$$

with

$$G(x, y) := -\Delta_x \delta(x - y) + 2g_0[n_c(x) + \tilde{n}(x)]\delta(x - y) \quad (2.38a)$$

$$M(x, y) := g_0[\Phi(x)^2 + \tilde{m}(x)]\delta(x - y). \quad (2.38b)$$

Moreover, the physical interpretation of  $(\Phi, \gamma, \sigma)$  (or equivalently,  $(\phi, \Gamma, \Lambda)$ ) is as follows: The one-particle wave function  $\Phi$  is called the condensate wave function which describes the behavior of the Bose-Einstein condensate (BEC). On the other hand,  $\gamma$  and  $\sigma$  describe the dynamics of sound waves, or quasiparticles, in the quasifree approximation. In particular,  $\tilde{n}(x) = \gamma(x, x)$  determines the density of the “thermal cloud” of atoms, i.e. the excitation density of the Bose gas. In the physics literature,  $\tilde{n}(x) = \gamma(x, x)$  is called the non-condensate density and  $\tilde{m}(x) = \sigma(x, x)$  is called the anomalous density.

*Remark 2.3.* It has been shown in [BBCFS16] that the evolution equations for the higher Fock marginal densities can be written in terms of an effective nonlinear quadratic Hamiltonian parametrized by  $(\phi, \Gamma, \Lambda)$  called the Hartree-Fock-Bogoliubov (HFB) Hamiltonian, denoted by  $\mathcal{H}_{\text{HFB}}$ ; more precisely, we have the evolution equation

$$\frac{1}{i} \frac{\partial}{\partial t} \mathcal{L}_{m,n} = \langle \mathcal{M}\Omega, [\mathcal{H}_{\text{HFB}}, \mathcal{P}_{m,n}] \mathcal{M}\Omega \rangle. \quad (2.39)$$

The HFB Hamiltonian is given explicitly by

$$\mathcal{H}_{\text{HFB}} = -N\mu + \sqrt{N}\mathcal{H}_1 + \mathcal{H}_2 \quad (2.40)$$

with

$$\mathcal{H}_1 = 2 \int dx \{ (v_N * |\phi|^2) \phi(x) a_x^* + (v_N * |\phi|^2) \bar{\phi}(x) a_x \} \quad (2.41a)$$

$$\mathcal{H}_2 = -\frac{1}{2} \int dx dy \{ 2g^T(x, y) a_x^* a_y + m(x, y) a_x^* a_y^* + \bar{m}(x, y) a_x a_y \} \quad (2.41b)$$

where

$$g(x, y) := -\Delta_x \delta(x - y) + (v_N * \text{diag } \Gamma)(x) \delta(x - y) + (v_N \Gamma)(x, y) \quad (2.42a)$$

$$m(x, y) := (v_N \Lambda)(x, y) = v_N(x - y) \Lambda(x, y). \quad (2.42b)$$

Also note, by (2.34b) and (2.39), we have the relation

$$\mathcal{H} + \mathcal{H}_{\text{HFB}} = \mathcal{M} \mathcal{H}_{\text{red}} \mathcal{M}^*. \quad (2.43)$$

Since we are interested in studying the uniform in  $N$  error bound of the quasifree approximation, it is then instructive to take the formal large particle limit,  $N \rightarrow \infty$ , to obtain the following equations

$$\mathbf{S}\phi(x_1) = -2g_0\{\Gamma(x_1, x_1) - |\phi(x_1)|^2\}\phi(x_1) - g_0\Lambda(x_1, x_1)\bar{\phi}(x_1) \quad (2.44a)$$

$$\begin{aligned} \mathbf{S}_{\pm}\bar{\Gamma}(x_1, x_2) &= -g_0\{\Lambda(x_1, x_1)\bar{\Lambda}(x_1, x_2) - \Lambda(x_1, x_2)\bar{\Lambda}(x_2, x_2)\} \\ &\quad - 2g_0\{\bar{\Gamma}(x_1, x_1) - \bar{\Gamma}(x_2, x_2)\}\bar{\Gamma}(x_1, x_2) \\ &\quad + 2g_0\{|\phi(x_1)|^2 - |\phi(x_2)|^2\}\phi(x_1)\bar{\phi}(x_2) \end{aligned} \quad (2.44b)$$

$$\begin{aligned}
\mathbf{S}\Lambda(x_1, x_2) = & -g_0\{\Lambda(x_1, x_1)\Gamma(x_1, x_2) + \Lambda(x_2, x_2)\bar{\Gamma}(x_1, x_2)\} \\
& - 2g_0\{\Gamma(x_1, x_1) + \Gamma(x_2, x_2)\}\Lambda(x_1, x_2) \\
& + 2g_0\{|\phi(x_1)|^2 + |\phi(x_2)|^2\}\phi(x_1)\phi(x_2)
\end{aligned} \tag{2.44c}$$

for some constant  $g_0$ . In many ways, system (2.44) serves as a guiding light in our studies of the uniform in  $N$  estimates for the TDHFB system. In particular, as remarked in [GM19], the solution to the system (2.44) is invariant under the scaling  $\lambda\phi(\lambda^2 t, \lambda x_1, \lambda x_2), \lambda^2\Gamma(\lambda^2 t, \lambda x_1, \lambda x_2), \lambda^2\Lambda(\lambda^2 t, \lambda x_1, \lambda x_2)$ , which makes the limiting system energy critical. However, the reader should also note that the scaling argument does not detect the collapse to the diagonal  $x_1 = x_2$  involved in these equations. More precisely, by replacing  $\Lambda(x_1, x_1)\bar{\Lambda}(x_1, x_2)$  with  $|\Lambda(x_1, x_2)|^2$ , the system would still scale the same way, but it is well known that going from  $|\Lambda(x_1, x_2)|^2$  to  $|\Lambda(x_1, x_1)|^2$  there is a natural loss of regularity; see for instance the standard sharp trace theorem or Lemma 5.1 in [GM17] for the collapsing estimates of the linear  $\Lambda$  and  $\Gamma$  equations. Thus, it is arguable that the system (2.44) is actually energy-supercritical, which illustrates the formidability of obtaining uniform in  $N$  global well-posedness and estimates for the TDHFB system (2.35) in energy space. Hence it seems desirable to first consider the well-posedness of  $\Gamma$  and  $\Lambda$  in  $H^{1+\varepsilon}$  Sobolev spaces for  $\varepsilon > 0$ . But even so, the situation is not completely obvious since the conserved energy of the system scales uniform in  $N$  like  $H^1$  for  $\Lambda$  which poses the classical questions regarding the growth of  $H^{1+\varepsilon}$  Sobolev norm of  $\Lambda$ . This question regarding the growth of the Sobolev norm of  $\Lambda$  was recently studied by the authors in a joint work with M. Grillakis and M. Machedon in [CGMZ20].

A more pressing question one might have is whether the TDHFB system is even uniform in  $N$  locally well-posed. Indeed, the uniform in  $N$  local well-posedness of the TDHFB system was proven in [GM19] for  $0 < \beta < 1$ . Let us summarize their result as follows.

We begin by defining the space of initial data. For every  $\alpha > 0$ , we define the space

$$\mathcal{X}^\alpha = \{(\phi, \Gamma, \Lambda) \in H^\alpha \times H_{\text{Herm}}^\alpha \times H_{\text{Sym}}^\alpha\} \tag{2.45}$$

with  $H^\alpha$  being the Sobolev space  $H^\alpha(\mathbb{R}^3)$ ,  $H_{\text{Herm}}^\alpha$  the Sobolev space  $H^\alpha(\mathbb{R}^3 \times \mathbb{R}^3)$  restricted to functions  $\Gamma$  such that  $\Gamma(x, y) = \overline{\Gamma(y, x)}$ , and  $H_{\text{Sym}}^\alpha(\mathbb{R}^3 \times \mathbb{R}^3)$  restricted to functions  $\Lambda$  such that  $\Lambda(x, y) = \Lambda(y, x)$ . Moreover,  $\mathcal{X}^\alpha$  is endowed with the norm

$$\begin{aligned}
\|(\phi, \Gamma, \Lambda)\|_{\mathcal{X}^\alpha} := & \|\langle \nabla \rangle^\alpha \phi\|_{L^2(\mathbb{R}^3)} + \left\| (\langle \nabla_x \rangle^2 \otimes 1 + 1 \otimes \langle \nabla_y \rangle^2)^{\frac{\alpha}{2}} \Gamma \right\|_{L^2(\mathbb{R}^3 \times \mathbb{R}^3)} \\
& + \left\| (\langle \nabla_x \rangle^2 \otimes 1 + 1 \otimes \langle \nabla_y \rangle^2)^{\frac{\alpha}{2}} \Lambda \right\|_{L^2(\mathbb{R}^3 \times \mathbb{R}^3)}
\end{aligned} \tag{2.46}$$

where  $\langle D \rangle = \sqrt{1 + |D|^2}$  is the standard bracket notation.

**Theorem 2.4** (Grillakis & Machedon '18). *Assume  $v$  is a Schwartz function with  $\hat{v}$  supported in the unit ball, such that  $|\hat{v}| \leq \hat{w}$  with  $w$  a Schwartz*

function. Let  $0 < \beta < 1$ . Fix  $\alpha > \frac{1}{2}$  so that  $2\alpha\beta < 1$  and choose  $R > 0$ . Then there exists  $T = T(\beta, R)$ , independent of  $N$ , and a corresponding spacetime function space  $\mathcal{X}_T$ , depending only on  $T$  and  $\alpha$ , such that for any given initial data

$$(\phi_0, \Gamma_0, \Lambda_0) \in \{(\phi, \Gamma, \Lambda) \in \mathcal{X}^\alpha \mid \|(\phi, \Gamma, \Lambda)\|_{\mathcal{X}^\alpha} < R\},$$

there exists a unique maximal mild solution to the time-dependent HFB system (2.35), with initial data  $(\phi_0, \Gamma_0, \Lambda_0)$ , satisfying

$$(\phi_t, \Gamma_t, \Lambda_t) \in C([0, T] \rightarrow \mathcal{X}^\alpha) \cap \mathcal{X}_T. \quad (2.47)$$

Theorem 2.4 is an improvement of their previous local well-posedness result in [GM17] for the case  $0 < \beta < \frac{2}{3}$ . The analysis of the case  $0 < \beta < 1$  is substantially more involved than the situation in [GM17]. However, the result of [GM19] does not cover the Fock space approximation which was addressed in [GM17], at least for small time. Let us also summarize their earlier findings.

**Theorem 2.5** (Grillakis & Machedon '17). *Assume  $v$  as in Theorem 2.4 and  $0 < \beta < \frac{2}{3}$ . Choose  $\alpha > \frac{1}{2}$  such that  $2\alpha\beta < 1$ . Then (2.35) is uniform in  $N$  locally well-posed in  $\mathcal{X}^\alpha$ . Moreover, suppose  $(\phi_t, \Gamma_t, \Lambda_t)$  is a solution to the HFB system with some smooth initial conditions  $(\phi_0, \Gamma_0, \Lambda_0)$  satisfying the following regularity condition uniformly in  $N$ : for  $0 \leq i \leq 1, 0 \leq j \leq 2$*

$$\begin{aligned} & \left\| \langle \nabla_x \rangle^\alpha \partial_t^i \nabla_x^j \phi(t, \cdot) \Big|_{t=0} \right\|_{L^2(dx)} \lesssim 1 \\ & \left\| \langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha \partial_t^i \nabla_{x+y}^j \Gamma(t, \cdot) \Big|_{t=0} \right\|_{L^2(dxdy)} \lesssim 1 \\ & \left\| \langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha \partial_t^i \nabla_{x+y}^j \Lambda(t, \cdot) \Big|_{t=0} \right\|_{L^2(dxdy)} \lesssim 1 \\ & \left\| \nabla_{x+y}^j \text{sh}(2k_0)(x, y) \right\|_{L^2(dxdy)} \lesssim 1. \end{aligned} \quad (2.48)$$

Then there exist  $T_0 > 0$  ( $T_0 \sim 1$ ) independent of  $N$ , a phase function  $\chi(t)$ , depending on  $N$ , and some constant  $C = C(T_0, \alpha, \beta)$  such that we have the Fock space estimate

$$\left\| e^{it\mathcal{H}} e^{-\sqrt{N}\mathcal{A}(\phi_0)} e^{-\mathcal{B}(k_0)} \Omega - e^{i\chi(t)} e^{-\sqrt{N}(\phi_t)} e^{-\mathcal{B}(k_t)} \Omega \right\|_{\mathcal{F}} \leq \frac{C}{N^{\frac{1}{6}}} \quad (2.49)$$

for all  $0 \leq t \leq T_0$ .

*Remark 2.6.* A crucial point to note in Theorem 2.5 is the dependence on the time derivative on the initial data (2.48). As pointed out in Remark 2.6 of [GM17], the assumption of time derivative dependence of the initial data rules out the case of coherent states, i.e.  $k(0, x, y) = 0$ .

**2.3. Main Results.** Since the well-posedness theory of (2.35) is stated in Strichartz-type space, let us briefly review some basic notations. The space-time function space endowed with the mixed norm

$$\|F\|_{L^q(dt)L^r(dx)L^s(dy)} := \left( \int_{-\infty}^{\infty} dt \left( \int_{\mathbb{R}^d} dx \|F(t, x, \cdot)\|_{L^s(\mathbb{R}^d)}^r \right)^{\frac{q}{r}} \right)^{\frac{1}{q}}$$

where the triplet  $(q, r, s)$  satisfies the admissible condition  $\frac{2}{q} + \frac{d}{r} + \frac{d}{s} = d$  for  $2 \leq q, r, s \leq \infty$  is called a Strichartz space.

Fix  $T > 0$  and let  $c(t)$  be the characteristic function<sup>9</sup> on the interval  $[0, T]$ . For our problem, we consider the function space  $\mathcal{X}_T$  for the triplet of functions  $X = (\phi, \Gamma, \Lambda)$  equipped with the norm that is a sum of the following norms

$$\mathbf{N}_T(\phi) := \|\langle \nabla_x \rangle^\alpha c(t) \phi\|_{L^\infty(dt)L^2(dx)} + \|\langle \nabla_x \rangle^\alpha c(t) \phi\|_{L^2(dt)L^6(dx)} \quad (2.50a)$$

$$\begin{aligned} \mathbf{N}_T(\Gamma) := & \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Gamma\|_{L^\infty(dt)L^2(dx dy)} \\ & + \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Gamma\|_{L^2(dt)L^6(dx)L^2(dy)} \\ & + \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Gamma\|_{L^2(dt)L^6(dy)L^2(dx)} \\ & + \sup_z \|\langle \nabla_x \rangle^{\alpha-\frac{1}{2}} |\nabla_x|^{\frac{1}{2}} c(t) \Gamma(t, x, x+z)\|_{L^2(dtdx)} \end{aligned} \quad (2.50b)$$

and

$$\begin{aligned} \mathbf{N}_T(\Lambda) := & \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Lambda\|_{L^\infty(dt)L^2(dx dy)} \\ & + \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Lambda\|_{L^2(dt)L^6(dx)L^2(dy)} \\ & + \|\langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha c(t) \Lambda\|_{L^2(dt)L^6(dy)L^2(dx)} \\ & + \sup_z \|\langle \nabla_x \rangle^\alpha c(t) \Lambda(t, x, x+z)\|_{L^2(dtdx)} \\ & + \sup_z \|\partial_t^{\frac{1}{4}} [c(t) \Lambda(t, x, x+w)]\|_{L^2(dtdx)}. \end{aligned} \quad (2.50c)$$

Moreover, let us denote the space of triplets  $X_t = (\phi_t, \Gamma_t, \Lambda_t)$  where the above norms are finite for any  $0 \leq T < \infty$  by  $\mathcal{X}_{\infty, \text{loc}}$ . Here, we also use

$$\mathbf{N}_T(X) := \mathbf{N}_T(\phi) + \mathbf{N}_T(\Gamma) + \mathbf{N}_T(\Lambda). \quad (2.51)$$

Let us state the first main result of our paper.

**Theorem 2.7.** *Assume  $v$  is a non-negative Schwartz function satisfying the condition that  $|\hat{v}| \leq \hat{w}$  for some Schwartz function  $w$ . Let  $0 < \beta < 1$ . Fix  $\alpha > \frac{1}{2}$  so that  $2\alpha\beta < 1$  and choose  $R > 0$ . Then for any*

$$(\phi_0, \Gamma_0, \Lambda_0) \in \{(\phi, \Gamma, \Lambda) \in \mathcal{X}^\alpha \mid \|(\phi, \Gamma, \Lambda)\|_{\mathcal{X}^\alpha} < R\},$$

*the corresponding local solution to the TDHFB system (2.35) given by Theorem 2.4 extends globally with  $X_t = (\phi_t, \Gamma_t, \Lambda_t) \in C([0, \infty) \rightarrow \mathcal{X}^\alpha) \cap \mathcal{X}_{\infty, \text{loc}}$ .*

---

<sup>9</sup>The reader should be warned that  $c$  has already been used to denote  $\text{ch}(k)$ , but we seldom use it. However, it should be clear from the context which function we are referring to. Nevertheless, we will make explicit the distinction when ambiguity arises.

Moreover, there exist constants  $C > 0$ , depending only on the initial data, such that the following uniform in  $N$  a-priori estimate

$$\mathbf{N}_T(X) \leq CT, \quad (2.52)$$

holds for all  $T > 0$ .

In view of Theorem 2.7, we can obtain a global-in-time Fock space estimate for the error of our quasifree approximation.

**Theorem 2.8.** *Assume  $v$  as in Theorem 2.7. Let  $0 < \beta < 1$ . Fix  $\alpha > \frac{1}{2}$  so that  $2\alpha\beta < 1$ . Let  $\phi, k$  be solutions to (2.35) with smooth initial conditions  $\phi_0, k_0$  satisfying the following regularity conditions: for  $0 \leq j \leq 2$*

$$\begin{aligned} & \| \langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha \nabla_{x+y}^j \Gamma_0 \|_{L^2(dxdy)} + \| \langle \nabla_x \rangle^\alpha \langle \nabla_y \rangle^\alpha \nabla_{x+y}^j \Lambda_0 \|_{L^2(dxdy)} \\ & + \| \langle \nabla_x \rangle^\alpha \nabla_x^j \phi_0 \|_{L^2(dx)} + \| \nabla_{x+y}^j \text{sh}(2k_0)(\cdot, \cdot) \|_{L^2(dxdy)} \lesssim 1. \end{aligned} \quad (2.53)$$

Then for some  $\varepsilon' > 0$ , there exists a real phase function  $\chi_0(t)$  depending on  $N$  and constants  $C = C(\beta, \varepsilon), \kappa = \kappa(\varepsilon) > 0$  such that

$$\| \psi_{\text{exact}}(t) - \tilde{\psi}_{\text{approx}}(t) \|_{\mathcal{F}} \leq \frac{C \exp(\kappa t)}{N^{\frac{1-\beta}{2}}} \quad (2.54)$$

for all  $t > 0$ .

*Remark 2.9.* Theorem 2.8 should be compared with Theorem 1.1 of [BCS17], which also provides a global-in-time Fock space for the error between the exact evolution and the effective evolution. In fact, our exponent for  $N$  is consistent with their exponent, which we believe to be sharp for the Fock space estimate. However, the nature of the two results are very different. In their work, the approximation scheme is given by

$$\psi_{\text{approx}}(t) = e^{i\chi_0(t)} e^{-\sqrt{N}\mathcal{A}(\phi_t)} e^{-\mathcal{B}(k_N(t))} U_{2,N}(t) \Omega \quad (2.55)$$

where the pair excitation function  $k_N(t, x, y)$  is given explicitly and  $U_{2,N}(t)$  is the dynamics associated the a quadratic generator which depends on  $k_N$ . Their form of  $k_N$  is inspired by the fact that when  $\beta = 1$  the ground state of (1.1) exhibits a short-range correlation structure, which means the ground state cannot be a simple tensor product. However, it is also known that for  $\beta \in (0, 1)$  the ground state factorizes in the large particle limit (cf. Appendix B in [ESY10]), which is not capture by (2.55).

In contrast, Theorem 2.8 is much more flexible. It lets us start with an uncorrelated system and study the dynamical formation of correlation structures in the system. Furthermore, another feature of our result is that by employing techniques from dispersive PDE theory we were able to get a better growth rate in time for the error term in comparison to the standard double exponential growth.



## 3. GLOBAL ESTIMATES FOR THE TIME-DEPENDENT HFB EQUATIONS

In this section we prove, for a sufficiently small  $\varepsilon > 0$ , the following estimates

$$\| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda(t) \|_{L^2(dxdy)} \leq C \quad (3.1a)$$

$$\| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma(t) \|_{L^2(dxdy)} \leq C \quad (3.1b)$$

$$\| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \phi(t) \|_{L^2(dx)} \leq C \quad (3.1c)$$

hold uniformly in  $N$  for any fixed time  $t > 0$ . The proof of estimates (3.1a)-(3.1c) relies on the conservation laws established in [GM13a]. For the reader's convenience, we restate the conservation laws for the time-dependent HFB system in the following proposition. Let us recall the total particle number and energy, which we denoted by  $\mathcal{N}$  and  $\mathcal{E}$  respectively, can be evaluated explicitly as follows:

$$\mathcal{N} = N \cdot \left\{ \int dx |\phi(x)|^2 + \frac{1}{N} \int dxdy |u(x, y)|^2 \right\} \quad (3.2a)$$

and

$$\begin{aligned} \mathcal{E} = N \cdot & \left\{ \int dx |\nabla \phi(x)|^2 + \frac{1}{2N} \int dxdy |\nabla_{x,y} u(x, y)|^2 \right. \\ & + \frac{1}{2N} \int dxdydz v_N(x-y) |\phi(x)u(y, z) + \phi(y)u(x, z)|^2 \\ & \left. + \frac{1}{4} \int dxdy v_N(x-y) \{ 2|\Lambda(x, y)|^2 + |\Gamma(x, y)|^2 + \Gamma(x, x)\Gamma(y, y) \} \right\}. \end{aligned} \quad (3.2b)$$

For the sake of compactness of notation, we have suppressed the dependence on the time variable since it only plays a passive role in our studies of the equations in this section.

**Proposition 3.1** (Conservation Quantities). *Suppose  $(\phi(t), \Gamma(t), \Lambda(t))$  is a smooth solution to (2.35) with  $v \in L^1(\mathbb{R}) \cap C^\infty(\mathbb{R})$ . Then the total-particle number and energy for the system are conserved.*

*Remark 3.2.* The reader should be aware of the fact that we are assuming that the energy per particle is constant and independent of  $N$ . More precisely, we make the assumption that  $\mathcal{N}$  and  $\mathcal{E}$  are proportional to  $N$  for some fixed  $N$ . In fact, we have that  $\mathcal{N} = N$  and  $\mathcal{E} \sim N$  or, equivalently,  $N^{-1}\mathcal{E} \sim 1$ .

As an immediate corollary of the conservation quantities, we prove estimate (3.1b) and (3.1c).

**Corollary 3.3.** *Let  $\phi(t)$  and  $\Gamma(t)$  be smooth solutions to the time-dependent HFB equations. Then, for any  $0 < \varepsilon \leq \frac{1}{2}$ , we have the estimates*

$$\begin{aligned} \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma(t) \|_{L^2(dxdy)} &\lesssim 1 \\ \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \phi(t) \|_{L^2(dx)} &\lesssim 1 \end{aligned}$$

which hold uniformly in  $N$  and independent of  $t$ .

*Proof.* It suffices to prove estimate (3.1b) since the prove of (3.1c) is similar. By Proposition 3.1 and Cauchy-Schwarz inequality, we obtain the estimate

$$\begin{aligned} \|\Gamma(t)\|_{L^2(dxdy)} &\leq \|\phi_t\|_{L^2(dx)}^2 + N^{-1} \|\overline{u}_t \circ u_t\|_{L^2(dxdy)} \\ &\leq \|\phi_t\|_{L^2(dx)}^2 + N^{-1} \|u_t\|_{L^2(dxdy)}^2 = 1 \end{aligned} \quad (3.3a)$$

uniformly in  $t$  and independent of  $N$ . Likewise, we see that

$$\|\nabla_x \nabla_y \Gamma(t)\|_{L^2(dxdy)} \leq \|\nabla_x \phi_t\|_{L^2}^2 + N^{-1} \|\nabla_x u_t\|_{L^2}^2 \lesssim 1. \quad (3.3b)$$

Hence interpolating (3.3a) and (3.3b) yields the desired result.  $\square$

In the remainder of this section, we shall prove that estimate (3.1a) holds. To this end, let us begin by making the observation that proving estimate (3.1a) is equivalent to establishing the estimate

$$N^{-1} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \text{sh}(2k_t)\|_{L^2(dxdy)} \lesssim C \quad (3.4)$$

for some sufficiently small  $\varepsilon > 0$ . Furthermore, to aid us in proving estimate (3.4), we apply the operator identity

$$\text{sh}(2k) = 2u \circ c = 2u + 2u \circ p \quad (3.5)$$

and the triangle inequality to obtain a preliminary estimate

$$\begin{aligned} &\|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \text{sh}(2k_t)\|_{L^2(dxdy)} \\ &\lesssim \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} u_t\|_{L^2(dxdy)} + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} u_t \circ p_t\|_{L^2(dxdy)} \\ &=: I_1(t) + I_2(t). \end{aligned}$$

Hence it remains to show that  $N^{-1} I_i(t) \lesssim C$  for  $i = 1, 2$ .

To estimate  $I_2(t)$ , we use the following lemma

**Lemma 3.4.** *We have the following estimates*

$$N^{-1} \left\| |\nabla_x|^{\frac{1}{2}} |\nabla_y|^{\frac{1}{2}} u_t \circ p_t \right\|_{L^2(dxdy)} \lesssim 1, \quad (3.6a)$$

$$N^{-1} \|\nabla_x u_t \circ \nabla_y p_t\|_{L^2(dxdy)} \lesssim 1 \quad (3.6b)$$

where both are independent of time  $t$ . In particular, by interpolating estimates (3.6a) and (3.6b), we obtain the estimate

$$N^{-1} \left\| |\nabla_x|^{\frac{1}{2}+\varepsilon} |\nabla_y|^{\frac{1}{2}+\varepsilon} u_t \circ p_t \right\|_{L^2(dxdy)} \lesssim 1 \quad (3.7)$$

for any  $0 \leq \varepsilon \leq \frac{1}{2}$ .

*Proof.* Using Plancherel identity and Cauchy-Schwarz inequality, we establish the estimate

$$\begin{aligned} &\left\| |\nabla_x|^{\frac{1}{2}} |\nabla_y|^{\frac{1}{2}} u_t \circ p_t \right\|_{L^2(dxdy)} \\ &\lesssim \|\nabla_x u_t\|_{L^2(dxdy)} \|p_t\|_{L^2(dxdy)} + \|u_t\|_{L^2(dxdy)} \|\nabla_y p_t\|_{L^2(dxdy)}. \end{aligned} \quad (3.8)$$

Next, taking derivatives of the kernel of the operator identity

$$\overline{u} \circ u = p \circ p + 2p \quad (3.9)$$

yields the operator identity

$$\overline{\nabla_x u} \circ \nabla_y u = \nabla_x p \circ \nabla_y p + 2\nabla_x \nabla_y p.$$

In particular, we have that

$$\|\nabla_x u\|_{L^2(dxdy)}^2 = \|\nabla_x p \circ \nabla_y p + 2\nabla_x \nabla_y p\|_{\text{tr}} \geq \|\nabla_x p\|_{L^2(dxdy)}^2$$

since both  $\nabla_x \nabla_y (p \circ p + 2p)$  and  $2\nabla_x \nabla_y p$  are positive trace class operators. Hence combining estimate (3.8) with the conservation laws, we obtain the estimate

$$N^{-1} \left\| |\nabla_x|^{\frac{1}{2}} |\nabla_y|^{\frac{1}{2}} u_t \circ p_t \right\|_{L^2(dxdy)} \lesssim N^{-1} \|\nabla_x u_t\|_{L^2(dxdy)} \|u_t\|_{L^2(dxdy)} \lesssim 1.$$

Likewise, we have shown that

$$N^{-1} \|\nabla_x u_t \circ \nabla_y p_t\|_{L^2(dxdy)} \lesssim N^{-1} \|\nabla_x u_t\|_{L^2}^2 \lesssim 1.$$

□

Next, to estimate  $I_1(t)$ , we state a useful global estimate for the  $\Lambda$ , which is established in our recent paper [CGMZ20]. See Theorem 1.1 and Theorem 1.2 in [CGMZ20].

Recall the equation for  $\Lambda(t)$  is given by

$$\begin{aligned} (\mathbf{S} + V) \Lambda = & - (v_N \Lambda) \circ \Gamma - \bar{\Gamma} \circ (v_N \Lambda) - (v_N \bar{\Gamma}) \circ \Lambda - \Lambda \circ (v_N \Gamma) \\ & + 2(v_N * |\phi|^2)(x) \phi(x) \phi(y) + 2(v_N * |\phi|^2)(y) \phi(y) \phi(x) =: F \end{aligned} \quad (3.10)$$

where  $v_N \Lambda = v_N(x - y) \Lambda(x, y)$  and

$$V := \frac{1}{N} v_N + (v_N * \text{diag } \Gamma)(x) + (v_N * \text{diag } \Gamma)(y).$$

For the convenience of the reader, let us also recall the theorem.

**Lemma 3.5** (Theorem 1.2 of [CGMZ20]). *Let  $\Lambda(t)$  be a solution to the time-dependent HFB equations. Then we have the following time-independent global estimate*

$$\|\nabla_x \nabla_y \Lambda(t)\|_{L^2(dxdy)} \lesssim \|\nabla_x \nabla_y \Lambda_0\|_{L^2(dxdy)} + N^{c_0}, \quad (3.11)$$

where  $c_0$  is a positive constant.

*Remark 3.6.* The positive power  $c_0$  is chosen to be  $\frac{11}{2}$  in [CGMZ20], which is not optimal. One may apply other methods to improve (lower) this power. But we will not explore this point in this paper since a positive power is sufficient for our problem. See Remark 3.9.

*Remark 3.7.* The proof of Lemma 3.5 tightly depends on a coordinate-flexible Strichartz estimate and a bootstrap argument. (See Theorem 2.3, 2.4 and 2.5 in [CGMZ20].) Other ingredients include an interaction Morawetz argument and a delicate analysis of the nonlinearity. We leave it to interested readers.

Moreover, by interpolation, we can obtain

**Lemma 3.8.** *There exists  $\varepsilon_0 > 0$  such that*

$$N^{-1} \| |\nabla_x|^{\frac{1}{2}+\varepsilon} |\nabla_y|^{\frac{1}{2}+\varepsilon} u \|_{L^2(dxdy)} \lesssim C \quad (3.12)$$

for  $0 < \varepsilon \leq \varepsilon_0$ .

*Proof.* Applying Lemma 3.5 and (3.5) give us the estimate

$$\begin{aligned} & N^{-1} \| \nabla_x \nabla_y u(t) \|_{L^2(dxdy)} \\ & \lesssim N^{-1} \| \nabla_x \nabla_y \text{sh}(2k) \|_{L^2(dxdy)} + N^{-1} \| \nabla_x u \circ \nabla_y p \|_{L^2(dxdy)} \\ & \lesssim \| \nabla_x \phi \|_{L^2(dx)}^2 + \| \nabla_x \nabla_y \Lambda \|_{L^2(dxdy)} + N^{-1} \| \nabla_x u \circ \nabla_y p \|_{L^2(dxdy)} \\ & \lesssim 1 + N^{c_0}. \end{aligned}$$

By interpolating the above inequality with the energy estimate

$$N^{-1} \| |\nabla_x|^{\frac{1}{2}} |\nabla_y|^{\frac{1}{2}} u \|_{L^2(dxdy)} \lesssim N^{-1} \| \nabla_x u \|_{L^2(dxdy)} \lesssim \frac{1}{\sqrt{N}}, \quad (3.13)$$

we have shown that there exists  $\varepsilon_0 > 0$  such that

$$N^{-1} \| |\nabla_x|^{\frac{1}{2}+\varepsilon_0} |\nabla_y|^{\frac{1}{2}+\varepsilon_0} \text{sh}(k_t) \|_{L^2(dxdy)} \lesssim C$$

where  $C$  is a constant independent of  $N$  and  $t$ . In fact, we see that for all  $0 < \varepsilon \leq \varepsilon_0 = \frac{1}{2(2c_0+1)}$  we have the estimate

$$N^{-1} \| |\nabla_x|^{\frac{1}{2}+\varepsilon} |\nabla_y|^{\frac{1}{2}+\varepsilon} u(t) \|_{L^2(dxdy)} \lesssim N^{-\frac{1}{2}+(2c_0+1)\varepsilon}. \quad (3.14)$$

□

*Remark 3.9.* Since we need  $2\beta(\frac{1}{2} + \varepsilon) < 1$ , we need the assumption that  $0 < \varepsilon \leq \min(\frac{1-\beta}{2\beta}, \frac{1}{2(2c_0+1)})$  in our proof of the global well-posedness of the TDHFB system. Noticing that we can choose  $\varepsilon > 0$  sufficiently small, a positive power  $c_0$  is enough.

#### 4. GLOBAL WELL-POSEDNESS OF THE TDHFB SYSTEM

Let us introduce some notation that we will use in the proof of the uniform in  $N$  global well-posedness of (2.35) for  $0 < \beta < 1$ .

**4.1. Dispersive Estimates.** For the sake of completeness, we recall some of the standard a-priori estimates for the  $\Gamma$  and  $\Lambda$  equations proven in [GM17, GM19]. However, we will be more keen on keeping track of the parameters used in [GM17, GM19] to make some of their proofs more transparent, which will be necessary when we prove our global well-posedness statement.

Recall the definition of the  $X^{s,b}$  space (or sometimes known as the Bourgain space). In this paper, we take  $s = 0$  and write  $X^b = X^{0,b}$ . Let us denote  $\mathbf{S}_\pm$  and  $\mathbf{S}$  as in (2.28). Then, we see that the spacetime symbol associated with  $\mathbf{S}_\pm$  is  $\tau + |\xi|^2 - |\eta|^2$  and the spacetime symbol associated with  $\mathbf{S}$  is either  $\tau + |\xi|^2 + |\eta|^2$  or  $\tau + |\xi|^2$ , depending on the context. We define the norms

$$\begin{aligned}\|f\|_{X_\pm^b} &= \left\| \langle \tau + |\xi|^2 - |\eta|^2 \rangle^b \tilde{f}(\tau, \xi, \eta) \right\|_{L^2(d\tau d\xi d\eta)}, \\ \|f\|_{X_S^b} &= \left\| \langle \tau + |\xi|^2 + |\eta|^2 \rangle^b \tilde{f}(\tau, \xi, \eta) \right\|_{L^2(d\tau d\xi d\eta)}\end{aligned}$$

when  $d = 6$  or that

$$\|f\|_{X_S^b} = \left\| \langle \tau + |\xi|^2 \rangle^b \tilde{f}(\tau, \xi) \right\|_{L^2(d\tau d\xi)}$$

when  $d = 3$ . Here,  $\tilde{f}$  denotes the spacetime Fourier transform defined by

$$\tilde{f}(\tau, \xi, \eta) = \int dt dx dy e^{-i(\tau t + \xi \cdot x + \eta \cdot y)} f(t, x, y). \quad (4.1)$$

See Chapter 2 of [Tao06] for an exposition of the topics.

Let us prove the following time localization lemma with characteristic cutoff function which is a generalization of the smooth time-localization of Lemma 2.11 in [Tao06].

**Lemma 4.1.** *Let  $c(t)$  be the characteristic function of  $[0, 1]$ . Suppose  $0 < T < 1$  and  $-\frac{1}{2} < b' \leq b < \frac{1}{2}$ , then we have*

$$\|c(t/T)u\|_{X^{b'}} \lesssim_{c,b,b'} T^{b-b'} \|u\|_{X^b}. \quad (4.2a)$$

Likewise, we have the estimate

$$\|c(t/T)u\|_{X_\pm^{b'}} \lesssim_{c,b,b'} T^{b-b'} \|u\|_{X_\pm^b}. \quad (4.2b)$$

*Proof.* Let us define the convolution operator  $I(f)(t) := (\widehat{c(\cdot/T)} * f)(\tau)$ , then it is clear that  $I$  is a  $L^2$ - $L^2$  bounded operator since

$$\|I(f)\|_{L^2(d\tau)} = \|c(\cdot/T)f\|_{L^2(dt)} \leq \|f\|_{L^2(dt)} = \|f\|_{L^2(d\tau)}.$$

Moreover, we see that  $|\partial^k \widehat{c(\cdot/T)}(\tau)| \lesssim |\tau|^{-1-k}$  for  $k = 0, 1$ . Then, by Corollary of Theorem 2 in section V.4.2 in Stein, we see have the estimate

$$\|I(f)\|_{L^2(\omega(\tau)d\tau)} \lesssim \|f\|_{L^2(\omega(\tau)d\tau)} \quad (4.3)$$

where  $\omega \in A_2$ ; see chapter V of Stein for definition of  $A_p$  spaces. In particular, it follows that

$$\|c(t/T)u\|_{X^b} \lesssim_{c,b} \|u\|_{X^b} \quad (4.4)$$

which is mainly what we need from this lemma.

Now to prove (4.2a), we decompose  $u$  into two parts

$$\begin{aligned} u &= P_{\langle \tau+|\xi|^2+|\eta|^2 \rangle \geq \frac{1}{T}} u + P_{\langle \tau+|\xi|^2+|\eta|^2 \rangle < \frac{1}{T}} u \\ &=: P_1(c(t/T)u) + P_2(c(t/T)u). \end{aligned}$$

Then it follows

$$\begin{aligned} \|c(t/T)P_1u\|_{X^{b'}}^2 &= \int \|\widehat{c(\cdot/T)} * \widetilde{P_1u}\|_{L^2(\langle \tau+|\xi|^2+|\eta|^2 \rangle^{2b'} d\tau)}^2 d\xi d\eta \\ &\lesssim \int \|\widetilde{P_1u}\|_{L^2(\langle \tau+|\xi|^2+|\eta|^2 \rangle^{2b'} d\tau)}^2 d\xi d\eta \\ &\lesssim T^{2(b-b')} \int \|\tilde{u}\|_{L^2(\langle \tau+|\xi|^2+|\eta|^2 \rangle^{2b} d\tau)}^2 d\xi d\eta \end{aligned}$$

where the second inequality follows from the remark of Stein chapter V.6.4. which states that  $|\tau|^{2b} \in A_2$  provided  $|b| < \frac{1}{2}$ .

Finally, we see that

$$\begin{aligned} \|\widehat{P_2u}(t)\|_{L^2(d\xi d\eta)} &\lesssim \left\| \int_{\langle \tau+|\xi|^2+|\eta|^2 \rangle \leq 1/T} d\tau |\tilde{u}(\tau, \xi, \eta)| \right\|_{L^2(d\xi d\eta)} \\ &\lesssim T^{b-\frac{1}{2}} \left\| \left( \int d\tau \langle \tau+|\xi|^2+|\eta|^2 \rangle^{2b} |\widetilde{P_2u}(\tau, \xi, \eta)|^2 \right)^{\frac{1}{2}} \right\|_{L^2(d\xi d\eta)} \\ &= T^{b-\frac{1}{2}} \|P_2u\|_{X^b} \end{aligned}$$

which means

$$\|c(t/T)P_2u\|_{X^0}^2 \leq \int_0^T dt \|P_2u(t)\|_{L^2(dx dy)}^2 \lesssim T^{2b} \|P_2u\|_{X^b}^2. \quad (4.5)$$

Interpolating (4.5) with (4.4) where  $u$  is replaced with  $P_2u$  yields the desired result.  $\square$

Here are some standard dispersive estimates which we will need later in the section. The proofs of the following two lemmas can be found in section 4 of [GM19].

**Lemma 4.2.** *Let  $\delta > \frac{1}{2}$  and  $q, r \geq 2$  satisfy the admissible condition. Then we have the estimates*

$$\|F\|_{L^q(dt)L^r(dx)L^2(dy)} \lesssim_\delta \|F\|_{X^\delta}, \text{ and its dual estimate} \quad (4.6a)$$

$$\|F\|_{X^{-\delta}} \lesssim_\delta \|F\|_{L^{q'}(dt)L^{r'}(dx)L^2(dy)}. \quad (4.6b)$$

**Lemma 4.3.** *Let  $0 < \delta < \frac{1}{2}$  and  $q, r > 2$  such that  $\frac{2}{q} + \frac{3}{r} = \frac{5-4\delta}{2}$ . Then we have*

$$\|F\|_{L^q(dt)L^r(dx)L^2(dy)} \lesssim_\delta \|F\|_{X^\delta}, \quad \text{and its dual estimate} \quad (4.7a)$$

$$\|F\|_{X^{-\delta}} \lesssim_\delta \|F\|_{L^{q'}(dt)L^{r'}(dx)L^2(dy)}. \quad (4.7b)$$

**4.2. Global Well-posedness.** Fix  $\varepsilon > 0$  small enough and consider  $c(t) = c_I(t)$  the characteristic function on the interval  $I = [T_0, T_1]$ . Then we define

$$\mathbf{N}_I(\Lambda) := \sup_z \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} c(t) \Lambda(t, x, x+z)\|_{L^2(dt)L^2(dx)} \quad (4.8a)$$

$$\begin{aligned} &+ \sup_z \|\partial_t^{\frac{1}{4}}[c(t) \Lambda(t, x, x+z)]\|_{L^2(dt)L^2(dx)} \\ &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Lambda\|_{L^\infty(dt)L^2(dx dy)} \\ &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Lambda\|_{L^2(dt)L^6(dx)L^2(dy)} \\ &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Lambda\|_{L^2(dt)L^6(dy)L^2(dx)} \end{aligned}$$

$$\mathbf{N}_I(\Gamma) := \sup_z \|\langle \nabla_x \rangle^{\frac{1}{2}} \langle \nabla_x \rangle^\varepsilon c(t) \Gamma(t, x, x+z)\|_{L^2(dt)L^2(dx)} \quad (4.8b)$$

$$\begin{aligned} &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Gamma\|_{L^\infty(dt)L^2(dx dy)} \\ &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Gamma\|_{L^2(dt)L^6(dx)L^2(dy)} \\ &+ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) \Gamma\|_{L^2(dt)L^6(dy)L^2(dx)} \end{aligned}$$

$$\mathbf{N}_I(\phi) := \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} c(t) \phi\|_{L^\infty(dt)L^2(dx)} + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} c(t) \phi\|_{L^2(dt)L^6(dx)}. \quad (4.8c)$$

Like before, we denote the sum of the three norms by

$$\mathbf{N}_I(X) := \mathbf{N}_I(\phi) + \mathbf{N}_I(\Gamma) + \mathbf{N}_I(\Lambda). \quad (4.9)$$

If  $I = [T_0, T_1] = [0, T]$  then we recover  $\mathbf{N}_T(X) = \mathbf{N}_I(X)$ . Moreover, we adopt the notation

$$\mathbf{N}_I(DX) := \mathbf{N}_I(D\phi) + \mathbf{N}_I(D\Gamma) + \mathbf{N}_I(D\Lambda) \quad (4.10)$$

for any differential operator  $D$ .

The goal of this section is to prove the global well-posedness of solutions for the time-dependent HFB equations. However, it suffices to prove an a-priori estimate of the form

$$\mathbf{N}_T(X) \lesssim F(T) \quad (4.11)$$

for some positive real-valued function  $F$  defined on all of  $[0, \infty)$ .

Let us now recall some propositions from [GM19], which are modified by Lemma 4.1. For the  $\phi$  and  $\Gamma$  equations, we have the following proposition.

**Proposition 4.4** (Proposition 3.7 in [GM19]). *Let  $c(t)$  be the characteristic function on  $I = [T_0, T_1]$  such that  $T_1 - T_0 < 1$ . Suppose  $\phi(t)$  and  $\Gamma(t)$  are*

solutions to

$$\begin{aligned}\mathbf{S}_\pm \Gamma &= F, \quad \Gamma(T_0) = \Gamma_0 \\ \mathbf{S}\phi &= f, \quad \phi(T_0) = \phi_0.\end{aligned}$$

Then for any  $\delta > 0$  we have the estimates

$$\begin{aligned}\mathbf{N}_I(\Gamma) &\lesssim_\delta \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma_0\|_{L^2(dxdy)} \\ &\quad + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t)F\|_{X_\pm^{-\frac{1}{2}+\delta}}\end{aligned}\tag{4.12a}$$

$$\mathbf{N}_I(\phi) \lesssim_\delta \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \phi_0\|_{L^2(dxdy)} + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} c(t)f\|_{X^{-\frac{1}{2}+\delta}}.\tag{4.12b}$$

For the  $\Lambda$  equations, with the potential  $\frac{1}{N}v_N(x-y)\Lambda$ , we have the following proposition, which is also the main result of [GM19].

**Proposition 4.5** (Theorem 3.8 in [GM19]). *Assume  $c(t)$  as in Proposition 4.4. Let  $0 < \beta < 1$ . Suppose  $\Lambda(t)$  is a solution to*

$$\left(\mathbf{S} + \frac{1}{N}v_N(x-y)\right)\Lambda = F, \quad \Lambda(0) = \Lambda_0.\tag{4.13}$$

Then for all  $\delta > 0$  sufficiently small, the following estimate

$$\begin{aligned}\mathbf{N}_I(\Lambda) &\lesssim_\delta \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda_0\|_{L^2(dxdy)} + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t)F\|_{X^{-\frac{1}{2}+\delta}} \\ &\quad + \min \left\{ \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^\varepsilon c(t)F\|_{X^{-\frac{1}{4}-\delta}}, \|\langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t)F\|_{X^{-\frac{1}{4}-\delta}} \right\}\end{aligned}\tag{4.14}$$

holds uniformly in  $N$ .

We begin by proving couple lemmas to aid us in establishing (4.11).

**Lemma 4.6** (sharp trace-type lemma). *Let  $f \in \mathcal{S}(\mathbb{R}^3 \times \mathbb{R}^3)$ . Then the following estimate holds*

$$\| |\nabla_x|^{\frac{1}{2}} \text{diag } f \|_{L^2(dx)} \lesssim \| \nabla_x \nabla_y f \|_{L^2(dxdy)}.\tag{4.15}$$

*Proof.* Note that

$$\widehat{\text{diag } f}(\xi) = \frac{1}{(2\pi)^3} \int d\eta \widehat{f}(\xi - \eta, \eta).$$

Then, by Plancherel and Cauchy Schwarz, we have the estimate

$$\begin{aligned}\| |\nabla_x|^{\frac{1}{2}} \text{diag } f \|_{L^2(dx)}^2 &\sim \int d\xi |\xi| \left| \int d\eta \widehat{f}(\xi - \eta, \eta) \right|^2 \\ &\lesssim \sup_\xi A(\xi) \int d\xi d\eta |\xi|^2 |\eta|^2 |\widehat{f}(\xi, \eta)|^2\end{aligned}$$

where

$$A(\xi) = \int d\eta \frac{|\xi|}{|\xi - \eta|^2 |\eta|^2}.$$



Hence, to complete the proof, it suffices to show  $\sup_{\xi} A(\xi) < \infty$ .

Let us evaluate the integral  $A(\xi)$  into each of the following three regions:  $R_1 = \{\frac{1}{2}|\xi| \leq |\eta| \leq 2|\xi|\}$ ,  $R_2 = \{\frac{1}{2}|\xi| \leq |\eta - \xi| \leq 2|\xi|\}$ , and  $R_3 = \{|\eta| > 2|\xi| \text{ or } |\eta - \xi| > 2|\xi|\}$ . Then, we see that

$$\int_{R_1} d\eta \frac{|\xi|}{|\xi - \eta|^2 |\eta|^2} \lesssim \frac{1}{|\xi|} \int_{|\xi - \eta| \leq 3|\xi|} \frac{d\eta}{|\xi - \eta|^2} \lesssim 1$$

and likewise for  $R_2$ . For the region  $R_3$ , we see that  $|\eta| > |\xi|$  and  $2|\xi - \eta| \geq |\xi - \eta| + |\xi| \geq |\eta|$ , which means

$$\int_{R_3} d\eta \frac{|\xi|}{|\xi - \eta|^2 |\eta|^2} \lesssim \int_{|\eta| > |\xi|} d\eta \frac{|\xi|}{|\eta|^4} \lesssim 1.$$

Thus, we have that  $\sup_{\xi} A(\xi) < \infty$ .  $\square$

**Lemma 4.7.** *Let  $(\phi(t), \Gamma(t), \Lambda(t))$  be a solution to (2.35) and  $I = [T_0, T_1]$ . Then there exists  $\delta > 0$  such that we have the following estimates*

$$\mathbf{N}_I(X) \lesssim (\text{data}) + |I|^{\delta} \mathbf{N}_I(X). \quad (4.16)$$

*Proof.* It suffices to consider the proof of estimate (4.16) for  $\Gamma$  and  $\Lambda$  since the proof for  $\phi$  is similar. Recall the equation for  $\Gamma$  is given by

$$\begin{aligned} \mathbf{S}_{\pm} \bar{\Gamma} &= - (v_N \Lambda) \circ \bar{\Lambda} + \Lambda \circ (v_N \bar{\Lambda}) - (v_N \bar{\Gamma}) \circ \bar{\Gamma} + \bar{\Gamma} \circ (v_N \bar{\Gamma}) \\ &\quad - (v_N * \text{diag } \Gamma) \cdot \bar{\Gamma} + \bar{\Gamma} \cdot (v_N * \text{diag } \Gamma) \\ &\quad + 2(v_N * |\phi|^2) \cdot |\phi\rangle \langle \phi| - 2|\phi\rangle \langle \phi| \cdot (v_N * |\phi|^2) =: F. \end{aligned} \quad (4.17)$$

Then, by Proposition 4.4 and Lemma 4.3, we have that for any  $0 < \delta < \frac{\varepsilon}{2}$  the following estimate holds

$$\begin{aligned} \mathbf{N}_I(\Gamma) &\lesssim \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma(T_0, \cdot)\|_{L^2(dx dy)} \\ &\quad + \left\| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) F \right\|_{X^{-\frac{1}{2}+\frac{\varepsilon-2\delta}{2}}} \\ &\lesssim \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma(T_0, \cdot)\|_{L^2(dx dy)} \\ &\quad + |I|^{\frac{\varepsilon-2\delta}{2}} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} c(t) F\|_{L^2(dt) L^{\frac{6}{5}+}(dx) L^2(dy)}. \end{aligned}$$

Here, the symbol  $\frac{6}{5}+$  denotes a fixed number slightly bigger than  $\frac{6}{5}$  with dependence on  $\varepsilon$ , in fact,  $\frac{6}{5}+ = \frac{6}{5-2\varepsilon}$ .

For the forcing term  $\Gamma \cdot (v_N * \text{diag } \Gamma)$ , we apply Young's convolution inequality, Sobolev inequality, and Corollary 3.3 to obtain the estimate

$$\begin{aligned} &\|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [\bar{\Gamma} \cdot (v_N * \text{diag } \Gamma)]\|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)} \\ &\lesssim \|\langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \bar{\Gamma}\|_{L^{\infty}(I) L^{3+}(dx) L^2(dy)} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \text{diag } \Gamma\|_{L^2(I) L^2(dx)} \\ &\quad + \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \bar{\Gamma}\|_{L^{\infty}(I) L^2(dx dy)} \|\text{diag } \Gamma\|_{L^2(I) L^{3+}(dx)} \\ &\lesssim \mathbf{N}_I(\Gamma), \end{aligned}$$

where  $3+ = \frac{3}{1-\varepsilon}$ . Next, for the forcing term  $\Lambda \circ (v_N \bar{\Lambda})$ , we apply Kato-Ponce, Sobolev, and estimate (3.1c) from the previous section to obtain the estimate

$$\begin{aligned}
& \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [\Lambda \circ (v_N \bar{\Lambda})] \|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)} \\
& \lesssim \int dz v_N(z) \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \bar{\Lambda}(x, x-z) \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda(x-z, y) \|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)} \\
& \quad + \int dz v_N(z) \| \bar{\Lambda}(x, x-z) \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda(x-z, y) \|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)} \\
& \lesssim \int dz v_N(z) \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \bar{\Lambda}(x, x-z) \|_{L^2(dt dx)} \| \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda \|_{L^\infty(I) L^{3+}(dx) L^2(dy)} \\
& \quad + \int dz v_N(z) \| \Lambda(x, x-z) \|_{L^2(dt) L^{3+}(dx)} \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda \|_{L^\infty(I) L^2(dx dy)} \\
& \lesssim \mathbf{N}_I(\Lambda).
\end{aligned}$$

The remaining nonlinear terms  $(v_N \Gamma) \circ \Gamma$  and  $(v_N * |\phi|^2) \cdot |\phi| \langle \phi |$  can be handled in a similar manner. Thus, we have shown

$$\mathbf{N}_I(\Gamma) \lesssim C_0(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \mathbf{N}_I(X).$$

Next, let us recall the equation for  $\Lambda$  given by

$$\begin{aligned}
\left( \mathbf{S} + \frac{1}{N} v_N \Lambda \right) &= - (v_N * \text{diag } \Gamma) \cdot \Lambda - \Lambda \cdot (v_N * \text{diag } \Gamma) \\
&\quad - (v_N \Lambda) \circ \Gamma - \bar{\Gamma} \circ (v_N \Lambda) - (v_N \bar{\Gamma}) \circ \Lambda - \Lambda \circ (v_N \Gamma) \\
&\quad + 2(v_N * |\phi|^2) \cdot \phi \otimes \phi - 2\phi \otimes \phi \cdot (v_N * |\phi|^2) =: G.
\end{aligned} \tag{4.18}$$

To estimate  $\Lambda$ , we employ Proposition 4.5 and Lemma 4.3 to get the estimate

$$\begin{aligned}
\mathbf{N}_I(\Lambda) &\lesssim \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda(T_0, \cdot) \|_{L^2(dx dy)} \\
&\quad + |I|^{\frac{\varepsilon-2\delta}{2}} \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} G \|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)} \\
&\quad + |I|^{\frac{\varepsilon-2\delta}{2}} \min \left( \| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} G \|_{L^{\frac{4}{3}}(I) L^2(dx dy)}, \right. \\
&\quad \left. \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^\varepsilon G \|_{L^{\frac{4}{3}}(I) L^2(dx dy)} \right).
\end{aligned}$$

Hence, to complete the proof, it suffices to estimate  $\langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} G$  in  $L^{\frac{4}{3}}(I) L^2(dx dy)$  since  $\| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} G \|_{L^2(I) L^{\frac{6}{5}+}(dx) L^2(dy)}$  can be handled in a similar manner like  $\Gamma$ .

Let us consider the forcing term  $(v_N * \text{diag } \Gamma) \cdot \Lambda$ . Then, by Kato-Ponce and Sobolev inequalities, we have that

$$\begin{aligned} & \| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [(v_N * \text{diag } \Gamma) \cdot \Lambda] \|_{L^{\frac{4}{3}}(I)L^2(dx dy)} \\ & \lesssim \| \langle \nabla_x \rangle^\varepsilon \text{diag } \Gamma \|_{L^2(I)L^{3-}(dx)} \| \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda \|_{L^4(I)L^{6+}(dx)L^2(dy)} \\ & \quad + \| \text{diag } \Gamma \|_{L^2(I)L^3(dx)} \| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda \|_{L^4(I)L^6(dx)L^2(dy)} \\ & \lesssim \| \langle \nabla_x \rangle^\varepsilon |\nabla_x|^{\frac{1}{2}-\varepsilon} \text{diag } \Gamma \|_{L^2(I)L^2(dx)} \| \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Lambda \|_{L^4(I)L^3(dx)L^2(dy)}. \end{aligned}$$

Finally, by Lemma 4.6 and Corollary 3.3, we obtain the estimate

$$\| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [(v_N * \text{diag } \Gamma) \cdot \Lambda] \|_{L^{\frac{4}{3}}(I)L^2(dx dy)} \lesssim |I|^{\frac{1}{2}} \mathbf{N}_I(\Lambda).$$

Next, let us consider the forcing term  $(v_N \Lambda) \circ \Gamma$ . By Kato-Ponce, Sobolev inequalities, and conservation of energy, we have that

$$\begin{aligned} & \| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [(v_N \Lambda) \circ \Gamma] \|_{L^{\frac{4}{3}}(I)L^2(dx dy)} \\ & \lesssim \int dz v_N(z) \| \langle \nabla_x \rangle^\varepsilon \Lambda(x, x-z) \|_{L^2(I)L^3(dx)} \| \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma \|_{L^4(I)L^6(dx)L^2(dy)} \\ & \quad + \int dz v_N(z) \| \Lambda(x, x-z) \|_{L^2(I)L^{3+}(dx)} \| \langle \nabla_x \rangle^\varepsilon \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \Gamma \|_{L^4(I)L^{6-}(dx)L^2(dy)} \\ & \lesssim |I|^{\frac{1}{4}} \mathbf{N}_I(\Lambda). \end{aligned}$$

The remaining forcing terms can be handled in a similar fashion. Hence it follows

$$\mathbf{N}_I(\Lambda) \lesssim C_0(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \mathbf{N}_I(X)$$

which completes the proof of the lemma.  $\square$

Next we aim to obtain a-priori estimates of the form (4.11).

**Proposition 4.8.** *Let  $T > 0$ . Assume  $(\phi(t), \Gamma(t), \Lambda(t))$  is a solution to the time-dependent HFB system, then we have the following a-priori estimate*

$$\mathbf{N}_T(X) \lesssim (\text{data}) + T. \quad (4.19)$$

*Proof.* Proposition 4.7 implies that for an interval with length similar to 1,

$$\mathbf{N}_I(X) \lesssim_{(\text{data})} 1. \quad (4.20)$$

We may split the interval  $[0, T]$  into  $M$  intervals  $I_i$  where  $M \sim T$  and the length of each interval  $I_i$  is less than  $\frac{1}{2}$ . Then we sum the norms  $\mathbf{N}_{I_i}(X)$  up which explains this proposition.  $\square$

Now Theorem 2.7 follows. In particular, we obtained the linear-in-time control for  $\mathbf{N}_T(X)$ .

**Lemma 4.9.** *Let  $(\phi(t), \Gamma(t), \Lambda(t))$  be a solution to (2.35) and  $I = [T_0, T_1]$ . Then there exists  $\delta > 0$  such that we have the following estimates*

$$\mathbf{N}_I(\nabla_{x+y} X) \lesssim C_1(T_0) + |I|^\delta \mathbf{N}_I(X) \mathbf{N}_I(\nabla_{x+y} X) \quad (4.21a)$$

where

$$C_1(T) := \|(\nabla_x \phi(T, \cdot), \nabla_{x+y} \Gamma(T, \cdot), \nabla_{x+y} \Lambda(T, \cdot))\|_{\mathcal{X}^\alpha}. \quad (4.21b)$$

Here, we use the notation  $\nabla_{x+y} := \nabla_x + \nabla_y$ .

*Proof.* Taking the  $\nabla_{x+y}$  derivative of (3.10) yields

$$\begin{aligned} (\mathbf{S} + N^{-1}v_N) \nabla_{x+y} \Lambda &= - (v_N \Lambda) \circ \nabla_{x+y} \Gamma - (v_N * \text{diag } \Gamma) \cdot \nabla_{x+y} \Lambda \\ &\quad + \text{similar terms} =: F. \end{aligned}$$

Note that we used the fact that  $\nabla_{x+y}$  commutes with  $N^{-1}v_N(x-y)$ , i.e.  $[\nabla_{x+y}, N^{-1}v_N(x-y)] = 0$ . By the same argument as in Lemma 4.7, we have the estimate

$$\mathbf{N}_I(\nabla_{x+y} \Lambda) \lesssim C_1(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} F\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)}.$$

We shall look at two generic cases, as stated above, to deduce (4.21a). In the first case, we estimate the term  $(v_N \Lambda) \circ \nabla_{x+y} \Gamma$ , which goes as follows

$$\begin{aligned} &\|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} (v_N \Lambda) \circ \nabla_{x+y} \Gamma\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\lesssim \int dz v_N(z) \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \Lambda(x, x-z) \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Gamma(x-z, y)\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\quad + \int dz v_N(z) \|\Lambda(x, x-z) \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Gamma(x-z, y)\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\lesssim \int dz v_N(z) \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \Lambda(x, x-z)\|_{L^2(I) L^2(dx)} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Gamma\|_{L^\infty(I) L^2(dx dy)} \\ &\quad + \int dz v_N(z) \|\Lambda(x, x-z)\|_{L^2(I) L^{3+(dx)}} \|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Gamma\|_{L^\infty(I) L^2(dx dy)} \\ &\lesssim \mathbf{N}_I(\Lambda) \mathbf{N}_I(\nabla_{x+y} \Gamma). \end{aligned}$$

In the second case, we estimate the term  $(v_N * \text{diag } \Gamma) \cdot \nabla_{x+y} \Lambda$  as follows

$$\begin{aligned} &\|\langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} [(v_N * \text{diag } \Gamma) \cdot \nabla_{x+y} \Lambda]\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\lesssim \|(v_N * \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \text{diag } \Gamma) \cdot \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Lambda\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\quad + \|(v_N * \langle \nabla_x \rangle^{\frac{1}{2}+\varepsilon} \text{diag } \Gamma) \cdot \langle \nabla_y \rangle^{\frac{1}{2}+\varepsilon} \nabla_{x+y} \Lambda\|_{L^2(I) L^{\frac{6}{5}+(dx)} L^2(dy)} \\ &\lesssim \mathbf{N}_I(\Gamma) \mathbf{N}_I(\nabla_{x+y} \Lambda). \end{aligned}$$

Hence, combining the above estimates yields

$$\begin{aligned} \mathbf{N}_I(\nabla_{x+y} \Lambda) &\lesssim C_1(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \{\mathbf{N}_I(\Lambda) \mathbf{N}_I(\nabla_{x+y} \Gamma) \\ &\quad + \mathbf{N}_I(\nabla_{x+y} \Lambda) \mathbf{N}_I(\Gamma) + \mathbf{N}_I(\nabla_{x+y} \phi) \mathbf{N}_I(\phi)\} \\ &\lesssim C_1(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \mathbf{N}_I(X) \mathbf{N}_I(\nabla_{x+y} X). \end{aligned}$$

Similarly, we can show

$$\mathbf{N}_I(\nabla_{x+y} \Gamma) \lesssim C_1(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \mathbf{N}_I(X) \mathbf{N}_I(\nabla_{x+y} X)$$

and

$$\mathbf{N}_I(\nabla_{x+y}\phi) \lesssim C_1(T_0) + |I|^{\frac{\varepsilon-2\delta}{2}} \mathbf{N}_I(X) \mathbf{N}_I(\nabla_{x+y}X).$$

Therefore, summing up the three inequalities yields (4.21a).  $\square$

Using the above lemma we could again prove an a-priori growth estimates for the norm of  $\nabla_{x+y}X$ .

**Proposition 4.10.** *Let  $T > 0$ . Suppose  $(\phi(t), \Gamma(t), \Lambda(t))$  is a solution to the TDHFB system, then we have the following uniform in  $N$  a-priori estimate*

$$\mathbf{N}_T(\nabla_{x+y}X) \lesssim \exp(\kappa T) \quad (4.22a)$$

for some  $\kappa > 0$ , which are independent of  $T$ .

*Proof.* Proposition 4.7 and Proposition 4.9 imply that for an interval  $I$  with length similar to 1,

$$\mathbf{N}_I(\nabla_{x+y}X) \lesssim_{(\text{data})} 1. \quad (4.23)$$

Here we note that for this case, the  $C_1(T)$  is not conserved, which is different from Proposition 4.8. Next we split the interval  $[0, T]$  into  $M'$  intervals  $I_i = [T_i, T_{i+1}]$  where  $M \sim T$  and the length of each interval  $I_i$  is less than  $\frac{1}{2}$ . On each interval,

$$\mathbf{N}_{I_i}(\nabla_{x+y}X) \lesssim C_1(T_i). \quad (4.24)$$

Summing them up, we have

$$\mathbf{N}_T(\nabla_{x+y}X) \lesssim \sum_i \mathbf{N}_{I_i}(\nabla_{x+y}X) \lesssim C_1(T_0) + \sum_i C_1(T_i). \quad (4.25)$$

By switching to the continuous  $T$ -variable, we obtain the estimate

$$\mathbf{N}_T(\nabla_{x+y}X) \lesssim C_1(T_0) + \int_0^T d\tau \mathbf{N}_\tau(\nabla_{x+y}X).$$

Finally, applying Gronwall's inequality yields

$$\mathbf{N}_T(\nabla_{x+y}X) \lesssim C_1(T_0) \exp(\alpha T),$$

which explains this proposition.  $\square$

Let us conclude this section with some a-priori estimates for the higher order derivatives of  $(\phi, \Gamma, \Lambda)$  which we will later use to estimate  $\text{sh}(2k)$ .

**Lemma 4.11.** *Let  $(\phi(t), \Gamma(t), \Lambda(t))$  be a solution to (2.35) and  $I = [T_0, T_1]$ . Then there exists  $\delta > 0$  such that for  $j \geq 2$  we have the estimates*

$$\mathbf{N}_I(\nabla_{x+y}^j X) \lesssim_j C_j(T_0) + |I|^\delta \sum_{i=0}^{[j/2]} \mathbf{N}_I(\nabla_{x+y}^i X) \mathbf{N}_I(\nabla_{x+y}^{j-i} X) \quad (4.26a)$$

where

$$C_j(T) = \|(\nabla_x^j \phi(T, \cdot), \nabla_{x+y}^j \Gamma(T, \cdot), \nabla_{x+y}^j \Lambda(T, \cdot))\|_{\mathcal{X}^\alpha}. \quad (4.27)$$

In particular, for  $|I|$  sufficiently small, we have that

$$\mathbf{N}_I(\nabla_{x+y}^j X) \lesssim C_j(T_0). \quad (4.28)$$

*Proof.* The proof is similar to the proof of Lemma 4.9.  $\square$

**Proposition 4.12.** *Let  $T > 0$ . Suppose  $(\phi(t), \Gamma(t), \Lambda(t))$  is a solution to the TDHFB system. Then there exists a constant  $\kappa > 0$ , depending on  $j$  and independent of  $T$ , such that we have the following uniform in  $N$  a-priori estimate*

$$\mathbf{N}_T(\nabla_{x+y}^j X) \lesssim_j \exp(\kappa T). \quad (4.29)$$

*Proof.* The proof is an induction argument. It is similar to the proof of Lemma 4.10 so we omit it.  $\square$

## 5. ESTIMATES FOR $\text{sh}(2k)$ AND $\text{ch}(2k)$

In order to obtain Fock space estimates for the error terms in our quasifree-approximation, we need to establish estimates for  $\text{sh}(2k)$  and  $\text{ch}(2k)$ .

Recall the equation for  $\text{sh}(2k)$  is given by

$$\begin{aligned} \mathbf{S}(\text{sh}(2k)) = & -2v_N(x-y)\Lambda - (v_N\Lambda) \circ p_2 - \bar{p}_2 \circ (v_N\Lambda) \\ & - ((v_N * \text{diag } \Gamma)(x) + (v_N * \text{diag } \Gamma)(y)) \text{sh}(2k) \\ & - (v_N\Gamma) \circ \text{sh}(2k) - \text{sh}(2k) \circ (v_N\Gamma). \end{aligned} \quad (5.1a)$$

Since  $\text{ch}(2k) = \delta(x-y) + p_2$  then we could simply write an equation for  $p_2$  instead, that is

$$\begin{aligned} \mathbf{S}_{\pm}(\bar{p}_2) = & - (v_N\Lambda) \circ \overline{\text{sh}(2k)} + \text{sh}(2k) \circ (v_N\Lambda) \\ & - ((v_N * \text{diag } \Gamma)(x) - (v_N * \text{diag } \Gamma)(y))\bar{p}_2 \\ & - (v_N\Gamma) \circ \bar{p}_2 + \bar{p}_2 \circ (v_N\Gamma). \end{aligned} \quad (5.1b)$$

*Remark 5.1.* The main difficulty one faces when handling (5.1a) and (5.1b) is in controlling the singular potential term  $v_N\Lambda$  of the  $\text{sh}(2k)$  equation. In a former version of the paper, we required a bounded on the time-derivative of the initial which greatly limited the kind of initial data that we could study. In particular, the requirement of bounded one time-derivative rules out the possibility of studying the case  $k(0) \equiv 0$ , i.e. the coherent state case (c.f. Remark 2.6 in [GM17]).

The following theorem is the main result of this section.

**Theorem 5.2.** *Let  $\text{sh}(2k)$  satisfies (5.1a) with initial conditions given by the Nonlinear Theorem. Then for  $j = 0, 1, 2$  we have the estimates*

$$\|\nabla_{x+y}^j \text{sh}(2k)(t, \cdot, \cdot)\|_{L^2(dx dy)} \lesssim \exp(\kappa T) \quad (5.2a)$$

$$\|\nabla_{x+y}^j \text{sh}(2k)(t, \cdot, \cdot)\|_{L^2([0, T])L^6(dx)L^2(dy)} \lesssim \exp(\kappa T) \quad (5.2b)$$

$$\sup_x \|\text{sh}(2k)(x, \cdot)\|_{L^2(dz)} \lesssim \exp(\kappa T) \quad (5.2c)$$

for some constant  $\kappa > 0$  and uniform in  $t$  on the interval  $[0, T]$ . Similar estimates hold for  $p_2$ .

As an immediate consequence, we have the following corollary.

**Corollary 5.3.** *Suppose  $(\phi(t), k(t))$  satisfies (2.30). Then there is some constant  $\kappa > 0$  such that the following estimates*

$$\|u(t, \cdot, \cdot)\|_{L^2(dx dy)} \lesssim \exp(\kappa T) \quad (5.3a)$$

$$\|p(t, \cdot, \cdot)\|_{L^2(dx dy)} \lesssim \exp(\kappa T) \quad (5.3b)$$

$$\sup_x \|u(t, x, \cdot)\|_{L^2(dy)} \lesssim \exp(\kappa T) \quad (5.3c)$$

$$\sup_x \|p(t, x, \cdot)\|_{L^2(dy)} \lesssim \exp(\kappa T) \quad (5.3d)$$

hold uniformly in  $N$  on the interval  $[0, T]$ .

*Proof.* Let us recall that  $\text{ch}(k) = \delta + p$  is an invertible operator since  $p$  is bounded (semi)positive-definite operator. In fact, we have

$$\|f\|_{L^2(dx)}^2 \lesssim \langle f, \text{ch}(k)f \rangle \implies \|\text{ch}(k)^{-1}\|_{\text{op}} \lesssim 1. \quad (5.4)$$

Hence, using the identity (3.5) and Theorem 5.2, it follows

$$\begin{aligned} \|u(t, \cdot, \cdot)\|_{L^2(dx dy)} &\lesssim \|\text{ch}(k)^{-1}\|_{\text{op}} \|\text{sh}(2k)(t, \cdot, \cdot)\|_{L^2(dx dy)} \\ &\lesssim \exp(\kappa T). \end{aligned}$$

Likewise, using identity (3.9), we see that

$$\|p(t, \cdot, \cdot)\|_{L^2(dx dy)} \lesssim \|u(t, \cdot, \cdot)\|_{L^2(dx dy)} \lesssim \exp(\kappa T).$$

To prove the remaining inequalities, we will use a duality argument. Observe we have that

$$\begin{aligned} \|u\|_{L^\infty(dx) L^2(dy)} &= \sup_{\|f\|_{L^1(dx) L^2(dy)}=1} \left| \int dx dy u(x, y) f(x, y) \right| \\ &\leq \sup_{\|f\|_{L^1(dx) L^2(dy)}=1} \left| \int dx dz dy \text{sh}(2k)(x, z) \text{ch}(k)^{-1}(z, y) f(x, y) \right| \\ &\leq \sup_{\|f\|_{L^1(dx) L^2(dy)}=1} \int dx \|\text{sh}(2k)(x, \cdot)\|_{L^2(dz)} \|\text{ch}(k)^{-1}\|_{\text{op}} \|f(x, \cdot)\|_{L^2(dy)} \\ &\leq \sup_x \|\text{sh}(2k)(x, \cdot)\|_{L^2(dz)}. \end{aligned}$$

Hence the result follows. The estimate for  $p$  follows from the identity  $p_2 \circ (\delta + \frac{1}{2}p)^{-1} = 4p$ .  $\square$

We also have the following corollary of the above corollary.

**Corollary 5.4.** *Let  $(\phi(t), k(t))$  solves (2.30). Then we have the following uniform in  $N$  endpoint Strichartz estimates*

$$\|u\|_{L^2([0, T]) L^6(dx) L^2(dy)} \lesssim \exp(\kappa T), \quad (5.5a)$$

$$\|p\|_{L^2([0, T]) L^6(dx) L^2(dy)} \lesssim \exp(\kappa T). \quad (5.5b)$$

The proof of Corollary 5.4 will be postponed till the end of the section since the crux of the argument follows from the proof of Theorem 5.2.

*Remark 5.5.* The reader should note that by interpolating (5.2a) and (5.2b) we recover all the Strichartz estimates of the form  $L^q L^r L^2$ , where  $(q, r)$  is an admissible pair, for  $\text{sh}(2k)$ . Likewise, we have all the Strichartz estimates for  $u$  and  $p$ .

**Proposition 5.6.** *Let  $s_a^0$  be the solution to*

$$\begin{aligned} \mathbf{S}(s_a^0) &= -2v_N(x-y)\Lambda \\ s_a^0(0, x, y) &= \text{sh}(2k)(0, x, y). \end{aligned} \quad (5.6)$$

*Then we have the estimate*

$$\|s_a^0\|_{L^\infty([0, T])L^2(dx dy)} + \|s_a^0\|_{L^2([0, T])L^6(dx)L^2(dy)} \lesssim 1 + T^2$$

*for some constant  $\kappa$  independent of  $T$ . Moreover, we also have*

$$\|\nabla_{x+y}^j s_a^0\|_{L^2(dx dy)} + \|\nabla_{x+y}^j s_a^0\|_{L^2([0, T])L^6(dx)L^2(dy)} \lesssim \exp(\kappa T) \quad (5.7)$$

*for  $j = 1, 2$ .*

**Lemma 5.7.** *Let  $b > 0$  and  $c(t)$  is the characteristic function on  $[0, T]$  where  $T > 1$ . Suppose we have*

$$\Lambda = \int_{-\infty}^{\infty} ds \, c(t-s) e^{i(t-s)\Delta_{x,y}} F(s) \quad (5.8)$$

*then it follows*

$$\|\Lambda\|_{X^b} \lesssim_b T \|F\|_{X^{b-1}}. \quad (5.9)$$

*Proof.* Let  $\mathcal{F}$  denote the spacetime Fourier transform, then we see that

$$\begin{aligned} &\mathcal{F} \left( \int_{-\infty}^{\infty} ds \, c(t-s) e^{i(t-s)\Delta_{x,y}} F(s) \right) \\ &= \int_{-\infty}^{\infty} dt \, e^{-it\tau} \int_{-\infty}^{\infty} ds \, c(t-s) e^{-i(t-s)(|\xi|^2 + |\eta|^2)} \widehat{F}(s) \\ &= \widehat{c}(\tau + |\xi|^2 + |\eta|^2) \widetilde{F}(\tau, \xi, \eta). \end{aligned}$$

In particular, we have the estimate

$$|\widehat{c}(\tau + |\xi|^2 + |\eta|^2)| \lesssim \left| \frac{\sin(\frac{1}{2}T(\tau + |\xi|^2 + |\eta|^2))}{\tau + |\xi|^2 + |\eta|^2} \right| \lesssim T \langle \tau + |\xi|^2 + |\eta|^2 \rangle^{-1}.$$

Then the result follows.  $\square$

To prove Proposition 5.6, it suffices to show the following proposition.

**Proposition 5.8.** *Suppose  $c(t)$  is the characteristic function of  $I = [0, T]$  and let*

$$E(t, x, y) = \int_{-\infty}^{\infty} c(t-s) e^{i(t-s)\Delta_{x,y}} (c(s) v_N(x-y) \Lambda(s)) \, ds. \quad (5.10)$$



Then we have the estimate

$$\begin{aligned} & \|E\|_{L^\infty(I)L^2(dxdy)} + \|E\|_{L^2(I)L^6(dx)L^2(dy)} \\ & \lesssim T \sup_z \left\| (|\partial_t|^{\frac{1}{4}+\varepsilon} + |\nabla_x|^{\frac{1}{2}+\varepsilon}) \Lambda(t, x, x+z) \right\|_{L^2(I)L^2(dx)}. \end{aligned} \quad (5.11)$$

It is helpful to rewrite

$$\begin{aligned} v_N(x-y)\Lambda(s) &= \int dz v_N(z) \delta(x-y-z) \Lambda\left(s, \frac{x+y+z}{2}, \frac{x+y-z}{2}\right) \\ &=: \int dz v_N(z) \delta(x-y-z) f_z(s, x+y). \end{aligned} \quad (5.12)$$

Then let us define

$$E_z(t, x, y) = \int_{-\infty}^{\infty} ds c(t-s) e^{i(t-s)\Delta} (c(s) \delta(x-y-z) f_z(s, x+y)) \quad (5.13)$$

and write

$$E(t, x, y) = \int dz v_N(z) E_z(t, x, y). \quad (5.14)$$

The following lemmas provide uniform in  $z$  estimates for  $E_z$ .

**Lemma 5.9.** *Fix  $z$  and  $\delta > 0$ . Then we have the estimates*

$$\left\| |\partial_t|^{\frac{1}{4}+\varepsilon} P_{|\xi-\eta| \gg |\tau+|\xi+\eta|^2|^{\frac{1}{2}}} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim_\delta T \left\| |\partial_t|^{\frac{1}{4}+\varepsilon} f_z \right\|_{L^2(I)L^2(dx)} \quad (5.15a)$$

$$\left\| |\nabla_x + \nabla_y|^{\frac{1}{2}+\varepsilon} P_{|\xi-\eta| \gg |\tau+|\xi+\eta|^2|^{\frac{1}{2}}} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim_\delta T \left\| |\nabla_x|^{\frac{1}{2}+\varepsilon} f_z \right\|_{L^2(I)L^2(dx)}. \quad (5.15b)$$

*Proof.* Applying Lemma 5.7 to  $|\partial_t|^{\frac{1}{4}+\varepsilon} P E_z$  yields

$$\left\| |\partial_t|^{\frac{1}{4}+\varepsilon} P_{1'} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim T \left\| |\partial_t|^{\frac{1}{4}+\varepsilon} (P_{1'} c \delta(x-y-z) f_z(s, x+y)) \right\|_{X^{-\frac{3}{4}-\delta}}$$

where  $P_{1'}$  is a short hand for the projection operator  $P_{|\xi-\eta| \gg |\tau+|\xi+\eta|^2|^{\frac{1}{2}}}$ .

Directly computing the spacetime Fourier transform of the RHS gives us

$$\begin{aligned} & \left\| |\partial_t|^{\frac{1}{4}+\varepsilon} (P_{1'} c \delta(x-y-z) f_z(s, x+y)) \right\|_{X^{-\frac{3}{4}-\delta}}^2 \\ & \lesssim \int_{|\xi-\eta| \gg |\tau+|\xi+\eta|^2|^{\frac{1}{2}}} d\tau d\xi d\eta \frac{|\tau|^{\frac{1}{2}+2\varepsilon} |\widetilde{c f_z}(\tau, \xi+\eta)|^2}{\langle \tau + |\xi+\eta|^2 + |\xi-\eta|^2 \rangle^{\frac{3}{2}+2\delta}} \\ & \lesssim \int d\tau d(\xi+\eta) |\tau|^{\frac{1}{2}+2\varepsilon} |\widetilde{c f_z}(\tau, \xi+\eta)|^2 \int_{2a}^{\infty} \frac{\rho^2 d\rho}{\langle \tau + |\xi+\eta|^2 + \rho^2 \rangle^{\frac{3}{2}+2\delta}} \end{aligned}$$

where  $a := |\tau + |\xi+\eta|^2|^{\frac{1}{2}}$ . Hence it remains to evaluate the integral in  $\rho$ . There are two cases: either  $\tau + |\xi+\eta|^2 > 0$  or  $\tau + |\xi+\eta|^2 < 0$ . Both cases

are essentially the same. In the latter case, we have

$$\int_{2a}^{\infty} \frac{\rho^2 d\rho}{(1 + |a^2 - \rho^2|)^{\frac{3}{2}+2\delta}} \lesssim \int_a^{\infty} \frac{\rho^2 d\rho}{(1 + \rho^2)^{\frac{3}{2}+2\delta}} \lesssim_{\delta} 1.$$

The proof of (5.15b) is similar.  $\square$

In the region  $|\xi - \eta| \lesssim a$ , we further subdivide into the cases  $\tau + |\xi + \eta|^2 > 0$  and  $\tau + |\xi + \eta|^2 < 0$ .

**Lemma 5.10.** *Fix  $z$  and  $\delta > 0$ . Then we have the estimate*

$$\left\| |\partial_t|^{\frac{1}{4}+\varepsilon} P_{\{|\xi-\eta|\lesssim a\}\cap\{\tau+|\xi+\eta|^2>0\}} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim T \left\| |\partial_t|^{\frac{1}{4}+\varepsilon} (cf_z) \right\|_{L^2(dtdx)}. \quad (5.16a)$$

$$\left\| |\nabla_x + \nabla_y|^{\frac{1}{2}+\varepsilon} P_{\{|\xi-\eta|\lesssim a\}\cap\{\tau+|\xi+\eta|^2>0\}} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim T \left\| |\nabla_x|^{\frac{1}{2}+\varepsilon} (cf_z) \right\|_{L^2(dtdx)}. \quad (5.16b)$$

*Proof.* The proof is similar to the proof of Lemma 5.9. It suffices to just bound the integral

$$\int_0^{2a} \frac{\rho^2 d\rho}{(1 + a^2 + \rho^2)^{\frac{3}{2}+2\delta}} \lesssim \int_0^{2a} \frac{\rho^2 d\rho}{(1 + a^2)^{\frac{3}{2}}} \lesssim 1.$$

Hence we have

$$\left\| |\partial_t|^{\frac{1}{4}+\varepsilon} P_{1''} E_z \right\|_{X^{\frac{1}{4}-\delta}} \lesssim T \left\| |\partial_t|^{\frac{1}{4}+\varepsilon} (cf_z) \right\|_{L^2(dtdx)}$$

where  $P_{1''} := P_{\{|\xi-\eta|\lesssim a\}\cap\{\tau+|\xi+\eta|^2>0\}}$ .  $\square$

**Lemma 5.11.** *Fix  $z$  and  $0 < \delta < \frac{1}{2}$ . Then we have the estimate*

$$\left\| P_{\{|\xi-\eta|\lesssim a\}\cap\{\tau+|\xi+\eta|^2<0\}} E_z \right\|_{X^{\frac{1}{2}+\delta}} \lesssim_{\delta} T \left\| (|\partial_t|^{\frac{1}{4}+\varepsilon} + |\nabla_x|^{\frac{1}{2}+\varepsilon}) (cf_z) \right\|_{L^2(dtdx)}. \quad (5.17)$$

*Proof.* Again, applying Lemma 5.7 yields

$$\| P_2 E_z \|_{X^{\frac{1}{2}+\delta}} \lesssim T \| P_2 c\delta(x - y - z) f_z(s, x + y) \|_{X^{-\frac{1}{2}+\delta}}$$

where  $P_2$  denotes the projection operator  $P_{\{|\xi-\eta|\lesssim a\}\cap\{\tau+|\xi+\eta|^2<0\}}$ .

Take the spacetime Fourier transform of the RHS gives us

$$\begin{aligned} & \| P_2 c\delta(x - y - z) f_z(s, x + y) \|_{X^{-\frac{1}{2}+\delta}}^2 \\ & \lesssim \int_{|\xi-\eta|\lesssim a} d\tau d\xi d\eta \frac{|\widetilde{cf_z}(\tau, \xi + \eta)|^2}{\langle \tau + |\xi + \eta|^2 + |\xi - \eta|^2 \rangle^{1-2\delta}} \\ & \lesssim \int d\tau d(\xi + \eta) |\widetilde{cf_z}(\tau, \xi + \eta)|^2 \int_0^{2a} \frac{\rho^2 d\rho}{(1 + |a^2 - \rho^2|)^{1-2\delta}}. \end{aligned}$$

Hence, it suffices to estimate the integral in  $\rho$ . Observe we have

$$\int_0^{2a} \frac{\rho^2 d\rho}{(1 + |a^2 - \rho^2|)^{1-2\delta}} \lesssim \int_0^a \frac{a^2 du}{(1 + au)^{1-2\delta}} \lesssim a^{1+4\delta}$$

which completes our proof of the lemma.  $\square$

*Proof of Proposition 5.8.* Write  $E = P_1 E + P_2 E$  where  $P_1 = P_{1'} + P_{1''}$ . Then, by Minkowski and Sobolev inequalities, we have

$$\begin{aligned} \|cE\|_{L^\infty(dt)L^2(dxdy)} &\lesssim \left( \int v \right) \sup_z \| |\partial_t|^{\frac{1}{4}+\varepsilon} c(t) P_1 E_z \|_{L^{4-\varepsilon}(dt)L^2(dxdy)} \\ &\quad + \left( \int v \right) \sup_z \| c(t) P_2 E_z \|_{L^\infty(dt)L^2(dxdy)}. \end{aligned}$$

Now, by Lemma 4.2-4.3 (c.f. Lemma 2.9 in [Tao06]), we have the estimate

$$\|cE\|_{L^\infty(dt)L^2(dxdy)} \lesssim \sup_z \| |\partial_t|^{\frac{1}{4}+\varepsilon} c P_1 E_z \|_{X^{\frac{1}{4}-\delta}} + \sup_z \| c P_2 E_z \|_{X^{\frac{1}{2}+\delta}}$$

where  $\delta = \frac{\varepsilon}{4(4-\varepsilon)}$ . Finally, combining Lemma 5.9-5.11, we obtain the desired estimate

$$\|cE\|_{L^\infty(dt)L^2(dxdy)} \lesssim T \sup_z \left\| |i\partial_t - \Delta_x|^{\frac{1}{4}+\varepsilon} c\Lambda(t, x, x+z) \right\|_{L^2(dtdx)}.$$

For the other endpoint, we use ‘‘Sobolev at an angle’’ (c.f. Lemma 8.3 in [GM19]) then it follows

$$\begin{aligned} \|cE\|_{L^2(dt)L^6(dx)L^2(dy)} &\lesssim \| \langle \nabla_x + \nabla_y \rangle^{\frac{1}{2}} c P_1 E \|_{L^2(dt)L^3(dx)L^2(dy)} \\ &\quad + \| c P_2 E \|_{L^2(dt)L^6(dx)L^2(dy)} \\ &\lesssim \| \langle \nabla_x + \nabla_y \rangle^{\frac{1}{2}} c P_1 E \|_{X^{\frac{1}{4}-\delta}} + \| c P_2 E \|_{X^{\frac{1}{2}+\delta}}. \end{aligned}$$

Then estimate (5.11) follows.  $\square$

Next, we include some potential terms from (5.1a) to (5.6). Let us start with the following lemma.

**Lemma 5.12.** *Consider the ‘‘potential’’ operator with kernel*

$$\begin{aligned} V(u)(x, y) &:= ((v_N * \text{diag } \Gamma)(x) + (v_N * \text{diag } \Gamma)(y))u(x, y) \\ &\quad + (v_N \Gamma) \circ u(x, y) + u \circ (v_N \Gamma)(x, y). \end{aligned} \quad (5.18)$$

Let us also define

$$\begin{aligned} (\partial_x^j V)(u) &:= ((v_N * \partial^j \text{diag } \Gamma)(x) + (v_N * \text{diag } \Gamma)(y))u \\ &\quad - v_N \Pi^j(\Gamma) \circ u - u \circ v_N \Gamma, \end{aligned} \quad (5.19a)$$

$$\begin{aligned} (\partial_y^j V)(u) &:= ((v_N * \text{diag } \Gamma)(x) + (v_N * \partial^j \text{diag } \Gamma)(y))u \\ &\quad - v_N \Gamma \circ u - u \circ v_N \Pi^j(\Gamma) \end{aligned} \quad (5.19b)$$

where  $\Pi^j(\Gamma)(x, y) := (\nabla_{x+y}^j \Gamma)(x, y)$ . Then, for  $j \geq 0$  and interval  $I = [T_0, T_1]$ , we have the following uniform in  $N$  estimates

$$\|(\partial^j V)(u)\|_{L^1(I)L^2(dxdy)} \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \|u\|_{L^2(I)L^6(dx)L^2(dy)}, \quad (5.20)$$

$$\|(\partial^j V)(u)\|_{L^2(I)L^{\frac{6}{5}}(dx)L^2(dy)} \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \|u\|_{L^\infty(I)L^2(dxdy)}. \quad (5.21)$$

*Proof.* It suffices to consider the case  $j = 1$ . Moreover, it also suffices to handle the terms  $(v_N * \partial^j \text{diag } \Gamma)(x)u$  and  $v_N \Pi^j(\Gamma) \circ u$ . For the first term, we apply Hölder, Young, and Sobolev inequalities to get

$$\begin{aligned} & \int_I dt \, \| (v_N * \partial^j \text{diag } \Gamma)(x)u \|_{L^2(dx dy)} \\ & \lesssim \int_I dt \, \| \nabla_x^{\frac{1}{2}} \text{diag}(\nabla_{x+y}^j \Gamma) \|_{L^2(dx)} \| u \|_{L^6(dx) L^2(dy)} \\ & \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \| u \|_{L^2(I) L^6(dx) L^2(dy)} \end{aligned}$$

For the other term, we have

$$\begin{aligned} & \int_I dt \, \| (v_N \Pi^j(\Gamma)) \circ u \|_{L^2(dx dy)} \\ & \lesssim \int_I dt \int dz \, v_N(z) \| \Pi^j(\Gamma)(x, x-z) \|_{L^3(dx)} \| u \|_{L^6(dx) L^2(dy)} \\ & \lesssim \int_I dt \int dz \, v_N(z) \| \nabla_x^{\frac{1}{2}} \Pi^j(\Gamma)(x, x-z) \|_{L^2(dx)} \| u \|_{L^6(dx) L^2(dy)} \\ & \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \| u \|_{L^2(I) L^6(dx) L^2(dy)} \end{aligned}$$

Thus, (5.20) follows immediately.

Similarly, by Hölder, Young, and Sobolev inequalities, we see that

$$\begin{aligned} & \| (v_N * \partial^j \text{diag } \Gamma)(x)u \|_{L^2(I) L^{\frac{6}{5}}(dx) L^2(dy)} \\ & \lesssim \| (v_N * \text{diag}(\nabla_{x+y}^j \Gamma)(x) \|_{L^2(I) L^3(dx)} \| u \|_{L^\infty(I) L^2(dx dy)} \\ & \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \| u \|_{L^\infty(I) L^2(dx dy)}. \end{aligned}$$

For the other term, we see that

$$\begin{aligned} & \| (v_N \Pi^j(\Gamma)) \circ u \|_{L^2(I) L^{\frac{6}{5}}(dx) L^2(dy)} \\ & \lesssim \int dz \, v_N(z) \| \Pi^j(\Gamma)(x, x-z)u(x-z, y) \|_{L^2(I) L^{\frac{6}{5}}(dx) L^2(dy)} \\ & \lesssim \mathbf{N}_I(\nabla_{x+y}^j \Gamma) \| u \|_{L^\infty(I) L^2(dx dy)}. \end{aligned}$$

Then (5.21) follows.  $\square$

**Proposition 5.13.** *Let  $s_a$  be a solution to*

$$\begin{aligned} \tilde{\mathbf{S}}(s_a) &= -2v_N(x-y)\Lambda \\ s_a(0, x, y) &= \text{sh}(2k)(0, x, y) \end{aligned} \tag{5.22}$$

where  $\tilde{\mathbf{S}} = \mathbf{S} + V$  and  $V$  as in the above lemma. Then the following estimate holds

$$\| s_a \|_{L^\infty([0, T]) L^2(dx dy)} \lesssim 1 + T^3. \tag{5.23}$$

Moreover, for every  $j \geq 1$ , there exists  $\kappa > 0$  such that we have the estimate

$$\| \nabla_{x+y}^j s_a \|_{L^2(dx dy)} \lesssim_j \exp(\kappa T). \tag{5.24}$$

*Proof.* The idea is to write  $s_a = s_a^0 + s_a^1$  where  $s_a^0$  is a solution to (5.6). Then we see that  $s_a^1$  satisfies the Cauchy problem

$$\begin{aligned}\tilde{\mathbf{S}}(s_a^1) &= -V(s_a^0) \\ s_a^1(0, x, y) &= 0.\end{aligned}\tag{5.25}$$

To obtain energy estimate for  $s_a^1$ , we employ the identity

$$\mathbf{S}_\pm(s_a^1 \circ \overline{s_a^1}) + V(s_a^1) \circ \overline{s_a^1} - s_a^1 \circ \overline{V(s_a^1)} = -V(s_a^0) \circ \overline{s_a^1} + s_a^1 \circ \overline{V(s_a^0)} \tag{5.26}$$

which means

$$\frac{d}{dt} \|s_a^1(t)\|_{L^2(dx dy)}^2 \leq \left| \text{Tr} \left( \mathbf{S}_\pm(s_a^1 \circ \overline{s_a^1}) + V(s_a^1) \circ \overline{s_a^1} - s_a^1 \circ \overline{V(s_a^1)} \right) \right|.$$

Then it follows that

$$\sup_{t \in [0, T]} \|s_a^1(t)\|_{L^2(dx dy)} \lesssim \int_0^T dt \|V(s_a^0(t))\|_{L^2(dx dy)}.$$

By Lemma 5.12, Proposition 4.8, and Proposition 5.6, we obtain (5.23).

The case of  $\nabla_{x+y}^j s_a$  is similar. We begin by writing the equation

$$\tilde{\mathbf{S}}(\nabla_{x+y}^j s_a^1) = [V, \nabla_{x+y}^j] s_a^1 - \nabla_{x+y}^j (V(s_a^0)). \tag{5.27}$$

Again, by energy method, we have the estimate

$$\begin{aligned}\|\nabla_{x+y}^j s_a^1\|_{L^\infty([0, T]) L^2(dx dy)} &\lesssim \int_0^T dt \left\| [V, \nabla_{x+y}^j] s_a^1 \right\|_{L^2(dx dy)} \\ &\quad + \int_0^T ds \left\| \nabla_{x+y}^j V(s_a^0) \right\|_{L^2(dx dy)}.\end{aligned}$$

The forcing term  $\nabla_{x+y}^j (V(s_a^0))$  is handled in the same manner as above. For the commutator term, we have the estimate

$$\int_0^T dt \left\| [V, \nabla_{x+y}^j] s_a^1 \right\|_{L^2(dx dy)} \lesssim \int_0^T dt \left\| (\partial^j V)(s_a^1) \right\|_{L^2(dx dy)}.$$

Hence, by Lemma 5.12 and (5.28) in the following corollary, we obtain the desired result.  $\square$

**Corollary 5.14.** *Suppose  $s_a$  is a solution to (5.22). Then the following estimate holds*

$$\|s_a\|_{L^2([0, T]) L^6(dx) L^2(dy)} \lesssim 1 + T^4. \tag{5.28}$$

Moreover, for  $j \geq 1$  we also have

$$\|\nabla_{x+y}^j s_a\|_{L^2([0, T]) L^6(dx) L^2(dy)} \lesssim_j \exp(\kappa T). \tag{5.29}$$

*Proof.* Let us once again reduce the problem to (5.25). By the endpoint Strichartz estimates, we have

$$\begin{aligned}\|s_a^1\|_{L^2([0, T]) L^6(dx) L^2(dy)} &\lesssim (\text{data}) + \|V(s_a^0)\|_{L^2([0, T]) L^{\frac{6}{5}}(dx) L^2(dy)} + \|V(s_a^1)\|_{L^2([0, T]) L^{\frac{6}{5}}(dx) L^2(dy)}.\end{aligned}$$

By Lemma 5.12, Proposition 5.6, and (5.23), we establish estimate (5.28). The proof for the higher derivatives are similar.  $\square$

*Proof of Theorem 5.2.* Let us write  $\text{sh}(2k) := s_2 = s_a + s_e$  where  $s_a$  satisfies (5.22). Then we have the coupled system

$$\tilde{\mathbf{S}}(s_e) = -(v_N \Lambda) \circ p_2 - \bar{p}_2 \circ (v_N \Lambda) \quad (5.30a)$$

$$\begin{aligned} \tilde{\mathbf{S}}_{\pm}(\bar{p}_2) &= -(v_N \Lambda) \circ \bar{s}_a + s_a \circ (v_N \Lambda) \\ &\quad - (v_N \Lambda) \circ \bar{s}_e + s_e \circ (v_N \Lambda) \\ &=: M - (v_N \Lambda) \circ \bar{s}_e + s_e \circ (v_N \Lambda) \end{aligned} \quad (5.30b)$$

where  $\tilde{\mathbf{S}}_{\pm} = \mathbf{S}_{\pm} + V_{\pm}$  and

$$V_{\pm}(p) = ((v_N * \text{diag } \Gamma)(x) - (v_N * \text{diag } \Gamma)(y))p + (v_N \Gamma) \circ p - p \circ (v_N \Gamma).$$

Let  $I = [0, T]$ . By Strichartz estimates, we have

$$\begin{aligned} \|p_2\|_{L^\infty(I)L^2(dx dy)} &\lesssim (\text{data}) + \|V_{\pm}(p_2)\|_{L^2(I)L^{\frac{6}{5}}(dx)L^2(dy)} + \|M\|_{L^2(I)L^{\frac{6}{5}}(dx)L^2(dy)} \\ &\quad + \left( \sup_z \int_I dt \|\Lambda(t, x - z, x - z)\|_{L^3(dx)}^2 \|s_e(t)\|_{L^2(dx dy)}^2 \right)^{\frac{1}{2}}. \end{aligned}$$

Applying Lemma 5.12, we have that

$$\begin{aligned} \|p_2\|_{L^\infty(I)L^2(dx dy)} &\lesssim (\text{data}) + T^4 \\ &\quad + \mathbf{N}_T(X) (\|p_2\|_{L^\infty(I)L^2(dx dy)} + \|s_e\|_{L^\infty(I)L^2(dx dy)}) \end{aligned} \quad (5.31)$$

since  $\|M\|_{L^2([0, T])L^{\frac{6}{5}}(dx)L^2(dy)} \lesssim 1 + T^4$ . Likewise, we see that

$$\begin{aligned} \|s_e\|_{L^\infty(I)L^2(dx dy)} &\lesssim (\text{data}) + \|V(s_e)\|_{L^2(I)L^{\frac{6}{5}}(dx)L^2(dy)} \\ &\quad + \left( \sup_z \int_I dt \|\Lambda(t, x - z, x - z)\|_{L^3(dx)}^2 \|p_2(t)\|_{L^2(dx dy)}^2 \right)^{\frac{1}{2}} \\ &\lesssim (\text{data}) + \mathbf{N}_T(X) (\|p_2\|_{L^\infty(I)L^2(dx dy)} + \|s_e\|_{L^\infty(I)L^2(dx dy)}) \end{aligned}$$

Next, define

$$E(t) = \|s_e\|_{L^\infty([0, t])L^2(dx dy)} + \|p_2\|_{L^\infty([0, t])L^2(dx dy)}.$$

Putting the estimates for  $p_2$  and  $s_e$  together, we have

$$E(T) \lesssim (\text{data}) + T^\alpha + \mathbf{N}_T(X)E(T), \quad (5.32)$$

where  $\alpha = 4$ . Then we will apply the above estimate on small intervals together with Proposition 4.8. For convenience, we use  $C_1 > 0$  to represent the implicit constant in the above estimate and we use  $C_2 > 0$  to represent the implicit constant in the estimate of Proposition 4.8. Here we divide the time interval  $[0, T]$  into  $l$  consequent subintervals  $\{I_i\}_i$  where  $I_i = [T_i, T_{i+1}]$  satisfying  $N_{I_i}(X) \leq \frac{1}{2C_1}$  and  $l \sim 2C_1 C_2 T$ .

For a certain interval (without loss of generality, we consider the first interval as an example), we have

$$E(T_2) \leq C_1 E(T_1) + C_1 + C_1 \cdot \frac{1}{2C_1} E(T_2). \quad (5.33)$$

This implies,

$$E(T_2) \leq 2C_1 E(T_1) + 2C_1. \quad (5.34)$$

Then we can iterate the arguments on all of the intervals, noticing the number of intervals are around  $l \sim 2C_1 C_2 T$ , we have, for some constant  $\kappa$ ,

$$\begin{aligned} E(T) = E(T_l) &\leq 2C_1 E(T_{l-1}) + 2C_1 \leq 2C_1 (2C_1 E(T_{l-2}) + 2C_1) + 2C_1 \\ &\leq \dots \leq (2C_1)^{2C_1 C_2 T} E(T_1) + \frac{(2C_1)^{2C_1 C_2 T+1} - 1}{2C_1 - 1} \lesssim e^{\kappa T}. \end{aligned}$$

The proof of (5.2b) follows from (5.2a).  $\square$

Let us complete this section with the proof of Corollary 5.4. We start by proving the following lemma.

**Lemma 5.15.** *Suppose  $(\phi(t), k(t))$  solves (2.30). Then there exists some  $\kappa > 0$  such that following estimates*

$$\| \tilde{\mathbf{W}}(\text{ch}(2k_t)) \|_{L^2(dx dy)} \lesssim \exp(\kappa T) \quad (5.35a)$$

$$\| \tilde{\mathbf{W}}(\text{ch}(k_t)) \|_{L^2(dx dy)} \lesssim \exp(\kappa T) \quad (5.35b)$$

holds for all  $t \in [0, T]$  and independent of  $N$ .

*Proof.* By (2.30c), Theorem 5.2, and Proposition 4.12, we have the estimate

$$\| \tilde{\mathbf{W}}(\text{ch}(2k)) \|_{L^2(dx dy)} \lesssim \mathbf{N}_T(\nabla_{x+y}^2 \Lambda) \| \text{sh}(2k) \|_{L^\infty([0, T]) L^2(dx dy)} \lesssim \exp(\kappa T).$$

To estimate  $\tilde{\mathbf{W}}(\text{ch}(k))$ , we invoke a variant of identity (30) in [GMM11], which is an immediate consequence of holomorphic functional calculus,

$$\tilde{\mathbf{W}}(\text{ch}(k)) = \frac{1}{4\pi i} \int_{\gamma} (u \circ \bar{u} - z)^{-1} \tilde{\mathbf{W}}(\text{ch}(2k)) (u \circ \bar{u} - z)^{-1} \sqrt{1+z} dz \quad (5.36)$$

where the contour  $\gamma$  encloses the spectrum of  $u \circ \bar{u}$  and contains inside the ball of radius  $C \|u\|_{L^2(dx dy)}^2$  for some  $C > 0$ , i.e.  $|z| \leq C \|u\|_{L^2(dx dy)}^2$ . Since  $(u \circ \bar{u} - z)^{-1}$  has bounded operator norm, then it follows from Corollary 5.3

$$\begin{aligned} \| \tilde{\mathbf{W}}(\text{ch}(k)) \|_{L^2(dx dy)} &\lesssim |z|^{\frac{3}{2}} \| (u \circ \bar{u} - z)^{-1} \|_{\text{op}}^2 \| \tilde{\mathbf{W}}(\text{ch}(2k)) \|_{L^2(dx dy)} \\ &\lesssim \exp(\alpha T) \end{aligned}$$

for some constant  $\alpha > 0$ .  $\square$

*Proof of Corollary 5.4.* Rewrite (5.30b) with  $p_2 = 2\bar{u} \circ u = 2p \circ p + 4p$  in the differential operator yields

$$\begin{aligned} \tilde{\mathbf{W}}(p) &= -\frac{1}{2} \tilde{\mathbf{W}}(p \circ p) - \frac{1}{4} [v_N \Lambda, \overline{\text{sh}(2k)}] \\ &= -\frac{1}{2} \tilde{\mathbf{W}}(p) \circ p - \frac{1}{2} p \circ \tilde{\mathbf{W}}(p) - \frac{1}{4} [v_N \Lambda, \overline{\text{sh}(2k)}]. \end{aligned} \quad (5.37)$$

Then, by the endpoint Strichartz estimates, we see that

$$\begin{aligned}
& \|p\|_{L^2([0,T])L^6(dx)L^2(dy)} \\
& \lesssim \|V_{\pm}(p)\|_{L^1([0,T])L^2(dxdy)} + \|\tilde{\mathbf{W}}(p) \circ p\|_{L^1([0,T])L^2(dxdy)} \\
& \quad + \|v_N \Lambda \circ \overline{\text{sh}(2k)}\|_{L^2([0,T])L^{\frac{6}{5}}(dx)L^2(dy)} \\
& \lesssim T \dot{\mathbf{N}}_T(\Gamma) \|p\|_{L^\infty([0,T])L^2(dxdy)} + T \|\tilde{\mathbf{W}}(p)\|_{L^\infty([0,T])L^2(dxdy)} \|p\|_{L^\infty([0,T])L^2(dxdy)} \\
& \quad + \mathbf{N}_T(\Lambda) \|\text{sh}(2k)\|_{L^\infty([0,T])L^2(dxdy)}.
\end{aligned}$$

Hence the desired estimate follows from Proposition 4.8, Theorem 5.2, Corollary 5.3 and Lemma 5.15.

Next, since  $\text{sh}(2k) = 2u + 2\bar{p} \circ u$ , then it follows

$$\begin{aligned}
\|u\|_{L^2([0,T])L^6(dx)L^2(dy)} & \lesssim \|\text{sh}(2k)\|_{L^2([0,T])L^6(dx)L^2(dy)} \\
& \quad + \|p\|_{L^2([0,T])L^6(dx)L^2(dy)} \|u\|_{L^\infty([0,T])L^2(dxdy)}
\end{aligned}$$

which again yields the desired estimate.  $\square$

## 6. FOCK SPACE ESTIMATES

The purpose of this section is to obtain global estimates for the error

$$\left\| \psi_{\text{exact}}(t) - e^{i \int X_0(t) dt} \psi_{\text{appr}}(t) \right\|_{\mathcal{F}}. \quad (6.1)$$

Using the identity

$$\left\| \psi_{\text{exact}}(t) - e^{i \int X_0(t) dt} \psi_{\text{appr}}(t) \right\|_{\mathcal{F}} = \left\| e^{-i \int X_0(t) dt} \psi_{\text{red}}(t) - \Omega \right\|_{\mathcal{F}}, \quad (6.2)$$

it suffices to study the error of the reduced dynamics which we denote by  $E := e^{-i \int X_0(t) dt} \psi_{\text{red}}(t) - \Omega$ .

Recall  $E$  satisfies the equation

$$\left( \frac{1}{i} \frac{\partial}{\partial t} - \mathcal{H}_{\text{red}} + X_0 \right) E = (0, 0, 0, X_3, X_4, 0, \dots) =: \tilde{X} \quad (6.3)$$

with initial condition  $E(0, \cdot) = 0$ . For convenience, we adopt the notation

$$\mathbf{S}_F := \frac{1}{i} \frac{\partial}{\partial t} - \mathcal{H}_{\text{red}} + X_0 =: \mathbf{S}_D - \mathcal{P} \quad (6.4)$$

where

$$\mathbf{S}_D := \frac{1}{i} \frac{\partial}{\partial t} - \int a_x^* \Delta a_x - \frac{1}{2N} \int dxdy \{v_N(x-y) a_x^* a_y^* a_x a_y\}. \quad (6.5)$$

Here we see that  $\mathcal{P}$  accounts for all the remaining terms, that is,  $\mathcal{P} = \mathcal{H}_{\text{red}} - X_0 - \mathcal{H}$ . Note that  $\mathcal{P}$  is not a diagonal Fock space operator.



**6.1. Explicit Form of  $\mathcal{P}$ .** Let us write down  $\mathcal{H}_{\text{red}}$  and  $\mathcal{P}$  in order to study (6.3). The reduced Hamiltonian was first calculated in section 5 of [GM13b] and was later rewritten using Wick's Theorem in section 7 of [GM17]. Let first us recall the conjugation relations

$$\begin{aligned} e^{\mathcal{B}(k)} a_x e^{-\mathcal{B}(k)} &= \int dy \{ \overline{\text{ch}(k)}(x, y) a_y + \text{sh}(k)(x, y) a_y^* \} \\ &= a(\overline{\text{ch}(k)}(x, \cdot)) + a^*(\text{sh}(k)(x, \cdot)) =: b_x \end{aligned} \quad (6.6a)$$

$$\begin{aligned} e^{\mathcal{B}(k)} a_x^* e^{-\mathcal{B}(k)} &= \int dy \{ \overline{\text{sh}(k)}(x, y) a_y + \text{ch}(k)(x, y) a_y^* \} \\ &= a(\overline{\text{sh}(k)}(x, \cdot)) + a^*(\text{ch}(k)(x, \cdot)) =: b_x^*. \end{aligned} \quad (6.6b)$$

We will also adopt the notation  $a_x^*(u) := a^*(\text{sh}(k)(x, \cdot))$  and  $a_x(\bar{u}) := a(\overline{\text{sh}(k)}(x, \cdot))$ . Then the reduced Hamiltonian is given by

$$\mathcal{H}_{\text{red}} = -N\mu(t) \quad (6.7a)$$

$$+ N^{\frac{1}{2}} \int dx \{ \tilde{h}(\phi(t, x)) b_x^* + \bar{\tilde{h}}(\phi(t, x)) b_x \} \quad (6.7b)$$

$$- \mathcal{H}_g - \text{Nor} \left\{ \mathcal{I} \begin{pmatrix} w^T & \bar{f} \\ -f & -w \end{pmatrix} \right\} \quad (6.7c)$$

$$- \frac{1}{\sqrt{N}} \text{Nor} \left\{ \int dx dy v_N(x-y) (\bar{\phi}(y) b_x^* b_x b_y + \phi(y) b_y^* b_x^* b_x) \right\} \quad (6.7d)$$

$$- \frac{1}{2N} \text{Nor} \left\{ \int dx dy v_N(x-y) b_x^* b_y^* b_x b_y \right\} \quad (6.7e)$$

where  $\text{Nor}\{\cdot\}$  refers to the normal ordering the the operators with respect to  $(a, a^*)$ . In the linear term (6.7b),  $\tilde{h}$  corresponds to the modified Hartree operator given by

$$\tilde{h}(\phi(t, x)) := \mathbf{S}\phi_t(x_1) + \int dy \left\{ (v_N \Lambda_t)(x_1, y) \bar{\phi}_t(y) + \frac{1}{N} \tilde{\alpha}^T(t, x_1, y) \phi_t(y) \right\}.$$

For the quadratic term (6.7c),  $\mathcal{I}(\cdot)$  is the Lie algebra isomorphism between symplectic matrices of integral operators and quadratic polynomials in  $(a, a^*)$  used in [GMM10, GMM11, GM13b], which is given by

$$\mathcal{I} \begin{pmatrix} w^T & \bar{f} \\ -f & -w \end{pmatrix} = -\frac{1}{2} \int dx dy \{ w(y, x) a_x a_y^* + w(x, y) a_x^* a_y - f(x, y) a_x^* a_y^* - \bar{f}(x, y) a_x a_y \} \quad (6.8)$$

where

$$f := \left( \tilde{\mathbf{S}}(\text{sh}(k)) + \overline{\text{ch}(k)} \circ (v_N \Lambda) \right) \circ \text{ch}(k) - \left( \tilde{\mathbf{W}}(\overline{\text{ch}(k)}) - \text{sh}(k) \circ \overline{(v_N \Lambda)} \right) \circ \text{sh}(k) \quad (6.9a)$$

$$w := \left( \tilde{\mathbf{W}}(\overline{\text{ch}(k)}) - \text{sh}(k) \circ \overline{(v_N \Lambda)} \right) \circ \overline{\text{ch}(k)} - \left( \tilde{\mathbf{S}}(\text{sh}(k)) + \overline{\text{ch}(k)} \circ (v_N \Lambda) \right) \circ \overline{\text{sh}(k)} \quad (6.9b)$$

and

$$\mathcal{H}_g = \int dxdy g(x, y) a_x^* a_y \quad (6.10)$$

where  $g$  is as in (2.42a). In particular, equations (2.21) correspond to  $\tilde{h}(\phi(t, x)) = 0$  and  $f = 0$ ; see Theorem 7.1 in [GM13a]. Hence it follows that

$$\mathcal{P} = -\mathcal{H}_g - \mathcal{H}_0 + \text{Nor} \left\{ \mathcal{I} \begin{pmatrix} w^T & 0 \\ 0 & -w \end{pmatrix} \right\} \quad (6.11a)$$

$$- \frac{1}{\sqrt{N}} \text{Nor} \left\{ \int dxdy v_N(x-y) (\bar{\phi}(y) b_x^* b_x b_y + \phi(y) b_y^* b_x^* b_x) \right\} \quad (6.11b)$$

$$- \frac{1}{2N} \text{Nor} \left\{ \int dxdy v_N(x-y) (b_x^* b_y^* b_x b_y - a_x^* a_y^* a_x a_y) \right\}. \quad (6.11c)$$

It is convenient to list out the terms in the normal ordering. Let us start with quadratic terms  $\mathcal{Q} := -(6.11a)$ . Using the fact that  $f = 0$ , we can rewrite  $\mathcal{Q}$  as follows

$$\mathcal{Q} = \int dxdy \{ (v_N * \rho_\Gamma)(t, x) \delta(x-y) - (v_N \Gamma)(t, x, y) \} a_x^* a_y \quad (6.12a)$$

$$\int dxdy w(x, y) a_x^* a_y \quad (6.12b)$$

where

$$w = \left( \tilde{\mathbf{W}}(\overline{\text{ch}(k)}) - \text{sh}(k) \circ \overline{(v_N \Lambda)} \right) \circ \overline{\text{ch}(k)}^{-1}. \quad (6.13)$$

Next, we write down the cubic terms  $\mathcal{C} := -(6.11b)$  into  $\mathcal{C} = \mathcal{C}_1 + \mathcal{C}_1^*$  where

$$\mathcal{C}_1 = \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x(\bar{u}) a_x(\bar{c}) a_y(\bar{c}) \quad (6.14a)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(c) a_x(\bar{c}) a_y(\bar{c}) \quad (6.14b)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(u) a_x^*(c) a_y(\bar{c}) \quad (6.14c)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(c) a_y^*(u) a_x(\bar{c}) \quad (6.14d)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(u) a_y^*(u) a_x^*(c) \quad (6.14e)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(u) a_y^*(u) a_x(\bar{u}) \quad (6.14f)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_x^*(u) a_x(\bar{u}) a_y(\bar{c}) \quad (6.14g)$$

$$+ \frac{1}{\sqrt{N}} \int dxdy v_N(x-y) \bar{\phi}(y) a_y^*(u) a_x(\bar{u}) a_x(\bar{c}), \quad (6.14h)$$

and  $\mathcal{C}_1^*$  denotes the adjoint operator of  $\mathcal{C}_1$ . Similarly, the quartic terms (6.11c) contains  $2^4 + 1$  normal ordered terms which we write (6.7e) =  $\mathcal{D} = \mathcal{D}_1 + \mathcal{D}_1^* + \mathcal{D}_2$  where

$$\mathcal{D}_1 = \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_y^*(c) a_x^*(u) a_y^*(u) \quad (6.15a)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_y^*(c) a_x^*(u) a_y(\bar{c}) \quad (6.15b)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_y^*(c) a_y^*(u) a_x(\bar{c}) \quad (6.15c)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_x^*(u) a_y^*(u) a_y(\bar{u}) \quad (6.15d)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(u) a_y^*(c) a_y^*(u) a_x(\bar{u}) \quad (6.15e)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_x^*(u) a_y(\bar{c}) a_y(\bar{u}) \quad (6.15f)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(u) a_y^*(c) a_y(\bar{c}) a_x(\bar{u}) \quad (6.15g)$$

and

$$\mathcal{D}_2 = \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(u) a_y^*(u) a_x(\bar{u}) a_y(\bar{u}) \quad (6.15h)$$

$$+ \frac{1}{2N} \int dxdy v_N(x-y) a_x^*(c) a_y^*(c) a_x(\bar{c}) a_y(\bar{c}) \quad (6.15i)$$

$$- \frac{1}{2N} \int dxdy v_N(x-y) a_x^* a_y^* a_x a_y. \quad (6.15j)$$

**6.2. Proof of Theorem 2.8.** Let us first study the solution to

$$\mathbf{S}_D u = f, \quad u(0, \cdot) = 0, \quad (6.16)$$

where  $u, f$  are Fock vectors. Since  $\mathbf{S}_D$  is a diagonal operator then it follows that (6.16) reduces to an inhomogeneous Schrödinger equation with potential on each sector. In order to study (6.16), we need to introduce Strichartz and dual Strichartz norm for Fock vectors.

Let  $T > 0$  and define the Strichartz norm on the time interval  $[0, T]$  by

$$\|u\|_S = \max \left\{ \|u\|_{L^\infty(dt) L^2(dx_1 \dots dx_n)}, \|u\|_{L^2(dt) L^6(d(x_1-x_2)) L^2(d(x_1+x_2) \dots dx_n)}, \right. \\ \left. \|u\|_{L^2(dt) L^3(d(x_1-x_2) d(x_1+x_2)) L^2(dx_3 \dots dx_n)}, \text{ and all other permutations} \right\}$$

and the dual Strichartz norm

$$\|u\|_{S'} = \min \left\{ \|u\|_{L^1(dt) L^2(dx_1 \dots dx_n)}, \|u\|_{L^2(dt) L^{\frac{6}{5}}(d(x_1-x_2)) L^2(d(x_1+x_2) \dots dx_n)}, \right. \\ \left. \|u\|_{L^2(dt) L^{\frac{3}{2}}(d(x_1-x_2) d(x_1+x_2)) L^2(dx_3 \dots dx_n)}, \text{ and all other permutations} \right\}.$$

Also, let  $X \in \mathcal{F}$  be a Fock vector such that all but finitely many of the components are zeros, say  $X_0, X_1, \dots, X_k$  are nonzero. Then we define the Strichartz norm on  $\mathcal{F}$  by

$$\|X\|_S = \max\{|X_0|, \|X_1\|_S, \dots, \|X_k\|_S\},$$

and similarly for the dual Strichartz norm

$$\|X\|_{S'} = \max\{|X_0|, \|X_1\|_{S'}, \dots, \|X_k\|_{S'}\}.$$

By standard arguments, we have the following lemma.

**Lemma 6.1.** *Let  $f$  be a Fock vector with zero entries past the  $k$ th sector. Assume  $u$  is a solution to*

$$\mathbf{S}_D u = f, \quad u(0, \cdot) = 0 \quad (6.17)$$

*then we have the estimate*

$$\|u\|_S \lesssim \|f\|_{S'}. \quad (6.18)$$

*In particular, it follows that  $\sup_{t \in [0, T]} \|u\|_{\mathcal{F}} \lesssim \|u\|_S$ .*

*Sketch of Proof.* Note for  $0 < \beta < 1$ , we see that

$$\begin{aligned} \frac{1}{N} \|v_N(x_1 - x_2)u\|_{S'} &\lesssim \frac{1}{N} \|v_N(x_1 - x_2)u\|_{L^2(dt)L^{\frac{6}{5}}(d(x_1-x_2))L^2(d(x_1+x_2)\cdots dx_n)} \\ &\lesssim \frac{1}{N} \|v_N\|_{L^{\frac{3}{2}}} \|u\|_S \lesssim N^{\beta-1} \|u\|_S. \end{aligned}$$

Hence the potential can be treated as a perturbation of the free Schrödinger case.  $\square$

The following is the main proposition of this section.

**Proposition 6.2.** *Let  $0 < \beta < 1$  and let  $E(t)$  be the solution to (6.16). Assume  $(\Lambda(t), \Gamma(t), \phi(t))$  solves the time-dependent HFB system. Then there exists  $\kappa = \kappa(\beta, \varepsilon)$  such that*

$$\|E(t)\|_{\mathcal{F}} \lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) \quad (6.19)$$

*for all  $t \in [0, T]$  when  $N$  is sufficiently large.*

*Remark 6.3.* The author believes it is more beneficial to first consider the heuristics behind the main argument of Proposition 6.2 before proceeding with the proof. The main idea behind the proof of the proposition is the fact that  $\mathcal{P}$  can be split into a sum  $\mathcal{P}_1 + \mathcal{P}_2$  where  $\mathcal{P}_2$  is a bounded operator on Fock space (for the first  $k$  sector) and  $\mathcal{P}_1$  satisfies  $\|\mathcal{P}_1 E\|_{\mathcal{F}} \lesssim N \exp(\kappa T) \|E\|_{\mathcal{F}}$ , which is the main obstacle we need to overcome, but, fortunately, we also have  $\|\mathcal{P}_1 E\|_{S'} \lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) \|E\|_S$ .

However, since  $\mathbf{S}_F$  is not a diagonal operator, it's not immediately obvious how one would solve (6.16) and estimate the solution. Nevertheless, we begin by considering

$$\mathbf{S}_D E_1 = \tilde{X}, \quad E_1(0, \cdot) = 0, \quad (6.20)$$

which can be readily solve using Lemma 6.1 and propositions in the following section. Also, note we have the estimate  $\|E_1\|_S \lesssim \|\tilde{X}\|_{S'} \lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T)$  since  $\frac{1}{\sqrt{N}} \|v_N\|_{\frac{6}{5}} \lesssim N^{\frac{\beta-1}{2}}$ . With  $E_1$ , we can construct

$$\begin{aligned} E_\infty &= E_1 + \mathbf{S}_D^{-1} \mathcal{P}_1 E_1 + (\mathbf{S}_D^{-1} \mathcal{P}_1)^2 E_1 + \dots + (\mathbf{S}_D^{-1} \mathcal{P}_1)^n E_1 + \dots \\ &= (\mathbf{S}_D - \mathcal{P}_1)^{-1} \tilde{X} \end{aligned}$$

which solves

$$(\mathbf{S}_D - \mathcal{P}_1) E_\infty = \tilde{X}, \quad E_\infty(0, \cdot) = 0. \quad (6.21)$$

Moreover, we see that

$$\begin{aligned} \|E_\infty\|_{\mathcal{F}} &\lesssim \|E_1\|_{\mathcal{F}} + \|\mathbf{S}_D^{-1} \mathcal{P}_1 E_1\|_{\mathcal{F}} + \dots + \|(\mathbf{S}_D^{-1} \mathcal{P}_1)^n E_1\|_{\mathcal{F}} + \dots \\ &\lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) + C(N^{\frac{\beta-1}{2}} \exp(\kappa T))^2 + \dots \\ &\lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) \end{aligned}$$

for some  $C > 0$  when  $N$  is sufficiently large. Finally, suppose  $E(t)$  is a solution to (6.3) then we see that  $E - E_\infty$  satisfies

$$(\mathbf{S}_D - \mathcal{P})(E - E_\infty) = \mathcal{P}_2 E_\infty. \quad (6.22)$$

By energy estimates, we see that

$$\begin{aligned} \|E(t) - E_\infty(t)\|_{\mathcal{F}} &\lesssim \int_0^t ds \|\mathcal{P}_2 E_\infty(s)\|_{\mathcal{F}} \\ &\lesssim \int_0^t ds \|E_\infty(s)\|_{\mathcal{F}} \lesssim N^{\frac{\beta-1}{2}} \exp(\alpha T) \end{aligned}$$

if  $\|\mathcal{P}_2 E_\infty(t)\|_{\mathcal{F}} \lesssim \|E_\infty(t)\|_{\mathcal{F}}$ . Then the result follows from the triangle inequality. Thus, to establish Proposition 6.2, it suffices to prove  $\|\mathcal{P}_2 E_\infty(t)\|_{\mathcal{F}} \lesssim \|E_\infty(t)\|_{\mathcal{F}}$ . Unfortunately, the claim is false as stated since we can only show that  $\mathcal{P}_2$  is a bounded operator on the first  $k$  sectors of  $\mathcal{F}$  not the entire  $\mathcal{F}$ .

To fix this problem, the proof will only consider the  $M$ -truncation (iteration) of the series  $E_\infty$ , i.e. we consider

$$E_M := E_1 + \mathbf{S}_D^{-1} \mathcal{P}_1 E_1 + (\mathbf{S}_D^{-1} \mathcal{P}_1)^2 E_1 + \dots + (\mathbf{S}_D^{-1} \mathcal{P}_1)^{M-1} E_1, \quad (6.23)$$

where  $M$  depends on  $\beta$ .

*Proof of Proposition 6.2.* Let  $0 < \beta < 1$ . Choose  $\alpha$  so that Theorem 3.3 in [GM19] holds and  $M \geq \lceil \frac{2}{1-\beta} \rceil + 1$ .

Following Remark 6.3, we let  $E_1$  be the solution to (6.20) then consider

$$E_M = E_1 + \mathbf{S}_D^{-1} \mathcal{P}_1 E_1 + (\mathbf{S}_D^{-1} \mathcal{P}_1)^2 E_1 + \dots + (\mathbf{S}_D^{-1} \mathcal{P}_1)^{M-1} E_1$$

which have at most  $k = 5 + 4 \times M$  nonzero sectors since  $\mathcal{P}$  is a fourth-order operator in  $a^*, a$ . Next, observe  $E_M$  satisfies

$$\mathbf{S}_D(I - \mathbf{S}_D^{-1} \mathcal{P}_1) E_M = \tilde{X} - \mathcal{P}_1 (\mathbf{S}_D^{-1} \mathcal{P}_1)^{M-1} E_1 \quad (6.24)$$

then, by Strichartz estimates, we have that

$$\begin{aligned} \|(I - \mathbf{S}_D^{-1}\mathcal{P}_1)E_M(t)\|_{\mathcal{F}} &\lesssim \|\tilde{X}\|_{S'} + \|\mathcal{P}_1(\mathbf{S}_D^{-1}\mathcal{P}_1)^{M-1}E_1\|_{S'} \\ &\lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) \end{aligned} \quad (6.25)$$

when  $N$  is sufficiently large. Using reverse triangle inequality on the LHS of (6.25), it follows

$$\|E_M(t)\|_{\mathcal{F}} \lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T). \quad (6.26)$$

Moreover, let  $E(t)$  be a solution to (6.3) then it follows  $E - E_M$  satisfies the equation

$$(\mathbf{S}_D - \mathcal{P})(E - E_M) = \mathcal{P}_2 E_{M-1} + \mathcal{P}(\mathbf{S}_D^{-1}\mathcal{P}_1)^{M-1}E_1. \quad (6.27)$$

Using energy estimate yields

$$\begin{aligned} \|E(t) - E_M(t)\|_{\mathcal{F}} &\lesssim \int_0^t ds \{ \|\mathcal{P}_2 E_M(s)\|_{\mathcal{F}} + \|\mathcal{P}(\mathbf{S}_D^{-1}\mathcal{P}_1)^{M-1}E_1(s)\|_{\mathcal{F}} \} \\ &\lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) + N^{1+M} N^{\frac{\beta-1}{2}} \exp(\kappa' T). \end{aligned}$$

Again, by the reverse triangle inequality, we arrive at the desired conclusion. In particular, we have proven Theorem 2.8.  $\square$

## 7. DISPERSIVE ESTIMATES FOR $\mathcal{P}$

In this section we obtain estimates for the Fock space operator  $\mathcal{P}$  which was used in the previous section. We start by splitting  $\mathcal{P}$  into  $\mathcal{P}_1 + \mathcal{P}_2$  and show that  $\mathcal{P}_2$  has uniform in  $N$  bounded operator norm (but grows as a function of  $T$ ) whereas  $\mathcal{P}_1$  can be controlled uniformly in  $N$  by Strichartz estimates on Fock space but has operator norm that grows in  $N$ . More precisely, we let  $\mathcal{P}_1 = \mathcal{C} + \mathcal{D}$  and  $\mathcal{P}_2 = \mathcal{Q}$ .

Let us consider estimates for  $\mathcal{P}_1$ . The proof is similar to the prove given in [GM17].

**Proposition 7.1.** *Fix  $k \in \mathbb{N}$ . If  $X$  is a Fock space vector that has nonzero entries only in the first  $k$  sectors, then we have the estimates*

$$\|\mathcal{C}X\|_{S'} \lesssim_k N^{\frac{\beta-1}{2}} \exp(\kappa T) \|X\|_S \quad (7.1a)$$

$$\|\mathcal{C}X\|_{\mathcal{F}} \lesssim_k N \exp(\kappa T) \|X\|_{\mathcal{F}}. \quad (7.1b)$$

*Proof.* Write  $\text{ch}(k) = \delta + p$ . Then letting (6.14a) acts on the Fock vector  $X = (0, \dots, 0, F(x_1, \dots, x_{n+3}), 0, 0, \dots)$  gives us the vector with

$$((6.14a)X)_n \sim \int dx dy dz \frac{v_N(x-y)}{\sqrt{N}} \bar{\phi}(y) \bar{u}(x, z) F(x, y, z, x_1, \dots, x_n) \quad (7.2a)$$

$$+ \int dx dy dz dx' \frac{v_N(x-y)}{\sqrt{N}} \bar{\phi}(y) \bar{u}(x, z) \bar{p}(x, x') F(y, z, x', \dots) \quad (7.2b)$$

$$+ \int dx dy dz dx' dy' \frac{v_N(x-y)}{\sqrt{N}} \bar{\phi}(y) \bar{u}(x, z) \bar{p}(x, x') \bar{p}(y, y') F(x', y', z, \dots) \quad (7.2c)$$

+ similar terms

on the  $n$ th sector and zero elsewhere.

Let us focus on establishing (7.1a) for the first term. Since  $x_1, \dots, x_n$  play only passive roles in the estimate, let us temporarily suppress them in the notation. We begin by writing  $F(x, y, z) = G(x - y, x + y, z)$ . Then, by Hölder's inequality and Sobolev embedding, we have the estimate

$$\begin{aligned} |(7.2a)| &\leq \int dz d(x-y) d(x+y) \frac{v_N(x-y)}{\sqrt{N}} |\bar{\phi}(y) \bar{u}(x, z) G(x-y, x+y, z)| \\ &\lesssim \int d(x-y) \frac{v_N(x-y)}{\sqrt{N}} \|\phi\|_{L^6(dx')} \|u\|_{L^3(dx')L^2(dz)} \|G(x-y, \cdot, \cdot)\|_{L^2(d(x+y)dz)} \\ &\lesssim \frac{1}{\sqrt{N}} \|v_N\|_{L^{\frac{6}{5}}} \|\nabla^{\frac{1}{2}} \phi\|_{L^3(dx)} \|u\|_{L^3(dx)L^2(dy)} \|F\|_{L^6(d(x-y))L^2(d(x+y)dz)}. \end{aligned}$$

Hence, by Corollary 5.4, we have the estimate

$$\begin{aligned} &\|(7.2a)\|_{L^1([0,T])L^2(dx_1 \dots dx_n)} \\ &\lesssim \frac{1}{\sqrt{N}} \|v_N\|_{L^{\frac{6}{5}}} \|\nabla^{\frac{1}{2}} \phi\|_{L^4([0,T])L^3(dx)} \|u\|_{L^4([0,T])L^3(dx)L^2(dy)} \|X\|_S \\ &\lesssim N^{\frac{\beta-1}{2}} \exp(\kappa T) \|X\|_S. \end{aligned}$$

For (7.2b), we see that

$$\begin{aligned} |(7.2b)| &\leq \int dx dy dz dx' \frac{v_N(x-y)}{\sqrt{N}} |\bar{\phi}(y) \bar{u}(x, z) \bar{p}(x, x') F(x', y, z)| \\ &\lesssim \int d(x-y) d(x+y) \frac{v_N(x-y)}{\sqrt{N}} |\phi(y)| \\ &\quad \times \|u(x, \cdot) p(x, \cdot)\|_{L^2(dz dx')} \|F(y, \cdot, \cdot)\|_{L^2(dz dx')} \\ &\lesssim N^{-\frac{1}{2}} \|\phi\|_{L^2(dx)} \|u\|_{L^\infty(dx)L^2(dy)} \|p\|_{L^\infty(dy)L^2(dz)} \|F\|_{L^2(dx dy dz)} \end{aligned}$$

which then, by Corollary 5.3, it follows

$$\begin{aligned} & \| (7.2b) \|_{L^1([0,T])L^2(dx_1 \dots dx_n)} \\ & \lesssim N^{-\frac{1}{2}} \| |\nabla|^{\frac{1}{2}+\varepsilon} \phi \|_{L^\infty(dt)L^2(dx)} \| u \|_{L^2([0,T])L^6(dx)L^2(dy)} \| p \|_{L^2([0,T])L^6(dx)L^2(dy)} \| X \|_S \\ & \lesssim N^{-\frac{1}{2}} \exp(\kappa T) \| X \|_S. \end{aligned}$$

Lastly, we see that

$$|(7.2c)| \lesssim N^{-\frac{1}{2}} \| \phi \|_{L^\infty(dx)} \| u \|_{L^\infty(dx)L^2(dy)} \| p \|_{L^\infty(dx)L^2(dy)}^2 \| F \|_{L^2(dxdydz)}$$

which again means

$$\| (7.2c) \|_{L^1([0,T])L^2(dx_1 \dots dx_n)} \lesssim N^{-\frac{1}{2}} \exp(\kappa T) \| X \|_S.$$

Thus, in general, the  $p$  term in  $\text{ch}(k)$  are lower order in  $N$  but they contribute to the growth in time in our estimates.

For the Fock space to Fock space estimate (7.1b) of (6.14a), let us first estimate (7.2a). By Cauchy-Schwarz inequality, it suffices to estimate

$$\begin{aligned} & \frac{1}{\sqrt{N}} \| v_N(x-y) \bar{\phi}(y) \bar{u}(x,z) \|_{L^2(dxdydz)} \\ & \lesssim \frac{1}{\sqrt{N}} \| v_N \|_{L^2} \| \phi \|_{L^\infty(dx)} \| u \|_{L^2(dxdy)} \lesssim N \exp(\kappa T) \end{aligned}$$

which follows from Theorem 5.2. The other two terms are handled in the same way with the only difference being the additional multiplication by  $\| p \|_{L^2(dxdy)}$ .

The estimates for the adjoint of (6.14a), contained in  $\mathcal{C}_1^*$ , is obtained by the dual estimate of (6.14a).

Next, let (6.14b) acts on  $X = (0, \dots, 0, F(x_1, \dots, x_{n+1}), 0, \dots)$  yields

$$\int dy \frac{v_N(x_1-y)}{\sqrt{N}} \bar{\phi}(y) F(y, x_1, \dots, x_n) \quad (7.3a)$$

$$+ \text{similar or lower order terms} \quad (7.3b)$$

on the  $n$  sector and zeroes elsewhere. Again, since  $x_2, \dots, x_n$  are passive variable, let us write  $F(y, x_1) = G(x_1 - y, x_1 + y)$ . Then it follows

$$\begin{aligned} \| (7.3a) \|_{L^2(dx_1)} & \lesssim \int dy \frac{v_N(y)}{\sqrt{N}} \| \bar{\phi}(x_1 - y) G(y, 2x_1 - y) \|_{L^2(dx_1)} \\ & \lesssim \int dy \frac{v_N(y)}{\sqrt{N}} \| \phi \|_{L^\infty(dx)} \| G(y, \cdot) \|_{L^2(dx)} \\ & \lesssim N^{\frac{\beta-1}{2}} \| \phi \|_{L^\infty(dx)} \| F \|_{L^6(d(x-y))L^2(d(x+y))} \end{aligned}$$

which means

$$\begin{aligned} & \| (7.3a) \|_{S'} \lesssim \| (7.3a) \|_{L^1(dt)L^2(dx_1 \dots dx_n)} \\ & \lesssim N^{\frac{\beta-1}{2}} \| |\nabla|^{\frac{1}{2}+\varepsilon} \phi \|_{L^2(dt)L^6(dx)} \| F \|_{L^2(dt)L^6(d(x_1-y))L^2(d(x_1+y)dx_2 \dots dx_n)} \\ & \lesssim N^{\frac{\beta-1}{2}} T \| X \|_S. \end{aligned}$$



The estimate for (7.3b) is obtained in a similar manner as above and the Fock space to Fock space estimate is clear. Moreover, the adjoint of (6.14b) is again obtained from the dual estimate.

The estimates for (6.14c)-(6.14h) are obtain in a similar manner.  $\square$

Next, we estimate the quartic terms.

**Proposition 7.2.** *Fix  $k \in \mathbb{N}$  and let  $0 < \beta < 1$ . If  $X$  is a Fock space vector that has nonzero entries only in the first  $k$  sectors, then we have the estimates*

$$\| \mathcal{D}X \|_{S'} \lesssim_k N^{\beta-1} \exp(\kappa T) \| X \|_S \quad (7.4a)$$

$$\| \mathcal{D}X \|_{\mathcal{F}} \lesssim_k \sqrt{N} \exp(\kappa T) \| X \|_{\mathcal{F}}. \quad (7.4b)$$

*Proof.* Letting (6.15a) act on  $F_{n-4}(x_1, \dots, x_{n-4})$  yields

$$(6.15a)F_{n-4} \sim \frac{1}{2N} v_N(x_1 - x_2) u(x_1, x_3) u(x_2, x_4) F_{n-5}(x_5, \dots, x_n) \quad (7.5a)$$

$$\frac{1}{2N} \int dy u(x_1, x_3) v_N(x_1 - y) p(y, x_2) u(y, x_4) F_{n-5}(x_5, \dots, x_n) \quad (7.5b)$$

$$\frac{1}{2N} \int dx dy u(x, x_3) p(x, x_1) v_N(x - y) p(y, x_2) u(y, x_4) F_{n-5}(x_5, \dots, x_n) \quad (7.5c)$$

$$+ \text{similar or lower order terms.} \quad (7.5d)$$

Since we have the estimate

$$\begin{aligned} & \frac{1}{2N} \| v_N(x_1 - x_2) u(x_1, x_3) u(x_2, x_4) \|_{L^{\frac{3}{2}}(d(x_1-x_2)d(x_1+x_2))L^2(dx_3dx_4)} \\ & \leq \frac{1}{2N} \| v_N \|_{L^{\frac{3}{2}}} \| u \|_{L^3(dx)}^2 L^2(dy) \end{aligned}$$

then it follows

$$\| (7.5a) \|_{S'} \lesssim N^{\beta-1} \| u \|_{L^4([0,T])L^3(dx)L^2(dy)}^2 \| X \|_S \lesssim N^{\beta-1} \exp(\kappa T) \| X \|_S.$$

To estimate (7.5b), observe we have the estimate

$$\begin{aligned} & \frac{1}{2N} \left\| \int dy v_N(x_1 - y) u(x_1, x_3) p(y, x_2) u(y, x_4) \right\|_{L^2(dx_1dx_2dx_3dx_4)} \\ & \lesssim \frac{1}{2N} \| v_N \|_{L^1} \| u \|_{L^2(dxdy)} \| u \|_{L^\infty(dx)L^2(dy)} \| p \|_{L^\infty(dx)L^2(dy)} \end{aligned}$$

which means

$$\| (7.5b) \|_{L^1(dt)L^2(dx_1 \dots dx_n)} \lesssim N^{-1} \exp(\kappa T) \| X \|_S.$$

Similar, we see that

$$\begin{aligned} & \| (7.5c) \|_{L^1(dt)L^2(dx_1 \dots dx_n)} \\ & \lesssim N^{-1} T \| u \|_{L^\infty([0,T])L^2(dx)L^2(dy)}^2 \| p \|_{L^\infty([0,T])L^\infty(dx)L^2(dy)} \| X \|_S \\ & \lesssim N^{-1} \exp(\kappa T) \| X \|_S. \end{aligned}$$

To obtain the Fock space to Fock space estimate, we see that

$$\begin{aligned} & \frac{1}{2N} \|v_N(x_1 - x_2)u(x_1, x_3)u(x_2, x_4)\|_{L^2(dx_1 dx_2 dx_3 dx_4)} \\ & \lesssim N^{-1} \|v_N\|_{L^2} \|u\|_{L^\infty(dx)L^2(dy)} \|u\|_{L^2(dxdy)} \end{aligned}$$

then it follows

$$\|(7.5a)\|_{L^2(dx_1 \dots dx_n)} \lesssim N^{\frac{1}{2}} \exp(\kappa T) \|X\|_{\mathcal{F}}.$$

The Fock space to Fock space estimate for (7.5b) and (7.5c) follow a similar calculation.

Next, let us consider the action of (6.15b) and (6.15c) on  $F_{n-2}(x_1, \dots, x_{n-2})$ . Observe we have

$$(6.15b)F_{n-2} \sim \frac{1}{2N} v_N(x_1 - x_2)u(x_1, x_3)F_{n-2}(x_2, x_4, \dots, x_n) \quad (7.6a)$$

$$+ \text{similar or lower order terms} \quad (7.6b)$$

It is similar for (6.15c). Since  $x_5, \dots, x_n$  are passive variables, let us suppress the notation. Then we see that

$$\begin{aligned} & \frac{1}{2N} \|v_N(x_1 - x_2)u(x_1, x_3)F(x_2, x_4)\|_{L^{\frac{3}{2}}(d(x_1-x_2)d(x_1+x_2))L^2(dx_3 dx_4)} \\ & \lesssim \frac{1}{2N} \|v_N\|_{L^{\frac{3}{2}}} \|u\|_{L^6(dx)L^2(dy)} \|F\|_{L^2(dxdy)} \end{aligned}$$

which leads to the desired estimate. The Fock space to Fock space estimate is similar to the above estimate for (7.5a).

The estimates for (6.15d), (6.15e), and (6.15h) follow from the same arguments as (7.5b) and (7.5c).

To estimate (6.15f), observe we have

$$(6.15f)F_n \sim \frac{1}{2N} \int dydz v_N(x_1 - y)u(x_1, x_2)\bar{u}(y, z)F_n(y, z, x_3, \dots, x_n) \quad (7.7a)$$

$$+ \text{similar or lower order terms.} \quad (7.7b)$$

Then it follows

$$\begin{aligned} & \|(7.7a)\|_{L^2(dx_1 dx_2)} \\ & \lesssim \frac{1}{2N} \int d\eta v_N(\eta) \left\| \int dz u(x_1, x_2)\bar{u}(x_1 - \eta, z)F_n(x_1 - \eta, z) \right\|_{L^2(dx_1 dx_2)} \\ & \lesssim N^{-1} \|u\|_{L^\infty(dx)L^2(dy)}^2 \|F\|_{L^2(dx_1 dx_2)} \end{aligned}$$

which yields the desired estimate. Likewise, we can estimate (6.15g).

Lastly, when we remove (6.15j) from (6.15i), the remainder parts can be handle in the same way as in the cases of (6.15b) and (6.15c).  $\square$

Let us now consider the estimates for the quadratic terms.

**Proposition 7.3.** *Fix  $k \in \mathbb{N}$ . Then  $\mathcal{Q}$ , defined by (6.11a), is bounded uniformly in  $N$  from the first  $k$  sectors of Fock space to Fock space. More precisely, let  $X$  be a Fock vector that has nonzero entries only in the first  $k$  sectors, then we have the following uniform in  $N$  estimate*

$$\| \mathcal{Q}X \|_{\mathcal{F}} \lesssim_{\varepsilon, k} \exp(\kappa T) \| X \|_{\mathcal{F}} \quad (7.8)$$

for some  $\kappa > 0$ .

*Proof.* Letting (6.12a) acts on the Fock vector  $X = (0, \dots, 0, F_n(x_1, \dots, x_n), 0, \dots, 0)$ , where  $n \leq k$ , yields

$$\begin{aligned} ((6.12a)X)_n &= \sum_{j=1}^n (v_N * \text{diag } \Gamma)(t, x_j) F_n(x_1, \dots, x_n) \\ &\quad \sum_{j=1}^n \int dy (v_N \Gamma)(t, x_j, y) F_n(y, x_1, \dots, \hat{x}_j, \dots, x_n) \end{aligned}$$

and zero elsewhere. It suffices to consider the first summand of the two summations. Also, since  $x_3, \dots, x_n$  are passive variables in the first summand, we will suppress them without further comment.

Observe, by Hölder, Sobolev, and Proposition 4.12, we have the estimate

$$\begin{aligned} &\| (v_N * \text{diag } \Gamma)(t, x_1) F(x_1, x_2) \|_{L^2(dx_1 dx_2)} \\ &\lesssim \dot{\mathbf{N}}_T(\nabla_{x+y}^2 \Gamma) \| F \|_{L^2(dx_1 dx_2)} \lesssim \exp(\kappa T) \| F \|_{L^2(dx_1 dx_2)} \end{aligned}$$

and likewise we have

$$\| \int dy v_N(x_1 - y) \Gamma(x_1, y) F(y, x_2) \|_{L^2(dx_1 dx_2)} \lesssim \exp(\kappa T) \| F \|_{L^2(dx_1 dx_2)}.$$

Next, letting (6.12b) acts on  $X$  yields

$$((6.12b)X)_n = \int dy w(x_1, y) F(y, x_2, \dots, x_n) + \text{similar terms}$$

and zero elsewhere. Then, by Cauchy-Schwarz inequality, we see that

$$\| \int dy w(x_1, y) F(y, x_2) \|_{L^2(dx_1 dx_2)} \lesssim \| w \|_{L^2(dx dy)} \| F \|_{L^2(dx_1 dx_2)}.$$

To estimate  $\| w \|_{L^2(dx dy)}$  uniformly in  $N$ , we divide the estimate into two parts. For the first part, we consider

$$\| \text{sh}(k) \circ \overline{(v_N \Lambda)} \circ \overline{\text{ch}(k)}^{-1} \|_{L^2(dx dy)} \quad (7.9a)$$

which can be estimated as follows

$$(7.9a) \lesssim \mathbf{N}_T(\nabla_{x+y}^2 \Lambda) \| \text{sh}(k) \|_{L^2(dx dy)} \lesssim \exp(\kappa T).$$

The estimate for the second part

$$\| \tilde{\mathbf{W}}(\overline{\text{ch}(k)}) \circ \overline{\text{ch}(k)}^{-1} \|_{L^2(dx dy)} \quad (7.9b)$$

follows from Lemma 5.15.  $\square$

## REFERENCES

- [BBCFS16] V. Bach, S. Breteaux, T. Chen, J. Fröhlich, and I. M. Sigal, *The time-dependent hartree-fock-bogoliubov equations for bosons*, arXiv preprint arXiv:1602.05171 (2016), 1–36.
- [BCS17] C. Boccato, S. Cenatiempo, and B. Schlein, *Quantum many-body fluctuations around nonlinear Schrödinger dynamics*, Annales Henri Poincaré **18** (2017), no. 1, 113–191.
- [BdOS15] N. Benedikter, G. de Oliveira, and B. Schlein, *Quantitative derivation of the Gross-Pitaevskii equation*, Communications on Pure and Applied Mathematics **68** (2015), no. 8, 1399–1482.
- [Ber12] F. A. Berazin, *The method of second quantization*, Pure and Applied Physics, vol. 24, Academic Press, 2012.
- [CGMZ20] J. Chong, M. Grillakis, M. Machedon, and Z. Zhao, *Global estimates for the hartree-fock-bogoliubov equations*, arXiv preprint, arXiv: 2008.01753 (2020), 1–47.
- [Che12] X. Chen, *Second order corrections to mean field evolution for weakly interacting bosons in the case of three-body interactions*, Archive for Rational Mechanics and Analysis **203** (2012), no. 2, 455–497.
- [Cho20] J. Chong, *Dynamics of large boson systems with attractive interaction and a derivation of the cubic focusing NLS in  $\mathbb{R}^3$* , arXiv preprint arXiv:1608.01615 (2020), 1–52.
- [CLS11] L. Chen, J.O. Lee, and B. Schlein, *Rate of convergence towards hartree dynamics*, Journal of Statistical Physics **144** (2011), no. 4, 872–903.
- [DG13] J. Dereziński and C. Gérard, *Mathematics of quantization and quantum fields*, Cambridge University Press, 2013.
- [ES09] L. Erdős and B. Schlein, *Quantum dynamics with mean field interactions: a new approach*, Journal of Statistical Physics **134** (2009), no. 5, 859–870.
- [ESY06] L. Erdős, B. Schlein, and H.-T. Yau, *Derivation of the Gross-Pitaevskii hierarchy for the dynamics of Bose-Einstein condensate*, Communications on Pure and Applied Mathematics **59** (2006), no. 12, 1659–1741.
- [ESY07] ———, *Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems*, Inventiones Mathematicae **167** (2007), no. 3, 515–614.
- [ESY09] ———, *Rigorous derivation of the gross-pitaevskii equation with a large interaction potential*, Journal of the American Mathematical Society **22** (2009), no. 4, 1099–1156.
- [ESY10] ———, *Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensate*, Annals of Mathematics **172** (2010), no. 1, 291–370.
- [EY01] L. Erdős and H.-T. Yau, *Derivation of the nonlinear Schrödinger equation from a many-body Coulomb system*, Advances in Theoretical and Mathematical Physics **5** (2001), no. 6, 1169–1205.
- [FKS09] J. Fröhlich, A. Knowles, and S. Schwarz, *On the mean-field limit of bosons with coulomb two-body interaction*, Communications in Mathematical Physics **288** (2009), no. 3, 1023–1059.
- [Fol89] G. B. Folland, *Harmonic analysis in phase space*, Annals of Mathematics Studies, no. 122, Princeton University Press, 1989.
- [GM13a] M. Grillakis and M. Machedon, *Beyond mean field: On the role of pair excitations in the evolution of condensates*, Journal of Fixed Point Theory and Applications **14** (2013), no. 1, 91–111.
- [GM13b] ———, *Pair excitations and the mean field approximation of interacting Bosons, I*, Communications in Mathematical Physics **324** (2013), no. 2, 601–636.

- [GM17] ———, *Pair excitations and the mean field approximation of interacting Bosons, II*, Communications in Partial Differential Equations **42** (2017), no. 1, 24–67.
- [GM19] ———, *Uniform in  $n$  estimates for a Bosonic system of Hartree–Fock–Bogoliubov type*, Communications in Partial Differential Equations **44** (2019), no. 12, 1431–1465.
- [GMM10] M. Grillakis, M. Machedon, and D. Margetis, *Second-order corrections to mean field evolution of weakly interacting Bosons. I.*, Communications in Mathematical Physics **294** (2010), no. 1, 273–301.
- [GMM11] ———, *Second-order corrections to mean field evolution of weakly interacting Bosons. II*, Advances in Mathematics **228** (2011), no. 3, 1788–1815.
- [GMM17] ———, *Evolution of the boson gas at zero temperature: Mean-field limit and second-order correction*, Quarterly of Applied Mathematics **75** (2017), no. 1, 69–104.
- [Gol16] François Golse, *On the dynamics of large particle systems in the mean field limit*, Macroscopic and Large Scale Phenomena: Coarse Graining, Mean Field Limits and Ergodicity, Springer, 2016, pp. 1–144.
- [GV79a] J. Ginibre and G. Velo, *The classical field limit of scattering theory for non-relativistic many-boson systems. I*, Communications in Mathematical Physics **66** (1979), no. 1, 37–76.
- [GV79b] ———, *The classical field limit of scattering theory for non-relativistic many-boson systems. II*, Communications in Mathematical Physics **68** (1979), no. 1, 45–68.
- [Hep74] K. Hepp, *The classical limit for quantum mechanical correlation functions*, Communications in Mathematical Physics **35** (1974), no. 4, 265–277.
- [KP10] Antti Knowles and Peter Pickl, *Mean-field dynamics: singular potentials and rate of convergence*, Communications in Mathematical Physics **298** (2010), no. 1, 101–138.
- [Kuz15] E. Kuz, *Rate of convergence to mean field for interacting Bosons*, Communications in Partial Differential Equations **40** (2015), no. 10, 1831–1854.
- [Kuz17] ———, *Exact evolution versus mean field with second-order correction for Bosons interacting via short-range two-body potential*, Differential and Integral Equations **30** (2017), no. 7/8, 587–630.
- [LGS15] T. Langen, R. Geiger, and J. Schmiedmayer, *Ultracold atoms out of equilibrium*, Annu. Rev. Condens. Matter Phys. **6** (2015), no. 1, 201–217.
- [LSSY05] E. H. Lieb, R. Seiringer, J. P. Philip Solovej, and J. Yngvason, *The mathematics of the bose gas and its condensation*, vol. 34, Springer Science & Business Media, 2005.
- [NN17] P. T. Nam and M. Napiórkowski, *Bogoliubov correction to the mean-field dynamics of interacting bosons*, Advances in Theoretical and Mathematical Physics **21** (2017), no. 3, 683–738.
- [RS09] I. Rodnianski and B. Schlein, *Quantum fluctuations and rate of convergence towards mean field dynamics*, Communications in Mathematical Physics **291** (2009), no. 1, 31–61.
- [Sol14] J. P. Solovej, *Many body quantum mechanics*, Lecture Notes. Summer (2014), 1–102.
- [Tao06] T. Tao, *Nonlinear dispersive equations: local and global analysis*, CBMS Regional Conference Series in Mathematics, no. 106, American Mathematical Society, 2006.

JACKY J. CHONG

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS, AUSTIN, U.S.A.

Email address: [jwchong@math.utexas.edu](mailto:jwchong@math.utexas.edu)

ZEHUA ZHAO

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MARYLAND, COLLEGE PARK, U.S.A.

*Email address:* `zzh@umd.edu`