

Regularization and Hierarchical Prior Distributions for Adjustment with Health Care Claims Data: Rethinking Comorbidity Scores

Jacob V Spertus¹, Samrachana Adhikari¹, and Sharon-Lise T Normand^{1,2}

1: Department of Health Care Policy, Harvard Medical School

2: Department of Biostatistics, Harvard TH Chan School of Public Health

Abstract

Health care claims data refer to information generated from interactions within health systems. They have been used in health services research for decades to assess effectiveness of interventions, determine the quality of medical care, predict disease prognosis, and monitor population health. While claims data are relatively cheap and ubiquitous, they are high-dimensional, sparse, and noisy, typically requiring dimension reduction. In health services research, the most common data reduction strategy involves use of a comorbidity index – a single number summary reflecting overall patient health. We discuss Bayesian regularization strategies and a novel hierarchical prior distribution as better options for dimension reduction in claims data. The specifications are designed to work with a large number of codes while controlling variance by shrinking coefficients towards zero or towards a group-level mean. A comparison of drug-eluting to bare-metal coronary stents illustrates approaches. In our application, regularization and a hierarchical prior improved over comorbidity scores in terms of prediction and causal inference, as evidenced by out-of-sample fit and the ability to meet falsifiability endpoints.

Keywords: Bayesian methods; comorbidity scores; propensity score weighting; regularization; coronary stent.

1 Introduction

Health services researchers focus on assessing the effectiveness of treatments and policy interventions, characterizing the quality of medical care delivered to specific populations, and developing payment systems, with the goal of improving health outcomes. A fundamental data source used in health services research is administrative billing claims data. Claims data refer to a wide range of administrative databases capturing health care utilization information for reimbursement purposes and are generated by routine interactions between patients, the health system, and insurance providers [12]. This type of data stands in contrast to other formats, like clinical registry data, which are prospectively collected for the explicit purpose of later analysis and are of considerably higher quality [6][27][12]. However claims data have many desirable features: they are generated during the normal work flow and thus avoid additional effort and expense; they provide an accounting of denominator populations to ensure rates are well-defined; and they have standard definitions

to help with consistency of the information collected across the nation and around the world.

In this article, we focus on claims data comprised of alpha-numeric codes for conditions from the International Classification of Diseases (9th Clinical Modification, [ICD-9]) system [9]. The structure of ICD-9 codes is a tree-like hierarchy with each code constructed as a binary variable. The codes contain 3-5 alpha-numeric strings where the first 3 characters define a group of diagnoses (or procedures) and the additional two characters provide further specificity. The ICD-9 codes include approximately 17,000 distinct codes at level 5, many of which represent rare conditions that are unlikely to be observed in any finite population. Coding practices can also vary across health delivery settings such that codes are noisy representations of true conditions. A revision of ICD-9 codes (ICD-10) took effect in October 2015 in the United States, and contains approximately 68,000 codes having a similar hierarchical structure to the ICD-9 system. Thus, while claims data are relatively cheap and ubiquitous, they are high-dimensional, sparse, and noisy. Consequently, some form of dimension reduction for inference is required.

Researchers have proposed a variety of dimension reduction approaches when using claims data. The most common strategies either use clinical expertise to select a small set of codes and include separate binary indicators for each discrete diagnosis or procedure, or implement standard variable selection approaches such as step-wise regression. Because these approaches may omit important variables, inclusion of a summary measure, called a comorbidity index or score, helps with adjusting for other health features [29]. Two of the most frequently used comorbidity indices include the Charlson and the Elixhauser [7][8]. These indices were constructed by selecting and validating ICD-9 codes based on a combination of clinical expertise and statistical analysis. The Elixhauser index has been shown to predict long-term clinical outcomes better than the Charlson index [29]. Both indices may be implemented as a set of dummy variables indicating the presence of groups of conditions (e.g. hypertension, diabetes, etc) if feasible or as a single continuous variable constructed from a weighted sum of the individual condition groups [7][39]. A recent paper by Gilbert et al argued that using a comorbidity index with fixed historical weights is as good as including individual claims coefficients in causal survival analysis [11]. However the assumptions associated with the strategy are rather strong, inferences may be compromised by ignoring uncertainty from the estimation of the weights, and the findings do not translate to other approaches like propensity score analyses or logistic regression for mortality prediction.

Our point of departure from typical approaches is to consider the dimension reduction task as one of regularization. Fundamentally some outside information, typically an assumption that some of the conditions have little or no effect on the outcome of interest, must be incorporated when the number of variables available exceeds the capacity of the data for appropriately precise estimation.

We note that other sophisticated methods have been developed to use high-dimensional claims data for various analysis goals. Syed et al proposed using a support vector machine to enhance risk prediction using current procedural terminology codes [36]. Perotte et al proposed exploiting the hierarchical nature of ICD-9 codes using a support vector machine for diagnosis code assignment [20]. Singh used the group lasso to improve outcome prediction by accounting for the ICD-9 hierarchy in a frequentist setting [30]. In the causal inference domain, Schneeweiss et al developed a frequentist algorithm to select claims confounders based on their prevalence and univariate associations with both treatment and

outcome [28].

Our work is motivated by determining the comparative effectiveness of coronary stents once released into the U.S. market. Stents are small scaffolds placed into the heart via a percutaneous coronary intervention (PCI) in order to treat arteries blocked by plaque and keep them clear and open. Drug-eluting stents (DESs) are coated in a drug designed to keep plaque from re-forming, while bare-metal stents (BMSs) have no such coating. Clinical trials have consistently shown that DESs are efficacious as they reduce the need for target vessel revascularization (TVR), i.e. follow-up procedures in the same artery, but have no benefit on survival [2][34]. However, DESs necessitate adherence to prolonged dual-antiplatelet therapies to prevent rare but fatal stent thrombosis. Several patient characteristics such as age, comorbidity, and socio-economic factors may impact adherence to dual-antiplatelet therapy, and thus such patients are likely to be implanted with BMSs [33]. Because these characteristics also tend to increase the risk of adverse outcomes, observed differences in average outcomes between BMS and DES treated patient groups are heavily confounded. Even after statistical adjustments various observational analyses have generally found that DES provides a survival benefit in usual care populations [14][37]. Although there has historically been speculation that this benefit was real but not observed in randomized trial populations, a recent quasi-randomization analysis and the results of a large pragmatic clinical trial make a strong case that any observed survival benefit is due to residual confounding [42][2]. As a result, it remains an open challenge for observational causal inference to accurately estimate the effects of DES versus BMS using observational data. It may be possible to achieve better results by including claims data in some form.

While we utilize a Bayesian propensity score framework, the ideas apply to other techniques such as frequentist propensity scores or Bayesian outcome modeling. A Bayesian approach is particularly well aligned with high-dimensional inference with claims data, as it provides a number of intuitive tools for model building and incorporates all major sources of uncertainty, which may ultimately lead to better inferential performance [31]. Our focus is on improving over a typical analysis using comorbidity indices and to this end we make three interdependent arguments: (1) administrative claims data can help analysts meet the ignorability assumption in observational studies, but standard comorbidity indices may be insufficient; (2) using a considerably larger and more granular set of claims data may lead to better results compared with a comorbidity index; and (3) regularization techniques are very useful for controlling variance when using a high-dimensional set of claims data. Combining these principles, we find that employing high-dimensional claims data with Bayesian regularization techniques leads to significant improvement over a comorbidity index in terms of treatment model fit and causal inference.

In section 2 we briefly remind the reader the necessary assumptions for causal inference with observational data and review propensity score approaches. Our proposed methods are presented in detail in section 3. Section 4 illustrates various strategies for dimension reduction by revisiting the DES vs BMS problem. We conclude with a discussion of the advantages of regularization over comorbidity indices for causal inference, as well as potential improvements and modifications for future research.

2 Causal Inference and Propensity Scores

2.1 Causal Inference

Causal inference involves various assumptions, three of which include stable unit treatment value assumption (SUTVA), positivity, and treatment assignment ignorability. We focus on binary treatments where a new treatment is compared to a standard treatment. In this setting, SUTVA implies that treatment does not vary within treatment groups and that the treatment received by one subject does not affect the outcomes of another. Positivity requires that every subject has a chance of receiving the new treatment and we assume that this probability is bounded away from 0 and 1. Positivity can be empirically tested in that blatant violations are apparent by examination or modeling of the data. Propensity scores help facilitate assessing the positivity assumption [21]. Finally, ignorability assumes that potential outcomes are independent of treatment assignment [24]. In observational designs, investigators must adjust for confounders, variables that affect both treatment assignment and outcome, to uncover the causal effect of a particular treatment. This task is among the most challenging in observational causal inference and has inspired many attempts to define methods, either statistical or expertise driven, to select confounders for adjustment [3][28][31]. High-dimensional datasets may provide important variables that help investigators meet the ignorability assumption. In particular, health care claims data provide an exciting source of potential confounders, presenting a host of opportunities as well as challenges for researchers.

2.2 Propensity Scores

Propensity scores are the probabilities of receiving the new treatment relative to the standard treatment. They are a commonly used tool for observational causal inference. One particularly important feature involves facilitating an *outcome free analysis* in that they separate outcome modeling from treatment modeling which can help to maintain objectivity [25]. This feature is often a requirement by regulators, such as the U.S. Food and Drug Administration, who place a premium on objective findings [13]. Propensity scores can be particularly useful in high-dimensional settings when event rates are low or when the number of confounders is large relative to number of events. As a result, many investigators interested in observational causal inference with high-dimensional data have focused on how to build propensity score models. Variable selection is the standard response to dealing with high-dimensional data, and researchers have historically focused on selecting important confounders based on subject matter expertise and rules of thumb [3][35]. More algorithmic variable selection techniques have also been explored, including the high-dimensional propensity score algorithm of Schneeweiss et al, but this uses outcome information to select variables [28]. Other approaches have proposed more technical variance reduction strategies through machine learning methods, such as elastic net regularization, tree-based algorithms, and learning ensembles [32][15][38]. Each of these methods has shown promise in estimating causal effects from large databases. We use a Bayesian framework because it propagates uncertainty from the propensity score model and provides intuitive and sophisticated variance reduction strategies via prior distributions, an advantage that we leverage through a hierarchical prior specification.

2.2.1 Bayesian Propensity Scores

We adopt a Bayesian treatment model and maintain an outcome free analysis, although jointly modeling the propensity score and outcomes has been proposed [16][26]. The particular algorithm we use resembles frequentist propensity score weighting [31]. The algorithm uses a conjugate prior-likelihood pair in the form of a beta-binomial: if Y represents the total number of subjects having an event out of n subjects, e.g., $Y \stackrel{\text{iid}}{\sim} \text{Bin}(n, p)$, then assuming p arises from $\text{Beta}(a, b)$ *a-priori*, yields *a-posteriori* $p \mid Y \sim \text{Beta}(a + Y, b + n - Y)$. We use this fact to integrate over draws from the propensity score model while conditionally drawing from the outcome model, so that uncertainty is incorporated from both the propensity score model and the outcome model.

Briefly, let Y_i denote binary outcome for subject i , X_i binary treatment, and π_i the propensity score, $\mathbb{P}(X_i = 1)$. The parameter $\boldsymbol{\pi}$ is the vector of propensity scores for all subjects, $\mathbf{X}$ is the vector of treatment indicators, $\mathbf{Y}$ the vector of outcomes, and $\mathbf{D} = \{\mathbf{X}, \mathbf{Y}\}$ their concatenation into an $n \times 2$ matrix. Together, $\boldsymbol{\pi}$ and $\mathbf{D}$ are sufficient to estimate an unbiased treatment effect under an ignorable treatment assignment mechanism. Let p_1 and p_0 be the marginal probability of the event in treatment group 1 and 0 respectively. These can be estimated in an unbiased fashion by weighting the original populations according to the inverse of their propensity scores.

We let $\tilde{Y}(1)$ denote the total number of potential events in treatment group 1 (all subjects for which $X_i = 1$), and $\tilde{Y}(0)$ the number of events in treatment group 0. By assuming a binomial likelihood for $\tilde{Y}(1)$ and $\tilde{Y}(0)$, and selecting flat priors p_0 and $p_1 \sim \text{Beta}(1, 1)$, we can take advantage of conjugacy to get closed form solutions for the posterior distributions for p_0 and p_1 . The priors are augmented by weighted counts of successes and failures in each treatment group to generate beta posteriors, $p_g \mid \mathbf{D}, \boldsymbol{\pi} \sim \text{Beta}(a_g^*, b_g^*)$ for $g = 0, 1$ where

$$\begin{aligned} a_g^* &= 1 + \gamma_g \left(g \sum_{i=1}^n \frac{X_i Y_i}{\pi_i} + (1 - g) \sum_{i=1}^n \frac{(1 - X_i) Y_i}{1 - \pi_i} \right), \\ b_g^* &= 1 + \gamma_g \left(g \sum_{i=1}^n \frac{X_i (1 - Y_i)}{\pi_i} + (1 - g) \sum_{i=1}^n \frac{(1 - X_i) (1 - Y_i)}{1 - \pi_i} \right) \end{aligned}$$

and $a_g^* (b_g^*)$ describes the weighted number of events (non-events) in treatment group g . The quantity γ_g is used to normalize the weighted population to its original size and is defined as $\gamma_g = g \sum_{i=1}^n \frac{(1 - X_i)(1 - Y_i)}{1 - \pi_i} + (1 - g) \sum_{i=1}^n \frac{(1 - X_i)(1 - Y_i)}{1 - \pi_i}$. The causal parameter, such as the average treatment effect, can be calculated as simple functions of draws from the respective posteriors of p_0 and p_1 - for example the average treatment effect defined by the posterior risk difference, $\Delta = p_1 - p_0$. Estimates can be obtained using Markov Chain Monte Carlo simulation and incorporate the uncertainty from both the propensity score and outcome models (see in Spertus et al [31]).

3 Proposed Risk Adjustment Approaches

We now consider estimating propensity scores $\pi_i = \mathbb{P}(X_i \mid \mathbf{B}_i, \mathbf{C}_i)$, where $\mathbf{B}_i$ is a vector of baseline confounders, for example demographic or clinical covariates from a clinical registry,

and $\mathbf{C}_i$ is a vector of 4-digit ICD-9 diagnosis codes. We use 4-digit rather than 5-digit codes for simplicity and without loss of generality. We define the propensity score model as

$$\pi_i = \text{logit}^{-1}\{\beta_0 + \beta_B \mathbf{B}_i + \beta_C \mathbf{C}_i\}. \quad (1)$$

The intercept β_0 and coefficients associated with the baseline confounders β_B are standard logistic regression parameters. We set a non-informative prior for the intercept as $\beta_0 \sim \mathcal{N}(0, 10)$. The 95% interval for β_0 is about $[-20, 20]$, which on the logit scale encompasses the vast majority of possible values for the marginal treatment rate. We assume that each component of β_B is distributed *a priori* as $t_5(0, 2.5)$ and components are mutually independent. This prior is referred to as “weakly informative” in the Bayesian data analysis literature [10]. It places the majority of prior mass within $[-5, 5]$ though heavy tails allow larger estimates if the data warrants. Weakly informative priors help produce a well-defined posterior but otherwise do little shrinkage compared to a maximum likelihood estimate. Having specified priors for the intercept and baseline variables, we turn to various approaches to handling the claims variables $\mathbf{C}_i$ using regularization.

3.1 Regularization via Prior Distributions

3.1.1 Weakly Informative Prior Distributions

A natural choice for the prior distribution of β_C is weakly informative such that $\beta_C \sim t_5(0, 2.5)$. These priors could be applied to groups of codes at the 3-digit level, or to more granular 4-digit codes. The latter is likely to have higher variance but less bias, and we explore both possibilities in our application. However, alternatives may be required because it is likely that the majority of the ICD-9 codes are rare and have no relationship with the treatment assignment, so they will introduce considerable variance.

3.1.2 Sparsity-Inducing Prior Distributions

An alternative *sparsity-inducing* prior is the horseshoe prior where for the j th component of β_C . We assume the associated coefficient is

$$\beta_{Cj} \sim \mathcal{N}(0, \lambda_j^2 \tau^2); \text{ where } \lambda_j \sim \text{Cauchy}^+(0, 1) \text{ and } \tau \sim \text{Cauchy}^+(0, 1). \quad (2)$$

The notation $\text{Cauchy}^+(0, 1)$ denotes a half-Cauchy distribution defined over the positive real line. The horseshoe performs global shrinkage through τ and local shrinkage through λ_j , allowing signals to remain near their maximum likelihood estimates, while shrinking noise variables aggressively towards zero. It has shown impressive predictive properties in simulations and applied problems [5][19]. For easier sampling, we replace the priors for λ_j with $t_3^+(0, 1)$ priors which users find have little effect on coefficient estimates in practice [22].

Under this specification, 50% of the prior mass on β_{Cj} is placed between -.4 and .4, which is quite close to zero. However the 95% prior interval is about $(-11, 11)$, reflecting the fact that the horseshoe places a lot of prior weight near zero but also has “fat tails” in order to avoid shrinking signals.

3.1.3 Hierarchical Prior Distributions

As an alternative to standard weakly informative or sparsity-inducing priors, we propose a hierarchical prior on β_C that takes advantage of the natural tree-like hierarchy in ICD-9 codes. Assuming p elements for C_i , each element falls into one of L 3-digit categories, where category l is associated with $p_l \geq 1$ codes. The codes in category l are denoted by C_{il} . The vector of coefficients for C_i has a special structure. It is partitioned into $\beta_C = \{\beta_{C_1}, \beta_{C_2}, \dots, \beta_{C_L}\}$ where β_{C_l} is the vector of coefficients for the grouped variables C_{il} . To emphasize the grouped structure in the regression, we rewrite equation 1 as:

$$\pi_i = \text{logit}^{-1}\{\beta_0 + \beta_B B_i + \sum_{l=1}^L \sum_{k=1}^{p_l} \beta_{C_{l_k}} C_{il_k}\}. \quad (3)$$

Capitalizing on the hierarchical structure, we define a hierarchical prior distribution as

$$\beta_{C_l} \sim \text{Laplace}(\mu_l, .5) \text{ where } \mu_l \sim t_5(0, 2.5). \quad (4)$$

The $t_5(0, 2.5)$ priors on μ_l are again weakly informative. The 4-digit ICD-9 code coefficients are shrunk to the 3-digit (group) mean μ_l according to a Laplace distribution with a relatively tight scale. For example, *a priori* each 4-digit code has a probability of 0.86 of being within 1 of the group mean coefficient μ_l . The choice of the Laplace distribution over the Normal distribution forces noisier 4-digit codes to be shrunk more aggressively to the 3-digit group mean, while fatter tails permits less shrinking of signals [10]. If there is only one 4-digit code in a 3-digit category, then we just assume $\beta_{C_l} = \mu_l$ and specify the weakly informative prior $\beta_{C_l} \sim t_5(0, 2.5)$.

To fix ideas, imagine the group of ICD-9 codes corresponds to the category "hypertensive heart disease" and that it has an overall effect of $\mu_l = 2$ (Figure 1). The 4-digit codes for this category include benign, unspecified, and malignant hypertensive heart disease. We might expect the benign condition to in fact have a small effect on stent selection such that $\beta_{C_{l_1}} = 0$; the malignant form of the condition may have a larger effect than the group mean, so $\beta_{C_{l_3}} = 3$; and the unspecified condition is more ambiguous and receives an estimate equal to the group mean, $\beta_{C_{l_2}} = 2$. The fundamental advantage is that some information is shared among similar conditions, while different forms and severities are able to receive larger or smaller effect estimates as the data warrant.

3.2 Regularization via Comorbidity Scoring

Comorbidity indices are the results of a dimension reduction strategy that have the related advantages of low variance and parsimonious interpretation [11]. In typical usage, comorbidity indices compress hundreds or thousands of binary-valued ICD-9 codes into a smaller set of indicator variables or into a scalar summary, either of which can then be used for prediction or risk adjustment. Either way, the resulting claims data set is much smaller, with many of the original diagnosis data discarded or aggregated. The Elixhauser comorbidity index, for example, reduces all ICD-9 codes into 30 indicators, which can then be combined further into a scalar score using fixed severity weights [8][39]. The severity weights are obtained from a regression model and are available in programs distributed by the developer of the index. In our setting of causal inference, the association of the




Figure 1: **Hierarchical prior on ICD-9 regression coefficients.** $t_5(0, 2.5)$ prior distribution for μ_l (on top) is weakly informative for the group-level mean. β_{C_l} , the coefficients for granular codes, are shrunk towards μ_l by a Laplace distribution.

Elixhauser comorbidities are nuisance parameters – their use is in meeting the ignorability of the treatment assignment assumption.

4 Application

We implement a variety of methods to model propensity scores for DES vs BMS using the Bayesian propensity score approach to compare inferences using a variety of regularization strategies. We utilize ICD-9 diagnosis codes that are present-on-admission for a cohort of adults undergoing coronary stenting in hospitals.

4.1 Claims Data and the Mass-DAC Registry

We used claims data from the Massachusetts Center for Health Information and Analysis (CHIA). The CHIA data included 15 free-response fields with Present On Admission (POA) ICD-9 codes. Because POA codes describe patients at hospital arrival, they are not impacted by treatment decisions. To set a lower bound on sparsity we considered all 4-digit POA ICD-9 for which 10 or more patients were coded as having the condition.

The claims data were supplemented with baseline demographic and clinical variables from the Mass-DAC registry, a state mandated database harvesting clinical information for all percutaneous coronary interventions (PCIs) performed in adults (age ≥ 18 years) in all

non-Federal Massachusetts’ hospitals annually. We identified 131 potential confounders in the Mass-DAC registry, including variables capturing demographic characteristics (e.g. age, race), pre-existing conditions (e.g. heart disease, diabetes), or procedure information (e.g. treated vessel, heart attack severity, hospital). The registry data are prospectively collected by hospital personnel who use the National Cardiovascular Data Registry instrument from the American College of Cardiology. A very small number of cells were missing data, and we imputed a single dataset for analysis.

We compared a number of outcomes between subjects implanted with a drug eluting stent (DES) and subjects implanted with bare metal stents (BMS) using Bayesian propensity score methods: 2-day mortality, 30-day mortality, 1-year mortality, and target vessel revascularization (TVR). Two-day and 30-day mortality are designed to serve as “falsifiability endpoints.” Because any difference in the stent performance would not manifest after such a short period, if we observe a difference on one of these outcomes it is almost certain it is due to residual confounding. We also generally expect no difference, or a very small difference, on 1-year mortality based on previous findings from RCTs and instrumental variable analyses [2][42]. We do expect a significant benefit to DES on TVR rates.

4.2 Applied Methods

In all methods, the Mass-DAC variables were included and their coefficients, β_B , were specified with weakly informative $t_5(0, 2.5)$ priors. Therefore only the modeling of claims data varied.

As baseline analyses, we fit two different Bayesian propensity score models to the log-odds of DES based on the Elixhauser index. We first grouped variables using the mapping of Quan et al from ICD-9 codes to the 30 original Elixhauser categories [8][23]. This mapping was accomplished in R using the package `medicalrisk` [17]. We then implemented a model with indicator variables for each category and used a weakly informative $t_5(0, 2.5)$ prior. We refer to this model as “Elixhauser indicator.” Second, we replaced the 30 binary indicators with a comorbidity score created by summing the product of each indicator with its fixed severity weight.[39] We refer to this model as the “Elixhauser continuous.”

We used the full claims set (defined as all 4-digit POA ICD-9 for which 10 or more patients were coded as having the condition) with weakly informative, sparsity-inducing, and hierarchical priors to estimate the propensity score. As standard approaches a logistic regression using 3-digit codes and $t_5(0, 2.5)$ priors and a logistic regression using 4-digit ICD-9 codes and $t_5(0, 2.5)$ priors were fit [18]. Using 3-digit rather than 4-digit codes is a typical dimension reduction technique. We have found $t_5(0, 2.5)$ priors to behave similarly to non-informative priors (analogous to maximum likelihood estimation) in this setting, but have slightly more posterior stability and easier sampling [10]. As an alternative, we used horseshoe priors for strong regularization on the full 4-digit ICD-9 code set. Finally, we place a hierarchical prior on the coefficients so that coefficients associated with 4-digit codes are shrunk together towards the coefficients of the 3-digit codes.

The propensity score models were fit using Stan for Hamiltonian Monte Carlo sampling from each posterior [4]. The convergence of the samplers was assessed by examining trace plots and by the Gelman-Rubin “r-hat” statistic, where a value less than 1.1 indicated sufficient convergence [10]. To assess the out-of-sample fit of different models, we calculated the leave-one-out information criterion (LOO-IC). LOO-IC is an approximation to leave-one-out cross validation that can be computed from a point-wise log-likelihood matrix

using Pareto-smoothed importance sampling, a procedure implemented in R through the `loo` package [40][41]. A lower LOO-IC indicates a model with better predictive accuracy.

To assess adjustment for measured confounders, we calculated how well the propensity scores balanced the observed covariates by calculating standardized differences, defined as the difference in means between the weighted BMS and DES populations over their pooled standard deviation. Because we have a different weighted population for each propensity score draw, this quantity is calculated once for each draw, giving a standardized difference distribution for each variable [1]. We assess the overall balance by determining the percentage of covariates with posterior mean standardized difference falling within $(-10,10)$ [1]. Ideally, the 95% intervals for standardized differences will also be contained within $(-10,10)$ for most variables.

We assessed how well each method met our falsifiability endpoints and their causal effect estimates for 2-day, 30-day, and 1-year TVR. We were particularly interested in the coverage and width of the 95% posterior intervals and whether they included zero.

4.3 Results

After eliminating diagnoses with fewer than 10 observed occurrences, we were left with a total of 334 4-digit ICD-9 confounders. There were fewer than 10 patients with individual codes in the “psychoses” Elixhauser category, so the Elixhauser indicator sets included 29 diagnoses rather than 30.

Grouping by the first 3-digits of the ICD-9 codes resulted in 69 groups containing at least 2 4-digit codes and 117 single 4-digit codes with no higher level group. The condition with the largest number of 4-digit codes in our sample was acute myocardial infarctions, i.e. heart attacks, and contained 10 4-digit codes. Thirty-seven of the 69 groups had only 2 4-digit codes. The complete list of all outcomes and confounders along with their prevalences (or means and standard deviations) appears in our appendix below.

Table 1 displays characteristics of the propensity score models including number of claims variables used, LOO-IC, and summaries of covariate balance. In terms of fit, Horseshoe regularization achieved the lowest LOO-IC score, indicating it had the best predictive accuracy. The Elixhauser indicator also had good LOO-IC, achieved through a considerable reduction in the dimension of the regression, though further reduction into a score degraded the quality of the fit considerably as indicated by a higher LOO-IC for Elixhauser continuous. The hierarchical prior provided lower LOO-IC than either the 4-digit or 3-digit $t_5(0, 2.5)$ prior approaches.

In terms of balance, the Elixhauser methods and 3-digit codes with weakly informative priors failed to bring posterior mean standardized difference for all variables to within $(-10,10)$. Regressions using the full claims set with weakly informative or hierarchical priors had many confounders for which the upper (or lower) 95% posterior interval limit is beyond 10% (or smaller than -10%) after weighting. In contrast, the confounder balance when using a horseshoe prior results in only 6 variables with standardized difference posterior intervals outside the 10% window.

Figure 2 illustrates the behavior of the different approaches to regularization when using the full claims set for two groups (e.g., two 3-digit diagnosis codes): cardiac dysrhythmias and drug/alcohol abuse. For cardiac dysrhythmias, the group mean for the hierarchical prior coefficients is very close to zero because the associations of the individual codes have opposite signs on the logit of the probability of receiving a DES. The hierarchical prior coef-

Approach	# Claims Variables	LOO-IC (SE)	Means $\notin$ (-10,10)	CIs $\notin$ (-10, 10)
Unadjusted*	0	11327 (54)	196	NA
<i>Elixhauser Methods</i>				
$t_5(0, 2.5)$ Prior Indicator Variables	29	9537 (94)	3	11
$t_5(0, 2.5)$ Prior Continuous Score	1	9640 (92)	4	12
<i>Full Claims Set</i>				
$t_5(0, 2.5)$ Prior 3-Digits	186	9662 (100)	1	59
$t_5(0, 2.5)$ Prior 4-Digits	334	9741 (107)	0	150
Horseshoe Prior 4-Digits	334	9468 (88)	0	6
Hierarchical Prior 3 and 4-Digits	334	9639 (104)	0	74

Table 1: **Characteristics of propensity score models.** Number of claims derived variables used in model, leave-one-out information criterion, and summaries of covariate balance for unadjusted analysis and each propensity score analysis. Column “Means $\notin$ (-10,10)” tabulates the number of variables with posterior mean standardized differences outside the interval -10 to 10, which is typically considered good balance. Column “CIs $\notin$ (-10, 10)” tabulates number of variables where some part of the 95% posterior interval lies outside (-10,10). LOO-IC = leave-one-out information criterion; SE = standard error. *: This is a model with an intercept only for the purpose of calculating LOO-IC and standardized differences reflect covariates in unweighted population.

ficients are slightly pooled towards their group mean, while coefficients under the horseshoe are pulled towards zero, in some cases quite sharply. The weakly informative coefficients are universally larger in magnitude than the others. A similar pattern emerges for drug/alcohol abuse coefficients. The group overall is associated with a lower probability of receiving a DES, with some granular codes exhibiting much smaller or larger effects. For both coefficient groups, a weakly informative prior using a single indicator for the 3-digit code category fails to capture the nuances apparent at lower levels.

Table 2 presents posterior summaries of causal effects in the Mass-DAC data under each method. The unadjusted difference in means between the two stent groups exhibits considerable confounding of both 30-day and 1-year mortality with large risk reductions for DES of 3.4% and 6.9% respectively. The Elixhauser approaches were the only models unable to adjust away the difference in 2-day mortality, though margins were slim. For 30-day mortality, only the full claims sets with $t_5(0, 2.5)$ or hierarchical priors estimated no significant effect. In contrast, aggressively regularizing using horseshoe priors yielded a 30-day mortality benefit to DES that is indicative of residual confounding. On one-year mortality, the model utilizing the hierarchical prior distribution for the ICD-9 coefficients achieved a considerable reduction from the large unadjusted difference, and improved over the comparably large effect estimates generated from the Elixhauser and horseshoe prior approaches. However no method included zero in its 95% credible interval, conflicting with findings from randomized trials. As expected, all methods produced a larger benefit to TVR than indicated by the unadjusted difference.

The Elixhauser indicator, Elixhauser continuous, and horseshoe approaches had the lowest uncertainty with posterior interval widths of 1.9, 1.8, and 2.0 respectively. The weakly informative 3-digit approach had an average width of 2.3, while the weakly informative 4-digit approach had an average of width of 3.0. The hierarchical prior approach had an average credible interval width of 2.5, about 20% tighter than the 4-digit approach. The weakly informative and hierarchical specifications thus incur more variance than the smaller or more regularized Elixhauser and horseshoe approaches, which is ultimately reflected in




Figure 2: **Selected posterior mean coefficients.** Values on log-odds scale for 4-digit ICD-9 coefficients with weakly informative priors (orange circle), horseshoe (red triangle), and hierarchical (blue diamond) specifications. Group mean defined by 3-digit code for hierarchical (blue dashed line) and 3-digit weakly informative coefficient estimate (green dashed line) are also shown. Left panel shows coefficients for code group 427, cardiac dysrhythmias; right panel shows coefficients for code group 305, drug or alcohol abuse.

the estimated distributions of causal parameters.

Approach	Mortality			
	2-Day	30-Day	1-Year	1-Year TVR
Unadjusted	-1.0 (-1.4, -0.6)	-3.4 (-4.1, -2.7)	-6.9 (-8.0, -5.7)	-2.4 (-3.7, -1.3)
<i>Elixhauser Methods</i>				
$t_5(0, 2.5)$ Prior Indicator Variables	-0.4 (-0.7, -0.1)	-1.2 (-2.0, -0.5)	-3.1 (-4.3, -1.8)	-4.3 (-5.7, -2.9)
$t_5(0, 2.5)$ Prior Continuous Score	-0.4 (-0.7, -0.1)	-1.3 (-2.0, -0.6)	-3.3 (-4.5, -2.1)	-4.5 (-5.9, -3.0)
<i>Full Claims Set</i>				
$t_5(0, 2.5)$ Prior 3-Digits	-0.3 (-0.7, 0.1)	-1.0 (-2.0, 0.2)	-2.7 (-4.1, -1.1)	-4.4 (-6.1, -2.8)
$t_5(0, 2.5)$ Prior 4-Digits	-0.3 (-0.7, 0.0)	-0.7 (-2.0, 1.2)	-2.5 (-4.5, -0.3)	-3.6 (-5.5, -1.8)
Horseshoe Prior 4-Digits	-0.3 (-0.7, 0.0)	-1.1 (-1.9, -0.3)	-3.0 (-4.3, -1.7)	-4.1 (-5.6, -2.6)
Hierarchical Prior 3 and 4-Digits	-0.3 (-0.7, 0.1)	-0.9 (-2.0, 0.5)	-2.6 (-4.2, -0.8)	-3.8 (-5.5, -2.0)

Table 2: **Casual effect estimates for selected outcomes.** Risk differences for various methods applied to the coronary stent study. Risk differences (95% Credible Intervals) estimated from the Mass-DAC data show benefit of DES compared to BMS in terms of reduced percent chance of given outcome ($\Delta = p_{\text{DES}} - p_{\text{BMS}}$). TVR = target vessel revascularization.

5 Discussion

In this paper, we proposed the use of regularizing prior distributions rather than comorbidity indices for data reduction strategies in high-dimensional causal inference problems in health care. We found that Bayesian priors improved inferences over comorbidity indices both in terms of out-of-sample fit or the ability to meet falsifiability endpoints. These prior distributions are able to work with a large number of claims variables while controlling variance.

To compare multiple methods with and without hierarchical modeling of ICD-9 codes, we estimated causal effects of DES compared to BMS on a number of outcomes. Specifically, we were looking for our methods to adjust away any difference in 30-day mortality and, ideally, 1-year mortality as well. RCTs have consistently shown no survival benefit to DES, but this has proved to be very difficult to replicate in observational analyses [2][42][34][14]. By using claims data and either a hierarchical ICD-9 or flat weakly informative propensity score model, we succeeded in removing the specious benefit shown to 2-day and 30-day mortality, which were very likely to be caused by residual confounding. In contrast, models using Elixhauser comorbidity indicators, a continuous Elixhauser score, or sparsity-inducing horseshoe priors on the full claims set did not manage to adjust away confounding on 30-day mortality. The fact that these dimension reducing approaches generated the best out-of-sample fit in terms of LOO-IC indicates the importance of using additional checks like falsifiability endpoints in causal inference problems, rather than relying solely on scores of predictive ability.

Unfortunately no method succeeded in removing all confounding of 1-year mortality, but the hierarchical and weakly informative prior distributions had comparably more success, estimating small effect sizes and 95% credible intervals close to 0. Residual confounding has plagued past observational studies of DES and BMS and it appears that, while claims data are able to account for important confounders not found in registries, we are still lacking the necessary variables to completely remove confounding [14]. Likely unmeasured confounders include socio-economic variables, which past studies have shown affect adherence to dual-antiplatelet therapy and ultimately a doctor’s decision to use a DES versus a BMS in usual care settings [33].

While generating similar posterior means, the hierarchical prior approach had considerably smaller variance than the unregularized approach, with about 20% tighter credible intervals on average. All methods showed, as expected, a considerable benefit to DES on the TVR outcome.

We restricted our attention to Bayesian propensity scores throughout, only changing the variable selection or regularization strategy for comparability. However, hierarchical ICD-9 code modeling is not limited to Bayesian propensity scores and could easily be adapted to Bayesian outcome modeling or frequentist techniques. Whatever the specifics, our work demonstrates that investigators performing causal inference in non-randomized settings could benefit by ignoring comorbidity indices and using more claims variables in their analyses. Comorbidity indices will continue to have utility for risk stratification and parsimonious assessments in clinical practice, but as we have shown, there are better alternatives for confounder adjustment.

6 Acknowledgments

Mr Spertus and Dr Normand were supported by a grant from the US Food and Drug Administration (U01FD004493). Mr Spertus, Dr Normand, and Dr Adhikari were supported by a grant from the National Institute of General Medical Sciences (1R01GM111339).

We are indebted to the Massachusetts Department of Public Health (MDPH) and the Massachusetts Center for Health Information and Analysis Case Mix Databases (CHIA) for the use of their data.

References

- [1] Peter C Austin and Elizabeth A Stuart. Moving towards best practice when using inverse probability of treatment weighting (iptw) using the propensity score to estimate causal treatment effects in observational studies. *Statistics in Medicine*, 34:3661–3679, 2015.
- [2] Kaare H. Bonaa, Jan Mannsverk, Rune Wiseth, Lars Aaberge, Yngvar Myreng, Ottar Nygard, Dennis W. Nilsen, Nils-Einar Klow, Michael Uchto, Thor Trovik, Bjorn Bendz, Sindre Stavnes, Reidar Bjornerheim, Alf-Inge Larsen, Morten Slette, Terje Steigen, Ole J. Jakobsen, Oyvind Bleie, Eigil Fossum, Tove A. Hanssen, Oystein Dahl-Eriksen, Inger Njolstad, Knut Rasmussen, Tom Wilsgaard, and Jan E. Nordrehaug. Drug-eluting or bare-metal stents for coronary artery disease. *The New England Journal of Medicine*, 2016.
- [3] M Alan Brookhart, Sebastian Schneeweiss, Kenneth J Rothman, Robert J Glynn, Jerry Avorn, and Til Sturmer. Variable selection for propensity score models. *Practice of Epidemiology*, 163:1149–1156, 2006.
- [4] Bob Carpenter, Andrew Gelman, Matt Hoffman, Daniel Lee, Ben Goodrich, Michael Betancourt, Michael A Brubaker, Jiqiang Guo, Peter Li, and Allen Riddell. Stan: a probabilistic programming language. *Journal of Statistical Software*, In Press, 2016.
- [5] Carlos M Carvahlo, Nicholas G Polson, and James G Scott. The horseshoe estimator for sparse signals. *Biometrika*, 97:465–480, 2010.
- [6] Massachusetts Data Analysis Center. Cardiac study annual reports. 2017.
- [7] Mary E Charlson, Peter Pompei, Kathy L Ales, and C Ronald MacKenzie. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Diseases*, 40:373–383, 1987.
- [8] Anne Elixhauser, Claudia Steiner, D Robert Harris, and Rosanna M Coffey. Comorbidity measures for use with administrative data. *Medical Care*, 36:8–27, 1998.
- [9] Centers for Disease Control and Prevention. International classification of diseases, ninth revision, clinical modification (icd-9-cm). <https://www.cdc.gov/nchs/icd/icd9cm.htm>.
- [10] Andrew Gelman, John B Carlin, Hal S Stern, David B Dunson, Aki Vehtari, and Donald B Rubin. *Bayesian Data Analysis: Third Edition*. CRC Press, 2014.
- [11] Elizabeth A Gilbert, Robert T Krafty, Richard J Bleicher, and Brian L Egleston. On the use of summary comorbidity measures for prognosis and survival treatment effect estimation. *Health Services Outcomes Research Methods*, <https://doi.org/10.1007/s10742-017-0171-2>, 2017.
- [12] Lisa Iezzoni. Assessing quality using administrative data. *Annals of Internal Medicine*, 127:666–674, 1997.

- [13] Mark S Levenson and Lilly Q Yue. Regulatory issues of propensity score methodology application to drug and device safety studies. *Journal of Biopharmaceutical Statistics*, 23:110–121, 2012.
- [14] Laura Mauri, Treacy S Silbaugh, Robert E Wolf, Katya Zelevinsky, Ann Lovett, Zheng Zhou, Frederic S Resnic, and Sharon-Lise T Normand. Long-term clinical outcomes after drug-eluting and bare-metal stenting in massachusetts. *Circulation*, 118:1817–1827, 2008.
- [15] Daniel F McCaffrey, Greg Ridgeway, and Andrew R Morral. Propensity score estimation with boosted regression for evaluating causal effects in observational studies. *Psychological Methods*, 9:403–425, 2004.
- [16] Lawrence C McCandless, Paul Gustafson, and Peter C Austin. Bayesian propensity score analysis for observational data. *Statistics in Medicine*, 28:94–112, 2009.
- [17] Patrick McCormick and Thomas Joseph. Package ‘medicalrisk’. 2016.
- [18] Sharon-Lise T Normand, Carl N Morris, Karen S Fung, Barbara J McNeil, and Arnold M Epstein. Development and validation of a claims based index for adjusting for risk of mortality: the case of acute myocardial infarction. *Journal of Clinical Epidemiology*, 48:229–243, 1995.
- [19] Tomi Peltola, Aki S Havulinna, Veikko Salomaa, and Aki Vehtari. Hierarchical bayesian survival analysis and projective covariate selection in cardiovascular event risk prediction. *Proceedings of the 11th Annual UAI Conference on Bayesian Modeling Applications Workshop*, 1218:79–88.
- [20] Adler Perotte, Rimma Pivovarov, Karthik Natarajan, Nicole Weiskopf, Frank Wood, and Noemie Elhadad. Diagnosis code assignment: models and evaluation metrics. *Journal of the American Medical Informatics Association*, 21:231–237, 2014.
- [21] Maya L Petersen, Kristin E Porter, Susan Gruber, Yue Wang, and Mark J van der Laan. Diagnosing and responding to violations in the positivity assumption. *Statistical Methods in Medical Research*, 21:31–54, 2010.
- [22] Juho Piironen and Aki Vehtari. Projection predictive variable selection using stan + r. 2015.
- [23] Hude Quan, Vijaya Sundarajan, Patricia Halfon, Andrew Fong, Bernard Burnand, Jean-Christophe Luthi, L Duncan Saunders, Cynthia A Beck, Thomas E Feasby, and William A Ghali. Coding algorithms for defining comorbidities in icd-9 and icd-10 administrative data. *Medical Care*, 43:1130–1139, 2005.
- [24] Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70:41–55, 1983.
- [25] Donald B Rubin. For objective causal inference, design trumps analysis. *The Annals of Applied Statistics*, 2:808–840, 2008.
- [26] Olli Saarela. On bayesian estimation of marginal structural models. *Biometrics*, 71:279–301, 2015.

- [27] Mary Vaughan Sarrazin and Gary E Rosenthal. Finding pure and simple truths with administrative data. *Journal of the American Medical Association*, 307:1433–1435, 2012.
- [28] Sebastian Schneeweiss, Jeremy A Rassen, Robert J Glynn, Jerry Avorn, Helen Mogun, and M Alan Brookhart. High-dimensional propensity score adjustment in studies of treatment effects using health care claims data. *Epidemiology*, 20:512–522, 2009.
- [29] Mansour TA Sharabiani, Paul Aylin, and Alex Bottle. Systematic review of comorbidity indices for administrative data. *Medical Care*, 50:1109–1118, 2012.
- [30] Anima Singh. Exploiting hierarchical and temporal information in building predictive models from ehr data. *PhD Dissertation for Massachusetts Institute of Technology*, 2015.
- [31] Jacob V Spertus and Sharon-Lise T Normand. Bayesian propensity scores for high-dimensional causal inference: A comparison of drug-eluting to bare-metal coronary stents. *Biometrics*, page In submission, 2018.
- [32] Jacob V Spertus, Sharon-Lise T Normand, Robert Wolf, Matt Cioffi, Ann Lovett, and Sherri Rose. Assessing hospital performance following percutaneous coronary intervention using big data. *Circulation: Cardiovascular Quality and Outcomes*, In submission, 2016.
- [33] John A Spertus, Richard Kettelkamp, Clifton Vance, Carole Decker, Philip G Jones, John S Rumsfeld, John C Messenger, Sanjaya Khanal, Eric D Peterson, Richard G Bach, Harlan M Krumholz, and David J Cohen. Prevalence, predictors, and outcomes of premature discontinuation of thienopyridine therapy after drug-eluting stent placement: Results from the premier registry. *Circulation*, 113:2803–2809.
- [34] Christoph Stettler, Simon Wandel, Sabin Allemann, Adnan Kastrati, Marie Claude Morice, Albert Schomig, Matthias E Pfisterer, Gregg W Stone, Martin B Leon, Jose Suarez de Leon, Jean-Jacques Goy, Seung-Jung Park, Manel Sabate, Maarten J Suttorp, Henning Kelbaek, Christian Spaulding, Maurizio Menichelli, Paul Vermeer-sch, Maurits T Dirksen, Pavel Cervinka, Anna Sonia Petronio, Alain J Nordmann, Peter Diem, Bernhard Meier, Marcel Zwahlen, Stephan Reichenbach, Sven Trelle, Stephan Windecker, and Peter Juni. Outcomes associated with drug-eluting and bare-metal stents: a collaborative network meta-analysis. *The Lancet*, 370:937–948, 2007.
- [35] Elizabeth A Stuart, Eva DuGoff, Michael Abrams, David Salkever, and Donald Steinwachs. Estimating causal effects in observational studies using electronic health data: Challenges and (some) solutions. *EGEMS*, 1, 2013.
- [36] Zeeshan Syed, Ilan Rubinfeld, Joe H Patton, Jennifer Ritz, Jack Jordan, Andrea Doud, and Vic Velanovich. Using procedural codes to supplement risk adjustment: a nonparametric learning approach. *Journal of the American College of Surgeons*, 212:1086–1093, 2011.
- [37] Jack V Tu, James Bowen, Maria Chiu, Dennis T Ko, Peter C Austin, Yaohua He, Robert Hopkins, Jean-Eric Tarride, Gord Blackhouse, Charles Lazzam, Eric A Cohen,

- and Ron Goeree. Effectiveness and safety of drug-eluting stents in ontario. *The New England Journal of Medicine*, 357:1393–1402, 2007.
- [38] Mark J. van der Laan and Sherri Rose. *Targeted Learning: Causal Inference for Observational and Experimental Data*. Springer, 2011.
- [39] Carl van Walraven, Peter C Austin, Alison Jennings, Hude Quan, and Alan J Forster. A modification of the elixhauser comorbidity measures into a point system for hospital death using administrative data. *Medical Care*, 47:626–633, 2009.
- [40] Aki Vehtari, Andrew Gelman, and Jonah Gabry. Practical bayesian model evaluation using leave-one-out cross-validation and waic. *Statistics and Computing*, 27:1413–1432, 2017.
- [41] Aki Vehtari, Andrew Gelman, Jonah Gabry, Juho Piironen, and Ben Goodrich. Package ‘loo’. 2017.
- [42] Lakshmi Venkitachalam, Yang Lei, Elizabeth A Magnuson, Paul S Chan, Joshua M Stolker, Kevin F Kennedy, Neal S Kleiman, and David J Cohen. Survival benefit with drug-eluting stents in observational studies: Fact or artifact? *Circulation*, 4:587–594, 2011.

A Covariate Tables

A.1 Registry Covariates

Table 3: Prevalence or mean (SD) of outcomes in the Mass-DAC registry. “Cases per Site” gives min, mean (SD), and max of numbers treated at the 25 hospitals, overall and in each treatment group. The 31 coronary artery segment indicators are not shown. HMO = health maintenance organization; CAD = coronary artery disease; CVD = cardiovascular disease; PAD = peripheral artery disease; CABG = coronary artery bypass graft; STEMI = ST-elevated myocardial infarction; LVSD = left ventricular systolic dysfunction; NYHA = New York heart association; LAD = left anterior descending (artery); RCA = right coronary artery.

	Overall	BMS	DES
Number Treated (n)	8718	3081	5637
Registry Outcomes			
% 2 Day Mortality	0.50	1.14	0.16
% 30 Day Mortality	2.01	4.19	0.82
% 1 Year TV Revascularization	7.38	8.96	6.51
% 1 Year Mortality	5.75	10.19	3.32
Registry Confounders			
% Male	69.39	68.16	70.05
Mean (SD) Age in Years	64.66 (12.5)	66.37 (11.7)	63.72 (13.5)
Mean (SD) Height in cm	170.67 (10.5)	170.41 (10.6)	170.81 (10.4)
Mean (SD) Weight in kg	86.48 (20.3)	84.89 (19.9)	87.34 (20.8)
Race			

% Caucasian	91.26	90.72	91.56
% Black	3.79	4.87	3.19
% Asian	2.31	2.43	2.24
% Native American	< 0.1	< 0.3	< 0.2
% Native Pacific	< 0.1	< 0.3	< 0.2
% Hispanic or Latino	4.55	4.8	4.42
Payor			
% Government	51.57	59.17	47.42
% None	2.56	4.12	1.7
% Non-US	< 0.1	< 0.3	< 0.2
% Private Commercial or HMO	45.82	36.68	50.82
% Smoker	74.68	70.85	76.78
% Hypertension	78.3	76.6	79.23
% Dyslipidemia	81.82	77.96	83.93
% Diabetes	32.16	30.38	33.14
% Family History CAD	28.96	24.08	31.63
% Chronic Lung Disease	14.21	16.42	13
% Current Dialysis	1.82	2.21	1.61
% Prior CVD	10.23	11.33	9.63
% Prior PAD	11.96	12.76	11.53
% Prior Myocardial Infarction	26.67	25.84	27.12
% Prior Heart Failure	10.61	12.46	9.6
% Prior Valve Surgery	1.56	2.34	1.14
% Prior PCI	27.56	19.77	31.83
% Prior CABG	12.3	11.46	12.76
% Prior Cardiogenic Shock	1.9	3.89	0.82
% Prior Cardiac Arrest	2.29	4.15	1.28
CAD Presentation			
% No Angina	5.32	5.78	5.07
% Symptom Unlikely to be Ischemic	1.00	1.17	0.90
% Stable Angina	11.88	5.81	15.2
% Unstable Angina	30.19	23.4	33.9
% Non-STEMI	27.23	28.17	26.72
% STEMI	24.37	35.67	18.2
% Thrombolytic Therapy	0.8	1.23	0.57
% Cardiomyopathy or LVSD	9.36	11.1	8.41
Anginal Canadian Classification			
% 0	9.64	11.65	8.53
% I	2.4	1.59	2.84
% II	12.2	7.66	14.69
% III	28.83	24.15	31.38
% IV	46.94	54.95	42.56
Anti-Anginal Medications			
% Beta Blockers	57.79	54.56	59.55
% Calcium Channel Blockers	11.44	11.39	11.46
% Long Acting Nitrates	11.49	9.41	12.63

% Ranolazine	0.62	0.45	0.71
% Other Agent	1.4	1.01	1.61
NYHA Class			
% 0	87.57	83.51	89.78
% I	0.91	0.78	0.98
% II	3.06	3.67	2.73
% III	4.16	5.39	3.49
% IV	4.3	6.65	3.02
% Compassionate Use	0.91	2.01	0.3
% Cardiogenic Shock	1.85	3.8	0.78
% Mechanical Ventricular Support	0.58	1.14	0.28
% Ejection Fraction < 30%	2.88	3.7	2.43
Coronary Anatomy			
% Left Dominant	8.30	8.08	8.42
% Right Dominant	86.19	85.88	86.36
% Left Dominant	5.51	6.04	5.22
% Left Main Disease	5.93	6.52	5.61
Mean Left Main Stenosis (SD)	7.98 (19.8)	8.74(19.3)	7.57(20.7)
Mean Proximal LAD Stenosis (SD)	36 (39.7)	35.1(39.7)	36.5(39.5)
Mean Mid-Distal LAD Stenosis (SD)	48.2 (39.7)	47.64(39.7)	48.51(39.6)
Mean Circumflex Stenosis (SD)	47.95 (40.8)	47.94(40.7)	47.96(40.8)
Mean RCA Stenosis (SD)	60.66 (39.9)	65.25(40)	58.15(39.2)
Status			
% Urgent	52.49	48.78	54.51
% Emergent	26.69	38.30	20.35
% Other	20.82	12.92	25.14
PCI Indication			
% Immediate PCI for STEMI	21.80	31.39	16.55
% STEMI (Unstable, > 12 hours)	1.67	2.66	1.14
% STEMI (Stable, > 12 hours)	0.57	0.94	0.37
% STEMI (Stable, thrombolytics)	0.44	0.58	0.35
% STEMI (Rescue, failed thrombolytics)	0.46	0.75	0.3
% High risk Non-STEMI	48.99	45.86	50.7
% Staged	0.86	0.45	1.08
% Other	25.21	17.36	29.5
Mean (SD) Thrombectomies	0.12 (0.3)	0.16(0.3)	0.09(0.4)
Mean (SD) Total Stents Used	1.46 (0.8)	1.41 (0.8)	1.5 (0.8)
Mean (SD) Lesions	1.3 (0.6)	1.24 (0.6)	1.33 (0.5)
Mean (SD) Lesion Length	18.08 (10)	17 (10.4)	18.68 (9.3)
% In-stent Restenosis	6.96	2.34	9.49
% Chronic Total Occlusion	1.73	1.3	1.97
Cases per Site (25 Total)			
Min # Treated	47	12	14
Mean # Treated (SD)	349 (269)	123 (99)	225 (187)
Max # Treated	896	352	605

Table 4: Prevalence of present on admission ICD-9 diagnosis codes used in analysis. As with the analysis using a hierarchical prior, codes are grouped together if there were two or more 4-digit codes sharing a 3-digit grouping. w/o = without.

Diagnosis Code	Description (Present on Admission)	Prevalence
041	Bacterial infection of unspecified site	0.5
0411	Staphylococcus infection, unspecified	0.2
0414	Shiga toxin-producing E. coli (STEC)	0.3
070	Viral hepatitis	1.1
0703	Viral hepatitis B w/o mention of hepatic coma	0.2
0705	Acute hepatitis C w/o mention of hepatic coma	0.4
0707	Unspecified viral hepatitis C w/o hepatic coma	0.5
242	Thyrotoxicosis with or w/o goiter	0.3
2420	Toxic diffuse goiter w/o mention of thyrotoxic crisis or storm	0.1
2429	Thyrotoxicosis w/o mention of goiter, thyrotoxic crisis, or storm	0.2
244	Acquired hypothyroidism	7.4
2440	Postsurgical hypothyroidism	0.2
2448	Other specified acquired hypothyroidism	0.2
2449	Unspecified acquired hypothyroidism	7
250	Diabetes mellitus	30.8
2500	Diabetes mellitus w/o mention of complication	27
2501	Diabetes with ketoacidosis	0.2
2504	Diabetes with renal manifestations	1.5
2505	Diabetes with ophthalmic manifestations	1.1
2506	Diabetes with neurological manifestations	1.9
2507	Diabetes with peripheral circulatory disorders	0.2
2508	Diabetes with other specified manifestations	0.3
272	Disorders of lipoid metabolism	69.9
2720	Pure hypercholesterolemia	13.7
2721	Pure hyperglyceridemia	1
2722	Mixed hyperlipidemia	0.1
2724	Other and unspecified hyperlipidemia	59.4
2729	Unspecified disorder of lipoid metabolism	0.1
274	Gout	3.5
2740	Gouty arthropathy, unspecified	0.3
2749	Gout, unspecified	3.2
275	Disorders of mineral metabolism	0.9
2750	Hereditary hemochromatosis	0.1
2752	Disorders of magnesium metabolism	0.3
2753	Disorders of phosphorus metabolism	0.2
2754	Unspecified disorder of calcium metabolism	0.3
276	Disorders of fluid, electrolyte, and acid-base balance	4.5
2760	Hyperosmolality and/or hyponatremia	0.1
2761	Hyposmolality and/or hyponatremia	1.2
2762	Acidosis	0.7
2764	Mixed acid-base balance disorder	0.1
2765	Volume depletion, unspecified	0.7

2766	Transfusion associated circulatory overload	0.1
2767	Hyperpotassemia	0.7
2768	Hypopotassemia	1.1
280	Iron deficiency anemias	1.4
2800	Iron deficiency anemia secondary to blood loss (chronic)	0.2
2809	Iron deficiency anemia, unspecified	1.2
281	Other deficiency anemias	0.3
2810	Pernicious anemia	0.1
2819	Unspecified deficiency anemia	0.1
285	Other and unspecified anemias	5.5
2851	Acute posthemorrhagic anemia	0.1
2852	Anemia in chronic kidney disease	1.5
2858	Other specified anemias	0.1
2859	Anemia, unspecified	3.8
287	Purpura and other hemorrhagic conditions	1.3
2874	Posttransfusion purpura	0.1
2875	Thrombocytopenia, unspecified	1.2
294	Other organic psychotic conditions (chronic)	1
2941	Dementia w/o behavioral disturbance	0.2
2948	Other persistent mental disorders	0.8
296	Affective psychoses	1.4
2962	Major depressive affective disorder, single episode	0.2
2963	Major depressive affective disorder, recurrent episode	0.1
2968	Bipolar disorder, unspecified	0.9
2969	Unspecified episodic mood disorder	0.1
300	Neurotic disorders	6.8
3000	Anxiety state, unspecified	4.6
3004	Dysthymic disorder	2.2
305	Nondependent abuse of drugs	21.9
3050	Alcohol abuse, unspecified	2.2
3051	Tobacco use disorder	20.4
3052	Cannabis abuse, unspecified	0.9
3055	Opioid abuse, unspecified	0.2
3056	Cocaine abuse, unspecified	0.7
3059	Other, mixed, or unspecified drug abuse, unspecified	0.2
333	Extrapyramidal disease and abnormal movement disorders	0.6
3331	Essential and other specified forms of tremor	0.2
3339	Unspecified extrapyramidal disease and abnormal movement	0.4
338	Pain, not elsewhere classified	2
3382	Chronic pain due to trauma	1.8
3384	Chronic pain syndrome	0.2
348	Other conditions of brain	0.5
3481	Anoxic brain damage	0.4
3483	Encephalopathy, unspecified	0.1
355	Mononeuritis of lower limb	0.4
3558	Mononeuritis of lower limb, unspecified	0.1

3559	Mononeuritis of unspecified site	0.3
362	Other retinal disorders	1.4
3620	Background diabetic retinopathy	1
3625	Macular degeneration (senile), unspecified	0.4
396	Diseases of mitral and aortic valves	0.7
3962	Mitral valve insufficiency and aortic valve stenosis	0.3
3963	Mitral valve insufficiency and aortic valve insufficiency	0.3
401	Essential hypertension	62.4
4010	Malignant essential hypertension	0.1
4011	Benign essential hypertension	0.7
4019	Unspecified essential hypertension	61.6
410	Acute myocardial infarction	54.3
4100	AMI of anterolateral wall	2.7
4101	AMI of other anterior wall	6.4
4102	AMI of inferolateral wall	2.2
4103	AMI of inferoposterior wall	2.4
4104	AMI of other inferior wall	9.5
4105	AMI of other lateral wall	0.9
4106	True posterior wall infarction	0.2
4107	Subendocardial infarction	28.7
4108	AMI of other specified sites	0.3
4109	AMI of unspecified site	1.1
411	Other acute and subacute form of ischemic heart disease	19.8
4111	Intermediate coronary syndrome	19.3
4118	Acute coronary occlusion w/o myocardial infarction	0.5
414	Other forms of chronic ischemic heart disease	86.7
4140	Coronary atherosclerosis of unspecified vessel	85.6
4141	Aneurysm of heart (wall)	0.6
4142	Chronic total occlusion of coronary artery	9.5
4148	Other specified forms of chronic ischemic heart disease	4.9
4149	Chronic ischemic heart disease, unspecified	0.2
424	Other diseases of endocardium	6.3
4240	Mitral valve disorders	3.5
4241	Aortic valve disorders	2.8
425	Cardiomyopathy	2.6
4254	Other primary cardiomyopathies	2.5
4255	Alcoholic cardiomyopathy	0.1
426	Conduction disorders	5.2
4260	Atrioventricular block, complete	0.8
4261	Atrioventricular block, unspecified	1.3
4262	Left bundle branch hemiblock	0.1
4263	Other left bundle branch block	1.4
4264	Right bundle branch block	1.5
4265	Bundle branch block, unspecified	0.4
427	Cardiac dysrhythmias	17.3
4270	Paroxysmal supraventricular tachycardia	0.2

4271	Paroxysmal ventricular tachycardia	2.4
4273	Atrial fibrillation	10
4274	Ventricular fibrillation	1.6
4275	Cardiac arrest	1.3
4276	Premature beats, unspecified	0.7
4278	Sinoatrial node dysfunction	4.3
428	Heart failure	15.5
4280	Congestive heart failure, unspecified	14.5
4282	Systolic heart failure, unspecified	5.6
4283	Diastolic heart failure, unspecified	3
4284	Combined systolic and diastolic heart failure, unspecified	1.6
429	Ill-defined descriptions and complications of heart disease	1.8
4293	Cardiomegaly	0.7
4298	Other disorders of papillary muscle	0.2
4299	Heart disease, unspecified	0.9
433	Occlusion and stenosis of precerebral arteries	1.8
4331	Occlusion/stenosis of carotid artery	1.8
4333	Occlusion/stenosis of multiple precerebral arteries	0.5
440	Atherosclerosis	2.6
4400	Atherosclerosis of aorta	0.3
4401	Atherosclerosis of renal artery	0.4
4402	Atherosclerosis of native arteries of the extremities	1.9
4408	Atherosclerosis of other specified arteries	0.2
441	Aortic aneurysm and dissection	1.2
4412	Thoracic aneurysm w/o mention of rupture	0.1
4414	Abdominal aneurysm w/o mention of rupture	1.1
443	Other peripheral vascular disease	5.6
4430	Raynaud's syndrome	0.3
4438	Peripheral angiopathy in diseases classified elsewhere	0.3
4439	Peripheral vascular disease, unspecified	5
444	Arterial embolism and thrombosis	0.2
4442	Arterial embolism and thrombosis of upper extremity	0.1
4448	Embolism and thrombosis of iliac artery	0.1
447	Other disorders of arteries and arterioles	0.5
4471	Stricture of artery	0.4
4477	Aortic ectasia, unspecified site	0.1
458	Hypotension	1.7
4580	Orthostatic hypotension	0.2
4582	Hypotension of hemodialysis	0.1
4588	Other specified hypotension	0.6
4589	Hypotension, unspecified	0.7
493	Asthma	4.9
4932	Chronic obstructive asthma, unspecified	1.4
4939	Asthma,unspecified type, unspecified	3.6
518	Other diseases of lung	3
5180	Pulmonary collapse	0.4

5184	Acute edema of lung, unspecified	0.1
5188	Acute respiratory failure	2.5
530	Diseases of esophagus	17.3
5301	Esophagitis, unspecified	0.3
5308	Esophageal reflux	17.1
536	Disorders of function of stomach	0.3
5363	Gastroparesis	0.1
5368	Dyspepsia and other specified disorders of function of stomach	0.2
564	Functional digestive disorders, not elsewhere classified	1.3
5640	Constipation, unspecified	0.8
5641	Irritable bowel syndrome	0.6
571	Chronic liver disease and cirrhosis	0.6
5715	Cirrhosis of liver w/o mention of alcohol	0.2
5718	Other chronic nonalcoholic liver disease	0.4
578	Gastrointestinal hemorrhage	0.4
5781	Blood in stool	0.1
5789	Hemorrhage of gastrointestinal tract, unspecified	0.2
584	Acute renal failure	2.8
5845	Acute kidney failure with lesion of tubular necrosis	0.3
5849	Acute kidney failure, unspecified	2.5
585	Chronic renal failure	9.9
5852	Chronic kidney disease, Stage II (mild)	0.4
5853	Chronic kidney disease, Stage III (moderate)	1.9
5854	Chronic kidney disease, Stage IV (severe)	0.6
5855	Chronic kidney disease, Stage V	0.2
5856	End stage renal disease	1.8
5859	Chronic kidney disease, unspecified	5.1
599	Other disorders of urethra and urinary tract	1.9
5990	Urinary tract infection, site not specified	1.5
5996	Urinary obstruction, unspecified	0.1
5997	Hematuria, unspecified	0.3
696	Psoriasis and similar disorders	0.8
6960	Psoriatic arthropathy	0.2
6961	Other psoriasis	0.6
707	Chronic ulcer of skin	0.8
7070	Pressure ulcer, unspecified site	0.3
7071	Ulcer of lower limb, unspecified	0.5
7072	Pressure ulcer, unspecified stage	0.3
710	Diffuse diseases of connective tissue	0.4
7100	Systemic lupus erythematosus	0.2
7101	Systemic sclerosis	0.2
715	Osteoarthritis and allied disorders	4
7153	Osteoarthritis, localized	0.8
7159	Osteoarthritis, unspecified	3.2
721	Spondylosis and allied disorders	0.7
7210	Cervical spondylosis w/o myelopathy	0.2

7213	Lumbosacral spondylosis w/o myelopathy	0.3
7219	Spondylosis of unspecified site, w/o mention of myelopathy	0.2
722	Intervertebral disc disorders	0.2
7221	Displacement of lumbar intervertebral disc w/o myelopathy	0.1
7225	Degeneration of thoracic or thoracolumbar intervertebral disc	0.1
723	Other disorders of cervical region	0.3
7231	Cervicalgia	0.2
7234	Brachial neuritis or radiculitis NOS	0.1
724	Other and unspecified disorders of back	3.4
7240	Spinal stenosis, unspecified region	0.9
7242	Lumbago	1.1
7243	Sciatica	0.2
7245	Backache, unspecified	1.3
729	Other disorders of soft tissues	1.3
7291	Myalgia and myositis, unspecified	0.8
7295	Pain in limb	0.1
7298	Swelling of limb	0.4
733	Other disorders of bone and cartilage	2.1
7330	Osteoporosis, unspecified	1.7
7339	Disorder of bone and cartilage, unspecified	0.4
780	General symptoms	3.5
7800	Coma	0.2
7802	Syncope and collapse	0.7
7803	Febrile convulsions (simple), unspecified	0.2
7804	Dizziness and giddiness	0.4
7805	Sleep disturbance, unspecified	1.4
7806	Fever, unspecified	0.3
7807	Chronic fatigue syndrome	0.2
7809	Fussy infant (baby)	0.3
785	Symptoms involving cardiovascular system	2.4
7850	Tachycardia, unspecified	0.2
7852	Undiagnosed cardiac murmurs	0.3
7855	Shock, unspecified	2
786	Symptoms involving chest or respiratory system	1.5
7860	Respiratory abnormality, unspecified	0.6
7862	Cough	0.1
7863	Hemoptysis	0.1
7865	Chest pain, unspecified	0.7
787	Symptoms involving digestive system	0.8
7870	Nausea with vomiting	0.2
7872	Dysphagia, unspecified	0.3
7879	Diarrhea	0.3
788	Symptoms involving urinary system	0.7
7882	Retention of urine, unspecified	0.3
7883	Urinary incontinence, unspecified	0.3
7884	Urinary frequency	0.1

790	Nonspecific findings on examination of blood	2.4
7902	Impaired fasting glucose	1.8
7904	Elevation of levels of transaminase or LDH	0.2
7906	Other abnormal blood chemistry	0.1
7909	Abnormal arterial blood gases	0.3
794	Nonspecific abnormal results of function studies	1.4
7943	Abnormal cardiovascular function study, unspecified	1.2
7948	Nonspecific abnormal results of function study of liver	0.2
996	Complications peculiar to certain specified procedures	5
9967	Other complications due to unspecified device, implant, and graft	4.8
9968	Complications of transplanted organ, unspecified	0.2
E87		0.5
E878	Transplant of whole organ causing abnormal reaction or complication	0.4
E879	Cardiac catheterization as the cause of reaction or complication	0.1
2780	Obesity, unspecified	12.3
4039	Hypertensive chronic kidney disease	9.1
4139	Other and unspecified angina pectoris	7.9
496	Chronic airway obstruction, unspecified	7.3
3272	Organic sleep apnea, unspecified	4.9
6000	Hypertrophy of prostate without urinary obstruction	4.9
311	Depressive disorder, unspecified	4.8
4168	Other chronic pulmonary heart diseases	1.9
3572	Polyneuropathy in diabetes	1.8
7140	Rheumatoid arthritis	1.5
7169	Arthropathy, unspecified, site unspecified	1.5
3659	Unspecified glaucoma	1.4
486	Pneumonia, organism unspecified	1.3
5621	Diverticulosis of colon	1.2
3970	Diseases of tricuspid valve	1.2
4912	Obstructive chronic bronchitis without exacerbation	1.1
2886	Leukocytosis, unspecified	1.1
3459	Epilepsy	1.1
5533	Diaphragmatic hernia	0.9
3569	Unspecified peripheral neuropathy	0.9
3039	Alcohol dependence	0.9
V498	Asymptomatic postmenopausal status	0.8
2689	Vitamin D deficiency	0.8
3469	Migraine	0.8
5838	Nephritis and nephropathy	0.8
4598	Venous insufficiency	0.7
5939	Unspecified disorder of kidney and ureter	0.7
4928	Other emphysema	0.7
7194	Pain in joint	0.6
515	Postinflammatory pulmonary fibrosis	0.6
3899	Unspecified hearing loss	0.6
185	Malignant neoplasm of prostate	0.5

2777	Dysmetabolic syndrome	0.5
725	Polymyalgia rheumatica	0.5
3098	Posttraumatic stress disorder	0.5
6078	Balanitis xerotica obliterans	0.4
5119	Unspecified pleural effusion	0.4
3320	Paralysis agitans	0.4
2662	Other B-complex deficiencies	0.4
2898	Primary hypercoagulable state	0.4
3694	Legal blindness, as defined in U.S.A.	0.4
7990	Asphyxia	0.4
9981	Hemorrhage complicating a procedure	0.4
4029	Hypertensive heart disease without heart failure	0.4
5569	Ulcerative colitis	0.4
412	Old myocardial infarction	0.4
5070	Pneumonitis due to inhalation of food or vomitus	0.4
7840	Headache	0.3
7823	Edema	0.3
5559	Regional enteritis of unspecified site	0.3
V08	Asymptomatic human immunodeficiency virus	0.3
3669	Unspecified cataract	0.3
9971	Cardiac complications, not elsewhere classified	0.3
5355	Gastritis and gastroduodenitis	0.3
5888	Secondary hyperparathyroidism	0.3
V458	Aortocoronary bypass status	0.3
2041	Chronic lymphoid leukemia	0.3
4779	Allergic rhinitis	0.3
E849	Home accidents	0.3
2028	Other malignant lymphomas	0.2
2520	Hyperparathyroidism	0.2
5965	Hypertonicity of bladder	0.2
2387	Essential thrombocythemia	0.2
3310	Alzheimer's disease	0.2
4239	Disease of pericardium	0.2
3140	Attention deficit disorder without mention of hyperactivity	0.2
5339	Peptic ulcer of unspecified site	0.2
6929	Contact dermatitis and other eczema	0.2
5920	Calculus of kidney	0.2
9959	Systemic inflammatory response syndrome	0.2
135	Sarcoidosis	0.2
2918	Alcohol withdrawal	0.2
340	Malignant neoplasm of right main bronchus	0.2
6826	Cellulitis and abscess of leg, except foot	0.2
7890	Abdominal pain	0.2
042	Human immunodeficiency virus	0.2
4049	Hypertensive heart and chronic kidney disease, unspecified	0.2
4465	Giant cell arteritis	0.2

7288	Interstitial myositis	0.2
4824	Pneumonia due to Staphylococcus, unspecified	0.2
2410	Nontoxic uninodular goiter	0.2
0389	Unspecified septicemia	0.1
2841	Antineoplastic chemotherapy induced pancytopenia	0.1
2959	Unspecified schizophrenia, unspecified	0.1
3040	Opioid type dependence, unspecified	0.1
3540	Carpal tunnel syndrome	0.1
570	Acute necrosis of liver	0.1
0084	Intestinal infection due to staphylococcus	0.1
1629	Malignant neoplasm of bronchus and lung	0.1
1985	Secondary malignant neoplasm of bone and marrow	0.1
2030	Multiple myeloma	0.1
4739	Unspecified sinusitis (chronic)	0.1
501	Asbestosis	0.1
5589	Noninfectious gastroenteritis and colitis	0.1
5771	Chronic pancreatitis	0.1
5790	Celiac disease	0.1
7832	Loss of weight	0.1
1889	Malignant neoplasm of bladder	0.1
2019	Hodgkin's disease	0.1
2536	Other disorders of neurohypophysis	0.1
2554	Glucocorticoid deficiency	0.1
2713	Intestinal disaccharidase deficiencies and malabsorption	0.1
2904	Vascular dementia, uncomplicated	0.1
3860	Meniere's disease	0.1
4059	Unspecified renovascular hypertension	0.1
4151	Iatrogenic pulmonary embolism and infarction	0.1
490	Bronchitis, acute/chronic unspecified	0.1
6953	Rosacea	0.1
2113	Benign neoplasm of colon	0.1
3429	Hemiplegia affecting unspecified side	0.1
4370	Cerebral atherosclerosis	0.1
5742	Calculus of gallbladder	0.1
591	Hydronephrosis	0.1
7367	Deformity of ankle and foot, acquired	0.1
7812	Abnormality of gait	0.1
V125	History of unspecified circulatory disease	0.1
V158	History of failed moderate sedation	0.1
