On central-max-point tolerance graphs and some subclasses of interval catch digraphs

Sanchita Paul^{*} [†]and Shamik Ghosh[‡]

Abstract

Max-point-tolerance graphs (MPTG) were studied by Catanzaro et al. in 2017 and the same class of graphs were introduced in the name of *p*-BOX(1) graphs by Soto and Caro in 2015. In our paper we consider *central-maxpoint tolerance graphs* (central MPTG) by taking the points of MPTG as center points of their corresponding intervals. In course of study on this class of graphs we show that the class of central MPTG is same as the class of unit max-tolerance graphs. We prove the class of *unit central max-point tolerance graphs* is same as that of *proper central max-point tolerance graphs* and both of them are equivalent to the class of proper interval graphs. Next we introduce 50% *max-tolerance graphs* and separate this class from unit max-tolerance graph whereas for min-tolerance graphs 50% and unit denote the same graph class.

Interval catch digraphs (ICD) was introduced by Maehera in 1984. In his introducing paper Maehera proposed a conjecture for the characterization of *central interval catch digraph* (central ICD) in terms of forbidden subdigraphs. In this paper, we disprove the conjecture by showing counter examples. We find close relation between a central MPTG and a central ICD. Also we characterize this digraph by defining a suitable mapping from the vertex set to the real line. Next we study *oriented interval catch digraphs* (oriented ICD). We characterize augmented adjacency matrix of an oriented ICD when it is a tournament. Also we characterize an oriented ICD whose underlying graph is a tree. Further we obtain characterization of the adjacency matrix of a *proper interval catch digraph*. Another important result is to provide forbidden subdigraph is characterization of those *proper oriented interval catch digraphs* whose underlying undirected graph is chordal. In last section we discuss relationships between these classes of graphs and digraphs.

Keywords: Interval graph, proper interval graph, tolerance graph, max-tolerance graph, max-point tolerance graph, interval catch digraph, oriented digraph, tournament.

2010 Mathematical Subject Classification: Primary: 05C62, 05C75

1 Introduction

Intersection graphs have many important applications in problems related to real-world situations. Because of its diverse applications in network science, different kinds of intersection graphs were introduced in modeling various geometric objects. Among them interval graph is the most important one. A simple graph G = (V, E)is an *interval graph* if one can map each vertex into an interval on the real line so that any two vertices are adjacent if and only if their corresponding intervals intersect. The class of interval graphs was initially posed by Hajös in 1957 [11]. In 1959, the molecular biological scientist Benzer [3] used the model of interval graphs to obtain a physical map from information on pairwise overlaps of the fragments of DNA. Since then interval graphs were well studied by many people in Computer Science and Discrete Mathematics for its wide application. Many combinatorial problems have been solved for interval graphs in linear time.

Due to their lot of applications in theories and practical situations the graph class was generalized to several variations. In one direction it went in developing concepts of probe interval graphs [10], circular-arc graphs

^{*}Department of Mathematics, Jadavpur University, Kolkata - 700 032, India. sanchitajumath@gmail.com

[†](Corresponding author)

[‡]Department of Mathematics, Jadavpur University, Kolkata - 700 032, India. ghoshshamik@yahoo.com

[2], interval digraphs [17]. On the other hand in 1982, Golumbic and Monma introduced the concept of min-tolerance graphs (commonly known as tolerance graphs) [8]. We denote the length of an interval I on the real line by |I|. A simple undirected graph G = (V, E) is a min-tolerance graph if each vertex $u \in V$ corresponds to a real interval I_u and a positive real number t_u , called tolerance, such that uv is an edge of G if and only if $|I_u \cap I_v| \ge \min \{t_u, t_v\}$. Golumbic in [9] introduced max-tolerance graphs where each vertex $u \in V$ corresponds to a real interval I_u and a positive real number t_u (known as tolerance) such that uv is an edge of G if and only if $|I_u \cap I_v| \ge \max \{t_u, t_v\}$. For max-tolerance graphs we may assume $t_u \le |I_u|$ for each $u \in V$ otherwise u becomes isolated. An max-tolerance graph is a unit-max-tolerance graph if $|I_u| = |I_v|$ for all $u, v \in V$. Some combinatorial problems like finding maximal cliques were obtained in polynomial time whereas the recognition problem was proved to be NP-hard [12] for max-tolerance graphs in 2006. Also a geometrical connection of max-tolerance graphs to semi-squares was obtained in [12]. For further details of tolerance graphs one can see [9].

In 2015 Soto and Caro [18] introduced a new graph class, namely p-BOX graphs where each vertex corresponds to a box and a point within it in the *d*-dimensional Euclidean space. Any two vertices are adjacent if and only if the intersection of their corresponding boxes contains both the corresponding points. When the dimension is one the graph class is denoted by p-BOX(1). In 2017 this dimension one graphs are studied independently by Catanzaro et al [6], but with a different name, max-point tolerance graphs (MPTG) where each vertex $u \in V$ corresponds to a pair of an interval and a point (I_u, p_u) , where I_u is an interval on the real line and $p_u \in I_u$, such that uv is an edge of G if and only if $\{p_u, p_v\} \subseteq I_u \cap I_v$. The graphs MPTG have many practical applications in human genome studies and modelling of telecommunication networks [6]. A graph G = (V, E) is called central-max-point-tolerance graph (central MPTG) if p_u is the center point of I_u for each $u \in V$. This graph class actually matches with the graph defined as c-p-BOX(1) graph in [18]. It is known that c-p-BOX(1) graphs are max-tolerance graphs. We use the terms MPTG and central MPTG for p-BOX(1) and c-p-BOX(1) graphs throughout the paper.

In 1984 [15] Maehera defined an analogous concept of intersection graphs for digraphs. He introduced catch digraph of F as a digraph G = (V, E) where $uv \in E$ if and only if $u \neq v$ and $p_v \in S_u$ where $F = \{(S_u, p_u) | u \in V\}$ is a family of pointed sets in a Euclidean space. The digraph G is said to be represented by F. Later on interval catch digraphs (in brief, ICD) was introduced by Erich Prisner in 1989 [16] where S_u is represented by interval I_u . This graph class has many applications in real world situations like networking and telecommunications. He characterized interval catch digraphs in terms of the absence of diasteroidal triple in the digraph. A digraph G = (V, E) is unilaterally connected if for each pair of distinct vertices $u, v \in V$, there is a directed path from u to v or from v to u (or both). The underlying graph of a digraph G = (V, E) is an undirected graph U(G) = (V', E') where V' = V and $E' = \{uv \mid uv \in E \text{ or } vu \in E\}$. An undirected graph G = (V, E) is a caterpillar if it is a tree with path (s_1, s_2, \ldots, s_k) , called the spine of G such that every vertex of G has at most distance one from the spine.

In our paper we prove that central MPTG graphs are same as unit max-tolerance graphs. Incidentally this settles a question raised in the book of Golumbic [9] that whether interval graphs are unit max tolerance graphs or not. Moreover we show that a *unit central MPTG* is same as a *proper central MPTG* and also is same as a proper interval graph. Looking at the definition and the fact that central max point tolerance graphs are also max-tolerance graphs, one may think that central max point tolerance graphs are same as max-tolerance graphs where tolerance values are half of their corresponding interval length. In this paper, we introduce 50% *max-tolerance* graphs analogous to the similar concept for min-tolerance graphs. In case of min-tolerance graphs, unit and 50% are defining the same class of graphs [4]. In our paper we show that for max-tolerance graphs they are not same. In fact, the classes central MPTG and 50% max-tolerance graphs are not comparable, although they contain various classes of graphs (for example $C_n, n \geq 3$, proper

intervals graphs) in common. Finally we find a close relation of central MPTG with central interval catch digraph (central ICD). Central ICD is an interval catch digraph where the points p_u are the center points of the intervals I_u . This digraph was introduced by Maehera [15] in name of *interval digraph*. We characterize this digraph by defining a suitable mapping from the vertex set to the real line. Moreover we disprove a conjecture proposed by Maehera in [15] by discussing some structural property of the augmented adjacency matrix of a central ICD and creating counter examples. We define oriented interval catch digraph (oriented ICD) as an interval catch digraph G = (V, E) where each edge of its underlying graph has exactly one direction in G. We prove that an oriented ICD is acyclic and study various properties of it. We also show augmented adjacency matrix of an oriented ICD takes a special form when it is a tournament. Next we characterize an oriented ICD whose underlying graph is a tree. A proper interval catch digraph (proper ICD) is an interval catch digraph where no interval contains other properly. We obtain characterization of the augmented adjacency matrix of a proper ICD. We also find an important result which characterize those proper oriented interval catch digraphs (proper oriented ICD) whose underlying undirected graph is chordal, in terms of forbidden subdigraphs. In Conclusion Section we show the relations between the subclasses of max-tolerance graph and central MPTG and the digraphs discussed in this paper and list major open problems in this area. Henceforth undirected graphs will be called simply graphs.

2 Preliminaries

The following characterizations is known for interval catch digraphs.

Theorem 2.1. [16] Let G = (V, E) be a simple directed graph. Then G is an interval catch digraph if and only if there exists an ordering " < " of V such that

for
$$x < y < z \in V$$
, $xz \in E \Longrightarrow xy \in E$ and $zx \in E \Longrightarrow zy \in E$. (2.1)

A matrix whose entries are only zeros and ones is a *binary* matrix. A binary matrix is said to satisfy consecutive 1's property for rows if its columns can be permuted in such a way that 1's in each row occur consecutively [7]. For a simple undirected graph G = (V, E), a matrix $A^*(G)$ is known as the augmented adjacency matrix of G if we replace all principal diagonal elements of the adjacency matrix of G by 1 [7]. Now one can check from [15] that with respect to the ordering described in Theorem 2.1, $A^*(G)$ satisfy consecutive 1's property along rows. We call this ordering an *ICD ordering* of V. This ordering is not unique for an ICD. For a simple digraph G = (V, E), we denote the set of (closed) neighbors $\{v \in V \mid uv \in E\} \cup \{u\}$ of u by $d^+[u]$ for each $u \in V$. It is clear that elements of $d^+[u]$ are consecutive in an ICD ordering for each $u \in V$ if G is an ICD. Moreover, if $\{(I_u, p_u) \mid u \in V\}$ is a pointed interval representation of an ICD G, then the points can be made distinct by slight adjustment and the increasing ordering of these points is an ICD ordering of the corresponding vertices. From [9] we get to know that $\{x_1, x_2, x_3\} \subseteq V$ form a diasteroidal triple if for every permutation σ of $\{1, 2, 3\}$ there is an $x_{\sigma(1)}$ -avoiding ${}^1 x_{\sigma(2)} - x_{\sigma(3)}$ chain in G.

Theorem 2.2. [16] A digraph is an interval catch digraph if and only if it does not contain any diasteroidal triple.

A proper interval graph G is an interval graph in which there is an interval representation of G such that no interval contains another properly. A unit interval graph is an interval graph in which there is an interval representation of G such that all intervals have the same length. Let G = (V, E) is a graph and $v \in V$. Then the set $N[v] = \{u \in V \mid u \text{ is adjacent to } v\} \cup \{v\}$ is the closed neighborhood of v in G. The reduced graph \tilde{G} is obtained from G by merging vertices having same closed neighborhood. G(n, r) is a graph with n vertices

¹A $v_2 - v_3$ chain is v_1 avoiding if no initial endpoint of an arc of the chain precedes v_1 [9]

 x_1, x_2, \ldots, x_n such that x_i is adjacent to x_j if and only if $0 < |i - j| \le r$, where r < n is a positive integer. Among many characterizations of proper interval graph we list the following which will serve our purpose.

Theorem 2.3. [5, 13, 14] (pg 387, ex 11.17) Let G = (V, E) be an interval graph. Then the following are equivalent:

- 1. G is a proper interval graph.
- 2. G is a unit interval graph.
- 3. There exist a linear ordering < on V such that for every choice of vertices u, v, wu < v < w and $uw \in E$ implies $uv, vw \in E$.
- 4. \tilde{G} is an induced subgraph of G(n,r) for some positive integers n,r with n > r.

The following characterization of MPTG is known:

Theorem 2.4. [6] Let G = (V, E) be a simple undirected graph. Then G is an MPTG if and only if there is an ordering of vertices of G such that the following condition holds:

For any
$$x < u < v < y$$
, $xv, uy \in E \Longrightarrow uv \in E$. (2.2)

Definition 2.5. Let $A = (a_{ij})$ and $B = (b_{ij})$ be two $n \times n$ binary matrices. We define $A \wedge B = (c_{ij})$ where $c_{ij} = a_{ij} \wedge b_{ij}$ with the rules: $0 \wedge 0 = 1 \wedge 0 = 0 \wedge 1 = 0$ and $1 \wedge 1 = 1$.

The above characterization leads to the following observations.

Observation 2.6. Let G be a simple undirected graph. Then following are equivalent:

- 1. G is an MPTG.
- 2. There is an ordering of vertices of G such that for any $u < v, u, v \in V$,

$$uv \notin E \Longrightarrow uw \notin E \text{ for all } w > v \text{ or, } wv \notin E \text{ for all } w < u.$$
 (2.3)

- 3. There exists an ordering of vertices such that every 0 above the principal diagonal of the augmented adjacency matrix A(G) has either all entries right to it are 0 or, all entries above it are 0.
- 4. There exists a binary matrix M with consecutive 1's property for rows such that the augmented adjacency matrix $A(G) = M \wedge M^T$.
- 5. There exist an interval catch digraph D such that $G = D \cap D^T$ where D^T is the digraph obtained from D by reversing direction of every arc.

Proof. The condition 2 is equivalent to (2.2) in the other way. Condition 3 is a matrix version of condition 2. Condition 4, 5 follows from definition of MPTG and ICD respectively.

Let G = (V, E) be a simple undirected graph and $\emptyset \neq X \subseteq V$. Then G[X] denotes the subgraph of G induced by X. In Proposition 6.7 of [6] it is proved that if G is an MPTG with non-adjacent vertices u and v, then $G[N(u) \cap N(v)]$ is an interval graph. Also in Proposition 7.1 of [12] it is shown that $\overline{C_n}$, n > 9 is not an max-tolerance graph. We show that these graphs are not MPTG as well.

Observation 2.7. The complement of a cycle of length greater than 9 is not an MPTG.

Proof. Suppose on contrary $\overline{C_n}$, n > 9 is an MPTG. Now as $\overline{C_n}$, n > 9 contain the graph G'_1 with vertices $\{1, 4, 5, n - 3, n - 2, n\}$ in Figure 1 (left) as induced subgraph where common neighbors of n - 2 and n - 3 form a chordless 4-cycle and so is not an interval graph. Thus G'_1 is not an MPTG. Since any subgraph of an MPTG must be an MPTG, hence the result follows.

In the following example we show that MPTG and max-tolerance graph are not same.

Example 2.8. The graph G'_2 in Figure 1 (right) with vertex set $V = \{v_i | 1 \le i \le 6\}$ is not an MPTG as above. But it is a max-tolerance graph with the following interval and tolerance representation. $I_1 = [10, 46], t_1 = 21, I_2 = [20, 50], t_2 = 18, I_3 = [18, 49.5], t_3 = 28.5, I_4 = [15, 60], t_4 = 31, I_5 = [21, 52], t_5 = 10, I_6 = [12, 50], t_6 = 30.$

Theorem 2.9. [18] The graph class central MPTG properly contains the class of interval graphs.

3 Central max point tolerance graphs

We begin with a trivial but important observation which will be used throughout the rest of the paper.

Observation 3.1. Let $\{I_u \mid u \in V\}$ be a collection of intervals, where $I_u = [\ell_u, r_u]$, $h_u = |I_u| = r_u - \ell_u$ and $c_u = \frac{\ell_u + r_u}{2}$. Then $\{c_u, c_v\} \subseteq I_u \cap I_v \iff |c_v - c_u| \leqslant \frac{1}{2}\min\{h_u, h_v\} \iff \ell_v \leqslant c_u \leqslant c_v \leqslant r_u$ (for $c_u \leqslant c_v$).

 $\begin{array}{l} Proof. \text{ We have } c_u \in I_v = [l_v, r_v] = [c_v - \frac{h_v}{2}, c_v + \frac{h_v}{2}] \Longleftrightarrow c_v - \frac{h_v}{2} \leq c_u \leq c_v + \frac{h_v}{2} \Longleftrightarrow -\frac{h_v}{2} \leq c_u - c_v \leq \frac{h_v}{2} \\ \Leftrightarrow |c_u - c_v| \leq \frac{h_v}{2}. \text{ Thus } \{c_u, c_v\} \subseteq I_u \cap I_v \iff |c_u - c_v| \leq \frac{1}{2} \min\{h_u, h_v\}. \text{ Also it is clear that } c_u \in I_v = [l_v, r_v] \iff l_v \leq c_u \leq r_v. \end{array}$

In the above section we observed that the graph classes of max-point-tolerance graphs and max tolerance graphs are not same. Now it is interesting to see that the classes of central MPTG and unit max-tolerance graphs are same.

Theorem 3.2. Let G be a simple undirected graph. Then G is a central MPTG if and only if G is a unit max-tolerance graph.

Proof. Let G = (V, E) be a central MPTG with a central MPTG representation (I_u, c_u) where $I_u = [l_u, r_u], c_u$ be the center point of I_u for each vertex $u \in V$. Let $h_u = r_u - l_u$ for all $u \in V$. Choose $h_0 > \max\{h_u \mid u \in V\}$. Define $t_u = \frac{h_0 - h_u}{2}, y_u = c_u + \frac{h_0}{2}$ and $T_u = [c_u, y_u]$. Note that $t_u > 0$ and $|T_u| = y_u - c_u = \frac{h_0}{2}$ which is a constant for all $u \in V$.



Figure 1: The graphs G'_1 in Observation 2.7 and G'_2 in Example 2.8

Suppose $uv \in E$ and $c_v \leq c_u$. Then $c_u - c_v \leq \frac{1}{2}\min\{h_u, h_v\} \leq \frac{h_0}{2}$. So $c_u \leq c_v + \frac{h_0}{2} = y_v$. This implies $c_v \leq c_u \leq y_v$. So $T_u \cap T_v = [c_u, y_v] \neq \emptyset$ and $|T_u \cap T_v| = y_v - c_u = c_v + \frac{h_0}{2} - c_u = \frac{h_0}{2} - (c_u - c_v) \geq \frac{h_0 - h_u}{2}, \frac{h_0 - h_v}{2}$. So $y_v - c_u \geq t_u, t_v$, i.e.,

$$|T_u \cap T_v| \ge \max\{t_u, t_v\}. \tag{3.1}$$

On the other hand, (3.1) implies $T_u \cap T_v \neq \emptyset$ and $c_v \leq c_u \leq y_v \leq y_u$. So $|T_u \cap T_v| = y_v - c_u$. Now $y_v - c_u \geq \max\{t_u, t_v\}$ implies $\frac{h_0}{2} - (c_u - c_v) \geq \max\{\frac{h_0 - h_u}{2}, \frac{h_0 - h_v}{2}\}$. Thus $c_u - c_v \leq \frac{1}{2}\min\{h_u, h_v\}$, i.e., $uv \in E$. Therefore G is a unit max-tolerance graph with interval representation $\{T_u = [c_u, y_u] | u \in V\}$ and tolerances $\{t_u | u \in V\}$ as defined above.

Conversely, let G = (V, E) be a unit max-tolerance graph with interval representation $\{T_u = [l_u, r_u] \mid u \in V\}$ and tolerances $\{t_u \mid u \in V\}$. Let $h = |T_u|$ for all $u \in V$. Define $I_u = [l_u - (h - t_u), l_u + (h - t_u)]$. Then c_u , the center of $I_u = l_u$ and $h_u = |I_u| = 2(h - t_u) < 2h$. Suppose $uv \in E$. Then $|T_u \cap T_v| \ge \max\{t_u, t_v\}$. Now for $l_u \le l_v$, $|T_u \cap T_v| = r_u - l_v$. Then $r_u - l_v \ge \max\{t_u, t_v\}$, i.e., $h + l_u - l_v \ge \max\{t_u, t_v\}$. This implies $l_v - l_u \le \min\{h - t_u, h - t_v\}$, i.e., $c_v - c_u \le \frac{1}{2}\min\{h_u, h_v\}$. Finally, the condition that $0 < c_v - c_u \le \frac{1}{2}\min\{h_u, h_v\} \Rightarrow l_v - l_u < h$ and $l_u \le l_v < l_u + h = r_u$. So $T_u \cap T_v \neq \emptyset$ and $|T_u \cap T_v| = r_u - l_v$. Then $l_v - l_u = c_v - c_u \le \frac{1}{2}\min\{h_u, h_v\}$ implies $|T_u \cap T_v| \ge \max\{t_u, t_v\}$, i.e., $uv \in E$. Thus $\{(I_u, c_u) \mid u \in V\}$ is a central MPTG representation of G.

Remark 3.3. In ([9], page 215) Golumbic wrote that "Every interval graph is a proper max-tolerance graph. It is not yet known if this can be strengthened to unit max-tolerance." The above theorem shows central MPTG and unit max-tolerance graphs denote the same graph class. Hence from Theorem 2.9 one can easily conclude that every interval graph is a unit max-tolerance graph. Thus we settle the above query posed in the book of Golumbic.

In the sequel we show that the class of max-tolerance graphs properly contains the class of central MPTG. We begin with the following definition which unfolds more insight in the structure of a central MPTG.

Definition 3.4. (*C*-order) Let G = (V, E) be a central MPTG with (distinct) center points $\{c_u \mid u \in V\}$ of the intervals $\{I_u \mid u \in V\}$ in its central MPTG representation $\{(I_u, c_u) \mid u \in V\}$. The *C*-order of the set *V* is the total order induced by the center points. For convenience abusing notation, henceforth we write u < v if and only if $c_u < c_v$.

In the following we present a necessary condition for central MPTG.

Theorem 3.5. Let G = (V, E) be a central MPTG. Then there is an ordering \prec^* of vertices of G such that the following condition holds:

For any
$$x \prec^* u \prec^* v \prec^* y$$
, $xv, uy \in E \Longrightarrow uv \in E$ and $(xu \in E \text{ or } vy \in E \text{ or } xu, vy \in E)$. (3.2)

Proof. Let G = (V, E) be a central MPTG with a central MPTG representation (I_u, c_u) for each $u \in V$. We arrange vertices according to the increasing order of center points (i.e., in *C*-order) of representing intervals. Suppose in this ordering we have x < u < v < y and $xv, uy \in E$. Then $c_v, c_x \in I_v \cap I_x$ and $c_u, c_y \in I_u \cap I_y$. Also we have $c_x < c_u < c_v < c_y$. Now $c_x, c_v \in I_v \Rightarrow c_u \in I_v$ and $c_u, c_y \in I_u \Rightarrow c_v \in I_u$. Therefore $uv \in E$. Again $c_x, c_v \in I_x \Rightarrow c_u \in I_x$ and $c_u, c_y \in I_y \Rightarrow c_v \in I_y$. Thus if $xu, vy \notin E$, then $c_x \notin I_u$ and $c_y \notin I_v$. But then $c_u - c_x > c_y - c_u$ as $c_x \notin I_u$ but $c_y \in I_u$, and $c_y - c_v > c_v - c_x$ as $c_y \notin I_v$ but $c_x \in I_v$. Combining these inequalities we have $c_v < \frac{c_x + c_y}{2} < c_u$ which is a contradiction. Therefore $xu \in E$ or, $vy \in E$ or, $xu, vy \in E$.




Figure 2: Relative positions of intervals described in the proof of Theorem 3.9

In [18], it is shown that every cycle C_n of length $n \ge 3$ is a central MPTG.

Definition 3.6. A cycle C_n is said to be *circularly consecutive* C-ordered if starting from a fixed vertex (say u) one can order all its vertices in a circularly consecutive way in clockwise (or anticlockwise) direction until u is reached in a C-order.

In particular, for n = 4, one can obtain a central MPTG representation of C_4 from [18] where vertices are circularly consecutive C-ordered. We state a stronger version in the following corollary.

Corollary 3.7. Any induced C_4 in central MPTG must be circularly consecutive C-ordered.

Proof. All other possible C-orderings of vertices will violate (3.2). Hence the proof follows.

Similarly one can obtain the following:

Corollary 3.8. Any induced P_4 in central MPTG must have vertex consecutive ending edges i.e; vertices corresponding to ending edges of P_4 occur consecutively in a C-order (up to permutations between them) at least in one end.

Theorem 3.9. $\overline{C_6}$ is a max-tolerance graph but it is not a central MPTG.

Proof. Let $\{v_i | 1 \le i \le 6\}$ be the vertices occurred in circularly consecutive way in clockwise (or anticlockwise) order in C_6 . We assign the following intervals and tolerances for all the vertices so that they satisfy maxtolerance representation in $\overline{C_6}$. $I_{v_1} = [0, 20], t_{v_1} = 10, I_{v_2} = [12, 24], t_{v_2} = 6, I_{v_3} = [0, 22], t_{v_3} = 11, I_{v_4} = [9.5, 19.5], t_{v_4} = 5, I_{v_5} = [7.5, 30.5], t_{v_5} = 11.5, I_{v_6} = [10.5, 21.5], t_{v_6} = 5.5.$

Now suppose $\overline{C_6} = (V, E)$ is a central MPTG with central MPTG representation (I_v, c_v) where $I_v = [a_v, b_v]$, c_v be the center point of I_v for each $v \in V$. Let $\{v_i | 1 \leq i \leq 6\}$ be the vertices occurred in circularly consecutive way in clockwise (or anticlockwise) order in C_6 . It is easy to check that the subgraph induced by deleting the vertices $\{v_2, v_5\}$ from $\overline{C_6}$ is a C_4 . Now from Corollary 3.7 we can conclude that the vertices in $C_4 = \{v_1, v_4, v_6, v_3\}$ are circularly consecutive C-ordered. Without loss of generality we can take $c_1 < c_4 < c_6 < c_3$. As $v_1v_5, v_3v_5 \in E$, $a_5 \leq c_1$ and $c_3 \leq b_5$. Thus we get $[c_1, c_3] \subseteq [a_5, b_5]$. Hence $c_4 \in [c_1, c_3]$ imply

 $c_4 \in I_5$. Below we will show $c_5 \in I_4$ (see Figure 2) which lead us to contradiction as v_4, v_5 are nonadjacent in $\overline{C_6}$.

As $v_3v_6 \in E$, $c_3 \leq b_6$. Again $v_1v_3 \in E$ imply $[c_1, c_3] \subseteq I_3$ as $c_1 < c_3$. Moreover $c_4 \in [c_1, c_3] \subseteq I_3$. Hence $c_3 \notin I_4$ as $v_3v_4 \notin E$, which imply $b_4 < c_3$ as $c_4 < c_3$. Combining we get

$$b_4 < c_3 \le b_6 \tag{3.3}$$

Since $v_1v_5, v_3v_5 \in E$, we get $[c_1, c_3] \subseteq I_5$ as $c_1 < c_3$. Now as $c_6 \in [c_1, c_3]$, $c_6 \in I_5$. Hence $c_5 \notin I_6$ as $v_5v_6 \notin E$. Hence $c_5 < a_6$ or $c_5 > b_6$. Again $c_5 \leq b_4$ (as $v_4v_5 \in E$) and hence c_5 can not be greater than b_6 as $b_4 < b_6$ from (3.3). Thus we get

$$c_5 < a_6 \tag{3.4}$$

Note that $v_1v_3 \in E$ imply $[c_1, c_3] \subseteq I_1$ as $c_1 < c_3$. Again $c_6 \in [c_1, c_3] \subseteq I_1$. But as $v_1v_6 \notin E$, $c_1 < a_6$ as $c_1 < c_6$. Again $v_1v_4 \in E$ imply $a_4 \leq c_1$. Hence combining we get $a_4 \leq c_1 < a_6$. Again $v_4v_6 \in E$ imply $a_6 \leq c_4$ and $c_6 \leq b_4$. Hence $a_6 \leq c_4 < c_6 \leq b_4$. Thus combining these inequalities and using (3.3) one can conclude $a_4 \leq c_1 < a_6 \leq c_4 < c_6 \leq b_4 < c_3 \leq b_6$. Thus we get

$$a_4 \le c_1 < a_6 < b_4 < c_3 \le b_6 \tag{3.5}$$

As $v_2v_4, v_2v_6 \in E$ imply $c_2 \in I_4 \cap I_6 = [a_6, b_4] \subseteq [c_1, c_3] \subseteq I_1, I_3$ (from (3.5)) which imply $c_2 \in I_1, I_3$. Now $v_1v_2 \notin E$ imply $a_2 > c_1$ as $c_1 < c_2$ from above. Again $v_2v_3 \notin E$ imply $b_2 < c_3$ as $c_2 < c_3$ from above. Thus we get $c_1 < a_2 \leq c_2 \leq b_2 < c_3$, i.e;

$$[a_2, b_2] \subseteq [c_1, c_3]. \tag{3.6}$$

We now show $c_5 \in I_4$. As $v_2v_5 \in E$, $c_5 \in [a_2, b_2] \subseteq [c_1, c_3] \subseteq [a_4, b_6]$ from (3.5) and (3.6). Now using (3.4) one can conclude that c_5 must belong to I_4 .

From above we can conclude now that no CMPTG representation of $\overline{C_6}$ can be found with respect to the above *C*-ordering. For other possible *C*-orderings following similar type argument one can reach to contradiction.

Now we present a sufficient condition for an MPTG to be a central MPTG.

Theorem 3.10. Let G = (V, E) be an MPTG with n vertices. Let the ordering $\{v_1, v_2, \ldots, v_n\}$ of vertices of G that satisfies (2.2) and each v_i corresponds to a natural number x_i such that $x_1 < x_2 < \cdots < x_n$ and the following conditions hold for all $i = 1, 2, \ldots, n$:

$$x_{i_2+1} - x_i > x_i - x_{i_1} \quad when \quad i_2 < n \tag{3.7}$$

$$x_i - x_{i_1-1} > x_{i_2} - x_i \quad when \quad i_1 > 1$$
(3.8)

where i_1 and i_2 be the least and the highest indices such that $i_1 = i$ or, $v_i v_{i_1} \in E$ and $i_2 = i$ or, $v_i v_{i_2} \in E$. Then G is a central MPTG.

Proof. Suppose the conditions hold. Define $r_i = \max\{x_i - x_{i_1}, x_{i_2} - x_i\}$ and $I_i = [x_i - r_i, x_i + r_i]$ for i = 1, 2, ..., n. We show that G = (V, E) is a central MPTG with an interval representation $\{I_{v_i} | i = 1, 2, ..., n\}$ where $V = \{v_1, v_2, ..., v_n\}$ and this ordering of vertices satisfies (2.2). Suppose $v_i v_j \in E$. Then by definition of i_1 and i_2 , we have $x_{i_1} \leq x_j \leq x_{i_2}$ and $x_{j_1} \leq x_i \leq x_{j_2}$. Then $x_i - x_{i_1} \geq x_i - x_j$ and $x_{i_2} - x_i \geq x_j - x_i$ which imply $|x_i - x_j| \leq r_i$ and so $x_j \in I_{v_i}$. Similarly $x_i \in I_{v_j}$. Hence $\{x_i, x_j\} \subseteq I_{v_i} \cap I_{v_j}$. Now let $v_i v_j \notin E$. Without loss of generality we assume i < j. Suppose $j_1 < i$ and $j < i_2$. Then we have $j_1 < i < j < i_2$ and

 $v_{j_1}v_j, v_iv_{i_2} \in E$. Then by (2.2), $v_iv_j \in E$, which is a contradiction. Thus either $i < j_1$ or $j > i_2$. Then $i \leq j_1 - 1$ or $j \geq i_2 + 1$. For the first inequality by (3.7), we have $x_j - x_i \geq x_j - x_{j_1-1} > x_{j_2} - x_j$. Also $x_j - x_i > x_j - x_{j_1}$, as $x_i < x_{j_1}$. Thus $x_j - x_i > r_j$ which implies $x_i \notin I_{v_j}$. Similarly $j \geq i_2 + 1$ implies $x_j \notin I_{v_i}$. Therefore G is a central MPTG.

Definition 3.11. A central MPTG G = (V, E) is called *proper* if it has an interval representation with the required condition such that no interval contains another properly. We call it *proper-central-max-point* tolerance graph (in brief, proper central MPTG). Similarly, a central MPTG G = (V, E) is called *unit* if it has an interval representation with the required condition such that every interval has unit (or, same) length. We call it *unit-central-max-point tolerance graph* (in brief, unit central MPTG).

Theorem 3.12. Let G be a simple undirected graph. Then the following are equivalent.

- 1. G is a proper central MPTG.
- 2. G is a unit central MPTG.
- 3. G is a proper interval graph.

Proof. $(\mathbf{1} \iff \mathbf{2})$: Let G be a proper central MPTG with respect to the representation (I_i, c_i) where $I_i = [a_i, b_i], c_i$ be the center point of I_i for each vertex $i \in V$. First we arrange the intervals according to increasing order of left end points. As no interval properly contains another, the right end points have the same order as left end points and so as the center points as well. We process the representation from left to right, adjusting all intervals to length l where l is the length of first interval (i.e; $|I_1| = l$). At each step until all intervals been adjusted I_x be the leftmost unadjusted interval.

Let I_j be an adjusted interval occurs before I_x , then one of following things happen.

- (1) $c_j \notin I_x$.
- (2) $c_j \in I_x, c_x \in I_j.$
- (3) $c_j \in I_x$ but $c_x \notin I_j$.

Let I_{j_1} and I_{j_2} be any two adjusted intervals referred in (2) and (3) respectively. Then I_{j_2} must occur before I_{j_1} otherwise the right end point of I_{j_2} would occur before c_x and so I_{j_2} would be properly contained in I_{j_1} as I_{j_1} contains c_x . But this is a contradiction.

Now if I_x does not contain center point of any adjusted intervals then take $\alpha = a_x$. If I_x contains center points of some adjusted intervals and I_i be the leftmost among them, then $c_l \in I_x$ for all $i \leq l \leq x$ as all of them have same length. Now if $c_x \in I_i$, then take $\alpha = c_i$. It follows from the last paragraph that $c_x \in I_l$ for all $i \leq l \leq x$ in this case. Now if $c_x \notin I_i$, $c_x \notin I_l$ for any l < i. Let I_j be the leftmost interval for which $c_x \in I_j$. Then i < j < x. Take $\alpha = c_j$ in this case. Now if no such I_j exists between I_i and I_x i.e; if $c_x \notin I_j$ for all $i \leq j < x$ then $c_x \notin I_l$ for any l < x. Take $\alpha = b_l$ in this case where b_l is the rightmost endpoint for which $c_l \in I_x, c_x \notin I_l$. Clearly $i \leq l < x$. We adjust the portion $[a_x, \infty)$ by shrinking or expanding $[a_x, b_x]$ to $[\alpha, \alpha + l]$ and scaling and shifting $[b_x, \infty)$ to $[\alpha + l, \infty)$. Iterating this operation produces the unit central MPTG representation.

Now it is sufficient to show adjusting I_x in above way does not affect the adjacency of vertex x with previous intervals. When $\alpha = a_x$, then $\alpha = a_x > c_l$ for all l < x. Hence $c_l \notin I_x$ after adjustment. When $\alpha = c_i$, then $c_i = \alpha \in I_x$ and $c_x = \alpha + \frac{l}{2} = c_i + \frac{l}{2} = b_i \in I_i$. Moreover for all i < l < x, $c_x \in I_l$ and $\alpha = c_i < c_l < b_i = \alpha + \frac{l}{2}$ (i.e; $c_l \in I_x$) and $c_x \notin I_l$ for all l < i after adjustment. When $\alpha = c_j$ then the arguments are similar as above. Again when $\alpha = b_l$, then $c_x = \alpha + \frac{l}{2} = b_l + \frac{l}{2} > b_l$ which imply $c_x \notin I_l$. Hence $c_x \notin I_k$ for all k < l.




Figure 3: The graph G in Example 3.15

Conversely, if G is a unit central MPTG then all intervals associated to the vertices of G must be of the same length. Thus none of them contains other properly and so G is a proper central MPTG with the same interval representation.

 $(\mathbf{3} \Rightarrow \mathbf{1})$: Let G = (V, E) be a proper interval graph. So the reduced graph $\hat{G} = (\hat{V}, \hat{E})$ is an induced subgraph of $G(n, r) = (V_n, E')$ for some $n, r \in \mathbb{N}$ with n > r, where $V_n = \{v_1, v_2, \ldots, v_n\}$ and $v_i \leftrightarrow v_j$ if and only if $|i - j| \leq r$ by condition 4 of Theorem 2.3. Let $\hat{V} = \{v_{i_1}, v_{i_2}, \ldots, v_{i_m}\} \subseteq V_n$. Now for each $u \in V$, define $p_u = i_j$ if u is a copy of v_{i_j} and $I_u = [p_u - r, p_u + r]$. Firstly all intervals I_u are of same length 2r and so none of them properly contains other.

Next let $u, v \in V$. Suppose $p_u = i_j$ and $p_v = i_k$. Then u is a copy of v_{i_j} and v is a copy of v_{i_k} . If $uv \in E$, then $v_{i_j}v_{i_k} \in \hat{E} \subseteq E'$. Therefore $|i_j - i_k| \leq r \Rightarrow |p_u - p_v| \leq r \Rightarrow p_v \in I_u$ and $p_u \in I_v \Rightarrow p_u, p_v \in I_u \cap I_v$. Finally, let $uv \notin E$. Then $v_{i_j}v_{i_k} \notin \hat{E}$. Since \hat{G} is an induced subgraph of G(n,r), we have $v_{i_j}v_{i_k} \notin E'$. Then $|i_j - i_k| > r \Rightarrow |p_u - p_v| > r \Rightarrow p_v \notin I_u$ and $p_u \notin I_v$. Thus G is a proper central MPTG.

 $(\mathbf{1} \Rightarrow \mathbf{3})$: let G = (V, E) be a proper central MPTG with a proper central MPTG representation (I_u, c_u) where $I_u = [l_u, r_u]$, c_u be the center point of I_u for each $u \in V$. We arrange vertices according to the increasing order of center points, $V = \{v_1, v_2, \ldots, v_n\}$. To prove that G is a proper interval graph we show that vertices of G satisfy condition 3 of Theorem 2.3 with respect to the above ordering.

Denote $I_{u_i} = [l_{u_i}, r_{u_i}]$ by $[l_i, r_i]$ and $c_i = \frac{l_i + r_i}{2}$ for i = 1, 2, ..., n. Let i < j < k and $u_i u_k \in E$. Then $c_i < c_j < c_k$. Now since G is a central MPTG, $c_k - c_i \le \min\{c_i - l_i, c_k - l_k\}$. Now $c_j - c_i < c_k - c_i \le c_i - l_i$. Now if $l_j > c_i$, then $l_k \le c_i < l_j < c_j < c_k$ as $c_k - c_i \le c_k - l_k$. So $[l_j, c_j] \subsetneq [l_k, c_k]$. But this implies $[l_j, r_j] \subsetneq [l_k, r_k]$ which contradicts the fact that G is a proper central MPTG. Thus $l_j \le c_i$. So we have $c_j - c_i \le c_j - l_j$. Hence $c_j - c_i \le \min\{c_i - l_i, c_j - l_j\}$. Therefore $u_i u_j \in E$, as required. Similarly, it can be shown that $u_j u_k \in E$. Thus G is a proper interval graph.

Next following observation 2.6 we have the following in a similar way:

Proposition 3.13. Let G be a simple undirected graph. Then G is a central MPTG if and only if there exists a central interval catch digraph D such that $G = D \cap D^T$ where D^T is the digraph obtained from D by reversing direction of every arc.

It is proved in [6] that if G is an MPTG with non-adjacent vertices u and v, then $G[N(u) \cap N(v)]$ is an interval graph. We found the following analogous result for central MPTG.

Proposition 3.14. If G is a central MPTG with non-adjacent vertices u and v, then $G[N(u) \cap N(v)]$ is a proper interval graph.

Proof. Let G = (V, E) be a central MPTG with an interval representation $\{I_i = [a_i, b_i] \mid i \in V\}$ where vertices are arranged according to C-order. Let c_i be the center point of I_i . Suppose u < v. Then vertices of $G[N(u) \cap N(v)]$ which occur between u, v form a clique from (2.2). Again if there exist a vertex of $G[N(u) \cap N(v)]$ occurs before u then no vertex can occur after v which belongs to $G[N(u) \cap N(v)]$ and conversely follows from (2.2) and the fact $uv \notin E$. Moreover vertices of $G[N(u) \cap N(v)]$ that occur before u form a clique. Let x < y < u < v such that $x, y \in G[N(u) \cap N(v)]$ then $c_x < c_y < c_u < c_v \le b_x$ as $vx \in E$. This implies $c_y \in [c_x, b_x] \subset I_x$. Now if $a_y > c_x$, then $a_u \le c_x < a_y < c_y < c_u$ (as $ux \in E$) which implies $c_y - a_y < c_u - a_u$. Hence $|I_y| < |I_u|$. Again as $a_y, c_y \in [a_u, c_u]$ from above we can conclude that $b_y \le b_u$. But $yv \in E$ implies $a_v < c_y < c_u < c_v \le b_y$ which imply $c_u \in [a_v, c_v] \subset I_v$. Also as u is not adjacent to $v, c_v > b_u$. Hence from above we get $b_u < c_v \le b_y$ which is a contradiction. Therefore $a_y \le c_x$. So $c_x \in [a_y, c_y] \subset I_y$. Hence $xy \in E$. Similarly one can show vertices of $G[N(u) \cap N(v)]$ which occur after v form a clique.

Now let $\{u_i | u_i < u\}$ be the vertices of $G[N(u) \cap N(v)]$ arranged in C-order and $\{x_i | u < x_i < v\}$ be the vertices of $G[N(u) \cap N(v)]$ arranged according to increasing order of left end points. From above observations it is clear that $u_i[x_i]$'s form clique for $u_i < u$ $[u < x_i < v]$. Let $u_k[x_m]$ be the last vertices occurred before u [between u and v] respectively. Now we show that $G[N(u) \cap N(v)]$ becomes a proper interval graph with respect to the ordering $\{u_1, \ldots, u_k, x_1, \ldots, x_m\}$. In this ordering by $p \prec q$ we mean p occurs before q. Infact we will show that the vertices satisfy condition 3 of Theorem 2.3 with respect to the ordering \prec . Let $u_l \prec$ $x_i \prec x_j$ where $1 \leq l \leq k, 1 \leq i, j \leq m$ such that $u_l x_j \in E$. Then $c_{u_l} < c_u < c_{x_i}, c_{x_j} < c_v < b_{u_l}$ as $vu_l \in E$. This implies $c_{x_i} \in [c_{u_l}, b_{u_l}] \subset I_{u_l}$. Now as $u_l x_j \in E$, $a_{x_j} \leq c_{u_l} < c_u < c_{x_i}$ implies $a_{x_i} < a_{x_j} \leq c_{u_l} < c_{x_i}$ (as $x_i \prec x_j \iff a_{x_i} < a_{x_j}$) which imply $c_{u_l} \in [a_{x_i}, c_{x_i}] \subset I_{x_i}$. Hence $u_l x_i \in E$. Let $u_i \prec u_j \prec x_l$ where $1 \leq i, j \leq k, 1 \leq l \leq m$ such that $u_i x_l \in E$. Then $u_i \prec u_j \prec x_l \prec v$ clearly. Now from (2.2) one can conclude $u_j x_l \in E$. Similarly one can show if there exists vertices of $G[N(u) \cap N(v)]$ that occurs after v, then with respect to the ordering $\{x_1, \ldots, x_m, v_1, \ldots, v_k\}$ (use $p \prec' q$ if and only if p occurs before q in this ordering) $G[N(u) \cap N(v)]$ forms a proper interval graph where $\{x_i | u < x_i < v\}$ are vertices of $G[N(u) \cap N(v)]$ arranged according to increasing order of right end points, and $\{v_i | v < v_i\}$ are vertices of $G[N(u) \cap N(v)]$ arranged in C-order.

The above proposition leads to a construction of the following forbidden graph for the class of central MPTG.

Example 3.15. By Proposition 3.14 we see that the graph G (see Figure 3) formed by taking $K_{1,3}$ together with two non-adjacent vertices (say, u, v) which are adjacent to each vertex of $K_{1,3}$ is not a central MPTG.

4 50% max-tolerance graphs

In this section we introduce a new type of max-tolerance graph, namely 50% max-tolerance graph similar to the concept of 50% min-tolerance graph [4].

Definition 4.1. A max-tolerance graph G = (V, E) is a 50% max-tolerance graph if $t_u = \frac{|I_u|}{2}$ for all $u \in V$ where t_u denotes the tolerance associated with the vertex $u \in V$.

We know that classes of unit (min) tolerance graphs and 50% (min) tolerance graphs are same [4]. So the very natural question which comes to our mind that whether unit and 50% max-tolerance graphs denote the same graph class. We settle down this question in the following theorem.

Theorem 4.2. Central MPTG (i.e., unit max-tolerance graphs) and the class of 50% max-tolerance graphs are not comparable.

Proof. The graph $K_{1,n}$ is an interval graph for any natural number n. So by Theorem 2.9 it is a central MPTG and hence a unit max-tolerance graph as well. In Lemma 4.3 we will show $K_{1,n}$, $n \ge 8$ is not a 50% max-tolerance graph. Again from Theorem 3.9 one can verify that $\overline{C_6}$ is a 50% max-tolerance graph but it is not a central MPTG. Thus $K_{1,n}$, $n \ge 8$ and $\overline{C_6}$ separate unit max-tolerance graph from 50%-max-tolerance graph. Hence these graph classes become incomparable.

We do the proof verification of the above theorem with the help of following Lemma.

Lemma 4.3. $K_{1,n}$ where $n \ge 8$ is not a 50% max-tolerance graph.

Proof. On the contrary lets assume that $K_{1,n}$, $n \ge 8$ has a 50% max-tolerance representation. By suitable scaling and shifting origin without loss of generality, we may assume that $I_u = [0, 1]$, interval corresponding to central vertex u and $I_{v_i} = [a_i, b_i]$, intervals corresponding to the pendant vertices v_i where $i \in \{1, \ldots, n\}$. We consider the intervals corresponding to any two pendants must be distinct as they are non-adjacent. Let [a, b] be an interval corresponding to a pendant vertex I_v . Then I_v must belong to one of the four sets, $S_1 = \{I_v | a \le 0 < 1 \le b\}$, $S_2 = \{I_v | a < 0, 0 < b < 1\}$, $S_3 = \{I_v | 0 < a < 1, b > 1\}$, $S_4 = \{I_v | 0 \le a < b \le 1\}$. We note that any pendant vertex cannot have [0, 1] as its interval representation. Hence the above sets are mutually exclusive. As every pendant vertex is adjacent to the central vertex u hence it follows that the four sets are also exhaustive. Now we show that

$$|S_1| \le 1$$
 and $|S_i| \le 2$ for $i \in \{2, 3, 4\}.$ (4.1)

• $|S_1| \le 1$.

Let $[a_1, b_1], [a_2, b_2]$ be the two intervals associated to two such pendant vertices (say v_1, v_2). Then the $[a_i, b_i] \supseteq [0, 1]$ for $i = \{1, 2\}$. Now $[a_i, b_i] \cap [0, 1] = [0, 1]$ clearly. As every pendant vertex is adjacent to u and I_u is of unit length, so each pendant vertex has tolerance at most 1 and therefore has length at most 2. But $[a_1, b_1] \cap [a_2, b_2] \supseteq [0, 1] \Rightarrow |[a_1, b_1] \cap [a_2, b_2] | \ge 1$. Hence $v_1 \leftrightarrow v_2$ which contradicts the fact that they are pendant vertices.

• $|S_2| \le 2.$

Let $[a_1, b_1], [a_2, b_2]$ and $[a_3, b_3]$ be three such intervals corresponding to the vertices (say v_1, v_2, v_3). First we will show that for any two of the three intervals, one must contain the other. On contrary, without loss of generality assume that $[a_1, b_1], [a_2, b_2]$ are two such intervals which are not containing each other and let $a_1 < a_2, b_1 < b_2$. It is sufficient to show for one case. Now let $[a_1, b_1] \cap [a_2, b_2] = [a_2, b_1]$. Now as $v_1 \leftrightarrow u, |[a_1, b_1] \cap [0, 1]| \ge \max\{\frac{1}{2}, \frac{b_1 - a_1}{2}\} \Rightarrow b_1 \ge \frac{1}{2}, \frac{b_1 - a_1}{2}$. This implies

$$b_1 \ge -a_1 \tag{4.2}$$

Similarly $b_2 \ge -a_2$.

Now $-a_1 \leq b_1 \Rightarrow -a_1 + b_1 \leq 2b_1 \Rightarrow -a_1 + b_1 < 2b_1 - 2a_2 \Rightarrow \frac{-a_1 + b_1}{2} < b_1 - a_2$. But $v_1 \nleftrightarrow v_2$. Hence $b_1 - a_2 < \frac{b_2 - a_2}{2} \Rightarrow 2b_1 - 2a_2 < b_2 - a_2 \Rightarrow 2b_1 - a_2 < b_2$. Hence $b_2 > 2b_1 - a_2 > 2b_1 \geq 1$ (Since $b_1 \geq \frac{1}{2}$ and $a_2 < 0$) which contradicts the fact $b_2 < 1$. Hence the three intervals form a well ordered set with inclusion as the ordering.

From the statement above without loss of generality we can assume that $[a_1, b_1] \supseteq [a_2, b_2] \supseteq [a_3, b_3]$. Now $[a_1, b_1] \cap [a_2, b_2] = [a_2, b_2]$. Therefore $2(b_2 - a_2) < b_1 - a_1$. Similarly $b_2 - a_2 > 2(b_3 - a_3)$. Hence $b_1 - a_1 > 4(b_3 - a_3) > 4b_3 \ge 2$ (since $b_3 \ge \frac{1}{2}$ as $v_3 \leftrightarrow u$). Now $b_1 - a_1 \le 2b_1 < 2$ from (4.2) and the fact $b_1 < 1$. Hence we are through.

• $|S_3| \le 2$.

This proof is same as the last case.

• $|S_4| \le 2.$

On the contrary we assume that there exists three such vertices with representations $[a_i, b_i] \subseteq [0, 1]$ for $i \in \{1, 2, 3\}$.

1. First we will show that for any two intervals, one must **not** contain the other.

If not, we assume $[a_i, b_i] \subseteq [a_j, b_j]$ for some $i, j \in \{1, 2, 3\}$. Then $v_i \leftrightarrow u \Rightarrow b_i - a_i \ge \frac{1}{2}$. Hence $|[a_i, b_i] \cap [a_j, b_j]| = |[a_i, b_i]| \ge \frac{1}{2}$. Again $v_i \leftrightarrow v_j \Rightarrow b_i - a_i < \frac{b_j - a_j}{2} \le \frac{1}{2}$ which is a contradiction. 2. Next we will show that no two intervals are disjoint.

If possible let $[a_1, b_1]$ and $[a_2, b_2]$ be disjoint. Without loss of generality we can assume $a_1 < a_2$. Hence since they are disjoint $b_1 < a_2$. But $b_1 \ge \frac{1}{2}$ as $v_1 \leftrightarrow u$. Hence $a_2 > \frac{1}{2}$. This implies $b_2 - a_2 < \frac{1}{2}$ which contradicts $v_2 \leftrightarrow u$.

Now without loss of generality we can assume $a_1 < a_2 < a_3$. Hence from 1 and 2 we conclude $a_1 < a_2 < a_3 \le b_1 < b_2 < b_3$. Under this situation we will show that there exist no choice of three such intervals. For this we first show $b_2 > \frac{3}{4}$. If $a_2 \le \frac{a_1 + b_1}{2}$ then $b_1 - a_2 \ge b_1 - \frac{a_1 + b_1}{2} = \frac{b_1 - a_1}{2}$. Then $b_1 - a_2 < \frac{b_2 - a_2}{2}$ since $v_1 \nleftrightarrow v_2$. Hence $b_2 > 2b_1 - a_2 \ge 2b_1 - \frac{a_1 + b_1}{2} = b_1 + \frac{b_1 - a_1}{2} \ge \frac{1}{2} + \frac{1/2}{2} = \frac{3}{4}$ (Since $v_1 \leftrightarrow u$, $b_1 - a_1 \ge \frac{1}{2}$ and hence $b_1 \ge \frac{1}{2}$). For other case, i.e., $a_2 > \frac{a_1 + b_1}{2} = \frac{b_1 - a_1}{2} + a_1 \ge \frac{b_1 - a_1}{2} \ge \frac{1/2}{2} = \frac{1}{4}$. Now $b_2 - a_2 \ge \frac{1}{2} \Rightarrow b_2 \ge a_2 + \frac{1}{2} > \frac{1}{4} + \frac{1}{2} = \frac{3}{4} \Rightarrow b_2 > \frac{3}{4}$.

For the remaining of the proof we split into two cases.

- 1. If $a_3 \leq \frac{a_2 + b_2}{2}$ then $b_2 a_3 \geq b_2 \frac{a_2 + b_2}{2} = \frac{b_2 a_2}{2}$. Then $b_2 a_3 < \frac{b_3 a_3}{2}$ as $v_2 \leftrightarrow v_3$. This implies $b_3 > 2b_2 - a_3 \geq 2b_2 - \frac{a_2 + b_2}{2} = b_2 + \frac{b_2 - a_2}{2} > \frac{3}{4} + \frac{1/2}{2} = 1$. Hence we arrive at a contradiction.
- 2. If $a_3 > \frac{a_2 + b_2}{2}$ then since $b_3 a_3 \ge \frac{1}{2}$ (as $v_3 \leftrightarrow u$), we have

$$b_3 > \frac{a_2 + b_2}{2} + \frac{1}{2} \tag{4.3}$$

Now if $a_2 \leq \frac{a_1 + b_1}{2}$ then $b_2 > 2b_1 - a_2$ as before. Hence $\frac{b_2 + a_2}{2} > b_1$. Hence using (4.3) we get $b_3 > b_1 + \frac{1}{2} \geq \frac{1}{2} + \frac{1}{2} = 1$ contradicting $b_3 \leq 1$. Now if $a_2 > \frac{a_1 + b_1}{2}$ then $b_2 > \frac{a_1 + b_1}{2} + \frac{1}{2}$ since $b_2 - a_2 \geq \frac{1}{2}$. Hence $a_2 + b_2 > a_1 + b_1 + \frac{1}{2}$. Again using (4.3) we get $b_3 > \frac{a_1 + b_1}{2} + \frac{1}{4} + \frac{1}{2} \geq \frac{1/2}{2} + \frac{1}{4} + \frac{1}{2} = 1$ (as $b_1 \geq \frac{1}{2}, a_1 \geq 0$). Hence we again arrive at a contradiction.

Hence we have established our claim.

Hence the maximum number of possible pendant vertices is 1 + 2 + 2 + 2 = 7. So we are done if we take at least 8 pendant vertices, i.e., $n \ge 8$.

In the following Propositions we will show that the class of proper interval graphs and any cycle of length n both are subclasses of 50% max-tolerance graphs.

Proposition 4.4. A proper interval graph is a 50% max-tolerance graph.

Proof. Let G = (V, E) be a proper interval graph. Now as from Theorem 3.12 it follows that proper interval graphs are same as proper central MPTG, we can take (I_u, c_u) to be a proper central MPTG representation

of G where $I_u = [a_u, b_u]$, c_u denotes the center point of I_u for each $u \in V$. We will show that G is a 50% maxtolerance graph with respect to the same interval representation. Suppose $uv \in E$. Then $I_u \cap I_v \supseteq \{c_u, c_v\}$. This implies $|I_u \cap I_v| \ge \frac{|I_u|}{2}$, $\frac{|I_v|}{2}$ as none of them contains other properly. Hence $|I_u \cap I_v| \ge \frac{1}{2} \max\{|I_u|, |I_v|\}$ which imply $uv \in E$ in its 50% max-tolerance representation. Again $uv \notin E$ implies $c_u \notin I_v$ or $c_v \notin I_u$. If $c_u \le c_v$ then $c_u \notin I_v \Rightarrow a_v > c_u \Rightarrow |I_u \cap I_v| < \frac{|I_u|}{2}$. Also $c_v \notin I_u \Rightarrow c_v > b_u \Rightarrow |I_u \cap I_v| < \frac{|I_v|}{2}$. From this we can conclude that $|I_u \cap I_v| < \frac{1}{2} \max\{|I_u|, |I_v|\}$. Hence $uv \notin E$ in its 50% max-tolerance representation. \Box **Proposition 4.5.** Any cycle $C_n, (n \ge 3)$ is a 50% max-tolerance graph.

Proof. We prove C_n to be a 50% max-tolerance graph by constructing interval representation $\{I_{v_i} = [a_i, b_i] | v_i \in V\}$ where the vertex set V is labelled clockwise starting in an arbitrary vertex. Let c_i be the center point of I_{v_i} and tolerance $t_{v_i} = \frac{|I_{v_i}|}{2}$.

- Let n = 3. We associate intervals $I_{v_1} = I_{v_2} = I_{v_3} = [1, 2]$.
- Let n = 4. We associate intervals $I_{v_1} = [1, 4.6], I_{v_2} = [2, 4], I_{v_3} = [2.9, 4.9], I_{v_4} = [2.7, 6.3].$
- Let n = 5. We associate intervals $I_{v_1} = [10, 30], I_{v_2} = [16, 28], I_{v_3} = [18, 24], I_{v_4} = [15, 21], I_{v_5} = [9, 21].$
- Let $n \ge 6$. We prove now in two cases considering n even and odd. We define

$$k = \frac{n}{2}$$
 when n is even, and $k = \frac{n+1}{2}$ when n is odd.

When *n* is even we define $I_{v_1} = [2 - n, n], I_{v_2} = [1, n], I_{v_i} = [1, 1 + \frac{k - 1}{2^{i - 4}}]$ for $i \in \{3, \dots, k\}, I_{v_{k+1}} = [1 - \frac{k - 1}{2^{k - 3}}, 1 + \frac{k - 1}{2^{k - 3}}], I_{v_j} = [1 - \frac{k - 1}{2^{n - j - 2}}, 1]$ for $j \in \{k + 2, \dots, n\}.$

We verify that the above representation indeed give 50% max-tolerance representation of $C_n, n \ge 6$.

Claim 1: We will show that v_1 is adjacent to only v_2, v_n .

We note $|I_{v_1} \cap I_{v_2}| = |I_{v_2}| = n-1 = \frac{|I_{v_1}|}{2}$. This implies $v_1 \leftrightarrow v_2$. Also since $n \ge 6, 5-2n < 2-n$. Hence $a_n < a_1$. Also $b_n = 1 < n = b_1$. So $|I_{v_1} \cap I_{v_n}| = |[a_1, b_n]| = |[2 - n, 1]| = n - 1 = \frac{|I_{v_1}|}{2} > n - 2 = \frac{|I_{v_n}|}{2}$. This implies $v_1 \leftrightarrow v_n$. Next we show $v_1 \leftrightarrow v_i$ for $i \in \{3, \ldots, k\}$. Clearly 2 - n < 1 implies $a_1 < a_i$. Also $n - 2 < (n - 1)2^{i-3}$ for $i \ge 3$. Hence

$$1 + \frac{k - 1}{2^{i - 4}} < n \tag{4.4}$$

Therefore $b_i < b_1$. Hence $|I_{v_1} \cap I_{v_i}| = |I_{v_i}| = \frac{k-1}{2^{i-4}} < n-1$ from $(4.4) = \frac{|I_{v_1}|}{2}$. Hence $v_1 \nleftrightarrow v_i$. Analogously $v_1 \nleftrightarrow v_j$ for $j \in \{k+2, \ldots, n-1\}$. Again $n-2 < (n-1)2^{k-2}$ implies

$$1 + \frac{k-1}{2^{k-3}} < n \tag{4.5}$$

Hence $a_{k+1} = 1 - \frac{k-1}{2^{k-3}} > 2 - n = a_1$. Similarly, $b_{k+1} = 1 + \frac{k-1}{2^{k-3}} < n = b_1$. Hence $|I_{v_1} \cap I_{v_{k+1}}| = |I_{v_{k+1}}| = \frac{k-1}{2^{k-4}} = \frac{n-2}{2^{k-3}} < n - 1 = \frac{|I_{v_1}|}{2}$. Hence $v_1 \nleftrightarrow v_{k+1}$.

Claim 2: We will show that v_2 is adjacent to only v_1, v_3 .

From previous case $v_2 \leftrightarrow v_1$. Note that $|I_{v_2} \cap I_{v_3}| = |[1, n-1]| = n-2 > \frac{n-1}{2} = \frac{|I_{v_2}|}{2}, \frac{n-2}{2} = \frac{|I_{v_3}|}{2}$. Hence $v_2 \leftrightarrow v_3$. Next to show $v_2 \nleftrightarrow v_i$ for $i \in \{4, \dots, k\}$ we note, $a_2 = a_i = 1$ and $b_i < b_1$ [from(4.4)] = $b_{2} \text{ for } i \in \{4, \dots, k\}. \text{ Hence } |I_{v_{2}} \cap I_{v_{i}}| = |I_{v_{i}}| = \frac{k-1}{2^{i-4}} = \frac{n-2}{2^{i-3}} < \frac{n-1}{2} = \frac{|I_{v_{2}}|}{2}. \text{ This implies } v_{2} \nleftrightarrow v_{i} \text{ for } i \in \{4, \dots, k\}. \text{ Again } |I_{v_{2}} \cap I_{v_{j}}| = \{1\} < \frac{n-1}{2} = \frac{|I_{v_{2}}|}{2}. \text{ Hence } v_{2} \nleftrightarrow v_{j} \text{ for } j \in \{k+2, \dots, n\}.$ Now clearly $a_{k+1} < 1 = a_{2}$ and $b_{k+1} < n = b_{2}$ from (4.5). Hence, $|I_{v_{2}} \cap I_{v_{k+1}}| = |[a_{2}, b_{k+1}]| = \frac{k-1}{2^{k-3}} = \frac{n-2}{2^{k-2}} < \frac{n-1}{2} = \frac{|I_{v_{2}}|}{2}. \text{ Hence } v_{2} \nleftrightarrow v_{k+1}.$

Claim 3: Vertex v_i where $i \in \{3, \ldots, k\}$ is adjacent to only v_{i-1}, v_{i+1} .

From previous case $v_3 \leftrightarrow v_2$. Let $v_i, v_{i'}$ be two vertices where $i, i' \in \{3, \dots, k\}$. Let i < i'. Then clearly $b_i > b_{i'}$. Hence $|I_{v_i} \cap I_{v_{i'}}| = |I_{v_{i'}}| = \frac{k-1}{2^{i'-4}}$. Now if i' = i+1, then $\frac{k-1}{2^{i'-4}} = \frac{k-1}{2^{i-3}} = \frac{|I_{v_i}|}{2} > \frac{|I_{v_{i'}}|}{2}$ clearly. Hence $v_i \leftrightarrow v_{i'}$ in this case. For i' > i+1, $|I_{v_i} \cap I_{v_{i'}}| = \frac{k-1}{2^{i'-4}} < \frac{k-1}{2^{i-3}} = \frac{|I_{v_i}|}{2}$. Hence $v_i \leftrightarrow v_{i'}$. Also one can check trivially that $a_{k+1} < 1 = c_{k+1} = a_i$ where $i \in \{3, \dots, k\}$. Moreover $|I_{v_k} \cap I_{v_{k+1}}| = |[c_{k+1}, b_{k+1}]| = |[1, 1 + \frac{k-1}{2^{k-3}}]| = \frac{k-1}{2^{k-3}} = \frac{|I_{v_k}|}{2}$. Hence $v_k \leftrightarrow v_{k+1}$. Again $a_i = c_{k+1} = 1$ and $b_{k+1} = 1 + \frac{k-1}{2^{k-3}} = c_k < b_k = 1 + \frac{k-1}{2^{k-4}} < 1 + \frac{k-1}{2^{i-4}} = b_i$ for $3 \le i < k$. Hence $|I_{v_i} \cap I_{v_{k+1}}| = \frac{k-1}{2^{k-3}} < \frac{k-1}{2^{i-3}} = \frac{|I_{v_i}|}{2}$. This shows $v_i \leftrightarrow v_{k+1}$ for $3 \le i < k$. Now $|I_{v_i} \cap I_{v_j}| = \{1\}$ for $i \in \{3, \dots, k\}, j \in \{k+2, \dots, n\}$. Hence $v_i \leftrightarrow v_j$.

Claim 4: Vertex v_{k+1} is adjacent to only v_k, v_{k+2} .

Claim 5: Vertex v_i where $i \in \{k + 2, ..., n\}$ is adjacent to only v_{i-1}, v_{i+1} .

Claim 4,5 can be shown similarly to the previous cases.

When *n* is odd we define $I_{v_1} = [2 - n, n], I_{v_2} = [1, n], I_{v_i} = [1, 1 + \frac{n-2}{2^{i-3}}]$ for $i \in \{3, \dots, k\}, I_{v_{k+1}} = [1 - \frac{3n-6}{2^{k-1}}, 1 + \frac{3n-6}{2^{k-1}}], I_{v_j} = [1 - \frac{3n-6}{2^{n+1-j}}, 1]$ for $j \in \{k+2, \dots, n\}$.

This proof is analogous to the case when n is even.

Remark 4.6. Combining Proposition 4.4 and Proposition 4.5 along with results of [18] we can easily conclude that proper intervals graphs and C_n both belong to the intersection class of central MPTG and 50% max-tolerance graphs.

5 Central interval catch digraphs

In 1984, Maehera posed the following conjecture:

Maehera's conjecture[15]: If a digraph G has no induced subdigraph isomorphic to one of the digraphs in Figure 4 and $A^*(G)$ satisfy the consecutive 1's property for rows, then G is a central ICD.

We show the digraphs G_1, G_2, G_3 in Example 5.2 disprove the conjecture. To see this we obtain the following necessary condition of a central ICD.

Proposition 5.1. Let G = (V, E) be a central ICD. Then there is an ordering of vertices $V = \{v_1, v_2, \ldots, v_n\}$ which satisfies (2.1) and the following condition:

for any
$$i < j$$
, either $i_1 \leq j_1$ or, $i_2 \leq j_2$, (5.1)

where i_1 and i_2 be the least and the highest numbers such that $i_1 = i$ or, $v_i v_{i_1} \in E$ and $i_2 = i$ or, $v_i v_{i_2} \in E$ for each i = 1, 2, ..., n.




Figure 4: Maehara's forbidden digraphs for central ICD

Proof. Let G = (V, E) be represented by $\{(I_i, c_i) | i \in V\}$ where $I_i = [a_i, b_i]$ is an interval and c_i be its center point. We arrange vertices of G according to the increasing order of these center points. It is easy to check G satisfy (2.1) with respect to this ordering. On contrary lets assume for some i < j, $i_1 > j_1$ and $i_2 > j_2$. Then $j_1 < i_1 \leq i < j \leq j_2 < i_2$. Now $v_j v_{j_1} \in E, v_i v_{j_1} \notin E$ imply $c_{j_1} \in I_j$ and $c_{j_1} \notin I_i$. Since $c_{j_1} \notin [a_i, b_i]$, either $c_{j_1} < a_i$ or $c_{j_1} > b_i$. If $c_{j_1} > b_i$, then $c_{i_1} \ge c_{j_1} > b_i$ as $i_1 > j_1$. But then $v_i v_{i_1} \notin E$ which is a contradiction. Thus $a_j \le c_{j_1} < a_i \le c_i \le c_j$. Hence $[a_i, c_i] \subset [a_j, c_j]$. Now as $v_i v_{i_2} \in E$, $c_{i_2} \in I_i \subset I_j$ which implies $v_j v_{i_2} \in E$. But then $i_2 \le j_2$ which is again a contradiction. Hence the proof follows.




Figure 5: Forbidden digraphs for central ICD when |V| = 4 and their augmented adjacency matrix.

Example 5.2. Consider the digraphs $G_k = (V_k, E_k)$ with vertex set V_k and edge set E_k for k = 1, 2, 3, 4 in Figure 5. Note that G_4 is the digraph (a) in Figure 4. From Theorem 2.1 one can easily verify that $v_1 < v_2 < v_3 < v_4$ and its reverse ordering are the only possible ICD ordering of V_k for each graph G_k (i.e., for which $A^*(G_k)$ satisfies the consecutive 1's property for rows). Now this augmented adjacency matrix (in Figure 5) shows a contradiction to (5.1) for i = 2, j = 3 for each k. Thus these graphs are not central ICD.

In the following we characterize central ICD. Let \mathbb{R}^+ be the set of all positive real numbers.

Theorem 5.3. Let G = (V, E) be a simple digraph. Then G is a central ICD if and only if there exist a distinct labeling $f : V \longrightarrow \mathbb{R}^+$ of vertices ² which satisfies the following condition:

$$d(i,j) < d(i,k) \text{ for all } i,j,k \in \{1,2,\dots,n\} \text{ such that } v_i v_j \in E \text{ and } v_i v_k \notin E$$

$$(5.2)$$

where $d(i, j) = |f(v_i) - f(v_j)|$ for all $i, j \in \{1, 2, ..., n\}$. (i.e., every out-neighbor distance is less than every non-out-neighbor distance from a vertex)

 $^{^{2}}$ Given a positive real number labeling, one can easily obtain a positive rational number labeling with slight adjustment and again those can be changed to positive integers by scaling as required. Thus we note that natural number labeling will produce the same class of digraphs.

Proof. Suppose G = (V, E) be a central ICD with $V = \{v_1, v_2, \dots, v_n\}$ and

 $\{(I_i, c_i) \mid i = 1, 2, ..., n\}$ be a central point-interval representation of G where v_i corresponds to (I_i, c_i) , i.e., $v_i v_j \in E$ if and only if $c_j \in I_i$ and assume that v_i 's are ordered according to increasing sequence of c_i 's (without loss of generality, we also assume that c_i 's are distinct). Define a labeling $f : V \longrightarrow \mathbb{R}^+$ by $f(v_i) = c_i$. This vertices are ordered according to increasing order of their labels. Now let $i, j, k \in \{1, 2, ..., n\}$ such that $v_i v_j \in E$ and $v_i v_k \notin E$. We consider following cases:

Case I: i < j < k or, k < j < i. Then d(i,k) = d(i,j) + d(j,k) > d(i,j) for d(j,k) > 0 as c_i 's are distinct.

Case II: i < k < j or, j < k < i. These cases are not possible by (2.1) as G is an ICD.

Case III: j < i < k or, k < i < j. Now $v_i v_j \in E$ and $v_i v_k \notin E$ imply $c_j \in I_i$ but $c_k \notin I_i$. Let $r = \frac{|I_i|}{2}$. Since c_i is the central point of I_i , we have $I_i = [c_i - r, c_i + r]$. Then $d(i, j) = |c_i - c_j| < r < |c_i - c_k| = d(i, k)$.

Conversely, suppose G satisfies (5.2) with a labeling f. Let us arrange vertices according to increasing order of their labels. For each i = 1, 2, ..., n, define $c_i = f(v_i)$. Let i_1 and i_2 be the least and the highest numbers such that $i_1 = i$ or, $v_i v_{i_1} \in E$ and $i_2 = i$ or, $v_i v_{i_2} \in E$. Note that $i_1 \leq i \leq i_2$. Define $r_i = \max \{d(i, i_1), d(i, i_2)\}$ and $I_i = [c_i - r_i, c_i + r_i]$. We show that $\{(I_i, c_i) \mid i = 1, 2, ..., n\}$ is a central point-interval representation of G, i.e., G is a central ICD. As c_i is the center point of I_i for each $i \in \{1, ..., n\}$, it is sufficient to prove that G is an ICD.

We verify (2.1) to show that G is an ICD. Let i < j < k and $v_i v_k \in E$. Now d(i,k) = d(i,j) + d(j,k) > d(i,j). So $v_i v_j \in E$. Let $v_k v_i \in E$. Again d(k,i) = d(k,j) + d(j,i) > d(k,j). So $v_k v_j \in E$ as required.

Example 5.4. Consider the digraph D_{10} in Figure 12. It can be easily checked that $v_1 < v_2 < v_3 < v_4$ is an ICD ordering of D_{10} and D_{10} is a central ICD with the labeling: $f(v_1) = 1$, $f(v_2) = 3$, $f(v_3) = 4$ and $f(v_4) = 6$.

Interestingly a family of (undirected) graphs satisfying the condition analogous to (5.2) becomes a well known class of graphs, namely, proper interval graphs.

Theorem 5.5. Let G = (V, E) is a graph. Then G is a proper interval graph if and only if there exist an ordering of vertices and a distinct labeling $f : V \to \mathbb{R}^+$ of vertices such that

$$d(u,v) < d(u,w) \text{ for all } u,v,w \in V \text{ such that } uv \in E \text{ but } uw \notin E,$$

$$(5.3)$$

where d(u, v) = |f(u) - f(v)| for all $u, v \in V$.

Proof. Let G = (V, E) be a proper interval graph. By Theorem 2.3 the reduced graph $\tilde{G} = (\tilde{V}, \tilde{E})$ of G is an induced subgraph of $G(n, r) = (V_n, E')$ for some $n, r \in N$ with n > r. Let $V_n = \{x_1, x_2, \ldots, x_n\}$ such that $x_i \leftrightarrow x_j$ in G(n, r) if and only if $0 < |i - j| \le r$. Let $\tilde{V} = \{x_{i_1}, x_{i_2}, \ldots, x_{i_m}\}$ such that $i_1 < i_2 < \ldots < i_m$. For convenience, we write $y_j = x_{i_j}$ for $j = 1, 2, \ldots, m$. Define $f : V \to \mathbb{R}^+$ by $f(u) = i_j + \frac{k}{z+1}$ if u is a k^{th} copy among z copies of y_j (for any but a fixed permutation of them). We arrange the vertices of V according to the increasing order of vertices in \tilde{V} keeping copies of same vertices together. Let $u, v, w \in V$ such that $u \leftrightarrow v$ and $u \nleftrightarrow w$. Let $f(u) = i_p, f(v) = i_q$ and $f(w) = i_t$ for some $p, q, t \in \{1, 2, \ldots, n\}$. Then $|i_p - i_q| \le r$ and $|i_p - i_r| > r$. So d(u, v) = |f(u) - f(v)| < |f(u) - f(w)| = d(u, w). Therefore d(u, v) < d(u, w) for all $u, v, w \in V$ such that $u \leftrightarrow v$ and $u \leftrightarrow w$.

Conversely, let G = (V, E) satisfies (5.3) with a labeling f. We arrange the vertices of G according to the increasing order of their labels.

Let i < j < k, where $i, j, k \in \{1, 2, ..., n\}$ and $v_i \leftrightarrow v_k$. Then $d(v_i, v_j) = d(v_i, v_k) - d(v_j, v_k) < d(v_i, v_k)$. Also $d(v_j, v_k) = d(v_i, v_k) - d(v_i, v_j) < d(v_i, v_k)$. Then by (5.3), $v_i \leftrightarrow v_j$ and $v_j \leftrightarrow v_k$. Thus G satisfies umbrella property and hence by Theorem 2.3, G becomes a proper interval graph.

6 Oriented interval catch digraph

We first note that the characterization of an oriented ICD is obvious. A simple digraph G = (V, E) is an oriented ICD if and only if G is oriented and there exists an ordering of vertices in V satisfying (2.1) or, equivalently, there exists an ordering of vertices in V such that $A^*(G) = (a_{i,j})$ satisfies consecutive 1's property for rows and for each pair $i \neq j$, $a_{i,j} = 1$ implies $a_{j,i} = 0$. Now we study some important properties of oriented ICD.

Theorem 6.1. An oriented ICD is a directed acyclic graph (DAG).

Proof. Let G be an oriented ICD. We first show that G has no induced directed cycles. Note that any induced subdigraph of an ICD is also an ICD by definition. Now it is easy to check that no induced directed cycles of length ≥ 3 satisfies (2.1) and so they are not ICD. Finally, since the digraph is oriented, there are no 2-cycles.

Now we use induction on the length of the cycle. Since G has no induced directed cycles, in particular, G has no directed cycles of length 3. Suppose there are no directed cycles of length less than k and G has a directed cycle C of length k > 3. Since C is not induced, there must be a chord. Interestingly, every such arc (chord) forms a smaller directed cycle on one side of it along with some arcs of C. This contradicts the induction hypothesis.

The following corollary is immediate from the above theorem.

Corollary 6.2. Let G = (V, E) be an oriented ICD. Then every induced 3-cycle of U(G) is transitively oriented in G.

In [15] Maehara proved that an acyclic ICD is a central ICD. Thus we have the following:

Corollary 6.3. Every oriented ICD is a central ICD.

Definition 6.4. Let G be an oriented digraph and C be an even chordless cycle of G which is not a directed cycle. Then C is said to be *alternatively oriented* if any two arcs of C with common end point have opposite directions.

Proposition 6.5. Let G = (V, E) be an oriented ICD and C be a chordless 4-cycle of G. Then C is alternatively oriented.

Proof. Let $C = (v_1, v_2, v_3, v_4, v_1)$ be a chordless 4-cycle in G. By Theorem 6.1, C is not a directed 4-cycle. Then there are only 3 other (non-isomorphic) options which are described in Figure 6. The first option T_1 is alternatively oriented. The two other options are not ICD. For example, in the case of T_2 , in order to find an ICD ordering we need the elements of each of the sets $\{v_1, v_2, v_4\}$, $\{v_2, v_3\}$ and $\{v_3, v_4\}$ to be consecutive. Since the ordering is linear and we require elements of sets $\{v_2, v_3\}$ and $\{v_3, v_4\}$ to be consecutive in that ordering. So we have either v_2, v_3, v_4 or v_4, v_3, v_2 consecutive in that ordering. But then $\{v_1, v_2, v_4\}$ cannot be consecutive. So T_2 is not ICD. Similarly, it can be shown that T_3 is not an ICD.

Now being an oriented acyclic digraph, an oriented ICD has some rich properties. For example, a unilaterally connected oriented ICD has unique source (vertex of indegree zero) and unique sink (vertex of outdegree




Figure 6: Orientation of 4-cycles which are not directed 4-cycles.

zero). Hence it possesses unique hamiltonian path (see [1]). In the following we characterize oriented ICDs which are tournaments.

A Ferrers digraph is a directed graph G = (V, E) whose successor sets are linearly ordered by inclusion, where the successor set of $u \in V$ is its set of out-neighbors $\{v \in V \mid uv \in E\}$. A (0, 1)-matrix M is a Ferrers matrix if 1's are clustered in a corner of M. A digraph G is Ferrers digraph if and only if there exists a permutation of vertices of G such that its adjacency matrix is a Ferrers matrix [2]. For a (0, 1)-matrix M, \overline{M} denotes the matrix obtained from M by interchanging 0's and 1's.

Theorem 6.6. Let G = (V, E) be an oriented ICD. Then G is a tournament if and only if there is an ordering of vertices of G with respect to which the augmented adjacency matrix $A^*(G)$ takes one of the following forms:

$$A^{*}(G) = \begin{bmatrix} M & | F \\ \hline F^{T} & | N \end{bmatrix} = \begin{bmatrix} v_{1} & v_{2} & \cdot & v_{l} & v_{l+1} & \cdot & v_{k} & v_{n} \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ v_{2} & 0 & 1 & 1 & 1 & 1 & 1 & 0 & 0 \\ \cdot & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ v_{l+1} & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ \cdot & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 \\ \cdot & v_{k} & 0 & 1 & 1 & 1 & 1 & 1 & 0 \\ v_{k} & 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & \cdot & v_{n} \\ v_{1} & 1 & 0 & 0 & 0 \\ \cdot & v_{n} & 1 & 1 & 1 & 1 & 1 & 0 \\ v_{k} & 0 & 1 & 1 & 1 & 1 & 1 & 1 \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & \cdot & v_{n} \\ v_{1} & 1 & 0 & 0 \\ \cdot & v_{n} & 1 & 1 & 1 & 1 & 1 \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ v_{1} & 1 & 0 & 0 \\ \cdot & v_{n} & 1 & 1 & 1 & 1 \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ v_{1} & 1 & 0 & 0 \\ \cdot & v_{n} & 1 & 1 & 1 \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ \cdot & v_{n} & v_{n} & v_{n} \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ \cdot & v_{n} & v_{n} & v_{n} \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ \cdot & v_{n} & v_{n} & v_{n} & v_{n} \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & \cdot & v_{n} \\ \cdot & v_{n} & v_{n} & v_{n} & v_{n} & v_{n} \end{bmatrix} orr A^{*}(G) = \begin{bmatrix} v_{1} & v_{n} & v_{n} & v_{n} \\ \cdot & v_{n} & v_{n} & v_{n} & v_{n} & v_{n} & v_{n} \\ \cdot & v_{n} \\ \cdot & v_{n} \\ \cdot & v_{n} & v_{n$$

where M is an upper triangular matrix with all entries on and above the principal diagonal are 1 and other enties are 0; N and P are lower triangular matrices with all entries on and below the principal diagonal are 1 and other entries are 0 and F is a Ferrers matrix.

Proof. Let G = (V, E) be an oriented ICD which is a tournament. We order the vertices of V according to the increasing value of the associated points in their corresponding intervals. Let $\{v_1, \ldots, v_n\}$ be the required ordering which is in fact, an ICD ordering, i.e., $\{v_1, \ldots, v_n\}$ satisfies (2.1). Hence $A = A^*(G)$ satisfies consecutive 1's property for rows with respect to this ordering. For each $1 \leq i \leq n$, let $(v_i)_1$ and $(v_i)_2$ be the column numbers where first and last 1 occur in the *i*-th row. We denote the (i, j)-th entry of the matrix A by $a_{i,j}$.

Case I: $(v_1)_2 = k > 1$.

In this case, $a_{1,2} = 1$ (as k > 1). So $a_{2,1} = 0$ as G is a tournament. Hence $(v_1)_1 = 1 < 2 = (v_2)_1$. Now the following subcases may happen.

- **Case I(a):** If $(v_2)_2 = 2$ then $2 = (v_2)_2 = (v_2)_1$ which implies the vertex v_2 has out degree 0. In this case we define l = 2 (note that the matrix M in the statement is of the order $l \times l$).
- **Case I(b):** If $(v_2)_2 > 2$, then we can show that $(v_2)_2 \le (v_1)_2$. If not, let $(v_2)_2 > (v_1)_2$. Then $a_{1,(v_2)_2} = 0$ which implies $a_{(v_2)_2,1} = 1$ as G is a tournament. Now $a_{2,(v_2)_2} = 1$ by definition. Then $a_{(v_2)_2,2} = 0$. But we

have $a_{(v_2)_2,1} = 1$ and $a_{(v_2)_2,(v_2)_2} = 1$. This contradicts the consecutive 1's property for the $(v_2)_2$ -th row as $1 < 2 < (v_2)_2$. Hence $(v_2)_2 \le (v_1)_2$.

Now since $(v_2)_2 > 2$, $a_{2,3} = 1$ and hence $a_{3,2} = 0$. Thus $(v_3)_1 = 3$. Let

$$S = \{i \mid (v_i)_1 = i, \ (v_i)_2 \le (v_{i-1})_2, \ 2 \le i \le n\}.$$

Already we have shown that $2 \in S$. This implies $S \neq \emptyset$. As |V| is finite S must have a maximum, say, l. First we will show that $\{i \mid 2 \le i \le l\} \subseteq S$. As $2 \in S$, applying induction we assume $\{2, 3, \ldots, j\} \subseteq S$. If $j + 1 \le l$, then we show that $j + 1 \in S$.

If $(v_j)_2 = j$ then $a_{j,l} = 0$ as l > j. Thus $a_{l,j} = 1$ and so $(v_l)_1 \leq j < l$ which contradicts $l \in S$. Therefore $(v_j)_2 > j$. But then $a_{j,j+1} = 1$ which implies $a_{j+1,j} = 0$ and so $(v_{j+1})_1 = j + 1$. Next we show $(v_{j+1})_2 \leq (v_j)_2$.

On contrary let $(v_{j+1})_2 > (v_j)_2$. Since $(v_j)_2 > j$, we have $(v_{j+1})_2 \ge j + 1$. Now if $(v_{j+1})_2 = j + 1$, then $(v_{j+1})_2 \le (v_j)_2$ as $(v_j)_2 > j$. So suppose $(v_{j+1})_2 > j + 1$. Now $a_{j,(v_{j+1})_2} = 0$ as we have assumed $(v_{j+1})_2 > (v_j)_2$. Thus $a_{(v_{j+1})_2,j} = 1$. Again $a_{j+1,(v_{j+1})_2} = 1$ implies $a_{(v_{j+1})_2,j+1} = 0$. But $a_{(v_{j+1})_2,(v_{j+1})_2} = 1$. This contradicts the consecutive 1's property for $(v_{j+1})_2$ -th row as $j < j + 1 < (v_{j+1})_2$. So we must have $(v_{j+1})_2 \le (v_j)_2$ and hence $j+1 \in S$. This completes the induction. Therefore $\{i \mid 2 \le i \le l\} \subseteq S$ and so

$$(v_i)_1 = i, (v_i)_2 \le (v_{i-1})_2$$
 for all *i* with $2 \le i \le l$.

Now we will show $l \leq k = (v_1)_2$ and $(v_l)_2 = l$. On contrary let l > k. Then $a_{1,l} = 0$ as $k = (v_1)_2$. This imply $a_{l,1} = 1$. Then $(v_l)_1 = 1 < l$ which contradicts $l \in S$. Therefore $l \leq k$.

Now if $(v_l)_2 > l$ then $a_{l,l+1} = 1$ which imply $a_{l+1,l} = 0$, i.e., $(v_{l+1})_1 = v_{l+1}$. But $l+1 \notin S$. This implies $(v_{l+1})_2 > (v_l)_2 > l$. Hence $a_{l,(v_{l+1})_2} = 0$ which imply $a_{(v_{l+1})_2,l} = 1$. But $a_{(v_{l+1})_2,l+1} = 0$ as $a_{l+1,(v_{l+1})_2} = 1$ and $a_{(v_{l+1})_2,(v_{l+1})_2} = 1$. Thus $(v_{l+1})_2 \neq l+1$ and so $(v_{l+1})_2 > l+1$. Then we have a contradiction in the consecutive 1's property for the $(v_{l+1})_2$ -th row as $l < l+1 < (v_{l+1})_2$. So $(v_l)_2 = l$. Already we have $(v_l)_1 = l$. Thus the *l*-th row contains only one 1 at the position (l, l).

Therefore the rows $\{1, 2, \ldots, l\}$ form upper triangular matrix M with all entries on and above diagonal as 1 and F becomes a Ferrers matrix formed by the rows $\{1, 2, \ldots, l\}$ and columns $\{l + 1, l + 2, \ldots, n\}$ as $(v_l)_2 = l \leq (v_{l-1})_2 \leq \ldots \leq (v_1)_2 = k$. As G forms a tournament, matrix formed by rows $\{l + 1, l + 2, \ldots, n\}$ and columns $\{1, 2, \ldots, l\}$ is $\overline{F^T}$.

Now the only remaining thing is to show $(v_i)_2 = i$ for all i > l. On contrary, let there exist some i where $l < i \le n$ for which $(v_i)_2 > i$. From this it follows $a_{i,i+1} = 1$ which implies $a_{i+1,i} = 0$ i.e., $(v_{i+1})_1 = i + 1$. Now since $(v_l)_2 = l$, $a_{l,i+1} = 0$ and so $a_{i+1,l} = 1$ which is a contradiction as $(v_{i+1})_1 = i + 1 > l$. So we have $(v_i)_2 = i$ for all i > l. Thus we get the required lower triangular matrix N formed by the vertices $\{v_{l+1}, v_{l+2}, \ldots, v_n\}$.

Case II: $(v_1)_2 = v_1$.

In this case $a_{1,j} = 0$ for all j such that $1 < j \le n$. Since G is a tournament this implies $a_{j,1} = 1$ for all such j. Then all the entries in the first column of A are 1. Also since A is the augmented adjacency matrix of G, $a_{j,j} = 1$ for all j = 1, 2, ..., n. Moreover, A satisfies consecutive 1's property for rows. Thus we have $a_{j,i} = 1$ for all i such that $1 \le i \le j$, for all j = 1, 2, ..., n. Finally since G is a tournament $a_{i,j} = 0$ for all i < j, for all j = 1, 2, ..., n. Therefore A is the lower triangular matrix P with all entries 1 on and below the principal diagonal.




Figure 7: D_1, D_2, K (left to right) (undirected edges can have any direction)

The converse part is obvious from the structure of $A^*(G)$ as both the matrices described in the statement have consecutive 1's property for rows and for each pair $i \neq j$, $a_{i,j} = 0$ if and only if $a_{j,i} = 1$. Thus G is an ICD which is a tournament.

Next we characterize an oriented ICD G for which its underlying undirected graph U(G) is a tree. We characterize the aforesaid digraphs in terms of forbidden subdigraphs.

Theorem 6.7. Let G = (V, E) be a oriented digraph such that U(G) is a tree. Then G is an oriented ICD if and only it does not contain any of D_1, D_2, K as an induced subdigraph.

Proof. Let G be a oriented ICD. As D_1 contains v, v_4, v_5, D_2 contains v, v_2, v_3 and K contains x, y, z as diasteroidal triple, from Theorem 2.2 it follows that G can not contain D_1, D_2, K as induced subdigraph.

Conversely let G be a oriented digraph which does not contain D_1, D_2, K as induced subdigraph. Let S be the set of all chains connecting every pair of vertices in G and $\vec{P} = (x = a_0, a_1, \dots, a_{l-1}, a_l = y)$ be a chain of maximum length (say l) between two vertices x, y (say) of V. As U(G) does not contain K as induced subdigraph, it becomes a caterpillar and $U(\vec{P})$ becomes the spine of the caterpillar. Let $a_i, i \neq 1, l-1$ be an internal vertex of $V(\vec{P})$. Then looking at D_1 and D_2 one can verify that a_i can not possess any inneighbour from $V \setminus V(\vec{P})$. Only a_1, a_{n-1} can have inneighbour from $V \setminus V(\vec{P})$ and that must also be one in number as D_2 is forbidden in G. Let $k_1(k_2)$ be the only possible inneighbour of $a_1(a_{n-1})$ outside $V(\vec{P})$ in G. Now if any of the vertices k_1, k_2 presents in G then we construct the modified chain $\hat{P} = (b_0, b_1, \dots, b_{l-1}, b_l)$ by only replacing the vertices a_0, a_l of $V(\vec{P})$ by k_1, k_2 respectively. Note that \vec{P} is of same length l as \vec{P} . We now attach a variable p_{v_i} to each vertex v_i of G. Then starting from p_{b_0} we place the points p_{b_i} corresponding to the vertices b_i of the chain \hat{P} in increasing order in a single sequence P_S (say) along the real line. From construction it is easy to observe that in the chain \hat{P} there is no internal vertex containing inneighbours from $V \setminus V(\hat{P})$. At *i* th step if b_i possesses outneighbours b_{i^1}, \ldots, b_{i^m} then we place $p_{b_{i^1}}, \ldots, p_{b_{i^m}}$ between $p_{b_i}, p_{b_{i+1}}$ in any order in P_S . Now we attach numbers starting from 1 to n with each point of P_S where |V| = n. Let $i_1(i_2)$ be the leftmost (rightmost) indices of the points, which are outneighbours of v_i in the sequence P_S (consider $i_1 \le i \le i_2$) and l_i, r_i, t_i be the numbers attached to p_{i_1}, p_{i_2}, p_i respectively. We assign the interval $I_i = [l_i, r_i]$ and the corresponding point $p_i = t_i$ as the value attached to p_{v_i} in P_S for each vertex $v_i \in V$. As outneighbours of each vertex occur consecutively right or left to v_i in P_S , one can easily check from Theorem 2.1 that the above representation actually gives us the oriented ICD representation of G. \Box

7 Proper interval catch digraph

Definition 7.1. An interval catch digraph G = (V, E) having a representation $\{(I_v, p_v) | v \in V\}$ where no interval contains other properly is called *proper ICD*.

													p_1	p_2	p_3	p_4	p_5			
	v_1	v_2	v_3	v_4	v_5		v_1	v_2	v_3	v_4	v_5		1.5	2	4	6	8			
v_1	1	1	0	0	0	v_1	1	1	0	0	0]	v_1	v_2	v_3	v_4	v_5			
v_2	0	1	1	1	0	v_3	1	1	1	0	0	v_1	1	2	3				[1, 3]	I_1
v_3	1	1	1	0	0	v_2	0	1	1	1	0	v_3			4	5			[1.5, 5]	I_3
v_4	0	0	0	1	1	v_5	0	0	1	1	1	v_2]		6	7		[2, 7]	I_2
v_5	0	0	1	1	1	v_4	0	0	0	1	1	v_5]		8	9	[4, 9]	I_5
						,						v_4						10	[6, 10]	I_4
													L					,		

Figure 8: Augmented adjacency matrix $A = A^*(G)$ of a proper ICD G (left), the row permuted matrix B obtained from A that satisfies MCA property (middle) and a proper ICD representation of G (right).

Let G = (V, E) be a simple digraph. Then the augmented adjacency matrix $A^*(G)$ has a monotone consecutive arrangement (MCA) [2] if and only if it has independent row and column permutations such that 1's appear consecutively in each row and $1_1 \leq 2_1 \leq \ldots \leq n_1$ and $1_2 \leq 2_2 \leq \ldots \leq n_2$ where |V| = n and the values i_1 and i_2 denote the initial column and final column containing 1's in the *i*-th row. Now one can seperate the 1's and 0's by drawing upper and lower stairs (polygonal path from top left to bottom right) as in Figure 8 (right).

In the following we give an adjacency matrix characterization of a proper ICD.

Theorem 7.2. Let G = (V, E) be a simple digraph. Then G is a proper ICD if and only if there exists a vertex ordering $\{v_1, v_2, \ldots, v_n\}$ of V with respect to which the augmented adjacency matrix $A^*(G)$ satisfies the following conditions:

- (1) $A^*(G)$ satisfies consecutive 1's property for rows.
- (2) For $i \neq j$, if $i_1 < j_1$ then $i_2 \leq j_2$ where i_1 and i_2 be the first and last column numbers containing 1 in the *i*-th row where $1 \leq i \leq n$ in $A^*(G)$.

Proof. Let G = (V, E) be a proper ICD with representation $\{(I_i, p_i) \mid i = 1, 2, ..., n\}$ where $V = \{v_1, v_2, ..., v_n\}$, $I_i = [a_i, b_i]$ and p_i 's are distinct for i = 1, 2, ..., n. Then arranging the vertices according to increasing order of p_i 's we have consecutive 1's property for rows in $A^*(G) = (a_{i,j})$. We note that for $i \neq j$, $a_{i,j} = 1$ if and only if $v_i v_j \in E$ if and only if $p_j \in I_i$ if and only if $i_1 \leq j \leq i_2$ by definition.

Now for $i \neq j$ suppose $i_1 < j_1$ and $i_2 > j_2$ hold simultaneously in $A^*(G)$. Then $p_{i_1} \notin I_j$ as $i_1 < j_1$. Again $i_1 < j_1 \leq j$ implies $p_{i_1} < p_{j_1} \leq p_j$. Since $p_j \in I_j$, the interval I_j lies entirely right to p_{i_1} . So we get $a_i < p_{i_1} < a_j$. As the intervals do not contain other properly we have $b_i < b_j$. Now as $p_{i_2} \in I_i$ and $p_{i_2} > p_j$ (for $i_2 > j_2 \geq j$) we get $a_j \leq p_j < p_{i_2} < b_i < b_j$. Then $p_{i_2} \in I_j$ which is a contraction as $i_2 > j_2$. Hence $i_2 \leq j_2$.³

Conversely, let G satisfy conditions (1) and (2). We permute the rows of $A = A^*(G) = (a_{i,j})$ according to the non-decreasing order of i_1 's keeping the columns intact (see Figure 8). If there exist pair of rows i, j for which $i_1 = j_1$, then place *i*-th row prior to *j*-th row when $i_2 < j_2$. If $i_1 = j_1$ and $i_2 = j_2$ then keep *i*-th row prior to *j*-th row only when i < j in A. Let us call the modified matrix by $B = (b_{i,j})$. As A satisfies (2), after permuting only the rows of A it is easy to check that B satisfies MCA. Moreover permutation of rows of A does not affect the adjacency of the graph G, i.e., if *i*-th row of A is shifted to k-th row in B, then

$$a_{i,j} = 1$$
 if and only if $b_{k,j} = 1.$ (7.1)

³Note that, it follows from the condition (2) that "for $i \neq j$, $i_2 < j_2$ implies $i_1 \leq j_1$ ". For otherwise, let $i_2 < j_2$ and $i_1 > j_1$. Then by the condition (2), we have $j_2 \leq i_2$ (as $j_1 < i_1$) which is a contradiction.

Step I: We show that the diagonal entries of the matrix *B* must be 1.

Suppose the *i*-th row of *B* was the *k*-th row of *A*. If k = i then $b_{i,i} = 1$ as $a_{i,i} = 1$. Suppose k > i in *A*. For each j = 1, 2, ..., i, $j_1 \leq j \leq i$. Now the *k*-th row of *A* moved upward to the *i*-th row position in *B*. This implies there exists at least one such *j* so that $k_1 \leq j_1$. Then $k_1 \leq i < k \leq k_2$. Hence $a_{k,i} = 1$ which implies $b_{i,i} = 1$. Again if k < i in *A* then there exists at least one $j \geq i$ such that $j_1 < k_1$ or, $j_1 = k_1$ and $j_2 < k_2$. Then by the condition (2), in either case, we have $j_2 \leq k_2$. Thus $k_1 \leq k < i \leq j \leq j_2 \leq k_2$. This implies $a_{k,i} = 1$ and hence $b_{i,i} = 1$.

Step II: Defining intervals $I_i = [l_i, r_i]$ for the *i*-th row of *B*.

Now we associate natural numbers in increasing order on the upper stair of B starting from the left top of it. Let b_i be the number on the stair in the *i*-th row where the stretch of consecutive 1's ends and a_i be the number on the stair in the *i*-th column where the stretch of consecutive 1's begins (e.g., in Figure 8, $a_i = 1, 2, 4, 6, 8$ and $b_i = 3, 5, 7, 9, 10$ for i = 1, 2, 3, 4, 5 respectively). Let i_o and i_e denote the first and last column numbers containing 1 in the *i*-th row of B (similar to the construction of i_1 and i_2 in A). Then by definition of a_i and b_i we have $b_i > a_j$ for all $i_o \le j \le i_e$.⁴ Thus

$$b_{i,j} = 1 \Longrightarrow b_i > a_j. \tag{7.2}$$

Let $k_i = \max \{j \mid j_o = i_o\}$. We assign interval $I_i = [l_i, r_i]$ where

$$l_i = a_{i_o} + \frac{i - i_o}{k_i - i_o + 1}$$
 and $r_i = b_i$.

By (7.2), $b_i > a_{i_o}$ and so $b_i \ge a_{i_o} + 1 > l_i$. Thus $l_i < r_i$ for all i = 1, 2, ..., n. Also since B satisfies MCA, one can check that $l_i < l_j$ and $r_i < r_j$ for all i < j. Therefore the set of intervals $[l_i, r_i]$ gives proper representation.

Step III: Defining points p_j for each column j of B (as well as of A).

Let $S_j = \{l_i \mid i \ge j, b_{i,j} = 1\}$. Note that $S_j \ne \emptyset$ as $b_{j,j} = 1$. Assign $p_j = \max\{a_j, \max S_j\}$.

Now $p_j \ge l_j$ as $l_j \in S_j$. Suppose for some $i \ge j$, $b_{i,j} = 1$. Then $i_o \le j \le i_e$ which implies $a_{i_o} \le a_j$. So $l_i < a_{i_o} + 1 \le a_j + 1 \le b_j$. Also since $b_{j,j} = 1$ by Step I, we have $a_j < b_j$ by (7.2). Thus $p_j < b_j$. Hence

$$p_j \in [l_j, r_j]. \tag{7.3}$$

Step IV: We show that $b_{i,j} = 1$ if and only if $p_j \in I_i$.

Case (i): $b_{i,j} = 1$ and i > j.

By definition of S_j , $l_i \in S_j$ and so $p_j \ge l_i$. Again by (7.3), $p_j \le r_j < r_i$ (as i > j). Thus $p_j \in [l_i, r_i] = I_i$.

Case (ii): $b_{i,j} = 1$ and i < j.

As i < j, $l_i < l_j \le p_j$. Also since $b_{i,j} = 1$, by (7.2), $r_i = b_i > a_j$ which implies $r_i \ge a_j + 1$. Moreover, for all t > j with $b_{t,j} = 1$, $l_t < a_j + 1$ as in Step III. Thus $p_j < a_j + 1 \le r_i$. Hence $p_j \in [l_i, r_i] = I_i$.

Case (iii): $b_{i,j} = 0$ and i > j.

We first show that $a_j \leq p_j < a_j + 1$. By definition of p_j , if $p_j = a_j$, then the claim is true. If $p_j = \max S_j$, then again by definition $p_j \geq a_j$. In this case, $p_j = l_x$ for some $x \geq j$ such that $b_{x,j} = 1$. Now $b_{x,j} = 1$ implies $x_o \leq j$ and so $a_{x_o} \leq a_j$. By definition $a_{x_o} \leq l_x < a_{x_o} + 1$. Then $p_j = l_x < a_{x_o} + 1 \leq a_j + 1$ as required.

In this case, the $\{i, j\}$ -th position is left to (or, below) the lower stair, i.e., $j < i_o$. So $a_j < a_{i_o}$ which implies $a_j + 1 \le a_{i_o}$. Then by the above argument, $p_j < a_j + 1 \le a_{i_o} \le l_i$ (by definition of l_i). Thus We have $p_j < l_i$. Hence $p_j \notin [l_i, r_i] = I_i$.

⁴For example, in Figure 8, for i = 2, $a_{2_o} = 1$, $a_{2_e} = 4$ and $b_2 = 5$; for i = 3, $a_{3_o} = 2$, $a_{3_e} = 6$ and $b_3 = 7$; etc.

Case (iv): $b_{i,j} = 0$ and i < j.

the position $\{i, j\}$ is right to (or, above) the upper stair. Hence $b_i < a_j$. But then $r_i = b_i < a_j \le p_j$. So p_j lies right to the interval $[l_i, r_i] = I_i$. Thus $p_j \notin I_i$.

Step V: Finally we show that p_j belongs to the interval corresponding to the vertex v_j .

Note that, if the *j*-th row of A is shifted to the k-th row of B, then the vertex v_j corresponds to the interval I_k . Now by (7.1), we have $b_{k,j} = 1$ as $a_{j,j} = 1$. Then $p_j \in I_k$ by Cases (i) and (ii) of Step IV.

This completes all the verifications. Therefore G is a proper ICD with respect to the above representation. \Box

An unit interval catch digraph (in brief, unit ICD) is an ICD where every interval has the same length. We note the following result without proof as it goes with the same line as the proof of $(1) \iff (2)$ in Theorem 2.3.

Proposition 7.3. Let G be an ICD. Then G is a proper ICD if and only if it is a unit ICD.

Finally we define a digraph as *proper oriented interval catch digraph* (in brief, proper oriented ICD) if it is an oriented ICD where no two intervals are contained in other properly. Hence from Theorem 2.1 and Theorem 7.2 we get the following:

Theorem 7.4. Let G = (V, E) be an oriented digraph. Then G is a proper oriented ICD if and only if there exists a vertex ordering which satisfy the following

for
$$u < v < w$$
, if $uw \in E$ then $uv, vw \in E$ and if $wu \in E$ then $wv, vu \in E$. (7.4)

Proof. Let G = (V, E) be a proper oriented ICD. Then by Theorem 7.2, there exists an ordering of V that satisfies conditions (1) and (2). Let u < v < w be three vertices of V in this ordering and $uw \in E$. Suppose u, v, w correspond to rows i, j, k respectively in $A^*(G) = (a_{i,j})$, where i < j < k. Since the matrix is augmented, $a_{i,i} = 1$. Also $uw \in E$ implies $a_{i,k} = 1$. Thus by consecutive 1's property for rows, we have $a_{i,j} = 1$ which implies $uv \in E$. Now the digraph is oriented. Thus $a_{j,i} = a_{k,i} = 0$. Then $j_1 > i \ge i_1$. Thus by condition (2), $j_2 \ge i_2$. But $a_{i,k} = 1$ and so $i_2 \ge k$. Then $j_2 \ge k$ which implies $a_{j,k} = 1$. Hence $vw \in E$. The other part can be proved similarly.

Conversely, let G = (V, E) be an oriented digraph satisfying the given condition. By Theorem 2.1, it follows that G is an ICD and so it satisfies the condition (1) of Theorem 7.2. Suppose in $A^*(G) = (a_{i,j})$, $i_1 < j_1$ for two distinct rows i, j. Now if we can show that $a_{j,i_2} = 1$, then it follows that $i_2 \leq j_2$ as required. If $i_2 \leq j$, then $i_2 \leq j_2$ as $j \leq j_2$. Let $i_2 > j$. If i < j, then $i < j < i_2$ and $a_{i,i_2} = 1$. Then by the given condition (first part), we have $a_{j,i_2} = 1$. Now if i > j, then $i_1 < j_1 \leq j < i$ and $a_{i,i_1} = 1$. Thus by the given condition (second part), we have $a_{j,i_1} = 1$. But this implies $j_1 \leq i_1$ which is a contradiction.

Remark 7.5. It follows from the proof of the converse part of the above theorem that in the case of a proper oriented ICD G, $A^*(G)$ itself satisfies MCA which may not be true for a proper ICD, in general. We call the ordering mentioned in (7.4) as proper ICD ordering of V.

Next we will give a forbidden subdigraph characterization to all those oriented interval catch digraphs which are proper and their corresponding underlying graphs are chordal.

Theorem 7.6. Let G = (V, E) be an oriented ICD such that U(G) is chordal. Then G is a proper oriented ICD if and only if it does not contain the digraphs of Figure 9 as induced subdigraph.

Proof. Let G be a proper oriented ICD with representation $\{(I_v = [a_v, b_v], p_v) | v \in V\}$ such that U(G) is chordal. Then from Theorem 7.4 there exist a vertex ordering (say \prec) of V satisfying (7.4). Lets assume on contrary that G contains each of the digraphs of Figure 9 as induced subdigraph. Without

loss of generality we assume $v_2 \prec v_4$. As v_2, v_4 are nonadjacent in D_3, D_6, D_7, D_8, D_9 it is easy to check that v_1, v_3 must occur within v_2, v_4 in the above ordering, i.e; $v_2 \prec v_1, v_3 \prec v_4$. Let $v_1 \prec v_3$. From Figure 9 one can observe that $\{v_2, v_1, v_3\}$, $\{v_1, v_3, v_4\}$ form tournaments in the above digraphs. Hence if $v_1v_3 \in E$ then $v_2v_3, v_2v_1, v_1v_4, v_3v_4 \in E$ as G is oriented and \prec satisfies (7.4). Similarly if $v_3v_1 \in E$ then $v_3v_2, v_1v_2, v_4v_1, v_4v_3 \in E$. Both of these cases imply existence of same proper oriented ICD \hat{D} in G (see Figure 10), which is not isomorphic to any of the digraphs D_3, D_6, D_7, D_8, D_9 . If $v_3 \prec v_1$ we get same contradiction.

Next for D_4 one can easily check that either $v_3 \prec v_2 \prec v_4$ or $v_4 \prec v_2 \prec v_3$. In both of these cases it is not possible to place v_1 anywhere in the above ordering as v_1 is adjacent (nonadjacent) to v_2 (v_3, v_4). Similarly in D_5 , $v_1 \prec v_2 \prec v_3$ or $v_3 \prec v_2 \prec v_1$ as v_1, v_3 are nonadjacent in $U(D_5)$. In these cases we can not place v_4 anywhere in the above ordering follows from its adjacency. Hence contradiction arises in both of these cases. Conversely, let G does not contain any of the digraphs of Figure 9 as induced subdigraph. Now as G is an ICD, it must possess an ICD-ordering $P = (v_1, v_2, \ldots, v_n)$ (say). We want to create a proper ICD ordering \hat{P} from P so that it satisfy (7.4) and G become a proper oriented ICD from Theorem 7.4. It is known from Theorem 6.1 that G is acyclic and it can not contain D_2 as induced subdigraph follows from Theorem 6.7. We denote the set of all outneighbours (inneighbours) of a vertex $v_i \in V$ by $N_O[v_i](N_I[v_i])$ (including v_i). From definition of ICD-ordering it is clear that vertices of $N_O[v_i]$ must occur consecutive along P. Below we list some lemmas which we repeatedly use throughout the proof.

Lemma 7.7. Let $v \in V$ and v_l, v_r be two out(in)-neighbours and v_k be an in(out)-neighbour of v in G. Then v_l and v_r must share an edge.

Proof. If v_l, v_r do not have an edge, then the vertices $\{v_l, v, v_k, v_r\}$ induce D_5 or D_4 or D_6 in G, which can not happen. In other case we also get contradiction as D_4, D_8 are forbidden.

Lemma 7.8. Let $v \in V$ and v_l, v_r be two nonadjacent out(in)-neighbours of v in G. Then there exist no common neighbour of v, v_l, v_r in U(G).

Proof. On contrary let v_k be the common neighbour of v, v_l, v_r in U(G). We first consider the case when v_k is an inneighbour of v. Now if $vv_l, vv_r \in E$ then from Lemma 7.7 there must exist an edge between v_l, v_r in U(G), which contradicts our assumption. Again if $v_lv, v_rv \in E$ then D_3 or D_7 or D_8 get induced in G, which is not true.

Next we consider the case when v_k is an out-neighbour of v. Now if $vv_l, vv_r \in E$ then D_6 or D_7 or D_9 gets formed by the vertices $\{v, v_l, v_k, v_r\}$. Again if $v_lv, v_rv \in E$ then from Lemma 7.7 it follows that v_l, v_r become adjacent in U(G). Hence in both of these cases contradiction arises. Thus the result holds.

Corollary 7.9. Let $v \in V$. Then $N_O[v](N_I[v])$ is either a tournament or is disjoint union of two tournaments in G.

Proof. Note that v can not possesses three nonadjacent out(in)-neighbours in G as D_5 (D_2) get induced in that case, which can not be true. Rest of the proof follows from Lemma 7.8.

Definition 7.10. A set A of vertices is said to form an *Out-Fountain* if there exists an ordering of vertices such that for each i < j, $v_i v_j \in E$. Similarly a set B of vertices is said to form an *In-Fountain* if there exists an ordering of vertices such that for each i > j, $v_i v_j \in E$. A set is said to be a *Fountain* if and only if it is either an *Out-Fountain* or an *In-Fountain*. Further we denote the first element by v_f and the last by v_l of any Fountain. Note that if we reverse the order of an Out-Fountain, it becomes an In-Fountain and vice versa. Moreover any Fountain must be a tournament.

Notation: By A we denote an Out-Fountain and B an In-Fountain. By $N_O(A \setminus \{v_f\})$ $(N_I(B \setminus \{v_f\}))$ we mean the set of all out(in)-neighbours of $A \setminus \{v_f\}$ $((B \setminus \{v_f\}))$ which lies outside A (B).

Lemma 7.11. If v_k be an out(in)-neighbour of some element (say v_i) of $A \setminus \{v_f\}$ ($B \setminus \{v_f\}$) then v_k is also an out(in)-neighbour of v_j for all $i \leq j \leq l$.

Proof. We assume on contrary that v_j , $i < j \le l$ be a vertex which does not have v_k as its out(in)-neighbour. Then the set $\{v_f, v_i, v_j, v_k\}$ induces D_3 or D_4 (D_4 or D_9), which can not happen.

Lemma 7.12. The vertices of $N_O(A \setminus \{v_f\})$ ($N_I(B \setminus \{v_f\})$) form a tournament.

Proof. Let $v_k, v_{k'} \in N_O(A \setminus \{v_f\})$ be two nonadjacent out-neighbours of v_i, v_j respectively where i < j and $v_i, v_j \in A$. Then from Lemma 7.11 it follows that both of them are out-neighbours of v_j . Then $\{v_f, v_j, v_k, v_{k'}\}$ induce D_5 . The proof for $N_I(B \setminus \{v_f\})$ follows similarly using the fact that D_2 is not an ICD.

Lemma 7.13. Let v_k be not an out(in)-neighbour of v_f . Then it cannot be both an in-neighbour and an out neighbour of A(B).

Proof. Let $v_i, v_j \in A$ where i < j. As oriented ICD is acyclic, $v_j v_k, v_k v_i \notin E$. Hence we assume on contrary that $v_i v_k, v_k v_j \in E$. Then by lemma 7.11, $v_j v_k \in E$ but this contradicts the fact that G is oriented. Similarly one can prove for the case of B when v_k is not an inneighbour of v_f .

Lemma 7.14. There is an ordering such that vertices of $A \cup N_O(A \setminus \{v_f\})(B \cup N_I(B \setminus \{v_f\}))$ satisfies (7.4). Moreover there exists a subset of the above set which forms an Out-Fountain (In-Fountain) under the induced order.

Proof. We arrange the vertices of $N_O(A \setminus \{v_f\})$ in decreasing order of their out-degrees restricted on the set. We place the elements of $N_O(A \setminus \{v_f\})$ to the right of v_l . Let \prec be the combined order on the set $A \cup N_O(A \setminus \{v_f\})$. Let v_1, v_2, v_3 be any three vertices. If all of the three lie in A then from Definition 7.10 they satisfy (7.4). Again if all of them come from $N_O(A \setminus \{v_f\})$, using Lemma 7.12 one can verify that they satisfy (7.4). In view of Lemma 7.11 this is also done if two of them lies in A. Now if $v_1 \in A$, $v_2, v_3 \in N_O(A \setminus \{v_f\})$ and $v_2 \prec v_3$, then let $v_1v_3 \in E$. Clearly $v_2v_3 \in E$ (from definition of \prec). It is sufficient to prove now $v_1v_2 \in E$. Let v_j be the first inneighbour of v_2 in A. Now if $v_1 \prec v_j$ in A, then $v_jv_3 \in E$ from Lemma 7.11 as $v_1v_3 \in E$. Moreover as $v_jv_2 \in E$, $v_2v_1 \notin E$ from Lemma 7.13. Therefore $\{v_1, v_j, v_2, v_3\}$ induce D_3 in G which is a contradiction. Hence $v_j \prec v_1$ and therefore from Lemma 7.11 we get $v_1v_2 \in E$.

Again in view of Lemma 7.11 there exists a vertex $v_i \in A$ such that each element in $N_O(A \setminus \{v_f\})$ is an outneighbour of v_i and hence for any $i \leq j \leq l$. Therefore in the given ordering \prec the set $\{v_i, v_{i+1}, \dots, v_l\} \cup N_O(A \setminus \{v_f\})$ forms an Out-Fountain due to Lemma 7.12.

Similarly one can prove $B \cup N_I(B \setminus \{v_f\})$ satisfies (7.4) if we arrange the vertices of $N_I(B \setminus \{v_f\})$ in decreasing order of their in-degrees restricted on the set. Moreover there exists a set $B' \subsetneq B$ for which $B' \cup N_I(B \setminus \{v_f\})$ form an In-Fountain.

Lemma 7.15. Let $v_k \notin A$ and $v_k v_i \in E$ for some $f \leq i < l$. If $v_k v_l \notin E$ then $v_k v_j \in E$ for all $f \leq j \leq i$ and $v_k v_j \in E$ for all $f \leq j \leq l$ if $v_k v_l \in E$ but $v_f v_k \notin E$.

(Let $v_k \notin B$ and $v_i v_k \in E$ for some $f \leq i < l$. If $v_l v_k \notin E$ then $v_j v_k \in E$ for all $f \leq j \leq i$ and $v_j v_k \in E$ for all $f \leq j \leq l$ if $v_l v_k \in E$ but $v_k v_f \notin E$)

Proof. As G is acyclic, $v_l v_k \notin E$. Again if $v_k v_l \notin E$ then we reverse the vertex ordering of A, then by definition A becomes an In-Fountain. Therefore by Lemma 7.11 the result follows. Now if $v_k v_l \in E$, then as $v_k v_i \in E$ and $v_f v_k \notin E$ using Lemma 7.8 for the vertices $\{v_f, v_l, v_i, v_k\}$ and Corollary 7.9 for $N_I[v_l]$, we

can conclude that $v_k v_f \in E$. Again as $v_k v_l \in E$, from Lemma 7.13 we get $v_k v_j \in E$ for all $f \leq j \leq l$ as G is chordal. The proof for the case of B is similar.

Definition 7.16. Let $A_0 = A$ be an Out-Fountain. We take $A_1 = A \cup N_O(A \setminus \{v_f\})$. For i > 1 let $A_{i+1} = A_i \cup N_O(A_i \setminus \{v_f\})$. Then clearly, A_i forms an increasing chain of subsets of V. As V is a finite set, this chain must be stable. So there is an integer N such that $A_N = A_{N+1}$. We call the set A_N as derived set of out-neighbours of A. Notationally we define it by Der(A). Analogously we can define derived set of in-neighbours (Der(B)) for an In-Fountain B.

Properties of Derived set:

- 1. No vertex of Der(A) except v_f has an out-neighbour outside Der(A) and no vertex of Der(B) except v_f has an in-neighbour outside Der(B).
- 2. There exists an ordering on Der(A) (Der(B)) satisfying (7.4) and it induces the natural order in A.

Proof. Follows from repeated application of Lemma 7.14.

As Der(A) is a ordered set from above property, by *boundary points* of Der(A) we mean the two extreme vertices of the set in the induced order. By *interior point* of Der(A) we mean any vertex of the set except the boundary points. For Der(B) these definitions are analogous.

3. If $v_k \notin Der(A)$ is an in-neighbour of some interior point of Der(A). Then it must also be an inneighbour of v_f . (If $v_k \notin Der(B)$ is an out-neighbour of some interior point of Der(B). Then it must also be an out-neighbour of v_f .)

Proof. We assume on contrary that $v_k v_i \in E$ but $v_k v_f \notin E$ where v_i is the first outneighbour of v_k in Der(A). Now if $v_i \in A \setminus \{v_l\}$ then from property (1) it is clear that $v_l v_k \notin E$. Hence applying Lemma 7.15 on vertices of A it follows that $v_k v_f \in E$, which contradicts our assumption.

Again if $v_i \notin A \setminus \{v_l\}$ then let $v_i \in A_{t+1} \setminus A_t$ for some t implies $v_i \in N_O[A_t]$ (say $N_O[v_j]$), then $v_j \prec v_i$ by our ordering. Also v_k is not an outneighbour of v_j from property (1). Hence if v_i is an interior point of $N_O[v_j]$ then from Lemma 7.15 we get $v_k v_j \in E$, which is not true from the choice of i. Next if v_i be the last end point of $N_O[v_j]$. Then we consider the sets $N_O[v_j]$ and $N_O[v_i]$. Again one can show that v_j, v_k are not adjacent in U(G). Hence D_4 or D_2 get induced by the vertices $\{v_j, v_i, v_k, v_{i_2}\}$ where v_{i_2} is the last vertex of $N_O[v_i]$, which is a contradiction. Thus the result holds. Similar proof will follow for the case of Der(B).

Lemma 7.17. Let $v \in V$ such that either v has no in(out)-neighbour and $N_O[v]$ $(N_I[v])$ is a single out(in)-neighbour tournament. Then there exists an ordering $(say \prec)$ on V satisfying (7.4) having v as the first element in \prec .

Proof. We proceed by induction on |V|. Clearly done for |V| = 1. If |V| > 1 then consider $A = N_O[v]$. Let $V' = (V \setminus Der(A)) \cup \{v_{l'}\}$ where $v_{l'}$ is the last element of Der(A). Then |V'| < |V|. Also $v_{l'}$ satisfies no out neighbour from property (1) and it can induce maximum one inneighbour tournament (say $R_{v_{l'}}$) whose vertices of are disjoint from Der(A) (follows from Corollary 7.9, property 1, 3 and the fact v has no inneighbour). Hence by induction there exists an ordering of V' satisfying (7.4) and let that be $v_{l'} \prec v_{l'+1} \prec \ldots \prec v_n$. Also Der(A) is an ordered set satisfying (7.4) follows from property 2 and hence $v = v_1 \prec v_2 \prec \ldots \prec v_{l'}$. We now order the elements of V as $v_1 \prec v_2 \prec \ldots \lor v_{l'+1} \prec \ldots \prec v_n$. This clearly satisfies (7.4). For the case of inneighbours we consider Der(B) where $B = N_I[v]$ and proceed similarly. \Box

Construction of \hat{P}

There are three cases which may occur while constructing \hat{P} . Below we list all of them. As v_1 is the first element of P one can easily check that $N_I[v_1]$ must be a single tournament. Further if $N_O[v_1] = A \cup B$ where A, B are disjoint outneighbour tournaments of v_1 , then v_1 can not have any inneighbour in G. On contrary if $v_k v_1 \in E$ then from Lemma 7.8 it follows that v_k can not be simultaneously adjacent to A and B in U(G). In other cases D_4 or D_5 get induced in G, which again introduce contradiction.

1. $N_I[v_1]$ is a single tournament and v_1 has no outneighbour.

Proof. Construction of \hat{P} follows from Lemma 7.17.

2. $N_O[v_1]$ is disjoint union of two tournaments and v_1 has no inneighbour.

Proof. Let $N_O[v_1] = A \cup B$. If we arrange the vertices of each sets individually according to decreasing outdegree restricted on the sets then both of them become Out-Fountains. Next we consider consider Der(A). If |Der(A)| > |A| then Der(B) = B follows from the fact that all outneighbours of v_1 occur consecutively in P right to v_1 . As $A \cap B = \emptyset$ and they are tournaments, one can easily verify that each vertex of B must occur prior every vertex of A in P. Therefore if any vertex (say v_b) of B possess inneighbour (say v_k) outside $V \setminus (A \cup B)$, then it must be an inneighbour of A (say v_a). Therefore $\{v_1, v_a, v_b, v_k\}$ form a 4-cycle in U(G). But as v_a, v_b are nonadjacent and G is chordal, $v_k v_1 \in E$. But this contradicts our assumption. Therefore using property 1 we can conclude now that no vertex of $B \setminus \{v_1\}$ have any neighbour outside the set B.

We place Der(A) right to v_1 in \hat{P} and place the vertices of B in reverse order left to v_1 so that it become an In-Fountain. Now applying Lemma 7.17 one can order the vertices of $(V \setminus Der(B)) \cup \{v_1\}$ so that they satisfy (7.4). Again as Der(B) satisfies (7.4) from property 2 it is easy to check now that \hat{P} satisfies (7.4). Now if Der(A) = A and Der(B) = B then the proof is trivial.

3. $N_O[v_1]$ is a single tournament.

If v_1 does not contain any inneighbour then \hat{P} is formed by Lemma 7.17.

Again if v_1 has an inneighbour (say v_k) then v_l can not have any outneighbour where v_l is the last element of the Out-Fountain $N_O[v_1]$. More specifically $Der(A) = N_O[v_1] = A$.

Note that v_k can not be an outneighbour of v_l as G is acyclic. On contrary let v_i be an outneighbour of v_l . Now in P as vertices of $N_O[v_1]$ occur consecutively right to $v_1, v_i, v_k > v_l$ in P. Hence $v_k v_l \in E$. Now if $v_k < v_i$ in P then $v_l v_k \in E$ which contradicts that G is oriented. Again if $v_k > v_i$ then $\{v_1, v_l, v_k, v_i\}$ induce D_9 which is a contradiction.

Now we will show $N_I[v_l]$ is a single tournament when $v_k v_1 \in E$.

Again if on contrary we assume v_l possesses an inneighbour v_i which is disjoint from v_1 , then $v_i \notin N_O[v_1]$ clearly. Now if $v_i < v_k$ in P, then the vertices $\{v_1, v_l, v_k, v_i\}$ induce D_7 , again if $v_k < v_i$, then they form D_3 , which are both contradiction. Hence the result follows.

We can now apply Lemma 7.17 on the In-Fountain $B = N_I[v_l]$ (say). Thus we get an ordering \dot{P} on V satisfying (7.4) having v_l as the first element.




Figure 9: oriented ICD but not proper oriented ICD (undirected edge can have any direction)



Figure 10: \hat{D}

8 Conclusion


It was proved in [18] that interval graphs \subset central MPTG \subseteq max-tolerance graph. Combining with these we establish the relations between some subclasses of max-tolerance graphs related to central MPTG in Figure 11.

Note that $C_4 \in$ unit max-tolerance \smallsetminus interval graphs. C_4 has a unit-max-tolerance representation having intervals [1,5], [2,6], [3,7], [4,8] and corresponding tolerances 1,3,3,1 for its consecutive vertices (clockwise or anticlockwise). But C_4 is not an interval graph. In Theorem 3.9 we have shown that $\overline{C_6}$ is a max-tolerance graph which is not a central MPTG. Also we note that $K_{2,3}$ is not a central MPTG (by Lemma 7 of [18]) but it is a MPTG (by Lemma 8 of [18]). Again $K_{1,3} \in 50\%$ max-tolerance graph \backsim proper interval graph. $K_{1,3}$ is an example of a graph which has a 50% max-tolerance representation having interval [1.9, 6.1] and tolerance 2.1 for its central vertex and the intervals [0, 8], [1.8, 4.3], [3.6, 5.9] and corresponding tolerances 4, 1.25, 1.15 for pendant vertices. But it is not a proper interval graph.

The following examples lead us to conclude that interval graphs and 50% max-tolerance graphs are incomparable.

 $C_4 \in 50\%$ max-tolerance graph $\$ interval graph. C_4 is an example of a graph which has 50% max-tolerance representation having intervals [1, 4.6], [2, 4], [2.9, 4.9], [2.7, 6.3] and corresponding tolerances 1.8, 1, 1, 1.8 for its consecutive vertices (clockwise or anticlockwise). But it is not an interval graph. Again $K_{1,n}$, $n \ge 8 \in$ interval Graphs $\$ 50% max-tolerance graph. $K_{1,n}$ is an interval graph having interval [1, 2n] for central vertex and intervals [2i - 1, 2i] for pendant vertices where $i \in \{1, \ldots, n\}$. But it has no 50% max-tolerance representation follows from Lemma 4.3.

Again from Theorem 4.2 we get to know that central MPTG and 50% max-tolerance graphs are not comparable. One can also conclude from above that $K_{1,n}$, $n \ge 8 \in$ max-tolerance graph. But it is not a 50% max-tolerance graph from Theorem 4.3.



proper interval graph=proper central MPTG=unit central MPTG

Figure 11: Hierarchy of subclasses of the class of max-tolerance graph

In this paper we consider three subclasses of the class of ICD, namely central ICD, oriented ICD and proper ICD. We obtain characterizations for central ICD, oriented ICD when it is a tournament and proper ICD. Also in terms of forbidden subdigraphs we characterize those oriented ICD for which their underlying graph is tree and those proper oriented ICD for which their underlying graph is chordal. In Figure 12 we show relationships between the digraphs discussed in this paper whose proofs follow from the mentioned characterization theorems. Several combinatorial optimization problems remain open for these digraphs, for example, to find the complete list of forbidden digraphs or to construct recognition algorithms for them. Finally we note the major unsolved problems in this area.

- 1. Recognition algorithm and forbidden subgraph characterization of max-point tolerance graphs.
- 2. Combinatorial characterization, adjacency matrix characterization, recognition algorithm and forbidden subgraph characterization of central max-point tolerance graphs and 50% max-tolerance graphs.
- 3. Find the complete list of forbidden digraphs of central ICD, proper ICD.




Figure 12: Relations between some subclasses of ICD

Acknowledgements:

This research is supported by UGC (University Grants Commission) NET fellowship (21/12/2014(ii)EU-V) of the first author.

References

- [1] Jorgen Bang-Jensen, Gregory Z.Gutin , Digraphs, SPRINGER MONOGRAPHS IN MATHEMATICS, 2008.
- [2] Asim Basu, Sandip Das, Shamik Ghosh and Malay Sen, Circular-arc bigraphs and its subclasses, J. Graph Theory 73 (2013), 361–376.
- [3] S. Benzer, On the topology of the genetic fine structure, Proc. Nat. Acad. Sci. USA 45 (1959), 1607–1620.
- [4] Kenneth P. Bogart, Peter C. Fishburn, Garth Isaak and Larry Langley, Proper and unit tolerance graphs, Discrete Applied Mathematics 60 (1995), 99–117.
- [5] Kenneth P. Bogart and Douglas B. West, A Short Proof that Proper = Unit, Discrete Mathematics 201 (1999), 21–23.
- [6] Daniele Catanzaroa, Steven Chaplick, Stefan Felsner, Bjarni V. Halldórssonc, Magnús M. Halldórssond, Thomas Hixonb and Juraj Stachoe, Max Point-Tolerance Graphs, Discrete Applied Mathematics 216 (2017), 84-97.
- [7] M. C. Golumbic, Algorithmic graph theory and perfect graphs, Annals of Disc. Math., 57, Elsevier Sci., USA, 2004.
- [8] M. C. Golumbic and C. L. Monma, A generalization of interval graphs with tolerances, Congress. Numer., 153 (1982), 97–106.
- [9] M. C. Golumbic and A. Trenk, Tolerance Graphs. Cambridge Studies in Advanced Mathematics 89, Cambridge University Press, 2004.
- [10] Shamik Ghosh, Maitry Podder, Malay K. Sen, Adjacency matrices of probe interval graphs, Discrete Applied Mathematics 158 (2010), 2004-2013.
- [11] G. Hajös, Über eine Art von Graphen. Internationale Mathematische Nachrichten 11 (1957), Problem 65.
- [12] Michael Kaufmann, Jan Kratochvil, Katharina A. Lehmann and Amarendran R. Subramanian, Max-Tolerance Graphs as Intersection Graphs: Cliques, Cycles, and Recognition SODA 2006, January 22-26, Miami, FL.
- [13] P. Looges, S. Olariu, Optimal greedy algorithms for indifference graphs, Comput. Math. Appl. 25 (1993) 15-25.
- [14] D. S. Malik, M. K. Sen and S. Ghosh, Introduction to Graph Theory, Cengage Learning Asia Pte Ltd, Singapore, 2014.
- [15] Hiroshi Maehara, A Digraph Represented by a Family of Boxes or Spheres, J. Graph Theory, 8 (1984), 431–439.
- [16] Erich Prisner, A Characterization of Interval Catch Digraphs, Discrete Mathematics 73 (1989), 285-289.
- [17] M. Sen, S. Das, A.B. Roy, D.B. West, Interval Digraphs: An Analogue of Interval Graphs, Journal of Graph Theory, Vol. 13, No. 2, (1989) 189-202
- [18] Maucorico Soto and Christopher Thraves Caro, p- Box: A new graph model, Journal of Discrete Mathematics and Theoretical Computer Science, DMTCS 17 (2015), 169–186.