

# ALMA Observations of a High-density Core, MC27/L1521F in Taurus: Dynamical Gas Interaction at the Possible Site of a Multiple Star Formation

Kazuki Tokuda

TOKUDA@P.S.OSAKAFU-U.AC.JP

Department of Physical Science, Graduate School of Science, Osaka Prefecture University, 1-1 Gakuen-cho, Naka-ku, Sakai, Osaka 599-8531, Japan

## Abstract

We present the results of ALMA observations of dust continuum emission and molecular rotational lines toward a dense core, MC27 (aka L1521F), which is considered to be very close to the first core phase. We revealed the spatial/velocity structures of the core are very complex and suggest that the initial condition of star formation is highly dynamical.

## 1. ALMA observations

Our ALMA observations reveal complex structures at the center of MC27/L1521F (Tokuda *et al.* 2014). We find a few starless high-density cores, one of which (MMS-2) has a very high number density of  $\sim 10^7 \text{ cm}^{-3}$ . The high angular resolution observations at 0.87 mm with a beam size of  $\sim 0.''74 \times 0.''32$  reveal that the MMS-2 has substructures in both dust and molecular emission (Fig. 1 lower panel). A very compact bipolar outflow with a dynamical timescale of a few hundred years was found toward the protostar. The molecular line observations show several cores are associated with an arc-like structure whose length is  $\sim 2000 \text{ AU}$ , possibly formed due to the dynamical gas interaction (Matsumoto *et al.* 2015). The observed column density structure ranges in scale from  $\sim 100$  to  $\sim 10,000 \text{ AU}$  as revealed by combining the 12 m array data with the 7 m array data as well as with the single dish MAMBO data (Fig. 1, see also Tokuda *et al.* 2016). The azimuthally averaged radial column density distribution of the inner part ( $r < 3000 \text{ AU}$ ) is  $N_{\text{H}_2} \sim r^{-0.4}$ , clearly flatter than that of the outer part,  $\sim r^{-1.0}$ . We detect the above-mentioned complex structures, such as the arc-like structures, inside the inner flatter region, which may reflect the dynamical status of the dense core (see also discussions in Tokuda *et al.* 2016).

## References

*Proceedings of the Star Formation in Different Environments*, ICISE, Quy Nhon, Vietnam, 2016. PSFDE: volume 01. Copyright 2016 by the author(s).

Matsumoto, T., *et al.* MNRAS, 449, L123 (2015)  
Tokuda, K., *et al.* ApJL, 789, L4 (2014).  
Tokuda, K., *et al.* ApJ, 826, 26 (2016).




Figure 1. Distribution of dust continuum and molecular line emission toward MC27/L1521F (Tokuda *et al.* 2016). Black cross in each panel represents the position of the protostar identified with *Spitzer* observations.