

# CUBIC LAGRANGE ELEMENTS SATISFYING EXACT INCOMPRESSIBILITY

JOHNNY GUZMÁN<sup>†</sup> AND L. RIDGWAY SCOTT<sup>‡</sup>

ABSTRACT. We prove that an analog of the Scott-Vogelius finite elements are inf-sup stable on certain nondegenerate meshes for piecewise cubic velocity fields. We also characterize the divergence of the velocity space on such meshes. In addition, we show how such a characterization relates to the dimension of  $C^1$  piecewise quartics on the same mesh.

[2010]65N30, 65N12, 76D07, 65N85

## 1. INTRODUCTION

In 1985 Scott and Vogelius [13] (see also [16]) presented a family of piecewise polynomial spaces in two dimensions that yield solutions to the Stokes equations with velocity approximations that are exactly divergence free. The velocity space consists of continuous piecewise polynomials of degree  $k \geq 4$ , and the pressure space is taken to be the divergence of the velocity space. Moreover, they proved stability of the method by establishing that the pair of spaces satisfy the so-called *inf-sup* condition assuming that the meshes are quasi-uniform and that the maximum mesh size is sufficiently small. In a recent paper [7] we gave an alternative proof of the inf-sup stability for  $k \geq 4$  on more general meshes, assuming only that they are non-degenerate (shape regular). One key aspect in the proof is to use the stability of the  $P^2 - P^0$  (or the Bernardi-Raugel [4]) finite element spaces. As a result the proof becomes significantly shorter. Here we utilize and extend the techniques to the case  $k = 3$ . The case  $k = 3$  has been considered earlier [11].

One key concept in this paper is the notion of a *local interpolating vertex*. Roughly speaking, this is an interior vertex  $z$  such that for every finite element pressure  $p$  we can find a discrete velocity  $\mathbf{v}$  in the finite element velocity space such that  $\operatorname{div} \mathbf{v}(z) = p(z)$  with support in the patch of  $z$  and such that  $\operatorname{div} \mathbf{v}(\sigma) = 0$  for all other vertices. Moreover, we require that  $\operatorname{div} \mathbf{v}$  has zero mean on each triangle. We then show that if all interior vertices are local interpolating vertices then the inf-sup stability holds; see Theorem 1. We generalize this result to show that if some interior vertices are local interpolating and that there are acceptable paths from any other vertex to one of the local interpolating vertices then the inf-sup condition holds; see Theorem 2. In [7] we showed that all interior vertices are local interpolating vertices if piecewise quartic velocities or higher are used. In this article, we show that a generic interior vertex is local interpolating if piecewise cubics are used for the velocity space. In particular, we show that singular vertices and vertices with odd number of triangles touching it are local interpolating vertices. In the case that a non-singular interior vertex has an even number of triangles touching it then we give sufficient conditions for it to be a local interpolating vertex. Although a generic

---

*Date:* April 1, 2024.

interior vertex is a local interpolating vertex, there are important meshes where no interior vertex is locally interpolating (e.g. a diagonal mesh).

It is known that the  $C^1$  piecewise quartic space which we denote by  $\widehat{S}_h^4$  is related to the piecewise cubic Lagrange space. Surprisingly, the dimension of  $\widehat{S}_h^4$  has not been verified for all meshes, but it has been verified for a large class of meshes; see [10, 1]. In the last sections of this paper we use the onto-ness of the divergence operator of piecewise cubics to verify the dimension of  $\widehat{S}_h^4$  for a large class of meshes. We also are able to verify the dimension of  $S_h^4$  which are  $C^1$  piecewise quartic space with functions vanishing on the boundary to second-order.

The paper is organized as follows. In the following section we begin with Preliminaries. In Section 3 we identify vertices at which we can interpolate pressure vertex values with the divergence of localized velocity fields. In Section 6 we prove the inf-sup stability for  $k = 3$  under some restrictive assumptions on the mesh. In Section 7 we characterize the divergence of piecewise cubics on a broader class of meshes than considered in Section 6. In Section 8 we compare our results with those of [11]. In Section 9 we relate our results to the dimension of  $C^1$  piecewise quartics.

## 2. PRELIMINARIES

We assume  $\Omega$  is a polygonal domain in two dimensions. We let  $\{\mathcal{T}_h\}_h$  be a non-degenerate (shape regular) family triangulation of  $\Omega$ ; see [5]. The set of vertices and the set of internal edges are denoted by

$$\begin{aligned}\mathcal{S}_h &= \{x : x \text{ is a vertex of } \mathcal{T}_h\}, \\ \mathcal{E}_h &= \{e : e \text{ is an edge of } \mathcal{T}_h \text{ and } e \not\subset \partial\Omega\}.\end{aligned}$$

We also let  $\mathcal{S}_h^{\text{int}}$  and  $\mathcal{S}_h^{\partial}$  denote all the interior vertices and boundary vertices, respectively.

Define the internal edges  $\mathcal{E}_h(z)$  and triangles  $\mathcal{T}_h(z)$  that have  $z \in \mathcal{S}_h$  as a vertex via

$$\mathcal{E}_h(z) = \{e \in \mathcal{E}_h : z \text{ is a vertex of } e\}, \quad \mathcal{T}_h(z) = \{T \in \mathcal{T}_h : z \text{ is a vertex of } T\}.$$

Finally, we define the patch

$$\Omega_h(z) = \bigcup_{T \in \mathcal{T}_h(z)} T.$$


The diameter of this patch is denoted by

$$(2.1) \quad h_z = \text{diam}(\Omega_h(z)).$$

To define the pressure space we must define singular and non-singular vertices. Let  $z \in \mathcal{S}_h$  and suppose that  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$ , enumerated so that  $T_j, T_{j+1}$  share an edge for  $j = 1, \dots, N-1$  and  $T_N$  and  $T_1$  share an edge in the case  $z$  is an interior vertex. If  $z$  is a boundary vertex then we enumerate the triangles such that  $T_1$  and  $T_N$  have a boundary edge. Let  $\theta_j$  denote the angle between the edges of  $T_j$  originating from  $z$ . Define

$$\Theta(z) = \begin{cases} \max\{|\sin(\theta_1 + \theta_2)|, \dots, |\sin(\theta_{N-1} + \theta_N)|, |\sin(\theta_N + \theta_1)|\} & \text{if } z \text{ is an interior vertex} \\ \max\{|\sin(\theta_1 + \theta_2)|, \dots, |\sin(\theta_{N-1} + \theta_N)|\} & \text{if } z \text{ is a boundary vertex} . \end{cases}$$

**Definition 2.1.** A vertex  $z \in \mathcal{S}_h$  is a singular vertex if  $\Theta(z) = 0$ . It is non-singular if  $\Theta(z) > 0$ .



**Figure 1.** Example of singular vertices  $z$ . Dashed edges denote boundary edges.

This is equivalent to the original definition given in [9, 16].

We denote all the non-singular vertices by

$$\mathcal{S}_h^1 = \{x \in \mathcal{S}_h : x \text{ is non-singular, that is, } \Theta(x) > 0\},$$

and all singular vertices by  $\mathcal{S}_h^2 = \mathcal{S}_h \setminus \mathcal{S}_h^1$ . We also define, for  $i = 1, 2$ ,  $\mathcal{S}_h^{i,\text{int}} = \mathcal{S}_h^i \cap \mathcal{S}_h^{\text{int}}$ , and  $\mathcal{S}_h^{i,\partial} = \mathcal{S}_h^i \cap \mathcal{S}_h^\partial$ .

Let  $q$  be a function such that  $q|_T \in C(\bar{T})$  for all  $T \in \mathcal{T}_h$ . For each vertex  $z \in \mathcal{S}_h^2$  define

$$(2.2) \quad A_h^z(q) = \sum_{j=1}^N (-1)^{j-1} q|_{T_j}(z).$$

The Scott-Vogelius finite element spaces for  $k \geq 1$  are defined by

$$\begin{aligned} \mathbf{V}_h^k &= \{v \in [C_0(\Omega)]^2 : v|_T \in [P^k(T)]^2, \forall T \in \mathcal{T}_h\}, \\ Q_h^{k-1} &= \{q \in L_0^2(\Omega) : q|_T \in P^{k-1}(T), \forall T \in \mathcal{T}_h, A_h^z(q) = 0 \text{ for all } z \in \mathcal{S}_h^2\}. \end{aligned}$$

Here  $P^k(T)$  is the space of polynomials of degree less than or equal to  $k$  defined on  $T$ . Also,  $L_0^2(\Omega)$  denotes the subspace of  $L^2$  of functions that have average zero on  $\Omega$ .

We also make the following definition

$$\mathbf{V}_{h,0}^k = \{\mathbf{v} \in \mathbf{V}_h^k : \int_T \operatorname{div} \mathbf{v} \, dx = 0, \quad \text{for all } T \in \mathcal{T}_h\}.$$

The definition of  $Q_h^{k-1}$  is based on the following result [14].

**Lemma 1.** For  $k \geq 1$ ,  $\operatorname{div} \mathbf{V}_h^k \subset Q_h^{k-1}$ .


The goal of this article is to prove the inf-sup stability of the pair  $\mathbf{V}_h^k, Q_h^{k-1}$  for  $k = 3$ , for certain meshes.

**Definition 2.2.** The pair of spaces  $\mathbf{V}_h^k, Q_h^{k-1}$  is inf-sup stable on a family of triangulations  $\{\mathcal{T}_h\}_h$  if there exists  $\beta > 0$  such that for all  $h$

$$(2.3) \quad \beta \|q\|_{L^2(\Omega)} \leq \sup_{\mathbf{v} \in \mathbf{V}_h^k, \mathbf{v} \neq 0} \frac{\int_\Omega q \operatorname{div} \mathbf{v} \, dx}{\|\mathbf{v}\|_{H^1(\Omega)}} \quad \forall q \in Q_h^{k-1}.$$

**2.1. Notation for piecewise linears.** For every  $z \in \mathcal{S}_h$  the function  $\psi_z$  is the continuous, piecewise linear function corresponding to the vertex  $z$ . That is, for every  $y \in \mathcal{S}_h$

$$(2.4) \quad \psi_z(y) = \begin{cases} 1 & \text{if } y = z, \\ 0 & \text{if } y \neq z. \end{cases}$$



**Figure 2.** Notation for single triangle

For  $z \in \mathcal{S}_h$  and  $e \in \mathcal{E}_h(z)$ , where  $e = \{z, y\}$ , let  $\mathbf{t}_e^z = |e|^{-1}(y - z)$  be a unit vector tangent to  $e$ , where  $|e|$  denotes the length of the edge  $e$ . Then

$$(2.5) \quad \mathbf{t}_e^z \cdot \nabla \psi_y = \frac{1}{|e|} \text{ on } e.$$

More generally, suppose that  $T \in \mathcal{T}_h(y)$  and let  $g$  be the edge of  $T$  that is opposite to  $y$ ; see Figure 2. Let  $\mathbf{n}_T^y$  be the unit normal vector to  $g$  that points out of  $T$ . Then

$$(2.6) \quad \nabla \psi_y|_T = -\frac{1}{h_T^y} \mathbf{n}_T^y,$$

where  $h_T^y$  is the distance of  $y$  to the line defined by the edge  $g$ . If  $z$  is another vertex of  $T$  and  $e = \{z, y\}$  then

$$(2.7) \quad h_T^y = |e| \sin \theta,$$

where  $\theta$  is the angle between the edges of  $T$  emanating from  $z$ .

**2.2. Preliminary stability results.** The following is a consequence of the stability of the Bernardi-Raugel [4] finite element spaces.

**Proposition 1.** *Let  $k \geq 1$ . There exists a constant  $\alpha_1$  such that for every  $p \in Q_h^{k-1}$  there exists a  $\mathbf{v} \in \mathbf{V}_h^2$  such that*

$$\int_T \operatorname{div} \mathbf{v} \, dx = \int_T p \, dx \quad \text{for all } T \in \mathcal{T}_h,$$

and

$$\|\mathbf{v}\|_{H^1(\Omega)} \leq \alpha_1 \|p\|_{L^2(\Omega)}.$$

The constant  $\alpha_1$  is independent of  $p$  and only depends on the shape regularity of the mesh and  $\Omega$ .

The next result is a simple consequence of Lemma 2.5 of [16] and is based on a simple counting argument.

**Proposition 2.** *Let  $k \geq 1$ . There exists a constant  $\alpha_2 > 0$  such that for every  $p \in Q_h^{k-1}$  with  $p(z) = 0$  for all  $z \in \mathcal{S}_h$  and  $\int_T p dx = 0$  for all  $T \in \mathcal{T}_h$  there exists  $\mathbf{v} \in \mathbf{V}_h^k$  such that*

$$\operatorname{div} \mathbf{v} = p \text{ on } \Omega,$$

and

$$\|\mathbf{v}\|_{H^1(\Omega)} \leq \alpha_2 \|p\|_{L^2(\Omega)}.$$

Using the above results we can prove inf-sup stability as long as we can interpolate pressure vertex values with the divergence of velocity fields. This is the subject of the next result.

**Lemma 2.** *Suppose there exists a constant  $\alpha_3 > 0$  such that for every  $p \in Q_h^2$  there exists a  $\mathbf{v} \in \mathbf{V}_{h,0}^3$  satisfying*

$$(2.8) \quad (\operatorname{div} \mathbf{v} - p)(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h,$$

with the bound

$$\|\mathbf{v}\|_{H^1(\Omega)} \leq \alpha_3 \|p\|_{L^2(\Omega)}.$$

Then, (2.3) holds for  $k = 3$  with  $\beta = \frac{1}{\alpha_1 + \alpha_3(1 + \alpha_1) + \alpha_2(1 + \alpha_3(1 + \alpha_1))}$ .

*Proof.* Let  $p$  be an arbitrary function in  $Q_h^2$ . First, we let  $\mathbf{v}_1$  be from Proposition 1. We define  $p_1 = p - \operatorname{div} \mathbf{v}_1$ . Then, from our hypothesis let  $\mathbf{v}_2 \in \mathbf{V}_{h,0}^3$  be such that  $(\operatorname{div}(\mathbf{v}_2) - p_1)(\sigma) = 0$  for all  $\sigma \in \mathcal{S}_h$ . Letting  $p_2 = p_1 - \operatorname{div} \mathbf{v}_2$  we see that  $p_2$  satisfies the hypotheses of Proposition 2 and we let  $\mathbf{v}_3$  be the resulting vector field. Then, we set  $\mathbf{w} = \mathbf{v}_1 + \mathbf{v}_2 + \mathbf{v}_3$ . Then we see that

$$\operatorname{div} \mathbf{w} = p \quad \text{on } \Omega,$$

and

$$(2.9) \quad \|\mathbf{w}\|_{H^1(\Omega)} \leq (\alpha_1 + \alpha_3(1 + \alpha_1) + \alpha_2(1 + \alpha_3(1 + \alpha_1))) \|p\|_{L^2(\Omega)}.$$

Hence,

$$\|p\|_{L^2(\Omega)}^2 = \int_{\Omega} p \operatorname{div} \mathbf{w} dx \leq \|\mathbf{w}\|_{H^1(\Omega)} \sup_{\mathbf{v} \in \mathbf{V}_h^3, \mathbf{v} \neq 0} \frac{\int_{\Omega} p \operatorname{div} \mathbf{v} dx}{\|\mathbf{v}\|_{H^1(\Omega)}}.$$

The result now follows after applying (2.9).  $\square$

Hence, one way of proving inf-sup stability is to establish the hypothesis of Lemma 2. This is going to be the task of the next sections.

### 3. LOCALLY INTERPOLATING VERTEX VALUES

In this section we will identify vertices at which we can interpolate pressure vertex values with the divergence of velocity fields. We first define the local spaces

$$\begin{aligned}\mathbf{V}_h^k(z) &= \{\mathbf{v} \in \mathbf{V}_h^k : \text{supp } \mathbf{v} \subset \Omega_h(z)\}, \\ \mathbf{V}_{h,0}^k(z) &= \{v \in \mathbf{V}_h^k(z) : \int_T \text{div } v \, dx = 0, \text{ for all } T \in \mathcal{T}_h(z)\}, \\ \mathbf{V}_{h,00}^k(z) &= \{v \in \mathbf{V}_{h,0}^k(z) : \text{div } v(\sigma) = 0, \text{ for all } \sigma \in \mathcal{S}_h, \sigma \neq z\}.\end{aligned}$$

Suppose that  $z \in \mathcal{S}_h$  and that  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$ , ordered as described following (2.1). Then in view of (2.2) we define

$$W(z) = \{(a_1, \dots, a_N) \in \mathbb{R}^N : \text{if } z \in \mathcal{S}_h^2, \text{ then } \sum_{j=1}^N (-1)^{j-1} a_j = 0\}.$$

Note that if  $z \in \mathcal{S}_h^1$  is non-singular then  $W(z) = \mathbb{R}^N$  and there is a constraint if  $z$  is singular.

**Definition 1.** Let  $z \in \mathcal{S}_h$  and suppose that  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$ . We say that  $z$  is a local interpolating vertex if for every  $(a_1, \dots, a_N) \in W(z)$  there exists a  $\mathbf{v} \in V_{h,00}^3(z)$  such that

$$(3.1) \quad \text{div } \mathbf{v}|_{T_j}(z) = a_j \quad \text{for all } 1 \leq j \leq N.$$

If  $z \in \mathcal{S}_h$  is a local interpolating vertex then, given  $a = (a_1, \dots, a_N) \in W(z)$  we define  $M_a = \{\mathbf{v} \in V_{h,00}^3(z) : \mathbf{v} \text{ satisfying (3.1)}\}$ . Also, we set

$$(3.2) \quad D_z = \max_{(a_1, \dots, a_N) \in W(z)} \min_{\mathbf{v} \in M_a} \frac{\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))}}{\max_{1 \leq j \leq N} |a_j|}.$$

We let  $\mathcal{L}_h$  be the collection of all local interpolating vertices. Then Definition 3.2 says that if  $z \in \mathcal{L}_h$  then given  $a \in W(z)$  there exists  $\mathbf{v} \in V_{h,00}^3(z)$  satisfying (3.1) and

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))} \leq D_z \max_{1 \leq j \leq N} |a_j|.$$

In the next section we identify local interpolating vertices. It will be useful to define fundamental vector fields used in [7]. For every  $z \in \mathcal{S}_h$  and  $e \in \mathcal{E}_h(z)$  with  $e = \{z, y\}$  define

$$(3.3) \quad \eta_e^z = \psi_z^2 \psi_y.$$

Let  $T_1$  and  $T_2$  be the two triangles that have  $e$  as an edge. Then we can easily verify the following (see also [7]):

$$(3.4a) \quad \text{supp } \eta_e^z \subset T_1 \cup T_2,$$

$$(3.4b) \quad \nabla \eta_e^z(\sigma) = 0 \quad \text{for } \sigma \in \mathcal{S}_h \text{ and } \sigma \neq z.$$

$$(3.4c) \quad \|\nabla \eta_e^z\|_{L^\infty(T_1 \cup T_2)} \leq \frac{C}{|e|}.$$

We then define the vector fields

$$(3.5) \quad \mathbf{w}_e^z = |e| \mathbf{t}_e^z \eta_e^z.$$

The following lemma is proved in [7].

**Lemma 3.** *Let  $z \in \mathcal{S}_h$  and  $e \in \mathcal{E}_h(z)$  with  $e = \{z, y\}$  and denote the two triangles that have  $e$  as an edge as  $T_1$  and  $T_2$ . Let  $\mathbf{w}_e^z$  be given by (3.5). Then*

$$(3.6a) \quad \mathbf{w}_e^z \in \mathbf{V}_{h,00}^3(z),$$

$$(3.6b) \quad \text{supp } \mathbf{w}_e^z \subset T_1 \cup T_2, \text{ and } \text{div } \mathbf{w}_e^z|_{T_s}(z) = 1 \quad \text{for } s = 1, 2$$

$$(3.6c) \quad \|\nabla \mathbf{w}_e^z\|_{L^\infty(T_1 \cup T_2)} \leq C.$$

The constant  $C$  only depends on the shape regularity of  $T_1$  and  $T_2$ .

**3.1. Schematic for interior vertex.** It will be useful to use the following notation for an interior vertex  $z \in \mathcal{S}_h^{\text{int}}$ . We assume that  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$ , enumerated as before so that  $T_j, T_{j+1}$  share an edge for  $j = 0, \dots, N-1$  and we identify  $T_0$  as  $T_N$  (indices modulo  $N$ ). For  $1 \leq i \leq N$  we let  $e_i$  be the edge shared by  $T_i$  and  $T_{i+1}$  and  $e_N$  is the edge shared by  $T_N$  and  $T_1$ . We let  $y_1, y_2, \dots, y_N$  be the vertices such that  $e_i = \{z, y_i\}$ . We set  $y_0 = z$ . Also,  $\mathbf{n}_i$  is unit-normal to  $e_i$  pointing out of  $T_i$  and  $\theta_i$  is the angle formed by the two edges of  $T_i$  emanating from  $z$ . Moreover, let  $\mathbf{t}_i$  be the unit-tangent vector to  $e_i$  pointing towards  $y_i$ . The edge opposite to  $z$  belonging to  $T_i$  is denoted by  $f_i$ . The unit-normal to  $f_i$  pointing out of  $T_i$  is denoted  $\mathbf{m}_i$ .

$$(3.7) \quad \mathbf{m}_i = |f_i|^{-1}(y_i - y_{i-1})^\perp = |y_i - y_{i-1}|^{-1}(y_i - y_{i-1})^\perp.$$

Finally,  $h_i = \text{dist}(z, f_i)$ . The notation  $\cdot^\perp$  denotes rotation by 90 degrees counter clockwise. See Figure 3 for an illustration.

Using this notation, we will use the shorthand notation  $\psi_i = \psi_{y_i}$  for  $0 \leq i \leq N$  and  $\mathbf{w}_i = \mathbf{w}_{e_i}^z$  for  $1 \leq i \leq N$ .

**3.2. Singular vertices are local interpolating vertices:**  $\mathcal{S}_h^2 \subset \mathcal{L}_h$ . In this section we recall that all singular vertices are local interpolating vertices. The proof can be found in [7], but we recall some of the details.

**Lemma 4.** *It holds  $\mathcal{S}_h^2 \subset \mathcal{L}_h$ . Moreover, there exists a constant  $C_{\text{sing}}$  such that*

$$D_z \leq C_{\text{sing}} \quad \text{for all } z \in \mathcal{S}_h^2,$$

where  $C_{\text{sing}}$  only depends on the shape regularity of the mesh.

*Proof.* We only consider interior singular vertices for simplicity (the proof for boundary singular vertices is similar). Suppose that  $z \in \mathcal{S}_h^{2,\text{int}}$  and we use the notation in Section 3.1. Note that  $N = 4$ . Let  $a = (a_1, \dots, a_4) \in W(z)$ . First define  $b_1 = a_1$  and inductively define

$$b_j = a_j - b_{j-1} \quad \text{for } j = 2, 3.$$

Then define

$$\mathbf{v} = b_1 \mathbf{w}_1 + b_2 \mathbf{w}_2 + b_3 \mathbf{w}_3.$$

By (3.6a),  $\mathbf{v} \in \mathbf{V}_{h,00}^3(z)$ . Using (3.6b) we see that


$$\text{div } \mathbf{v}|_{T_j}(z) = a_j \text{ for } 1 \leq j \leq 3.$$

We also have

$$\text{div } \mathbf{v}|_{T_4}(z) = b_3 = a_3 - a_2 + a_1 = a_4,$$

where we used that  $(a_1, a_2, a_3, a_4) \in W(z)$ . Moreover, using (3.6c) we have

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))} \leq C(b_1 + b_2 + b_3) \leq C \max_{1 \leq j \leq 4} |a_j|,$$



**Figure 3.** Example of  $\mathcal{T}_h(z)$  with  $N = 4$ .

where the constant  $C$  only depends on the shape regularity constant.  $\square$

**3.3. Interior vertices with odd number of triangles are local interpolating vertices.** In this section we prove that if  $z \in \mathcal{S}_h^{\text{int}}$  and  $\mathcal{T}_h(z)$  has an odd number of triangles then  $z \in \mathcal{L}_h$ .

**Lemma 5.** *Let  $z \in \mathcal{S}_h^{\text{int}}$  with  $\mathcal{T}_h(z) = \{T_1, \dots, T_N\}$  and suppose that  $N$  is odd. Then  $z \in \mathcal{L}_h$ . Moreover, there exists a constant  $C_{\text{odd}}$  such that*

$$D_z \leq C_{\text{odd}}.$$

Here  $C_{\text{odd}}$  is a fixed constant that only depends on the shape regularity of the mesh.

*Proof.* We use the notation in Section 3.1. We start by defining some auxiliary functions. First, define

$$\mathbf{v}_1 = \frac{1}{2} \sum_{j=1}^N (-1)^{j-1} \mathbf{w}_j.$$

We see that  $\mathbf{v}_1 \in \mathbf{V}_{h,00}^3(z)$ , by (3.6a). Moreover, using (3.6b) we see that

$$\text{div } \mathbf{v}_1|_{T_j}(z) = \delta_{1j} \text{ for } 1 \leq j \leq N,$$

where  $\delta_{ij}$  is the Kronecker delta function. Note that here we used crucially that  $N$  is odd. Moreover, we have by (3.6c) that

$$\|\nabla \mathbf{v}_1\|_{L^\infty(\Omega_h(z))} \leq C,$$



where  $C$  only depends on the shape regularity. Next, we define inductively

$$\mathbf{v}_i = \mathbf{w}_i - \mathbf{v}_{i-1} \quad \text{for } 2 \leq i \leq N.$$

Then, we easily see that  $\mathbf{v}_i \in \mathbf{V}_{h,00}^3(z)$  and

$$(3.8) \quad \operatorname{div} \mathbf{v}_i|_{T_j}(z) = \delta_{ij} \quad \text{for } 1 \leq i, j \leq N,$$

and, furthermore,

$$(3.9) \quad \|\nabla \mathbf{v}_i\|_{L^\infty(\Omega_h(z))} \leq C \quad \text{for } 1 \leq i, j \leq N,$$

where  $C$  only depends on the shape regularity. Now given  $a = (a_1, \dots, a_N) \in W(z)$  we set

$$\mathbf{v} = \sum_{j=1}^N a_j \mathbf{v}_j.$$

Then, using (3.8) we get (3.1). Moreover, using (3.9) we get

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))} \leq C_{\text{odd}} \max_{1 \leq j \leq N} |a_j|,$$

where  $C_{\text{odd}}$  is a fixed constant only depending on the shape regularity of the mesh.  $\square$

**3.4. Interior vertices with even number of triangles.** If  $z \in \mathcal{S}_h^{\text{int}}$  is non-singular and has an even number of triangles containing it, then it is not necessarily the case that  $z \in \mathcal{L}_h$ . In this section we give sufficient conditions for  $z \in \mathcal{L}_h$ . We use the notation in Section 3.1. To do this, in addition to the vector fields  $\mathbf{w}_i$ , we will need other vector fields. We start with

$$(3.10) \quad \boldsymbol{\chi}_i = \frac{12}{|e_i|} \eta_{e_i}^z \mathbf{n}_i = \frac{12}{|e_i|} \psi_z^2 \psi_{y_i} \mathbf{n}_i \quad \text{for } 1 \leq i \leq N.$$

In the following lemma, the indices are calculated modulo  $N$ .

**Lemma 6.** *It holds, for  $1 \leq i \leq N$*

$$(3.11a) \quad \operatorname{supp}(\boldsymbol{\chi}_i) \subset T_i \cup T_{i+1},$$

$$(3.11b) \quad (\operatorname{div} \boldsymbol{\chi}_i)(y_j) = 0 \quad \text{for } 1 \leq j \leq N,$$

$$(3.11c) \quad (\operatorname{div} \boldsymbol{\chi}_i)|_{T_i}(z) = \frac{12 \cot(\theta_i)}{|e_i|^2}, \quad (\operatorname{div} \boldsymbol{\chi}_i)|_{T_{i+1}}(z) = \frac{-12 \cot(\theta_{i+1})}{|e_i|^2},$$

$$(3.11d) \quad \int_{T_i} \operatorname{div} \boldsymbol{\chi}_i dx = 1, \quad \int_{T_{i+1}} \operatorname{div} \boldsymbol{\chi}_i dx = -1,$$

$$(3.11e) \quad \|\nabla \boldsymbol{\chi}_i\|_{L^\infty(T_i \cup T_{i+1})} \leq \frac{C}{|e_i|^2}.$$

*Proof.* It follows from the definition (3.3) of  $\eta_{e_i}^z$  that (3.11a) and (3.11b) hold. A simple calculation using (2.6) and (2.7) shows that

$$\nabla \psi_{y_i}|_{T_i} = \frac{1}{\sin(\theta_i)|e_i|} \mathbf{n}_{i-1}.$$

Thus

$$(\operatorname{div} \boldsymbol{\chi}_i)|_{T_i}(z) = \frac{12}{|e_i|} \psi_z^2(z) \nabla \psi_{y_i}|_{T_i} \cdot \mathbf{n}_i = \frac{12}{\sin(\theta_i)|e_i|^2} \mathbf{n}_{i-1} \cdot \mathbf{n}_i = \frac{12 \cot(\theta_i)}{|e_i|^2}.$$

Similarly, we can show that

$$(\operatorname{div} \chi_i)|_{T_{i+1}}(z) = \frac{-12 \cot(\theta_{i+1})}{|e_i|^2}.$$

To show (3.11d) we use integration by parts and use that  $\eta_{e_i}^z$  vanishes on  $\partial T_i \setminus e_i$  to get

$$\int_{T_i} \operatorname{div} \chi_i dx = 12 \int_{e_i} \eta_{e_i} ds = 1.$$

Similarly, we can show that

$$\int_{T_{i+1}} \operatorname{div} \chi_i dx = -1.$$

To prove (3.11e) we use the definition of  $\chi_i$  and the bound (3.4c).  $\square$

Note that  $\chi_i$  is not in  $\mathbf{V}_{h,0}^3(z)$  by (3.11d). However, again using (3.11d), we see that

$$\chi := \chi_1 + \chi_2 + \cdots + \chi_N,$$

does belong to  $\mathbf{V}_{h,0}^3(z)$ . In fact, using (3.11b) we have that  $\chi \in \mathbf{V}_{h,00}^3(z)$ . We collect it in the following result.

**Lemma 7.** *It holds that  $\chi \in \mathbf{V}_{h,00}^3(z)$  and*

$$(3.12) \quad (\operatorname{div} \chi)|_{T_j}(z) = 12 \cot(\theta_j) \left( \frac{1}{|e_j|^2} - \frac{1}{|e_{j-1}|^2} \right) \quad \text{for } 1 \leq j \leq N,$$

$$(3.13) \quad \|\nabla \chi\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{h_z^2}.$$

The inequality (3.13) follows from (3.11e).

So far, we have  $\mathbf{w}_1, \dots, \mathbf{w}_N$  and  $\chi$  that belong to  $\mathbf{V}_{h,00}^3(z)$ . Next we describe two more functions that also belong to the space.

We let  $\mathbf{E}_1 = [1, 0]^t$  and  $\mathbf{E}_2 = [0, 1]^t$  be canonical directions. We then define

$$\tilde{\xi}_i := \psi_z^2 \mathbf{E}_i \quad \text{for } i = 1, 2.$$

The following result can easily be proven.

**Lemma 8.** *It holds, for  $i = 1, 2$*

$$(3.14a) \quad \tilde{\xi}_i \in \mathbf{V}_h^3(z)$$

$$(3.14b) \quad (\operatorname{div} \tilde{\xi}_i)(y_j) = 0 \quad \text{for } 1 \leq j \leq N$$

$$(3.14c) \quad (\operatorname{div} \tilde{\xi}_i)|_{T_j}(z) = \frac{3}{|T_j|} b_{ji} \quad \text{for } 1 \leq j \leq N$$

$$(3.14d) \quad \int_{T_j} \operatorname{div} \tilde{\xi}_i dx = b_{ji} \quad \text{for } 1 \leq j \leq N,$$

where  $|T_j|$  denotes the area of  $T_j$ , and using (3.7) we get

$$(3.15) \quad b_{ji} = \frac{-|f_j| \mathbf{m}_j \cdot \mathbf{E}_i}{3} = \frac{-(y_j - y_{j-1})^\perp \cdot \mathbf{E}_i}{3} = \frac{-(y_j - y_{j-1}) \cdot \mathbf{E}_i^\perp}{3},$$

where  $\mathbf{v}^\perp$  denotes the rotation of  $\mathbf{v}$  by 90 degrees counter clockwise. Moreover, the following bound holds

$$(3.16) \quad \|\nabla \tilde{\boldsymbol{\xi}}_i\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{h_z}.$$

*Proof.* From the definition  $\psi_z$  we have that (3.14a) and (3.14b) hold. To show (3.14c) we have

$$(\operatorname{div} \tilde{\boldsymbol{\xi}}_i)|_{T_j}(z) = 2\psi_z(z)\nabla\psi_z|_{T_j} \cdot \mathbf{E}_i = \frac{-2\mathbf{m}_j \cdot \mathbf{E}_i}{h_j} = \frac{-|f_j|\mathbf{m}_j \cdot \mathbf{E}_i}{|T_j|}.$$

In the last equation we used that  $|T_j| = \frac{h_j|f_j|}{2}$ . Using that  $\operatorname{div} \tilde{\boldsymbol{\xi}}_i$  is linear in  $T_j$  and (3.14b), (3.14c) we have

$$\int_{T_j} \operatorname{div} \tilde{\boldsymbol{\xi}}_i dx = \frac{|T_j|}{3} (\operatorname{div} \tilde{\boldsymbol{\xi}}_i)|_{T_j}(z) = b_{ji}.$$

Finally, (3.16) follows by a simple computation.  $\square$

We note that  $\tilde{\boldsymbol{\xi}}_i$  does not belong to  $\mathbf{V}_{h,0}^3(z)$  by (3.14d). However, by integration by parts and using that  $\tilde{\boldsymbol{\xi}}_i$  vanishes on  $\partial\Omega_h(z)$  we do have that  $\int_{\Omega_h(z)} \operatorname{div} \tilde{\boldsymbol{\xi}}_i dx = 0$  and hence

$$(3.17) \quad b_{1i} + b_{2i} + \cdots + b_{Ni} = 0 \quad \text{for } i = 1, 2.$$

This also follows by summing (3.15).

We can now correct  $\tilde{\boldsymbol{\xi}}_i$  to make it belong to  $\mathbf{V}_{h,00}^3(z)$ . We define

$$\boldsymbol{\xi}_i := \tilde{\boldsymbol{\xi}}_i - c_{1i}\boldsymbol{\chi}_1 - c_{2i}\boldsymbol{\chi}_2 - \cdots - c_{N-1,i}\boldsymbol{\chi}_{N-1} \quad \text{for } i = 1, 2,$$

where

$$c_{ji} = b_{1i} + b_{2i} + \cdots + b_{ji} = \frac{1}{3}(y_N - y_j) \cdot \mathbf{E}_i^\perp,$$

for  $j = 1, \dots, N$ .

In the following result, indices are calculated mod  $N$ , and in particular  $c_{0,i} = c_{N,i} = 0$ .

**Lemma 9.** *It holds, for  $i = 1, 2$ ,  $\boldsymbol{\xi}_i \in \mathbf{V}_{h,00}^3(z)$  and*

$$(3.18) \quad \operatorname{div}(\boldsymbol{\xi}_i)|_{T_j}(z) = \frac{3}{|T_j|}b_{ji} - 12\cot(\theta_j)\left(\frac{c_{ji}}{|e_j|^2} - \frac{c_{j-1,i}}{|e_{j-1}|^2}\right) \quad \text{for } 1 \leq j \leq N,$$

$$(3.19) \quad \|\nabla \boldsymbol{\xi}_i\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{h_z}.$$

The bound (3.19) follows from (3.16) and (3.11e).

We define for  $i = 1, 2$

$$(3.20) \quad d_{ji} := (\operatorname{div} \boldsymbol{\xi}_i)|_{T_j}(z) = \frac{3}{|T_j|}b_{ji} - 12\cot(\theta_j)\left(\frac{c_{ji}}{|e_j|^2} - \frac{c_{j-1,i}}{|e_{j-1}|^2}\right) \quad \text{for } 1 \leq j \leq N,$$

and

$$(3.21) \quad d_{j0} := (\operatorname{div} \boldsymbol{\chi})|_{T_j}(z) = \cot(\theta_j)\left(\frac{1}{|e_j|^2} - \frac{1}{|e_{j-1}|^2}\right) \quad \text{for } 1 \leq j \leq N.$$

Using (3.19) and (3.13), we note that for all  $1 \leq j \leq N$

$$(3.22) \quad |d_{ji}| \leq \frac{C}{h_z^s} \quad \text{where } s = 1 \text{ if } i = 1, 2 \text{ and } s = 2 \text{ if } i = 0.$$

We can now prove the following important result.

**Lemma 10.** *Let  $z \in \mathcal{S}_h^{1,int}$  with  $\mathcal{T}_h(z) = \{T_1, \dots, T_N\}$  with  $N$  even. Assume that for at least one  $i = 0, 1, 2$*

$$(3.23) \quad \mathcal{D}_i := \sum_{j=1}^N (-1)^j d_{ji} \neq 0,$$

*then  $z \in \mathcal{L}_h$ . Moreover, in this case, there exists a constant  $C$  depending only on the shape regularity constant such that*

$$(3.24) \quad D_z \leq C \max_{0 \leq i \leq 2} \left( 1 + \frac{1}{|\mathcal{D}_i| h_z^s} \right),$$

*where  $s = 1$  if  $i = 1, 2$  and  $s = 2$  if  $i = 0$ .*

*Proof.* We set  $\xi_0 = \chi$ . Let  $i$  be such that (3.23) holds. We let  $s_1 = 0$  and define inductively

$$s_j = d_{ji} - s_{j-1} \quad \text{for } 2 \leq j \leq N.$$

Define

$$\mathbf{v}_1 = \frac{-1}{\mathcal{D}_i} (\xi_i - s_2 \mathbf{w}_2 - \dots - s_{N-1} \mathbf{w}_{N-1} - s_N \mathbf{w}_N).$$

We easily have that, using (3.20), (3.21)

$$\operatorname{div} \mathbf{v}_1|_{T_j}(z) = \frac{-1}{\mathcal{D}_i} (d_{ji} - (s_{j-1} + s_j)) = 0 \quad \text{for } 2 \leq j \leq N,$$

and

$$\operatorname{div} \mathbf{v}_1|_{T_1}(z) = \frac{-1}{\mathcal{D}_i} (d_{1i} - s_N).$$

We see that  $s_N = \sum_{j=2}^N (-1)^j d_{ji}$ , we have  $d_{1i} - s_N = -\sum_{j=1}^N (-1)^j d_{ji} = -\mathcal{D}_i$ . Hence,

$$\operatorname{div} \mathbf{v}_1|_{T_1}(z) = 1.$$

The following bound follows from (3.22), (3.19), (3.13) and (3.6c)

$$(3.25) \quad \|\nabla \mathbf{v}_1\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{|\mathcal{D}_i| h_z^s}.$$

Then, we define inductively

$$\mathbf{v}_\ell = \mathbf{w}_\ell - \mathbf{v}_{\ell-1} \quad \text{for } 2 \leq \ell \leq N.$$

We then easily see that

$$(3.26) \quad \operatorname{div} \mathbf{v}_\ell|_{T_j}(z) = \delta_{\ell j} \quad \text{for } 1 \leq \ell, j \leq N.$$

Moreover, from (3.6c) and (3.25) we have

$$\|\nabla \mathbf{v}_\ell\|_{L^\infty(\Omega_h(z))} \leq C \left( 1 + \frac{1}{|\mathcal{D}_i| h_z^s} \right).$$

For an arbitrary  $a = (a_1, a_2, \dots, a_N) \in W(z)$  we simply define

$$\mathbf{v} = \sum_{\ell=1}^N a_\ell \mathbf{v}_\ell.$$

Then,  $\mathbf{v}$  satisfies (3.1) and hence  $z \in \mathcal{L}_h$ . By (3.4) we have

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))} \leq C \left(1 + \frac{1}{|\mathcal{D}_i| h_z^s}\right) \max_\ell |a_\ell|,$$

which proves (3.24).  $\square$

**3.5. Simplification of condition (3.23).** In the case  $i = 0$ , we have

$$\begin{aligned} \mathcal{D}_0 &= \sum_{i=1}^N (-1)^j d_{j0} = \sum_{i=1}^N (-1)^j \cot(\theta_j) (|e_j|^{-2} - |e_{j-1}|^{-2}) \\ (3.27) \quad &= \sum_{i=1}^N (-1)^j (\cot(\theta_j) + \cot(\theta_{j+1})) |e_j|^{-2}. \end{aligned}$$

Thus we see that generically this is nonzero, since the lengths  $|e_j|$  can be chosen independently of the angles  $\theta_j$ . For  $i = 1, 2$ , we have

$$\begin{aligned} \sum_{j=1}^N (-1)^j \frac{3}{|T_j|} b_{ji} &= \sum_{j=1}^N (-1)^j \frac{1}{|T_j|} (y_{j-1} - y_j) \cdot \mathbf{E}_i^\perp \\ (3.28) \quad &= - \sum_{j=1}^N (-1)^j \left( \frac{1}{|T_j|} + \frac{1}{|T_{j+1}|} \right) y_j \cdot \mathbf{E}_i^\perp \end{aligned}$$

For  $i = 1, 2$ , we have

$$\begin{aligned} -12 \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{c_{j,i}}{|e_j|^2} - \frac{c_{j-1,i}}{|e_{j-1}|^2} \right) &= -12 \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{c_{j,i}}{|e_j|^2} - \frac{c_{j-1,i}}{|e_{j-1}|^2} \right) \\ &= 4 \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{y_j - y_N}{|e_j|^2} - \frac{y_{j-1} - y_N}{|e_{j-1}|^2} \right) \cdot \mathbf{E}_i^\perp \\ &= 4 \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{y_j}{|e_j|^2} - \frac{y_{j-1}}{|e_{j-1}|^2} \right) \cdot \mathbf{E}_i^\perp \\ (3.29) \quad &- 4 \left( \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{1}{|e_j|^2} - \frac{1}{|e_{j-1}|^2} \right) \right) y_N \cdot \mathbf{E}_i^\perp \\ &= 4 \sum_{j=1}^N (-1)^j \cot(\theta_j) \left( \frac{y_j}{|e_j|^2} - \frac{y_{j-1}}{|e_{j-1}|^2} \right) \cdot \mathbf{E}_i^\perp \\ &- 4 \left( \sum_{j=1}^N (-1)^j d_{j0} \right) y_N \cdot \mathbf{E}_i^\perp \\ &= 4 \sum_{j=1}^N (-1)^j \frac{\cot(\theta_j) + \cot(\theta_{j+1})}{|e_j|^2} y_j \cdot \mathbf{E}_i^\perp - 4 \mathcal{D}_0 y_N \cdot \mathbf{E}_i^\perp. \end{aligned}$$

Therefore, for  $i = 1, 2$ ,

$$(3.30) \quad \mathcal{D}_i = \sum_{j=1}^N (-1)^j \left( 4 \frac{\cot(\theta_j) + \cot(\theta_{j+1})}{|e_j|^2} - \left( \frac{1}{|T_j|} + \frac{1}{|T_{j+1}|} \right) \right) y_j \cdot \mathbf{E}_i^\perp - 4\mathcal{D}_0 y_N \cdot \mathbf{E}_i^\perp.$$

#### 4. MESHES WHERE (3.23) FAILS TO HOLD

Lemma 10 gives sufficient conditions for an interior vertex  $z$  with an even number of triangles to be a local interpolating vertex (i.e.,  $z \in \mathcal{L}_h$ ). We see that (3.23) is a mild constraint and that a generic vertex will satisfy (3.23), however, there are important examples of vertices which do not satisfy (3.23) and perhaps are not local interpolating vertices. Here we present some examples.

**4.1. Regular  $N$ -gon with  $N$  even.** Suppose that  $\Omega_h(z)$  is a triangulated regular  $N$ -gon with  $N$  even. More precisely, we assume that  $\Omega_h(z)$  is subdivided by  $N$  similar triangles, with edge lengths  $|e_1| = |e_2| = \dots = |e_N|$  and interior angles  $\theta_1 = \theta_2 = \dots = \theta_N$ . Then we can show that (3.23) does not hold.

First of all, the condition on the edge lengths alone implies that  $d_{j0} = 0$  for all  $j = 1, \dots, N$ . Thus  $\mathcal{D}_0 = 0$ .

Now consider  $i = 1, 2$ . The vertices of the regular  $N$ -gon can be written as

$$y_k = h e^{\iota 2\pi k/N},$$

for some  $h$ . Here we make the standard association  $e^{\iota\theta}$  with the vector  $[\cos \theta, \sin \theta]^t$ . We conclude that, for  $N$  even,

$$(4.1) \quad h^{-1} \sum_{j=1}^N (-1)^j y_j = \sum_{j=1}^{N/2} e^{\iota 4\pi j/N} - \sum_{j=1}^{N/2} e^{\iota 2\pi(2j-1)/N} = (1 - e^{-\iota 2\pi/N}) \sum_{j=1}^{N/2} e^{\iota 4\pi j/N} = \mathbf{0}.$$

Since all the triangles are the same, using (3.30) and that we showed  $\mathcal{D}_0 = 0$  we have for  $i = 1, 2$


$$\mathcal{D}_i = \left( \frac{8 \cot(\theta_1)}{|e_1|^2} - \frac{2}{|T_1|} \right) \sum_{j=1}^N (-1)^j y_j \cdot \mathbf{E}_i^\perp = 0,$$

where we used (4.1). Thus (3.23) fails for  $i = 1, 2$  as well, and Lemma 10 cannot be used.

**4.2. Three lines mesh.** For the three-lines mesh generating a regular hexagonal pattern as shown on the left in Figure 4, the condition (3.23) also fails for  $i = 0, 1, 2$  since each vertex is at the center of a regular hexagon.

#### 5. CROSSED TRIANGLES

Consider the mesh shown on the right in Figure 4. Half of the vertices are at the center of a regular 4-gon, but these are singular vertices, so these are all local interpolating vertices. For the other vertices, at the center of a non-regular 8-gon, we can argue as follows. Since the interior angles are all the same ( $\pi/4$ ), and  $\cot(\pi/4) = 1$ , we have using (3.27)



**Figure 4.** For the mesh on left no interior vertex satisfies (3.23). All interior vertices on the mesh on the right belong to  $\mathcal{L}_h$ .

$$(5.1) \quad \mathcal{D}_0 = 2 \sum_{j=1}^8 (-1)^j \frac{1}{|e_j|^2}.$$

Let  $L$  be the length of the smallest edge:  $L = \min \{|e_j| : j = 1, \dots, 8\}$ . Then the longest edge length is  $\sqrt{2}L$ , and the edge lengths  $|e_j|$  alternate  $L, \sqrt{2}L, L, \sqrt{2}L, \dots$ . Thus

$$\sum_{j=1}^8 (-1)^j \frac{1}{|e_j|^2} = \pm \frac{2}{L^2},$$

depending on where we start the counting. Therefore condition (3.23) holds at these vertices, and thus Lemma 10 can be applied to conclude that these vertices are also local interpolating vertices. Thus all of the vertices in the mesh shown on the right in Figure 4 are in  $\mathcal{L}_h$ .

## 6. INF-SUP STABILITY WHEN ALL INTERIOR VERTICES BELONG TO $\mathcal{L}_h$

In the previous section we have identified many local interpolating vertices. In particular, singular vertices and interior vertices with odd number of triangles are local interpolating vertices; see Lemmas 4 and 5. If  $z$  is an interior non-singular vertex with even number of triangles then Lemma 10 gives sufficient conditions for it to be a local interpolating vertex. None of the above examples address boundary vertices that are non-singular. In this section we will show how to interpolate at those vertices but in a non-local way. Then, using that result and assuming that  $\mathcal{S}_h^{\text{int}} \subset \mathcal{L}_h$  we will prove inf-sup stability. In the next section we will address  $\mathcal{S}_h^{\text{int}} \not\subset \mathcal{L}_h$ .

For vertices that are not local interpolating vertices we can still interpolate there but with a side effect of polluting a neighboring vertex. In other words, the vector field will not belong to  $\mathbf{V}_{h,00}^3(z)$ . To do this, we will need to define a piecewise cubic function that has average zero on edges. For every  $z \in \mathcal{S}_h^{\partial}$  and interior edge  $e \in \mathcal{E}_h$  with  $e = \{z, y\}$  we set

$$(6.1) \quad \kappa_e^z = \eta_e^z - \frac{1}{2} \psi_z \psi_y = \psi_z^2 \psi_y - \frac{1}{2} \psi_z \psi_y.$$

The function  $\kappa_e^z$  will play the same role as  $\gamma_e^z$  in [7] but the difference is that the added term in  $\kappa_e^z$  is piecewise quadratic. Let  $T_1, T_2 \in \mathcal{T}_h(z)$  be two triangles that have  $e$  as an edge, and let  $\theta_i$  be the angle between the edges of  $T_i$  emanating from  $z$ .

Then we can easily verify the following:

$$(6.2a) \quad \text{supp } \kappa_e^z \subset T_1 \cup T_2,$$

$$(6.2b) \quad (\nabla \kappa_e^z)(\sigma) = 0 \quad \text{for } \sigma \in \mathcal{S}_h, \sigma \neq z \text{ and } \sigma \neq y,$$

$$(6.2c) \quad \int_e \kappa_e^z ds = 0$$

$$(6.2d) \quad \nabla \kappa_e^z|_{T_i}(z) = \frac{1}{2} \nabla \psi_y|_{T_i}, \quad \nabla \kappa_e^z|_{T_i}(y) = -\frac{1}{2} \nabla \psi_z|_{T_i} \quad \text{for } i = 1, 2.$$

Using these functions we can prove the following result.

**Lemma 11.** *For every  $p \in Q_h^2$  and  $z \in \mathcal{S}_h^\partial$  there exists a  $\mathbf{v} \in \mathbf{V}_{h,0}^3(z)$  such the following properties hold:*

$$(6.3a) \quad \text{div } \mathbf{v}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h^\partial, \sigma \neq z,$$

$$(6.3b) \quad \text{div } \mathbf{v}|_T(z) = p|_T(z) \quad \text{for all } T \in \mathcal{T}_h(z).$$

If  $z$  is non-singular

$$(6.4) \quad \|\nabla \mathbf{v}\|_{L^2(\Omega_h(z))} \leq C \left( \frac{1}{\Theta(z)} + 1 \right) \|p\|_{L^2(\Omega_h(z))}.$$

If  $z$  is singular

$$(6.5) \quad \|\nabla \mathbf{v}\|_{L^2(\Omega_h(z))} \leq C \|p\|_{L^2(\Omega_h(z))}.$$

The constant  $C$  only depends only on the shape regularity.

*Proof.* If  $z$  is singular, the result follows from Lemma 4, so now assume that  $z$  is non-singular. Enumerate the triangles such that  $T_1$  and  $T_N$  each have a boundary edge, and  $T_j, T_{j+1}$  share an edge  $e_j = \{z, y_j\}$ , for  $j = 1, \dots, N-1$ . Let  $\theta_j$  denote the angle between the edges of  $T_j$  originating from  $z$ . Also let  $\mathbf{n}_j$  be the normal to  $e_j$  out of  $T_j$  and  $\mathbf{t}_j$  be tangent to  $e_j$  pointing away from  $z$ . Let  $1 \leq s \leq N-1$  be such that  $|\sin(\theta_s + \theta_{s+1})| = \Theta(z)$ . We will define vector fields  $\mathbf{v}_1, \dots, \mathbf{v}_N$ . We start by defining  $\mathbf{v}_s$ .

$$\mathbf{v}_s = \frac{2|e_s| \sin(\theta_s)}{\sin(\theta_s + \theta_{s+1})} \mathbf{t}_{s+1} \kappa_{e_s}^z.$$

Then, we can easily show that

$$\text{div } \mathbf{v}_s|_{T_i}(z) = \delta_{i,s} \text{ for } 1 \leq i \leq N,$$

where  $\delta_{i,s}$  is the Kronecker  $\delta$ . Indeed,

$$\text{div } \mathbf{v}_s|_{T_i}(z) = \frac{2|e_s| \sin(\theta_s)}{\sin(\theta_s + \theta_{s+1})} \mathbf{t}_{s+1} \cdot \nabla \kappa_{e_s}^z|_{T_i}(z) = \frac{|e_s| \sin(\theta_s)}{\sin(\theta_s + \theta_{s+1})} \mathbf{t}_{s+1} \cdot \nabla \psi_{y_s}|_{T_i}(z).$$

The result follows after using (2.5) and (2.7) to calculate  $\psi_{y_s}|_{T_i}$  and using basic trigonometry. Then we can define inductively for  $s+1 \leq j \leq N$ .

$$\mathbf{v}_j = \mathbf{w}_{e_j}^z - \mathbf{v}_{j-1}.$$

Also, for  $1 \leq j \leq s-1$  we define

$$\mathbf{v}_j = \mathbf{w}_{e_j}^z - \mathbf{v}_{j+1}.$$



Hence, we have the following property:

$$(6.6) \quad \operatorname{div} \mathbf{v}_j|_{T_i}(s) = \delta_{i,j} \text{ for } 1 \leq i, j \leq N.$$

We then define

$$\mathbf{v} = \sum_{j=1}^N p|_{T_j}(z) \mathbf{v}_j.$$

The stated conditions on  $\mathbf{v}$  are easily verified.  $\square$

We can then use the above lemma to prove the following result. We define

$$\Theta_{\min, \partial} = \min_{z \in \mathcal{S}_h^{1, \partial}} \Theta(z).$$

**Lemma 12.** *For every  $p \in Q_h^2$  there exists a  $\mathbf{v} \in \mathbf{V}_{h,0}^3$  such that*

$$(6.7) \quad (\operatorname{div} \mathbf{v} - p)(z) = 0 \quad \text{for all } z \in \mathcal{S}_h^\partial,$$

and

$$\|\nabla \mathbf{v}\|_{L^2(\Omega)} \leq C_b \left( \frac{1}{\Theta_{\min, \partial}} + 1 \right) \|p\|_{L^2(\Omega)}.$$

The constant  $C_b$  only depends on the shape regularity.

*Proof.* Let  $p \in Q_h^2$ . For every  $z \in \mathcal{S}_h^\partial$  let  $\mathbf{v}^z$  be the vector field from Lemma 11 then we set

$$\mathbf{v} = \sum_{z \in \mathcal{S}_h^\partial} \mathbf{v}^z.$$

Then, it is easy to verify the conditoin on  $\mathbf{v}$ .  $\square$

We can now prove the main result of the section.

**Theorem 1.** *Assume that  $\mathcal{S}_h^{\text{int}} \subset \mathcal{L}_h$ . Then, for every  $p \in Q_h^2$  there exists a  $\mathbf{v} \in V_{h,0}^3$  satisfying*

$$(6.8) \quad (\operatorname{div} \mathbf{v} - p)(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h,$$

with the bound

$$\|\mathbf{v}\|_{H^1(\Omega)} \leq C(1 + D) \left( 1 + \frac{1}{\Theta_{\min, \partial}} \right) \|p\|_{L^2(\Omega)}.$$

where  $D = \max_{z \in \mathcal{L}_h} D_z$ .

*Proof.* Let  $\mathbf{v}_1$  be from Lemma 12 and let  $p_1 = p - \operatorname{div} \mathbf{v}_1$  and we note that  $p_1 \in Q_h^2$  with  $p_1$  vanishing on boundary vertices. Since  $\mathcal{S}_h^{\text{int}} \subset \mathcal{L}_h$  for every  $z \in \mathcal{S}_h^{\text{int}}$  there exists a  $\mathbf{v}^z \in \mathbf{V}_{h,00}^3(z)$  such that


$$(\operatorname{div} \mathbf{v}^z - p_1)(z) = 0$$

with

$$\|\nabla \mathbf{v}^z\|_{L^\infty(\Omega_h(z))} \leq D_z \max_{T \in \mathcal{T}_h(z)} |p_1|_T(z)|.$$

Using an inverse estimate we can show

$$\|\nabla \mathbf{v}^z\|_{L^2(\Omega_h(z))} \leq CD_z \|p_1\|_{L^2(\Omega_h(z))}.$$



**Figure 5.** Illustration for Lemma 13, with  $N = 4$ .

If we set

$$\mathbf{v}_2 = \sum_{z \in \mathcal{S}_h^{\text{int}}} \mathbf{v}^z$$

and set  $\mathbf{v} = \mathbf{v}_1 + \mathbf{v}_2$ , then the desired conditions on  $\mathbf{v}$  are met.  $\square$

Using Lemma 2 we have the following corollary.

**Corollary 1.** Assume that  $\mathcal{S}_h^{\text{int}} \subset \mathcal{L}_h$ , then the inf-sup condition holds for  $Q_h^2 \times \mathbf{V}_h^3$  with constants given by Lemma 2 and Theorem 1.

## 7. INF-SUP STABILITY: THE GENERAL CASE

As mentioned above for most meshes  $\mathcal{S}_h^{\text{int}} \subset \mathcal{L}_h$  and hence by the previous section one can prove inf-sup stability. However, for very important meshes, such as the diagonal mesh, none of the interior vertices belong to  $\mathcal{L}_h$  (i.e.  $\mathcal{L}_h \cap \mathcal{S}_h^{\text{int}} = \emptyset$ ). However, if some interior nodes belong to  $\mathcal{L}_h$  then it might hold that  $\text{div } \mathbf{V}_h^3 = Q_h^2$  and we can give a bound for inf-sup constant. To do this we use a concept of a tree and paths. We consider the mesh as a graph and consider trees and paths that are subgraphs of the mesh. Precise statements are given below.

**7.1. Paths and trees in a mesh.** We will prove that if there is a tree of the mesh  $\mathcal{T}_h$  with root in  $\mathcal{L}_h$  satisfying certain mild conditions then we can interpolate a pressure on all the vertices of the tree. We start with some preliminary results. Let  $T_1, T_2 \in \mathcal{T}_h(z)$  be two triangles that have  $e$  as an edge and let  $\phi_i$  be the angle between the edges of  $T_i$  emanating from  $z$ . Then we define

$$(7.1) \quad M_e^z = \cot(\phi_1) + \cot(\phi_2).$$

Note that  $M_e^z = 0$  if  $\phi_1 + \phi_2 = \pi$ .

**Lemma 13.** Let  $z, y \in \mathcal{S}_h$  and  $e = \{z, y\} \in \mathcal{E}_h$  and suppose that  $M_e^z \neq 0$ . Let  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$  and suppose  $T_1$  and  $T_2$  are the two triangles that share  $e$  as an edge and let  $\theta_i$  be the angles of  $T_i$  originating from  $y$  for  $i = 1, 2$ ; see Figure 5. Let  $a = (a_1, a_2, \dots, a_N) \in \mathbb{R}^N$

and define the alternating sum  $s(a) = \sum_{j=1}^N (-1)^{j-1} a_j$ . Then, there exists a  $\mathbf{v} \in \mathbf{V}_{h,0}^3(z)$

$$(7.2a) \quad \operatorname{div} \mathbf{v}|_{T_i}(z) = a_i \quad \text{for all } 1 \leq i \leq N$$

$$(7.2b) \quad \operatorname{div} \mathbf{v}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq z, \sigma \neq y,$$

$$(7.2c) \quad \operatorname{div} \mathbf{v}|_{T_1}(y) = -s(a) \frac{\cot(\theta_1)}{M_e^z}, \quad \operatorname{div} \mathbf{v}|_{T_2}(y) = s(a) \frac{\cot(\theta_2)}{M_e^z}$$

$$(7.2d) \quad \|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{|M_e^z|} \max_{1 \leq i \leq N} |a_i|.$$

The constant  $C$  depends only on the shape regularity.

*Proof.* We prove the result in the case  $z$  is an interior vertex. The case  $z$  is a boundary vertex is similar. We use the notation from Section 3.1. We need to define some auxiliary vector fields. First let

$$\mathbf{r} = 2|e_1| \kappa_{e_1}^z \mathbf{n}_1,$$

where we recall that the definition of  $\kappa_{e_1}^z$  is given in (6.1). Using, (6.2a) and (6.2c) it is easy to show that  $\mathbf{r} \in \mathbf{V}_{h,0}^3(z)$ . Then, using (6.2b) we have

$$\operatorname{div} \mathbf{r}|_{T_i}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq z, \sigma \neq y.$$

Also, using (6.2d), (2.6), (2.7) we can show that

$$\operatorname{div} \mathbf{r}|_{T_1}(z) = \cot(\phi_1), \quad \operatorname{div} \mathbf{r}|_{T_2}(z) = -\cot(\phi_2),$$

and

$$\operatorname{div} \mathbf{r}|_{T_1}(y) = -\cot(\theta_1), \quad \operatorname{div} \mathbf{r}|_{T_2}(z) = \cot(\theta_2).$$

If we let

$$\mathbf{v}_1 = \frac{1}{M_e^z} (\mathbf{r} + \cot(\phi_2) \mathbf{w}_1),$$

then using the properties of  $\mathbf{w}_1$  (e.g. (3.6)) we have the following

$$\operatorname{div} \mathbf{v}_1|_{T_1}(z) = 1, \quad \operatorname{div} \mathbf{v}_1|_{T_2}(z) = 0,$$

$$\operatorname{div} \mathbf{v}_1|_{T_1}(y) = -\frac{\cot(\theta_1)}{M_e^z}, \quad \operatorname{div} \mathbf{v}_1|_{T_2}(y) = \frac{\cot(\theta_2)}{M_e^z},$$

and

$$\|\nabla \mathbf{v}_1\|_{L^\infty(\Omega_h(z))} \leq \frac{C}{|M_e^z|},$$

where the constant  $C$  only depends on the shape regularity constant.

Next, we define inductively

$$\mathbf{v}_j = \mathbf{w}_{j-1} - \mathbf{v}_{j-1} \quad \text{for } 2 \leq j \leq N.$$

Using the properties of  $\mathbf{v}_1$  just proved together with (3.6), we can show the following for all  $1 \leq j \leq N$ :

$$\begin{aligned}
(7.3a) \quad & \mathbf{v}_j \in \mathbf{V}_{h,0}^3(z), \\
(7.3b) \quad & \operatorname{div} \mathbf{v}_j(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq z, \sigma \neq y, \\
(7.3c) \quad & \operatorname{div} \mathbf{v}_j|_{T_i}(z) = \delta_{ij} \quad \text{for all } 1 \leq i, j \leq N, \\
(7.3d) \quad & \|\nabla \mathbf{v}_j\|_{L^\infty(\Omega_h(z))} \leq C \left(1 + \frac{1}{|M_e^z|}\right) \leq \frac{C}{|M_e^z|},
\end{aligned}$$

where we used that  $|M_e^z| \leq c$  is bounded where the constant  $c$  depends on the shape regularity. By the definition of  $\mathbf{v}_j$  and using that  $\operatorname{div} \mathbf{w}_j(y) = 0$ , we note that  $\operatorname{div} \mathbf{v}_j(y) = (-1)^{j-1} \operatorname{div} \mathbf{v}_1(y)$ .

We now set

$$\mathbf{v} = \sum_{j=1}^N a_j \mathbf{v}_j.$$

We easily see conditions (7.2) hold. □

We can apply the previous result repeatedly to generalize the result for a path.

**Definition 7.1.** Given  $z, y \in \mathcal{S}_h$   $P = \{y_0, y_1, \dots, y_L\}$  is a path between  $y_0 = z$ ,  $y_L = y$  if  $e_i = \{y_{i-1}, y_i\} \in \mathcal{E}_h$  and  $y_i \neq y_j$  for  $i \neq j$ . We say that the path is *acceptable* if  $M_{e_i}^{y_{i-1}} \neq 0$  for  $1 \leq i \leq L$ .

See Figure 7.6 for an illustration. For an acceptable path  $P$  as in Definition 7.1 we define for  $1 \leq j \leq L-1$

$$\tilde{\rho}_{z,y_j} := \frac{M_{e_1}^{y_1} M_{e_2}^{y_2} \cdots M_{e_j}^{y_j}}{M_{e_1}^{y_0} M_{e_2}^{y_1} \cdots M_{e_j}^{y_{j-1}}}.$$

We also let  $\tilde{\rho}_{z,z} = 1$ . Moreover, we define


$$(7.4) \quad \rho_{z,y_{j+1}} := \frac{\tilde{\rho}_{z,y_j}}{M_{e_{j+1}}^{y_j}}.$$

Finally, we let

$$(7.5) \quad \rho(P) = \max_{1 \leq j \leq L} |\rho_{z,y_j}|.$$

Also for any collection of vertices  $P$  we define  $\mathcal{T}_h(P) = \{T \in \mathcal{T}_h : T \in \mathcal{T}_h(y) \text{ for some } y \in P\}$ . We define  $\Omega_h(P) = \bigcup_{T \in \mathcal{T}_h(P)} T$ .

**Lemma 14.** Suppose that  $z, y \in \mathcal{S}_h$  and let  $\mathcal{T}_h(z) = \{T_1, T_2, \dots, T_N\}$ . Let  $P = \{y_0, y_1, \dots, y_L\}$  with  $y_0 = z$ ,  $y_L = y$  be an acceptable path. Also, we denote by  $K_1$  and  $K_2$  the two triangles that share  $e_L$  and  $\theta_i$  the corresponding angles. For every  $a = (a_1, \dots, a_N) \in \mathbb{R}^N$  there exists a



**Figure 6.** Illustration of acceptable path, with  $N = 6$ ,  $L=4$ .

$\mathbf{v} \in \mathbf{V}_{h,0}^3$  with support in  $\Omega_h(P)$  such that

$$(7.6a) \quad \operatorname{div} \mathbf{v}|_{T_i}(z) = a_i \quad \text{for all } 1 \leq i \leq N,$$

$$(7.6b) \quad \operatorname{div} \mathbf{v}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq z, \sigma \neq y,$$

$$(7.6c) \quad \operatorname{div} \mathbf{v}|_{K_1}(y) = \pm s(a) \tilde{\rho}_{z,y_{L-1}} \frac{\cot(\theta_1)}{M_{e_L}^{y_{L-1}}}, \quad \operatorname{div} \mathbf{v}|_{K_2}(y) = \mp s(a) \tilde{\rho}_{z,y_{L-1}} \frac{\cot(\theta_2)}{M_{e_L}^{y_{L-1}}}$$

$$(7.6d) \quad \operatorname{div} \mathbf{v}|_T(y) = 0 \quad \text{for all } T \in \mathcal{T}_h(y), T \neq K_1, T \neq K_2,$$

$$(7.6e) \quad \|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(P))} \leq C \rho(P) \max_{1 \leq i \leq N} |a_i|.$$

Here  $s(a) = \sum_{j=1}^N (-1)^j a_j$ . The constant  $C$  only depends on the shape regularity of the mesh.

*Proof.* Let  $a = (a_1, \dots, a_N) \in \mathbb{R}^N$  be given. We will use the following notation:  $e_j$  is the edge with vertices  $y_{j-1}, y_j$ . We assume that  $\mathcal{T}_h(y_j) = \{T_{1,j}, T_{2,j}, \dots, T_{N_j,j}\}$  and such that  $T_{1,j}, T_{2,j}$  share  $e_j$  as a common edge. The corresponding angles are denoted by  $\theta_{1,j}, \theta_{2,j}$ . Note that  $K_1 = T_{1,L}$  and  $K_2 = T_{2,L}$ . By Lemma 13 we have  $\mathbf{r}_1 \in \mathbf{V}_{h,0}^3(y_0)$

$$(7.7a) \quad \operatorname{div} \mathbf{r}_1|_{T_i}(y_0) = a_i \quad \text{for all } 1 \leq i \leq N$$

$$(7.7b) \quad \operatorname{div} \mathbf{r}_1(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq y_0, \sigma \neq y_1,$$

$$(7.7c) \quad \operatorname{div} \mathbf{r}_1|_{T_{1,1}}(y_1) = -s(a) \frac{\cot(\theta_{1,1})}{M_{e_1}^{y_0}}, \quad \operatorname{div} \mathbf{r}_1|_{T_{2,j}}(y_1) = s(a) \frac{\cot(\theta_{2,1})}{M_{e_1}^{y_0}}$$

$$(7.7d) \quad \operatorname{div} \mathbf{r}_1|_{T_{i,1}}(y_1) = 0 \quad \text{for all } 3 \leq i \leq N$$

$$(7.7e) \quad \|\nabla \mathbf{r}_1\|_{L^\infty(\Omega_h(y_0))} \leq \frac{C}{|M_{e_1}^{y_0}|} \max_{1 \leq i \leq N} |a_i|.$$

Now suppose that we have constructed  $\mathbf{r}_j \in \mathbf{V}_{h,0}^3(y_{j-1})$  for  $j = 2, \dots, \ell$  with the following properties

$$\begin{aligned}
(7.8a) \quad & (\operatorname{div} \mathbf{r}_j + \operatorname{div} \mathbf{r}_{j-1})(y_{j-1}) = 0, \\
(7.8b) \quad & \operatorname{div} \mathbf{r}_j(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq y_{j-1}, \sigma \neq y_j, \\
(7.8c) \quad & \operatorname{div} \mathbf{r}_j|_{T_{1,j}}(y_j) = \pm s(a) \tilde{\rho}_{z,y_{j-1}} \frac{\cot(\theta_{1,j})}{M_{e_j}^{y_{j-1}}}, \quad \operatorname{div} \mathbf{r}_j|_{T_{2,j}}(y_j) = \mp s(a) \tilde{\rho}_{z,y_{j-1}} \frac{\cot(\theta_{2,j})}{M_{e_j}^{y_{j-1}}} \\
(7.8d) \quad & \operatorname{div} \mathbf{r}_j|_{T_{i,j}}(y_j) = 0 \quad \text{for all } 3 \leq i \leq N_j, \\
(7.8e) \quad & \|\nabla \mathbf{r}_j\|_{L^\infty(\Omega_h(y_{j-1}))} \leq C |\rho_{z,y_j}| \max_{1 \leq i \leq N} |a_i|.
\end{aligned}$$

Setting  $\tilde{a}_i = \operatorname{div} \mathbf{r}_\ell|_{T_{i,\ell}}(y_\ell)$  for  $1 \leq i \leq N_\ell$  and using (7.8c) and (7.8d) we have

$$\begin{aligned}
(7.9) \quad s(\tilde{a}) &= \sum_{i=1}^{N_\ell} (-1)^{i-1} \tilde{a}_i = \tilde{a}_1 - \tilde{a}_2 = \pm s(a) \tilde{\rho}_{z,y_{\ell-1}} \frac{\cot(\theta_{1,\ell})}{M_{e_j}^{y_{\ell-1}}} \pm s(a) \tilde{\rho}_{z,y_{\ell-1}} \frac{\cot(\theta_{2,\ell})}{M_{e_\ell}^{y_{\ell-1}}}, \\
&= \pm s(a) \tilde{\rho}_{z,y_{\ell-1}} \frac{\cot(\theta_{1,\ell}) + \cot(\theta_{2,\ell})}{M_{e_j}^{y_{\ell-1}}} = \pm s(a) \tilde{\rho}_{z,y_{\ell-1}} \frac{M_{e_\ell}^{y_\ell}}{M_{e_\ell}^{y_{\ell-1}}} = \pm s(a) \tilde{\rho}_{z,y_\ell}.
\end{aligned}$$

Hence, using Lemma 13 we can find  $\mathbf{r}_{\ell+1} \in \mathbf{V}_{h,0}^3(y_\ell)$  such that

$$\begin{aligned}
(7.10a) \quad & \operatorname{div} \mathbf{r}_{\ell+1}|_{T_{i,\ell}}(y_\ell) = -\tilde{a}_i \quad \text{for all } 1 \leq i \leq N_\ell \\
(7.10b) \quad & \operatorname{div} \mathbf{r}_{\ell+1}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq y_\ell, \sigma \neq y_{\ell+1}, \\
(7.10c) \quad & \operatorname{div} \mathbf{r}_{\ell+1}|_{T_{1,\ell+1}}(y_{\ell+1}) = \pm s(\tilde{a}) \frac{\cot(\theta_{1,\ell+1})}{M_{e_{\ell+1}}^{y_\ell}}, \quad \operatorname{div} \mathbf{r}_{\ell+1}|_{T_{2,\ell+1}}(y_{\ell+1}) = \mp s(\tilde{a}) \frac{\cot(\theta_{2,\ell+1})}{M_{e_{\ell+1}}^{y_\ell}} \\
(7.10d) \quad & \|\nabla \mathbf{r}_{\ell+1}\|_{L^\infty(\Omega_h(y_\ell))} \leq \frac{C}{|M_{e_{\ell+1}}^{y_\ell}|} \max_{1 \leq i \leq N_\ell} |\tilde{a}_i| \leq \frac{C}{|M_{e_{\ell+1}}^{y_\ell}|} |s(a)| |\tilde{\rho}_{z,y_\ell}| = C |s(a)| |\rho_{z,y_{\ell+1}}|.
\end{aligned}$$

Hence, using (7.10) and (7.9) we get that (7.8) holds for  $j = \ell + 1$  and hence by induction (7.8) holds for  $2 \leq j \leq L$ .

We let  $\mathbf{v} = \mathbf{r}_1 + \mathbf{r}_2 + \dots + \mathbf{r}_L$ . We then easily see that (7.6a)-(7.6d) hold from (7.8) and (7.7). To prove (7.6e) we use that

$$\|\nabla \mathbf{v}\|_{L^\infty(\Omega_h(P))} \leq \max_{1 \leq i \leq L} \|\nabla \mathbf{r}_i\|_{L^\infty(\Omega_h(y_i))}.$$

Since finitely many  $\{\mathbf{r}_j\}$  have support in  $\Omega_h(y_i)$  the result follows from (7.8e) and (7.7e).  $\square$

The next task is to interpolate values on a tree of the mesh. We need some notation.

**Definition 2.** We say that  $\mathcal{R}(r) := Y \times E$  with  $Y = \bigcup_{0 \leq j \leq M} Y_j$  and  $E = \bigcup_{1 \leq j \leq M} E_j$  is a tree of  $\mathcal{T}_h$  with root  $r$  if the following hold

- 1)  $Y_0 = \{r\}$ ,
- 2) for every  $1 \leq j \leq M$ ,  $Y_j \subset \mathcal{S}_h$ ,  $E_j \subset \mathcal{E}_h$  and  $|Y_j| = |E_j|$ ,
- 3) for every  $y \in Y_j$  there is a unique  $e \in E_j$  such that  $e = \{y, z\}$  for some  $z \in Y_{j-1}$ ,
- 4)  $Y_j \cap Y_i = \emptyset$  for  $i \neq j$ .

If  $y \in Y_j$  and  $z \in Y_{j+s}$  for  $s \geq 1$  and there is a path in  $\mathcal{R}(r)$  connecting  $y$  to  $z$  then we say that  $z$  is a *descendant* of  $y$  and that  $y$  is an *ancestor* of  $z$ . If we let  $y_0 = z$ ,  $y_s = y$ , by path we mean  $P_{z,y} = \{y_0, y_1, \dots, y_s\}$  such that  $y_i \in Y_{j+s-i}$  for  $0 \leq i \leq s$  such that  $e_i = \{y_{i-1}, y_i\} \in E_{j+s-(i-1)}$  for  $1 \leq i \leq s$ . We let  $D(y)$  denote the set of all descendants of  $y$  and  $A(y)$  to be all the ancestors of  $y$ . We know that if  $z \in Y$  then there is a unique path  $P_{z,r}$  (which we denote by  $P_z$ ) from  $z$  to the root  $r$ . We say that the tree  $\mathcal{R}(r)$  is *acceptable* if  $P_z$  is acceptable for each  $z \in Y$ . Moreover, we define

$$(7.11) \quad \rho(\mathcal{R}(r)) = \max_{z \in Y} \rho(P_z),$$

where  $\rho(P_z)$  is defined in (7.5).

We can now state the following result.

**Lemma 15.** *Let  $\mathcal{R}(r) = Y \times E$  with  $Y = \bigcup_{0 \leq j \leq M} Y_j$  and  $E = \bigcup_{1 \leq j \leq M} E_j$  be an acceptable tree of  $\mathcal{T}_h$  with root  $r \in \mathcal{L}_h$ . Then, for any  $p \in Q_h^2$  there exist  $\mathbf{v} \in \mathbf{V}_{h,0}^3$  such that*

$$(7.12a) \quad \text{support}(\mathbf{v}) \subset \Omega_h(Y)$$

$$(7.12b) \quad (\text{div } \mathbf{v} - p)(\sigma) = 0 \quad \text{for all } \sigma \in Y$$

$$(7.12c) \quad \text{div } \mathbf{v}(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h \setminus Y.$$

If in addition  $\mathcal{T}_h$  is quasi-uniform the following bound holds

$$(7.13) \quad \|\nabla \mathbf{v}\|_{L^2(\Omega_h(Y))} \leq C \left(1 + D_r\right) \left(1 + \Upsilon(\mathcal{R}(r)) \rho(\mathcal{R}(r))\right) \|p\|_{L^2(\Omega_h(Y))},$$

where  $\Upsilon(\mathcal{R}(r))^2 := \max_{z \in Y} \left(\sum_{y \in A(z)} |D(y)|\right)$ . We recall that  $D_r$  is given in (3.2).

*Proof.* For every  $z \in Y$  with  $z \neq r$  there is a unique path  $P_z \subset Y$  that connects  $z$  to  $r$ . By Lemma 14 we can find  $\mathbf{v}_z \in \mathbf{V}_{h,0}^3$  such that

$$(7.14a) \quad \text{support}(\mathbf{v}_z) \subset \Omega_h(P_z),$$

$$(7.14b) \quad (\text{div } \mathbf{v}_z - p)(z) = 0$$

$$(7.14c) \quad \text{div } \mathbf{v}_z(\sigma) = 0 \quad \text{for all } \sigma \in \mathcal{S}_h, \sigma \neq z, \sigma \neq r,$$

$$(7.14d) \quad \|\nabla \mathbf{v}_z\|_{L^\infty(\Omega_h(P_z))} \leq C \rho(P_z) \max_{T \in \mathcal{T}_h(z)} |p|_T(z),$$

where  $C$  only depends on the shape regularity of the mesh.

Note that from (7.14d) and inverse estimates we get that for any  $y \in A(z)$ , assuming  $h_y \leq C h_z$  (which holds if the mesh is quasi-uniform) we have

$$(7.15) \quad \|\nabla \mathbf{v}_z\|_{L^2(\Omega_h(y))} \leq C \rho(P_z) \|p\|_{L^2(\Omega_h(z))}.$$

Then, we take

$$\mathbf{w} = \sum_{z \in Y, z \neq r} \mathbf{v}_z.$$

We then have the following properties of  $\mathbf{w}$

$$(7.16) \quad (\text{div } \mathbf{w} - p)(y) = 0 \quad \text{for all } y \in Y \setminus \{r\}$$

$$(7.17) \quad \text{support}(\mathbf{w}) \subset \Omega_h(Y)$$

$$(7.18) \quad \text{div } \mathbf{w}(\sigma) = 0, \quad \sigma \notin Y.$$

Since  $\mathbf{r} \in \mathcal{L}_h$  (using Definition 1) we can find  $\mathbf{r} \in \mathbf{V}_{h,00}^3(r)$  so that

$$(7.19) \quad (\operatorname{div} \mathbf{r} + \operatorname{div} \mathbf{w} - p)(r) = 0$$

$$(7.20) \quad \|\nabla \mathbf{r}\|_{L^\infty(\Omega_h(r))} \leq D_r \max_{T \in \mathcal{T}_h(r)} |(\operatorname{div} \mathbf{w} - p)|_T(r)|.$$

In this case, using inverse estimates, we can show that

$$(7.21) \quad \|\nabla \mathbf{r}\|_{L^2(\Omega_h(r))} \leq C D_r (\|\nabla \mathbf{w}\|_{L^2(\Omega_h(r))} + \|p\|_{L^2(\Omega_h(r))}).$$

We next set  $\mathbf{v} = \mathbf{w} + \mathbf{r}$  and we see that  $\mathbf{v} \in \mathbf{V}_{h,0}^3$  and (7.12a), (7.12b) and (7.12c) hold.

To get the bound (7.13) we assume that  $\mathcal{T}_h$  is quasi-uniform. Using the triangle inequality and (7.21)

$$\begin{aligned} \|\nabla \mathbf{v}\|_{L^2(\Omega_h(Y))} &\leq \|\nabla \mathbf{w}\|_{L^2(\Omega_h(Y))} + \|\nabla \mathbf{r}\|_{L^2(\Omega_h(Y))} \\ &\leq (1 + C D_r) (\|\nabla \mathbf{w}\|_{L^2(\Omega_h(Y))} + \|p\|_{L^2(\Omega_h(r))}). \end{aligned}$$

Next, we estimate  $\mathbf{w}$ :

$$\begin{aligned} \|\nabla \mathbf{w}\|_{L^2(\Omega_h(Y))}^2 &\leq \sum_{y \in Y} \|\nabla \mathbf{w}\|_{L^2(\Omega_h(y))}^2 \\ &\leq \sum_{y \in Y} \left\| \sum_{z \in \mathbf{D}(y)} \nabla \mathbf{v}_z \right\|_{L^2(\Omega_h(y))}^2 && \text{by (7.14a)} \\ &\leq \sum_{y \in Y} \left( \sum_{z \in \mathbf{D}(y)} \|\nabla \mathbf{v}_z\|_{L^2(\Omega_h(y))} \right)^2 && \text{by the triangle inequality} \\ &\leq \sum_{y \in Y} |\mathbf{D}(y)| \sum_{z \in \mathbf{D}(y)} \|\nabla \mathbf{v}_z\|_{L^2(\Omega_h(y))}^2 && \text{by Hölder's inequality} \\ &\leq C \rho(\mathcal{R}(r))^2 \sum_{y \in Y} |\mathbf{D}(y)| \sum_{z \in \mathbf{D}(y)} \|p\|_{L^2(\Omega_h(z))}^2 && \text{by (7.15)} \\ &\leq C \rho(\mathcal{R}(r))^2 \sum_{z \in Y} \|p\|_{L^2(\Omega_h(z))}^2 \sum_{y \in \mathbf{A}(z)} |\mathbf{D}(y)| && \text{interchange summation} \\ &\leq 3 C \rho(\mathcal{R}(r))^2 \Upsilon(\mathcal{R}(r))^2 \|p\|_{L^2(\Omega_h(Y))}^2 && \text{by definition of } \Upsilon(\mathcal{R}(r)). \end{aligned}$$

Taking square roots we get

$$\|\nabla \mathbf{w}\|_{L^2(\Omega_h(Y))} \leq C \rho(\mathcal{R}(r)) \Upsilon(\mathcal{R}(r)) \|p\|_{L^2(\Omega_h(Y))}.$$

The result now follows. □

The next result follows immediately from the previous lemma.

**Theorem 2.** *Suppose that we have  $\{r_1, \dots, r_t\} \in \mathcal{L}_h$  and corresponding acceptable trees  $\{\mathcal{R}(r_1), \mathcal{R}(r_2), \dots, \mathcal{R}(r_t)\}$ . If  $\mathcal{R}(r_i) = Z_i \times E_i$  we require that  $\bigcup_{i=1}^t Z_i = S_h$  and  $Z_i \cap Z_j = \emptyset$  for  $i \neq j$ . Then, for any  $p \in Q_h^2$  there exist  $\mathbf{v} \in \mathbf{V}_{h,0}^3$  such that*

$$(\operatorname{div} \mathbf{v} - p)(\sigma) = 0 \quad \text{for all } \sigma \in S_h$$



and if  $\mathcal{T}_h$  is quasi-uniform

$$\|\nabla \mathbf{v}\|_{L^2(\Omega)} \leq C(1 + \bar{D})(1 + \bar{\Upsilon}\bar{\rho})\|p\|_{L^2(\Omega)}.$$

where

$$(7.22) \quad \bar{D} = \max_{1 \leq i \leq t} D_{r_i}, \quad \bar{\Upsilon} = \max_{1 \leq i \leq t} \Upsilon(\mathcal{R}(r_i)), \quad \bar{\rho} = \max_{1 \leq i \leq t} \rho(\mathcal{R}(r_i)).$$

Using Theorem 2 and Lemma 2 we have that the inf-sup condition (2.3) holds for  $Q_h^2 \times \mathbf{V}_h^3$ .

**Corollary 2.** Assuming the hypothesis Theorem 2 then the estimate (2.3) holds for  $Q_h^2 \times \mathbf{V}_h^3$  with constants given by Lemma 2 and Theorem 2.

From Theorem 2 and Lemma 2 we deduce that if the hypotheses of Theorem 2 hold then  $Q_h^2 \times \mathbf{V}_h^3$  is inf-sup stable (see Corollary 2) and in fact that  $\operatorname{div} \mathbf{V}_h^3 = Q_h^2$ . If we do not care about the inf-sup constant in (2.3) and only care if  $\operatorname{div} \mathbf{V}_h^3 = Q_h^2$  then we can give weaker conditions. Inspecting the proof of Lemma 15 (and using Lemma 2), we can show the following.

**Theorem 3.** . If for every  $z \in \mathcal{S}_h$  there is exists an  $r \in \mathcal{L}_h$  such that there is an acceptable path  $P_{z,r}$  between them, then  $\operatorname{div} \mathbf{V}_h^3 = Q_h^2$ .

## 8. RELATIONSHIP TO QIN'S RESULT

Results concerning the pair of spaces  $V_h^k, Q_h^{k-1}$  were given in [11] for  $k \leq 3$ . Here we review the case  $k = 3$ . For the case  $k = 1$ , also see [12], and for the case  $k = 2$ , see [3]. Qin considered the mesh in Figure 10, which is called a Type I triangulation [8]. Of course, the upper-left and lower-right triangles are problematic, since the pressures will vanish at the corner vertices there. But more interestingly, Qin found an additional spurious pressure mode as indicated in Figure 7(a). We can relate this to the quantities  $\mathcal{D}_i$  in (3.23) by computing them for this mesh, as indicated in Figure 7(b). There are only two angles in this mesh,  $\pi/4$  and  $\pi/2$ , and  $\cot(\pi/4) = 1$  and  $\cot(\pi/2) = 0$ . Similarly, the edge lengths are  $L$  and  $L\sqrt{2}$ , for some  $L$ . Thus the quantities  $d_{j0}$  in (3.21) are of the form  $\pm A$  where  $A = 1/2L^2$  for the  $\pi/4$  angles, and 0 for the  $\pi/2$  angles, as indicated in Figure 7(b). Computing the alternating sum of terms in (3.21), we get

$$(8.1) \quad \mathcal{D}_0 = \sum_{j=1}^6 (-1)^j d_{j0} = A - (-A) + 0 - A + (-A) = 0.$$


Thus condition (3.23) is violated for  $i = 0$  for all the interior vertices in Figure 10.

Now let us compute  $\mathcal{D}_i$  for  $i = 1, 2$ . First, we note that the sequence of vertices  $y_k$  for a fixed interior vertex  $z$

$$\begin{aligned} y_1 &= L(1, 0) + z, \quad y_2 = L(1, 1) + z, \quad y_3 = L(0, 1) + z, \\ y_4 &= L(-1, 0) + z, \quad y_5 = L(-1, -1) + z, \quad y_6 = L(0, -1) + z. \end{aligned}$$

Thus,

$$(8.2) \quad \sum_{j=1}^6 (-1)^j y_j = L(-1 + 1 - 1 + 1, 1 - 1 + 1 - 1) + z \sum_{j=1}^6 (-1)^j = \mathbf{0}.$$



**Figure 7.** (a) A global spurious pressure mode on the mesh in Figure 10. (b) Computation of (8.1).

Letting  $t_j = \cot(\theta_j) + \cot(\theta_{j+1})$ , we easily can show

$$t_1 = 1, t_2 = 2, t_3 = 1, t_4 = 1, t_5 = 2, t_6 = 1.$$

Also, we have the sequence of values  $|e_j|^{-2}$  are

$$|e_1|^{-2} = \frac{1}{L^2}, |e_2|^{-2} = \frac{1}{2L^2}, |e_3|^{-2} = \frac{1}{L^2}, |e_4|^{-2} = \frac{1}{L^2}, |e_5|^{-2} = \frac{1}{2L^2}, |e_6|^{-2} = \frac{1}{L^2}.$$

Hence,  $\frac{(\cot(\theta_j) + \cot(\theta_{j+1}))}{|e_j|^2} = \frac{1}{L^2}$  for all  $j$ . Also,  $|T_j| = |T_1|$  for all  $j$ . Hence, using (3.30), (8.1), and (8.2) we have for  $i = 1, 2$

$$\mathcal{D}_i = \left( \frac{4}{L^2} - \frac{2}{|T_1|} \right) \sum_{j=1}^n (-1)^j y_j \cdot \mathbf{E}_i^\perp = 0.$$

Thus condition (3.23) is violated for all  $i = 0, 1, 2$  for all interior vertices in Figure 10. This suggests that the constraint (3.23) maybe required for inf-sup stability.

## 9. STRANG'S DIMENSION

For simplicity, let us assume that  $\Omega$  is simply connected. Then the space  $\mathbf{Z}_h^3 := \{\mathbf{v} \in \mathbf{V}_h^3 : \operatorname{div} \mathbf{v} = 0\}$  is the curl of the space  $S_h^4$  of  $C^1$  piecewise quartics on the same mesh, where the quartics must vanish to second order on the boundary. The dimension of the space  $\widehat{S}_h^4$  of  $C^1$  piecewise quartics, without boundary conditions, is known [10, 1] for a broad class of triangulations  $\mathcal{T}_h$  to be


$$(9.1) \quad \dim \widehat{S}_h^4 = E + 4V - V_0 + \sigma_i,$$

where  $T$  is the number of triangles in  $\mathcal{T}_h$ ,  $E$  (resp.,  $E_0$ ) is the number of edges (resp., interior edges) in  $\mathcal{T}_h$ ,  $V$  (resp.,  $V_0$ ) is the number of vertices (resp., interior vertices) in  $\mathcal{T}_h$ , and  $\sigma_i$  is the number of singular interior vertices in  $\mathcal{T}_h$ .

The dimension formula (9.1) is essentially the one conjectured by Gil Strang [15], so we refer to this as the *Strang dimension* of  $\widehat{S}_h^4$ :

$$(9.2) \quad \mathbb{D}(\widehat{S}_h^4) = E + 4V - V_0 + \sigma_i.$$

For  $C^1$  piecewise polynomials of degree  $k \geq 5$ , the Strang dimension was confirmed using an explicit basis [9]. But the Strang dimension has not yet been confirmed for arbitrary meshes for



**Figure 8.** (a) Nodal basis for quatics. (b) Nodes that determine value and gradient on an edge.

$k \leq 4$ . However, what is known is that it provides a lower bound [10]

$$(9.3) \quad \dim \hat{S}_h^4 \geq \mathbb{D}(\hat{S}_h^4).$$

**9.1. Computing  $\dim S_h^4$ .** Now let us compute  $\dim S_h^4 = \dim \mathbf{Z}_h^3$  under the assumption that the inf-sup condition holds. The space  $\mathbf{V}_h^3$  can be described in terms of Lagrange nodes:

$$(9.4) \quad \dim \mathbf{V}_h^3 = 2(T + 2E_0 + V_0) = 2T + 4E_0 + 2V_0.$$

We have  $\nabla \cdot \mathbf{V}_h^3 \subset Q_h^2$ , where the latter space consists of mean-zero piecewise quadratics that satisfy the alternating condition  $A_h^z(\operatorname{div} \mathbf{v}) = 0$  at singular vertices, where  $A_h^z$  is defined in (2.2):

$$\dim Q_h^2 = 6T - 1 - \sigma.$$

Thus  $\dim \nabla \cdot \mathbf{V}_h^3 = \dim Q_h^2 - K$  for some integer  $K \geq 0$ , and  $\nabla \cdot \mathbf{V}_h^3 = Q_h^2$  if and only if  $K = 0$ . Since

$$\dim \mathbf{V}_h^3 = \dim(\operatorname{image} \nabla \cdot \mathbf{V}_h^3) + \dim \mathbf{Z}_h^3,$$

we have

$$(9.5) \quad \begin{aligned} \dim \mathbf{Z}_h^3 &= \dim \mathbf{V}_h^3 - \dim(\operatorname{image} \nabla \cdot \mathbf{V}_h^3) \\ &= 2T + 4E_0 + 2V_0 - 6T + 1 + \sigma + K \\ &= -4T + 4E_0 + 2V_0 + 1 + \sigma + K. \end{aligned}$$

We have  $3T = (E - E_0) + 2E_0 = E + E_0$ . Thus

$$(9.6) \quad \dim \mathbf{Z}_h^3 = -T - (E + E_0) + 4E_0 + 2V_0 + 1 + \sigma + K.$$

By Euler's formula,  $1 = T - E + V = T - E_0 + V_0 = 1$ . Thus  $E_0 - V_0 = T - 1$ , and

$$(9.7) \quad \begin{aligned} \dim S_h^4 = \dim \mathbf{Z}_h^3 &= (V_0 - E_0) - (E + E_0) + 4E_0 + 2V_0 + \sigma + K \\ &= 2E_0 - E + 3V_0 + \sigma + K. \end{aligned}$$

Technically, we actually have

$$\dim S_h^4 = \min\{0, 2E_0 - E + 3V_0 + \sigma + K\},$$

since the dimension can never be negative. There are cases where the number of boundary edges  $E - E_0$  (which is the same as the number of boundary vertices) is larger than  $E_0 + 3V_0 + \sigma$ . For a domain consisting of only two triangles,  $E_0 = 1$ ,  $V_0 = 0$ ,  $\sigma = 2$ , and  $E - E_0 = 4$ , so the formula in (9.7) gives a negative number if  $K = 0$ . From now on, we assume that  $E_0 - (E - E_0) + 3V_0 + \sigma \geq 0$  for  $\mathcal{T}_h$ .

**Theorem 4.** *Let  $k = 3$  and suppose that  $\mathcal{T}_h$  is any triangulation satisfying  $E_0 + 3V_0 + \sigma \geq E - E_0$ . Then*

$$\nabla \cdot \mathbf{V}_h^3 = Q_h^2$$

*if and only if*

$$\dim S_h^4 = 2E_0 - E + 3V_0 + \sigma.$$

*More generally,*

$$\dim \nabla \cdot \mathbf{V}_h^3 = \dim Q_h^2 - K$$

*if and only if*

$$\dim S_h^4 = 2E_0 - E + 3V_0 + \sigma + K.$$

**9.2. Computing  $\dim \hat{S}_h^4$ .** Now let us relate the spaces  $S_h^4$  and  $\hat{S}_h^4$  by imposing boundary conditions on  $\hat{S}_h^4$  to yield the space  $S_h^4$ . Using the approach pioneered by Strang [15], it is natural to conjecture that this involves simply imposing constraints on the boundary. For example, a  $C^1$  piecewise quartic that vanishes to second order on  $\partial\Omega$  must vanish to second order at each boundary vertex (3 constraints per boundary vertex). In addition, the value at one point on each boundary edge must vanish, as well as the normal derivative at two points on each boundary edge.

To see why this is the right number of constraints, we pick special nodal variables for quartics as indicated in Figure 8(a). These are

- (1) the value and gradient at each vertex,
- (2) the value at edge midpoints, and
- (3) the second-order cross derivatives  $\partial_{e_i}\partial_{e_j}$  evaluated at the vertex  $\nu_{ij}$  at the intersection of  $e_i$  and  $e_j$ , where the  $e_k$ 's are the edges of the triangle.

More precisely,  $\partial_{e_i}\phi(\nu_{ij})$  is defined as the directional derivative of  $\phi$  in the direction of  $e_i$  away from  $\nu_{ij}$ . These nodal variables are unisolvent for quartics, as follows. Vanishing of nodal variables of type (1) and (2) guarantee vanishing on each edge; these are the standard nodal variables for Hermite quartics. Thus a quartic  $q$  with these nodal values zero is of the form  $q = L_1L_2L_3L$  where the non-trivial linear functions  $L_i$  vanish on  $e_i$ . But

$$\partial_{e_i}\partial_{e_j}q(\nu_{ij}) = (\partial_{e_i}L_j)(\partial_{e_j}L_i)L_k(\nu_{ij})L(\nu_{ij}),$$


where  $\{i, j, k\} = \{1, 2, 3\}$  and  $L_k(\nu_{ij}) \neq 0$ . Thus vanishing of the nodal variables of type (3) implies that  $L \equiv 0$ .

Moreover, similar arguments show that the nodal variables for  $q$  associated with a boundary edge, as indicated in Figure 8(b), determine  $q$  to second order on that edge. Thus satisfaction of second-order boundary conditions is guaranteed by setting these nodal values to zero. This reduces the dimension by at most  $3(V - V_0) + 3(E - E_0) = 6(E - E_0)$ , since  $E - E_0 = V - V_0$ . We will show that these conditions have (at least) one redundancy at singular boundary vertices. Thus

$$\dim S_h^4 \geq \dim \hat{S}_h^4 - 6(E - E_0) + \sigma_b,$$

where  $\sigma_b$  is the number of singular boundary vertices. Therefore

$$(9.8) \quad \dim S_h^4 + 6(E - E_0) - \sigma_b \geq \dim \hat{S}_h^4 \geq \mathbb{D}(\hat{S}_h^4) = E + 4V - V_0 + \sigma_i.$$



**Figure 9.** (a) Singular boundary vertex where two triangles meet. (b) Singular boundary vertex where three triangles meet.

Assume now that  $\nabla \cdot \mathbf{V}_h^3 = Q_h^2$ . Using Theorem 4 and  $E - E_0 = V - V_0$ , we find

$$\begin{aligned}
 \dim S_h^4 + 6(E - E_0) - \sigma_b &= E_0 - (E - E_0) + 3V_0 + \sigma + 6(E - E_0) - \sigma_b \\
 (9.9) \quad &= E_0 + 3V_0 + \sigma_i + 5(E - E_0) = E_0 + 3V + \sigma_i + 2(E - E_0) \\
 &= E + 3V + \sigma_i + (E - E_0) = E + 4V + \sigma_i - V_0.
 \end{aligned}$$

Combining (9.8) and (9.9) proves the following result.

**Theorem 5.** *Let  $k = 3$  and suppose that  $\mathcal{T}_h$  is a triangulation satisfying  $E_0 + 3V_0 + \sigma \geq E - E_0$  and suppose that  $\nabla \cdot \mathbf{V}_h^3 = Q_h^2$  on this triangulation. Then, the Strang dimension (9.1) is valid for  $\hat{S}_h^4$ ,  $\dim \hat{S}_h^4 = \mathbb{D}(\hat{S}_h^4)$ . Moreover, equality holds in (9.8), so the  $6(E - E_0) - \sigma_b$  constraints are nonredundant.*

To complete the proof of Theorem 5, we need to verify the redundancy of constraints at singular boundary vertices. This occurs because the second-order cross derivatives  $\partial_{e_i} \partial_{e_j}$  are linearly dependent at singular boundary vertices. For the case of a triangle with two boundary edges  $e_1$  and  $e_2$ , the vanishing of the nodal variables of type (1) and (2) on  $e_1$  and  $e_2$  already imply vanishing on both edges, so necessarily  $\partial_{e_1} \partial_{e_2}$  is already zero.

For the case where two triangles meet at a singular boundary vertex  $\nu$ , see Figure 9(a). Then  $e_1$  and  $e_3$  are parallel, and thus

$$(9.10) \quad \partial_{e_1} \partial_{e_2} \phi(\nu) = -\partial_{e_3} \partial_{e_2} \phi(\nu)$$

for any  $C^1$  piecewise quartic  $\phi$ . Thus setting one of them to zero sets the other; they are redundant.


For the case where three triangles meet at a singular boundary vertex  $\nu$ , see Figure 9(b). Equation (9.10) still holds, and in addition  $e_2$  and  $e_4$  are parallel, and thus

$$\partial_{e_3} \partial_{e_2} \phi(\nu) = -\partial_{e_3} \partial_{e_4} \phi(\nu)$$

for any  $C^1$  piecewise quartic  $\phi$ . Thus

$$\partial_{e_1} \partial_{e_2} \phi(\nu) = \partial_{e_3} \partial_{e_4} \phi(\nu),$$

and setting one of them to zero sets the other; they are redundant. This completes the proof of Theorem 5.



**Figure 10.** The Type I regular mesh studied in [11, Chapter 6]

Since Theorem 2 allows us to prove the inf-sup condition for quite general meshes, this confirms the Strang dimension (9.1) for  $\widehat{S}_h^4$  is correct for such meshes. In particular, if the hypothesis of Theorem 3 holds then (9.1) is correct.

**9.3. Connection to Qin's results.** There is a connection between Qin's results and dimension counting. Qin finds a spurious mode that suggests that  $\operatorname{div} \mathbf{V}_h^3 \neq Q_h^2$  on the right-triangle mesh in Figure 10. We conclude that the dimension of the space  $S_h^4$  of  $C^1$  quartics satisfying second-order boundary conditions on this mesh is at least one larger than the dimension for this space given in Theorem 4:  $K \geq 1$ . On the other hand, it is well known [8, 10] that the Strang dimension (9.1) is correct on Type I triangulations without boundary conditions. In view of (9.8), there is a further redundancy in the constraints enforcing boundary conditions. Unfortunately, the dimension of splines in two dimensions satisfying boundary conditions has had only limited study [6, 2] so far.

## REFERENCES

- [1] P. Alfeld, B. Piper, and L. L. Schumaker. An explicit basis for  $C^1$  quartic bivariate splines. *SIAM Journal on Numerical Analysis*, 24(4):891–911, 1987.
- [2] L. Anbo. On the dimension of spaces of pp functions with boundary conditions. *Approximation Theory and its Applications*, 5(4):19–29, 1989.
- [3] D. N. Arnold and J. Qin. Quadratic velocity/linear pressure stokes elements. *Advances in computer methods for partial differential equations*, 7:28–34, 1992.
- [4] C. Bernardi and G. Raugel. Analysis of some finite elements for the Stokes problem. *Mathematics of Computation*, pages 71–79, 1985.
- [5] S. C. Brenner and L. R. Scott. *The mathematical theory of finite element methods*, volume 15. Springer Science & Business Media, third edition, 2008.
- [6] C. Chui and L. Schumaker. On spaces of piecewise polynomials with boundary conditions. II. Type-1 triangulations. In Z. Ditzian, editor, *Second Edmonton Conference on Approximation Theory*, CMS Conf. Proc., 3. Amer. Math. Soc., Providence, R.I., 1983.
- [7] J. Guzmán and L. R. Scott. The Scott-Vogelius finite elements revisited. *Mathematics of Computation*, submitted, 2017.
- [8] M.-J. Lai and L. L. Schumaker. *Spline functions on triangulations*. Number 110 in Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2007.
- [9] J. Morgan and L. R. Scott. A nodal basis for  $C^1$  piecewise polynomials of degree  $n \geq 5$ . *Math. Comp.*, 29:736–740, 1975.
- [10] J. Morgan and L. R. Scott. The dimension of the space of  $C^1$  piecewise-polynomials. Research Report UH/MD 78, Dept. Math., Univ. Houston, 1990.

- [11] J. Qin. *On the convergence of some low order mixed finite elements for incompressible fluids*. PhD thesis, Penn State, 1994.
- [12] J. Qin and S. Zhang. Stability and approximability of the  $P1-P0$  element for Stokes equations. *International journal for numerical methods in fluids*, 54(5):497–515, 2007.
- [13] L. Scott and M. Vogelius. Norm estimates for a maximal right inverse of the divergence operator in spaces of piecewise polynomials. *RAIRO-Modélisation mathématique et analyse numérique*, 19(1):111–143, 1985.
- [14] L. R. Scott and M. Vogelius. Conforming finite element methods for incompressible and nearly incompressible continua. In *Large Scale Computations in Fluid Mechanics*, B. E. Engquist, et al., eds., volume 22 (Part 2), pages 221–244. Providence: AMS, 1985.
- [15] G. Strang. Piecewise polynomials and the finite element method. *Bulletin of the American Mathematical Society*, 79(6):1128–1137, 1973.
- [16] M. Vogelius. A right-inverse for the divergence operator in spaces of piecewise polynomials. *Numerische Mathematik*, 41(1):19–37, 1983.

<sup>†</sup> DIVISION OF APPLIED MATHEMATICS, BROWN UNIVERSITY, PROVIDENCE, RI 02912, USA  
*E-mail address:* johnny\_guzman@brown.edu

<sup>‡</sup>DEPARTMENTS OF COMPUTER SCIENCE AND MATHEMATICS, COMMITTEE ON COMPUTATIONAL AND APPLIED MATHEMATICS, UNIVERSITY OF CHICAGO, CHICAGO IL 60637, USA  
*E-mail address:* ridg@uchicago.edu