

# On the Calculus of Functions with the Cantor-Tartan Support

Alireza Khalili Golmankhaneh <sup>a\*</sup>

<sup>a</sup> Department of Physics, Urmia Branch, Islamic Azad University, Urmia, Iran

<sup>\*</sup>*E-mail address:* a.khalili@iaurmia.ac.ir

## Abstract

In this manuscript, integral and derivative of the functions with Cantor-Tartan spaces is defined. The generalization of standard calculus which is called  $F^\alpha$ -calculus utilized to obtain the integral and derivative of the functions on the Cantor-Tartan with different dimensions. Differential equation involving the new derivatives are solved. The illustrative examples are used to present the details. *Keywords:*

$F^\alpha$ -calculus; Staircase function ; Cantor-Tartan support; Fractional differential equation

MSC[2010]: 81Q35; 28A80;

The fractal shapes and objects are seen in the nature, e.g., clouds, mountains, coastlines, human body, and etc. The geometry of the fractals were studied [26]. The analysis on fractals were established, using different methods, such as fractional calculus, probability theory, measure theory, fractional spaces, and time scale theory by many researchers and found many applications [12, 25, 25, 6, 2, 7, 1, 31, 35, 36, 8]. The fractional derivatives have non-local property which are suitable to model the process with the memory effect, non-conservative systems [22, 23, 30, 39, 41, 24, 27, 28, 33, 18]. The fractional calculus, which involves derivatives with arbitrary orders, has applied on the process with fractals structures [37, 29, 9]. The anomalous diffusion on fractals was formulated which included sub- and super diffusion in view of different random walks [40, 11, 38]. Fractal antennas are small but have wide-band radiations which make them useful in microwave communications [21, 10]. Laminar flow of a fractal fluid in a cylindrical tube was studied

using the homogeneous flow in a fractional dimensional space [4]. On the Cantor cubes, the Maxwell's equation were obtained and, as an application, the electric field due to Cantor dust was obtained [5]. Recently,  $F^\alpha$ -calculus which is algorithmic was suggest and applied for modeling some physical process [32, 34]. As a pursuit of these researches,  $F^\alpha$ -calculus is generalized and utilized in optics and mechanics [13, 14, 15, 3]. The non-local integrals and derivatives are defined on the Cantor set which are used to model the fractal ideal solids and the fractal ideal fluids [16, 17].

This manuscript is built as follows:

In Section 1 we set up the notation and terminology of  $F^\alpha$ -calculus on the fractals that imbedded in  $\mathbb{R}^2$  [19, 20]. In Section 2 we study some examples on the Cantor-Tartan using suggested definitions. Section 3 is devoted to the conclusion.

## 1 Terminology and notations

In the section, we review and define the basic tools we need in our work.

Fractals are the sets with the self-similar properties such that their fractal dimension exceeds from theirs topological dimension. The calculus on the Cantor-Tartan  $\mathfrak{F} = F \cup F \subset \mathbb{R}^2$  where  $F = C \times \mathbb{R}$ ,  $C$  is Cantor set and  $\mathbb{R}$  is real line [12]. Let  $\mathfrak{F}$  be Cantor-Tartan that is subset of  $I = [a, b] \times [c, d]$ ,  $a, b, c, d \in \mathbb{R}$  (Real-line). We sketch in Figures [1] the Cantor-Tartan with different fractal box or similar dimensions. The Cantor-Tartan is established by Cartesian product of Cantor set and real line (F) and union of them. Here, we consider finite iteration which is approximation of fractals spaces.




Figure 1: The Cantor-Tartan with different dimensions are plotted.

The flag function for the Cantor-Tartan  $\mathfrak{F} = F \cup F \subset \mathbb{R}^2$ , which is denoted by  $\Theta(\mathfrak{F}, I)$ , is defined as follows

$$\Theta(\mathfrak{F}, I) = \begin{cases} 1, & \text{if } \mathfrak{F} \cap I \neq \emptyset, \\ 0, & \text{otherwise.} \end{cases} \quad (1)$$

The subdivision  $P_{[a,b]}$  is

$$P_{[a,b] \times [c,d]} = \{x_0 = a, x_1, x_2, \dots, x_n = b\} \times \{y_0 = c, y_1, y_2, \dots, y_m = d\}, \quad (2)$$

where  $\times$  denotes the Cartesian product.

The  $\gamma^\zeta(\mathfrak{F}, a, b, c, d)$  is defined

$$\gamma^\zeta(\mathfrak{F}, a, b, c, d) = \lim_{\delta \rightarrow 0} \inf_{P_{[a,b] \times [c,d]} : |P| \leq \delta} \sum_{j=1}^n \sum_{i=1}^n \frac{(x_i - x_{i-1})^\alpha}{\Gamma(\alpha + 1)} \frac{(y_j - y_{j-1})^\beta}{\Gamma(\beta + 1)} \Theta(\mathfrak{F}, [x_{i-1}, x_i]) \Theta(\mathfrak{F}, [y_{j-1}, y_j]), \quad (3)$$

where  $\zeta = \max\{\alpha + \beta\}$ ,  $\alpha = \beta = 1 + \text{Dim}(C)$  and  $|P|$  is

$$|P| = \max_{1 \leq i \leq n, 1 \leq j \leq n} (x_i - x_{i-1}) \times (y_j - y_{j-1}). \quad (4)$$

The integral staircase function  $S_{\mathfrak{F}}^\zeta(x, y)$  for Cantor-Tartan  $\mathfrak{F}$  is defined

$$S_F^\zeta(x, y) = \begin{cases} \gamma^\zeta(F, a_0, c_0, x, y), & \text{if, } x \geq a_0, y \geq c_0; \\ -\gamma^\zeta(F, a_0, c_0, x, y), & \text{otherwise,} \end{cases} \quad (5)$$

where  $a_0, c_0$  are arbitrary real numbers.




Figure 2: Staircase function corresponding to Cantor-Tartan with different dimensions are presented.

The integral staircase function for the different Cantor-Tartan with the different dimension are plotted in Figure [2].

The  $\zeta$ -dimension of  $\mathfrak{F} \cap [a, b] \times [c, d]$  is given by

$$\begin{aligned} \dim_\zeta(\mathfrak{F} \cap [a, b] \times [c, d]) &= \inf\{\zeta : \gamma^\zeta(\mathfrak{F}, a, c, b, d) = 0\}, \\ &= \sup\{\zeta : \gamma^\zeta(\mathfrak{F}, a, c, b, d) = \infty\}. \end{aligned} \quad (6)$$

A point  $(x, y)$  is called a point of change of a function  $f(x, y)$  if it is not constant over any open set  $[a, b] \times [c, d]$  involving  $(x, y)$ . The set of all points

of change of  $f(x, y)$  are indicated by  $Sch f$ .

The  $Sch(S_{\mathfrak{F}}^{\zeta}(x, y))$  is called  $\zeta$ -perfect if  $S_{\mathfrak{F}}^{\zeta}(x, y)$  is finite for all  $(x, y) \in \mathfrak{R}$ .

Let  $f(x, y)$  be a bounded function on  $\mathfrak{F}$  then we define

$$\begin{aligned} M[f, \mathfrak{F}, I] &= \sup_{(x, y) \in \mathfrak{F} \cap I} f(x, y), \quad \text{if } \mathfrak{F} \cap I \neq \emptyset. \\ &= 0, \quad \text{otherwise,} \end{aligned} \quad (7)$$

and similarly

$$\begin{aligned} m[f, \mathfrak{F}, I] &= \inf_{(x, y) \in \mathfrak{F} \cap I} f(x, y), \quad \text{if } \mathfrak{F} \cap I \neq \emptyset. \\ &= 0, \quad \text{otherwise.} \end{aligned} \quad (8)$$

Now, upper  $U^{\zeta}$ -sum and lower  $L^{\zeta}$ -sum for the function  $f(x, y)$  over the subdivision  $P$  are given as follows

$$U^{\zeta}[f, \mathfrak{F}, P] = \sum_{i=1}^n M[f, \mathfrak{F}, [(x_{i-1}, y_{i-1}), (x_i, y_i)]] (S_{\mathfrak{F}}^{\zeta}(x_i, y_i) - S_{\mathfrak{F}}^{\zeta}(x_{i-1}, y_{i-1})), \quad (9)$$

and

$$L^{\zeta}[f, \mathfrak{F}, P] = \sum_{i=1}^n m[f, \mathfrak{F}, [(x_{i-1}, y_{i-1}), (x_i, y_i)]] (S_{\mathfrak{F}}^{\zeta}(x_i, y_i) - S_{\mathfrak{F}}^{\zeta}(x_{i-1}, y_{i-1})). \quad (10)$$

The  $f(x, y)$  is called  $F^{\zeta}$ -integrable on Cantor-Tartan  $\mathfrak{F}$  if we have

$$\int_{(a, c)}^{(b, d)} f(x, y) d_F^{\alpha} x d_F^{\beta} y = \sup_{P_{[a, b]}} L^{\zeta}[f, \mathfrak{F}, P] = \overline{\int_{(a, c)}^{(b, d)} f(x, y) d_F^{\alpha} x d_F^{\beta} y} = \inf_{P_{[a, b]}} U^{\zeta}[f, \mathfrak{F}, P]. \quad (11)$$

The  $F^{\zeta}$ -integral is denoted by  $\int_{(a, b)}^{(b, d)} f(x, y) d_F^{\alpha} x d_F^{\beta} y$ .

Let  $\mathfrak{F}$  be  $\zeta$ -perfect set then we define  $F^{\alpha}$ -partial derivative of  $f(x, y)$  respect to  $x$  as

$${}_x D_{\mathfrak{F}}^{\alpha} f(x, y) = \begin{cases} \mathfrak{F} - \lim_{(x', y) \rightarrow (x, y)} \frac{f(x', y) - f(x, y)}{S_{\mathfrak{F}}^{\zeta}(x', y) - S_{\mathfrak{F}}^{\zeta}(x, y)}, & \text{if } (x, y) \in \mathfrak{F}, \\ 0, & \text{otherwise.} \end{cases} \quad (12)$$

if the limit exists.

## 2 The functions with Cantor-Tartan support

The task is now to apply definitions on examples. In this section, we give some examples to show more details.

## 2.1 Example 1.

Let us consider a function with Cantor-Tartan support with the different fractal dimensions

$$f(x, y) = \sin(\chi_F^\alpha x) \sin(\chi_F^\beta y), \quad (x, y) \in \mathfrak{F}, \quad (13)$$

where  $\chi_F^\alpha, \chi_F^\beta$  are characteristic function for fractal sets  $F$  [17, 32]. The graph of the  $f(x, y)$  is shown in Figures [3].




Figure 3: We plot the graph of function  $\sin(\chi_F^\alpha x) \sin(\chi_F^\beta y)$  with the different Cantor-Tartan support.

The fractal integral of  $f(x, y)$  on Cantor-Tartan  $\mathfrak{F} \subset [0, 1] \times [0, 1]$  is as follows:

$$\begin{aligned} g(x, y) \Big|_{(x=y=1)} &= \int_0^x \int_0^y \sin(\chi_F^\alpha x) \sin(\chi_F^\beta y) d_F^\alpha x' d_F^\beta y' \Big|_{(x=y=1)} = \\ &= \int_0^1 \cos(S_F^\beta(y')) \sin(\chi_F^\alpha x) \Big|_0^1 d_F^\alpha x' = \\ &= \int_0^1 [\cos(S_F^\beta(1)) \sin(\chi_F^\alpha x) - \cos(S_F^\beta(0)) \sin(\chi_F^\alpha x)] d_F^\alpha x' = \\ &= \cos(\Gamma(1 + \beta)) \cos(S_F^\alpha(x')) - \cos(S_F^\alpha(x')) \Big|_{S_F^\alpha(0)}^{S_F^\alpha(1)} = \\ &= \cos(\Gamma(1 + \beta)) \cos(\Gamma(1 + \alpha)) - \cos(\Gamma(1 + \alpha)) - \cos(\Gamma(1 + \beta)) + 1 \\ &= \begin{cases} 0.13, & \alpha = \beta = 0.6 \text{ With support Cantor-Tartan dimesion } 1.2 \\ 0.16, & \alpha = \beta = 0.8 \text{ With support Cantor-Tartan dimesion } 1.6 \end{cases} \end{aligned} \quad (14)$$

here  $S_F^\alpha(1) = \Gamma(1 + \alpha)$ ,  $S_F^\beta(1) = \Gamma(1 + \beta)$ ,  $S_F^\alpha(0) = S_F^\beta(0) = 0$ .




Figure 4: Fractal integral of  $\sin(\chi_F^\alpha x) \sin(\chi_F^\beta y)$  on the different Cantor-Tartan are sketched .

In Figure [4] we sketch the  $g(x, y)$  which is called fractal integral of  $f(x, y)$ .

## 2.2 Example 2.

Consider a function with Cantor-Tartan support as follows

$$f(x, y) = \sin(\chi_F^\alpha x + \chi_F^\beta y), \quad (x, y) \in \mathfrak{F} \quad (15)$$




Figure 5: We plot the graph of function  $\sin(\chi_F^\alpha x + \chi_F^\beta y)$  with the different Cantor-Tartan support.

We present in Figure [5] graph of Eq. (15). The fractal integral of Eq.

(15) is as follows

$$\begin{aligned}
g(x, y) \Big|_{(x=y=1)} &= \int_0^x \int_0^y \sin(\chi_F^\alpha x + \chi_F^\beta y) d_F^\alpha x' d_F^\beta y' \Big|_{(x=y=1)} = \\
&\int_0^1 -\cos(\chi_F^\alpha x + S_F^\beta(y)) \Big|_0^1 d_F^\alpha x' = \\
&\int_0^1 [-\cos(\chi_F^\alpha x + S_F^\beta(1)) + \cos(\chi_F^\alpha x)] d_F^\alpha x' = \\
&-\sin(S_F^\alpha(x') + \Gamma(1 + \beta)) + \sin(S_F^\alpha(x')) \Big|_0^1 = \\
&= -\sin(\Gamma(1 + \alpha) + \Gamma(1 + \beta)) + \sin(\Gamma(1 + \alpha)) + \sin(\Gamma(1 + \beta)) \\
&= \begin{cases} 0.581, & \alpha = \beta = 0.6 \text{ With support Cantor-Tartan dimension } 1.2 \\ 0.64, & \alpha = \beta = 0.8 \text{ With support Cantor-Tartan dimension } 1.6 \end{cases}
\end{aligned} \tag{16}$$

where  $S_F^\alpha(1) = \Gamma(1 + \alpha)$ ,  $S_F^\beta(1) = \Gamma(1 + \beta)$ ,  $S_F^\alpha(0) = S_F^\beta(0) = 0$ .




Figure 6: Fractal integral of  $\sin(\chi_F^\alpha x + \chi_F^\beta y)$  on different Cantor-Tartan are shown.

In Figure [6] we plot fractal integral of  $\sin(\chi_F^\alpha x + \chi_F^\beta y)$  versus to different Cantor-Tartan support with different dimensions.

### 2.3 Example 3.

Consider a function with the Cantor-Tartan as follows

$$f(x, y) = S_F^\alpha(x)^2 + S_F^\beta(y)^2. \tag{17}$$

The fractal partial derivatives of  $f(x, y)$  respect to  $x$  and  $y$  are

$${}^x D_F^\alpha f(x, y) = 2S_F^\alpha(x)\chi_F, \quad {}^y D_F^\beta f(x, y) = 2S_F^\beta(y)\chi_F. \tag{18}$$

In Figure [7] we plot Eq. (17) and it's partial derivative which is shown  ${}^x D_F^\alpha(x, y)$ .




Figure 7: We have plotted  $S_F^\alpha(x)^2 + S_F^\beta(y)^2$  and  ${}^x D_F^\alpha(x, y)$ .

**Remark:** One can obtain standard results by choosing  $\alpha = 1$ ,  $\beta = 1$ ,  $S = \Re$  which leads to  $S_F^\alpha(x) = x$ ,  $S_F^\beta(y) = y$ .

### 3 Conclusion

In this work, we define the local derivative and integral on Cantor-Tartan. The standard calculus can not be applied to integrate and differentiate for the function on this fractals. Therefore, we need a new type of calculus to calculate the physical properties and describe phenomena on fractals. As a result, the  $F^\alpha$ -calculus on Cantor-Tartan which has fractal dimension  $1 < \zeta < 2$  is given. More, we recall the standard calculus results which shows that suggested definitions are the generalization of standard calculus. Three illustrative examples were investigated and the corresponding graphs of the functions were drawn.

### References

- [1] Agarwal RP, Bohner M. Basic calculus on time scales and some of its applications. Results Math 1999; 35: 3-22.
- [2] Agarwal RP, Mahmoud RR, Saker SH, Tunc C. Acta Math Hung 2017; 152: 383-403.
- [3] Ashrafi S, Golmankhaneh A. K. Energy straggling function by  $F^\alpha$ -calculus. ASME, J Comput Nonlinear Dynam 2017, doi:10.1115/1.4035718.
- [4] Balankin AS, Mena B, Susarrey O, Samayoa D. Steady laminar flow of fractal fluids. Phys Lett A 2017; 381: 623-628.



- [5] Balankin AS, Mena B, Patino J, Morales D. Electromagnetic fields in fractal continua. *Phys Lett A* 2013; 377: 783-788.
- [6] Bohner M, Peterson A, *Dynamic Equations on Time Scales: An Introduction with Applications*. Birkhäuser, Boston 2001.
- [7] Bohner M, Wintz N. The linear quadratic tracker on time scales. *Int J Dyn Syst Differ Equ* 2011; 3: 423-447.
- [8] Brossard J, Carmona R. Can one hear the dimension of fractal?. *Comm Math Phys* 1986; 104: 103-122.
- [9] Butera S, Paola MD. A physically based connection between fractional calculus and fractal geometry. *Ann Phys-New York* 2014; 350: 146-158.
- [10] Cohen N. Fractal antenna applications in wireless telecommunications. *Electronics Industries Forum of New England, IEEE*, 1997.
- [11] Chen W, Sun HG, Zhang X, Koroak D. Anomalous diffusion modeling by fractal and fractional derivatives. *Comp Math Appl* 2010; 59: 1754-1758.
- [12] Falconer K. *Techniques in Fractal Geometry*. John Wiley and Sons, 1997.
- [13] Golmankhaneh AK, Baleanu D. Diffraction from fractal grating Cantor sets. *J Mod Optic* 2016; 63:1364-1369.
- [14] Golmankhaneh AK, Baleanu D. Fractal calculus involving Gauge function. *Commun Nonlinear Sci* 2016; 37: 125-130.
- [15] Golmankhaneh A K, Tunc C. On the Lipschitz condition in the fractal calculus. *Chaos Soliton Fract.* 2017; 95: 140-147.
- [16] Golmankhaneh AK, Baleanu D. New derivatives on the fractal subset of Real-line. *Entropy*, 2016; 18: 1-13.
- [17] Golmankhaneh AK, Baleanu D. Non-local Integrals and Derivatives on Fractal Sets with Applications. *Open Phys* 2016; 14: 542-548.
- [18] Golmankhaneh A K. *Investigations in Dynamics: With Focus on Fractional Dynamics*. LAP Lambert Academic Publishing, Saarbrücken, 2012.
- [19] Golmankhaneh AK. About Kepler's Third Law on fractal-time spaces. *Ain Shams Eng J* doi:/10.1016/j.asej.2017.06.005
- [20] Golmankhaneh A K. On the calculus of the parameterized fractal curves. *Turk J Phys* 2017; 41: 418-425.
- [21] Gianvittorio JP, Rahmat-Samii Y. Fractal antennas: A novel antenna miniaturization technique and applications. *IEEE Antennas Propag* 2002; 44: 20-36.

- [22] Herrmann R. Fractional calculus: an introduction for physicists. World Scientific, 2014.
- [23] Hilfer R ed. Applications of fractional calculus in physics. World Scientific, 2000.
- [24] Kilbas AA, Srivastava HH, Trujillo JJ. Theory and Applications of Fractional Differential Equations. Elsevier, The Netherlands, 2006.
- [25] Kigami J. Analysis on fractals. Volume 143 of Cambridge Tracts in Mathematics, Cambridge University Press, Cambridge, 2001.
- [26] Mandelbrot BB. The Fractal Geometry of Nature. Freeman and Company, 1977.
- [27] Magin RL. Fractional Calculus in Bioengineering. Begell House Publisher, Inc. Connecticut, 2006.
- [28] Malinowska AB, Torres DFM. Introduction to the fractional calculus of variations. Imperial College Press, London, 2012.
- [29] Nigmatullin RR, Zhang W, Gubaidullin I. Accurate relationships between fractals and fractional integrals: new approaches and evaluations. *Fract Calculus Appl Anal* 2017; 20: 1263-1280.
- [30] Nigmatullin RR, Evdokimov YK. The concept of fractal experiments: New possibilities in quantitative description of quasi-reproducible measurements. *Chaos Soliton Fract* 2016; 91: 319-328.
- [31] Naqvi QA, Fiaz MA. Electromagnetic behavior of a planar interface of non-integer dimensional spaces. *J Electromag Waves Appl* 2017; 31: 625-1637.
- [32] Parvate A, Gangal A D. Calculus on fractal subsets of real line II: Conjugacy with ordinary calculus. *Fractals* 2011; 19: 271-290.
- [33] Podlubny I. Fractional differential equations. Academic Press, New York, 1999.
- [34] Seema S, Gangal AD. Langevin Equation on Fractal Curves, *Fractals*. 2016; 24: 1650028.
- [35] Strichartz RS. Differential equations on fractals: a tutorial, Princeton University Press, 2006.
- [36] Tarasov VE, Fractional dynamics: applications of fractional calculus to dynamics of particles, fields and media, Springer Science, 2011.
- [37] Tatom FB. The relationship between fractional calculus and fractals. *Fractals* 1995; 3: 217-229.
- [38] Telcs A. The art of random walks. Springer, New York, 2006.

- [39] Uchaikin VV. Fractional Derivatives for Physicists and Engineers Vol. 1 Background and Theory. Vol 2. Application Springer, Berlin, 2013.
- [40] Von Kameke A, Huhn F, Fernández-García G, Muñuzuri, V. Pérez-Muñuzuri AP. Propagation of a chemical wave front in a quasi-two-dimensional superdiffusive flow. Phys Rev E 2010; 81: 066211.
- [41] Wu GC, Baleanu D, Xie HP, Chen FL. Chaos synchronization of fractional chaotic maps based on stability results. Physica A 2016; 460: 374-383.