

Open Problem Statement: Minimal Distortion Embeddings of Diversities in ℓ_1

David Bryant and Paul Tupper

March 14, 2022

1 Statement of Problem

A diversity is a pair (X, δ) where X is a set and δ is a function taking finite subsets of X to \mathbb{R} and satisfying the following axioms: [4]

- (i) $\delta(A) \geq 0$ for all $A \subset X$
- (ii) $\delta(A) = 0$ if and only if $|A| \leq 1$
- (iii) $\delta(A \cup B) \leq \delta(A \cup C) + \delta(B \cup C)$ whenever $C \neq \emptyset$.

Diversities are an extension of the concept of a metric space since for any diversity if we define $d(x, y) = \delta(\{x, y\})$ for $x, y \in X$ then (X, d) is a metric space [4]. Work on the mathematics of diversities has generated new theoretical results, applications, and unexpected links [7, 9, 11, 16, 15, 13, 12, 18].

Just as there is an ℓ_1 metric on \mathbb{R}^k there is an ℓ_1 diversity on \mathbb{R}^k defined as

$$\hat{\delta}(A) := \sum_{i=1}^k \max_{a, b \in A} |a_i - b_i|$$

for all finite subsets A of \mathbb{R}^k .

Let (X_1, δ_1) and (X_2, δ_2) be two diversities. Let ϕ be a map from X_1 to X_2 . We say ϕ has distortion c if there are $c_1, c_2 > 0$ such that $c = c_1 c_2$ and

$$\frac{1}{c_1} \delta_1(A) \leq \delta_2(\phi(A)) \leq c_2 \delta_1(A)$$

for all finite $A \subseteq X_1$ [5].

Problem: Find a bound $f(n)$ such that for any diversity (X, δ) with $|X| = n$ there is a map ϕ from (X, δ) into an ℓ_1 diversity $(\mathbb{R}^d, \hat{\delta})$ with distortion at most $f(n)$.

As we explain below, the best general bound we have so far is $\mathcal{O}(n)$, though this is likely far from optimal. We know that the bound cannot be better than $\Omega(\log n)$.

Our problem is the diversity analogue of a well-known problem in the theory of metric spaces: Find the minimal distortion embedding of a n -point metric space into $(\mathbb{R}^k, \|\cdot\|_1)$ for any k . Due to a result of Bourgain [3], it is known that the distortion bound in the metric spaces case is $\mathcal{O}(\log n)$ and that this can be attained with dimension $k = \mathcal{O}(\log n)$. This bound is tight [14].

We know that for the diversity case it is not possible to embed a general n -point diversities with distortion $\mathcal{O}(\log n)$ into ℓ_1 with dimension $k = \mathcal{O}(\log n)$ [5]. However, once we remove the restriction on the dimension, the $\mathcal{O}(\log n)$ distortion has not been ruled out yet.

The following argument, due to P. Wu, gives an upper bound for the worst distortion of $\mathcal{O}(n)$.

Theorem 1. *For all diversities (X, δ) where $|X| = n$ there is an embedding $\phi: X \rightarrow \ell_1^n$ with distortion at most n .*

Proof. Let (X, d) be the induced metric of (X, δ) . Letting $X = \{x_1, \dots, x_n\}$. Define ϕ by

$$\phi(x) = (d(x, x_1), d(x, x_2), \dots, d(x, x_n)).$$

Then for all $A \subseteq X$ we have

$$n\delta(A) \geq \sum_{i=1}^n \max_{a,b \in A} d(a, b) \geq \sum_i \max_{a,b \in A} d(a, x_i) - d(b, x_i) = \hat{\delta}(\phi(A)),$$

where the first inequality follows from the monotonicity of δ and the second follows from the triangle inequality for diversities. Also, for all $A \subseteq X$ we have

$$\hat{\delta}(\phi(A)) \geq \sum_{x \in A} \max_{a,b \in A} d(a, x) - d(b, x) = \sum_{x \in A} \max_{a \in A} d(a, x) \geq \sum_{x \in A} d(a_0, x) \geq \delta(A),$$

where a_0 is any element of A . These two inequalities give distortion at most n . \square

There are metric spaces that require $\Omega(\log n)$ distortion [14]. If we take a diversity that has this metric as its induced metric, then we cannot embed that diversity with less distortion.

In Section 2 we give a survey of special classes of diversities for which we have upper bounds better than $\mathcal{O}(n)$. In Section 3 we show how two of the techniques that were used for the metric embedding problem fail for diversities. Section 4 lists some ideas for possible directions to follow.

2 Special Cases

There are a few special cases where we have tighter upper bounds on the worst case distortion. See [5] for examples of diversities which can be embedded into ℓ_1 with no distortion, as well as characterisations of these diversities.

2.1 Diameter Diversities

Every metric space (X, d) induces a diversity $(X, \delta_{\text{diam}})$, where δ_{diam} is the diameter diversity defined by

$$\delta_{\text{diam}}(A) = \max_{a,b \in A} d(a, b).$$

$(X, \delta_{\text{diam}})$ is a diversity, and is in fact the minimal diversity that has d as its induced metric [4]

Let ℓ_1^k denote \mathbb{R}^k equipped with the ℓ_1 diversity $\hat{\delta}$. In [5, Prop. 12] we show that any diameter diversity on n points can be embedded into ℓ_1^k with distortion $\mathcal{O}(\log^2 n)$ where $k = \mathcal{O}(\log n)$. The result uses the ℓ_1 embedding result of [14] along with the fact that there is at most a ratio of k between the diameter diversity and the ℓ_1 diversity on ℓ_1^k .

One important example of a diameter diversity is the *discrete diversity*, given by

$$\delta_p(A) = \begin{cases} 1 & |A| > 1 \\ 0 & \text{otherwise.} \end{cases}$$

2.2 Steiner Diversities

Consider any metric space (X, d) . View the metric space as a complete weighted graph with vertices X and the weight of edge (u, v) to be $d(u, v)$. The Steiner diversity $\delta_S(A)$ of a finite set A is the minimal total weight of a tree in the graph whose leaves include the points in A . (X, δ_S) is known as the Steiner diversity of (X, d) , and is the maximal diversity that has (X, d) as its induced metric [5].

A simple argument based on a metric approximation result due to [10] gives the following. We note that the result of [10] was a refinement of an earlier result of Bartal [1, 2].

Theorem 2. *Let (X, d) be a metric space with $|X| = n$ and let (X, δ_S) be its Steiner diversity. Then (X, δ_S) has an ℓ_1 embedding with distortion $\mathcal{O}(\log n)$.*

Proof. By [10] there is a bijection ϕ from X to the vertices of a random tree τ with vertex set X such that for all $u, v \in X$ we have $d(u, v) \leq d_\tau(u, v)$ and $\mathbb{E}[d_\tau(u, v)] \leq (\log n) d(u, v)$. (Here \mathbb{E} denotes expectation.)

For all $A \subseteq X$, there is a tree $T = (X_{\phi(A)}, E_{\phi(A)})$ such that T is a subtree connecting $\phi(A)$ in τ and $\sum_{\{u, v\} \in E_{\phi(A)}} d_\tau(u, v) = \delta_\tau(A)$. Then

$$\delta_S(A) \leq \sum_{\{u, v\} \in E_{\phi(A)}} d(u, v) \leq \sum_{\{u, v\} \in E_{\phi(A)}} d_\tau(u, v) = \delta_\tau(A).$$

Likewise, for all $A \subseteq X$, there is a tree $T = (V_A, E_A)$ such that

$$\delta_S(A) = \sum_{\{u, v\} \in E_A} d(u, v)$$

(by the definition of the Steiner diversity) which gives

$$\begin{aligned} \mathbb{E}[\delta_\tau(A)] &\leq \mathbb{E} \left[\sum_{\{u, v\} \in E_A} d_\tau(u, v) \right] \\ &\leq \mathcal{O}(\log n) \sum_{\{u, v\} \in E_A} d(u, v) \\ &= \mathcal{O}(\log n) \delta_S(A). \end{aligned}$$

Hence $\mathbb{E}[\delta_\tau(A)] \leq \mathcal{O}(\log n) \delta_S(A)$.

For each τ we have that δ_τ can be embedded in ℓ_1 , see [4]. Hence so can the expectation,

$$\hat{\delta} = \mathbb{E} \delta_\tau$$

and from the above we have that

$$\delta(A) \leq \hat{\delta}(A) \leq \mathcal{O}(\log n) \delta(A)$$

for all A . □

2.3 Hypergraph Steiner

Let $\mathcal{H} = (V, \mathcal{E}, w)$ be a hypergraph with a weight function $w : \mathcal{E} \rightarrow \mathbb{R}$. The hypergraph Steiner diversity δ_H is defined as the minimum of

$$w(\mathcal{H}') = \sum_{U \in \mathcal{E}(\mathcal{H}')} w(U)$$

over all connected sub-hypergraphs of \mathcal{H} with $A \subseteq V(\mathcal{H}')$, with singletons defined to have diversity zero.

Every diversity on a finite set equals δ_H for some weighted hypergraph—just use $\mathcal{E} = P(V)$ and $w(U) = \delta(U)$. We bound the embedding distortion for δ_H in terms of the sizes of hyperedges in $\mathcal{E}(\mathcal{H})$.

Theorem 3. *Let δ_H be the Hypergraph Steiner diversity for $\mathcal{H} = (V, \mathcal{E}, w)$ and suppose that $k = \max\{|U| : U \in \mathcal{E}(\mathcal{H})\}$. Then δ_H can be embedded in ℓ_1 with $\mathcal{O}(k \log n)$ distortion.*

Proof. For each $U \in \mathcal{E}$ let T_U be a spanning tree of the connected graph with vertex set U , where each edge in T_U has weight $w(U)$. Construct the graph $G = (V, E)$ where E is the disjoint union

$$E = \bigsqcup_{U \in \mathcal{E}} E(T_U).$$

Every connected subgraph of G gives a connected sub-hypergraph of \mathcal{H} with lesser weight, and every connected sub-hypergraph of \mathcal{H} gives a connected subgraph of G with weight at most $(k - 1)$ times larger. It follows that

$$\delta_H(A) \leq \delta_S(A) \leq (k - 1)\delta_H(A)$$

and the result now follows from Theorem 2. □

2.4 Ball Diversities

Let (X, d) be any metric space. Define the diversity (X, δ_B) by letting $\delta_B(A)$ equal the diameter of the smallest ball containing A . We call this the *ball diversity*.

For any $A \subseteq X$ we have

$$\delta_{\text{diam}}(A) \leq \delta_B(A) \leq 2 \min_{a \in A} \max_{b \in A} d(a, b) \leq 2\delta_{\text{diam}}(A)$$

and so a bound of $\mathcal{O}(\log^2 n)$ distortion for embedding ball diversities in ℓ_1 follows from the diameter diversity case.

2.5 TSP Diversity

Let (X, d) be any metric space. Define the diversity (X, δ_{TSP}) where $\delta_{TSP}(A)$ is the length of the shortest tour through points in A . We then have that

$$\delta_S(A) \leq \delta_{TSP}(A) \leq 2\delta_S(A)$$

so δ_{TSP} can be embedded with $\mathcal{O}(\log n)$ distortion.

2.6 Partition diversities

Given a partition π of X we define δ_π by

$$\delta_\pi(A) = \begin{cases} 1 & A \text{ intersects at least two blocks of } \pi; \\ 0 & \text{otherwise.} \end{cases}$$

This reduces to a *split diversity* or *cut diversity* when π has two blocks [5]. In general, partition diversities are not ℓ_1 embeddable, however they are examples of diameter diversities. Consequently, any non-negative linear combination of partition diversities can be embedded into ℓ_1 with $\mathcal{O}(\log^2 n)$ distortion.

2.7 Symmetric Diversities

We say a diversity is symmetric if $\delta(A) = \delta(B)$ whenever $|A| = |B|$. Perhaps surprisingly, symmetric diversities can be embedded into ℓ_1 with a constant factor distortion [6]. A consequence of this result is that if $\frac{\delta(A)}{\delta(B)} \leq K$ whenever $|A| = |B| > 1$ then δ can be embedded in ℓ_1 with distortion $\mathcal{O}(K)$.

3 Failed Attempts

We have tried two different approaches to the problem, one based on the techniques used in Bourgain's proof, and the other based on Bartal's techniques. Both were unsuccessful, but we will explain where the problem lies in each case.

3.1 Bourgain's Method

To summarize Bourgain's method at a high level [3, 8], suppose we are given a metric (X, d) with $|X| = n$ and we want to embed it in ℓ_1^k for some k . We select a collection of maps $\phi_{d,A} : X \rightarrow \mathbb{R}$ for $A \subseteq X$, and a collection of weights $c_{d,A} \geq 0$. Then the ℓ_1 metric is defined as

$$\hat{d}(x, y) = \sum_{A \subseteq X} c_{d,A} |\phi_{d,A}(x) - \phi_{d,A}(y)|.$$

The space (X, \hat{d}) can be embedded in ℓ_1^k where k is the number of non-zero $c_{d,A}$. The $\phi_{d,A}$ in the metric case is chosen as $\phi_{d,A}(x) = d(x, A)$, and the constants $c_{d,A}$ are chosen independently of d and are 0 unless $|A| = 2^s$ for some integer s [8, Thm 10.1.2].

We can imagine following a similar method for diversities. Given a diversity (X, δ) with $|X| = n$, we select mappings $\phi_{\delta,A} : X \rightarrow \mathbb{R}$, and constants $c_{\delta,A} \geq 0$. Then we define

$$\hat{\delta}(B) = \sum_{A \subseteq X} c_{\delta,A} \max_{b_1, b_2 \in B} |\phi_{\delta,A}(b_1) - \phi_{\delta,A}(b_2)|,$$

which is an ℓ_1 diversity. The goal is then to define $\phi_{\delta,A}$ and $c_{\delta,A}$ to make $\hat{\delta}$ close to δ . We now describe some of the difficulties with making this work for diversities.

Here are three choices we could try for $\phi_{\delta,A}$.

1. $\phi_{\delta,A} = \delta_{A|\bar{A}}$ the split diversity for the split $A|\bar{A}$ [5].
2. $\phi_{\delta,A} = d(A, x) = \min_{a \in A} \delta(\{x, a\})$.

$$3. \phi_{\delta,A} = \delta(A \cup \{x\}).$$

We consider each in turn.

(1). Any ℓ_1 diversity can be written as a non-negative linear combination of the split diversities, so this choice of $\phi_{\delta,A}$ is just a restatement of the problem.

(2). This choice of $\phi_{\delta,A}$ only depends on δ through the values $\delta(\{x, a\}), x, a \in X$. In other words, the $\phi_{\delta,A}$ only depend on δ through its induced metric. In general $c_{\delta,A}$ may also depend on δ , and in this is, as far as we know, a possible approach to the problem. But if we restrict ourselves to constants independent of δ , c_A for $A \subseteq X$, have a stronger restriction. With this choice, any two diversities with the same induced metric will yield the same embedded diversity $\hat{\delta}$. This puts a constraint on how good the worst case distortion between δ and $\hat{\delta}$ can be. To see this, let δ_ρ be the discrete diversity on n points that assigns value 1 to all sets with two or more elements. Let δ_c be given by $\delta_2(A) = |A| - 1$ for $|A| \geq 2$. δ_ρ and δ_c have the same induced metric, and so $\hat{\delta}$ would be identical for them. But $\delta_\rho(X) = 1$ and $\delta_c(X) = n - 1$ so $\hat{\delta}(X)$ must be off at least one of them by a factor of $\sqrt{n-1}$ or greater. So the best worst case distortion for this method is at least $\mathcal{O}(\sqrt{n})$. We don't know if this is attainable.

(3). The choice $\phi_{\delta,A}(x) = \delta(A \cup \{x\})$ looks promising in that it potentially uses all the information about the diversity. We get

$$\hat{\delta}(B) = \sum_{A \subseteq X} c_A \max_{b_1, b_2 \in B} \delta(A \cup \{b_1\}) - \delta(A \cup \{b_2\}).$$

To get an idea of how this works, we apply it to the diversity δ_ρ from the previous case. Most terms are 0 and we get

$$\begin{aligned} \hat{\delta}(B) &= \sum_{a \in X} c_{\delta_\rho, \{a\}} \max_{b_1, b_2 \in B} \delta_\rho(\{a, b_1\}) - \delta_\rho(\{a, b_2\}) \\ &= \sum_{a \in X} c_{\delta_\rho, \{a\}} \mathbf{1}_{a \in B, |B| \geq 2} \\ &= \mathbf{1}_{|B| \geq 2} \sum_{a \in B} c_{\delta_\rho, \{a\}} = \delta(B) \sum_{a \in B} c_{\delta_\rho, \{a\}}. \end{aligned}$$

Suppose we have

$$\hat{\delta}(B) \leq \delta(B) \leq g(n) \hat{\delta}(B).$$

Taking $B = X$ this gives:

$$\sum_{a \in X} c_{\delta_\rho, \{a\}} \leq 1.$$

Taking $B = \{a_1, a_2\}$, $a_1 \neq a_2$ gives

$$1 \leq g(n)(c_{\delta_\rho, \{a_1\}} + c_{\delta_\rho, \{a_2\}})$$

Summing this over $\lfloor n/2 \rfloor$ pairs of distinct points in X gives

$$\lfloor n/2 \rfloor \leq g(n) \sum_{a \in X} c_{\delta_\rho, \{a\}} \leq g(n).$$

So $g(n)$ is at least $\mathcal{O}(n)$. So this choice of $\phi_{\delta,A}$ cannot give us better results than the bound we already have.

3.2 Bartal's Method

Bartal's method [1, 2], and as later refined in [10], relies on embedding a metric space (X, d) in a random tree metric. Since tree metrics are ℓ_1 , a random tree metric is also ℓ_1 (being a positive linear combination of tree metrics). These proofs work by randomly and hierarchically building a tree corresponding to the set X such that tree metric dominates d and, for each pair of nodes x, y , the expected distance is within $\mathcal{O}(\log n)$ of $d(x, y)$.

The problem with this approach for diversities can be seen easily. We again use the diversity δ_ρ from above, where $\delta_\rho(A) = 1$ whenever $|A| \geq 2$. Since δ_ρ is a diameter diversity, we know it can be approximated with an ℓ_1 diversity with distortion $\mathcal{O}(\log^2 n)$. But the best we can do with a phylogenetic diversity (the equivalent of a tree diversity) is much worse.

Proposition 1. *Let (X, δ_ρ) be the diversity with $|X| = n$ and $\delta_\rho(A) = 1$ for all $|A| \geq 2$. Let δ_τ be a random phylogenetic diversity with*

$$\delta_\rho(A) \leq \delta_\tau(A), \text{ and } \mathbb{E}\delta_\tau(A) \leq g(n)\delta_\rho(A)$$

for all $A \subseteq X$. Then $g(n) \geq \lfloor n/2 \rfloor$.

Proof. Let L_t be the length of the tree δ_t . By Corollary 5.18 and Theorem 5.6.2 in [17] there are $m = \lfloor n/2 \rfloor$ pairs of vertices such that the paths joining each pair are disjoint. Denoting these pairs $(a_1, b_1), \dots, (a_m, b_m)$, we have that $L_t \geq \sum_{i=1}^m d(a_i, b_i) = \sum_{i=1}^m \delta_t(\{a_i, b_i\}) \geq m$. On the other hand,

$$m \leq \mathbb{E}L_t = \mathbb{E}\delta_t(X) \leq g(n)\delta_\rho(X) = g(n).$$

So $g(n) \geq \lfloor n/2 \rfloor$. □

4 Other ideas

1. Every diversity (X, δ) with a given induced metric (X, d) satisfies the inequalities

$$\delta_{\text{diam}}(A) \leq \delta(A) \leq \delta_S(A).$$

We have shown that Bourgain's approach embeds δ_{diam} with small distortion, but applying the metric result directly does appear to give an embedding for δ_S . On the other hand, Bartal's approach embeds δ_S , but does not successfully embed all diameter diversities. Is some hybrid approach possible, perhaps one which applies Bartal's approach but switches to a Bourgain approach for some subproblems?

2. Expander graphs have proven to be a good source for lower bounds on the distortion required to embed metrics in ℓ_1 . Expander hypergraphs are far less understood, but perhaps analogous arguments could be used to obtain lower bounds on the distortion of diversities, particularly given the connections between diversity embedding and hypergraph sparsest cut established in [5].
3. The embedding used to prove Theorem 1 was derived completely from the induced metric. We know that we cannot obtain a distortion bound in this way that is better than $\Omega(\sqrt{n})$, as we showed in Part 2 of our discussion of Bourgain's approach. We conjecture that there exists an embedding based solely on the induced metric which nevertheless achieves an $\mathcal{O}(\sqrt{n})$ distortion bound for all diversities.

References

- [1] Yair Bartal. Probabilistic approximation of metric spaces and its algorithmic applications. In *Foundations of Computer Science, 1996. Proceedings., 37th Annual Symposium on*, pages 184–193. IEEE, 1996.
- [2] Yair Bartal. On approximating arbitrary metrics by tree metrics. In *Proceedings of the thirtieth annual ACM symposium on Theory of computing*, pages 161–168. ACM, 1998.
- [3] J. Bourgain. On Lipschitz embedding of finite metric spaces in Hilbert space. *Israel J. Math.*, 52(1-2):46–52, 1985.
- [4] David Bryant and Paul F Tupper. Hyperconvexity and tight-span theory for diversities. *Advances in Mathematics*, 231(6):3172–3198, 2012.
- [5] David Bryant and Paul F. Tupper. Diversities and the geometry of hypergraphs. *Discrete Math. Theor. Comput. Sci.*, 16(2):1–20, 2014.
- [6] David Bryant and Paul F Tupper. Constant distortion embeddings of symmetric diversities. *Analysis and Geometry in Metric Spaces*, 4(1), 2016.
- [7] Elena Deza, Michel Deza, and Mathieu Dutour Sikirič. *Generalizations of finite metrics and cuts*. World Scientific, 2016.
- [8] Michel Marie Deza and Monique Laurent. *Geometry of cuts and metrics*, volume 15 of *Algorithms and Combinatorics*. Springer-Verlag, Berlin, 1997.
- [9] Rafael Espínola and Aurora Fernández-León. Fixed point theory in hyperconvex metric spaces. In *Topics in Fixed Point Theory*, pages 101–158. Springer, 2014.
- [10] Jittat Fakcharoenphol, Satish Rao, and Kunal Talwar. A tight bound on approximating arbitrary metrics by tree metrics. *Journal of Computer and System Sciences*, 69:485–497, 2004.
- [11] Sven Herrmann and Vincent Moulton. Trees, tight-spans and point configurations. *Discrete Mathematics*, 312(16):2506–2521, 2012.
- [12] William Kirk and Naseer Shahzad. Diversities. In *Fixed Point Theory in Distance Spaces*, pages 153–158. Springer, 2014.
- [13] Pawan Kumar, Nikos Komodakis, and Nikos Paragios. (Hyper)-Graphs Inference via Convex Relaxations and Move Making Algorithms: Contributions and Applications in artificial vision. Technical report, Inria, 2015.
- [14] Nathan Linial, Eran London, and Yuri Rabinovich. The geometry of graphs and some of its algorithmic applications. *Combinatorica*, 15(2):215–245, 1995.
- [15] Bożena Piątek. On the gluing of hyperconvex metrics and diversities. *Annales Universitatis Paedagogicae Cracoviensis. Studia Mathematica*, 13(1):65–76, 2014.
- [16] Andrew Poelstra. On the topological and uniform structure of diversities. *Journal of Function Spaces and Applications*, 2013.

- [17] Charles Semple and Mike A Steel. *Phylogenetics*, volume 24. Oxford University Press on Demand, 2003.
- [18] Mike Steel. Tracing evolutionary links between species. *The American Mathematical Monthly*, 121(9):771–792, 2014.