

# On the counting function for numerical monoids

Francisco Franco Munoz

## Abstract

We provide explicit expressions for the counting function and main terms for numerical monoids.

## 1 Introduction

The study of numerical monoids is a huge area of research (see [2] and references therein). A numerical monoid  $M$  is a subset of  $\mathbb{N}$  containing 0 that is closed under addition and finite complement. In other words, a finite complement submonoid of  $(\mathbb{N}, +)$ .

The aim of this paper is to calculate in an elementary way the main term of the counting function of a monoid (Theorems 5, 13, 14) which is a natural object. We'll avoid the use of more advanced treatments such as quasi-polynomial functions and integer points of polytopes ([1]) that provide a more comprehensive view of the subject. The paper is essentially self-contained and requires no prior knowledge of the subject.

### 1.1 Notation

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ , the natural numbers

$\mathbb{Z} = -\mathbb{N} \cup \mathbb{N}$ , the integers

$\mathbb{N}_+ = \{1, 2, 3, \dots\}$ , the positive integers

## 2 Setting

Let  $M = \langle a_1, \dots, a_{n-1}, a_n \rangle$  be a submonoid of  $(\mathbb{N}, +)$ . By definition  $M$  is called a numerical monoid if its complement is finite. Denote by  $M_{\widehat{a_n}} = \langle a_1, \dots, a_{n-1} \rangle$  the submonoid obtained by dropping the generator  $a_n$ .

**Definition 1.** For  $x \in \mathbb{R}$ ,  $a \in \mathbb{N}_+$ , define  $\{x; a\} = x - a \left\lfloor \frac{x}{a} \right\rfloor$ .

With this definition, we have  $\{x\} = \{x; 1\}$  the fractional part of  $x$ .

**Proposition 1.** Let  $x, k \in \mathbb{Z}$ ,  $a, b \in \mathbb{N}_{>0}$

1.  $\{x; a\}$  is the residue of  $x$  modulo  $a$
2.  $0 \leq \{x; a\} < a$
3.  $\{x; a\} = 0 \iff a|x$ .
4.  $\{x + ka; a\} = \{x; a\}$
5.  $\{x; b\} = \{\{x; ab\}; b\}$

*Proof.* Immediate. ■

**Lemma 2.** For  $y \in \mathbb{N}$ , if  $y < a_n$ , then  $y \in M \iff y \in M_{\widehat{a_n}}$ . In particular, for  $x \in \mathbb{N}$ ,  $\{x; a_n\} \in M \iff \{x; a_n\} \in M_{\widehat{a_n}}$ .

*Proof.* This is clear. ■

**Definition 2.** We define the counting function

$$T_M(x) = \#\{(x_1, \dots, x_n) \in \mathbb{N}^n \mid x_1 a_1 + \dots + x_n a_n = x\}$$

The Theory of Erhart polynomials [1] provides ways to determine  $T_M$ . Instead we'll address them directly in an elementary way.

**Theorem 3.**

$$T_M(x) = T_M(x - a_n) + T_{M_{\widehat{a_n}}}(x)$$

*Proof.* This is immediate since the set  $\{(x_1, \dots, x_n) \in \mathbb{N}^n \mid x_1 a_1 + \dots + x_n a_n = x\}$  can be partitioned as  $\{(x_1, \dots, x_n) \in \mathbb{N}^n \mid x_1 a_1 + \dots + x_n a_n = x\} =$

$$\begin{aligned} &= \{(x_1, \dots, x_n) \in \mathbb{N}^n \mid x_n > 0, x_1 a_1 + \dots + x_n a_n = x\} \sqcup \{(x_1, \dots, x_n) \in \mathbb{N}^n \mid x_n = 0, x_1 a_1 + \dots + x_n a_n = x\} \\ &= \{(x_1, \dots, x_{n-1}, y_n) \in \mathbb{N}^n \mid y_n = x_n - 1 \geq 0, x_1 a_1 + \dots + x_{n-1} a_{n-1} + y_n a_n = x - a_n\} \sqcup \\ &\quad \{(x_1, \dots, x_{n-1}, 0) \in \mathbb{N}^n \mid x_n = 0, x_1 a_1 + \dots + x_{n-1} a_{n-1} + 0 = x\} \text{ and the counting for each is } \\ &\quad T_M(x - a_n) + T_{M_{\widehat{a_n}}}(x). \end{aligned}$$
■

**Corollary 4.**  $T_M(x) = \sum_{k=0}^{\lfloor \frac{x}{a_n} \rfloor} T_{M_{\widehat{a_n}}}(x - k a_n) = \sum_{l=0}^{\lfloor \frac{x}{a_n} \rfloor} T_{M_{\widehat{a_n}}}(\{x; a_n\} + l a_n)$

*Proof.*  $T_M(x) - T_M\left(x - a_n \left\lfloor \frac{x}{a_n} \right\rfloor\right) = \sum_{k=0}^{\lfloor \frac{x}{a_n} \rfloor - 1} [T_M(x - k a_n) - T_M(x - (k+1) a_n)]$

$$= \sum_{k=0}^{\lfloor \frac{x}{a_n} \rfloor - 1} T_{M_{\widehat{a_n}}}(x - k a_n). \text{ Also } T_M\left(x - a_n \left\lfloor \frac{x}{a_n} \right\rfloor\right) = T_M(\{x; a_n\}) = T_{M_{\widehat{a_n}}}(\{x; a_n\}) = T_{M_{\widehat{a_n}}}\left(x - a_n \left\lfloor \frac{x}{a_n} \right\rfloor\right)$$

by Lemma 2 and we conclude that  $T_M(x) = \sum_{k=0}^{\lfloor \frac{x}{a_n} \rfloor} T_{M_{\widehat{a_n}}}(x - k a_n)$  and making the change of variable  $k = \left\lfloor \frac{x}{a_n} \right\rfloor - l$  we get the last equality. ■

### 3 Dimension $n = 2$

$M = \langle a, b \rangle$  and assume that  $\gcd(a, b) = 1$  in this subsection.

**Theorem 5.** We have

$$T_{\langle a, b \rangle}(x) = \left\lfloor \frac{x}{ab} \right\rfloor + \epsilon_{\langle a, b \rangle}(\{x; ab\})$$

where for  $0 \leq y < ab$ ,  $\epsilon(y) = \delta(y \in \langle a, b \rangle)$  is the indicator function.

*Proof.* Applying Corollary 4 with  $n = 2$  and  $a_2 = b$ , we have  $T_{\langle a, b \rangle}(x) = \sum_{l=0}^{\lfloor \frac{x}{b} \rfloor} T_{M_b}(\{x; b\} + lb) = \sum_{l=0}^{\lfloor \frac{x}{b} \rfloor} T_{\langle a \rangle}(\{x; b\} + lb)$ . Now, it's clear that  $T_{\langle a \rangle}(y)$  is 1 or 0 according to whether  $a|y$  or not and so  $T_{\langle a \rangle}(y + ka) = T_{\langle a \rangle}(y)$ . Let  $y = \lfloor \frac{x}{b} \rfloor$ . By Proposition 1,  $\lfloor \frac{\lfloor \frac{x}{b} \rfloor}{a} \rfloor = \lfloor \frac{x}{ab} \rfloor$ , so  $y = \{y; a\} + a \lfloor \frac{x}{ab} \rfloor$ . We split the sum  $\sum_{l=0}^{\lfloor \frac{x}{b} \rfloor} T_{\langle a \rangle}(\{x; b\} + lb) = \sum_{i=0}^{\lfloor \frac{x}{ab} \rfloor - 1} \sum_{j=0}^{a-1} T_{\langle a \rangle}(\{x; b\} + (ai + j)b) + \sum_{j=0}^{\{y; a\}} T_{\langle a \rangle}(\{x; b\} + (a \lfloor \frac{x}{ab} \rfloor + j)b) = \sum_{i=0}^{\lfloor \frac{x}{ab} \rfloor - 1} \sum_{j=0}^{a-1} T_{\langle a \rangle}(\{x; b\} + jb) + \sum_{j=0}^{\{y; a\}} T_{\langle a \rangle}(\{x; b\} + jb) = \lfloor \frac{x}{ab} \rfloor \sum_{j=0}^{a-1} T_{\langle a \rangle}(\{x; b\} + jb) + \sum_{j=0}^{\{y; a\}} T_{\langle a \rangle}(\{x; b\} + jb)$  and the proof follows from the next two lemmas. ■

**Lemma 6.** For any  $y \in \mathbb{N}$ ,  $\sum_{j=0}^{a-1} T_{\langle a \rangle}(y + jb) = 1$

*Proof.* Since  $\gcd(a, b) = 1$ , there's unique  $j, 0 \leq j \leq a - 1$  such that  $a \mid (y + jb)$ . ■

**Lemma 7.** For any  $x \in \mathbb{N}$ , then  $\sum_{j=0}^{\{y; a\}} T_{\langle a \rangle}(\{x; b\} + jb) = \epsilon_{\langle a, b \rangle}(\{x; ab\})$  where  $y = \lfloor \frac{x}{b} \rfloor$ .

*Proof.* Omitted. ■

### 3.1 A different proof

**Lemma 8.** If  $x = \alpha a + \beta b \in M = \langle a, b \rangle$ , then  $T_M(x) = \lfloor \frac{\alpha}{b} \rfloor + \lfloor \frac{\beta}{a} \rfloor + 1$

*Proof.* Fix a representation  $x = \alpha a + \beta b$ . We need to count representations  $x = \alpha_1 a + \beta_1 b$ . But  $\alpha a + \beta b = \alpha_1 a + \beta_1 b \iff (\alpha - \alpha_1)a = (\beta_1 - \beta)b$  and since  $\gcd(a, b) = 1$  this happens  $\iff a \mid (\beta_1 - \beta), b \mid (\alpha - \alpha_1)$  and  $\beta_1 - \beta = ja, \alpha - \alpha_1 = jb$ . So we need to count the possibilities for  $j \in \mathbb{Z}$ . We have  $\beta_1 = ja + \beta, \alpha_1 = -jb + \alpha$ . And the condition is exactly  $\beta_1 \geq 0$  and  $\alpha_1 \geq 0$ , which happens  $\iff ja + \beta \geq 0, -jb + \alpha \geq 0 \iff \frac{-\beta}{a} \leq j \leq \frac{\alpha}{b} \iff \lceil \frac{-\beta}{a} \rceil \leq j \leq \lfloor \frac{\alpha}{b} \rfloor$ . Now we have  $\lceil \frac{-\beta}{a} \rceil = -\lfloor \frac{\beta}{a} \rfloor$  hence there are  $\lfloor \frac{\alpha}{b} \rfloor + \lfloor \frac{\beta}{a} \rfloor + 1$  such  $j$ . ■

**Corollary 9.** For  $x < ab$ ,  $T_M(x) = 1 \iff x \in M$ .

*Proof.* It's enough to show that if  $x \in M$  and  $x < ab$ , then  $T_M(x) = 1$ . By Lemma 8, this holds iff  $\lfloor \frac{\alpha}{b} \rfloor = \lfloor \frac{\beta}{a} \rfloor = 0$  where  $x = \alpha a + \beta b$  is some representation. But since  $x < ab$ , then  $\alpha a \leq x < ab$  so  $\alpha < b$  and  $\beta b \leq x < ab$  so  $\beta < a$  so  $\lfloor \frac{\alpha}{b} \rfloor = \lfloor \frac{\beta}{a} \rfloor = 0$ . ■

**Lemma 10.** If  $x \in M$ , then for all  $k \in \mathbb{N}$ ,  $T_M(x + kab) = T_M(x) + k$

*Proof.* By Lemma 8, take  $x = \alpha a + \beta b \in M$  then  $x + kab \in M$  and  $x + kab = (\alpha + kb)a + \beta b$  and so  $T_M(x + kab) = \lfloor \frac{\alpha + kb}{b} \rfloor + \lfloor \frac{\beta}{a} \rfloor + 1 = k + \lfloor \frac{\alpha}{b} \rfloor + \lfloor \frac{\beta}{a} \rfloor + 1 = T_M(x) + k$ . ■

**Lemma 11.** If  $x \notin M$ ,  $0 \leq x \leq ab - 1$ , then  $T_M(x + ab) = 1$

*Proof.* Since  $x+ab \geq ab$  and the Frobenius number of  $M$  is  $ab-a-b$  ([2]),  $x+ab$  belongs to  $M$ , so  $T_M(x+ab) \geq 1$ . Now, as in Corollary 9, it's enough to show that in any representation of  $x+ab = \alpha a + \beta b$ , both  $\alpha < b$  and  $\beta < a$ . If not, say  $\alpha \geq b$ , then  $x = \alpha a + \beta b - ab = (\alpha - b)a + \beta b$  which says that  $x \in M$ , contradiction. In the same way,  $\beta < a$  and we're done. ■

**Theorem 12.** *For all  $x \in \mathbb{N}$ ,  $T_M(x+ab) = T_M(x) + 1$  and Theorem 5 is valid.*

*Proof.* By contradiction, take  $x$  minimal such that this formula is not valid for  $x$ . Notice that by Lemma 11, we have  $x \geq ab$ , and by Lemma 10, we have  $x \notin M$ . Let  $y = x - ab < x$  then by minimality, the theorem is valid for  $y$  and so  $T_M(x) = T_M(y+ab) = T_M(y) + 1 \geq 1$  in particular  $x \in M$ , contradiction.

Since  $\forall x \in \mathbb{N}$ ,  $T_M(x+ab) = T_M(x) + 1$ , by induction  $\forall x, k \in \mathbb{N}$ ,  $T_M(x+kab) = T_M(x) + k$ , and so  $T_M(x) = T_M(\{x; ab\} + ab \lfloor \frac{x}{ab} \rfloor) = \lfloor \frac{x}{ab} \rfloor + T_M(\{x; ab\}) = \lfloor \frac{x}{ab} \rfloor + \epsilon_{\langle a, b \rangle}(\{x; ab\})$  the last equality since  $\{x; ab\} < ab$  and using Corollary 9. This proves the theorem. ■

## 4 Dimension $n = 3$

Assume  $\gcd(a, b) = \gcd(a, c) = \gcd(b, c) = 1$ . These conditions are stronger than necessary, but the theory of numerical monoids permits one to reduce to this case in any dimension [2].

Let  $M = \langle a, b, c \rangle$ ,  $M_{\widehat{c}} = \langle a, b \rangle$  dimension 3 and 2 numerical monoids.

$$\begin{aligned}
\text{Recall } T_M(x) &= \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} T_{\langle a, b \rangle}(\{x; c\} + lc) \\
\text{Let } y &= \lfloor \frac{x}{c} \rfloor. \text{ Then } y = \{y; ab\} + ab \lfloor \frac{y}{ab} \rfloor. \text{ By Theorem 5, } T_{\langle a, b \rangle} \text{ splits as sum, consequently} \\
\text{we split the sum } \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} T_{\langle a, b \rangle}(\{x; c\} + lc) &= \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} \left\lfloor \frac{\{x; c\} + lc}{ab} \right\rfloor + \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} \epsilon_{\langle a, b \rangle}(\{x; c\} + lc; ab) = S1 + S2 \\
S1 &= \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} \left\lfloor \frac{\{x; c\} + lc}{ab} \right\rfloor = \sum_{i=0}^{\lfloor \frac{x}{abc} \rfloor - 1} \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + (abi+j)c}{ab} \right\rfloor + \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + (ab \lfloor \frac{y}{ab} \rfloor + j)c}{ab} \right\rfloor \\
&= \sum_{i=0}^{\lfloor \frac{x}{abc} \rfloor - 1} \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \sum_{i=0}^{\lfloor \frac{x}{abc} \rfloor - 1} \sum_{j=0}^{ab-1} ic + \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{x}{abc} \right\rfloor c \\
&= \left\lfloor \frac{x}{abc} \right\rfloor \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) \left\lfloor \frac{x}{abc} \right\rfloor + \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + (\{y; ab\} + 1) \left\lfloor \frac{x}{abc} \right\rfloor c \\
S2 &= \sum_{l=0}^{\lfloor \frac{x}{c} \rfloor} \epsilon_{\langle a, b \rangle}(\{x; c\} + lc; ab) = \sum_{i=0}^{\lfloor \frac{x}{abc} \rfloor - 1} \sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{x; c\} + (abi+j)c; ab) + \sum_{j=0}^{\{y; ab\}} \epsilon_{\langle a, b \rangle}(\{x; c\} + \\
&\quad (ab \lfloor \frac{y}{ab} \rfloor + j)c; ab) = \sum_{i=0}^{\lfloor \frac{x}{abc} \rfloor - 1} \sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{x; c\} + jc; ab) + \sum_{j=0}^{\{y; ab\}} \epsilon_{\langle a, b \rangle}(\{x; c\} + jc; ab) \\
&= \left\lfloor \frac{x}{abc} \right\rfloor \sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{x; c\} + jc; ab) + \sum_{j=0}^{\{y; ab\}} \epsilon_{\langle a, b \rangle}(\{x; c\} + jc; ab)
\end{aligned}$$

Using Hermite reciprocity, since  $\gcd(ab, c) = 1$ :

$$\begin{aligned} \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor &= \sum_{j=0}^{c-1} \left\lfloor \frac{\{x; c\} + j ab}{c} \right\rfloor = \sum_{j=0}^{c-1} \left\lfloor \frac{\{x; c\} + \{jab; c\} + c \lfloor \frac{j ab}{c} \rfloor}{c} \right\rfloor = \sum_{j=0}^{c-1} \left\lfloor \frac{\{x; c\} + \{jab; c\}}{c} \right\rfloor + \\ \sum_{j=0}^{c-1} \left\lfloor \frac{j ab}{c} \right\rfloor &= \sum_{l=0}^{c-1} \left\lfloor \frac{\{x; c\} + l}{c} \right\rfloor + \sum_{j=0}^{c-1} \left\lfloor \frac{j ab}{c} \right\rfloor = \left\lfloor c \frac{\{x; c\}}{c} \right\rfloor + \frac{(ab-1)(c-1)}{2} = \{x; c\} + \frac{(ab-1)(c-1)}{2} \end{aligned}$$

Hence we obtain  $S1 =$

$$\begin{aligned} &\sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \left\lfloor \frac{x}{abc} \right\rfloor \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) \left\lfloor \frac{x}{abc} \right\rfloor + (\{y; ab\} + 1) \left\lfloor \frac{x}{abc} \right\rfloor c \\ &= \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \left\lfloor \frac{x}{abc} \right\rfloor \left( \{x; c\} + \frac{(ab-1)(c-1)}{2} \right) + \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) \left\lfloor \frac{x}{abc} \right\rfloor + \\ &(\{y; ab\} + 1) \left\lfloor \frac{x}{abc} \right\rfloor c = \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) + \{x; c\} + \frac{(ab-1)(c-1)}{2} + (\{y; ab\} + 1)c \right) \\ &= \sum_{j=0}^{\{y; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + S3, \text{ where } S3 \text{ is the sum} \\ S3 &= \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) + \{x; c\} + \frac{(ab-1)(c-1)}{2} + (\{y; ab\} + 1)c \right) \\ &= \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) + \{x; c\} + \frac{(ab-1)(c-1)}{2} + (\left\lfloor \frac{x}{c} \right\rfloor; ab) + 1)c \right) \\ &= \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left( \left\lfloor \frac{x}{abc} \right\rfloor - 1 \right) + x - c \left\lfloor \frac{x}{c} \right\rfloor + \frac{(ab-1)(c-1)}{2} + \left( \left\lfloor \frac{x}{c} \right\rfloor - ab \left\lfloor \frac{x}{abc} \right\rfloor + 1 \right)c \right) \\ &= \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor - \frac{abc}{2} + x - c \left\lfloor \frac{x}{c} \right\rfloor + \frac{abc - ab - c + 1}{2} + c \left\lfloor \frac{x}{c} \right\rfloor - abc \left\lfloor \frac{x}{abc} \right\rfloor + c \right) \\ &= \left\lfloor \frac{x}{abc} \right\rfloor \left( x - \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor + \frac{c - ab + 1}{2} \right) = \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor + \{x; abc\} + \frac{c - ab + 1}{2} \right) \end{aligned}$$

$$\begin{aligned} \text{Finally we get } S1 &= \left\lfloor \frac{x}{abc} \right\rfloor \left( \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor + \{x; abc\} + \frac{c - ab + 1}{2} \right) + \sum_{j=0}^{\{\lfloor \frac{x}{c} \rfloor; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor \\ &= \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor^2 + \left\lfloor \frac{x}{abc} \right\rfloor \left( \{x; abc\} + \frac{c - ab + 1}{2} \right) + \sum_{j=0}^{\{\lfloor \frac{x}{c} \rfloor; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor \end{aligned}$$

Notice that the main term is  $\frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor^2$  which has order  $\sim \frac{x^2}{2abc}$ .

The entire result is then  $S1 + S2$

$$\begin{aligned} &= \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor^2 + \left\lfloor \frac{x}{abc} \right\rfloor \left( \{x; abc\} + \frac{c - ab + 1}{2} \right) + \sum_{j=0}^{\{\lfloor \frac{x}{c} \rfloor; ab\}} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor + \\ &\left\lfloor \frac{x}{abc} \right\rfloor \sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{\{x; c\} + jc; ab\}) + \sum_{j=0}^{\{y; ab\}} \epsilon_{\langle a, b \rangle}(\{\{x; c\} + jc; ab\}). \end{aligned}$$

Each of the sums have easy to find bounds:

- $\{x; abc\} + \frac{c - ab + 1}{2} < abc + \frac{c - ab + 1}{2}.$

- $\sum_{j=0}^{\lfloor \frac{x}{c} \rfloor; ab} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor \leq \sum_{j=0}^{ab-1} \left\lfloor \frac{\{x; c\} + jc}{ab} \right\rfloor = \{x; c\} + \frac{(ab-1)(c-1)}{2} \leq c + \frac{(ab-1)(c-1)}{2}.$
- $\sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{\{x; c\} + jc; ab\}) \leq \sum_{j=0}^{ab-1} 1 = ab$
- $\sum_{j=0}^{\{y; ab\}} \epsilon_{\langle a, b \rangle}(\{\{x; c\} + jc; ab\}) \leq \sum_{j=0}^{ab-1} \epsilon_{\langle a, b \rangle}(\{\{x; c\} + jc; ab\}) \leq ab$

Hence we obtain:

**Theorem 13.**  $n = 3$ ,  $M = \langle a, b, c \rangle$ ,  $\gcd(a, b) = \gcd(a, c) = \gcd(b, c) = 1$ . We have

$$T_M(x) = \frac{abc}{2} \left\lfloor \frac{x}{abc} \right\rfloor^2 + r_M(x)$$

where  $|r_M(x)| \leq Kx + L$  for absolute constants  $K, L$  (depending only on  $a, b, c$ ).

## 5 Dimension $n$ arbitrary

The technique above generalizes, by summing up the leading terms using Corollary 4 obtaining:

**Theorem 14.** For  $M = \langle a_1, \dots, a_d \rangle$ ,  $\gcd(a_i, a_j) = 1$  for all  $i \neq j$ . The counting function  $T_M(x)$  has main order

$$T_M(x) \sim \frac{(a_1 \dots a_d)^{d-2}}{(d-1)!} \left\lfloor \frac{x}{a_1 \dots a_d} \right\rfloor^{d-1}$$

meaning that  $T_M(x) = \frac{(a_1 \dots a_d)^{d-2}}{(d-1)!} \left\lfloor \frac{x}{a_1 \dots a_d} \right\rfloor^{d-1} + r_M(x)$ , where  $|r_M(x)|$  is bounded by a polynomial in  $x$  (depending only on  $M$ ) of degree at most  $d-2$ .

## References

- [1] Beck, M.; De Loera, J. A.; Develin, M.; Pfeifle, J.; Stanley, R. P. (2005), “Coefficients and roots of Ehrhart polynomials”, Integer points in polyhedra- geometry, number theory, algebra, optimization, Contemp. Math., 374, Providence, RI: Amer. Math. Soc., pp. 15-36
- [2] J. C. Rosales, J. C and P. A. García-Sánchez, P. A. Numerical semigroups. Developments in Mathematics, 20. Springer, New York, 2009

Francisco Franco Munoz, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WASHINGTON, SEATTLE, WA 98195

E-mail address: `ffm1@uw.edu`