# On binomial coefficients modulo squares of primes

## Darij Grinberg

March 14, 2022

**Abstract.** We give elementary proofs for the Apagodu-Zeilberger-Stanton-Amdeberhan-Tauraso congruences

$$\sum_{n=0}^{p-1} {2n \choose n} \equiv \eta_p \bmod p^2;$$

$$\sum_{n=0}^{rp-1} {2n \choose n} \equiv \eta_p \sum_{n=0}^{r-1} {2n \choose n} \bmod p^2;$$

$$\sum_{n=0}^{rp-1} \sum_{m=0}^{sp-1} {n+m \choose m}^2 \equiv \eta_p \sum_{m=0}^{r-1} \sum_{n=0}^{s-1} {n+m \choose m}^2 \bmod p^2,$$

where p is an odd prime, r and s are nonnegative integers, and

$$\eta_p = \begin{cases} 0, & \text{if } p \equiv 0 \mod 3; \\ 1, & \text{if } p \equiv 1 \mod 3; \\ -1, & \text{if } p \equiv 2 \mod 3 \end{cases}$$

### **Contents**

1.	Introduction		
	1.1.	Binomial coefficients	2
	1.2.	Classical congruences	3
	1.3.	The three modulo- $p^2$ congruences	4
		proofs	Ę
	2.1.	Identities and congruences from the literature	5
	2.2.	Variants and consequences of Vandermonde convolution	7
	2.3.	A congruence of Bailey's	11
	2.4.	Two congruences for polynomials	13

2.5.	Proving Theorem 1.8	15
2.6.	Proving Theorem 1.9	18
2.7.	Proving Theorem 1.10	20
2.8	Acknowledgments	25

#### 1. Introduction

In this note, we prove that any odd prime p and any  $r,s \in \mathbb{N}$  satisfy

$$\sum_{n=0}^{p-1} \binom{2n}{n} \equiv \eta_p \bmod p^2 \qquad \text{(Theorem 1.8)};$$

$$\sum_{n=0}^{rp-1} \binom{2n}{n} \equiv \eta_p \sum_{n=0}^{r-1} \binom{2n}{n} \bmod p^2 \qquad \text{(Theorem 1.9)};$$

$$\sum_{n=0}^{rp-1} \sum_{m=0}^{sp-1} \binom{n+m}{m}^2 \equiv \eta_p \sum_{m=0}^{r-1} \sum_{n=0}^{s-1} \binom{n+m}{m}^2 \bmod p^2 \qquad \text{(Theorem 1.10)},$$

where

$$\eta_p = \begin{cases} 0, & \text{if } p \equiv 0 \mod 3; \\ 1, & \text{if } p \equiv 1 \mod 3; \\ -1, & \text{if } p \equiv 2 \mod 3 \end{cases}$$

These three congruences are (slightly extended versions of) three of the "Super-Conjectures" (namely, 1, 1" and 4′) stated by Apagodu and Zeilberger in [ApaZei16]<sup>1</sup>. Our proofs are more elementary than previous proofs by Stanton [Stanto16] and Amdeberhan and Tauraso [AmdTau16].

#### 1.1. Binomial coefficients

Let us first recall the definition of binomial coefficients:<sup>2</sup>

**Definition 1.1.** Let  $n \in \mathbb{N}$  and  $m \in \mathbb{Z}$ . Then, the *binomial coefficient*  $\binom{m}{n}$  is a rational number defined by

$$\binom{m}{n} = \frac{m(m-1)\cdots(m-n+1)}{n!}.$$

<sup>&</sup>lt;sup>1</sup>In the arXiv preprint version of [ApaZei16] (arXiv:1606.03351v2), these congruences appear as "Super-Conjectures" 1, 1" and 5′, respectively.

<sup>&</sup>lt;sup>2</sup>We use the notation  $\mathbb{N}$  for the set  $\{0, 1, 2, \ldots\}$ .

**Definition 1.2.** Let n be a negative integer. Let  $m \in \mathbb{Z}$ . Then, the *binomial coefficient*  $\binom{m}{n}$  is a rational number defined by  $\binom{m}{n} = 0$ .

(This is the definition used in [GrKnPa94] and [Grinbe17b]. Some authors follow other conventions instead.)

The following proposition is well-known (see, e.g., [Grinbe17b, Proposition 1.9]):

**Proposition 1.3.** We have 
$$\binom{m}{n} \in \mathbb{Z}$$
 for any  $m \in \mathbb{Z}$  and  $n \in \mathbb{Z}$ .

Proposition 1.3 shows that  $\binom{m}{n}$  is an integer whenever  $m \in \mathbb{Z}$  and  $n \in \mathbb{Z}$ . We shall tacitly use this below, when we study congruences involving binomial coefficients.

One advantage of Definition 1.2 is that it makes the following hold:

**Proposition 1.4.** For any  $n \in \mathbb{Z}$  and  $m \in \mathbb{Z}$ , the binomial coefficient  $\binom{n}{m}$  is the coefficient of  $X^m$  in the formal power series  $(1+X)^n \in \mathbb{Z}[[X]]$ . (Here, the coefficient of  $X^m$  in any formal power series is defined to be 0 when m is negative.)

#### 1.2. Classical congruences

The behavior of binomial coefficients modulo primes and prime powers is a classical subject of research; see [Mestro14, §2.1] for a survey of much of it. Let us state two of the most basic results in this subject:

**Theorem 1.5.** Let p be a prime. Let a and b be two integers. Let c and d be two elements of  $\{0,1,\ldots,p-1\}$ . Then,

$$\binom{ap+c}{bp+d} \equiv \binom{a}{b} \binom{c}{d} \mod p.$$

Theorem 1.5 is known under the name of *Lucas's theorem*, and is proven in many places (e.g., [Mestro14, §2.1] or [Hausne83, Proof of §4] or [AnBeRo05, proof of Lucas's theorem] or [GrKnPa94, Exercise 5.61]) at least in the case when *a* and *b* are nonnegative integers. The standard proof of Theorem 1.5 in this case uses generating functions (specifically, Proposition 1.4); this proof applies (mutatis mutandis) in the general case as well. See [Grinbe17b, Theorem 1.11] for an elementary proof of Theorem 1.5.

Another fundamental result is the following:

**Theorem 1.6.** Let p be a prime. Let a and b be two integers. Then,

$$\binom{ap}{bp} \equiv \binom{a}{b} \bmod p^2.$$

Theorem 1.6 is a known result, perhaps due to Charles Babbage. It appears with proof in [Grinbe17b, Theorem 1.12]; again, many sources prove it for nonnegative a and b (for example [Stanle11, Exercise 1.14 c] or [GrKnPa94, Exercise 5.62]). Notice that if  $p \ge 5$ , then the modulus  $p^2$  can be replaced by  $p^3$  or (depending on a, b and p) by even higher powers of p; see [Mestro14, (22) and (23)] for the details. See also [SunTau11, Lemma 2.1] for another strengthening of Theorem 1.6.

## 1.3. The three modulo- $p^2$ congruences

**Definition 1.7.** For any  $p \in \mathbb{Z}$ , we define an integer  $\eta_p \in \{-1,0,1\}$  by

$$\eta_p = \begin{cases}
0, & \text{if } p \equiv 0 \mod 3; \\
1, & \text{if } p \equiv 1 \mod 3; \\
-1, & \text{if } p \equiv 2 \mod 3
\end{cases}$$

Notice that  $\eta_p$  is the so-called *Legendre symbol*  $\left(\frac{p}{3}\right)$  known from number theory. We are now ready to state three conjectures by Apagodu and Zeilberger, which we shall prove in the sequel. The first one is [ApaZei16, Super-Conjecture 1]:<sup>3</sup>

**Theorem 1.8.** Let *p* be an odd prime. Then,

$$\sum_{n=0}^{p-1} \binom{2n}{n} \equiv \eta_p \bmod p^2.$$

The next one ([ApaZei16, Super-Conjecture 1"]) is a generalization:

**Theorem 1.9.** Let p be an odd prime. Let  $r \in \mathbb{N}$ . Set

$$\alpha_r = \sum_{n=0}^{r-1} \binom{2n}{n}.$$

<sup>&</sup>lt;sup>3</sup>To be precise (and boastful), our Theorem 1.8 is somewhat stronger than [ApaZei16, Super-Conjecture 1], since we only require p to be odd (rather than  $p \ge 5$ ). Of course, in the case of Theorem 1.8, this extra generality is insignificant, since it just adds the possibility of p = 3, in which case Theorem 1.8 can be checked by hand. However, for Theorems 1.9 and 1.10 further below, we gain somewhat more from this generality.

Then,

$$\sum_{n=0}^{rp-1} \binom{2n}{n} \equiv \eta_p \alpha_r \operatorname{mod} p^2.$$

Theorem 1.8 and Theorem 1.9 both have been proven by Dennis Stanton [Stanto16] using Laurent series (in the case when  $p \ge 5$ ), and by Liu [Liu16, (1.3)] using harmonic numbers. We shall reprove them elementarily.

The third conjecture that we shall prove is [ApaZei16, Super-Conjecture 5']:

**Theorem 1.10.** Let p be an odd prime. Let  $r \in \mathbb{N}$  and  $s \in \mathbb{N}$ . Set

$$\epsilon_{r,s} = \sum_{m=0}^{r-1} \sum_{n=0}^{s-1} {n+m \choose m}^2.$$

Then,

$$\sum_{n=0}^{rp-1} \sum_{m=0}^{sp-1} \binom{n+m}{m}^2 \equiv \eta_p \epsilon_{r,s} \bmod p^2.$$

A proof of Theorem 1.10 has been found by Amdeberhan and Tauraso, and was outlined in [AmdTau16, §6]; we give a different, elementary proof.

## 2. The proofs

## 2.1. Identities and congruences from the literature

Before we come to the proofs of Theorems 1.8, 1.9 and 1.10, let us collect various well-known results that will prove useful.

The following properties of binomial coefficients are well-known (see, e.g., [Grinbe17, §3.1] and [Grinbe17b, §1]):

**Proposition 2.1.** We have 
$$\binom{m}{0} = 1$$
 for every  $m \in \mathbb{Z}$ .

**Proposition 2.2.** We have 
$$\binom{m}{n} = 0$$
 for every  $m \in \mathbb{N}$  and  $n \in \mathbb{N}$  satisfying  $m < n$ .

**Proposition 2.3.** We have 
$$\binom{m}{n} = \binom{m}{m-n}$$
 for any  $m \in \mathbb{N}$  and  $n \in \mathbb{N}$  satisfying  $m \ge n$ .

**Proposition 2.4.** We have  $\binom{m}{m} = 1$  for every  $m \in \mathbb{N}$ .

**Proposition 2.5.** We have

$$\binom{m}{n} = (-1)^n \binom{n-m-1}{n}$$

for any  $m \in \mathbb{Z}$  and  $n \in \mathbb{N}$ .

**Proposition 2.6.** We have

$$\binom{m}{n} = \binom{m-1}{n-1} + \binom{m-1}{n}$$

for any  $m \in \mathbb{Z}$  and  $n \in \mathbb{Z}$ .

**Proposition 2.7.** For every  $x \in \mathbb{Z}$  and  $y \in \mathbb{Z}$  and  $n \in \mathbb{N}$ , we have

$$\binom{x+y}{n} = \sum_{k=0}^{n} \binom{x}{k} \binom{y}{n-k}.$$

Proposition 2.7 is the so-called *Vandermonde convolution identity*, and is a particular case of [Grinbe17, Theorem 3.29].

**Corollary 2.8.** For each  $n \in \mathbb{N}$ , we have

$$\sum_{i=0}^{n-1} (-1)^i \binom{n-1-i}{i} = (-1)^n \cdot \begin{cases} 0, & \text{if } n \equiv 0 \mod 3; \\ -1, & \text{if } n \equiv 1 \mod 3; \\ 1, & \text{if } n \equiv 2 \mod 3 \end{cases}$$

Corollary 2.8 is [Grinbe17, Corollary 8.63]. Apart from that, Corollary 2.8 can be easily derived from [GrKnPa94, §5.2, Problem 3], [BenQui03, Identity 172] or [BenQui08].

Another simple identity (sometimes known as the "absorption identity") is the following:

**Proposition 2.9.** Let 
$$n \in \mathbb{Z}$$
 and  $k \in \mathbb{Z}$ . Then,  $k \binom{n}{k} = n \binom{n-1}{k-1}$ .

Proposition 2.9 appears in [GrKnPa94, (5.6)], and is easily proven just from the definition of binomial coefficients.

Finally, we need the following result from elementary number theory:

**Theorem 2.10.** Let p be a prime. Let  $k \in \mathbb{N}$ . Assume that k is not a positive multiple of p-1. Then,

$$\sum_{l=0}^{p-1} l^k \equiv 0 \bmod p.$$

Theorem 2.10 is proven, e.g., in [Grinbe17b, Theorem 3.1] and (in a slightly rewritten form) in [MacSon10, Theorem 1].

#### 2.2. Variants and consequences of Vandermonde convolution

We are now going to state a number of identities that are restatements or particular cases of the Vandermonde convolution identity (Proposition 2.7). We begin with the following one:

**Corollary 2.11.** Let  $u \in \mathbb{Z}$  and  $l \in \mathbb{N}$  and  $w \in \mathbb{N}$ . Then,

$$\sum_{m=0}^{l} \binom{u}{w+m} \binom{l}{m} = \binom{u+l}{w+l}.$$

*Proof of Corollary 2.11.* Proposition 2.7 (applied to x = u, y = l and n = w + l)

yields

This proves Corollary 2.11.

Let us also state another corollary of Proposition 2.7:

 $=\sum_{n=0}^{l}\binom{u}{w+m}\binom{l}{m}.$ 

**Corollary 2.12.** Let  $x \in \mathbb{Z}$  and  $y \in \mathbb{N}$  and  $n \in \mathbb{Z}$ . Then,

$$\binom{x+y}{n} = \sum_{i=0}^{y} \binom{x}{n-i} \binom{y}{i}.$$

See [Grinbe17b, Corollary 2.2] for a proof of Corollary 2.12.

**Lemma 2.13.** Let  $u \in \mathbb{Z}$  and  $w \in \mathbb{N}$  and  $l \in \mathbb{N}$ . Then,

$$\binom{u+2l}{w+l} = \binom{u}{w} \binom{2l}{l} + \sum_{i=1}^{l} \left( \binom{u}{w+i} + \binom{u}{w-i} \right) \binom{2l}{l-i}.$$

*Proof of Lemma 2.13.* Corollary 2.12 (applied to x = u, y = 2l and n = w + l) yields

$$\binom{u+2l}{w+l} = \sum_{i=0}^{2l} \binom{u}{w+l-i} \binom{2l}{i} = \sum_{i=-l}^{l} \binom{u}{w+i} \binom{2l}{l-i}$$

(here, we have substituted l - i for i in the sum)

$$= \sum_{\substack{i \in \{-l, -l+1, \dots, l\}; \\ i \neq 0}} \binom{u}{w+i} \binom{2l}{l-i} + \binom{u}{w} \binom{2l}{l}$$

(here, we have split off the addend for i = 0 from the sum). Hence,

In other words,

$$\binom{u+2l}{w+l} = \binom{u}{w} \binom{2l}{l} + \sum_{i=1}^{l} \left( \binom{u}{w+i} + \binom{u}{w-i} \right) \binom{2l}{l-i}.$$

This proves Lemma 2.13.

**Lemma 2.14.** Let  $p \in \mathbb{N}$ . Let  $c \in \mathbb{Z}$ . Let  $l \in \{0, 1, ..., p - 1\}$ . Then,

$$\binom{cp+2l}{l} = \sum_{k=0}^{p-1} \binom{cp+l}{k} \binom{l}{k}.$$

*Proof of Lemma 2.14.* Corollary 2.12 (applied to x = cp + l, y = l and n = l) yields

(here, we have substituted k for l - i in the sum)

$$=\sum_{k=0}^{l}\binom{cp+l}{k}\binom{l}{k}.$$

Comparing this with

$$\sum_{k=0}^{p-1} \binom{cp+l}{k} \binom{l}{k} = \sum_{k=0}^{l} \binom{cp+l}{k} \binom{l}{k} + \sum_{k=l+1}^{p-1} \binom{cp+l}{k} \underbrace{\binom{l}{k}}_{=0}$$
(by Proposition 2.2 (applied to  $m=l$  and  $n=k$ )
(since  $l < k$ )

(here, we have split the sum at k = l, since  $0 \le l \le p - 1$ )

$$=\sum_{k=0}^{l}\binom{cp+l}{k}\binom{l}{k}+\sum_{k=l+1}^{p-1}\binom{cp+l}{k}0=\sum_{k=0}^{l}\binom{cp+l}{k}\binom{l}{k},$$

we obtain  $\sum_{k=0}^{p-1} {cp+l \choose k} {l \choose k} = {cp+l+l \choose l} = {cp+2l \choose l}$ . This proves Lemma 2.14.

**Lemma 2.15.** Let  $p \in \mathbb{N}$ . Let  $l \in \mathbb{N}$ . Then,

$$\sum_{i=1}^{l} \binom{p}{i} \binom{2l}{l-i} = \binom{p+2l}{l} - \binom{2l}{l}.$$

*Proof of Lemma 2.15.* Proposition 2.7 (applied to x = p, y = 2l and n = l) yields

$$\binom{p+2l}{l} = \sum_{k=0}^{l} \binom{p}{k} \binom{2l}{l-k} = \sum_{i=0}^{l} \binom{p}{i} \binom{2l}{l-i}$$

(here, we have renamed the summation index k as i)

$$=\underbrace{\binom{p}{0}}_{=1}\underbrace{\binom{2l}{l-0}}_{=\binom{2l}{l}} + \sum_{i=1}^{l} \binom{p}{i} \binom{2l}{l-i} = \binom{2l}{l} + \sum_{i=1}^{l} \binom{p}{i} \binom{2l}{l-i}.$$

Thus,

$$\sum_{i=1}^{l} \binom{p}{i} \binom{2l}{l-i} = \binom{p+2l}{l} - \binom{2l}{l}.$$

This proves Lemma 2.15.

#### 2.3. A congruence of Bailey's

Next, we shall prove a modulo- $p^2$  congruence for certain binomial coefficients that can be regarded as a counterpart to Theorem 1.6:

**Theorem 2.16.** Let p be a prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{Z}$  and  $i \in \{1, 2, ..., p-1\}$ . Then:

(a) We have

$$\binom{Np}{Kp+i} \equiv N\binom{N-1}{K}\binom{p}{i} \bmod p^2.$$

(b) We have

$$\binom{Np}{Kp-i} \equiv N \binom{N-1}{K-1} \binom{p}{i} \bmod p^2.$$

(c) We have

$$\binom{Np}{Kp+i} + \binom{Np}{Kp-i} \equiv N\binom{N}{K}\binom{p}{i} \bmod p^2.$$

Theorem 2.16 (a) is essentially the result [Bailey91, Theorem 4] by Bailey (see also [Mestro14, (26)]); in fact, it transforms into [Bailey91, Theorem 4] if we rewrite  $N\binom{N-1}{K}$  as  $(K+1)\binom{N}{K+1}$  (using Proposition 2.9). We shall nevertheless give our own proof.

*Proof of Theorem* 2.16. From  $i \in \{1, 2, ..., p-1\}$ , we conclude that both i-1 and p-i are elements of  $\{0, 1, ..., p-1\}$ . Notice also that i is not divisible by p (since  $i \in \{1, 2, ..., p-1\}$ ); hence, i is coprime to p (since p is a prime). Therefore, i is also coprime to  $p^2$ .

(a) Proposition 2.9 (applied to n = Np and k = Kp + i) yields

Troposition 2.5 (applied to 
$$N = Np$$
 and  $N = Np$ ) yields
$$(Kp+i) \binom{Np}{Kp+i} = Np \binom{Np-1}{Kp+i-1} = Np \underbrace{\binom{(N-1)p+(p-1)}{Kp+(i-1)}}_{\text{$Kp+(i-1)$}}$$

$$\equiv \binom{N-1}{K} \binom{p-1}{i-1} \mod p$$
(by Theorem 1.5, applied to  $a=N-1, b=K, c=p-1 \text{ and } d=i-1$ )
$$\equiv Np \binom{N-1}{K} \binom{p-1}{i-1} \mod p^2 \tag{1}$$

(notice that the presence of the p factor has turned a congruence modulo p into a congruence modulo  $p^2$ ). Thus,

$$(Kp+i) \binom{Np}{Kp+i} \equiv Np \binom{N-1}{K} \binom{p-1}{i-1} \equiv 0 \mod p,$$

so that  $0 \equiv \underbrace{(Kp+i)}_{\equiv i \bmod p} \binom{Np}{Kp+i} \equiv i \binom{Np}{Kp+i} \bmod p$ . We can cancel i from this

congruence (since i is coprime to p), and thus obtain  $0 \equiv \binom{Np}{Kp+i} \mod p$ . Hence,  $\binom{Np}{Kp+i}$  is divisible by p. Thus,  $p\binom{Np}{Kp+i}$  is divisible by  $p^2$ . In other words,

$$p\binom{Np}{Kp+i} \equiv 0 \bmod p^2. \tag{2}$$

Now,

$$(Kp+i) \binom{Np}{Kp+i} = K \underbrace{p \binom{Np}{Kp+i}}_{\equiv 0 \bmod p^2} + i \binom{Np}{Kp+i} \equiv i \binom{Np}{Kp+i} \bmod p^2.$$

Hence,

$$i\binom{Np}{Kp+i} \equiv (Kp+i)\binom{Np}{Kp+i} \equiv Np\binom{N-1}{K}\binom{p-1}{i-1} \qquad \text{(by (1))}$$

$$= N\binom{N-1}{K} \quad p\binom{p-1}{i-1} = N\binom{N-1}{K}i\binom{p}{i} \bmod p^{2}.$$

$$= i\binom{p}{i}$$
(by Proposition 2.9)

We can cancel i from this congruence (since i is coprime to  $p^2$ ), and thus obtain

$$\binom{Np}{Kp+i} \equiv N\binom{N-1}{K}\binom{p}{i} \bmod p^2.$$

This proves Theorem 2.16 (a).

**(b)** We have  $i \in \{1, 2, ..., p-1\}$  and thus  $p-i \in \{1, 2, ..., p-1\}$ . Hence, Theorem 2.16 **(a)** (applied to K-1 and p-i instead of K and i) yields

$$\binom{Np}{(K-1)p+(p-i)} \equiv N\binom{N-1}{K-1} \qquad \underbrace{\binom{p}{p-i}}_{=\binom{p}{i}} = N\binom{N-1}{K-1}\binom{p}{i} \bmod p^{2}.$$
(by Proposition 2.3)

In view of (K-1) p + (p-i) = Kp - i, this rewrites as

$$\binom{Np}{Kp-i} \equiv N \binom{N-1}{K-1} \binom{p}{i} \bmod p^2.$$

This proves Theorem 2.16 (b).

(c) We have

$$\underbrace{\begin{pmatrix} Np \\ Kp+i \end{pmatrix}}_{\text{K}p+i} + \underbrace{\begin{pmatrix} Np \\ Kp-i \end{pmatrix}}_{\text{K}p-i}$$

$$\equiv N \binom{N-1}{K} \binom{p}{i} \mod p^2 \quad \equiv N \binom{N-1}{K-1} \binom{p}{i} \mod p^2$$
(by Theorem 2.16 (a)) (by Theorem 2.16 (b))
$$\equiv N \binom{N-1}{K} \binom{p}{i} + N \binom{N-1}{K-1} \binom{p}{i}$$

$$= N \underbrace{\binom{N-1}{K} + \binom{N-1}{K}}_{\text{K}} \binom{p}{i} = N \binom{N}{K} \binom{p}{i} \mod p^2.$$

$$= \binom{N}{K}$$
(by Proposition 2.6)

This proves Theorem 2.16 (c).

#### 2.4. Two congruences for polynomials

Now, we recall that  $\mathbb{Z}[X]$  is the ring of all polynomials in one indeterminate X with integer coefficients.

**Lemma 2.17.** Let 
$$p$$
 be a prime. Let  $c \in \mathbb{Z}$ . Let  $P \in \mathbb{Z}[X]$  be a polynomial of degree  $< 2p - 1$ . Then,  $\sum_{l=0}^{p-1} (P(cp+l) - P(l)) \equiv 0 \mod p^2$ .

*Proof of Lemma* 2.17. WLOG assume that  $P = X^k$  for some  $k \in \{0, 1, ..., 2p - 2\}$  (since the congruence we are proving depends  $\mathbb{Z}$ -linearly on P). If k = 0, then Lemma 2.17 is easily checked (because in this case, P is constant). Thus, WLOG assume that  $k \neq 0$ . Hence, k is a positive integer (since  $k \in \mathbb{N}$ ). Thus,  $k - 1 \in \mathbb{N}$ .

Each  $l \in \{0, 1, \dots, p-1\}$  satisfies

$$P(cp+l) = (cp+l)^{k} \qquad \left(\text{since } P = X^{k}\right)$$

$$= \sum_{i=0}^{k} \binom{k}{i} (cp)^{i} l^{k-i} \qquad \text{(by the binomial formula)}$$

$$= \underbrace{(cp)^{0} l^{k-0}}_{=l^{k}} + k \underbrace{(cp)^{1}}_{=cp} l^{k-1} + \sum_{i=2}^{k} \binom{k}{i} \underbrace{(cp)^{i}}_{\text{since } i \geq 2} l^{k-i}$$

$$\equiv l^{k} + kcpl^{k-1} + \sum_{i=2}^{k} \binom{k}{i} 0l^{k-i} = l^{k} + kcpl^{k-1} \bmod p^{2}$$

and  $P(l) = l^k$  (since  $P = X^k$ ). Thus,

$$\sum_{l=0}^{p-1} \left( \underbrace{P(cp+l)}_{\equiv l^k + kcpl^{k-1} \bmod p^2} - \underbrace{P(l)}_{=l^k} \right) \equiv \sum_{l=0}^{p-1} \underbrace{\left(l^k + kcpl^{k-1} - l^k\right)}_{=kcpl^{k-1}} = kcp \sum_{l=0}^{p-1} l^{k-1} \bmod p^2.$$

The claim of Lemma 2.17 now becomes obvious if k=p (because if k=p, then kcp is already divisible by  $p^2$ ); thus, we WLOG assume that  $k\neq p$ . Hence,  $k-1\neq p-1$ . If k-1 was a positive multiple of p-1, then we would have k-1=p-1 (since  $k\in\{0,1,\ldots,2p-2\}$ ), which would contradict  $k-1\neq p-1$ . Hence, k-1 is not a positive multiple of p-1. Thus, Theorem 2.10 (applied to k-1 instead of k) yields  $\sum_{l=0}^{p-1} l^{k-1} \equiv 0 \bmod p.$  Thus,  $p\sum_{l=0}^{p-1} l^{k-1} \equiv 0 \bmod p^2$ , so that

$$\sum_{l=0}^{p-1} \left( P\left( cp+l \right) - P\left( l \right) \right) \equiv kc \, \underbrace{p \sum_{l=0}^{p-1} l^{k-1}}_{\equiv 0 \, \mathrm{mod} \, p^2} \equiv 0 \, \mathrm{mod} \, p^2.$$

This proves Lemma 2.17.

**Lemma 2.18.** Let p, a and b be three integers such that a - b is divisible by p. Then,  $a^2 - b^2 \equiv 2 (a - b) b \mod p^2$ .

*Proof of Lemma 2.18.* The difference  $(a^2 - b^2) - 2(a - b)b = (a - b)^2$  is divisible by  $p^2$  (since a - b is divisible by p). In other words,  $a^2 - b^2 \equiv 2(a - b)b \mod p^2$ . Lemma 2.18 is proven.

**Lemma 2.19.** Let p be an odd prime. Let  $c \in \mathbb{Z}$ . Let  $P \in \mathbb{Z}[X]$  be a polynomial of degree  $\leq p-1$ . Then,

$$\sum_{l=0}^{p-1} \left( P\left( cp+l \right) - P\left( l \right) \right) P\left( l \right) \equiv 0 \bmod p^2.$$

*Proof of Lemma* 2.19. Fix  $l \in \mathbb{Z}$ . We have  $P \in \mathbb{Z}[X]$ . Thus, P(u) - P(v) is divisible by u - v whenever u and v are two integers<sup>4</sup>. Applying this to u = cp + l and v = l, we conclude that P(cp + l) - P(l) is divisible by (cp + l) - l = cp, and thus also divisible by p.

Hence, Lemma 2.18 (applied to a = P(cp + l) and b = P(l)) shows that

$$(P(cp+l))^2 - (P(l))^2 \equiv 2(P(cp+l) - P(l))P(l) \bmod p^2.$$
 (3)

Now, forget that we fixed l. We thus have proven (3) for each  $l \in \mathbb{Z}$ .

The polynomial P has degree  $\leq p-1$ . Hence, the polynomial  $P^2$  has degree  $\leq 2(p-1) < 2p-1$ . Thus, Lemma 2.17 (applied to  $P^2$  instead of P) shows that

$$\sum_{l=0}^{p-1} \left( P^2 \left( cp + l \right) - P^2 \left( l \right) \right) \equiv 0 \operatorname{mod} p^2.$$

Thus,

$$0 \equiv \sum_{l=0}^{p-1} \underbrace{\left(P^2\left(cp+l\right) - P^2\left(l\right)\right)}_{=(P(cp+l))^2 - (P(l))^2} \equiv 2 \sum_{l=0}^{p-1} \left(P\left(cp+l\right) - P\left(l\right)\right) P\left(l\right) \bmod p^2.$$

$$= 2(P(cp+l) - P(l)) P(l) \bmod p^2$$
(by (3))

We can cancel 2 from this congruence (since p is odd), and conclude that

$$0 \equiv \sum_{l=0}^{p-1} \left( P\left( cp + l \right) - P\left( l \right) \right) P\left( l \right) \bmod p^{2}.$$

This proves Lemma 2.19.

#### 2.5. Proving Theorem 1.8

Now, let us prepare for the proofs of our results by showing several lemmas.

<sup>&</sup>lt;sup>4</sup>This is a well-known fact. It can be proven as follows: WLOG assume that  $P = X^k$  for some  $k \in \mathbb{N}$  (this is a valid assumption, since the claim is  $\mathbb{Z}$ -linear in P); then,  $P(u) - P(v) = u^k - v^k = (u - v) \sum_{i=0}^{k-1} u^i v^{k-i}$  is clearly divisible by u - v.

**Lemma 2.20.** Let p be an odd prime. Let  $c \in \mathbb{Z}$ . Let  $k \in \{0, 1, ..., p-1\}$ . Then,

$$\sum_{l=0}^{p-1} \left( \binom{cp+l}{k} - \binom{l}{k} \right) \binom{l}{k} \equiv 0 \bmod p^2.$$

*Proof of Lemma* 2.20. Notice that k! is coprime to p (since  $k \le p - 1$ ), and thus  $k!^2$  is coprime to  $p^2$ .

Define a polynomial  $P \in \mathbb{Z}[X]$  by

$$P = X(X-1)\cdots(X-k+1).$$

Then, *P* has degree  $k \le p - 1$ . Thus, Lemma 2.19 yields

$$\sum_{l=0}^{p-1} \left( P\left( cp+l \right) - P\left( l \right) \right) P\left( l \right) \equiv 0 \operatorname{mod} p^{2}.$$

Since each  $n \in \mathbb{Z}$  satisfies  $P(n) = n(n-1)\cdots(n-k+1) = k!\binom{n}{k}$ , this rewrites as

$$\sum_{l=0}^{p-1} \left( k! \binom{cp+l}{k} - k! \binom{l}{k} \right) k! \binom{l}{k} \equiv 0 \bmod p^2.$$

We can cancel  $k!^2$  from this congruence (since  $k!^2$  is coprime to  $p^2$ ), and thus obtain

$$\sum_{l=0}^{p-1} \left( \binom{cp+l}{k} - \binom{l}{k} \right) \binom{l}{k} \equiv 0 \bmod p^2.$$

This proves Lemma 2.20.

**Lemma 2.21.** Let p be an odd prime. Let  $c \in \mathbb{Z}$ . Then,

$$\sum_{l=0}^{p-1} \left( \binom{cp+2l}{l} - \binom{2l}{l} \right) \equiv 0 \bmod p^2.$$

*Proof of Lemma 2.21.* For each  $l \in \{0, 1, ..., p-1\}$ , we have

$$\underbrace{\begin{pmatrix} cp+2l \\ l \end{pmatrix}}_{l} - \underbrace{\begin{pmatrix} 2l \\ l \end{pmatrix}}_{l}$$

$$= \sum_{k=0}^{p-1} \binom{cp+l}{k} \binom{l}{k} = \sum_{k=0}^{p-1} \binom{l}{k} \binom{l}{k}$$
(by Lemma 2.14) (by Lemma 2.14, applied to 0 instead of c)
$$= \sum_{k=0}^{p-1} \binom{cp+l}{k} \binom{l}{k} - \sum_{k=0}^{p-1} \binom{l}{k} \binom{l}{k} = \sum_{k=0}^{p-1} \binom{cp+l}{k} - \binom{l}{k} \binom{l}{k}.$$

Summing these equalities over all  $l \in \{0, 1, ..., p - 1\}$ , we find

$$\sum_{l=0}^{p-1} \left( \binom{cp+2l}{l} - \binom{2l}{l} \right) = \sum_{l=0}^{p-1} \sum_{k=0}^{p-1} \left( \binom{cp+l}{k} - \binom{l}{k} \right) \binom{l}{k}$$

$$= \sum_{k=0}^{p-1} \sum_{l=0}^{p-1} \left( \binom{cp+l}{k} - \binom{l}{k} \right) \binom{l}{k} \equiv \sum_{k=0}^{p-1} 0 = 0 \mod p^2.$$

$$\stackrel{\equiv 0 \bmod p^2}{\text{(by Lemma 2.20)}}$$

This proves Lemma 2.21.

*Proof of Theorem 1.8.* Lemma 2.21 (applied to c = -1) yields

$$\sum_{l=0}^{p-1} \left( \binom{-p+2l}{l} - \binom{2l}{l} \right) \equiv 0 \bmod p^2.$$

Thus,

$$0 \equiv \sum_{l=0}^{p-1} \left( \binom{-p+2l}{l} - \binom{2l}{l} \right) = \sum_{l=0}^{p-1} \binom{-p+2l}{l} - \sum_{l=0}^{p-1} \binom{2l}{l} \bmod p^2,$$

so that

$$\sum_{l=0}^{p-1} {2l \choose l} \equiv \sum_{l=0}^{p-1} {-p+2l \choose l} \bmod p^2. \tag{4}$$

Now,

$$\sum_{n=0}^{p-1} \binom{2n}{n} = \sum_{l=0}^{p-1} \binom{2l}{l} \equiv \sum_{l=0}^{p-1} \underbrace{\begin{pmatrix} -p+2l \\ l \end{pmatrix}}_{\text{(by Proposition 2.5)}}$$

$$= \sum_{l=0}^{p-1} (-1)^l \underbrace{\begin{pmatrix} l - (-p+2l) - 1 \\ l \end{pmatrix}}_{\text{(by Proposition 2.5)}} = \sum_{l=0}^{p-1} (-1)^l \binom{p-1-l}{l}$$

$$= \sum_{i=0}^{p-1} (-1)^i \binom{p-1-i}{i} = \underbrace{\begin{pmatrix} -1 \end{pmatrix}^p}_{\text{(since $p$ is odd)}} \cdot \begin{cases} 0, & \text{if $p \equiv 0 \bmod 3$;} \\ -1, & \text{if $p \equiv 1 \bmod 3$;} \\ 1, & \text{if $p \equiv 1 \bmod 3$;} \end{cases} = \begin{cases} 0, & \text{if $p \equiv 0 \bmod 3$;} \\ -1, & \text{if $p \equiv 1 \bmod 3$;} \\ 1, & \text{if $p \equiv 1 \bmod 3$;} \end{cases} = \begin{cases} 0, & \text{if $p \equiv 0 \bmod 3$;} \\ -1, & \text{if $p \equiv 1 \bmod 3$;} \\ -1, & \text{if $p \equiv 1 \bmod 3$;} \end{cases} = \begin{cases} 1, & \text{if $p \equiv 1 \bmod 3$;} \\ -1, & \text{if $p \equiv 2 \bmod 3$} \end{cases}$$

This proves Theorem 1.8.

#### 2.6. Proving Theorem 1.9

**Lemma 2.22.** Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{N}$ . Let p be a prime. Let  $l \in \{0, 1, ..., p-1\}$ . Then,

$$\binom{Np+2l}{Kp+l} - \binom{N}{K} \binom{2l}{l} \equiv N \binom{N}{K} \left( \binom{p+2l}{l} - \binom{2l}{l} \right) \bmod p^2.$$

*Proof of Lemma 2.22.* Theorem 1.6 yields  $\binom{Np}{Kp} \equiv \binom{N}{K} \mod p^2$ .

Lemma 2.13 (applied to u = Np and w = Kp) yields

$$\binom{Np+2l}{Kp+l} = \binom{Np}{Kp} \binom{2l}{l} + \sum_{i=1}^{l} \underbrace{\binom{Np}{Kp+i} + \binom{Np}{Kp-i}}_{Kp-i} \binom{2l}{l-i}$$

$$\equiv \binom{N}{K} \mod p^{2} \qquad \qquad \equiv N\binom{N}{K} \binom{p}{i} \mod p^{2}$$

$$\equiv \binom{N}{K} \binom{2l}{l} + \sum_{i=1}^{l} N\binom{N}{K} \binom{p}{i} \binom{2l}{l-i}$$

$$= \binom{N}{K} \binom{2l}{l} + N\binom{N}{K} \sum_{i=1}^{l} \binom{p}{i} \binom{2l}{l-i}$$

$$= \binom{N}{K} \binom{2l}{l} + N\binom{N}{K} \binom{p}{k} \binom{2l}{l-i}$$

$$= \binom{N}{K} \binom{2l}{l} + N\binom{N}{K} \binom{p+2l}{l} - \binom{2l}{l} \pmod p^{2}.$$

Subtracting  $\binom{N}{K}\binom{2l}{l}$  from both sides of this congruence, we obtain

$$\binom{Np+2l}{Kp+l} - \binom{N}{K} \binom{2l}{l} \equiv N \binom{N}{K} \left( \binom{p+2l}{l} - \binom{2l}{l} \right) \bmod p^2.$$

This proves Lemma 2.22.

**Lemma 2.23.** Let p be an odd prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{N}$ . Then,

$$\sum_{l=0}^{p-1} \binom{Np+2l}{Kp+l} \equiv \binom{N}{K} \eta_p \operatorname{mod} p^2.$$

*Proof of Lemma 2.23.* For any  $l \in \{0, 1, ..., p-1\}$ , we have

$$\binom{Np+2l}{Kp+l} \equiv \binom{N}{K} \binom{2l}{l} + N \binom{N}{K} \left( \binom{p+2l}{l} - \binom{2l}{l} \right) \bmod p^2$$

(by Lemma 2.22). Summing these congruences over all  $l \in \{0, 1, ..., p-1\}$ , we find

$$\sum_{l=0}^{p-1} {Np+2l \choose Kp+l} \equiv \sum_{l=0}^{p-1} {\binom{N}{K}} {2l \choose l} + N {\binom{N}{K}} {\binom{p+2l}{l} - \binom{2l}{l}}$$

$$= {\binom{N}{K}} \sum_{l=0}^{p-1} {2l \choose l} + N {\binom{N}{K}} \sum_{l=0}^{p-1} {\binom{p+2l}{l} - \binom{2l}{l}}$$

$$\equiv {\binom{N}{K}} \sum_{l=0}^{p-1} {2l \choose l} + N {\binom{N}{K}} \sum_{l=0}^{p-1} {\binom{p+2l}{l} - \binom{2l}{l}}$$

$$\equiv {\binom{N}{K}} \sum_{l=0}^{p-1} {2l \choose l} \equiv {\binom{N}{K}} \eta_p \mod p^2.$$

$$= \sum_{n=0}^{p-1} {2n \choose n} \equiv \eta_p \mod p^2$$
(by Theorem 1.8)

This proves Lemma 2.23.

Proof of Theorem 1.9. The map

$$\{0,1,\ldots,p-1\} \times \{0,1,\ldots,r-1\} \to \{0,1,\ldots,rp-1\},\ (l,K) \mapsto Kp+l$$

is a bijection (since each element of  $\{0, 1, ..., rp-1\}$  can be uniquely divided by p with remainder, and said remainder will belong to  $\{0, 1, ..., r-1\}$ ). Thus, we can substitute Kp + l for n in the sum  $\sum_{n=0}^{rp-1} {2n \choose n}$ . This sum thus rewrites as follows:

$$\sum_{n=0}^{rp-1} {2n \choose n} = \underbrace{\sum_{\substack{(l,K) \in \{0,1,\dots,p-1\} \times \{0,1,\dots,r-1\} \\ = \sum\limits_{K=0}^{r-1} \sum\limits_{l=0}^{p-1}}} \underbrace{\begin{pmatrix} 2 (Kp+l) \\ Kp+l \end{pmatrix}} = \underbrace{\sum_{k=0}^{r-1} \sum\limits_{l=0}^{p-1} \binom{2Kp+2l}{Kp+l}} = \underbrace{\begin{pmatrix} 2Kp+2l \\ Kp+l \end{pmatrix}} = \underbrace{\begin{pmatrix} 2K \\ K \end{pmatrix} \eta_p \operatorname{mod} p^2}_{\text{(by Lemma 2.23, applied to } N=2K)}$$

$$\equiv \underbrace{\sum_{k=0}^{r-1} \binom{2K}{K}}_{n=0} \eta_p = \alpha_r \eta_p = \eta_p \alpha_r \operatorname{mod} p^2.$$

$$= \underbrace{\sum_{k=0}^{r-1} \binom{2R}{K}}_{n=0} \eta_p = \alpha_r \eta_p = \eta_p \alpha_r \operatorname{mod} p^2.$$

This proves Theorem 1.9.

## 2.7. Proving Theorem 1.10

**Lemma 2.24.** Let p be an odd prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{N}$ . Then,

$$\sum_{l=0}^{p-1} \sum_{m=0}^{l} \left( \binom{Np+l}{Kp+m} - \binom{N}{K} \binom{l}{m} \right) \binom{l}{m} \equiv 0 \bmod p^2.$$

Proof of Lemma 2.24. We have

This proves Lemma 2.24.

**Lemma 2.25.** Let p be an odd prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{N}$ . Then,

$$\sum_{l=0}^{p-1} \sum_{m=0}^{l} {Np+l \choose Kp+m}^2 \equiv {N \choose K}^2 \eta_p \operatorname{mod} p^2.$$

*Proof of Lemma* 2.25. Fix  $l \in \{0, 1, ..., p-1\}$  and  $m \in \{0, 1, ..., p-1\}$ . Then, Theorem 1.5 (applied to a = N, b = K, c = l and d = m) yields that  $\binom{Np+l}{Kp+m} \equiv \binom{N}{K} \binom{l}{m} \mod p$ . In other words,  $\binom{Np+l}{Kp+m} - \binom{N}{K} \binom{l}{m}$  is divisible by p. Hence,

Lemma 2.18 (applied to 
$$a = \binom{Np+l}{Kp+m}$$
 and  $b = \binom{N}{K}\binom{l}{m}$ ) shows that

$${\binom{Np+l}{Kp+m}}^2 - {\binom{N}{K}} {\binom{l}{m}}^2$$

$$\equiv 2 {\binom{Np+l}{Kp+m}} - {\binom{N}{K}} {\binom{l}{m}} {\binom{N}{K}} {\binom{l}{m}} \bmod p^2.$$
(5)

Now, forget that we fixed l and m. We thus have proven (5) for all  $l \in \{0, 1, ..., p-1\}$  and  $m \in \{0, 1, ..., p-1\}$ . Now,

$$\sum_{l=0}^{p-1} \sum_{m=0}^{l} {Np+l \choose Kp+m}^2 - \sum_{l=0}^{p-1} \sum_{m=0}^{l} {\binom{N}{K} \binom{l}{m}}^2$$

$$= \sum_{l=0}^{p-1} \sum_{m=0}^{l} {\binom{Np+l}{Kp+m}}^2 - {\binom{N}{K} \binom{l}{m}}^2$$

$$\equiv 2 {\binom{Np+l}{Kp+m}} - {\binom{N}{K} \binom{l}{m}} {\binom{N}{K} \binom{l}{m}} \mod p^2$$

$$(by (5))$$

$$\equiv 2 {\binom{N}{K}} \sum_{l=0}^{p-1} \sum_{m=0}^{l} {\binom{Np+l}{Kp+m}} - {\binom{N}{K} \binom{l}{m}} \binom{l}{m} \equiv 0 \mod p^2.$$

$$\equiv 0 \mod p^2$$

$$(by Lemma 2.24)$$

Thus,

$$\sum_{l=0}^{p-1} \sum_{m=0}^{l} {Np+l \choose Kp+m}^{2} \equiv \sum_{l=0}^{p-1} \sum_{m=0}^{l} {\binom{N}{k}} {l \choose m}^{2} = {\binom{N}{k}}^{2} \sum_{l=0}^{p-1} \sum_{m=0}^{l} {\binom{l}{m}}^{2}$$

$$= \sum_{m=0}^{l} {\binom{l}{m}} {\binom{l}{m}} = {\binom{2l}{l}}$$
(by Corollary 2.11, applied to  $u=l$  and  $w=0$ )
$$= {\binom{N}{k}}^{2} \sum_{l=0}^{p-1} {\binom{2l}{l}} \equiv {\binom{N}{k}}^{2} \eta_{p} \bmod p^{2}.$$
(by Theorem 1.8)

This proves Lemma 2.25.

**Lemma 2.26.** Let p be a prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{Z}$ . Let u and v be two elements of  $\{0,1,\ldots,p-1\}$  satisfying  $u+v \geq p$ . Then,  $p \mid \binom{Np+u+v}{Kp+u}$ .

*Proof of Lemma 2.26.* We have  $u + v \ge p$ . Thus, u + v = p + c for some  $c \in \mathbb{N}$ . Consider this c. From  $v \in \{0,1,\ldots,p-1\}$ , we obtain v < p. Thus,  $c + p = p + c = u + \underbrace{v}_{< p} < u + p$ , so that  $c < u \le p - 1$  (since  $u \in \{0,1,\ldots,p-1\}$ ). Thus,

 $c \in \{0, 1, \dots, p-1\}$  (since  $c \in \mathbb{N}$ ). Also, c < u. Hence, Proposition 2.2 (applied to m = c and n = u) yields  $\binom{c}{u} = 0$ .

Now, u + v = p + c, so that Np + u + v = Np + p + c = (N + 1)p + c. Hence,

$$\binom{Np+u+v}{Kp+u} = \binom{(N+1)p+c}{Kp+u} \equiv \binom{N+1}{K} \underbrace{\binom{c}{u}}_{=0}$$

(by Theorem 1.5, applied to a = N + 1, b = K and d = u) =  $0 \mod p$ .

In other words,  $p \mid \binom{Np+u+v}{Kp+u}$ . This proves Lemma 2.26.

**Lemma 2.27.** Let p be an odd prime. Let  $N \in \mathbb{Z}$  and  $K \in \mathbb{N}$ . Then,

$$\sum_{u=0}^{p-1} \sum_{v=0}^{p-1} {Np+u+v \choose Kp+u}^2 \equiv {N \choose K}^2 \eta_p \bmod p^2.$$

*Proof of Lemma* 2.27. If u and v are two elements of  $\{0,1,\ldots,p-1\}$  satisfying  $v \ge p-u$ , then

5

<sup>&</sup>lt;sup>5</sup>Proof of (6): Let u and v be two elements of  $\{0, 1, ..., p-1\}$  satisfying  $v \ge p-u$ . From  $v \ge p-u$ , we obtain  $u+v \ge p$ . Thus, Lemma 2.26 yields  $p \mid \binom{Np+u+v}{Kp+u}$ . Hence,  $p^2 \mid \binom{Np+u+v}{Kp+u}^2$ . This proves (6).

Hence, any  $u \in \{0, 1, ..., p - 1\}$  satisfies

$$\sum_{v=0}^{p-1} {\binom{Np+u+v}{Kp+u}}^2 = \sum_{v=0}^{p-u-1} {\binom{Np+u+v}{Kp+u}}^2 + \sum_{v=p-u}^{p-1} \underbrace{\binom{Np+u+v}{Kp+u}}^2_{=0 \bmod p^2 \text{(by (6))}}$$

(here, we have split the sum at v = p - u)

$$\equiv \sum_{v=0}^{p-u-1} {Np+u+v \choose Kp+u}^2 = \sum_{l=u}^{p-1} {Np+l \choose Kp+u}^2 \bmod p^2$$

(here, we have substituted l for u+v in the sum). Summing up these congruences for all  $u \in \{0, 1, ..., p-1\}$ , we obtain

$$\sum_{u=0}^{p-1} \sum_{v=0}^{p-1} {Np + u + v \choose Kp + u}^{2}$$

$$\equiv \underbrace{\sum_{u=0}^{p-1} \sum_{l=u}^{p-1}}_{l=u} {Np + l \choose Kp + u}^{2} = \sum_{l=0}^{p-1} \sum_{u=0}^{l} {Np + l \choose Kp + u}^{2} = \sum_{l=0}^{p-1} \sum_{m=0}^{l} {Np + l \choose Kp + u}^{2}$$

$$= \sum_{l=0}^{p-1} \sum_{v=0}^{l}$$

(here, we have renamed the index u as m in the second sum)

$$\equiv \binom{N}{K}^2 \eta_p \bmod p^2$$

(by Lemma 2.25). This proves Lemma 2.27.

*Proof of Theorem 1.10.* First, let us observe that

$$\epsilon_{r,s} = \sum_{m=0}^{r-1} \sum_{n=0}^{s-1} {n+m \choose m}^2 = \sum_{n=0}^{s-1} \sum_{m=0}^{r-1} {n+m \choose m}^2 = \sum_{K=0}^{s-1} \sum_{L=0}^{r-1} {K+L \choose L}^2$$

$$= \sum_{K=0}^{s-1} \sum_{L=0}^{r-1} {K+L \choose K}^2 \tag{7}$$

(since Proposition 2.3 yields  $\binom{K+L}{L} = \binom{K+L}{K}$  for all  $K \in \mathbb{N}$  and  $L \in \mathbb{N}$ ).

Each  $n \in \mathbb{N}$  satisfies

$$\sum_{m=0}^{sp-1} {n+m \choose m}^2 = \sum_{u=0}^{p-1} \sum_{K=0}^{s-1} {n+Kp+u \choose Kp+u}^2$$

(here, we have substituted Kp + u for m in the sum, since the map

$$\{0,1,\ldots,p-1\} \times \{0,1,\ldots,s-1\} \to \{0,1,\ldots,sp-1\},\ (u,K) \mapsto Kp+u$$

is a bijection). Summing up this equality over all  $n \in \{0, 1, ..., rp - 1\}$ , we obtain

$$\sum_{n=0}^{rp-1} \sum_{m=0}^{sp-1} {n+m \choose m}^2 = \sum_{n=0}^{rp-1} \sum_{u=0}^{p-1} \sum_{K=0}^{s-1} {n+Kp+u \choose Kp+u}^2$$
$$= \sum_{v=0}^{p-1} \sum_{L=0}^{r-1} \sum_{u=0}^{p-1} \sum_{K=0}^{s-1} {Lp+v+Kp+u \choose Kp+u}^2$$

(here, we have substituted Lp + v for n in the sum, since the map

$$\{0,1,\ldots,p-1\} \times \{0,1,\ldots,r-1\} \to \{0,1,\ldots,rp-1\},\ (v,L) \mapsto Lp+v$$

is a bijection).

Thus,

$$\sum_{n=0}^{rp-1} \sum_{m=0}^{sp-1} {n+m \choose m}^{2} = \sum_{v=0}^{p-1} \sum_{L=0}^{r-1} \sum_{u=0}^{p-1} \sum_{K=0}^{s-1} \underbrace{\left( Lp + v + Kp + u \right)^{2}}_{Kp + u}^{2}$$

$$= \sum_{K=0}^{s-1} \sum_{L=0}^{r-1} \sum_{u=0}^{p-1} \sum_{v=0}^{p-1} = \underbrace{\left( (K+L) p + u + v \right)^{2}}_{Kp + u}^{2}$$

$$= \sum_{K=0}^{s-1} \sum_{L=0}^{r-1} \sum_{u=0}^{p-1} \sum_{v=0}^{p-1} \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u + v \right)^{2}_{Kp + u}^{2}$$

$$= \left( (K+L) p + u +$$

This proves Theorem 1.10.

## 2.8. Acknowledgments

Thanks to Doron Zeilberger and Roberto Tauraso for alerting me to [AmdTau16] and [SunTau11].

#### References

[AmdTau16] Tewodros Amdeberhan, Roberto Tauraso, *Two triple binomial sum su-percongruences*, Journal of Number Theory **175** (2017), pp. 140–157. A preprint is arXiv:1607.02483v1.

- [AnBeRo05] Peter G. Anderson, Arthur T. Benjamin and Jeremy A. Rouse, *Combinatorial Proofs of Fermat's*, *Lucas's*, *and Wilson's Theorems*, The American Mathematical Monthly, Vol. **112**, No. 3 (Mar., 2005), pp. 266–268.
- [ApaZei16] Moa Apagodu, Doron Zeilberger, *Using the "Freshman's Dream" to Prove Combinatorial Congruences*, The American Mathematical Monthly, Vol. **124**, No. 7 (August-September 2017), pp. 597–608.

  (A preprint can be found at arXiv:1606.03351v2, but is less up-to-date and uses a different numbering of the conjectures.)
- [Bailey91] D. F. Bailey, Some Binomial Coefficient Congruences, Applied Mathematics Letters, Volume 4, Issue 4, 1991, pp. 1–5. https://doi.org/10.1016/0893-9659(91)90043-U
- [BenQui03] Arthur T. Benjamin and Jennifer J. Quinn, *Proofs that Really Count: The Art of Combinatorial Proof*, The Mathematical Association of America, 2003.
- [BenQui08] Arthur T. Benjamin and Jennifer J. Quinn, *An Alternate Approach to Alternating Sums: A Method to DIE for*, The College Mathematics Journal, Volume 39, Number 3, May 2008, pp. 191-202(12).
- [GrKnPa94] Ronald L. Graham, Donald E. Knuth, Oren Patashnik, *Concrete Mathematics*, *Second Edition*, Addison-Wesley 1994.
- [Grinbe17] Darij Grinberg, Notes on the combinatorial fundamentals of algebra, 4
  October 2018.

  https://github.com/darijgr/detnotes/releases/tag/2018-10-04
  See also http://www.cip.ifi.lmu.de/~grinberg/primes2015/sols.pdf
  for a version that is getting updates.
- [Grinbe17b] Darij Grinberg, *The Lucas and Babbage congruences*, 3 October 2018. http://www.cip.ifi.lmu.de/~grinberg/lucascong.pdf
- [Hausne83] Melvin Hausner, *Applications of a Simple Counting Technique*, The American Mathematical Monthly, Vol. 90, No. 2 (Feb., 1983), pp. 127–129.
- [Liu16] Ji-Cai Liu, On two conjectural supercongruences of Apagodu and Zeilberger, arXiv:1606.08432v3.
- [MacSon10] Kieren MacMillan and Jonathan Sondow, *Proofs of Power Sum and Binomial Coefficient Congruences Via Pascal's Identity*, arXiv:1011.0076v1. Published in: The American Mathematical Monthly, Vol. **118** (2011), pp. 549–551.
- [Mestro14] Romeo Meštrović, Lucas' theorem: its generalizations, extensions and applications (1878–2014), arXiv:1409.3820v1.

- [Stanle11] Richard Stanley, Enumerative Combinatorics, volume 1, Second edition, version of 15 July 2011. Available at http://math.mit.edu/~rstan/ec/
- [Stanto16] Dennis Stanton, Addendum to "Using the "Freshman's Dream" to Prove

  Combinatorial Congruences",

  http://www.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/freshmanDennisStant
- [SunTau11] Zhi-Wei Sun, Roberto Tauraso, *On some new congruences for binomial coefficients*, International Journal of Number Theory, Vol. 7, No. 3 (2011), pp. 645–662. A preprint is arXiv:0709.1665v10.