

HORIZONTAL VARIATION OF TATE–SHAFAREVICH GROUPS

ASHAY A. BURUNGALÉ, HARUZO HIDA, AND YE TIAN

ABSTRACT. Let E be an elliptic curve over \mathbb{Q} . Let p be an odd prime and $\iota : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ an embedding. Let K be an imaginary quadratic field and H_K the corresponding Hilbert class field. For a class group character $\chi \in \widehat{\text{Cl}}_K$, let $\mathbb{Q}(\chi)$ be the field generated by $\text{Im}(\chi)$ and \mathfrak{p}_χ the prime of $\mathbb{Q}(\chi)$ above p determined via ι_p . Under mild hypotheses, we show that the number of $\chi \in \widehat{\text{Cl}}_K$ such that the χ -isotypic Tate–Shafarevich group $\text{III}(E/H_K)^\chi$ is finite with trivial \mathfrak{p}_χ -part increases with the absolute value of the discriminant of K .

We also show an analogous result simultaneously for $\text{III}(E/H_K)^\chi$ and the analytic Sha of a GL_2 -type abelian variety over \mathbb{Q} . Here analytic Sha refers to the order of the conjecturally finite Tate–Shafarevich group as predicted by the Birch–Swinnerton-Dyer conjecture.

The approach is based on the arithmetic of automorphic forms on a quaternion algebra along with Euler system of toric periods and Heegner points. An underlying role is played by Michel’s Equidistribution of special points on a Shimura set arising from a definite quaternion algebra and Zariski density of special points on a Shimura curve arising from an indefinite quaternion algebra.

CONTENTS

1. Introduction	1
Acknowledgement	7
Notation	7
2. Toric Periods	7
2.1. Setup	7
2.2. Non-vanishing	9
2.3. Tate–Shafarevich Groups I	12
2.4. Analytic Sha I	15
3. Heegner Points	18
3.1. Setup	18
3.2. Shimura curve	20
3.3. Non-vanishing	21
3.4. p -indivisibility of Heegner points	23
3.5. Tate–Shafarevich groups II	25
3.6. Analytic Sha II	26
4. Main results	29
4.1. Torus embedding	29
4.2. Main results	30
References	32

1. INTRODUCTION

Tate–Shafarevich group is a mysterious arithmetic invariant associated to a motive \mathcal{M} over a number field F . It measures the failure of local to global principle for \mathcal{M} and is conjectured to be finite. In view of the Bloch–Kato conjecture, the conjectural finite size is expected to be closely related to a special value or a derivative of the L -function associated to \mathcal{M} . As the motive varies in a well-controlled family, variation of the Tate–Shafarevich group in the family seems to be a fundamental question.

An instructive setup arises from twists of \mathcal{M} by Hecke characters over K for a finite extension K/F . Families arise in numerous ways. For example, we may fix K and consider Hecke characters with fixed infinity type but allow the conductor to grow. This is usually referred as an Iwasawa or a vertical variation and it continues to be investigated in various situations. On the other hand, we may vary K with fixed degree and consider Hecke characters with fixed infinity type and bounded conductor. We

may refer this as a horizontal variation and it seems to be investigated in only few situations. In any of these families, we can ask for variation of the p -part of the Tate–Shafarevich group with p a fixed prime.

For a fixed GL_2 -type abelian variety over the rationals and a prime p , we hope to initiate the study of horizontal variation of the p -part of Tate–Shafarevich groups. In this article, we consider the GL_2 -case for a specific horizontal family. The family arises from the twists of the abelian variety by ideal class characters over imaginary quadratic fields.

Let us begin with a brief discussion of the GL_1 -case regarding ideal class groups. We first recall some history. Kummer defined p to be a regular prime if $p \nmid |\mathrm{Cl}(\mathbb{Q}(\zeta_p))|$ for the class group $\mathrm{Cl}(\mathbb{Q}(\zeta_p))$ of the p^{th} -cyclotomic field $\mathbb{Q}(\zeta_p)$. Kummer showed that p is an irregular prime if and only if p divides the numerator of the Bernoulli number B_k for some even integer k with $2 \leq k \leq p-3$. After several decades, Herbrand and Ribet showed a refined version incorporating $\mathrm{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{Q})$ -action on the class group $\mathrm{Cl}(\mathbb{Q}(\zeta_p))$. They showed that the ω_p^{1-k} -isotypic class group $\mathrm{Cl}(\mathbb{Q}(\zeta_p))^{\omega_p^{1-k}}$ has non-trivial p -part if and only if p divides the numerator of the Bernoulli number B_k . Here ω_p denotes the Teichmüller character. As $p \nmid |\mathrm{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{Q})|$, the action of $\mathrm{Gal}(\mathbb{Q}(\zeta_p)/\mathbb{Q})$ on $\mathrm{Cl}(\mathbb{Q}(\zeta_p))$ is semi-simple. In these results, the Bernoulli number B_k can be replaced with the special zeta-value $\zeta(1-k)$.

We introduce some notation to state the results. Let G be an abelian group and \widehat{G} the group of $\overline{\mathbb{Q}}^\times$ -valued characters of G . For $\chi \in \widehat{G}$, let $\mathbb{Z}[\chi]$ be the ring generated over \mathbb{Z} by the image of χ and $\mathbb{Q}(\chi)$ its fraction field. For any $\mathbb{Z}[G]$ -module M , let M^χ denote the χ -isotypic part of $M \otimes_{\mathbb{Z}} \mathbb{Z}[\chi]$. Fix embeddings $\iota_\infty : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$, $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$ and an isomorphism $\iota : \mathbb{C} \rightarrow \mathbb{C}_p$ such that $\iota_p = \iota \circ \iota_\infty$. Let v_p be the valuation on \mathbb{C}_p with $v_p(p) = 1$. For $\chi \in \widehat{G}$, let \mathfrak{p}_χ be the prime ideal of $\mathbb{Q}(\chi)$ determined by ι_p .

We now introduce an analogue of Kummer’s notion of regularity for Dirichlet characters. Let χ be a $\overline{\mathbb{Q}}^\times$ -valued Dirichlet character. Let \mathbb{Q}_χ be the field cut out by χ and $\mathrm{Cl}(\mathbb{Q}_\chi)$ its ideal class group with the action of $\mathrm{Gal}(\mathbb{Q}_\chi/\mathbb{Q})$. Suppose that $p \nmid [\mathbb{Q}_\chi : \mathbb{Q}]$. Following Kummer, we say that

χ is a p -regular Dirichlet character if the χ -isotypic class group $\mathrm{Cl}(\mathbb{Q}_\chi)^\chi$ has trivial \mathfrak{p}_χ -part.

By Mazur–Wiles ([31]) and Wiles ([51]), it is equivalent to the Dirichlet L -value $L(0, \chi^{-1})$ being \mathfrak{p}_χ -indivisible or equivalently to the p -indivisibility of the Bernoulli number $B_{1, \chi^{-1}}$. Recently, Burungale–Sun obtained results towards the number of p -regular Dirichlet characters ([11]). We refer to [11, §1] for their main results. In the Iwasawa setup, the results go back to Washington ([49]) and Sinnott ([42]).

We now turn to GL_2 -type abelian varieties over the rationals. In the introduction, we restrict to elliptic curves for simplicity.

Let K be an imaginary quadratic field and D_K the discriminant. Let $K_\mathbb{A}$ be the ring of adeles. Let G_K denote the absolute Galois group over K . Let Cl_K be the ideal class group of K and h_K the ideal class number. Let H_K be the Hilbert class field of K .

Let E be an elliptic curve over \mathbb{Q} . There is a natural action of Cl_K on the Tate–Shafarevich group $\mathrm{III}(E/H_K)$ via Artin’s reciprocity isomorphism $\mathrm{Gal}(H_K/K) \simeq \mathrm{Cl}_K$. A basic question is the following.

(Q) For a prime p , how does the $\mathbb{Z}[\mathrm{Cl}_K]$ -module $\mathrm{III}(E/H_K)[p^\infty]$ vary as K -varies over imaginary quadratic fields?

A fundamental conjecture of Tate asserts that $\mathrm{III}(E/L)$ is finite for a number field L . We are not aware of any conjecture regarding the distribution of the above horizontal variation on III . The horizontal variation seems to be complementary to the variation considered in Delaunay ([16]) and Bhargava–Kane–Lenstra–Poonen–Rains ([5]). The question is closely related to the variation of the $\mathbb{Z}[\chi]$ -module $\mathrm{III}(E/H_K)^\chi[p^\infty]$ as the pair (K, χ) -varies with $\chi \in \widehat{\mathrm{Cl}}_K$.

Recall that $\mathfrak{p}_\chi \subset \mathbb{Z}[\chi]$ denotes the prime above p of $\mathbb{Q}(\chi)$ determined via the embedding ι_p . We consider the horizontal variation of the \mathfrak{p}_χ -primary part of $\mathrm{III}(E/H_K)^\chi$ in this article. This seems to be a slightly approachable question than the original (Q).

One of our main results is the following.

Theorem 1.1. *Let E be a non-CM elliptic curve over \mathbb{Q} . Then, there exists an explicit finite subset Σ_E of primes only dependent on E , such that for any fixed prime $p \notin \Sigma_E$ and $\epsilon > 0$ we have*

$$\#\left\{ \chi \in \widehat{\mathrm{Cl}}_K \mid \mathrm{III}(E/H_K)^\chi \text{ is finite with trivial } \mathfrak{p}_\chi\text{-part} \right\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic fields such that

- (i) $p \nmid h_K$, and
- (ii) either p splits in K or the root number $\epsilon(E, \chi_0)$ of the Rankin–Selberg convolution $L(s, E, \chi_0)$ equals $+1$ for some $\chi_0 \in \widehat{\mathrm{Cl}}_K$.

Remark 1.2. (1). Under mild hypothesis, a precise description of the exceptional set Σ_E is given in Theorem 1.5.

(2). Cohen-Lenstra heuristics predicts that the imaginary quadratic fields satisfying the hypotheses in the theorem have positive density. It is known that there exist infinitely many imaginary quadratic fields K satisfying the hypotheses ([7] and [51]). For a recent quantitative refinement of these results, we refer to [1].

As for the horizontal variation, we may ask the same question for analytic Sha. Recall that the BSD conjecture predicts the size for the conjecturally finite Tate–Shafarevich groups in terms of the behaviour of L-function around the center and arithmetic invariants of the underlying abelian variety. We refer to this quantity as analytic Sha. In this setup, relevant abelian variety turns out to be the following. For $\chi \in \widehat{\text{Cl}}_K$, let $E_\chi = E_K \otimes_{\mathbb{Z}} \mathbb{Z}[\chi]$ denote the Serre tensor where the absolute Galois group G_K acts on $\mathbb{Z}[\chi]$ via χ ([30]). Then E_χ is an abelian variety over K with the following properties:

$$\text{III}(E_\chi/K) \otimes_{\mathbb{Z}} \mathbb{Z}[h_K^{-1}] \cong \text{III}(E/H_K)^\chi \otimes_{\mathbb{Z}} \mathbb{Z}[h_K^{-1}], \quad L(s, E_\chi) = L(s, E, \chi) \in \mathbb{C} \otimes_{\mathbb{Q}} \mathbb{Q}(\chi).$$

Let $\mathcal{L}(E_\chi)$ be the analytic Sha of E_χ (part (2) of Conjecture 2.11), which is conjectured to be a non-zero integral ideal of $\mathbb{Q}(\chi)$ as predicted by the BSD conjecture for the abelian variety E_χ over K .

Definition 1.3. A character $\chi \in \widehat{\text{Cl}}_K$ is said to be p -regular for E if the following conditions hold.

- $\text{rank}_{\mathbb{Z}[\chi]} E_\chi(K) = \text{ord}_{s=1} L(s, E_\chi/K)$, and
- Let $\mathfrak{p}_\chi | p$ be the prime ideal of $\mathbb{Z}[\chi]$ induced via the fixed embedding ι_p , then
 - (a) $\text{III}(E_\chi)$ is finite with trivial \mathfrak{p}_χ -part, and
 - (b) $\mathcal{L}(E_\chi)$ has trivial \mathfrak{p}_χ -part.

In particular, for a p -minimal $\chi \in \widehat{\text{Cl}}_K$, the \mathfrak{p} -part of the full BSD conjecture holds for E_χ .

Our result regarding p -minimality is the following.

Theorem 1.4. Let E be a non-CM elliptic curve over \mathbb{Q} . Then, there exists an explicit finite subset Σ_E of primes only dependent on E , such that for any prime $p \notin \Sigma_E$ and $\epsilon > 0$ we have

$$\#\left\{ \chi \in \widehat{\text{Cl}}_K \mid \chi \text{ is } p\text{-regular for } E \right\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic fields satisfying

- (i) $p \nmid h_K$, and
- (ii) either p splits in K or the root number $\epsilon(E, \chi_0)$ of the Rankin–Selberg convolution $L(s, E, \chi_0)$ equals $+1$ for some $\chi_0 \in \widehat{\text{Cl}}_K$.

Note that the above theorem is stronger than Theorem 1.1 in appearance.

The set Σ_E is intrinsically related to the arithmetic of E . More precisely, it encodes the Tamagawa numbers of E , Galois image of the p -adic Galois representation $\rho_{E,p}$ and the local Galois representations arising from $\rho_{E,p}$ at primes dividing the conductor N of E . For $\ell | N$, let c_ℓ be the Tamagawa number of E at ℓ . Let $d_\ell \geq 1$ be the greatest common divisor of $[M_\ell : \mathbb{Q}_\ell]$ for all extensions M_ℓ/\mathbb{Q}_ℓ such that the base change of E to M_ℓ has either good or multiplicative reduction. For example, if $\ell | N$, then $d_\ell = 1$. An explicit version of Theorem 1.4 is the following.

Theorem 1.5. Let E/\mathbb{Q} be a non-CM elliptic curve over \mathbb{Q} with conductor N . Let $p \nmid 6N \prod_{\ell | N} (\ell^2 - 1) c_\ell d_\ell$ be a prime such that the $\rho_{E,p}$ is surjective. Then for any $\epsilon > 0$,

$$\#\left\{ \chi \in \widehat{\text{Cl}}_K \mid \chi \text{ is } p\text{-regular for } E \right\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic extensions satisfying

- (i) $p \nmid h_K$,
- (ii) $(N, D_K) = 1$,
- (iii) E is semi-stable at any prime factor of N inert in K ;
- (iv) either p splits in K or the number of prime factors of N inert in K is odd.

In the article, we also prove analogous results for GL_2 -type abelian varieties over the rationals (§4). By Khare–Wintenberger ([27]), such abelian varieties are modular. In turn, modular forms are integral to our approach.

We now describe the approach. It is based on the arithmetic of automorphic forms on a quaternion algebra along with Euler system of toric periods and Heegner points. The Euler system controls both

the Tate–Shafarevich group and analytic Sha. Our point is to first establish p -indivisibility of the Euler system. A finer analysis of the relation of the Euler system with Tate–Shafarevich group and analytic Sha then leads to the main result. An underlying role is played by the following.

- Michel’s equidistribution of special points arising from varying imaginary quadratic fields on a Shimura set arising from a definite quaternion algebra ([32]).
- Zariski density of CM points arising from varying imaginary quadratic fields on a Shimura curve arising from an indefinite quaternion algebra.

As $\chi \in \widehat{\text{Cl}}_K$, the Rankin–Selberg convolution $L(s, E, \chi)$ corresponding to the pair (E, χ) is self-dual. Let $\epsilon(E, \chi)$ denote the global root number. Let

$$\mathfrak{X}_K^\pm = \left\{ \chi \in \widehat{\text{Cl}}_K \mid \epsilon(E, \chi) = \pm 1 \right\}.$$

The approach crucially relies on the root number. Let S be the set of places of \mathbb{Q} dividing $N\infty$. For each character $\chi = \otimes \chi_v$ of Cl_K viewed as a Hecke character of $K_\mathbb{A}^\times$ via class field theory, let $\chi_S = \otimes_{v \in S} \chi_v : \prod_{v \in S} K_v^\times \rightarrow \overline{\mathbb{Q}}^\times$ be its S -component. Some of the notation used here and in the sketch below is not followed in the rest of the article.

The set \mathfrak{X}_K^\pm admits a finite partition according to the S -local component of characters in \mathfrak{X}_K^\pm . For a fixed type $\chi_{0,S}$, there is a definite quaternion algebra B/\mathbb{Q} and a so-called $\chi_{0,S}$ -toric line V in π_B , such that $L(1, E, \chi) \neq 0$ if and only if the period integral

$$P_f(\chi) := \int_{K_\mathbb{A}^\times / K^\times \mathbb{A}^\times} f(t) \chi(t) dt$$

is non-vanishing for any non-zero vector $f \in V$. This is an inexplicit version of Waldspurger formula ([48] and [52]). Here π_B denotes the Jacquet–Langlands transfer of π to $B_\mathbb{A}^\times$ for $B_\mathbb{A} = B \otimes \mathbb{A}$ with \mathbb{A} the adèles over \mathbb{Q} . Moreover, we fix an embedding of K into B dependent on the S -component.

The space V contains non-zero vectors and we fix such a form f from now. Let U be an open compact subgroup of $B_\mathbb{A}^{(\infty), \times}$ such that f is U -invariant. Here $B_\mathbb{A}^{(\infty)}$ denotes the finite part. Let X_U be the corresponding definite Shimura set with level U . We can choose U such that for any pair (K, χ) with the fixed type $\chi_{0,S}$, the chosen embedding $K_\mathbb{A}$ into $B_\mathbb{A}$ induces a map $\varphi_K : \text{Cl}_K \rightarrow X_U$. The image of the map is referred as the special points arising from the class group Cl_K . Let $\mathfrak{p}_0|p$ be the prime of $\mathbb{Z}[\chi_{0,S}]$ induced via the embedding ι_p . Without loss of generality, we may suppose that the image of $f : X_U \rightarrow \mathbb{Z}[\chi_{0,S}]$ generates an ideal prime to \mathfrak{p}_0 .

A key point is to study the non-vanishing of toric periods $P_f(\chi)$ modulo \mathfrak{p}_χ as the pair (K, χ) -varies with S -component $\chi_{0,S}$ (thus $\mathfrak{p}_\chi | \mathfrak{p}_0$). The \mathfrak{p}_χ -indivisibility of $P_f(\chi)$ begins with Fourier analysis on Cl_K and its relation with the induced map $\varphi_K : \text{Cl}_K \rightarrow X_U$. Recall that X_U is a finite set whereas the class number grows with the discriminant of K . Based on Shimura’s reciprocity law, we reduce the \mathfrak{p}_χ -indivisibility to equidistribution of special points on the Shimura set X_U . The equidistribution concerns the distribution of the special points

$$\left\{ \varphi_K(H_K) \mid H_K \subset \text{Cl}_K \right\} \subset X_U$$

with $H_K \subset \text{Cl}_K$ a subgroup such that $[\text{Cl}_K : H_K] \ll |D_K|^\delta$ for some $\delta > 0$. Such an equidistribution is essentially due to Michel ([32]). It is based on the subconvex bound due to Michel–Venkatesh [34] and the explicit Waldspurger formula in Cai–Shu–Tian ([13]). The hypothesis on the p -indivisibility of the class number of the imaginary quadratic fields seems essential for the Fourier analysis.

In view of the the work on the Euler system of toric periods due to Bertolini–Darmon [2] as refined by Nekovář [37], the non-vanishing of the toric period $P_f(\chi)$ implies the finiteness of $\text{III}(E/H_K)^\chi$. Moreover, there exists a finite set of primes Σ_E such that the size of the \mathfrak{p}_χ -part $\text{III}(E/H_K)^\chi[\mathfrak{p}_\chi^\infty]$ is controlled by the \mathfrak{p}_χ -divisibility of the toric period $P_f(\chi)$ for $p \notin \Sigma_E$. More precisely, we have

$$P_f(\chi) \cdot \mathfrak{p}_\chi^{C_{\chi, \mathfrak{p}_\chi}} \cdot \text{III}(E/H_K)^\chi[\mathfrak{p}_\chi^\infty] = 0$$

for a non-negative integer $C_{\chi, \mathfrak{p}_\chi}$. Our point is to investigate horizontal variation of the constants $C_{\chi, \mathfrak{p}_\chi}$. We show the existence of a finite set of primes $\Sigma_{E,1}$ such that the constant $C_{\chi, \mathfrak{p}_\chi}$ vanishes as (K, χ) -varies for $p \notin \Sigma_{E,1}$. The hypothesis on the p -indivisibility of the class number of the imaginary quadratic fields seems essential for this control. Along with the \mathfrak{p}_χ -indivisibility of toric periods $P_f(\chi)$, this leads to the proof Theorem 1.1 in the setup.

The situation is analogous for analytic Sha. Based on the explicit Waldspurger formula ([13, §1.3]), we have $\mathcal{L}(E, \chi) \in \mathbb{Q}(\chi)^\times$ as long as the analytic rank equals zero. More precisely, we have

$$\mathcal{L}(E_\chi) = D_\chi \cdot P_f(\chi)^2$$

for a fractional ideal $D_\chi \subset \mathbb{Z}[\chi]$. Our point is to investigate horizontal variation of the ideals D_χ . We show the existence of a finite set of primes $\Sigma_{E,2}$ such that $v_p(D_\chi) = 0$ for $p \notin \Sigma_{E,2}$. Along with the \mathfrak{p}_χ -indivisibility of toric periods and the horizontal variation of the constants $C_{\chi, \mathfrak{p}_\chi}$, this leads to the proof of Theorem 1.4 in the setup.

Under the hypotheses of Theorem 1.5, a finer analysis of the above approach especially that of the constants $C_{\chi, \mathfrak{p}_\chi}$ and the ideals D_χ , leads to the proof of Theorem 1.5 in the setup.

The lower bound in Theorem 1.1 and Theorem 1.5 is based on Shimura's reciprocity law and an elementary group theoretic lemma. Roughly speaking, the reciprocity law implies that the set S_K of p -regular characters is invariant under a Galois action. The Fourier analysis implies that the subgroup generated by S_K has size bounded below by $|\mathrm{Cl}_K|^{\delta'}$ for some $\delta' > 0$. An elementary argument then shows that the size of S_K is bounded below by $(\log(|\mathrm{Cl}_K|))^{1-\epsilon}$ for $\epsilon > 0$.

The p -adic Waldspurger formula due to Bertolini–Darmon–Prasanna ([3]), Brooks ([6]) and Liu–Zhang–Zhang ([29]) allows an analogous approach in the rank one case.

The set \mathfrak{X}_K^- admits a finite partition according to the S -local component of characters in \mathfrak{X}_K^- . For a fixed type $\chi_{0,S}$, there is an indefinite quaternion algebra B/\mathbb{Q} and a so-called $\chi_{0,S}$ -toric line V in π_B , such that $L'(1, E, \chi) \neq 0$ if and only if the Heegner point

$$P_f(\chi) := \int_{K_\mathbb{A}^\times / K^\times \mathbb{A}^\times} f(t) \chi(t) dt$$

is non-torsion for any non-zero vector $f \in V$. This is an inexplicit version of Gross–Zagier formula ([19] and [52]). Here π_B denotes the Jacquet–Langlands transfer of π to $B_\mathbb{A}^\times$. Moreover, we fix an embedding of K into B dependent on the S -component. In the above expression for the Heegner point, f is viewed as a modular parametrisation $f : X_U \rightarrow E_{\chi_{0,S}}$ for Shimura curve X_U arising from B with level U specified below and the Serre tensor $E_{\chi_{0,S}}$.

The space V contains non-zero vectors and we fix such a form f from now. Let U be an open compact subgroup of $B_\mathbb{A}^{(\infty), \times}$ such that f is U -invariant. Let X_U be the corresponding Shimura curve with level U . We can choose U such that for any pair (K, χ) with the fixed type $\chi_{0,S}$, the chosen embedding $K_\mathbb{A}$ into $B_\mathbb{A}$ induces a map $\varphi_K : \mathrm{Cl}_K \rightarrow X_U$. The image of the map is referred as the CM points arising from the class group Cl_K . Without loss of generality, we may suppose that f is a non-zero p -primitive form.

A key point is to study the non-vanishing of toric periods $P_\chi(g)$ modulo \mathfrak{p}_χ as the pair (K, χ) -varies with S -component $\chi_{0,S}$. Here g is a weight zero p -adic modular form given by $d^{-1}(f^{(p)})$ for d the Katz p -adic differential operator and $f^{(p)}$ the p -depletion of f . In view of the p -adic Waldspurger formula, the p -adic logarithm of the Heegner point $P_f(\chi)$ is a p -unit if and only if a normalisation of the toric period $P_g(\chi)$ is a p -unit. In this sense, the \mathfrak{p}_χ -indivisibility of the Heegner point $P_f(\chi)$ is closely related to the p -indivisibility of the toric period $P_g(\chi)$.

Strictly speaking, $P_g(\chi)$ is a toric period on an Igusa tower over the Shimura curve. Let $\pi : Ig \rightarrow X_U$ be the p -ordinary Igusa tower. Recall that Ig is a formal scheme defined over W the Witt ring of the algebraic closure \mathbb{F} of \mathbb{F}_p arising from the embedding ι_p . We also recall that g is a function on the Igusa tower Ig . The p -indivisibility of $P_g(\chi)$ begins with Fourier analysis on Cl_K and its relation with the induced map $\varphi'_K : \mathrm{Cl}_K \rightarrow Ig$. Here φ'_K is a lift of φ_K . Based on Shimura's reciprocity law, we reduce the p -indivisibility to Zariski density of the mod p reduction of special points on the special fiber $Ig_{\mathbb{F}}$. The Zariski density concerns the density of the special points

$$\left\{ \varphi'_K(H_K) \mid H_K \subset \mathrm{Cl}_K \right\} \subset Ig_{\mathbb{F}}$$

with $H_K \subset \mathrm{Cl}_K$ a subgroup of bounded index as $|D_K| \rightarrow \infty$. The Zariski density is a consequence of CM theory and the geometry of Igusa tower modulo p . The hypothesis on the p -indivisibility of the class number of the imaginary quadratic fields seems essential for the Fourier analysis.

In view of the work on the Euler system of Heegner points due to Kolyvagin [28] as refined by Nekovář [36], the non-triviality of the Heegner point $P_\chi(f)$ implies the finiteness of $\mathrm{III}(E/H_K)^\chi$. Moreover, there exists a finite set of primes Σ'_E such that the size of the \mathfrak{p}_χ -part $\mathrm{III}(E/H_K)^\chi[\mathfrak{p}_\chi^\infty]$ is controlled by the \mathfrak{p}_χ -divisibility of the Heegner point $P_f(\chi)$ for $p \notin \Sigma'_E$. More precisely, we have

$$[E(H_K)^\chi : \mathbb{Z}[\chi] \cdot P_f(\chi) + E(H_K)_{\mathrm{tor}}^\chi] \cdot \mathfrak{p}_\chi^{C'_{\chi, \mathfrak{p}_\chi}} \cdot \mathrm{III}(E/H_K)^\chi[\mathfrak{p}_\chi^\infty] = 0$$

for a non-negative integer $C'_{\chi, \mathfrak{p}_\chi}$. Our point is to investigate horizontal variation of the constants $C'_{\chi, \mathfrak{p}_\chi}$. We show that there exists a finite set of primes $\Sigma'_{E,1}$ such that the constant $C'_{\chi, \mathfrak{p}_\chi}$ vanishes as (K, χ) -varies for $p \notin \Sigma'_{E,1}$. The hypothesis on the p -indivisibility of the class number of the imaginary quadratic fields seems essential for this control. Along with Galois image result for E and the \mathfrak{p}_χ -indivisibility of the Heegner points $P_f(\chi)$, this leads to the proof Theorem 1.1 in the setup.

The situation is analogous for analytic Sha. Based on an explicit Gross–Zagier formula in Cai–Shu–Tian ([13, §1.3]), we have $\mathcal{L}(E_\chi) \in \mathbb{Q}(\chi)^\times$ as long as the analytic rank equals one. More precisely, we have

$$\mathcal{L}(E_\chi) = D'_\chi \cdot [E(H_K)^\chi : \mathbb{Z}[\chi] \cdot P_f(\chi) + E(H_K)^\chi_{\text{tor}}]^2$$

for a fractional ideal $D'_\chi \subset \mathbb{Z}[\chi]$. Our point is to investigate horizontal variation of the ideals D'_χ . We show that there exists a finite set of primes $\Sigma'_{E,2}$ such that $v_p(D'_\chi) = 0$ for $p \notin \Sigma'_{E,2}$. Along with Galois image result for E , the \mathfrak{p}_χ -indivisibility of Heegner points and the horizontal variation of the constants $C'_{\chi, \mathfrak{p}_\chi}$, this leads to the proof of Theorem 1.4 in the setup.

Under the hypotheses of Theorem 1.5, a finer analysis of the above approach especially that of the constants $C'_{\chi, \mathfrak{p}_\chi}$ and the ideals D'_χ , leads to the proof of Theorem 1.5 in the setup.

As before, the lower bound in Theorem 1.1 and Theorem 1.5 is based on Shimura’s reciprocity law and the elementary group theoretic lemma.

As evident from the sketch, there is a visible analogy among the case of root number 1 and the case of root number -1 . The analogy seem to resonate resemblance among the Waldspurger formula and the p -adic Waldspurger formula. On the other hand, there are striking differences. Here we only mention that the equidistribution of special points relies on a subconvex bound for Rankin–Selberg L-functions whereas the Zariski density of CM points relies on the geometry of Igusa tower modulo p .

Based on Jochnowitz congruence ([46] and [2]), the case of root number -1 can be deduced from the case of root number 1 for variation over imaginary quadratic fields satisfying extra hypothesis that there exists a fixed prime inert in them.

Our main results can be considered as a mod p analogue of the characteristic zero non-vanishing results for Rankin–Selberg L-functions due to Michel–Venkatesh ([33] and [34]) and Templier ([43]). As far as we know, these are the first results towards a mod p analogue. In the case of root number 1 (resp. -1), these article obtain quantitative results towards the number of class group characters with the corresponding Rankin–Selberg central L-value (resp. central derivative of the L-function) non-vanishing. The approach in these articles seems to have more analytic/ ergodic flavour. In Burungale–Tian ([12]), non-quantitative results are obtained in the case of root number -1 based on an algebro-geometric approach which relies on André–Oort conjecture. It seems instructive to compare our current approach with the ones in [33], [34], [43], [10] and [12]. Here we only mention that Michel’s equidistribution also plays an underlying role for the characteristic zero non-vanishing result due to Michel–Venkatesh ([34]). It seems striking that the characteristic zero and mod p non-vanishing share a key ingredient in the proof.

In the Iwasawa setup, the non-vanishing goes back to Cornut ([15]) and Vatsal ([46], [47]). From the consideration of approach to non-vanishing, there seem to be fundamental differences in Cornut–Vatsal and our horizontal setup. To begin with, Iwasawa algebra plays a pivotal role in Cornut–Vatsal to reduce the generic non-vanishing to equidistribution of a class of special points on a definite Shimura set. The equidistribution is proven based on Ratner’s theory. The equidistribution also crucially relies on Iwasawa algebra as it leads to an action of a p -adic Lie group on the class of special points. In the horizontal setup, there does not seem to be a naive analogue of the Iwasawa algebra as the number fields in consideration vary horizontally. Moreover, the special points do not seem to admit a group action.

The article is organised as follows. In §2, we consider the root number $+1$ case and analyse certain aspects of toric periods. In §2.1, we introduce the setup and fix a definite quaternion algebra B . In §2.2, we consider the horizontal mod p non-vanishing of toric periods of a toric form on B . In §2.3, we consider the horizontal variation of the constants appearing in the Euler system method to bound the Tate–Shafarevich group. In §2.4, we consider the horizontal variation of the constants appearing in the explicit Waldspurger formula alluded to above. In §3, we consider the root number -1 case and analyse certain aspects of Heegner points. In §3.1, we introduce the setup and fix an indefinite quaternion algebra B . In §3.2, we recall generalities regarding CM points on Shimura curves arising from B . In §3.3, we consider the horizontal mod p non-vanishing of toric periods of a weight zero p -adic modular form g alluded to above. In §3.4, we recall the p -adic Waldspurger formula. In §3.5, we consider the horizontal variation of the constants appearing in the Euler system method to bound the Tate–Shafarevich group. In §3.6, we consider the horizontal variation of the constants appearing in the explicit Gross–Zagier

formula alluded to above. In §4, we prove the main results based on §2 and §3 along with the existence of suitable torus embeddings into the underlying quaternion algebras.

ACKNOWLEDGEMENT

We thank Li Cai for his assistance and Hae-Sang Sun for stimulating conversations regarding horizontal non-vanishing. We also thank Henri Darmon, Najmuddin Fakhruddin, Dimitar Jetchev, Mahesh Kakde, Chandrashekhhar Khare, Chao Li, Philippe Michel, C.-S. Rajan, Nicolas Templier, Jacques Tilouine, Vinayak Vatsal, Xinyi Yuan, Shou-Wu Zhang and Wei Zhang for instructive conversations about the topic.

NOTATION

We use the following notation unless otherwise stated.

For a finite abelian group G , let \widehat{G} denote $\overline{\mathbb{Q}}^\times$ -valued character group of G . For a \mathbb{Z} -module A , let $\widehat{A} = A \otimes_{\mathbb{Z}} \widehat{\mathbb{Z}}$ for $\widehat{\mathbb{Z}} = \varprojlim_n \mathbb{Z}/n$.

For a number field L , let \mathcal{O}_L be the corresponding integer ring and D_L the discriminant. Let L_+ denote the totally positive elements in L . Let $G_L = \text{Gal}(\overline{\mathbb{Q}}/L)$ denote the absolute Galois group over L . Let \mathbb{A}_L denote the adeles over L . For a finite subset S of places in L , let $\mathbb{A}_L^{(S)}$ denote the adeles outside S and $\mathbb{A}_{L,S}$ the S -part. When L equals the rationals or the underlying totally real field, we drop the subscript L . For a \mathbb{Q} -algebra C , let $C_{\mathbb{A}} = C \otimes_{\mathbb{Q}} \mathbb{A}$. Let $\widehat{C}^{(S)}$ (resp. C_S) denote the part outside S (resp. S -part) of $C_{\mathbb{A}}$.

For a place v of F for a totally real field F and a quadratic extension K_v/F_v , let η_v denote the corresponding quadratic character. For a quaternion algebra B_v/F_v , let $\epsilon(B_v)$ denote the corresponding local invariant. For an CM quadratic extension K/F and an integral ideal \mathfrak{c} of F , let $H_{K,\mathfrak{c}}$ be the ring class field with conductor \mathfrak{c} and $\text{Pic}_{K/F}^{\mathfrak{c}}$ the relative ring class group with conductor \mathfrak{c} . Let h_K (resp. $h_{K,\mathfrak{c}}$) be the relative ideal class number of K (resp. $H_{K,\mathfrak{c}}$) over F .

2. TORIC PERIODS

In this section, we consider toric periods of a modular form on a definite quaternion algebra over the rationals. In §2.1, we introduce the setup. In §2.2., we prove horizontal mod p non-vanishing of toric periods. In §2.3 and §2.4, we deduce the consequences of the mod p non-vanishing for Tate–Shafarevich group and analytic Sha, respectively.

2.1. Setup. In this subsection, we introduce the setup.

Let $\phi \in S_2(\Gamma_0(N))$ be a newform with weight 2 and level $\Gamma_0(N)$ for $N \geq 3$. In particular, we consider newforms with trivial central characters. Let L_ϕ denote the Hecke field corresponding to the newform ϕ .

Let B be a definite quaternion algebra over \mathbb{Q} such that there exists an irreducible automorphic representation π_B on $B_{\mathbb{A}}^\times$ whose Jacquet–Langlands transfer is the automorphic representation of $\text{GL}_2(\mathbb{A})$ associated to ϕ . Recall $B_{\mathbb{A}} = B \otimes_{\mathbb{Q}} \mathbb{A}$.

Let $S = \text{Supp}(N\infty)$. Let $K_{0,S} \subset B_S$ be a \mathbb{Q}_S -subalgebra such that $K_{0,\infty} = \mathbb{C}$ and $K_{0,v}/\mathbb{Q}_v$ is semi-simple quadratic. As in the introduction, B_S (resp. \mathbb{Q}_S) denotes the S -part of $B_{\mathbb{A}}$ (resp. \mathbb{A}). For any $v \in S$, we say that v is non-split if $K_{0,v}$ is a field and split otherwise. Let

$$U_{0,S} := \prod_{v \in S, v \text{ split}} \mathcal{O}_{K_{0,v}}^\times \times \prod_{v \in S, v \text{ non-split}} K_{0,v}^\times.$$

Suppose we are given a finite order character $\chi_{0,S} : U_{0,S} \rightarrow \overline{\mathbb{Q}}^\times$ with conductor one such that the following holds.

$$(LC1) \quad \omega \cdot \chi_{0,S} \big|_{\mathbb{Q}_S^\times \cap U_{0,S}} = 1.$$

$$(LC2) \quad \epsilon(\phi, \chi_{0,v}) \chi_{0,v} \eta_v(-1) = \epsilon(B_v) \text{ for all places } v|N\infty \text{ with the local root number } \epsilon(\phi, \chi_{0,v}) \text{ corresponding to the Rankin–Selberg convolution.}$$

Fix a maximal order $R^{(S)}$ of $B_{\mathbb{A}}^{(S)} \cong M_2(\mathbb{A}^{(S)})$. Let $U^{(S)} = R^{(S)\times}$. Note that $U^{(S)}$ is a maximal compact subgroup of $B_{\mathbb{A}}^{(S)\times} \cong \text{GL}_2(\mathbb{A}^{(S)})$.

Let p be an odd prime. As in the introduction, we fix an embedding $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$. Let v_p be a p -adic valuation. Let \mathbb{F} be an algebraic closure of \mathbb{F}_p .

We introduce underlying imaginary quadratic fields.

Definition 2.1. Let Θ_S denote the set of imaginary quadratic fields K such that

- (i) $p \nmid h_K$,
- (ii) *there exists an embedding $\iota_K : K \rightarrow B$ with $K_S = K_{0,S}$,*
- (iii) $\mathbb{A}_K^{(S)} \cap R^{(S)} = \widehat{\mathcal{O}}_K^{(S)}$ *under the embedding and*
- (iv) $K \neq \mathbb{Q}(i), \mathbb{Q}(\omega)$ *for ω a primitive third root of unity.*

There exist infinitely many imaginary quadratic extensions K/\mathbb{Q} with $K \in \Theta_S$ ([7], [51] and [1]). For $K \in \Theta_S$, we fix such an embedding ι_K .

We introduce underlying Hecke characters over imaginary quadratic fields.

Definition 2.2. *For each $K \in \Theta$, let $\mathfrak{X}_{K,\chi_{0,S}}$ denote the set of finite order characters χ over K such that*

- (i) $\chi|_{\mathbb{A}^\times} \cdot \omega = 1$,
- (ii) $\chi_S = \chi_{0,S}$ *via the embedding ι_K , and*
- (iii) χ *is unramified outside S .*

Note that the conductor of $\chi \in \mathfrak{X}_{K,\chi_{0,S}}$ equals one. Moreover, we have

$$(\text{RN}) \quad \epsilon(\phi, \chi) = 1.$$

Here $\epsilon(\phi, \chi)$ denotes the global root number of the Rankin–Selberg convolution corresponding to the pair (ϕ, χ) .

In the rest of the section, we let $\Theta = \Theta_S$ and $\mathfrak{X}_K = \mathfrak{X}_{K,\chi_{0,S}}$.

Lemma 2.3. *The set \mathfrak{X}_K is non-empty for all but finitely many imaginary quadratic fields K with $K \in \Theta$. Moreover, it is a homogenous space for the class group Cl_K .*

Proof. Note that for all but finitely many imaginary quadratic fields K , we have that $\mathcal{O}_K^\times = \mathbb{Z}^\times$. We fix such an imaginary quadratic field from now.

In view of class field theory, the class group is given by $\text{Cl}_K = \mathbb{A}_K^\times / K^\times K_\infty^\times \mathbb{A}^\times U$. From the structure of U , there exists a non-trivial Hecke character ϵ of \mathbb{A}_K^\times with conductor one, trivial on K_∞^\times and \mathbb{A}^\times for all but finitely many imaginary quadratic fields K ([25]).

Let $\chi'_{0,S} = \chi_{0,S} \cdot \epsilon|_{U_{0,S}}$. Then, there exists a character χ' of $\mathbb{A}_K^\times / K^\times K_\infty^\times \mathbb{A}^\times U$ extending $\chi'_{0,S}$. The character $\chi' \cdot \epsilon^{-1}$ is a desired one. This finishes the proof of first part.

For $\chi \in \mathfrak{X}_K$, note that $\chi' \epsilon^{-1} \chi^{-1}$ factors through Cl_K . □

Let \mathcal{O} be the ring of integers of the field $\mathbb{Q}(\phi, \chi_{0,S})$ generated over \mathbb{Q} by the Hecke eigenvalues of the newform ϕ and the values of the local character $\chi_{0,S}$. Let \mathfrak{p} be the prime ideal of \mathcal{O} corresponding to ι_p . Let $\mathcal{O}_{(\mathfrak{p})}$ be the localisation of the integer ring \mathcal{O} at the prime ideal \mathfrak{p} .

We have the following existence of toric test vectors.

Lemma 2.4. *There exists a non-zero $\mathcal{O}_{(\mathfrak{p})}$ -valued form $f \in \pi_B$ such that the ideal generated by the image of f is prime to \mathfrak{p} and f satisfies the following.*

- (F1) *The subgroup $U_{0,S}$ acts on f via $\chi_{0,S}$ and*
- (F2) $f \in \pi^{U^{(S)}}$.

Proof. In view of (RN), there exists a non-zero $\ell \in \text{Hom}_{\mathbb{A}_K^\times}(\pi_B \otimes \chi, \mathbb{C})$ ([41], [44] and [45]). Here \mathbb{A}_K^\times acts trivially on \mathbb{C} .

From [13], we thus have $\varphi \in \pi_B^{U^{(S)}}$ such that $\ell(\varphi) \neq 0$. We further choose φ to be \mathfrak{p} -integral. For $t \in U_{0,S}$, note that

$$\ell(\chi_{0,S}(t)\varphi) = \chi_{0,S}(t)\varphi.$$

Recall that

$$\dim_{\mathbb{C}} \text{Hom}_{K_{\mathbb{A}}^\times}(\pi_B \otimes \chi, \mathbb{C}) = 1.$$

We can thus take f to be the automorphic form given by

$$g \mapsto \int_{U_{0,S}/\mathbb{Q}_S^\times \cap U_{0,S}} \chi_{0,S}(t)\varphi(gt)dt.$$

We further normalise f to be \mathfrak{p} -primitive by requiring that the ideal generated by the image of f is prime to \mathfrak{p} . □

Note that f is spherical outside S .

Let U be an open compact subgroup of $B_{\mathbb{A}}^{(\infty),\times}$ such that $U = U_S U^{(S)}$ with $U_S \subset B_S^\times$ and $U_S = \prod_{v \in S} U_v$ such that

- (L1) $f \in \pi_B^U$ and
(L2) $\mathcal{O}_{0,v}^\times \subset U_v$ for all $v \in S \setminus \{\infty\}$.

Let X_U be the corresponding Shimura set of level U given by

$$X_U = B^\times \backslash B_{\mathbb{A}}^{(\infty), \times} / U.$$

Let

$$\overline{f} : X_U \rightarrow \mathbb{F}$$

be the reduction of f modulo \mathfrak{p} .

Let \mathfrak{p}_0 be the prime above p in the Hecke field L_ϕ determined via the embedding ι_p . Let $\rho_{\phi, \mathfrak{p}_0} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(L_{\phi, \mathfrak{p}_0})$ be the corresponding p -adic Galois representation. Let $\overline{\rho}_{\phi, \mathfrak{p}_0}$ be the reduction modulo \mathfrak{p}_0 .

We have the following non-constancy regarding the reduction \overline{f} .

Lemma 2.5. *Assume that the mod p Galois representation $\overline{\rho}_{\phi, \mathfrak{p}_0}$ associated to the newform ϕ is absolutely irreducible. Let f be a toric form as above with non-zero reduction modulo \mathfrak{p} . Then, the mod \mathfrak{p} reduction \overline{f} is non-constant.*

Proof. Suppose it is not the case. From the definition of f , it is a Hecke eigenform outside the Hecke operators arising from primes in S . Moreover, the Hecke eigensystem is the same as that of ϕ . From the constancy, the eigensystem outside S is Eisenstein. This contradicts the irreducibility assumption. \square

We identify \mathbb{A}_K^\times as a subgroup of $B_{\mathbb{A}}^\times$ under the embedding $\iota_K : K \hookrightarrow B$. As $\iota_K(\mathcal{O}_K) \subset U$, the choice of embedding ι_K gives rise to a map

$$\varphi_K : \mathrm{Cl}_K \rightarrow X_U.$$

For $\sigma \in \mathrm{Cl}_K$, let $x_\sigma \in X_U$ be the corresponding point on the Shimura set. This is usually referred as a special point on the Shimura set. In what follows, the map φ_K plays an underlying role.

For $\chi \in \widehat{\mathrm{Cl}}_K$, let $P_f(\chi)$ be the toric period given by

$$(2.1) \quad P_f(\chi) = \frac{1}{h_K} \cdot \sum_{\sigma \in \mathrm{Cl}_K} \chi(\sigma)^{-1} f(x_\sigma).$$

As f is \mathfrak{p} -integral and $p \nmid h_K$, the period is \mathfrak{p} -integral. More precisely, $P_f(\chi) \in \mathcal{O}_{(\mathfrak{p})}[\chi]$. Here $\mathcal{O}_{(\mathfrak{p})}[\chi]$ denotes the extension of $\mathcal{O}_{(\mathfrak{p})}$ obtained by adjoining the values of χ .

The period possibly depends on the choice of the torus embedding ι_K .

Remark 2.6. (1). For $\chi \in \widehat{\mathrm{Cl}}_K$, the non-vanishing of $P_f(\chi)$ implies that $\chi \in \mathfrak{X}_K$. This follows from f being $\chi_{0,S}$ -toric. The observation will be used in the proof of Theorem 2.7.

(2). We restrict to newforms with trivial central character and unramified Hecke characters over the imaginary quadratic fields for simplicity.

2.2. Non-vanishing. In this subsection, we prove the horizontal mod p non-vanishing of toric periods of a modular form on a definite quaternion algebra.

Let the notation and assumptions be as in §2.1. In particular, ϕ is a weight two newform and L_ϕ the corresponding Hecke field. Let \mathfrak{p}_0 be the prime above p in L_ϕ determined via the embedding ι_p . Recall that $\rho_{\phi, \mathfrak{p}_0} : G_{\mathbb{Q}} \rightarrow \mathrm{GL}_2(L_{\phi, \mathfrak{p}_0})$ denotes the corresponding p -adic Galois representation. Moreover, $\overline{\rho}_{\phi, \mathfrak{p}_0}$ denotes the reduction modulo \mathfrak{p}_0 .

Our main result regarding the mod p non-vanishing of toric periods is the following.

Theorem 2.7. *Let $\phi \in S_2(\Gamma_0(N))$ be a newform. Let $p \nmid N$ be a prime such that*

(irr) *the mod p Galois representation $\overline{\rho}_{\phi, \mathfrak{p}_0}$ is absolutely irreducible.*

Let f be a toric test vector on a definite quaternion algebra as above. Let $\Theta = \Theta_S$ be the set of imaginary quadratic extensions as in Definition 2.1. For $K \in \Theta$, let \mathfrak{X}_K be the set of finite order Hecke character over K as in Definition 2.2. For $\chi \in \mathfrak{X}_K$, let $P_f(\chi)$ be the toric period corresponding to the pair (f, χ) .

For $\epsilon > 0$, we have

$$\#\left\{ \chi \in \mathfrak{X}_K \mid v_p(P_f(\chi)) = 0 \right\} \gg_\epsilon \log(|D_K|)^{1-\epsilon}.$$

as K varies in Θ .

Proof. Recall that $f : X_U \rightarrow \mathcal{O}_{(\mathfrak{p})}$ is a function on the Shimura set X and \bar{f} its mod \mathfrak{p} reduction. In view of the map $\varphi_K : \text{Cl}_K \rightarrow X_U$ of special points, we may regard f and \bar{f} as functions on the abelian group Cl_K .

The approach is based on the Fourier analysis on Cl_K and its relation to the Shimura set X_U .

To begin with, the toric periods $P_f(\chi)$ modulo \mathfrak{p}_χ equal the χ -th Fourier coefficients of the function

$$\sigma \mapsto f(x_\sigma) \bmod \mathfrak{p}$$

on the finite abelian group Cl_K . Here \mathfrak{p}_χ (resp. \mathfrak{p}) is the prime above p in $L_{\phi,\chi}$ (resp. L'_ϕ) determined via the embedding ι_p . Here L'_ϕ denotes the Hecke field corresponding to the pair $(\phi, \chi_{0,S})$ and $L_{\phi,\chi}$ its extension obtained by adjoining the values of χ .

In what follows, we thus consider Fourier analysis of the function

$$\bar{f} : \text{Cl}_K \rightarrow \mathbb{F}$$

and its variation for $K \in \Theta$ with \mathbb{F} an algebraic closure of \mathbb{F}_p . As $p \nmid h_K$, the Fourier inversion works as usual.

Let us first consider the non-vanishing of at least one twist. In view of the Fourier inversion, it suffices to show that

$$(2.2) \quad \bar{f}(x_\sigma) \neq 0$$

for at least one $\sigma \in \text{Cl}_K$ for all but finitely many K .

By a result of Iwaniec ([24]), the special points

$$\left\{ x_\sigma \mid \sigma \in \text{Cl}_K \right\} \subset X_U$$

become equidistributed on X_U with respect to the probability measure on the finite set X_U as $|D_K| \rightarrow \infty$. Strictly speaking, the equidistribution is a consequence of Iwaniec's main result in [24] as noticed in Michel ([32]).

In view of the p -primitivity of f (Lemma 2.4), the equidistribution readily implies (2.2).

We now consider the growth in the number of non-vanishing twists. Suppose that there exists a positive integer ℓ_K such that exactly $\ell_K \geq 1$ of the toric periods $P_f(\chi)$ are non-vanishing modulo \mathfrak{p}_χ , for K with sufficiently large discriminant. Let

$$\Xi_K = \{\chi_{K,1}, \dots, \chi_{K,\ell_K}\}$$

be the set of the non-vanishing modulo \mathfrak{p}_χ twists.

Note that

$$\overline{P_f(\chi)} \neq 0 \iff \overline{P_f(\chi^\sigma)} \neq 0$$

for $\sigma \in \text{Gal}(\mathbb{F}/\mathbb{F}_p(\overline{\phi}, \overline{\chi_{0,S}}))$. Here “ $\overline{}$ ” denotes the reduction modulo a prime ideal above p determined via the embedding ι_p . As $\text{Im}(\bar{f}) \subset \mathbb{F}_p(\overline{\phi}, \overline{\chi_{0,S}})$, the above version of reciprocity law is evident in the setup. We conclude that Ξ_K is invariant under the action of $\text{Gal}(\mathbb{F}/\mathbb{F}_p(\overline{\phi}, \overline{\chi_{0,S}}))$.

Let $C_K \subset \widehat{\text{Cl}}_K$ the subgroup generated by these characters and let $H_K \subset \text{Cl}_K$ be the orthogonal complement of C_K . The orthogonal complement is with respect to the pairing on $\widehat{\text{Cl}}_K$ and its character group. Recall that $|H_K| = |\text{Cl}_K|/|C_K|$.

In view of the Fourier inversion on Cl_K and the hypothesis on the periods, we have

$$(2.3) \quad \bar{f}(x_\sigma) = \sum_{i=1}^{\ell_K} c_{K,i} \chi_{K,i}(\sigma).$$

Here $c_{K,i} \in \mathbb{F}$ are the Fourier coefficients possibly dependent on K and determined by the pairs $(\bar{f}, \chi_{K,i})$. From the above formula on H_K (2.3), we note that f is constant on H_K .

We now recall a refinement of Iwaniec's equidistribution. By a result of Michel ([32, §6]), the special points

$$\left\{ x_\sigma \mid \sigma \in H'_K, [\text{Cl}_K : H'_K] \leq |D_K|^\delta \right\} \subset X_U$$

become equidistributed on X_U with respect to the probability measure as $|D_K| \rightarrow \infty$ for an absolute constant $\delta > 0$ (remark below).

It thus follows that

$$|C_K| \gg |D_K|^\delta.$$

Otherwise, this contradicts the non-constancy in Lemma 2.5.

From the Galois-stability of Ξ_K and an elementary Lemma 2.9 below, we conclude

$$\ell_K \gg_\epsilon \log(|\text{Cl}_K|)^{1-\epsilon}$$

for any $\epsilon > 0$.

This finishes the proof. \square

Remark 2.8. (1). In [32, §6], the equidistribution is proven for the special points arising from the maximal order \mathcal{O}_K . In this case, δ can be taken to be $1/2115$. The hypothesis seems to be made for simplicity and partly arises from the availability of explicit Waldspurger formula at the time. In [13], the formula has been established unconditionally.

(2). The approach only requires surjectivity of the map

$$\varphi_K : H_K \rightarrow X_U$$

with H_K as in the proof for a sufficiently large discriminant.

(3). Under the assumptions in Theorem 3.9, it seems tempting to predict

$$\#\left\{\chi \in \mathfrak{X}_K \mid v_p(P_f(\chi)) = 0\right\} \gg_\epsilon |\text{Cl}_K|^{1-\epsilon}$$

for any $\epsilon > 0$.

(4). For related equidistribution results on a quaternionic Shimura variety, we refer to [53].

The following elementary lemma is used in the above proof.

Lemma 2.9. *Let \mathbb{F}_q be a finite field with characteristic p . Let G be a finite abelian group and \widehat{G} the \mathbb{F}^\times -valued character group. Suppose that $p \nmid |G|$. Let $S \subset \widehat{G}$ be a Galois-stable subset i.e.*

(St) *If $\chi \in S$, then the Galois conjugates $\chi^\sigma \in S$ for any $\sigma \in \text{Gal}(\overline{\mathbb{F}}/\mathbb{F}_q)$.*

Let $\widehat{H}_S \subset \widehat{G}$ be the subgroup generated by S . Consider a sequence (G_i, S_i) as above with $|G_i| \rightarrow \infty$. Suppose that there exists δ with $0 < \delta < 1$ such that $|\widehat{H}_{S_i}| \gg |G_i|^\delta$. For $\epsilon > 0$, we then have

$$|S_i| \gg_\epsilon \ln(|G_i|)^{1-\epsilon}.$$

Proof. As H_S is abelian, say

$$H_S \simeq \mathbb{Z}/(n_1) \times \mathbb{Z}/(n_2) \times \dots \times \mathbb{Z}/(n_d)$$

with $n_1 \geq n_2 \geq \dots \geq n_d$.

It follows that there exists an element χ_1 in the generating set S with order n_1 . The number of Galois conjugates of χ_1 over \mathbb{F}_q equals the degree $[\mathbb{F}_q(\mu_{n_1}) : \mathbb{F}_q]$. Here μ_l denotes a primitive l^{th} root of unity over \mathbb{F}_p . By duality, the subgroup generated by χ_1 corresponds to a quotient $H_S \rightarrow C_1$. Note that

$$\widehat{H}_S / \widehat{C}_1 \simeq \mathbb{Z}/(n_2) \times \dots \times \mathbb{Z}/(n_d).$$

We conclude that there exists an element χ_2 in the generating set S with order n_2 .

Repeating the process, we have

$$|S| \geq \sum_i [\mathbb{F}_q(\mu_{n_i}) : \mathbb{F}_q] \geq \sum_i \frac{\varphi(n_i)}{(n_i, q-1)}.$$

As

$$\sum_i \frac{\varphi(n_i)}{(n_i, q-1)} \gg_\epsilon \left(\sum_i \ln(n_i) \right)^{1-\epsilon}$$

for any $\epsilon > 0$, this finishes the proof. The last estimate follows from [40]. \square

Remark 2.10. (1). Considering arbitrary finite products of $\mathbb{Z}/(2)$, the estimate is optimal in general.

(2). For a sequence $(G_i)_i$ as in Lemma 2.9 with the number of prime factors $\nu(|G_i|)$ being bounded, the proof shows that the lower bound for a minimal Galois-stable subset of the character group can be improved significantly.

2.3. Tate–Shafarevich Groups I. In this subsection, we describe bounds for the Tate–Shafarevich group in the analytic rank zero case in terms of a toric period based on an analysis Euler system of toric periods due to Bertolini–Darmon ([2]) as refined by Nekovář ([36]).

For convenience, let us briefly recall the Birch and Swinnerton-Dyer (BSD) conjecture.

In the beginning of this subsection, let A be an abelian variety over a number field K such that $\mathcal{O}_L \subset \text{End}(A)$ for a number field L . Let A^\vee be the dual abelian variety. Let $L(s, A)$ be the L-series arising from the Tate module of A at the finite primes of $\mathcal{O} = \mathcal{O}_L$. In particular, it is without Euler factor at infinite places.

For a torsion \mathcal{O} -module M , let $\text{Fitt}_{\mathcal{O}}(M)$ denote the Fitting ideal of M .

We have the following fundamental

Conjecture 2.11 (BSD). *Let $L(s, A)$ be the L-series for an abelian variety A over a number field K such that $\mathcal{O} = \mathcal{O}_L \subset \text{End}(A)$ for a number field L . Then, the following holds.*

- (1). *The L-series has analytic continuation to \mathbb{C} .*
- (2). *We have $\text{ord}_{s=1} L(s, A) = \text{rank}_{\mathcal{O}} A(K)$.*
- (3). *Let $L^*(1, A) \in L \otimes_{\mathbb{Q}} \mathbb{C}$ be the leading coefficient of the L-series. Then,*

$$\mathcal{L}(A) := \frac{L^*(1, A)}{\Omega(A/K)R(A/K)} \cdot \prod_v \text{Fitt}_{\mathcal{O}}(c_v(A))^{-1} \cdot \text{Fitt}_{\mathcal{O}}(A(K)_{\text{tor}}) \text{Fitt}_{\mathcal{O}}(A^\vee(K)_{\text{tor}}) \subset L \otimes \mathbb{C}$$

is a fractional ideal of \mathcal{O} for $r = \text{ord}_{s=1} L(s, A) = \text{rank}_{\mathcal{O}} A(K)$, period $\Omega(A/K)$, regulator $R(A/K)$ ([8]) and Tamagawa number c_v . Moreover,

$$\mathcal{L}(A) = \text{Fitt}_{\mathcal{O}}(\text{III}(A/K)).$$

For a prime \mathfrak{p} of \mathcal{O}_L , we refer to the asserted equality in (3) for the \mathfrak{p} -parts as the \mathfrak{p} -part of the BSD ($\text{BSD}_{\mathfrak{p}}$).

We now consider our setup. Let the notation be as in §2.2. In particular, L_ϕ denotes the Hecke field of the newform $\phi \in S_2(\Gamma_0(N))$. Let π the cuspidal automorphic representation of $\text{GL}_2(\mathbb{A})$ corresponding to the newform ϕ so that we have the equality

$$L(s, \pi) = L(s, \phi)$$

of complex L-functions. Let A be a GL_2 -type abelian variety over \mathbb{Q} associated to ϕ such that $\mathcal{O}_{L_\phi} \subset \text{End}(A)$.

To state the results, we introduce some notions. Let us begin with some results on the image of a p -adic Galois representation associated to a modular GL_2 -type abelian variety over the rationals ([36, App. B]). For simplicity, we suppose that A does not have CM.

Let $\Gamma \subset \text{Aut}(L_\phi/\mathbb{Q})$ be the group of inner twists of π . Recall that an inner twist of π is a pair (σ, χ) with σ an embedding of L_ϕ and χ a Dirichlet character such that there exists an isomorphism

$$\sigma\pi \simeq \pi \otimes \chi$$

with $\sigma\pi$ the σ -conjugate. For such a σ , we have a unique Dirichlet character χ_σ such that the above isomorphism holds ([37, B.3]). Let L^Γ be the fixed subfield of L_ϕ corresponding to Γ . For $\sigma \in \Gamma$, we thus have an inner twist arising from χ_σ . Let $\mathbb{Q}_\Gamma/\mathbb{Q}$ be the extension corresponding to $\bigcap_{\sigma \in \Gamma} \ker(\chi_\sigma)$. Let $\mathfrak{p}_\phi|p$ be a prime in L_ϕ and \mathfrak{p}_Γ the corresponding prime in L^Γ .

From the \mathfrak{p} -adic Tate-module of A , we have the \mathfrak{p}_ϕ -adic Galois representation $\rho_{\mathfrak{p}_\phi} : G_{\mathbb{Q}} \rightarrow \text{GL}_2(\mathcal{O}_{L_\phi, \mathfrak{p}_\phi})$. It induces a Galois representation

$$\rho_{\mathfrak{p}_\phi} : G_{\mathbb{Q}_\Gamma} \longrightarrow \text{GL}_2(\mathcal{O}_{L^\Gamma, \mathfrak{p}_\Gamma}).$$

In view of the work of Ribet ([39]) and Momose ([35]),

$$\rho_{\mathfrak{p}_\phi}(G_{\mathbb{Q}_\Gamma}) \subseteq H_\Gamma := \left\{ x \in \text{GL}_2(\mathcal{O}_{L^\Gamma, \mathfrak{p}_\Gamma}) \mid \det(x) \in \mathbb{Z}_p^\times \right\}$$

is an open subgroup (for example, see [37, B.5.2]). Moreover, the equality holds for all but finitely many primes \mathfrak{p} .

We say that $\rho_{\mathfrak{p}_\phi}$ has maximal image if the equality holds in the above inclusion. Here we only mention that $\rho_{\mathfrak{p}_\phi}$ is maximal for all but finitely many primes \mathfrak{p} . If $\rho_{\mathfrak{p}_\phi}$ is maximal, then we evidently have

$$\text{GL}_2(\mathbb{Z}_p) \subset \text{Im}(\rho_{\mathfrak{p}_\phi}).$$

For $\ell|N$, let π_ℓ be the corresponding local representation of $\mathrm{GL}_2(\mathbb{Q}_\ell)$. Let $d_\ell \geq 1$ be such that $d_\ell\mathbb{Z}$ is generated by $[M_\ell : \mathbb{Q}_\ell]$ where M_ℓ varies over extensions of \mathbb{Q}_ℓ such that the base change of π_ℓ to M_ℓ is either a spherical or a Steinberg representation of $\mathrm{GL}_2(M_\ell)$. Note that, $d_\ell = 1$ if $\ell \nmid N$.

Let B be a definite quaternion algebra and $\chi \in \mathfrak{X}_K$ a character over an imaginary quadratic field K as in §2.1. Recall that f is a toric form on $B_\mathbb{A}^\times$ and $U \subset (B_\mathbb{A}^{(\infty)})^\times$ the toric level corresponding to f . Let I be an ideal of \mathcal{O}_{L_ϕ} given by

$$I = \langle \ell + 1 - a_\ell \mid \ell \text{ prime}, \ell \nmid N, \ell \in \ker(\widehat{\mathbb{Q}}^\times \rightarrow \mathbb{Q}_+^\times \backslash \widehat{\mathbb{Q}}^\times / \mathrm{Nrd}(U)) \rangle.$$

Here a_ℓ denotes the T_ℓ -eigenvalue of the newform ϕ and Nrd the reduced norm map.

In the rest of this subsection, let $L = L_{\phi, \chi}$. Let $A_\chi = A_K \otimes_{\mathcal{O}_{L_\phi}} \mathcal{O}_L$ be the Serre tensor where the absolute Galois group G_K acts on \mathcal{O}_L via χ . In regards to certain aspects of the arithmetic corresponding the pair (A, χ) , it turns out that A_χ is a relevant abelian variety.

The main result of this subsection regarding the Tate–Shafarevich group of the Serre tensor A_χ is the following.

Proposition 2.12. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform. Let K/\mathbb{Q} be an imaginary quadratic extension and χ an unramified finite order Hecke character over K as above. Let p be a prime and \mathfrak{p}_ϕ (resp. \mathfrak{p}) the prime above p of the coefficient fields L_ϕ (resp. L) corresponding to the fixed embedding ι_p . Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_{L_\phi} \subset \mathrm{End}(A)$. For $\ell|N$, let c_ℓ be the corresponding Tamagawa number. Suppose that the following holds.*

- (1) *The p -adic Galois representation $\rho_{\mathfrak{p}_\phi}$ has maximal image,*
- (2) *$p \nmid 6N \cdot (\prod_\ell c_\ell d_\ell) \cdot I \cdot h_K$ and*
- (3) *The toric period $P_f(\chi)$ corresponding to the pair (f, χ) is a \mathfrak{p} -unit with f the p -primitive toric form on a definite quaternion algebra as in last subsection.*

Then,

$$\mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0 \quad \text{for } |D_K| \gg 0.$$

As $p \nmid h_K$, we also have

$$\mathrm{III}(A/H_K)^\chi[\mathfrak{p}^\infty] \cong \mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0 \quad \text{for } |D_K| \gg 0.$$

Proof. When the toric period $P_f(\chi)$ does not vanish, the Bertolini–Darmon method to bound $\mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty]$ shows that

$$\mathfrak{p}^{C_\chi + v_p(P_f(\chi))} \cdot \mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0$$

for a non-negative integer C_χ ([37, 2.9.6]). The constant C_χ possibly depends on p .

Under the hypotheses (1) and (2), we show that

$$C_\chi = 0$$

for all $\chi \in \mathfrak{X}_K$ as $|D_K| \gg 0$.

In the following exposition, we suppose some familiarity with [36, §2].

In view of the description of C_χ , we consider the following.

- (i) As the Galois representation $\rho_{A, \mathfrak{p}_\phi}$ has maximal image, only finitely many imaginary quadratic extensions are contained in $\mathbb{Q}(A[\mathfrak{p}^\infty])$. Let K be an imaginary quadratic field with $|D_K| \gg 0$.

For $p \geq 5$, let $\begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix} \in \mathrm{GL}_2(\mathbb{Z}_p)$ such that $\lambda_1 = \pm 1$ and $\lambda_2^2 - 1$ is a p -adic unit. Since the image is maximal, there exists $g_K \in G_{\mathbb{Q}}$ with trivial action on \mathbb{Q}_χ such that

$$- \rho_{\mathfrak{p}_\phi}(g_K) = \begin{pmatrix} \lambda_1 & \\ & \lambda_2 \end{pmatrix}.$$

– g_K does not act trivially on K .

In the notation of [37, §2], we thus have $C_2 = 0$ for $|D_K| \gg 0$.

- (ii) We now show that

$$\mathrm{Im}(\mathcal{O}_{L, \mathfrak{p}}[G_{\mathbb{Q}}] \rightarrow \mathrm{End}_{\mathcal{O}_{L, \mathfrak{p}}}(A_\chi[\mathfrak{p}^\infty])) = M_2(\mathcal{O}_{L, \mathfrak{p}}).$$

As $A_\chi[\mathfrak{p}]$ is absolutely irreducible $G_{\mathbb{Q}}$ -module, by Burnside’s lemma

$$\mathrm{Im}(\mathcal{O}_{L, \mathfrak{p}}[G_{\mathbb{Q}}] \rightarrow \mathrm{End}_{\mathcal{O}_{L, \mathfrak{p}}}(A_\chi[\mathfrak{p}])) = \mathrm{End}_{\mathcal{O}_{L, \mathfrak{p}}}(A_\chi[\mathfrak{p}]).$$

The assertion thus readily follows from Nakayama lemma.

In the notation of [37, §2], we thus have $C_4 = 0$.

(iii) We show that

$$\mathrm{Im}(O_{L,\mathfrak{p}}[G_{H_\chi}] \rightarrow \mathrm{End}_{O_{L,\mathfrak{p}}}(A_\chi[\mathfrak{p}^\infty])) = M_2(O_{L,\mathfrak{p}})$$

for $p > 3$ and $|D_K| \gg 0$. Here H_χ denotes the extension of K cut out by the finite order Hecke character χ over K .

In fact, we show the following. Suppose that K is not a subfield of $\mathbb{Q}(A_\chi[\mathfrak{p}^\infty])$. Then,

$$O_{L,\mathfrak{p}}[\rho_{\mathfrak{p}}(G_{H_\chi})] = M_2(O_{L,\mathfrak{p}}).$$

Let M denote the intersection of $\mathbb{Q}(A_\chi[\mathfrak{p}^\infty])$ and H_χ . As K is not a subfield of $\mathbb{Q}(A_\chi[\mathfrak{p}^\infty])$, the extension M/\mathbb{Q} is abelian. Moreover, the corresponding Galois group has exponent 2 as $\mathrm{Gal}(M/\mathbb{Q}) \simeq \mathrm{Gal}(MK/K)$ and the latter extension is both cyclotomic and anticyclotomic. It follows that

$$O_{L,\mathfrak{p}}[H_\Gamma^2] \subset O_{L,\mathfrak{p}}[\rho_{\mathfrak{p}}(G_{H_\chi})].$$

Here H_Γ is as in the description of maximal image. Here and in what follows, for a subgroup $H' \subset \mathrm{GL}_2(O_{L,\mathfrak{p}})$, let H'^2 denotes the subgroup generated by the square of elements in H' .

For $p > 3$, note that $\mathbb{Z}_p[\mathrm{GL}_2(\mathbb{Z}_p)^2] = \mathbb{Z}_p[\mathrm{GL}_2(\mathbb{Z}_p)]$. As $\mathrm{GL}_2(\mathbb{Z}_p) \subset H_\Gamma$, we conclude that

$$M_2(O_{L,\mathfrak{p}}) = O_{L,\mathfrak{p}}[\mathrm{GL}_2(\mathbb{Z}_p)^2] \subset O_{L,\mathfrak{p}}[\rho_{\mathfrak{p}}(G_{H_\chi})].$$

In the notation of [37, §2], we thus have $C_8 = 0$ for $|D_K| \gg 0$.

(iv) Let m be a sufficiently large positive integer and $n = m + C_0$ for the constant C_0 in [37, 2.7.9]. Let $H_{n,\chi} = \mathbb{Q}(T/\mathfrak{p}^n)H_\chi$ for $T = A_\chi[\mathfrak{p}^\infty]$.

Let $g_n \in \mathrm{Gal}(H_{n,\chi}/\mathbb{Q})$ be the restriction of g_K to $H_{n,\chi}$. Here g_K is an element as in the paragraph preceeding the vanishing of C_2 . Let $H'_{n,\chi}$ be the subfield of $H_{n,\chi}$ fixed by $\langle g_n^2 \rangle$. Let $\mathrm{Res} : H^1(H_\chi, T/\mathfrak{p}^m) \rightarrow H^1(H_{n,\chi}, T/\mathfrak{p}^m)$ and $\mathrm{Res}' : H^1(H_\chi, T/\mathfrak{p}^m) \rightarrow H^1(H'_{n,\chi}, T/\mathfrak{p}^m)$ be the restriction maps. We show that

$$\ker(\mathrm{Res}') = \ker(\mathrm{Res}) = 0$$

for $p > 3$ and $|D_K| \gg 0$.

This is based on the proof of [20, Prop. 9.1]. By inflation–restriction, we have

$$\ker(\mathrm{Res}) = H^1(\mathrm{Gal}(H_{n,\chi}/H_\chi), A_\chi[\mathfrak{p}^m]).$$

The Galois action gives rise to an embedding $\rho : \mathrm{Gal}(H_{n,\chi}/H_\chi) \hookrightarrow \mathrm{GL}(A_\chi[\mathfrak{p}^n])$. Suppose that K is not a subfield of $\mathbb{Q}(A_\chi[\mathfrak{p}^\infty])$ and let M be as in the discussion preceding the vanishing of C_8 . As $\mathrm{Gal}(M/\mathbb{Q})$ is abelian with exponent two, $\rho((\mathrm{Gal}(\mathbb{Q}(A_\chi[\mathfrak{p}^n])/\mathbb{Q}))^2)$ is a subgroup of $\rho(\mathrm{Gal}(H_{n,\chi}/H_\chi))$, and therefore

$$Z_0 \subset Z(\mathrm{Im}(\rho)) \subset \mathrm{Im}(\rho),$$

where Z_0 is the image of $(\mathbb{F}_p^\times)^2$ under the Teichmüller lifting. The existence follows from the maximality of the image. As $p > 3$, the subgroup Z_0 is non-trivial with order prime to p . It follows that

$$H^i(Z_0, A_\chi[\mathfrak{p}^m]) = 0$$

for $i \geq 0$. The spectral sequence

$$H^i(\mathrm{Gal}(H_{n,\chi}/H_\chi)/\rho^{-1}(Z_0), H^j(Z_0, A_\chi[\mathfrak{p}^m])) \implies H^{i+j}(\mathrm{Gal}(H_{n,\chi}/H_\chi), A_\chi[\mathfrak{p}^m])$$

thus implies the desired vanishing of $\ker(\mathrm{Res})$.

Recall that the elements g_n^2 does not belong to the center $Z(\mathrm{Im}(\rho))$. Going modulo $\langle g_n^2 \rangle$, the above argument thus also applies for $\ker(\mathrm{Res}')$.

In the notation of [37, §2], we thus have $C_7 = 0$ for $|D_K| \gg 0$.

(v). In view of our hypothesis $p \nmid \prod_\ell c_\ell$ and the description of Tamagawa numbers in [26, Prop. 3], we have $C_6 = 0$ in the notation of [37, §2].

(vi). In view of our hypothesis $p \nmid \prod_\ell d_\ell$, we have $C_5 = 0$ in the notation of [37, §2].

We now recall

$$C_\chi = 3C_2 + 12C_4 + C_5 + C_6 + C_7 + C_8 + v_p(I \cdot [H_\chi : K]) + v_p([H_\chi : K])$$

for all $\chi \in \mathfrak{X}_K$ ([37, 2.9.6]). As $p \nmid h_K$, we thus conclude

$$C_\chi = 0$$

for all $\chi \in \mathfrak{X}_K$ such that $|D_K| \gg 0$ ([37, 2.9.6]). This finishes the proof for $\mathrm{III}(A_\chi)$.

In view of [30], the rest follows for $\mathrm{III}(A/H_K)$ from the result for the Serre tensor A_χ . \square

Remark 2.13. Based on an analysis of the Euler system method for toric periods, it may be checked that there exists a constant C such that

$$\#\mathrm{III}(A_\chi)[\mathfrak{p}^\infty] \Big| p^{C+v_p(P_f(\chi))}$$

for all $\chi \in \mathfrak{X}_K$ as $|D_K| \gg 0$. In this sense, the toric periods control the size of the Tate–Shafarevich groups.

2.4. Analytic Sha I. In this subsection, we describe bounds for analytic Sha in the analytic rank zero case in terms of a toric period based on an analysis of explicit Waldspurger formula due to Cai–Shu–Tian ([13]). The consideration along with the main result in §2.3 leads us to establish the p -part of corresponding Birch and Swinnerton-Dyer conjecture for a class of primes p .

We first introduce a definition.

Definition 2.14. For an abelian variety A over a number field L , we refer to the ideal $\mathcal{L}(A)$ in the formula part of the BSD conjecture for A over L (Conjecture 2.11) as the analytic Sha.

We now introduce our setup. Let the notation and assumptions be as in §2.3. In particular, A denotes a GL_2 -type abelian variety over the rationals corresponding to a weight two newform $\phi \in S_2(\Gamma_0(N))$ and χ a finite order Hecke character over an imaginary quadratic field K such that

$$\epsilon(A, \chi) = 1$$

((RN)). Moreover, A_χ denotes the Serre tensor of A by χ which is an abelian variety over K with $\mathcal{O}_L \subset \mathrm{End}(A_\chi)$.

As above, we refer to $\mathcal{L}(A_\chi)$ as the analytic Sha of A_χ .

In this subsection, we consider the conjecture in the analytic rank zero case. We thus suppose that

$$L(1, A_\chi) \neq 0.$$

To introduce an explicit Waldspurger formula, we begin with some notation. Let B be a definite quaternion algebra as in §2.1. Let $R \subset B$ be an order as in [13, §1.1]. Recall that f is a toric form on $B_{\mathbb{A}}^\times$ and $U \subset (B_{\mathbb{A}}^{(\infty)})^\times$ the level corresponding to f . Let X_U be the corresponding Shimura set. Let $N = N^+ N^-$ for N^+ (resp. N^-) precisely divisible by split (resp. non-split) primes in the extension K/\mathbb{Q} . Recall that S denotes the set of prime divisors of N^∞ .

For any $f_i = \sum_x f_i(x) \cdot x \in \mathbb{C}[X_U]$ with $i = 1, 2$, we put

$$\langle f_1, f_2 \rangle = \sum_x w_x^{-1} f_1(x) f_2(x \tau_B).$$

Here w_x is the cardinality of $(B^\times \cap g \widehat{R}^\times g^{-1}) / \{\pm 1\}$ with $g \in \widehat{B}^\times$ any representative of x and $\tau_B \in N_{\widehat{B}^\times}(\widehat{R}^\times)$ such that

- for $\ell | N^+$, $\tau_{B, \ell} \in N_{B_\ell^\times}(R_\ell^\times) \setminus \mathbb{Q}_\ell^\times R_\ell^\times$ the Atkin–Lehner operator;
- for $\ell | N^-$, $\tau_{B, \ell} t_\ell = \overline{t}_\ell \tau_{B, \ell}$ for any $t_\ell \in K_\ell^\times$ where \overline{t}_ℓ is the Galois conjugate of t_ℓ .
- for $\ell \nmid N$, $\tau_{B, \ell} = 1$.

We consider Hermitian pairing $(\ , \)_{\widehat{R}^\times}$ on the space $\mathbb{C}[X_U]$ of \mathbb{C} -valued functions on the Shimura set X_U is given by

$$(f_1, f_2)_{\widehat{R}^\times} = \sum_x w_x^{-1} f_1(x) \overline{f_2(x)}$$

with $\overline{}$ being the complex conjugation.

From [13, Thm. 1.8], we have an explicit Waldspurger formula given by

$$L(1, A_\chi) = 2^{-\#\Sigma_D+2} \cdot \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|^2}} \cdot \frac{P_{f_1}(\chi) P_{f_2}(\chi^{-1})}{(f_1, f_2)_{\widehat{R}^\times}}.$$

Here $\Sigma_D = \{\ell | \ell(N, D_K)\}$, $f_1 \in V(\pi_B, \chi)$ and $f_2 \in V(\pi_B, \chi^{-1})$ are non-zero vectors. For the definition of $V(\pi_B, \chi)$, we refer to [13, Def. 1.7]. Here we only mention that f_1 can be taken to be the toric form f as in §2.1.

In the analytic rank zero case, the analytic Sha of A_χ is thus given by

$$\mathcal{L}(A_\chi) = \frac{\pi^2(\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2} \Omega(A_\chi/K)} \cdot \frac{P_{f_1}(\chi) P_{f_2}(\chi^{-1})}{(f_1, f_2)_{\widehat{R}^\times}} \cdot \frac{\mathrm{Fitt}(A_\chi(K)_{\mathrm{tor}}) \mathrm{Fitt}(A_{\chi^{-1}}^\vee(K)_{\mathrm{tor}})}{2^{\#\Sigma_D-5} \cdot [\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \cdot \prod_v \mathrm{Fitt}(c_v(A_\chi))}.$$

We have the following key proposition regarding the variation of the \mathfrak{p} -part of analytic Sha in terms of toric periods.

Proposition 2.15. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_\phi \subset \text{End}(A)$.*

There exists a finite set Σ_A of primes of \mathcal{O}_ϕ only dependent on A , such that for any prime $\mathfrak{p}_0 \notin \Sigma_A$ of \mathcal{O}_ϕ and any $\chi \in \mathfrak{X}$ with $L(1, A_\chi) \neq 0$, the ideal

$$\frac{\mathcal{L}(A_\chi)}{P_f(\chi)^2}$$

is coprime to \mathfrak{p}_0 . Here f is the $\chi_{0,S}$ -toric vector as in last subsection. In particular, if $P_f(\chi)$ is a \mathfrak{p} -unit, then $\mathcal{L}(A_\chi)$ is also a \mathfrak{p} -unit as long \mathfrak{p} lies above some $\mathfrak{p}_0 \notin \Sigma_A$.

Proof. Let $f'_1 = f$ be a \mathfrak{p} -minimal toric test vector satisfying (F1) and (F2) in §2.1. Let $J \in B$ be as in [14, §2.1] and f'_2 the J -translate of f'_1 given by

$$f'_2(x) = f'_1(xJ).$$

Locally, f'_2 differs from f'_1 only at the places dividing N .

Note that

$$P_{f'_2}(\chi^{-1}) = P_{f'_1}(\chi).$$

We now recall the S -version of Waldspurger formula in [13, Thm. 1.9]. It states that

$$L(1, \phi, \chi) = 2^{-\#\Sigma_D+2} \cdot C_\infty \cdot \frac{\langle \phi^0, \bar{\phi}^0 \rangle_{U_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|}} \cdot \frac{P_{f'_1}(\chi) P_{f'_2}(\chi^{-1})}{\langle f'_1, f'_2 \rangle_{\widehat{R}^\times}} \cdot \prod_{v \in S \setminus \{\infty\}} \frac{\beta^0(f_{1,v}, f_{2,v})}{\beta^0(f'_{1,v}, f'_{2,v})}.$$

Here ϕ^0 is a normalised new vector fixed by level $U_1(N)$ and the pairing $\langle \cdot, \cdot \rangle_{U_0(N)}$ is as in [13, §1.3]. Further, $f_1 = f \in V(\pi_B, \chi)$ and $f_2 \in V(\pi_B, \chi^{-1})$ are non-zero vectors, the local pairing $\beta^0(\cdot, \cdot)$ is as in [13, Thm. 1.6] and C_∞ denotes the constant in [13, Thm. 1.8].

It follows that the analytic Sha is given by

$$\mathcal{L}(A_\chi) = \frac{C_\infty \cdot (\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2} \Omega(A_\chi/K)} \cdot \frac{P_f(\chi)^2}{(f_1, f_2)_{\widehat{R}^\times}} \cdot \frac{\text{Fitt}(A_\chi(K)_{\text{tor}}) \text{Fitt}(A_{\chi^{-1}}^\vee(K)_{\text{tor}})}{2^{\#\Sigma_D-5} [\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \cdot \prod_v \text{Fitt}(c_v(A_\chi))} \cdot \prod_{v \in S \setminus \{\infty\}} \frac{\beta^0(f_{1,v}, f_{2,v})}{\beta^0(f'_{1,v}, f'_{1,J,v})}.$$

We now discuss horizontal variation of the terms in the right hand side of the above expression except the toric period. It suffices to show that the terms have bounded prime divisors as the pair (K, χ) -varies.

By strong approximation, we can choose

$$J \in \left(\prod_{v|N} B_v^\times \right) \cdot \text{GL}_2(\widehat{\mathbb{Z}}^{(N)}) \cap B^\times.$$

Since the S -type of K is fixed, it thus follows that

$$\prod_{v \in S \setminus \{\infty\}} \frac{\beta^0(f_{1,v}, f_{2,v})}{\beta^0(f'_{1,v}, f'_{1,J,v})}$$

is a fixed rational number as K varies.

We have

$$\Omega(A_\chi/K) = \Omega(A/K)$$

as fractional ideals up to ideals dividing the level N ([8, §4.4.1]).

In view of the definition of the period $\Omega(A/K)$ and the description of the Tamagawa numbers for the Serre tensor A_χ , it follows that there exists a nonzero ideal \mathfrak{a}_1 such that

$$\mathfrak{a}_1 \subseteq \frac{C_\infty \cdot (\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2} \Omega(A_\chi/K)} \cdot \prod_v \text{Fitt}(c_v(A_\chi)).$$

Moreover, the support of prime divisors of \mathfrak{a}_1 is bounded as K varies.

From Proposition 2.16, there exists a nonzero ideal \mathfrak{a}_2 such that

$$\mathfrak{a}_2 \subseteq \text{Fitt}(A_\chi(K)_{\text{tor}}) \text{Fitt}(A_{\chi^{-1}}^\vee(K)_{\text{tor}})$$

with support of prime divisors bounded as K varies.

In view of the earlier expression for the analytic Sha, this finishes the proof. \square

The following finiteness is used in the proof.

Proposition 2.16. *Let A be a non-CM GL_2 -type abelian variety over the rationals. Then, the set $\bigcup_K A(H_K)_{\mathrm{tor}}$ is finite as K varies imaginary quadratic fields ([12, Prop. 3.13]).*

Along with Proposition 2.12, we have the following immediate consequence for the BSD.

Corollary 2.17. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_\phi \subset \mathrm{End}(A)$.*

There exists a finite set Σ'_A of primes of \mathcal{O}_ϕ only dependent on A , such that the following holds. For any prime $\mathfrak{p}_0 \notin \Sigma'_A$ of \mathcal{O}_ϕ and for $\chi \in \mathfrak{X}$ with the toric period $P_f(\chi)$ being a \mathfrak{p} -unit for \mathfrak{p} above \mathfrak{p}_0 , the \mathfrak{p} -part of BSD holds for the Serre tensor A_χ as long as $p \nmid h_K$ for $(p) = \mathfrak{p}_0 \cap \mathbb{Q}$.

The exceptional set of primes Σ_A in Proposition 2.15 admits an explicit description under mild hypothesis.

Proposition 2.18. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let K be an imaginary quadratic field and χ an unramified finite order anticyclotomic Hecke character over K as above. Let $N = N^+ N^-$ where $l|N^+$ (resp. $l|N^-$) if and only if $l|N$ and split in K (resp. non-split in K). Let p be a prime and $\mathfrak{p}_0|p$ a prime of L_ϕ . Suppose that the following holds.*

- (1) $(N, D_K) = 1$,
- (2) N^- is square-free with odd number of prime factors,
- (3) $p \nmid 6ND_K$ and
- (4) The Galois representation ρ_{A, \mathfrak{p}_0} has maximal image and the residual representation $\overline{\rho}_{A, \mathfrak{p}_0}$ is ramified at $\ell|N^-$ with $\ell^2 \equiv 1 \pmod{p}$.

For all primes $\mathfrak{p}|\mathfrak{p}_0$ of $L = L_{\phi, \chi}$, we then have

$$\mathrm{ord}_{\mathfrak{p}} \mathcal{L}(A_\chi) = 2 \cdot \mathrm{ord}_{\mathfrak{p}}(P_\chi(f)) - 2 \cdot \sum_{\ell|N^+, \chi|_{D_\ell}=1} \mathrm{ord}_{\mathfrak{p}}(c_\ell).$$

Proof. We first describe an explicit form of Waldspurger formula under the hypotheses that N^- is square-free with odd number of prime factors.

We have

$$L(1, \phi, \chi) = \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|}} \frac{P_f(\chi)^2}{\langle f, f \rangle} \chi_{N^+}^{-1}(N^+).$$

Here the notation is as in the paragraphs following Conjecture 2.11. This follows from an analysis of an explicit version in [13, §1.3]. We indicate the sketch below.

Let $J \in B^\times$ such that $Jt = \bar{t}J$ for any $t \in K$ as in the proof of Proposition 2.15. We have

$$(2.4) \quad \frac{P_{f_J}(\chi)}{(f, f)} = \frac{P_f(\chi)}{\langle f, f \rangle} \chi_{N^+}^{-1}(N^+)$$

for f_J being the J -translate as in the proof of Proposition 2.15.

The equality can be deduced as follows. We may assume

$$J \in \iota_{N^+}^{-1} \left[\begin{pmatrix} N^+ & 0 \\ 0 & 1 \end{pmatrix} \right] \cdot \tau_B \cdot \widehat{R}^\times.$$

For $\ell|N^+$, we thus have

$$f(tJ_\ell) = \varepsilon_\ell(\pi_\ell) f \left[t^{(\ell)} \iota_\ell^{-1} \begin{pmatrix} t_\ell N_\ell & 0 \\ 0 & t_{\bar{\ell}} \end{pmatrix} \right], \quad t \in \widehat{K}^\times$$

and

$$P_{f_{J_\ell}}(\chi) = \chi_\ell^{-1}(N_\ell) P_{f_{\tau_B, \ell}}(\chi).$$

As f_{τ_B} and f both in the one-dimensional space $V(\pi_B, \chi)$, there exists a constant $c \in S^1$ such that

$$f_{\tau_B} = cf.$$

We thus have

$$\frac{P_{f_J}(\chi)}{(f, f)} = \frac{P_{f_{\tau_B}}(\chi)}{(f, f)} \chi_{N^+}^{-1}(N^+) = \frac{c P_f(\chi)}{(f, f)} \chi_{N^+}^{-1}(N^+) = \frac{P_f(\chi)}{\langle f, f \rangle} \chi_{N^+}^{-1}(N^+).$$

It thus follows that

$$\mathcal{L}(A_\chi) = \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2} \Omega(A_\chi/K)} \cdot \frac{P_f(\chi)^2}{\langle f, f \rangle} \cdot \frac{\mathrm{Fitt}(A_\chi(K)_{\mathrm{tor}}) \mathrm{Fitt}(A_{\chi^{-1}}^\vee(K)_{\mathrm{tor}})}{\prod_v \mathrm{Fitt}(c_v(A_\chi))} \cdot \chi_{N^+}^{-1}(N^+).$$

We now consider \mathfrak{p} -adic valuation of the terms in the right hand side of the above expression. Recall that we have

$$\Omega(A_\chi/K) = \Omega(A/K)$$

as fractional ideals up to primes dividing N .

In view of the hypothesis (4) on Galois image, both $A_\chi(K)$ and $A_{\chi^{-1}}^\vee(K)$ have no non-trivial \mathfrak{p}_0 -torsion point.

From ([38, (12)]), we recall that

$$\mathrm{ord}_{\mathfrak{p}_0} \left(\frac{8\pi^2(\phi, \phi)}{\Omega_A(f, f)} \right) = \mathrm{ord}_{\mathfrak{p}_0} \left(\prod_{\ell|N^-} c_\ell \right).$$

The proposition thus follows by noting that

$$\mathrm{ord}_{\mathfrak{p}_0} \left(\prod_v c_v(A_\chi) \right) = \mathrm{ord}_p \left(\prod_{\ell|N^-} c_\ell \prod_{\ell|N^+, \chi|_{D_\ell}=1} c_\ell^2 \right).$$

□

We have the following immediate consequence.

Corollary 2.19. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let K be an imaginary quadratic field and χ an unramified finite order anticyclotomic Hecke character over K as above. Let $N = N^+ N^-$ where $l|N^+$ (resp. $l|N^-$) if and only if $l|N$ and split in K (resp. non-split in K). Let p be a prime and $\mathfrak{p}_0|p$ a prime of L_ϕ . Suppose that the following holds.*

- (1) $(N, D_K) = 1$,
- (2) N^- is square-free with odd number of prime factors,
- (3) $p \nmid 6ND_K \cdot h_K$ and
- (4) The Galois representation ρ_{A, \mathfrak{p}_0} has maximal image and the residual representation $\bar{\rho}$ is ramified at $\ell|N^-$ with $\ell^2 \equiv 1 \pmod{p}$.

For all primes $\mathfrak{p}|\mathfrak{p}_0$ of $L = L_\phi(\chi)$, the \mathfrak{p} -part of BSD holds for the Serre tensor A_χ whenever the toric period $P_f(\chi)$ is a \mathfrak{p} -unit.

3. HEEGNER POINTS

In this section, we consider Heegner points arising from a Shimura curve associated to an indefinite quaternion algebra over the rationals. In §3.1, we introduce the setup. In §3.2, we recall generalities regarding CM points on a Shimura curve. In §3.3-3.4., we prove horizontal p -indivisibility of Heegner points. In §3.5 and §3.6, we deduce the consequences of the p -indivisibility for Tate–Shafarevich group and analytic Sha, respectively.

3.1. Setup. In this subsection, we introduce the setup regarding Heegner points.

Let $\phi \in S_2(\Gamma_0(N))$ be a newform with weight 2 and level $\Gamma_0(N)$ for $N \geq 3$. In particular, we consider newforms with trivial central character. Let L_ϕ denote the Hecke field corresponding to the newform ϕ .

Let B be an indefinite quaternion algebra over \mathbb{Q} such that there is an irreducible automorphic representation π_B on $B_\mathbb{A}^\times$ whose Jacquet-Langlands correspondence is the automorphic representation associated to ϕ . Recall $B_\mathbb{A} = B \otimes_\mathbb{Q} \mathbb{A}$.

Let $S = \mathrm{Supp}(N\infty)$. Let $K_{0,S} \subset B_S$ be a \mathbb{Q}_S -subalgebra such that $K_{0,\infty} = \mathbb{C}$ and $K_{0,v}/\mathbb{Q}_v$ is separable quadratic. For any $v \in S$, we say that v is non-split if $K_{0,v}$ is a field and split otherwise. Let

$$U_{0,S} := \prod_{v \in S, v \text{ split}} \mathcal{O}_{K_{0,v}}^\times \times \prod_{v \in S, v \text{ non-split}} K_{0,v}^\times.$$

Suppose we are given a finite order character $\chi_{0,S} : U_{0,S} \longrightarrow \overline{\mathbb{Q}}^\times$ with conductor one such that the following holds.

- (LC1) $\omega \cdot \chi_{0,S}|_{\mathbb{Q}_S^\times \cap U_{0,S}} = 1$.
- (LC2) $\epsilon(\phi, \chi_{0,v})\chi_{0,v}\eta_v(-1) = \epsilon(B_v)$ for all places $v|N$ with the local root number $\epsilon(\phi, \chi_{0,v})$ corresponding to the Rankin–Selberg convolution.

Fix a maximal order $R^{(S)}$ of $B_{\mathbb{A}}^{(S)} \cong M_2(\mathbb{A}^{(S)})$. Let $U^{(S)} = R^{(S)\times}$. Note that $U^{(S)}$ is a maximal compact subgroup of $B_{\mathbb{A}}^{(S)\times} \cong \mathrm{GL}_2(\mathbb{A}^{(S)})$.

Let p be an odd prime such that $p \nmid N$. As in the introduction, we fix an embedding $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$. Let v_p be a p -adic valuation. Let \mathbb{F} be an algebraic closure of \mathbb{F}_p .

Let Θ_S denote the set of imaginary quadratic extensions as in Definition 2.1. Recall that there exist infinitely many imaginary quadratic extensions K/\mathbb{Q} with $K \in \Theta_S$ ([7], [51] and [1]). For $K \in \Theta_S$, we fix a torus embedding ι_K as in §2.1.

For each $K \in \Theta_S$, let $\mathfrak{X}_{K, \chi_{0,S}}$ denote the set of finite order characters χ over K as in Definition 2.2. Recall that the conductor of χ equals one. Moreover, we have

$$(\mathrm{RN}) \quad \epsilon(\phi, \chi) = -1.$$

Here $\epsilon(\phi, \chi)$ denotes the global root number of the Rankin–Selberg convolution corresponding to the pair (ϕ, χ) .

In the rest of the section, we let $\Theta = \Theta_S$ and $\mathfrak{X}_K = \mathfrak{X}_{K, \chi_{0,S}}$.

Lemma 3.1. *The set \mathfrak{X}_K is non-empty for all but finitely many imaginary quadratic fields K with $K \in \Theta$. Moreover, it is a homogenous space for the class group Cl_K .*

Proof. The same argument as in the proof of Lemma 2.3 applies. \square

Let $A = A_\phi$ be an abelian variety over \mathbb{Q} associated to ϕ such that $\mathcal{O}_{L_\phi} \subset \mathrm{End}(A)$.

As in [52], we have the representation of $B_{\mathbb{A}}^{(\infty), \times}$ over the field $M := \mathrm{End}^0(A/\mathbb{Q})$ given by

$$\pi^B = \varinjlim_{U \subset B_{\mathbb{A}}^{(\infty), \times}} \mathrm{Hom}_{\xi}^0(X_U, A).$$

Here ξ is a Hodge class on $X = \varprojlim_U X_U$. Recall that

$$\pi^B \otimes_M \mathbb{C} \simeq \pi_{B,f}$$

for $\pi_{B,f}$ being the finite part of π_B .

Let \mathcal{O} be the ring of integers of the field $\mathbb{Q}(\phi, \chi_{0,S})$ generated over \mathbb{Q} by the Hecke eigenvalues of ϕ and the values of $\chi_{0,S}$. Let \mathfrak{p} be the prime ideal of \mathcal{O} corresponding to ι_p . Let $\mathcal{O}_{(\mathfrak{p})}$ be the localisation of the integer ring \mathcal{O} at the prime ideal \mathfrak{p} .

We have the following existence of toric test vectors.

Lemma 3.2. *There exists a non-zero form $f \in \pi_B$ defined over $\mathcal{O}_{(\mathfrak{p})}$ satisfying the following.*

- (F1) *The subgroup $U_{0,S}$ acts on f via $\chi_{0,S}$ and*
- (F2) *$f \in \pi_B^{U^{(S)}}$.*

Proof. The same argument as in the proof of Lemma 2.4 applies. \square

Note that, f is spherical outside S .

We further normalise f to be \mathfrak{p} -primitive in the sense arising from geometric definition of modular forms (§3.2.2).

Let U be an open compact subgroup of $B_{\mathbb{A}}^{(\infty), \times}$ such that $U = U_S U^{(S)}$ with $U_S \subset B_S^\times$ and $U_S = \prod_{v \in S} U_v$ such that

- (L1) $f \in \pi_B^U$ and
- (L2) $\mathcal{O}_{0,v}^\times \subset U_v$ for all $v \in S \setminus \{\infty\}$.

Let X_U be the corresponding Shimura set of level U . As a modular parametrisation, f corresponds to a morphism

$$f : X_U \rightarrow A \otimes_{\mathcal{O}_\phi} \mathcal{O}$$

and it maps the Hodge class ξ to the identity element. Here $A \otimes_{\mathcal{O}_\phi} \mathcal{O}$ denotes the Serre-tensor. We may also regard f as a $\overline{\mathbb{Z}}_{(p)}$ -valued weight two modular form at CM points on the Shimura curve X_U . For generalities on CM points, we refer to §3.2.

Let $f^{(p)}$ be the corresponding p -depletion. Let d be the Katz p -adic differential operator. In view of the p -adic Waldspurger formula ([3] and [29]), the CM values of weight zero p -adic modular form $d^{-1}(f^{(p)})$ are closely related to p -adic logarithm of Heegner points arising from CM points on the Shimura curve.

3.2. Shimura curve. In this subsection, we briefly describe generalities on Shimura curves, modular forms, CM points and the underlying Igusa tower. For a detailed treatment, we refer to [23, Ch. 7] and [9, §2].

As in §3.1, p denotes an odd prime and \mathbb{F} an algebraic closure of \mathbb{F}_p . Let W denote the Witt ring $W(\mathbb{F})$. Let \mathcal{W} be the strict Henselisation of $\mathbb{Z}_{(p)}$ given by $\iota_p^{-1}(W)$.

3.2.1. Shimura curve. We describe generalities regarding Shimura curves.

Let B/\mathbb{Q} be an indefinite quaternion algebra as in §3.1. In particular, $p \nmid \text{disc}(B)$. Let $(X_U)_U$ be the corresponding tower of Shimura curves indexed by open compact subgroups $U \subset B_{\mathbb{A}}^{(\infty), \times}$ and $X = \varprojlim_U X_U$ the underlying Shimura variety.

We recall the following ([17]).

Theorem 3.3. *Let the notation and assumptions be as above.*

- (1). *For a sufficiently small level U , the Shimura curve X_U admit a smooth canonical model over the rationals.*
- (2). *For a sufficiently small level U prime to p , the Shimura curve $X_{U/\mathbb{Q}}$ admits a smooth p -integral model $X_{U/\mathbb{Z}_{(p)}}$.*

We also recall that X_U represents a moduli functor classifying a class of elliptic curves or abelian surfaces with level U -structure. The difference in the moduli functors depends on whether B is split. For a description of the moduli problem, we refer to [9, §2.2]. Let $f_U : \mathcal{A}_U \rightarrow X_{U/\mathbb{Z}_{(p)}}$ be the universal abelian scheme.

For level U as in §3.1, the Shimura curve $X_{U/\mathbb{Q}}$ admits a p -integral model smooth at CM points with complex multiplication by \mathcal{O}_K . This follows from the hypothesis (L2) and part (iv) of Definition 2.1.

In what follows, we fix a sufficiently small level U prime to p unless otherwise stated.

3.2.2. Modular forms. We briefly describe the notion of geometric modular forms.

For an odd prime p , let A be a $\mathbb{Z}_{(p)}$ -algebra. A weight two, p -integral modular form with level U over A is an element of $H^0(X_{U/A}, \mathcal{L}_{U/A})$ for a line bundle $\mathcal{L}_{U/\mathbb{Z}_{(p)}}$ on the Shimura curve $X_{U/\mathbb{Z}_{(p)}}$ ([21] and [22]).

When A is nothing but $\mathcal{O}_{(\mathfrak{p})}$ for \mathcal{O} an integer ring of a number field and \mathfrak{p} a prime above p in \mathcal{O} , we say that a modular form over A is \mathfrak{p} -primitive if its image in $H^0(X_{U/\mathbb{F}_{\mathfrak{p}}}, \mathcal{L}_{U/\mathbb{F}_{\mathfrak{p}}})$ is non-zero. Here $\mathbb{F}_{\mathfrak{p}}$ denotes the residue field of \mathcal{O} at the prime \mathfrak{p} .

3.2.3. CM points. We describe generalities on CM points.

We identify \mathbb{A}_K^{\times} as a subgroup of $B_{\mathbb{A}}^{\times}$ under the embedding $\iota_K : K \hookrightarrow B$ for an imaginary quadratic field K as in §3.1. Let U be an open compact subgroup of $B_{\mathbb{A}}^{(\infty), \times}$ as in §3.1 satisfying (L1) and (L2).

As $\iota_K(\mathcal{O}_K) \subset U$, the choice of embedding ι_K gives rise to a map

$$\varphi_K : \text{Cl}_K \rightarrow X_{U/\overline{\mathbb{Q}}}.$$

For $\sigma \in \text{Cl}_K$, let $x_{\sigma} \in X_{U/\overline{\mathbb{Q}}}$ be the corresponding point on the Shimura curve. This is usually referred as a CM point on the Shimura curve. In what follows, the map φ_K plays an underlying role.

We now suppose that K is an imaginary quadratic field as above such that

(ord) p splits in K .

Based on the CM theory of Shimura–Taniyama–Weil and the hypothesis (ord), we have the following ([9, §2.4]).

Lemma 3.4. *Let the notation and assumptions be as above. Then, the CM points x_{σ} correspond to p -ordinary CM abelian varieties. Moreover, the CM points x_{σ} are defined over the Henselisation \mathcal{W} .*

In view of the lemma, we can consider the mod p reduction $x_{\sigma/\mathbb{F}} \in X_{U/\mathbb{F}}$ of the CM point x_{σ} . We have the following key proposition based on the Serre–Tate deformation theory.

Proposition 3.5. *Let the notation and assumptions be as above. For $\sigma, \tau \in \text{Cl}_K$, we have*

$$x_{\sigma/\mathbb{F}} = x_{\tau/\mathbb{F}} \iff \sigma = \tau.$$

Proof. For simplicity, let us consider the case when the quaternion B is split.

The condition (ord) guarantees a categorical equivalence between ordinary CM elliptic curves over the Witt ring W and ordinary elliptic curves over \mathbb{F} ([23, §6.3]). The equivalence is via the canonical lift in the deformation space over W .

For $\sigma \in \text{Cl}_K$, recall that the CM point x_σ is defined over \mathcal{W} , in particular over W . It is thus the canonical lift of $x_{\sigma/\mathbb{F}}$. This finishes the proof. \square

3.2.4. Igusa tower. We describe generalities on the Igusa tower.

Let U be a sufficiently small level prime to p as in §3.2.1 and $X_{U/\mathbb{Z}_{(p)}}$ corresponding Shimura curve. Let $X_{U/W}^{\text{ord}}$ be the p -ordinary locus. Adding p^∞ -level structure to the moduli problem corresponding to the Shimura curve on the ordinary locus, we obtain an étale Igusa tower

$$\pi : Ig \rightarrow X_{U/W}^{\text{ord}}$$

([9, §2.5]). Here we only mention that this amounts to trivialise the connected component $\mathcal{A}_U[p^\infty]^\circ$ or equivalently the étale quotient $\mathcal{A}_U[p^\infty]^{\text{ét}}$ of the p -divisible group $\mathcal{A}_U[p^\infty]$.

For an imaginary quadratic field K satisfying (ord), recall that $x_\sigma \in X_{U/W}^{\text{ord}}$ for $\sigma \in \text{Cl}_K$. In view of (ord), the \mathfrak{p}_K^∞ -divisible subgroup of the p -divisible group corresponding to x_σ gives rise to a canonical point in the Igusa tower above x_σ . Here \mathfrak{p}_K denotes the prime of K above p determined via the embedding ι_p . By abuse of notation, let x_σ also denote a point in the Igusa tower lying over the CM point.

For a W -algebra B , recall that a p -adic modular form g over B with tame level U is given by

$$g \in H^0(Ig/B, \mathcal{O}_{Ig/B}).$$

Here we mention that the geometric theory of p -adic modular forms plays an essential role in the p -indivisibility of Heegner points.

Regarding non-constancy of a mod p modular form on the Igusa tower, we have the following immediate consequence of Proposition 3.5 and the Brauer-Siegel lower bound for the size of class groups.

Corollary 3.6. *Let g be a non-constant p -adic modular form over \mathbb{F} with weight zero and tame level U . For $c \in \mathbb{F}$, the size of the CM fiber*

$$\left\{ x_\sigma \mid \sigma \in \text{Cl}_K, g(x_\sigma) = c \right\}$$

is bounded independent of K .

This corollary plays a key role in the mod p non-vanishing of toric periods of a p -adic modular form.

3.3. Non-vanishing. In this subsection, we prove the horizontal mod p non-vanishing of toric periods of a p -adic modular form on a Shimura curve.

Let the notation and assumptions be as in §3.1. In particular, f is a weight two modular form on the Shimura curve X_U and $f^{(p)}$ the p -depletion. Let g the weight zero p -adic modular form given by

$$g = d^{-1}(f^{(p)}).$$

Let Θ be the set of imaginary quadratic fields as in §3.1. Let $K \in \Theta$. For $\sigma \in \text{Cl}_K$, let $x_\sigma \in Ig/W$ be the corresponding CM point as in §3.2.4. For $\chi \in \widehat{\text{Cl}}_K$, let $P_g(\chi)$ be the toric period given by

$$(3.1) \quad P_g(\chi) = \frac{1}{h_K} \cdot \sum_{\sigma \in \text{Cl}_K} \chi(\sigma)^{-1} g(x_\sigma).$$

Note that g is defined over a finite flat discrete valuation ring W' over the Witt ring $W = W(\mathbb{F})$ as a p -adic modular form. Here \mathbb{F} denotes an algebraic closure of \mathbb{F}_p . Let \overline{g} denote the reduction of g modulo the maximal ideal of the local ring W' .

As g is p -integral and $p \nmid h_K$, the period $P_g(\chi)$ is p -integral. More precisely, $P_g(\chi) \in W'[\chi]$ as the underlying CM points are defined over W . Here $W'[\chi]$ denotes the extension of W' obtained by adjoining the values of χ . Note that the period possibly depends on the choice of the torus embedding ι_K .

Remark 3.7. (1). For $\chi \in \widehat{\text{Cl}}_K$, the non-vanishing of $P_g(\chi)$ implies that $\chi \in \mathfrak{X}_K$. This follows from g being $\chi_{0,S}$ -toric (part (1) of Lemma 3.8). The observation will be used in the proof of Theorem 3.9.

(2). We restrict to newforms with trivial central character and unramified Hecke characters over the imaginary quadratic fields for simplicity.

Let \mathfrak{p}_0 be the prime above p in the Hecke field L_ϕ corresponding to the newform determined via the embedding ι_p . Let $\rho_{\phi, \mathfrak{p}_0} : G_{\mathbb{Q}} \rightarrow \text{GL}_2(L_{\phi, \mathfrak{p}_0})$ be the corresponding p -adic Galois representation. Let $\overline{\rho}_{\phi, \mathfrak{p}_0}$ be the reduction modulo \mathfrak{p}_0 .

We have the following lemma regarding g .

Lemma 3.8. *Let the notation and assumptions be as above.*

- (1). The p -adic modular form g is $\chi_{0,S}^{-1}$ -invariant under the action of $K_{0,S}^\times$.
- (2). Assume that the mod p Galois representation $\overline{\rho}_{\phi,p_0}$ associated to the newform ϕ is absolutely irreducible. Then the weight zero p -adic modular form g is non-constant modulo p .

Proof. For the first part, note that the p -depletion only changes the Hecke action at p . In particular, the action of $K_{0,S}$ is unchanged. As

$$d^{-1}(f^{(p)}) = \lim_{j \rightarrow 0} d^{-1+j}(f^{(p)})$$

and each $d^{-1+j}(f^{(p)})$ is $\chi_{0,S}^{-1}$ -invariant under $K_{0,S}^\times$, this finishes the proof.

For the second part, we first recall the non-constancy of the p -depletion $f^{(p)}$ under irreducibility of the underlying mod p Galois representation ([9, Prop. 5.3]). Recall that the p -adic differential operator acts as $t \frac{d}{dt}$ on the t -expansion of a p -adic modular form around a CM point. The non-constancy of $d^{-1}(f^{(p)})$ thus follows. Here t denotes the Serre–Tate co-ordinate of the deformation space around the CM point ([9, §3.1]). \square

Our main result regarding the mod p non-vanishing of toric periods is the following.

Theorem 3.9. *Let $\phi \in S_2(\Gamma_0(N))$ be a newform. Let $p \nmid N$ be a prime such that*

(irr) the mod p Galois representation $\overline{\rho}_{\phi,p}$ is absolutely irreducible.

Let f be a toric test vector on an indefinite quaternion algebra as above, $f^{(p)}$ the corresponding p -depletion and $g = d^{-1}(f^{(p)})$ with d being the Katz p -adic differential operator. Let $\Theta = \Theta_S$ be the set of imaginary quadratic fields as above. For $K \in \Theta$, let \mathfrak{X}_K be the set of finite order Hecke character over K as in Definition 2.2. For $\chi \in \mathfrak{X}_K$, let $P_g(\chi)$ be the toric period corresponding to the pair (g, χ) .

For $\epsilon > 0$, we have

$$\#\left\{ \chi \in \mathfrak{X}_K \mid v_p(P_g(\chi)) = 0 \right\} \gg_\epsilon \log(|D_K|)^{1-\epsilon}.$$

as $K \in \Theta$ varies over imaginary quadratic fields with

(ord) p split.

Proof. Recall that g is a function on the Igusa tower Ig defined over W' and \overline{g} its reduction modulo the maximal ideal of W' . Note that \overline{g} is a function on the mod p Igusa tower $Ig_{\mathbb{F}}$. In view of the map $\varphi_K : \text{Cl}_K \rightarrow Ig$, we may regard g and \overline{g} as functions on the abelian group Cl_K .

The approach is based on the Fourier analysis on Cl_K and its relation to the mod p Igusa tower $Ig_{\mathbb{F}}$.

To begin with, the toric periods $P_g(\chi)$ modulo \mathfrak{m}_χ equal the χ -th Fourier coefficients of the function

$$\sigma \mapsto g(x_\sigma) \bmod \mathfrak{m}'$$

on the finite abelian group Cl_K . Here \mathfrak{m}_χ (resp. \mathfrak{m}') is the maximal ideal of $W'[\chi]$ (resp. W'). Here W' denotes a finite flat extension of W and $W'[\chi]$ its extension obtained by adjoining the values of χ as above. Let \mathfrak{m} denote the maximal ideal of W .

In what follows, we thus consider Fourier analysis of the function

$$\overline{g} : \text{Cl}_K \rightarrow \mathbb{F}$$

and its variation for $K \in \Theta$. As $p \nmid h_K$, the Fourier inversion works as usual.

Let us first consider the non-vanishing of at least one twist. In view of the Fourier inversion, it suffices to show that

$$(3.2) \quad \overline{g}(x_\sigma) \neq 0.$$

for at least one $\sigma \in \text{Cl}_K$ for all but finitely many $K \in \Theta$. By abuse of notation, x_σ denotes the reduction of the CM point x_σ modulo \mathfrak{m} .

In view of Proposition 3.5 and the Brauer–Siegel lower bound, the number of CM points

$$\left\{ x_\sigma \mid \sigma \in \text{Pic}_{K,c} \right\} \subset Ig_{\mathbb{F}}$$

increases with the discriminant of K . Thus, the p -primitivity of g readily implies 3.2 (part (2) of Lemma 3.8).

We now consider the growth in the number of non-vanishing twists. Suppose that there exists a positive integer ℓ_K such that exactly $\ell_K - 1$ of the toric periods $P_g(\chi)$ are non-vanishing modulo \mathfrak{m}_χ , for K with sufficiently large discriminant. Let

$$\Xi_K = \{\chi_{K,1}, \dots, \chi_{K,\ell_K-1}\}$$

be the set of the non-vanishing modulo \mathfrak{m}_χ twists.

From Shimura's reciprocity law,

$$\overline{P_g(\chi)} \neq 0 \iff \overline{P_g(\chi^\sigma)} \neq 0$$

for $\sigma \in \text{Gal}(\mathbb{F}/\mathbb{F}_p(\overline{\phi}, \overline{\chi_0, s}))$. Here “ $\overline{}$ ” denotes the reduction modulo a prime ideal above p determined via the embedding ι_p . We conclude that Ξ_K is invariant under the action of $\text{Gal}(\mathbb{F}/\mathbb{F}_p(\overline{\phi}, \overline{\chi_0, s}))$.

Let $C_K \subset \widehat{\text{Cl}}_K$ the subgroup generated by these characters and let $H_K \subset \text{Cl}_K$ be the orthogonal complement of C_K . The orthogonal complement is with respect to the pairing on $\widehat{\text{Cl}}_K$ and its character group. Recall that $|H_K| = |\text{Cl}_K|/|C_K|$.

In view of the Fourier inversion on Cl_K and the hypothesis on the periods, we have

$$(3.3) \quad \overline{g}(x_\sigma) = \sum_{i=1}^{\ell_K-1} c_{K,i} \chi_{K,i}(\sigma).$$

Here $c_{K,i} \in \mathbb{F}$ are the Fourier coefficients possibly dependent on K and determined by the pairs $(\overline{g}, \chi_{K,i})$.

From the above formula on H_K , it follows that g is constant on H_K . In view of part (2) of 3.8 and Corollary 3.6, there exists a constant c such that

$$|H_K| < c$$

as $|D_K| \rightarrow \infty$.

In particular,

$$|C_K| \gg_{\epsilon'} |D_K|^{\frac{1}{2}-\epsilon'}$$

for $\epsilon' > 0$.

From the Galois-stability of Ξ_K and Lemma 2.9, we conclude

$$\ell_K \gg_{\epsilon} \log(|D_K|)^{1-\epsilon}$$

for any $\epsilon > 0$.

This finishes the proof. \square

Remark 3.10. Under the assumptions in Theorem 3.9, it seems tempting to predict

$$\#\left\{\chi \in \mathfrak{X}_K \mid v_p(P_g(\chi)) = 0\right\} \gg_{\epsilon} |\text{Cl}_K|^{1-\epsilon}$$

for any $\epsilon > 0$.

3.4. p -indivisibility of Heegner points. In this subsection, we prove the horizontal p -indivisibility of Heegner points based on the horizontal mod p non-vanishing of toric periods.

Let the notation and assumptions be as in §3.1. In particular, we have modular parametrisation $f : X_U \rightarrow A_\phi \otimes_{\mathcal{O}_\phi} \mathcal{O}$ for the Shimura curve X_U and the abelian variety A_ϕ associated to a newform $\phi \in S_2(\Gamma_0(N))$. For choice of a base point for Albanese map of the Shimura curve X_U into its Jacobian, we follow [54, Rem. 6]. Here we only remark that existence of such a normalisation follows from maximality of the p -adic Galois representation $\rho_{\phi, \mathfrak{p}}$ associated to ϕ .

In what follows, we choose the parametrisation f to be $(\mathcal{O}_\phi, \mathfrak{p}_\phi)$ -optimal as in [54, §3.7]. Here \mathfrak{p}_ϕ denotes the prime of \mathcal{O}_ϕ above p arising from the embedding ι_p .

Recall, Θ denotes the set of imaginary quadratic fields in §3.1. For $K \in \Theta$, \mathfrak{X}_K denotes the set of finite order Hecke character over K as in Definition 2.2. For $\sigma \in \text{Cl}_K$, we have the same CM point $x_\sigma \in X_U/\overline{\mathbb{Q}}$ (§3.2.3). Let $\chi \in \mathfrak{X}_K$. Associated to the pair (f, χ) , we have Heegner point $P_f(\chi)$ given by

$$(3.4) \quad P_f(\chi) := \sum_{\sigma \in \text{Cl}_K} \chi(\sigma)^{-1} f(x_\sigma)$$

We have the following immediate.

Lemma 3.11. *Let the notation and assumptions be as above. Then,*

$$P_f(\chi) \in A(H_\chi)$$

for the extension H_χ/K cut out by χ and $A = A_\phi \otimes_{\mathcal{O}_\phi} \mathcal{O}$.

Fundamental p -adic invariants associated to a Heegner point are its p -adic height and p -adic logarithm. Regarding the p -adic logarithm of the Heegner point, we have the following fundamental p -adic Waldspurger formula due to Bertolini–Darmon–Prasanna ([3], [4]), Brooks ([6]) and Liu–Zhang–Zhang ([29]).

Theorem 3.12. *Let $\phi \in S_2(\Gamma_0(N))$ be a newform. Let $p \nmid N$ be a prime and $a_p(\phi)$ the corresponding Hecke eigenvalue. Let f be a toric test vector on an indefinite quaternion algebra as above and $g = d^{-1}(f^{(p)})$ with d being the Katz p -adic differential operator. Let K/\mathbb{Q} be an imaginary quadratic extension with p -split. Let \mathfrak{p}_K be the prime above p in K determined via the embedding ι_p and $\bar{\mathfrak{p}}_K$ its conjugate. Let χ be a finite order unramified Hecke character over K as above. Let $P_f(\chi)$ be the Heegner point corresponding to the pair (f, χ) .*

Then, we have

$$h_K \cdot P_g(\chi) = (1 - \chi^{-1}(\bar{\mathfrak{p}}_K)p^{-1}a_p(\phi) + \chi^{-2}(\bar{\mathfrak{p}}_K)p^{-1}) \log_{\omega_f}(P_f(\chi))$$

for the p -adic logarithm $\log_{\omega_f}(\cdot)$.

Proof. This is essentially proven in [3]. We recall the steps and refer to [3, §3] for details.

Note that

$$(3.5) \quad T_p(f) = a_p(\phi)f, \langle p \rangle(f) = f$$

for $\langle p \rangle$ being the diamond operator associated with p . Indeed, this follows from the construction of the toric test vector ϕ in §3.1.

Let F_f the Coleman primitive of the differential ω_f associated with f ([3, §3.7 and §3.8]). In view of [3, Lem. 3.23 and Prop. 3.24], we have

$$F_f(x_\sigma) = g(x_\sigma) - a_p(\phi)g(\mathfrak{p}_K * x_\sigma) + \frac{1}{p}g(\mathfrak{p}_K^2 * x_\sigma).$$

For the “ $*$ ”-action on the set of CM points, we refer to [3, (1.4.8)].

From [3, Lem. 3.23] and [4, Thm. 3.12], we have

$$F_f(x_\sigma) = \log_{\omega_f}(f(x_\sigma)).$$

The summation of the above identity over $\sigma \in \text{Cl}_K$ along with 3.5 and [3, Proof of Thm. 5.13] finishes the proof. \square

Remark 3.13. Based on Waldspurger formula, the left hand side of the above formula turns out to be a value of an anticyclotomic Rankin–Selberg p -adic L-function outside its interpolation range ([3], [6] and [29]). However, the above version only relies on Coleman integration and it suffices for our application to p -indivisibility of Heegner points.

In view of the p -adic Waldspurger formula and the p -indivisibility of toric periods (Theorem 3.9), we have the following p -indivisibility of Heegner points.

Theorem 3.14. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform. Let $p \nmid N$ be a prime such that*

(irr) the mod p Galois representation $\bar{\rho}_{\phi, \mathfrak{p}}$ is absolutely irreducible.

Let A_ϕ be the corresponding GL_2 -type abelian variety such that $\mathcal{O}_{L_\phi} \subset \text{End}(A)$. Let K be an imaginary quadratic field with $K \in \Theta$ for Θ as above. Let χ be a finite order Hecke character over K as above. Let \mathfrak{p} the prime above p of the coefficient field L corresponding to the fixed embedding ι_p . Let $P_f(\chi)$ be the Heegner point corresponding to the pair (f, χ) .

For $\epsilon > 0$, we have

$$\# \left\{ \chi \in \mathfrak{X}_K \mid P_f(\chi) \text{ } \mathfrak{p}\text{-indivisible in } A(H_\chi) \right\} \gg_\epsilon \log(|D_K|)^{1-\epsilon}$$

as K varies over imaginary quadratic fields with

(ord) p split.

Proof. Recall that p is unramified in the anticyclotomic extension H_χ . In view of Lemma 3.11 and the p -integrality of normalised p -adic logarithm, the p -indivisibility thus follows from Theorem 3.9 and Theorem 3.12. \square

Remark 3.15. Based on Jochnowitz congruence ([46] and [2]), above theorem can be deduced from Theorem 2.7 for variation over imaginary quadratic fields satisfying extra hypothesis that there exists a fixed prime inert in them.

3.5. Tate–Shafarevich groups II. In this subsection, we describe bounds for the Tate–Shafarevich group in the analytic rank one case in terms of a Heegner points based on an analysis Euler system of Heegner points due to Kolyvagin ([28]) as refined by Nekovář ([37]).

Let the notation be as in §3.1. In particular, L_ϕ denotes the Hecke field of the newform $\phi \in S_2(\Gamma_0(N))$. Let π the cuspidal automorphic representation of $\mathrm{GL}_2(\mathbb{A})$ corresponding to the newform ϕ so that we have the equality

$$L(s, \pi) = L(s, \phi)$$

of complex L-functions. Let A be a GL_2 -type abelian variety over \mathbb{Q} associated to ϕ such that $\mathcal{O}_{L_\phi} \subset \mathrm{End}(A)$.

Let B be a indefinite quaternion algebra and $\chi \in \mathfrak{X}_K$ a character over an imaginary quadratic field K as in §2.1. Recall that f is a toric form on $B_{\mathbb{A}}^\times$ and $U \subset (B_{\mathbb{A}}^{(\infty)})^\times$ the level corresponding to f . We have the underlying Heegner point $P_f(\chi) \in A(H_\chi)$.

In the rest of this subsection, let $L = L_{\phi, \chi}$. Let $A_\chi = A_K \otimes_{\mathcal{O}_{L_\phi}} \mathcal{O}_L$ be the Serre tensor where the absolute Galois group G_K acts on \mathcal{O}_L via χ . In regards to certain aspects of the arithmetic corresponding the pair (A, χ) , it turns out that A_χ is a relevant abelian variety.

The main result of this subsection regarding the Tate–Shafarevich group of the Serre tensor A_χ is the following.

Proposition 3.16. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform. Let K/\mathbb{Q} be an imaginary quadratic extension and χ an unramified finite order Hecke character over K as above. Let p be a prime and \mathfrak{p}_ϕ (resp. \mathfrak{p}) the prime above p of the coefficient fields L_ϕ (resp. L) corresponding to the fixed embedding ι_p . Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_{L_\phi} \subset \mathrm{End}(A)$. For $\ell|N$, let c_ℓ be the corresponding Tamagawa number. Suppose that the following holds.*

- (1) *The p -adic Galois representation $\rho_{A, \mathfrak{p}_\phi}$ has maximal image,*
- (2) *$p \nmid 6N \cdot (\prod_\ell c_\ell) \cdot h_K$,*
- (3) *There exists a polarisation $\varphi : A \rightarrow A^\vee$ with $p \nmid \deg(\varphi)$ for A^\vee the dual abelian variety.*
- (4) *The Heegner point $P_f(\chi)$ is \mathfrak{p} -indivisible with f the p -primitive toric form in last subsection.*

Then,

$$\mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0 \quad \text{for } |D_K| \gg 0.$$

As $p \nmid h_K$, we also have

$$\mathrm{III}(A/H_K)^\chi[\mathfrak{p}^\infty] \cong \mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0 \quad \text{for } |D_K| \gg 0.$$

Proof. When the Heegner point $P_f(\chi)$ is non-torsion and \mathfrak{p}_χ -indivisible, the Kolyvagin method to bound $\mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty]$ shows that

$$\mathfrak{p}^{C_\chi} \mathrm{III}(A_\chi/K)[\mathfrak{p}^\infty] = 0$$

for a non-negative integer C_χ ([36, 7.5.3 & 7.6.5]). The constant C_χ possibly depends on p .

Under the hypotheses (1)–(3), we show that

$$C_\chi = 0$$

for all $\chi \in \mathfrak{X}_K$ as $|D_K| \gg 0$.

In the following exposition, we suppose some familiarity with [36, §2].

Strictly speaking, the approach is slightly different which we now describe. Let $\mathcal{O} = \mathcal{O}_{\mathfrak{p}}$ and M a positive integer with $h_L|M$.

Let $\mathrm{Sel}_{M, \chi}$ be the classical Selmer group $\mathrm{Sel}(A_\chi/H_\chi, \mathfrak{p}^M)$ for \mathfrak{p}^M -descent on the abelian variety A_χ . Multiplication by \mathfrak{p}^M along with a choice of generator of \mathfrak{p}^M gives rise to the Kummer exact sequence

$$0 \rightarrow A_\chi(H_\chi)/\mathfrak{p}^M \xrightarrow{\delta} \mathrm{Sel}_{M, \chi} \rightarrow \mathrm{III}(A_\chi/H_\chi)[\mathfrak{p}^M] \rightarrow 0.$$

Let $\varpi \in \mathcal{O}_L$ be a uniformiser corresponding to \mathfrak{p} . Let

$$\kappa = \varpi^{C_1} \delta(P_f(\chi)) \in \mathrm{Sel}_{M, \chi}$$

for a non-negative integer C_1 considered below. Then, there exists a non-negative integer C_χ such that

$$\mathfrak{p}^{C_\chi} (\mathrm{Sel}_{M, \chi}/\mathcal{O} \cdot \kappa) = 0.$$

Under the hypotheses (1)–(3), we show that

$$C_1 = C_\chi = 0$$

for all $\chi \in \mathfrak{X}_K$ as $|D_K| \gg 0$. It would then follow

$$\mathrm{III}(A_\chi/H_\chi)[\mathfrak{p}^\infty] = 0.$$

In view of the description of C_1 and C_χ , we consider the following.

- (i) For a finite place v of H_χ , let $\widetilde{A_{\chi,v}}$ be the special fiber of the Néron model of A_χ over \mathcal{O}_{H_χ} at v . Let $\pi_0(\widetilde{A_{\chi,v}})$ be the \mathcal{O}_L -module of the connected components of $\widetilde{A_{\chi,v}}$. Let $C_{1,v}$ be smallest non-negative integer such that

$$\mathfrak{p}^{C_{1,v}} \pi_0(\widetilde{A_{\chi,v}})_{\mathfrak{p}} = 0.$$

Let

$$C_1 = \max_v C_{1,v}.$$

In view of the hypothesis (2), we have

$$C_1 = 0$$

([36, 7.2.1]).

- (ii) As in the proof of Proposition 2.12, we also have

$$\mathrm{Im}(\mathcal{O}_{L,\mathfrak{p}}[G_{H_\chi}] \rightarrow \mathrm{End}_{\mathcal{O}_{L,\mathfrak{p}}}(A_\chi[\mathfrak{p}^\infty])) = M_2(\mathcal{O}_{L,\mathfrak{p}})$$

for $p > 3$ and $|D_K| \gg 0$.

In the notation of [36], we thus have $C_3 = 0$.

- (iii) Let M_0 be the constant in [36, 5.1]. Let $H_{M,\chi} = \mathbb{Q}(A[\mathfrak{p}^{M+M_0}])H_\chi$ and $U_M = \mathrm{Gal}(H_{M,\chi}/H_\chi)$. Let $\mathrm{Res} : H^1(H_\chi, A[\mathfrak{p}^M]) \rightarrow H^1(H_{M,\chi}, A[\mathfrak{p}^M])^{U_M}$ be the restriction map. Based on the proof of (iv) in the proof of Proposition 2.12, we have

$$\ker(\mathrm{Res}) = 0.$$

for $p > 3$ and $|D_K| \gg 0$.

In the notation of [36], we thus have $C_3 = 0$.

- (iv) As $p \nmid h_K$, we have $C_4 = C_5 = 0$ in the notation of [36].
- (v) In view of our hypothesis $p \nmid \deg(\varphi)$, we have $C_6 = 0$ in the notation of [36].

We now recall

$$C_\chi = 2C_1 + 4C_2 + 4C_3 + C_5 + C_6.$$

if $\chi^2 = 1$ and

$$C_\chi = 4C_1 + 7C_2 + 7C_3 + 5C_4 + 2C_5 + 2C_6.$$

otherwise ([36, 7.5.3 & 7.6.5]).

We thus conclude

$$C_\chi = 0$$

for all $\chi \in \mathfrak{X}_K$ such that $|D_K| \gg 0$ ([37, 2.9.6]). This finishes the proof for $\mathrm{III}(A_\chi)$.

In view of [30], the rest follows for $\mathrm{III}(A/H_K)$ from the result for the Serre tensor A_χ . \square

Remark 3.17. Based on an analysis of Euler system method for Heegner points, it may be checked that there exists a constant C such that

$$\#\mathrm{III}(A_\chi)[\mathfrak{p}^\infty] \Big| p^{C+v_p(P_f(\chi))}$$

for all $\chi \in \mathfrak{X}_K^-$ as $|D_K| \gg 0$. Here $v_p(P_f(\chi))$ measures the \mathfrak{p} -divisibility of the Heegner point $P_f(\chi)$. In this sense, the Heegner points control the size of the Tate-Shafarevich groups.

3.6. Analytic Sha II. In this subsection, we describe bounds for analytic Sha in the analytic rank one case in terms of a Heegner point based on an analysis of explicit Waldspurger formula due to Cai-Shu-Tian ([13]). The consideration along with the main result in §3.5 leads us to establish the p -part of corresponding Birch and Swinnerton-Dyer conjecture for a class of primes p .

Let the notation and assumptions be as in §3.1. In particular, A denotes a GL_2 -type abelian variety over the rationals corresponding to a weight two newform $\phi \in S_2(\Gamma_0(N))$ and χ a finite order Hecke character over an imaginary quadratic field K such that

$$\epsilon(A, \chi) = -1$$

((RN)). Moreover, A_χ denotes the Serre tensor of A by χ which is an abelian variety over K with $\mathcal{O}_L \subset \mathrm{End}(A_\chi)$.

Recall that, we refer to $\mathcal{L}(A_\chi)$ in §2.3 as the analytic Sha of A_χ .

In this subsection, we consider the BSD conjecture in the analytic rank one case. We thus suppose that

$$L'(1, A_\chi) \neq 0.$$

To introduce an explicit Gross–Zagier formula, we begin with some notation. Let B be a definite quaternion algebra as in §3.1. Let $R \subset B$ be an order as in [13, §1.1]. Recall that f is a toric form on $B_\mathbb{A}^\times$ and $U \subset (B_\mathbb{A}^{(\infty)})^\times$ the level corresponding to f . Let X_U be the corresponding Shimura curve. Let $N = N^+ N^-$ for N^+ (resp. N^-) precisely divisible by split (resp. non-split) primes in the extension K/\mathbb{Q} . Recall that S denotes the set of prime divisors of $N\infty$.

For any $f_i \in \pi^B$ factoring through X_U with $i = 1, 2$ (§3.1), we put

$$\langle f_1, f_2 \rangle = \frac{1}{\text{vol}(X_U)} \deg((f_2 \circ \tau_B)^\vee \circ f_1).$$

Here a canonical polarisation is used on the Jacobian and $\tau_B \in N_{\widehat{B}^\times}(\widehat{R}^\times)$ such that

- for $\ell | N^+$, $\tau_{B,\ell} \in N_{B_\ell^\times}(R_\ell^\times) \setminus \mathbb{Q}_\ell^\times R_\ell^\times$ the Atkin–Lehner operator;
- for $\ell | N^-$, $\tau_{B,\ell} t_\ell = \overline{t}_\ell \tau_{B,\ell}$ for any $t_\ell \in K_\ell^\times$ where \overline{t}_ℓ is the Galois conjugate of t_ℓ .
- for $\ell \nmid N$, $\tau_{B,\ell} = 1$.

We consider another pairing $(\ , \)_{\widehat{R}^\times}$ is given by

$$(f_1, f_2)_{\widehat{R}^\times} = \frac{1}{\text{vol}(X_U)} \deg(f_2^\vee \circ f_1).$$

From [13, Thm. 1.5], we have an explicit Gross–Zagier formula given by

$$L'(1, A_\chi) = 2^{-\#\Sigma_D+2} \cdot \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|}} \cdot \frac{\langle P_{f_1}(\chi), P_{f_2}(\chi^{-1}) \rangle}{(f_1, f_2)_{\widehat{R}^\times}}.$$

Here $\Sigma_D = \{\ell | \ell | (N, D_K)\}$, $f_1 \in V(\pi_B, \chi)$ and $f_2 \in V(\pi_B, \chi^{-1})$ are non-zero vectors. For the definition of $V(\pi_B, \chi)$, we refer to [13, Def. 1.7]. Here we only mention that f_1 can be taken to be the toric form f as in §3.1.

In the analytic rank one case, the analytic Sha of A_χ is thus given by

$$\mathcal{L}(A_\chi) = \frac{\pi^2(\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2} \Omega(A_\chi/K)} \cdot \frac{\langle P_{f_1}(\chi), P_{f_2}(\chi^{-1}) \rangle}{(f_1, f_2)_{\widehat{R}^\times}} \cdot \frac{\text{Fitt}(A_\chi(K)_{\text{tor}}) \text{Fitt}(A_{\chi^{-1}}^\vee(K)_{\text{tor}})}{2^{\#\Sigma_D-5} \cdot [\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \cdot \prod_v \text{Fitt}(c_v(A_\chi))}.$$

We have the following key proposition regarding the variation of the \mathfrak{p} -part of analytic Sha in terms of Heegner points.

Proposition 3.18. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_\phi \subset \text{End}(A)$.*

There exists a finite set Σ_A of primes of \mathcal{O}_ϕ only dependent on A , such that for any prime $\mathfrak{p}_0 \notin \Sigma_A$ of \mathcal{O}_ϕ and any $\chi \in \mathfrak{X}$ with $L'(1, A_\chi) \neq 0$, the ideal

$$\frac{\mathcal{L}(A_\chi)}{[A_\chi(K) : \mathcal{O}_{\phi,\chi} P_f(\chi)]^2}$$

is coprime to \mathfrak{p}_0 . Here f is the $\chi_{0,S}$ -toric vector as in last subsection. In particular, if $P_f(\chi)$ is a \mathfrak{p} -indivisible, then $\mathcal{L}(A_\chi)$ is a \mathfrak{p} -unit as long \mathfrak{p} lies above some $\mathfrak{p}_0 \notin \Sigma_A$.

Proof. Let $f'_1 = f$ be a \mathfrak{p} -minimal toric test vector satisfying (F1) and (F2) in §3.1. Let $J \in B$ be analogue of the construction in [14, §2.1]. Let f'_2 the J -translate of f'_1 given by

$$f'_2(x) = f'_1(xJ).$$

Locally, f'_2 differs from f'_1 only at the places dividing N . We have

$$P_{f'_1}(\chi), P_{f'_2}(\chi^{-1}) \in A_\chi(K).$$

Let $\alpha : A \rightarrow A^\vee$ be an isogeny. It gives rise to an isogeny $\alpha : A_\chi \rightarrow A_{\chi^{-1}}^\vee$. Note that

$$P_{\alpha \circ f'_2}(\chi^{-1}) = \alpha \circ P_{f'_1}(\chi).$$

We now recall the S -version of Gross–Zagier formula in [13, Thm. 1.6]. It states that

$$L'(1, \phi, \chi) = 2^{-\#\Sigma_D+2} \cdot C_\infty \cdot \frac{\langle \phi^0, \overline{\phi}^0 \rangle_{U_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|}} \cdot \frac{\langle P_{f'_1}(\chi), P_{f'_2}(\chi^{-1}) \rangle}{\langle f'_1, f'_2 \rangle_{\widehat{R}^\times}} \cdot \prod_{v \in S \setminus \{\infty\}} \frac{\beta^0(f_{1,v}, f_{2,v})}{\beta^0(f'_{1,v}, f'_{2,v})}.$$

Here ϕ^0 is a normalised new vector fixed by level $U_1(N)$ and the pairing $\langle \cdot, \cdot \rangle_{U_0(N)}$ is as in [13, §1.3]. Further, $f_1 = f \in V(\pi_B, \chi)$ and $f_2 \in V(\pi_B, \chi^{-1})$ are non-zero vectors, the local pairing $\beta^0(\cdot, \cdot)$ is as in [13, Thm. 1.6] and C_∞ denotes the constant in [13, Thm. 1.8].

It follows that the analytic Sha is given by

$$\mathcal{L}(A_\chi) = \frac{C_\infty \cdot (\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2 \Omega(A_\chi/K)}} \cdot \frac{\langle P_{f'_1}(\chi), P_{f'_1}(\chi) \rangle_{A_\chi}}{(f_1, f_2)_{\widehat{R}^\times}} \cdot \frac{\text{Fitt}(A_\chi(K)_{\text{tor}}) \text{Fitt}(A_{\chi^{-1}}^\vee(K)_{\text{tor}})}{2^{\#\Sigma_D - 5} [\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \cdot \prod_v \text{Fitt}(c_v(A_\chi))} \cdot \prod_{v \in S \setminus \{\infty\}} \frac{\beta^0(f_{1,v}, f_{2,v})}{\beta^0(f'_{1,v}, f'_{1,v})}.$$

Here $\langle \cdot, \cdot \rangle_A$ is the pairing given by $\langle P, Q \rangle_A = \langle P, \alpha(Q) \rangle$.

We now discuss horizontal variation of the terms in the right hand side of the above expression except the Heegner point. It suffices to show that the terms have bounded prime divisors as the pair (K, χ) -varies.

Note that the terms are identical to the ones appearing in the proof of Proposition 2.15 for analytic Sha in the analytic rank zero case.

In view of the expression for the analytic Sha, we thus finish the proof. \square

Along with Proposition 3.16, we have the following immediate consequence for the BSD.

Corollary 3.19. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_\phi \subset \text{End}(A)$.*

There exists a finite set Σ'_A of primes of \mathcal{O}_ϕ only dependent on A , such that the following holds. For any prime $\mathfrak{p}_0 \notin \Sigma'_A$ of \mathcal{O}_ϕ and for $\chi \in \mathfrak{X}$ with the Heegner point $P_f(\chi)$ being a \mathfrak{p} -indivisible for \mathfrak{p} above \mathfrak{p}_0 , the \mathfrak{p} -part of BSD holds for the Serre tensor A_χ as long as $\mathfrak{p} \nmid h_K$ for $(p) = \mathfrak{p}_0 \cap \mathbb{Q}$.

The exceptional set of primes Σ_A in Proposition 3.18 admits an explicit description under mild hypothesis.

Proposition 3.20. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let K be an imaginary quadratic field and χ an unramified finite order anticyclotomic Hecke character over K as above. Let $N = N^+ N^-$ where $l|N^+$ (resp. $l|N^-$) if and only if $l|N$ and split in K (resp. non-split in K). Let p be a prime and $\mathfrak{p}_0|p$ a prime of L_ϕ . Suppose that the following holds.*

- (1) $(N, D_K) = 1$,
- (2) N^- is square-free with even number of prime factors,
- (3) $p \nmid 6ND_K$,
- (4) There exists an isogeny $\varphi : A \rightarrow A^\vee$ with $p \nmid \deg(\varphi)$ and
- (5) The Galois representation ρ_{A, \mathfrak{p}_0} has maximal image and the residual representation $\overline{\rho}_{A, \mathfrak{p}_0}$ is ramified at $\ell|N^-$ with $\ell^2 \equiv 1 \pmod{p}$.

For all primes $\mathfrak{p}|\mathfrak{p}_0$ of $L = L_{\phi, \chi}$, we then have

$$\text{ord}_{\mathfrak{p}} \mathcal{L}(A_\chi) = 2 \cdot \text{ord}_{\mathfrak{p}} [A_\chi(K) : \mathcal{O}_{\phi, \chi} P_{f_1}(\chi)] - 2 \cdot \sum_{\ell|N^+, \chi|_{D_\ell}=1} \text{ord}_{\mathfrak{p}}(c_\ell).$$

Proof. We first describe an explicit form of Gross–Zagier formula under the hypotheses that N^- is square-free with odd number of prime factors.

We have

$$L'(1, \phi, \chi) = \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{[\mathcal{O}_K^\times : \mathbb{Z}^\times]^2 \sqrt{|D_K|}} \cdot \frac{\langle P_f(\chi), P_f(\chi) \rangle_{A_\chi}}{\langle f, f \rangle} \chi_{N^+}^{-1}(N^+).$$

Here the notation is as in the earlier part of this subsection. This follows from an analysis of an explicit version in [13, §1.3]. As the proof is analogous to the proof of explicit version of the Waldspurger formula in Proposition 2.18, we skip the details.

It thus follows that

$$\mathcal{L}(A_\chi) = \frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{\sqrt{|D_K|^2 \Omega(A_\chi/K)}} \cdot \frac{\langle P_f(\chi), P_f(\chi) \rangle_{A_\chi}}{\langle f, f \rangle} \cdot \deg(\varphi) \cdot \frac{\text{Fitt}(A_\chi(K)_{\text{tor}}) \text{Fitt}(A_{\chi^{-1}}^\vee(K)_{\text{tor}})}{\prod_v \text{Fitt}(c_v(A_\chi))} \cdot \chi_{N^+}^{-1}(N^+).$$

We now consider \mathfrak{p} -adic valuation of the terms in the right hand side of the above expression.

Under the hypotheses, the test vector f is nothing but the new vector. From [54, Proof of Lem. 10.1], we thus have

$$\text{ord}_{\mathfrak{p}_0} \left(\frac{8\pi^2(\phi, \phi)_{\Gamma_0(N)}}{\Omega(A/K)(f, f)} \right) = \text{ord}_{\mathfrak{p}_0} \left(\prod_{\ell|N^-} c_\ell \right).$$

The other terms in the above expression for analytic Sha except the one involving Heegner point are identical to the ones appearing in the proof of Proposition 2.18. The proposition thus follows by noting that

$$\text{ord}_{\mathfrak{p}_0} \left(\prod_v c_v(A_\chi) \right) = \text{ord}_p \left(\prod_{\ell|N^-} c_\ell \prod_{\ell|N^+, \chi|_{D_\ell}=1} c_\ell^2 \right).$$

□

Remark 3.21. As in [13], there is a striking resemblance among explicit Waldspurger formula and explicit Gross–Zagier formula in our setup. Essentially, the terms other than the ones involving toric period and Heegner point turn out to be identical.

We have the following immediate consequence.

Corollary 3.22. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let K be an imaginary quadratic field and χ an unramified finite order anticyclotomic Hecke character over K as above. Let $N = N^+ N^-$ where $l|N^+$ (resp. $l|N^-$) if and only if $l|N$ and split in K (resp. non-split in K). Let p be a prime and $\mathfrak{p}_0|p$ a prime of L_ϕ . Suppose that the following holds.*

- (1) $(N, D_K) = 1$,
- (2) N^- is square-free with even number of prime factors,
- (3) $p \nmid 6ND_K \cdot h_K$,
- (4) There exists an isogeny $\varphi : A \rightarrow A^\vee$ with $p \nmid \deg(\varphi)$ and
- (5) The Galois representation ρ_{A, \mathfrak{p}_0} has maximal image and the residual representation $\bar{\rho}_{A, \mathfrak{p}_0}$ is ramified at $\ell|N^-$ with $\ell^2 \equiv 1 \pmod{p}$.

For all primes $\mathfrak{p}|\mathfrak{p}_0$ of $L = L_\phi(\chi)$, the \mathfrak{p} -part of BSD holds for the Serre tensor A_χ whenever the toric period $P_f(\chi)$ is \mathfrak{p} -indivisible.

4. MAIN RESULTS

In this section, we describe the main results.

4.1. Torus embedding. In this subsection, we describe a result on the existence of a suitable torus embedding into a quaternion algebra. The embedding plays a role in the proof of the main results in §5.2.

Let us first introduce the setup.

Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform and L_ϕ the corresponding Hecke field. Let π the cuspidal automorphic representation of $\text{GL}_2(\mathbb{A})$ corresponding to the newform ϕ so that we have the equality

$$L(s, \pi) = L(s, \phi)$$

of complex L-functions. Let A the abelian variety corresponding to ϕ such that $\mathcal{O}_{L_\phi} \subset \text{End}(A)$.

Let B be a quaternion algebra over \mathbb{Q} such that there exists an irreducible automorphic representation π_B on $B_{\mathbb{A}}^\times$ whose Jacquet-Langlands transfer is the automorphic representation π of $\text{GL}_2(\mathbb{A})$ associated to ϕ .

Let $S = \text{Supp}(N\infty)$. Let $K_{0,S} \subset B_S$ be a \mathbb{Q}_S -subalgebra such that $K_{0,\infty} = \mathbb{C}$ and $K_{0,v}/\mathbb{Q}_v$ is semi-simple quadratic. Fix a maximal order $R^{(S)}$ of $B_{\mathbb{A}}^{(S)} \cong M_2(\mathbb{A}^{(S)})$. Let $U^{(S)} = R^{(S)\times}$. Note that $U^{(S)}$ is a maximal compact subgroup of $B_{\mathbb{A}}^{\times(S)} \cong \text{GL}_2(\mathbb{A}^{(S)})$.

Lemma 4.1. *Let K be an imaginary quadratic field with $K_S \cong K_{0,S}$. Then, there exists an embedding $\iota : K \hookrightarrow B$ such that*

- (i). $\iota(K_S) = K_{0,S}$ and
- (ii). $\iota(\mathbb{A}_K^S) \cap U = \iota(\mathcal{O}_K^S)$.

Proof. Let $\iota_0 : K \hookrightarrow B$ be an embedding such that

$$\iota(\mathbb{A}_K^S) \cap U = \iota(\widehat{\mathcal{O}}_K^S).$$

For $v \in S$, we have $\iota(K_v) \simeq K_v$. Thus, there exists $b_v \in B_v^\times$ such that

$$b_v^{-1} \iota(K_v) b_v = K_{0,v}.$$

Let $b_S = (b_v)_{v \in S} \in B_S \subset B_{\mathbb{A}}^\times$.

In view of the choice of level U , strong approximation for the subgroup $B^1 \subset B^\times$ consisting of elements with identity norm implies that

$$B_{\mathbb{A}}^\times = B^\times \cdot U \prod_{v \in S} K_{0,v}.$$

There exists $b \in \widehat{B}^\times$ such that $b_S = bu$ with $b \in B^\times$ and $u \in U \prod_{v \in S} K_{0,v}$. We can take ι to be $\iota_0^b := b^{-1} \iota_0 b$. \square

Let p be an odd prime.

Definition 4.2. Let Θ_S denote the set of imaginary quadratic fields K such that

- (i) $K_S \cong K_{0,S}$ and
- (ii) $p \nmid h_K$.

For any imaginary quadratic field K with $K_S \cong K_{0,S}$, we fix an embedding ι_K as above from now.

4.2. Main results. In this section, we describe and prove the main results. The proof is essentially a compilation of the results in §2, §3 and §4.1.

Let the notation and assumptions be as in §3.1. In particular, $\phi \in S_2(\Gamma_0(N))$ is a non-CM newform and L_ϕ the Hecke field. Let A be the corresponding abelian variety with $\mathcal{O}_{L_\phi} \subset \text{End}(A)$.

Let p be an odd prime. As in the introduction, we fix the embeddings $\iota_\infty : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}$ and $\iota_p : \overline{\mathbb{Q}} \hookrightarrow \mathbb{C}_p$. For Θ in §4.1, let $K \in \Theta$. For $\chi \in \widehat{\text{Cl}}_K$, let \mathfrak{p}_χ be the prime above p of $L_{\phi,\chi}$ determined via the embedding ι_p .

For $\chi \in \widehat{\text{Cl}}_K$, let $A_\chi = A_K \otimes_{\mathbb{Z}} \mathbb{Z}[\chi]$ denote the Serre tensor where the absolute Galois group G_K acts on $\mathbb{Z}[\chi]$ via χ ([30]). Then A_χ is an abelian variety over K with the following properties:

$$\text{III}(A_\chi/K) \otimes_{\mathbb{Z}} \mathbb{Z}[h_K^{-1}] \cong \text{III}(A/H_{K,c})^\chi \otimes_{\mathbb{Z}} \mathbb{Z}[h_K^{-1}], \quad L(s, A_\chi) = L(s, A, \chi) \in \mathbb{C} \otimes_{\mathbb{Q}} L_{\phi,\chi}.$$

Let $\mathcal{L}(A_\chi)$ be the analytic Sha of A_χ (part (2) of Conjecture 2.11), which is conjectured to be a non-zero integral ideal of $L_\phi(\chi)$ as predicted by the BSD conjecture for the abelian variety A_χ over K .

Definition 4.3. A character $\chi \in \widehat{\text{Cl}}_K$ is said to be p -regular for A if the following conditions hold.

- $\text{rank}_{\mathcal{O}_{L_{\phi,\chi}}} A_\chi(K) = \text{ord}_{s=1} L(s, A_\chi/K)$, and
- Let $\mathfrak{p}_\chi|p$ be the prime ideal of $\mathcal{O}_{L_{\phi,\chi}}$ induced by the fixed embedding ι_p , then
 - (a) $\text{III}(A_\chi)$ is finite with trivial \mathfrak{p}_χ -part, and
 - (b) $\mathcal{L}(A_\chi)$ has trivial \mathfrak{p}_χ -part.

In particular, for a p -regular $\chi \in \widehat{\text{Cl}}_K$, the \mathfrak{p}_χ -part of the BSD conjecture holds for the Serre tensor A_χ .

Our result regarding p -regular characters is the following.

Theorem 4.4. Let A be a non-CM GL_2 -type abelian variety over the rationals. Then, there exists an explicit finite subset Σ_A of primes only dependent on A , such that for any prime $p \notin \Sigma_A$ and $\epsilon > 0$ we have

$$\#\left\{ \chi \in \widehat{\text{Cl}}_K \mid \chi \text{ is } p\text{-regular for } A \right\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic fields satisfying

- (i) $p \nmid h_K$, and
- (ii) either p splits in K or the root number $\epsilon(A, \chi_0)$ of the Rankin–Selberg convolution $L(s, A, \chi_0)$ equals $+1$ for some $\chi_0 \in \widehat{\text{Cl}}_K$.

Proof. Let $S = \text{supp}(N\infty)$ as before.

Let $K_{0,S}$ be a \mathbb{Q}_S -subalgebra such that $K_{0,\infty} = \mathbb{C}$ and $K_{0,v}/F_v$ is semi-simple quadratic. Let $U_{0,S}$ be as in §2.1. Suppose that there exists a finite order character $\chi_{0,S} : U_{0,S} \rightarrow \overline{\mathbb{Q}}^\times$ with conductor one satisfying (LC1) in §2.1 such that

$$\epsilon(\pi, \chi_{0,v}) \chi_{0,v} \eta_{0,v}(-1) = \pm 1$$

for $v \in S$.

From now, let $K \in \Theta_S$ (Definition 4.2). Let $\mathfrak{X}_{K,\chi_{0,S}}$ denote the set of finite order Hecke characters over K satisfying the conditions (i) and (iii) in Definition 2.2. We have

$$\mathfrak{X}_{K,\chi_{0,S}} = \mathfrak{X}_{K,\chi_{0,S}}^+ \bigcup \mathfrak{X}_{K,\chi_{0,S}}^-.$$

Here $\mathfrak{X}_{K,\chi_{0,S}}^+$ (resp. $\mathfrak{X}_{K,\chi_{0,S}}^-$) denotes the subset of Hecke characters χ over K such that $\epsilon(\pi, \chi) = 1$ (resp. $\epsilon(\pi, \chi) = -1$).

Note that

$$\Theta_S = \Theta_S^+ \cup \Theta_S^-.$$

Here Θ_S^+ (resp. Θ_S^-) denotes the subset of imaginary quadratic fields K such that there exists $\chi \in \mathfrak{X}_{K,\chi_{0,S}}^+$ (resp. $\mathfrak{X}_{K,\chi_{0,S}}^-$) with $\chi_S = \chi_{0,S}$ for some $\chi_{0,S}$. The above union need not be disjoint.

Note that there are only finitely many choices for $\chi_{0,S}$. For any such $\chi_{0,S}$, we consider a quaternion algebra B over \mathbb{Q} with $\text{ram}(B) \subset S$ such that

$$\epsilon(\pi, \chi_{0,v}) \chi_{0,v} \eta_{0,v}(-1) = \epsilon(B_v)$$

for $v \in S$.

As K varies over imaginary quadratic extensions satisfying the hypotheses in the Theorem, there are only finitely many choices for the characters $\chi_{0,S}$. So, it suffices to consider the case of a fixed character $\chi_{0,S}$.

In the case of root number 1, the result thus follows from Lemma 4.1, Theorem 2.7, Proposition 2.12 and Proposition 2.15.

In the case of root number -1 , the result thus follows from Lemma 4.1, Theorem 3.14, Proposition 3.16 and Proposition 3.18. \square

Remark 4.5. In view of the proof, the set of exceptional primes Σ_A arises from the ones in Proposition 2.12 (resp. Proposition 3.16) and Proposition 2.15 (resp. Proposition 3.18). Note that the set Σ_A can be larger than the set of exceptional primes for the p -minimality of III or analytic Sha when considered individually.

Based on the proof, we have the following explicit version of p -minimality in individual cases.

Corollary 4.6. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform with N square-free and L_ϕ the corresponding Hecke field. Let p be a prime and $\mathfrak{p}_\phi|p$ a prime of L_ϕ . Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_{L_\phi} \subset \text{End}(A)$. Suppose that the following holds.*

- (1) *The Galois representation $\rho_{A,\mathfrak{p}_\phi}$ has maximal image and*
- (2) *$p \nmid 6N \cdot (\prod_\ell c_\ell)$ with c_ℓ the Tamagawa number at ℓ .*

For any $\epsilon > 0$, we have

$$\#\{\chi \in \widehat{\text{Cl}}_K \mid \chi \text{ is } p\text{-regular for } A\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic fields satisfying

- (i) $p \nmid h_K$
- (ii) $(N, D_K) = 1$,
- (iii) N^- has odd number of prime factors and $\text{ord}_\ell(N^-) = 1$ for any prime $\ell|N$. Here $N = N^+ N^-$ where $\ell|N^+$ (resp. $\ell|N^-$) if and only if $\ell|N$ and split in K (resp. non-split in K).
- (iv) $p \nmid 6ND_K$ and
- (v) *The residual representation $\bar{\rho}_{A,\mathfrak{p}_\phi}$ is ramified at $\ell|N^-$ with $\ell^2 \equiv 1 \pmod{p}$.*

Proof. This follows from the proof of Theorem 4.4 along with Proposition 2.12 and Proposition 2.18. \square

Corollary 4.7. *Let $\phi \in S_2(\Gamma_0(N))$ be a non-CM newform with N square-free and L_ϕ the corresponding Hecke field. Let p be a prime and $\mathfrak{p}_\phi|p$ a prime of L_ϕ . Let A be an abelian variety corresponding to ϕ such that $\mathcal{O}_{L_\phi} \subset \text{End}(A)$. Suppose that the following holds.*

- (1) *The Galois representation $\rho_{A,\mathfrak{p}_\phi}$ has maximal image,*
- (2) *$p \nmid 6N \cdot (\prod_\ell c_\ell)$ with c_ℓ the Tamagawa number at ℓ and*
- (3) *There exists a polarisation $\varphi : A \rightarrow A^\vee$ with $p \nmid \deg(\varphi)$.*

For any $\epsilon > 0$, we have

$$\#\{\chi \in \widehat{\text{Cl}}_K \mid \chi \text{ is } p\text{-regular for } A\} \gg_\epsilon (\ln |D_K|)^{1-\epsilon}.$$

Here K varies over imaginary quadratic fields satisfying

- (i) $p \nmid h_K$
- (ii) p split in K and $(N, D_K) = 1$,
- (iii) N^- has even number of prime factors. Here $N = N^+ N^-$ where $\ell|N^+$ (resp. $\ell|N^-$) if and only if $\ell|N$ and split in K (resp. non-split in K).

- (iv) $p \nmid 6ND_K$ and
- (v) The residual representation $\bar{\rho}_{A, \mathfrak{p}, \phi}$ is ramified at $\ell | N^-$ with $\ell^2 \equiv 1 \pmod{p}$.

Proof. This follows from the proof of Theorem 4.4 along with Proposition 3.16 and Proposition 3.20. \square

We end with results in the introduction.

Remark 4.8. (1). Theorem 1.1 and Theorem 1.4 follow from Theorem 4.4.

(2). Theorem 1.5 follows from Corollary 4.6 and Corollary 4.7.

REFERENCES

- [1] O. Beckwith, *Indivisibility of class numbers of imaginary quadratic fields*, Preprint, arXiv:1612.04443.
- [2] M. Bertolini and H. Darmon, *Iwasawa's main conjecture for elliptic curves over anticyclotomic \mathbb{Z}_p -extensions*, Ann. of Math. (2) 162 (2005), no. 1, 1–64.
- [3] M. Bertolini, H. Darmon and K. Prasanna, *Generalised Heegner cycles and p -adic Rankin L -series*, Duke Math. J. 162 (2013), no. 6, 1033–1148.
- [4] M. Bertolini, H. Darmon and K. Prasanna, *p -adic Rankin L -series and rational points on CM elliptic curves*, Pacific Journal of Mathematics, Vol. 260, No. 2, 2012. 261–303.
- [5] M. Bhargava, B. Kane, H. Lenstra, B. Poonen and E. Rains, *Modeling the distribution of ranks, Selmer groups, and Shafarevich-Tate groups of elliptic curves*, Camb. J. Math. 3 (2015), no. 3, 275–321.
- [6] E. Brooks, *Shimura curves and special values of p -adic L -functions*, Int. Math. Res. Not. IMRN 2015, no. 12, 4177–4241.
- [7] J. Bruinier, *Nonvanishing modulo l of Fourier coefficients of half-integral weight modular forms*, Duke Math. J. 98 (1999), no. 3, 595–611.
- [8] D. Burns, D. Macias Castillo and C. Wuthrich, *On Mordell-Weil groups and congruences between derivatives of twisted Hasse-Weil L -functions*, to appear in J. reine u. angew. Math.
- [9] A. Burungale, *On the non-triviality of the p -adic Abel-Jacobi image of generalised Heegner cycles modulo p , II: Shimura curves*, J. Inst. Math. Jussieu 16 (2017), no. 1, 189–222.
- [10] A. Burungale and H. Hida, *André-Oort conjecture and non-vanishing of central L -values over Hilbert class fields*, Forum of Mathematics, Sigma 4 (2016), e8, 26 pp.
- [11] A. Burungale and H.-S. Sun, *Quantitative non-vanishing of Dirichlet L -values modulo p* , preprint, 2017.
- [12] A. Burungale and Y. Tian, *Horizontal non-vanishing of Heegner points and toric periods*, preprint, 2017.
- [13] L. Cai, J. Shu, Y. Tian, *Explicit Gross-Zagier and Waldspurger formulae*, Algebra and Number Theory 8 (2014), no. 10, 2523–2572.
- [14] M. Chida and M.-L. Hsieh, *Special values of anticyclotomic L -functions for modular forms*, to appear in J. reine angew. Math.
- [15] C. Cornut, *Mazur's conjecture on higher Heegner points*, Invent. Math. 148 (2002), no. 3, 495–523.
- [16] C. Delaunay, *Heuristics on class groups and on Tate-Shafarevich groups: the magic of the Cohen-Lenstra heuristics, Ranks of elliptic curves and random matrix theory*, London Math. Soc. Lecture Note Ser., vol. 341, Cambridge Univ. Press, Cambridge, 2007, pp. 323–340.
- [17] P. Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 2, 247–289, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979.
- [18] B. Ferrero and L. Washington, *The Iwasawa invariant μ_p vanishes for abelian number fields*, Ann. of Math. (2) 109 (1979) 377–395.
- [19] B. Gross and D. Zagier, *Heegner points and derivatives of L -series*, Invent. Math. 84 (1986), no. 2, 225–320.
- [20] B. Gross, *Kolyvagin's work on modular elliptic curves, L -functions and arithmetic* (Durham, 1989), 235–256, London Math. Soc. Lecture Note Ser., 153, Cambridge Univ. Press, Cambridge, 1991.
- [21] M. Harris, *Arithmetic vector bundles and automorphic forms on Shimura varieties. I*, Invent. Math. 82 (1985), no. 1, 151–189.
- [22] H. Hida, *Hilbert modular forms and Iwasawa theory*, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006. xiv+402 pp.
- [23] H. Hida, *Elliptic Curves and Arithmetic Invariants*, Springer Monographs in Mathematics, Springer, New York, 2013. xviii+449 pp.
- [24] H. Iwaniec, *Fourier coefficients of modular forms of half-integral weight*, Invent. Math. 87 (1987), no. 2, 385–401.
- [25] C. Khare and I. Kiming, *Mod pq Galois representations and Serre's conjecture*, J. Number Theory 98 (2003), no. 2, 329–347.
- [26] C. Khare, *On isomorphisms between deformation rings and Hecke rings*, Invent. Math. 154 (2003), no. 1, 199–222.
- [27] C. Khare and J.-P. Wintenberger, *Serre's modularity conjecture. I.*, Invent. Math. 178 (2009), no. 3, 485–504.
- [28] V. Kolyvagin, *Euler systems*, The Grothendieck Festschrift, Vol. II, 435–483, Progr. Math., 87, Birkhäuser Boston, Boston, MA, 1990.
- [29] Y. Liu, S. Zhang and W. Zhang, *On p -adic Waldspurger formula*, preprint, 2014, available at <http://www.math.mit.edu/liuyf/>.
- [30] B. Mazur, K. Rubin and A. Silverberg, *Twisting commutative algebraic groups*, J. Algebra 314 (2007), no. 1, 419–438.
- [31] B. Mazur and A. Wiles, *Class fields of abelian extensions of \mathbb{Q}* , Invent. Math. 76 (1984), no. 2, 179–330.
- [32] P. Michel, *The subconvexity problem for Rankin-Selberg L -functions and equidistribution of Heegner points*, Ann. of Math. (2) 160 (2004), no. 1, 185–236.

- [33] P. Michel and A. Venkatesh, *Heegner points and non-vanishing of Rankin/Selberg L -functions*, Analytic number theory, 169–183, Clay Math. Proc., 7, Amer. Math. Soc., Providence, RI, 2007.
- [34] P. Michel and A. Venkatesh, *The subconvexity problem for GL_2* , Publ. Math. Inst. Hautes des Sci. No. 111 (2010), 171–271.
- [35] F. Momose, *On the l -adic representations attached to modular forms*, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 28 (1981), no. 1, 89–109.
- [36] J. Nekovář, *The Euler system method for CM points on Shimura curves*, L-functions and Galois representations, 471–547, London Math. Soc. Lecture Note Ser., 320, Cambridge Univ. Press, Cambridge, 2007.
- [37] J. Nekovář, *Level raising and anticyclotomic Selmer groups for Hilbert modular forms of weight two*, Canad. J. Math. 64 (2012), no. 3, 588–668.
- [38] R. Pollack and T. Weston, *On anticyclotomic μ -invariants of modular forms*, Compos. Math. 147 (2011), no. 5, 1353–1381.
- [39] K. Ribet, *Twists of modular forms and endomorphisms of abelian varieties*, Math. Ann. 253 (1980), no. 1, 43–62.
- [40] G. Robin, *Irrégularités dans la distribution des nombres premiers dans les progressions arithmétiques*, Ann. Fac. Sci. Toulouse Math. (5) 8 (1986/87), no. 2, 159–173.
- [41] H. Saito, *On Tunnell’s formula for characters of $GL(2)$* , Compositio Math. 85 (1993), no. 1, 99–108.
- [42] W. Sinnott, *On a theorem of L. Washington*, Journées arithmétiques (Besançon 1985), Asterisque 147–148, Societe Mathematique de France, Paris (1987), 209–224.
- [43] N. Templier, *Sur le rang des courbes elliptiques sur les corps de classes de Hilbert*, Compos. Math. 147 (2011), no. 4, 1087–1104.
- [44] J. Tunnell, *On the local Langlands conjecture for $GL(2)$* , Invent. Math. 46 (1978), 179–200.
- [45] J. Tunnel, *Local ϵ -factors and characters of $GL(2)$* , Amer. J. Math. 105 (1983), no. 6, 1277–1307.
- [46] V. Vatsal, *Uniform distribution of Heegner points*, Invent. Math. 148, 1–48 (2002).
- [47] V. Vatsal, *Special values of L -functions modulo p* , International Congress of Mathematicians. Vol. II, 501–514, Eur. Math. Soc., Zürich, 2006.
- [48] J.-L. Waldspurger, *Sur les valeurs de certaines fonctions L automorphes en leur centre de symétrie*, Compositio Math. 54 (1985), no. 2, 173–242.
- [49] L. Washington, *The non- p -part of the class number in a cyclotomic \mathbb{Z}_p -extension*, Invent. Math. 49 (1978), 87–97.
- [50] A. Wiles, *The Iwasawa conjecture for totally real fields*, Ann. of Math. (2) 131 (1990), no. 3, 493–540.
- [51] A. Wiles, *On class groups of imaginary quadratic fields*, J. Lond. Math. Soc. (2) 92 (2015), no. 2, 411–426.
- [52] X. Yuan, S.-W. Zhang and W. Zhang, *The Gross–Zagier Formula on Shimura Curves*, Annals of Mathematics Studies, Princeton University Press, 2012.
- [53] S.-W. Zhang, *Equidistribution of CM-points on quaternion Shimura varieties*, Int. Math. Res. Not. 2005, no. 59, 3657–3689.
- [54] W. Zhang, *Selmer groups and the indivisibility of Heegner points*, Cambridge Journal of Math., Vol. 2 (2014), No. 2, 191–253.

ASHAY A. BURUNGAL: INSTITUTE FOR ADVANCED STUDY, EINSTEIN DRIVE, PRINCETON NJ 08540
E-mail address: ashayburungale@gmail.com

HARUZO HIDA: DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095-1555, U.S.A.
E-mail address: hida@math.ucla.edu

YE TIAN: ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, MORNINGSIDE CENTER OF MATHEMATICS, CHINESE ACADEMY OF SCIENCES, BEIJING 100190 AND
 SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 10049
E-mail address: ytian@math.ac.cn