

A totally bounded Uniformity on coarse metric Spaces

Elisa Hartmann

March 2, 2022

Abstract

This paper presents a new version of boundary on coarse spaces. The space of ends functor maps coarse metric spaces to uniform topological spaces and coarse maps to uniformly continuous maps.

Contents

0	Introduction	1
0.1	Background and related Theories	2
0.2	Main Contributions	3
0.3	Outline	4
1	Metric Spaces	4
2	Totally Bounded Uniformity	6
3	Definition	10
4	Properties	13
5	Side Notes	18
6	Remarks	20

0 Introduction

Coarse Geometry of metric spaces studies the large scale properties of a metric space. Meanwhile uniformity of metric spaces is about small scale properties.

Our purpose is to pursue a new version of duality between the coarse geometry of metric spaces and uniform spaces. We present a notion of boundary on coarse metric spaces which is a totally bounded separating uniform space. The methods are very basic and do not require any deep theory.

Note that the topology of metric spaces is well understood and there are a number of topological tools that can be applied on coarse metric spaces which have not been used before. The new discovery may lead to new insight on the topic of coarse geometry.

0.1 Background and related Theories

There are quite number of notions for a boundary on a metric space. In this chapter we are going to discuss properties for three of them. The first paragraph is devoted to the *Higson corona*, in the second paragraph the *space of ends* is presented and in the last paragraph we study the *Gromov boundary*.

First we present the Higson corona. If X is a proper metric space the *Higson corona* νX is the boundary of the Higson compactification hX of X which is a compact topological space that contains the underlying topological space of X as a dense open subset.

If $C(X)$ denotes the bounded continuous functions on X then the so called Higson functions are a subset of $C(X)$. This subset determines a compactification which is called the Higson compactification. By a comment on [1, p. 31] the Higson corona can be defined for any coarse space. The same does not work for the Higson compactification¹. The [1, Proposition 2.41] implies that the Higson corona is a covariant functor that sends coarse maps modulo closeness to continuous maps. Thus ν is a functor:

$$\nu : \text{Coarse} \rightarrow \text{Top}$$

The topology of νX has been studied in [2]. It was shown in [2, Theorem 1] that for every σ -compact subset $A \subseteq \nu X$ the closure \bar{A} of A in νX is equivalent to the Stone-Ćech compactification of A . The topology of νX is quite complicated, especially if X is a metric space. It has been noted in [1, Exercise 2.49] that the topology of νX for X an unbounded proper metric space is never second countable. In [3, Theorem 1.1] and [4, Theorem 7.2] it was shown that if the asymptotic dimension $\text{asdim}(X)$ of X is finite then

$$\text{asdim}(X) = \dim(\nu X)$$

where the right side denotes the topological dimension of νX . Note that one direction of the proof uses the notion of coarse covers².

Now we present the *space of ends*. If Y is a locally connected, connected and locally compact Hausdorff space then the space of ends ΩY of Y is the boundary of the Freudenthal compactification εY . It is totally disconnected and every other compactification of Y that is totally disconnected factors uniquely through εY by [5, Theorem 1]. The points of ΩY are called *end-points* or *ends*.

Now [5, Theorem 5] shows that if Y is a connected locally finite countable CW-complex every endpoint of Y can be represented by a proper map

$$a : \mathbb{R}_+ \rightarrow Y.$$

Two proper maps $a_1, a_2 : \mathbb{R}_+ \rightarrow Y$ represent the same endpoint if they are connected by a proper homotopy. Denote by \mathbf{pTop} the category of topological spaces and proper continuous maps. Then the association $\Omega \cdot$ is a functor:

$$\Omega : \mathbf{pTop} \rightarrow \text{Top}$$

If Y is a locally compact Hausdorff space then ΩY can be constructed using a proximity relation which is a relation on the subsets of Y . See [6] for that one.

This section studies the Gromov boundary. If X is a proper Gromov hyperbolic metric space then the *Gromov boundary* ∂X consists of equivalence classes of *sequences that converge to infinity* in X . The topology on ∂X is generated by a basis of open neighborhoods. Loosely

¹for which the topology of X needs to be locally compact which is given if the metric is proper.

²but under a different name.

speaking two points on the boundary are close if the sequences that represent them stay close for a long time.

By [7, Proposition 2.14] the topological spaces ∂X and $\partial X \cup X$ are compact and by [7, Theorem 2.1] the topology on ∂X is metrizable. If $f : X \rightarrow Y$ is a *quasi-isometry* between proper Gromov hyperbolic groups then it extends to a homeomorphism

$$\partial f : \partial X \rightarrow \partial Y$$

by [7, Proposition 2.20]. In [8] is studied a notion of morphisms for which the Gromov boundary is a functor: If $f : X \rightarrow Y$ is a *visual function* between proper Gromov hyperbolic metric spaces then there is an induced map

$$\partial f : \partial X \rightarrow \partial Y$$

which is continuous by [8, Theorem 2.8].

Now is there a notion of boundary on metric space which is both a functor on coarse spaces and coarse maps and has nice properties such as being Hausdorff and locally compact. As it turns out there is one such functor which is going to be designed in the course of this study.

0.2 Main Contributions

In the course of this study we will define a functor that associates to every coarse metric space a space at infinity which is a topological space.

Based on the observation that twisted coarse cohomology with $\mathbb{Z}/2\mathbb{Z}$ -coefficients of \mathbb{Z}^n is the same as singular cohomology of S^{n-1} with $\mathbb{Z}/2\mathbb{Z}$ coefficients we considered notions of boundary which reflect that observation.

In Definition 8 we define a relation on subsets of a metric space. As it turns out this relation is almost but not quite a proximity relation as noted in Remark 11. The proof of Proposition 10 uses that X is a metric space, it does not work for general coarse spaces. That is why we restrict our study to metric spaces.

Note that while constructing the functor we presuppose which kind of spaces we want to distinguish. Indeed there is a certain class of metric spaces for which the local structure looks boring. The functor that we are going to define, the space of ends functor, is well suited for metric spaces that are coarsely proper coarsely geodesic. That class includes all Riemannian manifolds and finitely generated groups.

While the topology of the space at infinity is immediately defined using coarse covers there are two choices of points which are both solid: If X is a metric space

(A): the endpoints of X are images of coarse maps $\mathbb{Z}_+ \rightarrow X$ modulo finite Hausdorff distance
or

(B): the points at infinity are subsets of X modulo finite Hausdorff distance.

Note that choice B has been implemented in [9]. The space at infinity with choice A contains strictly less points than choice B. The [10, Proposition 93] guarantees that for choice A there exists at least one endpoint if the space X is coarsely proper coarsely geodesic. The proof of [10, Proposition 93] is similar to the one of Königs Lemma in graph theory.

The space at infinity functor with choice B reflects isomorphisms by [11, Proposition 2.18] and the space at infinity functor with choice A is representable.

In the course of this article and in Definition 21 we use choice A, endpoints are images of coarse maps $\mathbb{Z}_+ \rightarrow X$. Then we define the topology of the space of ends, $O(X)$, via surroundings of the diagonal in Definition 23. The uniformity on $O(X)$ is generated by a basis $(Du)_u$ of

entourages over coarse covers \mathcal{U} of X . If $f : X \rightarrow Y$ is a coarse map then it induces a uniformly continuous map $O(f) : O(X) \rightarrow O(Y)$ between spaces of ends. That way the space of ends $O(\cdot)$ is a functor, we obtain the following result:

Theorem A. *If $mCoarse$ denotes the category of metric spaces and coarse maps modulo closeness and Top the category of topological spaces and continuous maps then O is a functor*

$$O : mCoarse \rightarrow Top.$$

If $Uniform$ denotes the category of uniform spaces and uniformly continuous maps then O is a functor

$$O : mCoarse \rightarrow Uniform.$$

It was nontrivial to show that a subspace in the domain category gives rise to a subspace in the image category. Proposition 33 shows if $Z \subseteq Y$ is a subspace then the inclusion $i : Z \rightarrow Y$ induces a uniform embedding $O(i) : O(Z) \rightarrow O(Y)$.

The functor $O(\cdot)$ preserves coproducts by Lemma 39. The uniformity on $O(X)$ is totally bounded by Lemma 41 and separating by Proposition 40.

Theorem B. *If X is a metric space then $O(X)$ is totally bounded and separating.*

We still lack a good study including the most basic properties of the new space of ends functor like compact and metrizable probably because the proofs are more difficult.

0.3 Outline

The outline of this paper is as follows:

- Chapter 1 introduces metric spaces and the coarse geometry of them. We recall basic notation used in this study.
- The Chapter 2 creates the basic tool sets which are going to be used in this study.
- Finally Chapter 3 is about the space of ends functor. We give the definition and prove that the space of ends is a functor.
- Chapter 4 studies a few basic properties. We prove that the space of ends is totally bounded and separated.
- The study closes with Chapter 5 which sets the space of ends functor in context with other notions.

1 Metric Spaces

Definition 1. Let (X, d) be a metric space.

- Then the *bounded coarse structure associated to d* on X consists of those subsets $E \subseteq X^2$ for which

$$\sup_{(x,y) \in E} d(x,y) < \infty.$$

We call an element of the coarse structure *entourage*.

- The *bounded cocoarse structure associated to d* on X consists of those subsets $C \subseteq X^2$ such that every sequence $(x_i, y_i)_i$ in C is either bounded (which means both of the sequences $(x_i)_i$ and $(y_i)_i$ are bounded) or the set $d(x_i, y_i)_i$ is not bounded. for $i \rightarrow \infty$. We call an element of the cocoarse structure coentourage.
- In what follows we assume the metric d to be finite for every $(x, y) \in X^2$.

Remark 2. Note that there is a more general notion of coarse spaces. By [1, Theorem 2.55] a coarse structure on a coarse space X is the bounded coarse structure associated to some metric d on X if and only if the coarse structure has a countable base.

Remark 3. If X is a metric space a subset $B \subseteq X$ is *bounded* if the set B^2 is an entourage in X .

Remark 4. Note the following duality:

- A subset $F \subseteq X^2$ is an entourage if and only if for every coentourage $C \subseteq X^2$ there is a bounded set $A \subseteq X$ such that

$$F \cap C \subseteq A^2.$$

- A subset $D \subseteq X^2$ is a coentourage if and only if for every entourage $E \subseteq X^2$ there is a bounded set B such that

$$E \cap D \subseteq B^2.$$

Definition 5. A map $f : X \rightarrow Y$ between metric spaces is called *coarse* if

- $E \subseteq X^2$ being an entourage implies that $f^{\times 2}E$ is an entourage (*coarsely uniform*);
- and if $A \subseteq Y$ is bounded then $f^{-1}(A)$ is bounded (*coarsely proper*).

Or equivalently

- $B \subseteq X$ being bounded implies that $f(B)$ is bounded;
- and if $D \subseteq Y^2$ is a coentourage then $(f^{\times 2})^{-1}(D)$ is a coentourage.

Two maps $f, g : X \rightarrow Y$ between metric spaces are called *close* if

$$f \times g(\Delta_X)$$

is an entourage in Y . Here Δ_X denotes the diagonal in X .

Notation 6. A map $f : X \rightarrow Y$ between metric spaces is called

- *coarsely surjective* if there is an entourage $E \subseteq Y^2$ such that

$$E[\text{im } f] = Y$$

- *coarsely injective* if
 1. for every entourage $F \subseteq Y^2$ the set $(f^{\times 2})^{-1}(F)$ is an entourage in X .
 2. or equivalently if for every coentourage $C \subseteq X^2$ the set $f^{\times 2}(C)$ is a coentourage in Y .
- two subsets $A, B \subseteq X$ are called *coarsely disjoint* if $A \times B$ is a coentourage.

Remark 7. We study metric spaces up to coarse equivalence. A coarse map $f : X \rightarrow Y$ is a *coarse equivalence* if

- There is a coarse map $g : Y \rightarrow X$ such that $f \circ g$ is close to id_Y and $g \circ f$ is close to id_X .
- or equivalently if f is both coarsely injective and coarsely surjective.

2 Totally Bounded Uniformity

Definition 8. (close relation) Let X be a metric space. Two subsets $A, B \subseteq X$ are called *close* if they are not coarsely disjoint. We write

$$A \wedge B.$$

Then \wedge is a relation on the subsets of X .

Lemma 9. *In every metric space X :*

1. *if B is bounded, $B \not\wedge A$ for every $A \subseteq X$*
2. *$U \wedge V$ implies $V \wedge U$*
3. *$U \wedge (V \cup W)$ if and only if $U \wedge V$ or $U \wedge W$*

Proof. 1. easy.

2. easy.

3. easy. □

Proposition 10. *Let X be a metric space. Then for every subspaces $A, B \subseteq X$ with $A \not\wedge B$ there are subsets $C, D \subseteq X$ such that $C \cap D = \emptyset$ and $A \not\wedge (X \setminus C)$, $B \not\wedge (X \setminus D)$.*

Proof. Note this is the same as [11, Proposition 4.5] where the same statement was proven in a similar fashion. Let $E_1 \subseteq E_2 \subseteq \dots$ be a symmetric basis for the coarse structure of X . Then for every $x \in A^c \cap B^c$ there is a least number $n_1(x)$ such that $x \in E_{n_1(x)}[A]$ and a least number $n_2(x)$ such that $x \in E_{n_2(x)}[B]$. Define:

$$V_1 = \{x \in A^c \cap B^c : n_1(x) \leq n_2(x)\}$$

and

$$V_2 = A^c \cap B^c \setminus V_1.$$

Now for every n

$$E_n[V_1] \cap B \subseteq E_{2n}[A] \cap B$$

because for every $x \in V_1$, if $x \in E_n[B]$ then $x \in E_n[A]$. Now define

$$C = A \cup V_1$$

and

$$D = B \cup V_2. \quad \square$$

Remark 11. Compare \wedge with the notion of proximity relation [12, chapter 40, pp. 266]. By Lemma 9 and Proposition 10 the close relation satisfies [12, P-1), P-3)-P-5) of Definition 40.1] but not P-2).

Remark 12. If $f : X \rightarrow Y$ is a coarse map then whenever $A \wedge B$ in X then $f(A) \wedge f(B)$ in Y .

We recall [10, Definition 45]:

Definition 13. (coarse cover) If X is a metric space and $U \subseteq X$ a subset a finite family of subsets $U_1, \dots, U_n \subseteq U$ is said to *coarsely cover* U if

$$U^2 \cap \left(\bigcup_i U_i^2 \right)^c$$

is a coentourage in X .

Remark 14. Note that coarse covers determine a Grothendieck topology on X . If $f : X \rightarrow Y$ is a coarse map between metric spaces and $(V_i)_i$ a coarse cover of $V \subseteq Y$ then $(f^{-1}(V_i))_i$ is a coarse cover of $f^{-1}(V) \subseteq X$.

Lemma 15. *Let X be a metric space. A finite family $\mathcal{U} = \{U_\alpha : \alpha \in A\}$ is a coarse cover if and only if there is a finite cover $\mathcal{V} = \{V_\alpha : \alpha \in A\}$ of X as a set such that $V_\alpha \not\ll U_\alpha^c$ for every α .*

Proof. Suppose \mathcal{U} is a coarse cover of X . We proceed by induction on the index of \mathcal{U} :

- $n = 1$: a subset U coarsely covers X if and only if U^c is bounded if and only if $U^c \not\ll X$.
- two subsets U_1, U_2 coarsely cover X if and only if $U_1^c \not\ll U_2^c$. Now by Proposition 10 there are $C, D \subseteq X$ with $C^c \cup D^c = X$ and $C^c \not\ll U_1^c$ and $D^c \not\ll U_2^c$. Define $V_1 = C^c, V_2 = D^c$.
- $n + 1 \rightarrow n + 2$: Subsets U, V, U_1, \dots, U_n coarsely cover X if and only if U, V coarsely cover $U \cup V$ and $U \cup V, U_1, \dots, U_n$ coarsely cover X at the same time.

Suppose U, V, U_1, \dots, U_n coarsely cover X . By induction hypothesis there is a cover of sets V'_1, V'_2 of $U \cup V$ such that $V'_1 \not\ll U^c \cap V$ and $V'_2 \not\ll V^c \cap U$ and there is a cover of sets W, V_1, \dots, V_n such that $W \not\ll (U \cup V)^c$ and $V_i \not\ll U_i^c$ for every i . Then $V'_1 \cap W \not\ll U^c, V'_2 \cap W \not\ll V^c$. Now

$$B := (U \cup V)^c \cap W$$

is bounded. Then

$$V'_1 \cap W, V'_2 \cap W, V_1, \dots, V_n \cup B$$

is a finite cover of X with the desired properties.

Suppose $(V_\alpha)_\alpha$ cover X as sets and $V_\alpha \not\ll U_\alpha^c$ for every α . Let $E \subseteq X^2$ be an entourage. Then $E[U_\alpha^c] \cap V_\alpha$ is bounded for every α . Then

$$\begin{aligned} \bigcap_\alpha E[U_\alpha^c] &= \bigcap_\alpha E[U_\alpha^c] \cap \left(\bigcup_\alpha V_\alpha \right) \\ &= \bigcup_\alpha (V_\alpha \cap \bigcap_\beta E[U_\beta^c]) \end{aligned}$$

is bounded. By [10, Proposition 54] we can conclude that $(U_\alpha)_\alpha$ is a coarse cover of X . \square

Remark 16. The [12, Theorem 40.15] states that every proximity relation on a set is induced by some totally bounded uniformity on it. Note that a coarse cover on a metric space X does not precisely need to cover X as a set. Except for that the collection of all coarse covers of a metric space satisfies [12, a), b) of Theorem 36.2]. We can compare coarse covers of X with a base for a totally bounded uniformity on X : the collection of all sets $\bigcup_i U_i^2$ for $(U_i)_i$ a coarse cover satisfies [12, b)-e) of Definition 35.2] but not a). Note that by [12, Definition 39.7] a diagonal uniformity is totally bounded if it has a base consisting of finite covers.

Lemma 17. (separation cover) If U_1, U_2 coarsely cover a metric space X (or equivalently if U_1^c, U_2^c are coarsely disjoint) then there exists a coarse cover V_1, V_2 of X such that $V_1 \not\ll U_1^c$ and $V_2 \not\ll U_2^c$.

Proof. By Proposition 10 there are subsets $C, D \subseteq X$ such that $C \cap D = \emptyset$, $U_1^c \not\ll C^c$ and $U_2^c \not\ll D^c$. Thus U_1, C is a coarse cover of X such that $C \not\ll U_2^c$.

By Proposition 10 there are subsets $A, B \subseteq X$ such that $A \cap B = \emptyset$, $A^c \not\ll U_1^c$ and $B^c \not\ll C^c$. Then B, C are a coarse cover of X such that $B \not\ll U_1^c$.

Then $V_1 = B$ and $V_2 = C$ have the desired properties. \square

Notation 18. (coarse star refinement) Let $\mathcal{U} = (U_i)_{i \in I}$ be a coarse cover of a metric space X .

1. If $S \subseteq X$ is a subset then

$$\text{cst}(S, \mathcal{U}) = \bigcup \{U_i : S \wedge U_i\}$$

is called the *coarse star* of S .

2. Let $\mathcal{V} = (V_j)_{j \in J}$ be a coarse cover of X . Then \mathcal{V} is a *coarse barycentric refinement* of \mathcal{U} if for every $j_1, \dots, j_k \in J$ and entourage $E \subseteq X^2$ with

$$\bigcap_k E[V_{j_k}]$$

not bounded there is some $i \in I$ and an entourage $F \subseteq X^2$ such that

$$\bigcup_k V_{j_k} \subseteq F[U_i].$$

3. A coarse cover $\mathcal{V} = (V_j)_{j \in J}$ of X is called a *coarse star refinement* of \mathcal{U} if for every $j \in J$ there is some $i \in I$ and entourage $E \subseteq X^2$ such that

$$\text{cst}(V_j, \mathcal{V}) \subseteq E[U_i].$$

Lemma 19. If $\mathcal{V} = (V_i)_i$ is a coarse star refinement of a coarse cover $\mathcal{U} = (U_i)_i$ of a metric space X then

- if $S \subseteq X$ is a subset then there is an entourage $E \subseteq X^2$ such that

$$\text{cst}(\text{cst}(S, \mathcal{V}), \mathcal{V}) \subseteq E[\text{cst}(S, \mathcal{U})];$$

- if $f : X \rightarrow Y$ is a coarse map between metric spaces, $(U_i)_i$ a coarse cover of Y and $S \subseteq X$ a subset then

$$f(\text{cst}(S, f^{-1}(\mathcal{U}))) \subseteq \text{cst}(f(S), \mathcal{U}).$$

Proof. • Suppose $E \subseteq X^2$ is an entourage such that for every V_j there is an U_i such that $\text{cst}(V_j, \mathcal{V}) \subseteq E[U_i]$. Note that $S \wedge V_j$ implies $S \wedge U_i$ in that case. Then

$$\begin{aligned} \text{cst}(\text{cst}(S, \mathcal{V}), \mathcal{V}) &= \text{cst}(\bigcup \{V_i : V_i \wedge S\}, \mathcal{V}) \\ &= \bigcup_{V_i \wedge S} \text{cst}(V_i, \mathcal{V}) \\ &\subseteq \bigcup_{S \wedge U_j} E[U_j] \\ &= E[\text{cst}(S, \mathcal{U})]. \end{aligned}$$

•

$$\begin{aligned}
f(\text{cst}(S, f^{-1}(\mathcal{U}))) &= \bigcup \{f \circ f^{-1}(U_i) : S \wedge f^{-1}(U_i)\} \\
&\subseteq \bigcup \{f \circ f^{-1}(U_i) : f(S) \wedge f \circ f^{-1}(U_i)\} \\
&\subseteq \bigcup \{U_i : f(S) \wedge U_i\} \\
&= \text{cst}(f(S), \mathcal{U}).
\end{aligned}$$

□

Lemma 20. *If \mathcal{U} is a coarse cover of a metric space X then there exists a coarse cover \mathcal{V} of X that coarsely star refines \mathcal{U} .*

Proof. There are three steps:

If $\mathcal{V} = (V_j)_j$ is a coarse barycentric refinement of \mathcal{U} and $\mathcal{W} = (W_k)_k$ is a coarse barycentric refinement of \mathcal{V} then \mathcal{W} is a coarse star refinement of \mathcal{U} :

fix W_k and denote $J = \{j : W_k \wedge W_j\}$.

Then for every $j \in J$ there is some V_j and entourage $E_j \subseteq X^2$ such that $W_k \cup W_j \subseteq E_j[V_j]$.

Define $E = \bigcup_j E_j$. Then $\bigcap_j E[V_j] \supseteq W_k$. Thus there is some U_i and entourage $F \subseteq X^2$ such that $\bigcup_j V_j \subseteq F[U_i]$.

For every $j \in J$:

$$\begin{aligned}
W_j &\subseteq E[V_j] \\
&\subseteq E \circ F[U_i].
\end{aligned}$$

Thus $\text{cst}(W_k, \mathcal{W}) \subseteq E \circ F[U_i]$.

We show there is a coarse barycentric refinement $\mathcal{V} = (V_i)_i$ of \mathcal{U} : First we show if U_1, U_2 is a coarse cover of X then there is a coarse barycentric refinement V_1, V_2, V_3 of U_1, U_2 :

By Lemma 17 there is a coarse cover W_1, W_2 of X such that $W_1 \not\ll U_1^c$ and $W_2 \not\ll U_2^c$.

Then W_1^c, U_1 and W_2^c, U_2 are coarse covers of X .

By Proposition 10 there are $C, D \subseteq X$ such that $C \cap D = \emptyset$, $D^c \not\ll U_2^c$, $C^c \not\ll W_2$.

Also there are $A, B \subseteq X$ such that $A \cap B = \emptyset$, $A^c \not\ll U_1^c$, $B^c \not\ll W_1$.

Then

$$V_1 = W_1, V_2 = C \cap B, V_3 = W_2$$

has the desired properties:

$(V_i)_i$ is a coarse cover:

Note that by $B^c \not\ll W_1$ and $W_1^c \not\ll W_2^c$ the sets W_2, B are a coarse cover of X .

Note that by $C^c \not\ll W_2$ and $W_1^c \not\ll W_2^c$ the sets W_1, C coarsely cover X .

Note that $(W_1 \cap W_2) \not\ll (C^c \cup B^c)$. Then, combining items i,ii, we get that

$$W_1, W_2, B \cap C$$

is a coarse cover as required.

$(V_i)_i$ is a coarse barycentric refinement of U_1, U_2 :

There is some entourage $E \subseteq X^2$ such that $V_1 \cup V_2 \subseteq E[U_1]$: For W_1 we use that $W_1 \not\ll U_1^c$. For $C \cap B$ we use that $A^c \not\ll U_1^c$ and $B \subseteq A^c$.

There is an entourage $E \subseteq X^2$ such that $V_2 \cup V_3 \subseteq E[U_2]$: For W_2 we use that $W_2 \not\ll U_2^c$. For $C \cap B$ we use that $D^c \not\ll U_2^c$ and $C \subseteq D^c$.

$V_1 \not\ll V_3$: We use $W_1^c \not\ll W_2^c$.

Now we show the general case: Suppose $U_i \subseteq X$ are subsets such that $\mathcal{U} = (U_i)_i$. We show there is a coarse barycentric refinement \mathcal{V} of \mathcal{U} .

For every i the sets $U_i, \bigcup_{j \neq i} U_j$ coarsely cover X . By Lemma 17 there are subsets W_1^i, W_2^i that coarsely cover X such that $W_1^i \not\ll U_i^c$ and $W_2^i \not\ll (\bigcup_{j \neq i} U_j)^c$.

Then there is a coarse barycentric refinement V_1^i, V_2^i, V_3^i of W_1^i, W_2^i for every i .

Then we define

$$\mathcal{V} := (\bigcap_i V_{\sigma(i)}^i)_\sigma,$$

here $\sigma(i) \in \{1, 2, 3\}$ is all possible permutations.

We show \mathcal{V} is a coarse cover of X that is a coarse barycentric refinement of \mathcal{U} :

\mathcal{V} is a coarse cover: by design.

\mathcal{V} is a coarse barycentric refinement of \mathcal{U} : Suppose there is an entourage $E \subseteq X^2$ and a subindex $(\sigma_k)_k$ such that $\bigcap_{\sigma_k} E[\bigcap_{\sigma_k} V_{\sigma_k(i)}^i]$ is not bounded. Then

$$\bigcap_{i, \sigma_k} E[V_{\sigma_k(i)}^i]$$

is not bounded. Then there is an entourage $F \subseteq X^2$ such that for every i :

$$\bigcup_{\sigma_k} V_{\sigma_k(i)}^i \subseteq F[W_{l_i}^i]$$

where l_i is one of 1, 2. Then

$$\begin{aligned} \bigcup_{\sigma_k} \bigcap_i V_{\sigma_k(i)}^i &\subseteq \bigcap_i \bigcup_{\sigma_k} V_{\sigma_k(i)}^i \\ &\subseteq \bigcap_i F[W_{l_i}^i] \end{aligned}$$

if $l_i = 1$ for one i then we are done. Otherwise

$$\bigcup_{\sigma_k} \bigcap_i V_{\sigma_k(i)}^i \subseteq \bigcap_i F[W_2^i]$$

and $F[W_2^i] \not\ll (\bigcup_{j \neq i} U_j)^c$ implies

$$\bigcap_i F[W_2^i] \not\ll \bigcup_i \bigcup_{j \neq i} U_j^c$$

which implies that $\bigcap_i F[W_2^i]$ is bounded, a contradiction. \square

3 Definition

We introduce the space of ends of a coarse space which is a functor O from the category of coarse metric spaces to the category of uniform spaces.

Definition 21. (endpoint) Let X be a metric space,

- two coarse maps $\phi, \psi : \mathbb{Z}_+ \rightarrow X$ are said to represent the same *endpoint in X* if there is an entourage $E \subseteq X^2$ such that

$$E[\psi(\mathbb{Z}_+)] = \phi(\mathbb{Z}_+).$$

- if $\mathcal{U} = (U_i)_i$ is a coarse cover of X and p, q are two endpoints in X which are represented by $\phi, \psi : \mathbb{Z}_+ \rightarrow X$. Then q is said to be in a \mathcal{U} -neighborhood of p , denoted $q \in \mathcal{U}[p]$, if there is an entourage $E \subseteq X^2$ such that

$$E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})] \supseteq \psi(\mathbb{Z}_+)$$

and

$$E[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{U})] \supseteq \phi(\mathbb{Z}_+).$$

Lemma 22. *If $\mathcal{V} \leq \mathcal{U}$ is a refinement of a coarse cover of a metric space X then for every two endpoints p, q of X the relation $q \in \mathcal{V}[p]$ implies the relation $q \in \mathcal{U}[p]$.*

Proof. Suppose $\mathcal{V} = (V_i)_i$ and $\mathcal{U} = (U_i)_i$. If p, q are represented by $\phi, \psi : \mathbb{Z}_+ \rightarrow X$ then

$$\begin{aligned} \text{cst}(\phi(\mathbb{Z}_+), \mathcal{V}) &= \bigcup \{V_i : \phi(\mathbb{Z}_+) \wedge V_i\} \\ &\subseteq \bigcup \{U_i : \phi(\mathbb{Z}_+) \wedge U_i\} \\ &= \text{cst}(\phi(\mathbb{Z}_+), \mathcal{U}) \end{aligned}$$

in the same way $\text{cst}(\psi(\mathbb{Z}_+), \mathcal{V}) \subseteq \text{cst}(\psi(\mathbb{Z}_+), \mathcal{U})$. Then if $q \in \mathcal{V}[p]$ there is some entourage $E \subseteq X^2$ such that

$$\begin{aligned} \psi(\mathbb{Z}_+) &\subseteq E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V})] \\ &\subseteq E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})] \end{aligned}$$

and $\phi(\mathbb{Z}_+) \subseteq E[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{U})]$. Thus $q \in \mathcal{U}[p]$. □

Definition 23. (space of ends) Let X be a metric space. As a set the *space of ends* $O(X)$ of X consists of the endpoints in X . A subset $U \subseteq O(X)$ is open if for every $p \in U$ there is a coarse cover \mathcal{U} of X such that

$$\mathcal{U}[p] \subseteq U.$$

This defines a topology on $O(X)$.

Remark 24. The topology on the set of endpoints $O(X)$ is generated by a uniformity: If \mathcal{U} is a coarse cover of X then

$$D_{\mathcal{U}} = \{(p, q) : q \in \mathcal{U}[p]\}$$

is the *entourage associated to \mathcal{U}* . Then $(D_{\mathcal{U}})_{\mathcal{U}}$ over coarse covers \mathcal{U} of X are a base for a diagonal uniformity on $O(X)$.

Lemma 25. *If X is a metric space then $O(X)$ is indeed a uniform space. Coarse covers of X give rise to a base for the uniform structure.*

Proof. We check that $(D_{\mathcal{U}})_{\mathcal{U}}$ over coarse covers are a base for a uniformity on $O(X)$:

1. If \mathcal{U} is a coarse cover of X then $\Delta \subseteq D_{\mathcal{U}}$, where $\Delta = \{(p, p) : p \in O(X)\}$: $p \in \mathcal{U}[p]$.
2. If \mathcal{U}, \mathcal{V} are coarse covers of X then $D_{\mathcal{U}} \cap D_{\mathcal{V}}$ is an entourage: Suppose $\mathcal{U} = (U_i)_i, \mathcal{V} = (V_i)_i$ then define

$$\mathcal{U} \cap \mathcal{V} := (U_i \cap V_j)_{ij}.$$

Suppose p, q are represented by $\phi, \psi : \mathbb{Z}_+ \rightarrow X$. Then $q \in (\mathcal{U} \cap \mathcal{V})[p]$ implies there is an entourage $E \subseteq X^2$ such that

$$\begin{aligned} \psi(\mathbb{Z}_+) &\subseteq E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U} \cap \mathcal{V})] \\ &= E[\bigcup \{U_i \cap V_i : \phi(\mathbb{Z}_+) \wedge U_i \cap V_i\}] \\ &\subseteq \bigcup_{U_i \cap V_i \wedge \phi(\mathbb{Z}_+)} E[U_i] \cap E[V_i] \\ &\subseteq (\bigcup_{U_i \wedge \phi(\mathbb{Z}_+)} E[U_i]) \cap (\bigcup_{V_i \wedge \phi(\mathbb{Z}_+)} E[V_i]) \\ &= E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})] \cap E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V})]. \end{aligned}$$

Thus $q \in \mathcal{U}[p] \cap \mathcal{V}[p]$. This way we have proven:

$$D_{\mathcal{U} \cap \mathcal{V}} \subseteq D_{\mathcal{U}} \cap D_{\mathcal{V}}.$$

3. If \mathcal{U} is a coarse cover of X then there is a coarse cover \mathcal{V} of X such that $D_{\mathcal{V}} \circ D_{\mathcal{V}} \subseteq D_{\mathcal{U}}$: By Lemma 20 there is a coarse star refinement \mathcal{V} of \mathcal{U} . And by Lemma 32 item 2 the uniform cover $(\mathcal{V}[p])_p$ star refines the uniform cover $(\mathcal{U}[p])_p$ thus the result.
4. If \mathcal{U} is a coarse cover then $D_{\mathcal{U}} = D_{\mathcal{U}}^{-1}$.

A subset $D \subseteq E(X)^2$ is an entourage of the uniform structure of $O(X)$ if there is a coarse cover \mathcal{U} of X such that

$$D_{\mathcal{U}} \subseteq D.$$

□

Theorem 26. *If $f : X \rightarrow Y$ is a coarse map between metric spaces then the induced map*

$$\begin{aligned} O(f) : O(X) &\rightarrow O(Y) \\ [\varphi] &\mapsto [f \circ \varphi] \end{aligned}$$

is a continuous map between topological spaces.

Proof. We show $O(f)$ is well defined: if $\phi, \psi : \mathbb{Z}_+ \rightarrow X$ represent the same endpoint in X then there is some entourage $E \subseteq X^2$ such that $E[\psi(\mathbb{Z}_+)] = \phi(\mathbb{Z}_+)$. But then

$$\begin{aligned} f^2(E)[f \circ \psi(\mathbb{Z}_+)] &\supseteq f(E[\psi(\mathbb{Z}_+)]) \\ &= f \circ \phi(\mathbb{Z}_+). \end{aligned}$$

Thus $f \circ \phi, f \circ \psi$ represent the same endpoint in Y .

We show $O(f)$ continuous: For that we show that the inverse image of an open set is an open set.

Let $U \subseteq O(Y)$ be open and $p \in O(f)^{-1}(U)$ be a point. Suppose that p is represented by a coarse map $\phi : \mathbb{Z}_+ \rightarrow X$. Then $f \circ \phi$ represents $O(f)(p) \in U$. Now there is a coarse cover $\mathcal{U} = (U_i)_i$ of Y such that $\mathcal{U}[O(f)(p)] \subseteq U$. Then $f^{-1}(\mathcal{U}) = (f^{-1}(U_i))_i$ is a coarse cover of X .

If $q \in f^{-1}(\mathcal{U})[p]$ we show that $O(f)(q) \in \mathcal{U}[O(f)(p)]$: Suppose that q is represented by a coarse map $\psi : \mathbb{Z}_+ \rightarrow X$. Then there is some entourage $F \subseteq X^2$ such that

$$F[\text{cst}(\phi(\mathbb{Z}_+), f^{-1}(\mathcal{U}))] \supseteq \psi(\mathbb{Z}_+)$$

and

$$F[\text{cst}(\psi(\mathbb{Z}_+), f^{-1}(\mathcal{U}))] \supseteq \phi(\mathbb{Z}_+).$$

By Lemma 19:

$$\begin{aligned} f^2(F)[\text{cst}(f \circ \phi(\mathbb{Z}_+), \mathcal{U})] &\supseteq f^2(F)[f(\text{cst}(\phi(\mathbb{Z}_+), f^{-1}(\mathcal{U})))] \\ &\supseteq f(F[\text{cst}(\phi(\mathbb{Z}_+), f^{-1}(\mathcal{U}))]) \\ &\supseteq f \circ \psi(\mathbb{Z}_+) \end{aligned}$$

and

$$\begin{aligned} f^2(F)[\text{cst}(f \circ \psi(\mathbb{Z}_+), \mathcal{U})] &\supseteq f^2(F)[f(\text{cst}(\psi(\mathbb{Z}_+), f^{-1}(\mathcal{U})))] \\ &\supseteq f(F[\text{cst}(\psi(\mathbb{Z}_+), f^{-1}(\mathcal{U}))]) \\ &\supseteq f \circ \phi(\mathbb{Z}_+). \end{aligned}$$

Now $f \circ \psi$ represents $O(f)(q)$ which by the above is in $\mathcal{U}[O(f)(p)]$. \square

Remark 27. The proof of Theorem 26 uses the following: if $f : X \rightarrow Y$ is a coarse map and $D_{\mathcal{U}}$ the entourage of $O(Y)$ associated to a coarse cover \mathcal{U} of Y then there is an entourage $D_{f^{-1}(\mathcal{U})}$ of $O(X)$ associated to the coarse cover $f^{-1}(\mathcal{U})$ of X such that $(p, q) \in D_{f^{-1}(\mathcal{U})}$ implies $(O(f)(p), O(f)(q)) \in D_{\mathcal{U}}$. Thus $O(f)$ is a uniformly continuous map between uniform spaces $O(X)$ and $O(Y)$.

Lemma 28. *If two coarse maps $f, g : X \rightarrow Y$ are close then $O(f) = O(g)$.*

Proof. Let $p \in O(X)$ be a point that is represented by φ . Now f, g are close thus $H := f \times g(\Delta_X)$ is an entourage. But then

$$H[g \circ \varphi(\mathbb{Z}_+)] \supseteq f \circ \varphi(\mathbb{Z}_+)$$

thus $O(f)(p) = O(g)(p)$. \square

Corollary 29. *If f is a coarse equivalence then $O(f)$ is a homeomorphism between topological spaces $O(X)$ and $O(Y)$. In fact $O(f)$ is a uniform isomorphism between uniform spaces $O(X)$ and $O(Y)$.*

Corollary 30. *If $mCoarse$ denotes the category of metric spaces and coarse maps modulo closeness and Top the category of topological spaces and continuous maps then O is a functor*

$$O : mCoarse \rightarrow Top.$$

If $Uniform$ denotes the category of uniform spaces and uniformly continuous maps then O is a functor

$$O : mCoarse \rightarrow Uniform.$$

Example 31. $O(\mathbb{Z}_+)$ is a point.

4 Properties

Lemma 32. *If X is a metric space*

- *and \mathcal{V} is a coarse star refinement of a coarse cover \mathcal{U} of X then $q \in \mathcal{V}[p]$ and $r \in \mathcal{V}[q]$ implies $r \in \mathcal{U}[p]$.*

- if \mathcal{V} coarsely star refines \mathcal{U} then $(\mathcal{V}[p])_p$ star refines $(\mathcal{U}[p])_p$

Proof. • Suppose p is represented by $\phi : \mathbb{Z}_+ \rightarrow X$, q is represented by $\psi : \mathbb{Z}_+ \rightarrow X$ and r is represented by $\rho : \mathbb{Z}_+ \rightarrow X$. Then $E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V}) \supseteq \psi(\mathbb{Z}_+)]$ and $E[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{V}) \supseteq \phi(\mathbb{Z}_+)]$, $E[\text{cst}(\rho(\mathbb{Z}_+), \mathcal{V}) \supseteq \psi(\mathbb{Z}_+)]$ and $E[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{V}) \supseteq \rho(\mathbb{Z}_+)]$. By Lemma 19 there is an entourage $F \subseteq X^2$ such that $\text{cst}(\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V}), \mathcal{V}) \subseteq F[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})]$. Then

$$\begin{aligned} E^{\circ 2} \circ F[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})] &\supseteq E^{\circ 2}[\text{cst}(\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V}), \mathcal{V})] \\ &\supseteq E[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{V})] \\ &\supseteq \rho(\mathbb{Z}_+) \end{aligned}$$

the other direction works the same way.

- Fix $p \in O(X)$. Then

$$\text{st}(\mathcal{V}[p], (\mathcal{V}[p])_p) \subseteq \mathcal{U}[p]$$

because if $q \in \mathcal{V}[p]$ and $q \in \mathcal{V}[r]$ then $r \in \mathcal{U}[p]$ by Item 1

□

Proposition 33. *If $i : Z \rightarrow Y$ is an inclusion of metric spaces then $O(i) : O(Z) \rightarrow O(Y)$ is a uniform embedding.*

Proof. That $O(i)$ is injective is easy to see.

Define a map

$$\begin{aligned} \Phi : O(i)(O(Z)) &\rightarrow O(Z) \\ O(i)(p) &\mapsto p. \end{aligned}$$

We show Φ is a uniformly continuous map:

If $\mathcal{U} = (U_i)_i$ is a coarse cover of Z we show there is a coarse cover \mathcal{V} of Y such that for every $p, q \in O(Z)$: the relation $O(i)(q) \in \mathcal{V}[O(i)(p)]$ implies $q \in \mathcal{U}[p]$.

Note that for every i the sets $U_i^c \setminus (\bigcup_{j \neq i} U_j)^c$ are coarsely disjoint in Y . By Lemma 17 there are subsets $W_1^i, W_2^i \subseteq Y$ that coarsely cover Y and $W_1^i \not\ll U_i^c$ and $W_2^i \not\ll (\bigcup_{j \neq i} U_j)^c$. Now define

$$\mathcal{V} := (\bigcap_i W_{\sigma(i)}^i)_\sigma$$

with $\sigma(i) \in \{1, 2\}$ all possible permutations. Note that there is some entourage $E \subseteq Z^2$ such that for every σ there is some U_j such that

$$\bigcap_i W_{\sigma(i)}^i \cap Z \subseteq E[U_j].$$

Let $p, q \in O(Z)$ such that $O(i)(q) \in \mathcal{V}[O(i)(p)]$. Suppose p, q are represented by $\phi, \psi : \mathbb{Z}_+ \rightarrow Z$. Then there is some entourage $F \subseteq Y^2$ such that

$$F[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V})] \supseteq \psi(\mathbb{Z}_+)$$

and

$$F[\text{cst}(\psi(\mathbb{Z}_+), \mathcal{V})] \supseteq \phi(\mathbb{Z}_+).$$

Then

$$\begin{aligned} F \circ E[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{U})] &\supseteq F[\text{cst}(\phi(\mathbb{Z}_+), \mathcal{V}) \cap Z] \\ &\supseteq \psi(\mathbb{Z}_+). \end{aligned}$$

The other direction works the same way.

Then $q \in \mathcal{U}[p]$ as we wanted to show. \square

Remark 34. By Proposition 33 and Corollary 29 every coarsely injective coarse map $f : X \rightarrow Y$ induces a uniform embedding. We identify $O(X)$ with its image $O(f)(O(X))$ in $O(Y)$.

Example 35. There is a coarsely surjective coarse map $\omega : \mathbb{Z}_+ \rightarrow \mathbb{Z}^2$. Now $O(\omega) : O(\mathbb{Z}_+) \rightarrow O(\mathbb{Z}^2)$ is not a surjective map obviously.

Lemma 36. *If $f : X \rightarrow Y$ is a coarse map between metric spaces, Y is geodesic proper and $O(f) : O(X) \rightarrow O(Y)$ is surjective then f is already coarsely surjective.*

Proof. Assume the opposite. Then $(\text{im } f)^c \subseteq Y$ contains a countable subset $(s_i)_i$ that is coarsely disjoint to $\text{im } f$. Then by [10, Proposition 93] there is a coarse ray $\rho : \mathbb{Z}_+ \rightarrow Y$, an unbounded subsequence $(s_{i_k})_k$ and an entourage $E \subseteq Y^2$ such that

$$(s_{i_k})_k \subseteq E[\rho(\mathbb{Z}_+)]$$

Now ρ represents a point $r \in O(Y)$ and $O(f)$ is surjective. Thus there is some $p \in O(X)$ such that $O(f)(p) = r$. Suppose p is represented by $\varphi : \mathbb{Z}_+ \rightarrow X$ then $f \circ \varphi$ represents r and has image in $\text{im } f$. Thus there is an entourage $F \subseteq Y^2$ such that $\rho(\mathbb{Z}_+) \subseteq F[\text{im } f]$. Then

$$(s_{i_k})_k \subseteq E \circ F[\text{im } f]$$

a contradiction to the assumption. \square

Remark 37. Note that the coarse map

$$\begin{aligned} \mathbb{Z}_+ &\rightarrow \mathbb{Z}_+ \\ n &\mapsto \lfloor \sqrt{n} \rfloor \end{aligned}$$

is not coarsely injective. Since every map $O(\mathbb{Z}_+) \rightarrow O(\mathbb{Z}_+)$ is an isomorphism we cannot conclude that the functor $O(\cdot)$ reflects isomorphisms.

Lemma 38. *If two subsets U, V coarsely cover a metric space X then*

$$O(U \cap V) = O(U) \cap O(V).$$

Proof. The inclusion $O(U \cap V) \subseteq O(U) \cap O(V)$ is obvious.

We show the reverse inclusion: if $p \in O(U) \cap O(V)$ then it is represented by $\phi : \mathbb{Z}_+ \rightarrow U$ in $O(U)$ and $\psi : \mathbb{Z}_+ \rightarrow V$ in $O(V)$. Then there is an entourage $E \subseteq X^2$ such that

$$E[\psi(\mathbb{Z}_+)] = \phi(\mathbb{Z}_+).$$

Denote by F the set of indices $i \in \mathbb{Z}_+$ such that for each $j \in \mathbb{Z}_+$ the inclusion

$$(\phi(i), \psi(j)) \subseteq E \cap (V^c \times U^c).$$

holds. Since $E \cap (V^c \times U^c)$ is bounded the set F is finite. Now we construct a coarse map $\varphi : \mathbb{Z}_+ \rightarrow U \cap V$: for every $i \in \mathbb{Z}_+ \setminus F$ do:

1. if $\phi(i) \in V$ then define $\varphi(i) := \phi(i)$;
2. if $\phi(i) \notin V$ then there exists some $j \in \mathbb{Z}_+$ such that $(\phi(i), \psi(j)) \in E$. Now define $\varphi(i) := \psi(j)$.

Fix a point $x_0 \in U \cap V$ then for every $i \in F$ define: $\varphi(i) = x_0$. Then φ represents p in $O(U \cap V)$. \square

Lemma 39. *The functor $O(\cdot)$ preserves finite coproducts.*

Proof. Let $X = A \sqcup B$ be a coarse disjoint union of metric spaces. Without loss of generality we assume that A, B cover X as sets. Fix a point $x_0 \in X$. Then there is a coarse map

$$r : X \rightarrow \mathbb{Z}$$

$$x \mapsto \begin{cases} d(x, x_0) & x \in A \\ -d(x, x_0) & x \in B. \end{cases}$$

Note that $O(\mathbb{Z}) = \{-1, 1\}$ is a space which consists of two points with the discrete uniformity. Then $O(r)(O(A)) = 1$ and $O(r)(O(B)) = -1$. Thus $O(X)$ is the uniform disjoint union of $O(A), O(B)$. \square

Proposition 40. *Let X be a metric space. The uniformity $O(X)$ is separated.*

Proof. If $p \neq q$ are two points in $O(X)$ we show there is a coarse cover \mathcal{U} such that

$$q \notin \mathcal{U}[p].$$

Suppose p is represented by $\phi : \mathbb{Z}_+ \rightarrow X$ and q is represented by $\psi : \mathbb{Z}_+ \rightarrow X$. Now there is one of two cases:

1. there is a subsequence $(i_k)_k \subseteq \mathbb{N}$ such that $\phi(i_k)_k \not\sim \psi(\mathbb{Z}_+)$.
2. there is a subsequence $(j_k)_k \subseteq \mathbb{N}$ such that $\psi(j_k)_k \not\sim \phi(\mathbb{Z}_+)$.

Without loss of generality we can assume the first case holds. By Lemma 17 there is a coarse cover $\mathcal{U} = \{U_1, U_2\}$ of X such that $U_1 \not\sim \phi(i_k)_k$ and $U_2 \not\sim \psi(\mathbb{Z}_+)$. Then $q \notin \mathcal{U}[p]$.

Now

$$q \notin \mathcal{U}[p]$$

$$= st(p, (\mathcal{U}[r])_r).$$

Thus the result. \square

Lemma 41. *If X is a metric space,*

- $\mathcal{U} = (U_i)_{i \in I}$ is a coarse cover of X and $p \in O(X)$ is represented by $\phi : \mathbb{Z}_+ \rightarrow X$ then define

$$I(p) := \{i \in I : \phi(\mathbb{Z}_+) \wedge U_i\}.$$

If $S \subseteq I$ is a subset then define

$$U(S) = \{p \in O(X) : \phi(\mathbb{Z}_+) \subseteq E[\bigcup_{i \in S} U_i]\}$$

here $\phi : \mathbb{Z}_+ \rightarrow X$ represents p and $E \subseteq X^2$ is an entourage. If $q \in O(X)$ then $q \in \mathcal{U}[p]$ if and only if $q \in U(I(p))$ and $p \in U(I(q))$.

- Define

$$\mathcal{U}(S) = \{p \in O(X) : p \in U(S), I(p) \supseteq S\}.$$

Then $q \in \mathcal{U}[p]$ if and only if there is some $S \subseteq I$ such that $p, q \in \mathcal{U}(S)$. The uniform cover

$$(\mathcal{U}(S))_{S \subseteq I}$$

associated to $D_{\mathcal{U}}$ is a finite cover.

- The uniform space $O(X)$ is totally bounded.

Proof. • easy.

- We just need to show: if $q \in \mathcal{U}[p]$ then $p \in \mathcal{U}(I(p) \cap I(q))$. For that it is sufficient to show if $\phi : \mathbb{Z}_+ \rightarrow X$ represents p then there is an entourage $E \subseteq X^2$ such that $\phi(\mathbb{Z}_+) \subseteq E[\bigcup_{i \in I(q)} U_i]$. Assume the opposite: there is some subsequence $(i_k)_k \subseteq \mathbb{Z}_+$ such that

$$\phi(i_k)_k \not\subseteq \bigcup_{i \in I(p) \cap I(q)} U_i.$$

Now $\phi(\mathbb{Z}_+) \not\subseteq \bigcup_{i \notin I(p)} U_i$ thus

$$\phi(i_k)_k \not\subseteq \left(\bigcup_{i \in I(p) \cap I(q)} U_i \right) \cup \left(\bigcup_{i \notin I(p)} U_i \right).$$

And thus $\phi(i_k)_k \not\subseteq \bigcup_{i \in I(q)} U_i$ a contradiction to the assumption.

- easy

□

Notation 42. If $A, B \subseteq X$ are two subsets of a metric space and $x_0 \in X$ a point then define

$$\begin{aligned} \chi_{A,B} : \mathbb{N} &\rightarrow \mathbb{R}_+ \\ i &\mapsto d(A \setminus B(x_0, i), B \setminus B(x_0, i)) \end{aligned}$$

Now $A \wedge B$ if and only if $\chi_{A,B}$ is bounded. And if $A_1, A_2, B \subseteq X$ are subsets with $A_1 = E[A_2]$ then the function

$$i \mapsto |\chi_{A_1,B}(i) - \chi_{A_2,B}(i)|$$

is bounded.

Definition 43. Let X be a metric space. If two endpoints $p, q \in O(X)$ are represented by coarse maps $\phi, \psi : \mathbb{Z}_+ \rightarrow X$ then the distance of p to q is at least $f \in \mathbb{R}_+^{\mathbb{N}}$, written $d(p, q) \geq f$, if there is a subsequence $(i_k)_k \subseteq \mathbb{Z}_+$ such that one of the following holds

1. $\phi(\mathbb{Z}_+) \not\subseteq \psi(i_k)_k$ and $\chi_{\phi(\mathbb{Z}_+), \psi(i_k)_k} + c \geq f$ for some $c \geq 0$;
2. $\psi(\mathbb{Z}_+) \not\subseteq \phi(i_k)_k$ and $\chi_{\psi(\mathbb{Z}_+), \phi(i_k)_k} + c \geq f$ for some $c \geq 0$.

We define $d(p, q) = 0$ if and only if $p = q$.

Lemma 44. If X is a metric space and \mathcal{U} a coarse cover of X then there is an unbounded function $f \in \mathbb{R}_+^{\mathbb{N}}$ such that for every two endpoints $p, q \in O(X)$ the relation $q \notin \mathcal{U}[p]$ implies $d(p, q) \geq f$.

Proof. By Lemma 41 the uniform space $O(X)$ is totally bounded. Without loss of generality we can fix an endpoint $p \in O(X)$ and study the endpoints $q \in O(X)$ for which $q \notin \mathcal{U}[p]$.

We will define a function $f \in \mathbb{R}_+^{\mathbb{N}}$ as the minimum of a finite collection $f_0, \dots, f_n \geq 0$ of numbers.

If $q \notin \mathcal{U}[p]$ there are 2 cases:

1. $p \notin U(I(q))$: There is a subset $(i_k)_k \subseteq \mathbb{Z}_+$ such that

$$\phi(i_k)_k \not\subseteq \bigcup_{i \in I(q)} U_i.$$

Now for every $S \subseteq I$ if $p \notin U(S)$: then there is some subset $(i_k)_k \subseteq \mathbb{Z}_+$ such that $\phi(i_k)_k \not\subseteq \bigcup_{i \in S} U_i$. Thus if $I(q) = S$ then $d(p, q) > \chi_{\phi(i_k)_k, \bigcup_{i \in S} U_i}$. Define

$$f_S := \chi_{\phi(i_k)_k, \bigcup_{i \in S} U_i}$$

in this case.

2. $q \notin U(I(p))$: There is some $(i_k)_k \subseteq \mathbb{Z}_+$ such that

$$\psi(i_k)_k \not\subseteq \bigcup_{i \in I(p)} U_i.$$

Now $\psi(i_k)_k \subseteq \bigcup_{i \notin I(p)} U_i$ and $\phi(\mathbb{Z}_+) \not\subseteq \bigcup_{i \notin I(p)} U_i$. Then define

$$f_b := \chi_{\phi(\mathbb{Z}_+), \bigcup_{i \notin I(p)} U_i}.$$

Now define

$$f(i) := \min_S (f_S(i), f_b(i))$$

for every i . Then f has the desired properties. \square

Proposition 45. *If X is a metric space then the uniformity on $O(X)$ is coarser than the uniformity τ_d , induced by d .*

Proof. By Lemma 44 every entourage in $O(X)$ is a neighborhood of an entourage of τ_d on $O(X)$. \square

5 Side Notes

Lemma 46. (Higson corona) *If X is a metric space then the C^* -algebra that determines the Higson corona is a sheaf. That means exactly that the association*

$$U \mapsto C(\nu U) = B_h(U)/B_0(U)$$

for every subset $U \subseteq X$ is a sheaf with values in \mathbf{CStar} . By a sheaf we mean a sheaf on the Grothendieck topology determined by coarse covers on subsets of a coarse space.

Proof. We recall a few definitions which can be found in [1, p.29, 30].

- The algebra of bounded functions that satisfy the Higson condition is denoted by B_h .

- A bounded function $f : X \rightarrow \mathbb{C}$ satisfies the Higson condition if for every entourage $E \subseteq X^2$ the function

$$\begin{aligned} df|_E : E &\rightarrow \mathbb{C} \\ (x, y) &\mapsto f(y) - f(x) \end{aligned}$$

tends to 0 at infinity.

- the ideal of bounded functions that tend to 0 at infinity is called B_0 .
- A function $f : X \rightarrow \mathbb{C}$ tends to 0 at infinity if for every $\varepsilon > 0$ there is a bounded subset $B \subseteq X$ such that $|f(x)| \geq \varepsilon$ implies $x \in B$.

We check the sheaf axioms:

1. global axiom: if U_1, U_2 coarsely cover a subset $U \subseteq X$ and $f \in B_h(X)$ such that $f|_{U_1} \in B_0(U_1)$ and $f|_{U_2} \in B_0(U_2)$ we show that $f \in B_0(U)$ already. Let $\varepsilon > 0$ be a number. Then there are bounded subsets $B_1 \subseteq U_1$ and $B_2 \subseteq U_2$ such that $|f|_{U_i}(x)| \geq \varepsilon$ implies $x \in B_i$ for $i = 1, 2$. Now

$$B := B_1 \cup B_2 \cup (U_1 \cup U_2)^c$$

is a bounded subset of U . Then $|f(x)| \geq \varepsilon$ implies $x \in B$. Thus $f \in B_0(U)$.

2. gluing axiom: if U_1, U_2 coarsely cover a subset $U \subseteq X$ and $f_1 \in B_h(U_1), f_2 \in B_h(U_2)$ are functions such that

$$f_1|_{U_2} = f_2|_{U_1} + g$$

where $g \in B_0(U_1 \cap U_2)$. We show there is a function $f \in B_h(U)$ which restricts to f_1 on U_1 and $f_2 + g$ on U_2 . Define:

$$\begin{aligned} f : U &\rightarrow \mathbb{C} \\ x &\mapsto \begin{cases} f_1(x) & x \in U_1 \\ f_2(x) + g & x \in U_2 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

then f is a bounded function. We show f satisfies the Higson condition: Let $E \subseteq U^2$ be an entourage and $\varepsilon > 0$ be a number. Then there are bounded subsets $B_1 \subseteq U_1$ and $B_2 \subseteq U_2$ such that $|df_i|_{E \cap U_i^2}(x, y)| \geq \varepsilon$ implies $x \in B_i$ for $i = 1, 2$. There is a bounded subset $B_3 \subseteq U$ such that

$$E \cap (U_1^2 \cup U_2^2)^c \cap U^2 \subseteq B_3^2.$$

Define

$$B := B_1 \cup B_2 \cup B_3$$

then $|df|_E(x)| \geq \varepsilon$ implies $x \in B$. Thus f has the desired properties. □

Proposition 47. *If X is a proper geodesic metric space denote by \sim the relation on $O(X)$ of belonging to the same uniform connection component in $O(X)$. Then there is a continuous bijection*

$$O(X)/\sim \rightarrow \Omega(X)$$

where the right side denotes the space of ends of X as a topological space.

Proof. There are several different definitions for the space of ends of a topological space. We use [13, Definition 8.27].

An end in X is represented by a proper continuous map $r : [0, \infty) \rightarrow X$. Two such maps r_1, r_2 represent the same end if for every compact subset $C \subseteq X$ there is some $N \in \mathbb{N}$ such that $r_1[N, \infty), r_2[N, \infty)$ are contained in the same path component of $X \setminus C$.

If $r : [0, \infty) \rightarrow X$ is an end then there is a coarse map $\varphi : \mathbb{Z}_+ \rightarrow X$ and an entourage $E \subseteq X^2$ such that

$$E[r[0, \infty)] = \varphi(\mathbb{Z}_+).$$

We construct φ inductively:

1. $\varphi(0) := r(0)$
2. if $\varphi(i-1) = r(t_{i-1})$ is already defined then $t_i := \min\{t > t_{i-1} : d(r(t_{i-1}), r(t)) = 1\}$. Set $\varphi(i) := r(t_i)$.

By the above construction φ is coarsely uniform. The map φ is coarsely proper because r is proper and X is proper.

Note that every geodesic space is also a length space. If for some compact subset $C \subseteq X$ the space $X \setminus C$ has two path components X_1, X_2 then for every $x_1 \in X_1, x_2 \in X_2$ a path (in particular the shortest) joining x_1 to x_2 contains a point $c \in C$. Thus

$$d(x_1, x_2) = \inf_{c \in C} (d(x_1, c) + d(x_2, c))$$

Then X is the coarse disjoint union of X_1, X_2 . On the other hand if X is the coarse disjoint union of subspaces X_1, X_2 then there is a bounded and in particular because X is proper compact subset $C \subseteq X$ such that

$$X \setminus C = X'_1 \sqcup X'_2$$

is a path disjoint union and $X'_1 \subseteq X_1, X'_2 \subseteq X_2$ differ only by bounded sets.

Now we show the association is continuous:

We use [13, Lemma 8.28] in which $\mathcal{G}_{x_0}(X)$ denotes the set of geodesic rays issuing from $x_0 \in X$. Then [13, Lemma 8.28] states that the canonical map

$$\mathcal{G}_{x_0} \rightarrow \Omega(X)$$

is surjective. Fix $r \in \mathcal{G}_{x_0}$. Then $\tilde{V}_n \subseteq \mathcal{G}_{x_0}$ denotes the set of proper rays $r' : \mathbb{R}_+ \rightarrow X$ such that $r'(n, \infty), r(n, \infty)$ lie in the same path component of $X \setminus B(x_0, n)$. Now [13, Lemma 8.28] states the sets $(V_n = \{[r'] : r' \in \tilde{V}_n\})_n$ form a neighborhood base for $[r] \in \Omega(X)$.

Now to every n we denote by U_1^n the path component of $X \setminus B(x_0, n)$ that contains $r(\mathbb{R}_+)$ and we define $U_2^n := X \setminus U_1^n$. For every $n \in \mathbb{N}$ the sets U_1^n, U_2^n are a coarse cover of X .

Suppose $\rho : \mathbb{Z}_+ \rightarrow X$ is a coarse map associated to r and represents $\tilde{r} \in O(X)$. If $s \in \mathcal{G}_0$ suppose $\sigma : \mathbb{Z}_+ \rightarrow X$ is the coarse map associated to s and represents $\tilde{s} \in O(X)$. If $[s] \notin V_n$ then $\sigma(\mathbb{Z}_+) \not\subset U_1^n$. This implies $\tilde{s} \notin \{U_1^n, U_2^n\}[\tilde{r}]$.

Thus for every $n \in \mathbb{N}$ there is an inclusion $\{U_1^n, U_2^n\}[\tilde{r}] / \sim \subseteq V_n$ by the association. \square

6 Remarks

The starting point of this research was an observation in the studies of [10]: coarse cohomology with twisted coefficients looked like singular cohomology on some kind of boundary. We tried to find a functor from the coarse category to the category of topological spaces that would reflect that observation.

And then we noticed that two concepts play an important role: One is the choice of topology on the *space of ends* and one is the choice of points. The points were designed such that

- coarse maps are mapped by the functor to maps of sets
- and the space \mathbb{Z}_+ is mapped to a point

If the metric space is Gromov hyperbolic then coarse rays represent the points of the Gromov boundary, thus the Gromov boundary is a subset of the space of ends. The topology was trickier to find. We looked for the following properties:

- coarse maps are mapped to continuous maps
- coarse embeddings are mapped to topological embeddings

Now a proximity relation on subsets of a topological space helps constructing the topology on the space of ends of Freudenthal. We discovered that coarse covers on metric spaces give rise to a totally bounded uniformity and thus used that a uniformity on a space gives rise to a topology.

Finally, after a lucky guess, we came up with the uniformity on the set of endpoints. In which way does the space of ends functor reflect isomorphism classes will be studied in a paper that follows.

It would be possible, conversely, after a more thorough examination to find more applications. Coarse properties on metric spaces may give rise to topological properties on metrizable uniform spaces.

We wonder if this result will be of any help with classifying coarse spaces up to coarse equivalence. However, as of yet, the duality has not been studied in that much detail.

References

- [1] J. Roe, *Lectures on coarse geometry*, vol. 31 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2003. <http://dx.doi.org/10.1090/ulect/031>.
- [2] J. Keesling, “Subcontinua of the Higson corona,” *Topology Appl.* **80** no. 1-2, (1997) 155–160. [http://dx.doi.org/10.1016/S0166-8641\(97\)00009-6](http://dx.doi.org/10.1016/S0166-8641(97)00009-6).
- [3] A. N. Dranishnikov, J. Keesling, and V. V. Uspenskij, “On the Higson corona of uniformly contractible spaces,” *Topology* **37** no. 4, (1998) 791–803. [http://dx.doi.org/10.1016/S0040-9383\(97\)00048-7](http://dx.doi.org/10.1016/S0040-9383(97)00048-7).
- [4] A. N. Dranishnikov, “Asymptotic topology,” *Uspekhi Mat. Nauk* **55** no. 6(336), (2000) 71–116. <http://dx.doi.org/10.1070/rm2000v055n06ABEH000334>.
- [5] G. Peschke, “The theory of ends,” *Nieuw Arch. Wisk. (4)* **8** no. 1, (1990) 1–12.
- [6] B. Krön and E. Teufl, “Ends–Group-theoretical and topological aspects,” *Preprint* (2009) .
- [7] I. Kapovich and N. Benakli, “Boundaries of hyperbolic groups,” in *Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001)*, vol. 296 of *Contemp. Math.*, pp. 39–93. Amer. Math. Soc., Providence, RI, 2002. <http://dx.doi.org/10.1090/conm/296/05068>.

- [8] J. Dydak and v. Virk, “Inducing maps between Gromov boundaries,” *Mediterr. J. Math.* **13** no. 5, (2016) 2733–2752.
<http://dx.doi.org/10.1007/s00009-015-0650-z>.
- [9] P. Grzegrzolka and J. Siegert, “Coarse Proximity and Proximity at Infinity,” *ArXiv e-prints* (Apr., 2018) , [arXiv:1804.10263](https://arxiv.org/abs/1804.10263) [math.MG].
- [10] E. Hartmann, “Coarse Cohomology with twisted Coefficients,” *ArXiv e-prints* (Sept., 2017) , [arXiv:1710.06725](https://arxiv.org/abs/1710.06725) [math.AG].
- [11] S. Kalantari and B. Honari, “Asymptotic resemblance,” *Rocky Mountain J. Math.* **46** no. 4, (2016) 1231–1262. <https://doi.org/10.1216/RMJ-2016-46-4-1231>.
- [12] S. Willard, *General topology*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1970.
- [13] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, vol. 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1999.
<http://dx.doi.org/10.1007/978-3-662-12494-9>.