A NOTE ON CONSTRUCTION OF DUAL-HAMILTONIAN GRAPHS

JOÃO PAULO COSTALONGA¹

ABSTRACT. A connected simple graph is said dual-hamiltonian if its vertex set has a 2-coloring such that each color class induces a tree. We call such a coloring a hamiltonian coloring. We prove that if *G* is a graph with a certain type of hamiltonian coloring and *T* is a tree, then $G \times T$ is also dual-hamiltonian having the same certain type of hamiltonian coloring. This result is used to constructed a class of dual-hamiltonian graphs, which includes the hypercubes and other multidimensional grids.

Key words: dual-hamiltonian, Yutsis graph, induced tree, cocircuit, bond.

A **bond** in a connected graph is a minimal set with the property that its deletion results in a disconnected graph. A bond in a connected graph *G* has at most ||G|| - |G| + 2 edges; when a bond has this size it is a **hamiltonian bond**. A graph is **dual-hamiltonian** if it has a hamiltonian bond. Some authors call dual-hamiltonian graphs **Yutsis graphs**. An alternative definition for dual-hamiltonian graphs is provided by the following result:

Theorem 1. (*Jaeger* [3]) A subset $B \subseteq E(G)$ is a hamiltonian bond of a connected graph G if and only if G\B has two induced trees of G as connected components.

If *G* is a planar graph with dual graph G^* , then each hamiltonian bond of *G* corresponds to a hamiltonian cycle of G^* . Determining if a graph is dual-hamiltonian is an NP-complete problem. Indeed, it is in the NP class by the certificate provided by Theorem1 and it is NP-complete because determining if a planar graph is hamiltonian is an NP-complete problem [2].

The most notorious problem about dual-hamiltonian graphs is probably Jaeger's conjecture [3] that all cyclically 4-connected cubic graphs are dual-hamiltonian. Whitney [5] proved that the planar case of this conjecture is equivalent to the Four Colors Theorem. Another motivation to study dual-hamiltonian graphs is their application in quantum mechanics (see [4] for more references).

Some necessary conditions for dual-hamiltonianicity are established in [1]. We can add one more that follows from Theorem 1 with a straightforward calculation:

Corollary 2. If G is a dual-hamiltonian graph, then each induced subgraph H of G satisfies $||H|| \le |H| - 2 + |H|^2/4$.

To the best of our knowledge, in opposition to hamiltonianicity, not many sufficient conditions for dual-hamiltonianicity are established in literature. This make establishing if some important classes of graphs are dual-hamiltonian a bit hard. Our objective here is to provide some tools for constructing dual-hamiltonian graphs using cartesian product of graphs. Our results were first motivated by the conjecture of Haidong Wu and Melissa Flynn (private communication) that all hypercubes are dual-hamiltonian; the *n*-dimensional hypercube Q_n may be defined as the graph with $\{0, 1\}^n$ as vertex-set and such that two vertices are adjacent if and only if they differ by exactly one coordinate. We will establish that Flynn's and Wu's conjecture is true in Corollary 4.

Before stating the main result, some definitions are necessary. Let G be a graph and let C be a 2-coloring of V(G). We say that C is a **hamiltonian coloring** of G if each color class

¹Departamento de Matemática, Universidade Federal do Espírito Santo. Av. Fernando Ferrari, 514; Campus de Goiabeiras, 29075-910, Vitória, ES, Brazil. e-mail: joaocostalonga@gmail.com.

of *C* induces a tree in *G*. By Theorem 1, a 2-coloring *C* of *G* is hamiltonian if and only if the set of edges with edvertices of different colors is a hamiltonian bond of *G*. If $X \subseteq V(G)$ and *C* is a 2-coloring of V(G), we denote by $C\Delta X$ the 2-coloring obtained from *C* by switching the colors of the vertices in *X*. If *C* is a hamiltonian coloring of *G* and *I* and *J* are disjoint 2-subsets of V(G), we say that (I, J) is a **quartet** of *C* if:

- (Q1) both *I* and *J* have one vertex of each color,
- (Q2) $C\Delta I$ is a hamiltonian coloring of G, and
- (Q3) each color class of $C\Delta J$ induces a forest with exactly two connected components: one meeting *I* and the other meeting *J*.

The **cartesian product** $G \times H$ of two graphs G and H is defined as the graph with vertex set $V(G) \times V(H)$ and edge-set as the union of $\{(u, x)(v, x) : uv \in E(G) \text{ and } x \in V(H)\}$ and $\{(u, x)(u, y) : u \in V(G) \text{ and } xy \in E(H)\}$. The next theorem is our main result:

Theorem 3. Let G be a connected graph and T be a tree. If G has a hamiltonian coloring with a quartet, then $G \times T$ also has a hamiltonian coloring with a quartet.

The variation of Theorem 3 where the hamiltonian colorings do not necessarily have quartets is not valid. Indeed, let T_1 and T_2 be disjoint trees, each one having a simple path with three edges, and let G be the graph obtained from $T_1 \cup T_2$ by adding all possible edges linking a vertex of T_1 to a vertex of T_2 . Note that { $V(T_1), V(T_2)$ } is the unique 2-partition of V(G) such that both members induce forests in G. This implies that $G \times T$ is not dual-hamiltonian for each tree T with at least two vertices. This paper contains only one more section, which is dedicated to proving Theorem 3.

From Theorem 3 we derive the next corollary. This result was proved for $n \le 5$ and conjectured in general in an unpublished undergraduate thesis written by Melissa Flynn under supervision of Haidong Wu.

Corollary 4. For $n \ge 2$, the *n*-dimensional hypercube Q_n is dual-hamiltonian. Moreover, for $n \ge 3$, Q_n has a hamiltonian coloring with a quartet.

We denote by P_n the path graph with n edges. It is straightforward to check that Q_1 and Q_2 are dual-hamiltonian. Since $Q_n \cong P_1 \times \cdots \times P_1$, the second part of Corollary 4 follows by induction using Theorem 3 with the next example as initial case.

Example 5. Let $n \ge 1$. The coloring of $P_1 \times P_1 \times P_n$ in Figure 1 is hamiltonian with $((i_1, i_2), (j_1, j_2))$ as quartet. In particular this proves that the 3-dimensional cube has a hamiltonian coloring with a quartet.




FIGURE 1. The figure for Example 5

Similarly to the class of hypercubes, using the next example we may prove that a larger class of multidimensional grids are dual-hamiltonian using the graphs in th next example as initial cases. But we do not know if all the multidimensional grids are dual-hamiltonian. We conjecture so, but we conjecture that some do not have hamiltonian colorings with quartets.

Example 6. For $m \ge 2$ and $n \ge 3$, the grid $P_m \times P_n$ has a hamiltonian coloring with a quartet $((i_1, i_2), (j_1, j_2))$. The coloring is exemplified in Figure 2 for odd values of n and in Figure 3 for even values.




FIGURE 3. A hamiltonian coloring with a quartet for $P_5 \times P_7$.

1. Proofs

quartet for $P_5 \times P_6$.

In this section we prove Theorem 3. Figures 4, 5 and 6 illustrate the construction used in the proof.

Consider a graph *G* and a coloring $C : G \rightarrow \{\text{red}, \text{blue}\}$. We say that an edge of *G* is a **red** (resp. **blue**) edge of *C* if its both endvertices have color red (resp. blue) in *C*. An edge of *G* is **monochromatic** if it is red or blue.

Proof of Theorem 3: We may assume that $|T| \ge 2$. For each $x \in V(T)$, we denote by G_x the graph $G \times T[\{x\}]$ and, for $A \subseteq V(G)$, $A_x := A \times \{x\}$.

Let *C* be a hamiltonian coloring of *G* with a quartet (*I*, *J*) and colors in {red, blue}. We write $I = \{i_{red}, i_{blue}\}$ and $J = \{j_{red}, j_{blue}\}$, where the vertices are colored by *C* with the color in their indices.

For each $x \in V(T)$, let C_x be the coloring of G_x such that $C_x(v, x) := C(v)$ for all $v \in V(G)$. Note that C_x is a hamiltonian coloring of G_x with quartet (I_x, J_x) .

Consider a leaf *r* as root for *T* and, for each $x \in V(T)$, let d(x) denote the usual distance from *r* to *x* in *T*. Now, for each $x \in V(T)$, define $D_x := C_x$ if d(x) is even and $D_x := C_x \Delta(V(G_x) - I_x)$ if d(x) is odd. Let *D* be the 2-coloring of $G \times T$ such that, for each $x \in V(T)$, the restriction of *D* to G_x is equal to D_x . Note that for all $x \in V(T)$, $D(i_{red}, x) = red$ and $D(i_{blue}, x) = blue$. Let *l* be a leaf of *T* other than *r*. We shall prove that *D* is a hamiltonian coloring of $G \times T$ with (J_r, I_l) as quartet. Next we check:

Claim 1. For each $\kappa \in \{\text{red}, \text{blue}\}\)$ and each $xy \in E(T)$, $(i_{\kappa}, x)(i_{\kappa}, y)$ is the unique edge of color κ linking a vertex of G_x to a vertex of G_y .

Indeed, by the definition of $G \times T$, such an edge must be of the form (v, x)(v, y) with $v \in V(G)$. We may assume without loosing generality that d(x) is even and d(y) is odd. So $D_x = C_x$ and $D_y = C_y \Delta(V(G_y) - I_y)$. Thus the edge (v, x)(v, y) is monochromatic if and only if $v \in I$. Moreover, this edge has color κ if and only if $C(v) = \kappa$ and this implies Claim 1.

Denote $H := G \times T$ and let H_{red} be the subgraph of H induced by the red vertices of D. Since T is a tree, Claim 1 implies:

12




FIGURE 4. A hamiltonian coloring C of a graph with a quartet (I, J).

FIGURE 5. A tree.




FIGURE 6. A construction for the hamiltonian coloring *D* of $G \times T$ as in the proof of Theorem 3 with quartet (J_r, I_l) , where *G* is the graph in Figure 4 and *T* is the tree in Figure 5.

Claim 2. Let $xy \in E(G)$, then $(i_{red}, x)(i_{red}, y)$ is a bridge of H_{red} .

By Claim 2, if *Z* is a cycle of *R*, then *Z* is a cycle of G_x for some $x \in V(T)$. But, as C_x is a hamiltonian coloring of G_x , it follows by (Q2) that D_i is also a hamiltonian coloring of G_x . Thus $H_{\text{red}} \cap G_x$ is a tree and H_{red} has no cycles. Note $G[\{i_{\text{red}}\}] \times T$ is a connected subgraph of H_{red} because it is isomorphic to *T*. Each vertex of H_{red} is in a subgraph of the form $H_{\text{red}} \cap G_x$, which is a tree with a vertex in common with $G[\{i_{\text{red}}\}] \times T$. Thus H_{red} is connected and, therefore, H_{red} is a tree. Analogously the same holds for the subgraph of *H* induced by the blue vertices of *D*. This proves that *D* is a hamiltonian coloring of *H*.

It is left to prove that (J_r, I_l) is a quartet for *D*. For this we have to prove (Q1)-(Q3). By construction, both J_l and I_r have vertices of different colors. So (Q1) holds.

Consider the coloring $D' := D\Delta J_r$. Denote by R' the set of red vertices of D' and let $H'_{red} := H[R']$. Note that $H_1 := H'_{red} - (V(G_r) \cap V(H'_{red})) = H_{red} - (V(G_r) \cap V(H_{red}))$. Since H_{red} is a tree, it follows from Claim 2 that H_1 is a tree. Now consider the restriction D'_x of D' to G_x . As $D_i = C_i$ is a hamiltonian coloring of G_r with (I_r, J_r) as quartet, it follows from (Q3) that the red vertices of $D'_r = C_r\Delta J_r$ induce in G_x a forest with exactly two components T_I and T_J such that $V(T_I) \cap I_r = \{(i_{red}, r)\}$ and $V(T_J) \cap J_r = \{(j_{blue}, r)\}$. Let s be the neighbor of r in T. It follows from Claim 1 that $(i_{red}, r)(i_{red}, s)$ and $(j_{blue}, r)(j_{blue}, s)$ are the unique edges that link $G_x \cap H'_{red}$ to H_1 . This implies that H'_{red} is obtained from the disjoint trees H_1 , T_J and T_J by adding $(i_{red}, r)(i_{red}, s)$ linking T_i to H_1 and adding $(j_{blue}, r)(j_{blue}, s)$ linking T_j to H_1 and,

5

therefore, H'_{red} is a tree. Analogously, the blue vertices of D' also induce a tree in H and this proves (Q2).

Consider now $D'' := D\Delta I_l$. Let R'' be the set of red vertices of D'' and $H''_{red} := H[R'']$. Similarly to $H_1, H_2 := H''_{red} - (V(G_l) \cap V(H''_{red})) = H_{red} - (V(G_l) \cap V(H_{red}))$ is a tree. By Claim 1 there is no monochromatic edge of D'' from H_2 to G_l . The restriction D''_l of D'' to G_l is equal to $C_l\Delta I_l$ if d(l) is even and equal to $C_l\Delta V(G_l)$ if d(l) is odd. Thus D''_l is a hamiltonian coloring of G_l and $H''_{red} \cap G_l$ is a tree. Therefore, H''_{red} is a forest with H_2 and $H''_{red} \cap G_l$ as connected components. Note that H_2 has a vertex of J_r and $H''_{red} \cap G_l$ has a vertex of I_l . Since similar arguments are valid for the blue vertices, this implies that (Q3) holds and the theorem is valid.

ACKNOWLEDGMENTS

The Author thanks to Haindong Wu for the helpful discussions on the subject.

REFERENCES

- [1] R.E.L. Aldred, D. Van Dyck, G. Brinkmann, V.Fack, B.D. McKay, *Graph structural properties of non-Yutsis graphs allowing fast recognition*, Disc. Appl. Math. 157 (2009) pp. 377–386.
- [2] M. R. Garey, D. S. Johnson and R. Emdre Tarjan, *The planar hamiltonian circuit probem is NP-complete*, SIAM J. Comput. 5 (1976) 704–714.
- [3] F. Jaeger, *On vertex-induced forests in cubic graphs*, Proceedings 5th Southeastern Conference, Congressus Numerantium (1974) 501–512.
- [4] D. Van Dyck, G. Brinkmann, V. Fack, B. McKay, To be or not to be Yutsis: algorithms for the decision problem, Comput. Phys. Commun. 173 (2005) 61–70.
- [5] H. Whitney, A theorem on graphs, Ann. of Math., 32 (1931), 378–390.