

CONTINUOUS \ast -G-FRAME IN HILBERT C^\ast -MODULES

MOHAMED ROSSAFI AND SAMIR KABBAJ

ABSTRACT. In this paper, we introduce the concept of Continuous \ast -g-Frame in Hilbert C^\ast -Modules and we establish some results. We also discuss the stability problem for Continuous \ast -g-Frame.

1. INTRODUCTION AND PRELIMINARIES

The concept of frames in Hilbert spaces has been introduced by Duffin and Schaeffer [8] in 1952 to study some deep problems in nonharmonic Fourier series, after the fundamental paper [7] by Daubechies, Grossman and Meyer, frame theory began to be widely used, particularly in the more specialized context of wavelet frames and Gabor frames [10].

Traditionally, frames have been used in signal processing, image processing, data compression and sampling in sampling theory. A discrete frame is a countable family of elements in a separable Hilbert space which allows for a stable, not necessarily unique, decomposition of an arbitrary element into an expansion of the frame elements. The concept of a generalization of frames to a family indexed by some locally compact space endowed with a Radon measure was proposed by G. Kaiser [12] and independently by Ali, Antoine and Gazeau [1]. These frames are known as continuous frames. Gabardo and Han in [11] called these frames frames associated with measurable spaces, Askari-Hemmat, Dehghan and Radjabalipour in [2] called them generalized frames and in mathematical physics they are referred to as coherent states [1].

In this paper, we introduce the notion of Continuous \ast -g-Frame which are generalization of \ast -g-Frame in Hilbert C^\ast -Modules introduced by A.Alijani [5] and we establish some results.

The paper is organized as follows, we continue this introductory section we briefly recall the definitions and basic properties of C^\ast -algebra, Hilbert C^\ast -modules. In Section 2, we introduce the Continuous \ast -g-Frame, the Continuous pre- \ast -g-frame operator and the Continuous \ast -g-frame operator. In Section 3, we discuss the stability problem for Continuous \ast -g-Frame.

In the following we briefly recall the definitions and basic properties of C^\ast -algebra, Hilbert \mathcal{A} -modules. Our reference for C^\ast -algebras is [9, 6]. For a C^\ast -algebra \mathcal{A} if $a \in \mathcal{A}$ is positive we write $a \geq 0$ and \mathcal{A}^+ denotes the set of positive elements of \mathcal{A} .

2010 *Mathematics Subject Classification.* Primary 41A58; Secondary 42C15.

Key words and phrases. Continuous Frame, \ast -g-frame, Continuous \ast -g-Frame, C^\ast -algebra, Hilbert \mathcal{A} -modules.

Definition 1.1. [6]. If \mathcal{A} is a Banach algebra, an involution is a map $a \rightarrow a^*$ of \mathcal{A} into itself such that for all a and b in \mathcal{A} and all scalars α the following conditions hold:

- (1) $(a^*)^* = a$.
- (2) $(ab)^* = b^*a^*$.
- (3) $(\alpha a + b)^* = \bar{\alpha}a^* + b^*$.

Definition 1.2. [6]. A C^* -algebra \mathcal{A} is a Banach algebra with involution such that :

$$\|a^*a\| = \|a\|^2$$

for every a in \mathcal{A} .

Example 1.3. $\mathcal{B} = B(\mathcal{H})$ the algebra of bounded operators on a Hilbert space, is a C^* -algebra, where for each operator A , A^* is the adjoint of A .

Definition 1.4. [13]. Let \mathcal{A} be a unital C^* -algebra and \mathcal{H} be a left \mathcal{A} -module, such that the linear structures of \mathcal{A} and \mathcal{H} are compatible. \mathcal{H} is a pre-Hilbert \mathcal{A} -module if \mathcal{H} is equipped with an \mathcal{A} -valued inner product $\langle \cdot, \cdot \rangle : \mathcal{H} \times \mathcal{H} \rightarrow \mathcal{A}$, such that is sesquilinear, positive definite and respects the module action. In the other words,

- (i) $\langle x, x \rangle \geq 0$ for all $x \in \mathcal{H}$ and $\langle x, x \rangle = 0$ if and only if $x = 0$.
- (ii) $\langle ax + y, z \rangle = a\langle x, y \rangle + \langle y, z \rangle$ for all $a \in \mathcal{A}$ and $x, y, z \in \mathcal{H}$.
- (iii) $\langle x, y \rangle = \langle y, x \rangle^*$ for all $x, y \in \mathcal{H}$.

For $x \in \mathcal{H}$, we define $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$. If \mathcal{H} is complete with $\|\cdot\|$, it is called a Hilbert \mathcal{A} -module or a Hilbert C^* -module over \mathcal{A} . For every a in C^* -algebra \mathcal{A} , we have $|a| = (a^*a)^{\frac{1}{2}}$ and the \mathcal{A} -valued norm on \mathcal{H} is defined by $|x| = \langle x, x \rangle^{\frac{1}{2}}$ for $x \in \mathcal{H}$.

The following lemmas will be used to prove our mains results

Lemma 1.5. [15]. Let \mathcal{H} be Hilbert \mathcal{A} -module. If $T \in \text{End}_{\mathcal{A}}^*(\mathcal{H})$, then

$$\langle Tx, Tx \rangle \leq \|T\|^2 \langle x, x \rangle, \forall x \in \mathcal{H}.$$

Lemma 1.6. [3]. Let \mathcal{H} and \mathcal{K} two Hilbert \mathcal{A} -modules and $T \in \text{End}^*(\mathcal{H}, \mathcal{K})$. Then the following statements are equivalent:

- (i) T is surjective.
- (ii) T^* is bounded below with respect to norm, i.e., there is $m > 0$ such that $\|T^*x\| \geq m\|x\|$ for all $x \in \mathcal{K}$.
- (iii) T^* is bounded below with respect to the inner product, i.e., there is $m' > 0$ such that $\langle T^*x, T^*x \rangle \geq m'\langle x, x \rangle$ for all $x \in \mathcal{K}$.

Lemma 1.7. [4]. Let \mathcal{H} and \mathcal{K} two Hilbert \mathcal{A} -modules and $T \in \text{End}^*(\mathcal{H}, \mathcal{K})$. Then:

- (i) If T is injective and T has closed range, then the adjointable map T^*T is invertible and

$$\|(T^*T)^{-1}\|^{-1} \leq T^*T \leq \|T\|^2.$$

- (ii) If T is surjective, then the adjointable map TT^* is invertible and

$$\|(TT^*)^{-1}\|^{-1} \leq TT^* \leq \|T\|^2.$$

2. CONTINUOUS *-G-FRAME IN HILBERT C^* -MODULES

Let X be a Banach space, (Ω, μ) a measure space, and function $f : \Omega \rightarrow X$ a measurable function. Integral of the Banach-valued function f has defined Bochner and others. Most properties of this integral are similar to those of the integral of real-valued functions. Because every C^* -algebra and Hilbert C^* -module is a Banach space thus we can use this integral and its properties.

Let (Ω, μ) be a measure space, let U and V be two Hilbert C^* -modules, $\{V_w : w \in \Omega\}$ is a sequence of subspaces of V , and $End_{\mathcal{A}}^*(U, V_w)$ is the collection of all adjointable \mathcal{A} -linear maps from U into V_w . We define

$$\oplus_{w \in \Omega} V_w = \left\{ x = \{x_w\} : x_w \in V_w, \left\| \int_{\Omega} |x_w|^2 d\mu(w) \right\| < \infty \right\}.$$

For any $x = \{x_w : w \in \Omega\}$ and $y = \{y_w : w \in \Omega\}$, if the \mathcal{A} -valued inner product is defined by $\langle x, y \rangle = \int_{\Omega} \langle x_w, y_w \rangle d\mu(w)$, the norm is defined by $\|x\| = \|\langle x, x \rangle\|^{\frac{1}{2}}$, the $\oplus_{w \in \Omega} V_w$ is a Hilbert C^* -module.

Definition 2.1. We call $\{\Lambda_w \in End_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ a continuous *-g-frame for Hilbert C^* -module U with respect to $\{V_w : w \in \Omega\}$ if

- for any $x \in U$, the function $\tilde{x} : \Omega \rightarrow V_w$ defined by $\tilde{x}(w) = \Lambda_w x$ is measurable;
- there exist two strictly nonzero elements A and B in \mathcal{A} such that

$$A \langle x, x \rangle A^* \leq \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \leq B \langle x, x \rangle B^*, \forall x \in U. \quad (2.1)$$

The elements A and B are called continuous *-g-frame bounds. If $A = B$ we call this continuous *-g-frame a continuous tight *-g-frame, and if $A = B = 1_{\mathcal{A}}$ it is called a continuous Parseval *-g-frame. If only the right-hand inequality of (2.1) is satisfied, we call $\{\Lambda_w : w \in \Omega\}$ a continuous *-g-Bessel for U with respect to $\{\Lambda_w : w \in \Omega\}$ with Bessel bound B .

We mentioned that the set of all continuous g-frames in Hilbert C^* -Modules can be considered as a subset of continuous *-g-frame. To illustrate this, let $\{\Lambda_w : w \in \Omega\}$ be a continuous g-frames for Hilbert C^* -Modules U with real continuous g-frames bounds A and B . note that for all $x \in U$,

$$(\sqrt{A})1_{\mathcal{A}} \langle x, x \rangle (\sqrt{A})1_{\mathcal{A}} \leq \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \leq (\sqrt{B})1_{\mathcal{A}} \langle x, x \rangle (\sqrt{B})1_{\mathcal{A}}.$$

Therefore, every continuous g-frames in Hilbert C^* -Modules U with real bounds A and B is a continuous *-g-frame in U with \mathcal{A} -valued bounds $(\sqrt{A})1_{\mathcal{A}}$ and $(\sqrt{B})1_{\mathcal{A}}$.

Theorem 2.2. Let $\{\Lambda_w \in End_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ be a continuous *-g-frame for U , with lower and upper bounds A and B , respectively. The continuous *-g-frame transform $T : U \rightarrow \oplus_{w \in \Omega} V_w$ defined by: $Tx = \{\Lambda_w x : w \in \Omega\}$ is injective, closed range adjointable and $\|T\| \leq \|B\|$. The adjoint operator T^* is surjective and it given by: $T^*x = \int_{\Omega} \Lambda_w^* x_w d\mu(w)$ where $x = \{x_w\}_{w \in \Omega}$.

Proof. Let $x \in U$, by the definition of a continuous $*$ -g-frame for U we have

$$A\langle x, x \rangle A^* \leq \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \leq B\langle x, x \rangle B^*.$$

So

$$A\langle x, x \rangle A^* \leq \langle Tx, Tx \rangle \leq B\langle x, x \rangle B^*. \quad (2.2)$$

If $Tx = 0$ then $\langle x, x \rangle = 0$ so $x = 0$ i.e. T is injective.

We now show that the range of T is closed. Let $\{Tx_n\}_{n \in \mathbb{N}}$ be a sequence in the range of T such that $\lim_{n \rightarrow \infty} Tx_n = y$.

By (2.2) we have, for $n, m \in \mathbb{N}$,

$$\|A\langle x_n - x_m, x_n - x_m \rangle A^*\| \leq \|\langle T(x_n - x_m), T(x_n - x_m) \rangle\| = \|T(x_n - x_m)\|^2.$$

Since $\{Tx_n\}_{n \in \mathbb{N}}$ is Cauchy sequence in U ,

$$\|A\langle x_n - x_m, x_n - x_m \rangle A^*\| \rightarrow 0, \text{ as } n, m \rightarrow \infty.$$

Note that for $n, m \in \mathbb{N}$,

$$\|\langle x_n - x_m, x_n - x_m \rangle\| = \|A^{-1}A\langle x_n - x_m, x_n - x_m \rangle A^*(A^*)^{-1}\|$$

$$\leq \|A^{-1}\|^2 \|A\langle x_n - x_m, x_n - x_m \rangle A^*\|.$$

Therefore the sequence $\{x_n\}_{n \in \mathbb{N}}$ is Cauchy and hence there exists $x \in U$ such that $x_n \rightarrow x$ as $n \rightarrow \infty$. Again by (2.2) we have $\|T(x_n - x)\|^2 \leq \|B\|^2 \|\langle x_n - x, x_n - x \rangle\|$.

Thus $\|Tx_n - Tx\| \rightarrow 0$ as $n \rightarrow \infty$ implies that $Tx = y$. It concludes that the range of T is closed.

For all $x \in U$, $y = \{y_w\} \in \oplus_{w \in \Omega} V_w$, we have

$$\langle Tx, y \rangle = \int_{\Omega} \langle \Lambda_w x, y_w \rangle d\mu(w) = \int_{\Omega} \langle x, \Lambda_w^* y_w \rangle d\mu(w) = \left\langle x, \int_{\Omega} \Lambda_w^* y_w d\mu(w) \right\rangle.$$

Then T is adjointable and $T^*y = \int_{\Omega} \Lambda_w^* y_w d\mu(w)$.

By (2.2) we have $\|Tx\|^2 \leq \|B\|^2 \|x\|^2$ so $\|T\| \leq \|B\|$.

By (2.2) we have $\|Tx\| \geq \|A^{-1}\|^{-1} \|x\| \forall x \in U$ so by lemma 1.6 T^* is surjective. This completes the proof. \square

Now we define the continuous $*$ -g-frame operator and studies some of its properties.

Definition 2.3. Let $\{\Lambda_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ be a continuous $*$ -g-frame for U . Define the continuous $*$ -g-frame operator S on U by: $Sx = T^*Tx = \int_{\Omega} \Lambda_w^* \Lambda_w x d\mu(w)$.

Theorem 2.4. The continuous $*$ -g-frame operator S is a bounded, positive, self-adjoint, invertible and $\|A^{-1}\|^{-2} \leq \|S\| \leq \|B\|^2$.

Proof. First we show, S is a selfadjoint operator. By definition we have $\forall x, y \in U$

$$\begin{aligned}\langle Sx, y \rangle &= \left\langle \int_{\Omega} \Lambda_w^* \Lambda_w x d\mu(w), y \right\rangle \\ &= \int_{\Omega} \langle \Lambda_w^* \Lambda_w x, y \rangle d\mu(w) \\ &= \int_{\Omega} \langle x, \Lambda_w^* \Lambda_w y \rangle d\mu(w) \\ &= \left\langle x, \int_{\Omega} \Lambda_w^* \Lambda_w y d\mu(w) \right\rangle \\ &= \langle x, Sy \rangle.\end{aligned}$$

Then S is a selfadjoint.

By Lemma 1.7 and Theorem 2.2, S is invertible. Clearly S is positive.

By definition of a continuous *-g-frame we have

$$A\langle x, x \rangle A^* \leq \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \leq B\langle x, x \rangle B^*.$$

So

$$A\langle x, x \rangle A^* \leq \langle Sx, x \rangle \leq B\langle x, x \rangle B^*.$$

This give

$$\|A^{-1}\|^{-2} \|x\|^2 \leq \|\langle Sx, x \rangle\| \leq \|B\|^2 \|x\|^2, \forall x \in U.$$

If we take supremum on all $x \in U$, where $\|x\| \leq 1$, then $\|A^{-1}\|^{-2} \leq \|S\| \leq \|B\|^2$. \square

Theorem 2.5. Let $\{\Lambda_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ be a continuous *-g-frame for U , with lower and upper bounds A and B , respectively and with the continuous *-g-frame operator S . Let $T \in \text{End}_{\mathcal{A}}^*(U)$ be invertible. Then $\{\Lambda_w T : w \in \Omega\}$ is a continuous *-g-frame for U with continuous *-g-frame operator $T^* S T$.

Proof. We have

$$A\langle Tx, Tx \rangle A^* \leq \int_{\Omega} \langle \Lambda_w Tx, \Lambda_w Tx \rangle d\mu(w) \leq B\langle Tx, Tx \rangle B^*, \forall x \in U. \quad (2.3)$$

Using Lemma , we have $\|(T^* T)^{-1}\|^{-1} \langle x, x \rangle \leq \langle Tx, Tx \rangle$, $\forall x \in U$. Or $\|T^{-1}\|^{-2} \leq \|(T^* T)^{-1}\|^{-1}$. This implies

$$\|T^{-1}\|^{-1} A\langle x, x \rangle (\|T^{-1}\|^{-1} A)^* \leq A\langle Tx, Tx \rangle A^*, \forall x \in U. \quad (2.4)$$

And we know that $\langle Tx, Tx \rangle \leq \|T\|^2 \langle x, x \rangle$, $\forall x \in U$. This implies that

$$B\langle Tx, Tx \rangle B^* \leq \|T\| B\langle x, x \rangle (\|T\| B)^*, \forall x \in U. \quad (2.5)$$

Using (2.3), (2.4), (2.5) we have

$$\|T^{-1}\|^{-1} A\langle x, x \rangle (\|T^{-1}\|^{-1} A)^* \leq \int_{\Omega} \langle \Lambda_w Tx, \Lambda_w Tx \rangle d\mu(w) \leq B\|T\| \langle x, x \rangle (B\|T\|)^*, \forall x \in U. \quad (2.6)$$

So $\{\Lambda_w T : w \in \Omega\}$ is a continuous *-g-frame for U .

Moreover for every $x \in U$, we have

$$T^*STx = T^* \int_{\Omega} \Lambda_w^* \Lambda_w T x d\mu(w) = \int_{\Omega} T^* \Lambda_w^* \Lambda_w T x d\mu(w) = \int_{\Omega} (\Lambda_w T)^* (\Lambda_w T) x d\mu(w).$$

This completes the proof. \square

Corollary 2.6. *Let $\{\Lambda_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ be a continuous $*$ -g-frame for U , with continuous $*$ -g-frame operator S . Then $\{\Lambda_w S^{-1} : w \in \Omega\}$ is a continuous $*$ -g-frame for U .*

Proof. Result the next theorem by taking $T = S^{-1}$. \square

3. STABILITY PROBLEM FOR CONTINUOUS $*$ -G-FRAME IN HILBERT C^* -MODULES

The question of stability plays an important role in various fields of applied mathematics. The classical theorem of the stability of a base is due to Paley and Wiener. It is based on the fact that a bounded operator T on a Banach space is invertible if we have: $\|I - T\| < 1$.

Theorem 3.1 ([14] Paley-Wiener). *Let $\{f_i\}_{i \in \mathbb{N}}$ be a basis of a Banach space X , and $\{g_i\}_{i \in \mathbb{N}}$ a sequence of vectors in X . If there exists a constant $\lambda \in [0, 1)$ such that*

$$\left\| \sum_{i \in \mathbb{N}} c_i (f_i - g_i) \right\| \leq \lambda \left\| \sum_{i \in \mathbb{N}} c_i f_i \right\|$$

for all finite sequence $\{c_i\}_{i \in \mathbb{N}}$ of scalars, then $\{g_i\}_{i \in \mathbb{N}}$ is also a basis for X .

Theorem 3.2. *Let $\{\Lambda_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ be a continuous $*$ -g-frame for U , with lower and upper bounds A and B , respectively. Let $\Gamma_w \in \text{End}_{\mathcal{A}}^*(U, V_w)$ for any $w \in \Omega$. Then the following are equivalent:*

- (1) $\{\Gamma_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ is a continuous $*$ -g-frame for U .
- (2) There exists a constant $M > 0$, such that for any $x \in U$, one has

$$\begin{aligned} & \left\| \int_{\Omega} \langle (\Lambda_w - \Gamma_w)x, (\Lambda_w - \Gamma_w)x \rangle d\mu(w) \right\| \\ & \leq M \min \left(\left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|, \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\| \right). \end{aligned} \quad (3.1)$$

Proof. (1) \Rightarrow (2). Suppose that $\{\Gamma_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ is a continuous $*$ -g-frame for U with lower and upper bounds C and D , respectively. Then for

any $x \in U$, we have

$$\begin{aligned}
\left\| \int_{\Omega} \langle (\Lambda_w - \Gamma_w)x, (\Lambda_w - \Gamma_w)x \rangle d\mu(w) \right\|^{\frac{1}{2}} &= \left\| \{(\Lambda_w - \Gamma_w)x\}_{w \in \Omega} \right\| \\
&\leq \left\| \{\Lambda_w x\}_{w \in \Omega} \right\| + \left\| \{\Gamma_w x\}_{w \in \Omega} \right\| \\
&= \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&\leq \|B\| \|\langle x, x \rangle\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&\leq \|B\| \|C^{-1}\| \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&= \left(\|B\| \|C^{-1}\| + 1 \right) \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}}.
\end{aligned}$$

Similary we have

$$\left\| \int_{\Omega} \langle (\Lambda_w - \Gamma_w)x, (\Lambda_w - \Gamma_w)x \rangle d\mu(w) \right\|^{\frac{1}{2}} \leq \left(\|D\| \|A^{-1}\| + 1 \right) \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}}.$$

Let $M = \min \left\{ \left(\|B\| \|C^{-1}\| + 1 \right)^2, \left(\|D\| \|A^{-1}\| + 1 \right)^2 \right\}$, then the inequality (3.1)

holds.

(2) \Rightarrow (1). Suppose that the inequality (3.1) holds. For any $x \in U$, we have

$$\begin{aligned}
\|A^{-1}\|^{-1} \|\langle x, x \rangle\|^{\frac{1}{2}} &\leq \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&\leq \left\| \int_{\Omega} \langle (\Lambda_w - \Gamma_w)x, (\Lambda_w - \Gamma_w)x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&\leq M^{\frac{1}{2}} \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
&= (1 + M^{\frac{1}{2}}) \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}}.
\end{aligned}$$

Also we obtain

$$\begin{aligned}
& \left\| \int_{\Omega} \langle \Gamma_w x, \Gamma_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
& \leq \left\| \int_{\Omega} \langle (\Lambda_w - \Gamma_w)x, (\Lambda_w - \Gamma_w)x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
& \leq M^{\frac{1}{2}} \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} + \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
& = (1 + M^{\frac{1}{2}}) \left\| \int_{\Omega} \langle \Lambda_w x, \Lambda_w x \rangle d\mu(w) \right\|^{\frac{1}{2}} \\
& \leq (1 + M^{\frac{1}{2}}) \|B\| \|\langle x, x \rangle\|^{\frac{1}{2}}.
\end{aligned}$$

So $\{\Gamma_w \in \text{End}_{\mathcal{A}}^*(U, V_w) : w \in \Omega\}$ is a continuous $*$ -g-frame for U . \square

REFERENCES

1. S. T. Ali, J. P. Antoine, J. P. Gazeau, *Continuous frames in Hilbert spaces*, Annals of Physics 222 (1993), 1-37.
2. A. Askari-Hemmat, M. A. Dehghan, M. Radjabalipour, *Generalized frames and their redundancy*, Proc. Amer. Math. Soc. 129 (2001), no. 4, 1143-1147.
3. L. Arambašić, *On frames for countably generated Hilbert C^* -modules*, Proc. Amer. Math. Soc. 135 (2007) 469-478.
4. A. Alijani, M. Dehghan, *$*$ -frames in Hilbert C^* modules*, U. P. B. Sci. Bull. Series A 2011.
5. A. Alijani, *Generalized Frames with C^* -Valued Bounds and their Operator Duals*, Filomat 29:7, 14691479 DOI 10.2298/FIL1507469A (2015).
6. J. B. Conway, *A Course In Operator Theory*, AMS, V.21, 2000.
7. I. Daubechies, A. Grossmann, and Y. Meyer, *Painless nonorthogonal expansions*, J. Math. Phys. 27 (1986), 1271-1283.
8. R. J. Duffin, A. C. Schaeffer, *A class of nonharmonic fourier series*, Trans. Amer. Math. Soc. 72 (1952), 341-366.
9. F. R. Davidson, *C^* -algebra by example*, Fields Ins. Monog. 1996.
10. D. Gabor, *Theory of communications*, J. Elec. Eng. 93 (1946), 429-457.
11. J. P. Gabardo and D. Han, *Frames associated with measurable space*, Adv. Comp. Math. 18 (2003), no. 3, 127-147.
12. G. Kaiser, *A Friendly Guide to Wavelets*, Birkhauser, Boston, 1994.
13. I. Kaplansky, *Modules over operator algebras*, Amer. J. Math. 75 (1953), 839-858.
14. R. Paley, N. Wiener, *Fourier Transforms in Complex Domains*. MAS Colloquium Publications 19, 1934.
15. W. Paschke, *Inner product modules over B^* -algebras*, Trans. Amer. Math. Soc., (182)(1973), 443-468.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF IBN TOFAIL
B.P.133 KENITRA, MOROCCO
E-mail address: rossafimohamed@gmail.com

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF IBN TOFAIL
B.P.133 KENITRA, MOROCCO
E-mail address: samkabbaj@yahoo.fr