

FREE-BOOLEAN INDEPENDENCE WITH AMALGAMATION

WEIHUA LIU AND PING ZHONG

ABSTRACT. In this paper, we develop the notion of free-Boolean independence in an amalgamation setting. We construct free-Boolean cumulants and show that the vanishing of mixed free-Boolean cumulants is equivalent to our free-Boolean independence with amalgamation. We also provide a characterization of free-Boolean independence by conditions in terms of mixed moments. In addition, we study free-Boolean independence over a C^* -algebra and prove a positivity property.

1. INTRODUCTION

Free probability theory is a probability theory that studies noncommutative random variables with highest noncommutativity. This theory, due to Voiculescu, is based on the notion of free independence which is an analogue of the classical independence. In [18], Voiculescu generalized his notion of free independence to free independence with amalgamation over an arbitrary algebra in details. To be specific, moments of random variables are no longer scalar numbers but elements from a given algebra. On the other aspect, Voiculescu started to study pairs of random variables simultaneously thereby generalized the notion of free independence to a notion of bi-free independence [19]. Further more, the notion of bi-free independence with amalgamation, defined by Voiculescu [19], was fully developed in [2]. there are exactly two unital universal independence relations, namely Voiculescu's free independence relation, the classical independence relation [15]. It was mentioned that we would obtain more independence relations by decreasing the number of axioms for universal products [8]. For instance, people introduced Boolean independence [17], monotone independence [10], conditionally independence [1] in various contexts. Their operator-valued generalization were studied as well [9, 12, 13]. Recently, their corresponding independence relations for pairs of random variables, analog of Voiculescu's bi-free theory, were introduced and studied [4, 3, 6]. Furthermore, the conditionally bi-free independence with amalgamation is studied in [5].

In [8], the first-named author introduced a notion of mixed independence relations for pairs of random variables, where random variables in different faces exhibit different kinds of noncommutative independence. In particular, the combinatorics of free-Boolean independence relation was fully developed. In this paper, we generalize the notion of free-Boolean independence to an amalgamation setting. Relevant combinatorial tools are extended to study this new independence. Beyond the corresponding combinatorial results, we address the positivity of free-Boolean independence with amalgamation. Therefore, it is possible to study the relation in topological probability spaces but not only algebraic probability spaces. For instance, we can study our

free-Boolean independence with amalgamation over a C^* -algebra, which is a suitable framework to address some probabilistic questions. We plan to study probabilistic results such as operator-valued infinitely divisible laws in a forthcoming paper.

The paper is organized as follows: Besides this introduction, in Section 2, we give the definition of free-Boolean independence with amalgamation over an algebra. In Section 3, we review some relevant combinatorial tools. In Section 4, we demonstrate that free-Boolean independence can be characterized by the property of the vanishing of mixed free-Boolean cumulants. In Section 5, we prove an operator-valued version of free-Boolean central limit law. In Section 6, we provide an equivalent characterization of free-Boolean independence by certain moments-conditions. In Section 7, we study the positivity property for free-Boolean independence relation.

2. PRELIMINARIES AND NOTATIONS

In this section, we give the motivation and the definition for free-Boolean independence relation with amalgamation over an algebra.

Definition 2.1. A B - B -bimodule with a specified projection is a triple $(\mathcal{X}, \overset{\circ}{\mathcal{X}}, p)$, where $\mathcal{X}$ is a direct sum of B - B -bimodules $\mathcal{X} = B \oplus \overset{\circ}{\mathcal{X}}$, and $p : \mathcal{X} \rightarrow B$ is the projection

$$p(b \oplus \eta) = b.$$

Denote by $\mathcal{L}(\mathcal{X})$ the algebra of linear operators with respect to the B - B -bimodule structure. The *expectation* from $\mathcal{L}(\mathcal{X})$ onto B is the linear map $\mathbb{E}_{\mathcal{L}(\mathcal{X})} : \mathcal{L}(\mathcal{X}) \rightarrow B$ defined by

$$\mathbb{E}_{\mathcal{L}(\mathcal{X})}(a) = p(a(1_B \oplus 0)).$$

We now recall the definition of the reduced free product of B - B -bimodules with specified projections [16, 18]. Let $\{(\mathcal{X}_i, \overset{\circ}{\mathcal{X}}_i, p_i)\}_{i \in \mathcal{I}}$ be a family of B - B -bimodules with specified projections. The *reduce free product of $\{(\mathcal{X}_k, \overset{\circ}{\mathcal{X}}_k, p_k)\}_{k \in \mathcal{I}}$ with amalgamation over B* is defined to be the B - B -bimodule with a specified projection $(\mathcal{X}, \overset{\circ}{\mathcal{X}}, p)$, where $\mathcal{X} = B \oplus \overset{\circ}{\mathcal{X}}$ and $\overset{\circ}{\mathcal{X}}$ is the B - B -bimodule defined by

$$\overset{\circ}{\mathcal{X}} = \bigoplus_{n \geq 1} \bigoplus_{i_1 \neq i_2 \neq \dots \neq i_n} \overset{\circ}{\mathcal{X}}_{i_1} \otimes_B \dots \otimes_B \overset{\circ}{\mathcal{X}}_{i_n}.$$

For each $i \in \mathcal{I}$, we denote by

$$\mathcal{X}(i) = B \oplus \bigoplus_{n \geq 1} \bigoplus_{\substack{i_1 \neq i_2 \neq \dots \neq i_n \\ i_1 \neq i}} \overset{\circ}{\mathcal{X}}_{i_1} \otimes_B \dots \otimes_B \overset{\circ}{\mathcal{X}}_{i_n},$$

and let V_i be the natural isomorphism of bimodules

$$V_i : \mathcal{X} \rightarrow \mathcal{X}_i \otimes_B \mathcal{X}(i).$$

For each $i \in \mathcal{I}$, $\lambda_i : \mathcal{L}(\mathcal{X}_i) \rightarrow \mathcal{L}(\mathcal{X})$ is a unital homomorphism defined by

$$\lambda_i(a) = V_i^{-1}(a \otimes I)V_i$$

and $\beta_i : \mathcal{L}(\mathcal{X}_i) \rightarrow \mathcal{L}(\mathcal{X})$ is a linear map defined by

$$\beta_i(a) = P_i \lambda_i(a) P_i,$$

where $P_i : \mathcal{X} \rightarrow B \oplus \mathring{\mathcal{X}}_i$ is the natural projection onto $B \oplus \mathring{\mathcal{X}}_i$ and vanishes on the other direct summands.

Proposition 2.2. *For any $a \in \mathcal{L}(\mathcal{X}_i)$, we have $P_i \lambda_i(a) = \lambda_i(a) P_i$.*

Proof. Notice that the reduced free product $\mathcal{X}$ of B - B -bimodules with specified projections can be decomposed as

$$\mathcal{X} = (B \oplus \mathring{\mathcal{X}}_i) \oplus \mathcal{X}'_i, \quad \text{where} \quad \mathcal{X}'_i = \bigoplus_{j \neq i} \mathring{\mathcal{X}}_j \oplus \bigoplus_{n \geq 2} \bigoplus_{i_1 \neq i_2 \neq \dots \neq i_n} \mathring{\mathcal{X}}_{i_1} \otimes_B \dots \otimes_B \mathring{\mathcal{X}}_{i_n}.$$

The space $B \oplus \mathring{\mathcal{X}}_i$ is invariant under $\lambda_i(a)$ for any $a \in \mathcal{L}(\mathcal{X}_i)$. We can check directly that the space $\mathcal{X}'_i$ is also invariant under $\lambda_i(a)$ for any $a \in \mathcal{L}(\mathcal{X}_i)$ by the definition of λ_i . Hence the result follows. $\square$

The preceding result implies the next corollary.

Corollary 2.3. The map $\beta_i : \mathcal{L}(\mathcal{X}_i) \rightarrow \mathcal{L}(\mathcal{X})$ is a homomorphism.

Definition 2.4. A B -valued probability space is a pair $(\mathcal{A}, \mathbb{E})$ consisting of an algebra $\mathcal{A}$ over B and an B - B -bimodule map $\mathbb{E} : \mathcal{A} \rightarrow B$, i.e. a linear map such that

$$\mathbb{E}(b_1 a b_2) = b_1 \mathbb{E}(a) b_2$$

for all $a \in \mathcal{A}$ and $b_1, b_2 \in B$.

Definition 2.5. Let $(\mathcal{A}, \mathbb{E})$ be a B -valued probability space. A family of B -faces of $\mathcal{A}$ is a family $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ of (not necessarily unital) subalgebras of $\mathcal{A}$ such that C_i, D_i are B - B -bimodules for each $i \in \mathcal{I}$. The family of B -faces $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ is said to be free-Boolean with amalgamation over B if

- $\{C_i | i \in \mathcal{I}\}$ are unital algebras
- there are B - B -bimodules with specified projections $\{(\mathcal{X}_i, \mathring{\mathcal{X}}_i, p_i)\}_{i \in \mathcal{I}}$ such that there are unital homomorphisms $\gamma_i : C_i \rightarrow \mathcal{L}(\mathcal{X}_i)$, (not necessarily unital) homomorphisms $\delta_i : D_i \rightarrow \mathcal{L}(\mathcal{X}_i)$,
- Let $(\mathcal{X}, \mathring{\mathcal{X}}, p)$ be the reduce free product of $\{(\mathcal{X}_i, \mathring{\mathcal{X}}_i, p_i)\}_{i \in \mathcal{I}}$, so that the joint distribution of the family $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ in $(\mathcal{A}, \mathbb{E})$ is equal to the joint distribution of the family of operators $\{(\lambda_i(\gamma_i(C_i)), \beta_i(\delta_i(D_i)))\}_{i \in \mathcal{I}}$ in the probability space $(\mathcal{L}(\mathcal{X}), \mathbb{E}_{\mathcal{L}(\mathcal{X})})$. That is,

$$\mathbb{E}_{\mathcal{L}(\mathcal{X})}(\lambda_{i_1}(\gamma_{i_1}(c_1)) \beta_{i_1}(\delta_{i_1}(d_1)) \cdots \lambda_{i_n}(\gamma_{i_n}(c_n)) \beta_{i_n}(\delta_{i_n}(d_n))) = \mathbb{E}(c_1 d_1 \cdots c_n d_n),$$

where $c_k \in C_{i_k}, d_k \in D_{i_k}, 1 \leq k \leq n$.

3. INTERVAL-NONCROSSING PARTITIONS

In this section, we review some combinatorial tools which will be used

to define operator-valued free-Boolean cumulants. We give a characterization of free-Boolean independence with amalgamation thereby generalizes results in [8] to the operator-valued framework. In noncommutative probability theory, non-crossing partitions are used in the combinatorics of free probability and the interval partitions are used in the combinatorics of Boolean independence. It turns out the partitions used in the combinatorics of free-Boolean independence are so-called interval-noncrossing partitions introduced in [8]. All results without proof in this section are taken from [8].

3.1. Interval-noncrossing partitions. Throughout this section, we let $n \in \mathbb{N}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$ and $\epsilon : \{1, \dots, n\} \rightarrow \mathcal{I}$, for some fixed index set $\mathcal{I}$. We will denote by $[n]$ the set $\{1, \dots, n\}$ for $n \in \mathbb{N}$.

Definition 3.1. Let S be a linearly ordered set. A partition π of the set S consists of a collection disjoint, nonempty sets $\{V_1, \dots, V_p\}$ whose union is S . The sets $V_1, \dots, V_p$ are called the blocks of π . Given $v_1, v_2 \in S$, we write $v_1 \sim v_2$ if the two elements v_1, v_2 are in the same block.

1. A partition π is called *noncrossing* if there is no quadruple (v_1, v_2, w_1, w_2) such that $v_1 < w_1 < v_2 < w_2$, $v_1, v_2 \in V_s$, and $w_1, w_2 \in V_t$, where V_s, V_t are two disjoint blocks of π . The set of all noncrossing partitions of $[n]$ will be denoted by $NC(n)$.
2. A block V_s of π is called *interval* if for any $v_1, v_2 \in V_s$ and $v_1 < w < v_2$, we have $w \in V_s$. A partition $\pi = \{V_1, \dots, V_p\}$ is called an *interval* partition if all blocks V_s are interval blocks.
3. A block V_s of a partition π is said to be *inner* if there is another block $V_t \in \pi$ and $v_1, v_2 \in V_t$ such that $v_1 < w < v_2$ for all $w \in V_s$. A block is *outer* if it is not inner.
4. Let $\epsilon : [n] \rightarrow \mathcal{I}$. We denote by $\ker(\epsilon)$ the partition whose blocks are the sets $\omega^{-1}(i), i \in \mathcal{I}$.
5. Given two partitions σ and π , we say $\sigma \leq \pi$ if each block of σ is contained in a block of π . This relation is called the *reversed refinement order*.
6. We denote by 0_n the partition of $[n]$ consists of n blocks and by 1_n the partition of $[n]$ consists of exactly one block.

Definition 3.2. Let $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$. A partition π of $[n]$ is said to be *interval-noncrossing* with respect to χ if π is noncrossing, and v_1, v_2, w are in the same block whenever $v_1 < w < v_2$, $v_1 \sim v_2$ and $\chi(w) = \mathcal{B}$. We denote by $INC(\chi)$ the set of all interval-noncrossing partitions of the set $\{1, 2, \dots, n\}$ with respect to χ .

Remark 3.3. The set $INC(\chi)$ does not depend on the value of χ at 1 and n . In particular, when $\chi^{-1}(\mathcal{B}) \cap [2, n-1] = \emptyset$, we have $INC(\chi) = NC(n)$.

For example, let $\chi : \{1, \dots, 7\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$ such that $\chi^{-1}(\mathcal{B}) = \{2, 6, 7\}$. Given two noncrossing $\pi_1 = \{\{1, 4\}, \{2\}, \{3\}, \{5, 7\}, \{6\}\}$ and $\pi_2 = \{\{1, 2, 4\}, \{3\}, \{5, 6, 7\}\}$ of the set $\{1, \dots, 7\}$, then $\pi_1 \notin INC(\chi)$ and $\pi_2 \in INC(\chi)$. In pictures below, we use " $\bullet$ " to denote elements in $\chi^{-1}(\mathcal{F})$

and "o" to denote elements in $\chi^{-1}(\mathcal{B})$.



Diagram of π_1 .




Diagram of π_2 .

Assume now $\chi^{-1}(\mathcal{B}) \cap [2, n-1] = \{l_1 < \dots < l_{m-1}\}$ and set $l_0 = 0, l_m = n$. We denote by $[l_i, l_{i+1}]$ the interval $\{l_i, l_i + 1, \dots, l_{i+1}\}$. For each $i = 1, \dots, m$, we denote by $\alpha_i(\pi)$ the restriction of π to the interval $[l_{i-1}, l_i]$. We also denote by $\alpha'(\pi)$ the restriction of π to the interval $[l_1, n]$ and χ' the restriction of χ to the interval $[l_1, n]$. Note that each $\alpha_i(\pi)$ can be any noncrossing partition of the set $[l_{i-1}, l_i]$, since there is no $l_{i-1} < w < l_i$ such that $\chi(w) = \mathcal{B}$.

Proposition 3.4. Let $\alpha'_1 : INC(\chi) \rightarrow NC([1, l_1]) \times INC(\chi')$ be defined by

$$\alpha'_1(\pi) = (\alpha_1(\pi), \alpha'(\pi))$$

and $\alpha : INC(\chi) \rightarrow NC([1, l_1]) \times NC([l_1, l_2]) \times \dots \times NC([l_{m-1}, n])$ be defined by

$$\alpha(\pi) = (\alpha_1(\pi), \dots, \alpha_m(\pi)).$$

Then α'_1 and α are isomorphisms of partial ordered sets. The set $INC(\chi)$ is a lattice with respect to the reverse refinement order $\leq$ on partitions.

We provide pictures below to illustrate the preceding proposition. Let $n = 10$, $\chi^{-1}(\mathcal{B}) = \{1, 3, 7, 8, 9, 10\}$ and $\pi = \{\{1, 3, 4, 7\}, \{2\}, \{5, 6\}, \{9, 8\}, \{10\}\}$ which is an interval-noncrossing of the set $\{1, 2, \dots, 10\}$ with respect to χ as shown in the following diagram.





Diagram of π .

In the above diagram, $l_0 = 1, l_1 = 3, l_2 = 7, l_3 = 8, l_4 = 9, l_5 = 10$. Therefore, $\alpha_1(\pi) = \{\{1, 3\}, \{2\}\}$, $\alpha_2(\pi) = \{\{3, 4, 7\}, \{5, 6\}\}$, $\alpha_3(\pi) = \{\{7\}, \{8\}\}$, $\alpha_4(\pi) = \{\{8, 9\}\}$ and $\alpha_5(\pi) = \{\{9\}, \{10\}\}$ are illustrated in the following diagrams:



Proposition 3.5. Let $\pi = \{V_1, \dots, V_p\} \in INC(\chi)$ and let $\sigma \in NC(n)$ such that $\sigma \leq \pi$, i.e. each block of σ is contained in a block of π . Then $\sigma \in INC(\chi)$ if and only if $\sigma|_{V_s} \in INC(\chi|_{V_s})$ for all $s = 1, \dots, p$.

Proposition 3.6. Let $\pi = \{V_1, \dots, V_p\} \in INC(\chi)$. Denote by $[0_n, \pi]$ the set of all $\sigma \in INC(\chi)$ such that $\sigma \leq \pi$. Then

$$[0_n, \pi] \cong INC(\chi|_{V_1}) \times \dots \times INC(\chi|_{V_p}).$$

3.2. Möbius functions on interval-noncrossing partitions. One can define the convolution for functions on the lattice following the standard procedure for partially ordered sets (see [14]). Once the map χ is fixed, the lattice structure of $INC(\chi)$ caputred from the lattice of the product of noncrossing partitions according to the natural isomorphism described in Proposition 3.4.

Let $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$. Given two complex-valued functions defined on the set $\{(\sigma, \pi) | \sigma, \pi \in INC(\chi), \sigma \leq \pi\}$. The convolution of f and g is given by

$$f * g(\sigma, \pi) = \sum_{\substack{\rho \in INC(\chi) \\ \sigma \leq \rho \leq \pi}} f(\sigma, \rho) g(\rho, \pi).$$

The delta function defined as follows:

$$\delta_{INC}(\sigma, \pi) = \begin{cases} 1, & \text{if } \sigma = \pi, \\ 0, & \text{otherwise.} \end{cases}$$

We then define the zeta function by

$$\zeta_{INC}(\sigma, \pi) = \begin{cases} 1, & \text{if } \sigma \leq \pi, \\ 0, & \text{otherwise.} \end{cases}$$

and the Möbius function μ_{INC} is the inverse of the zeta function in the following sense:

$$\mu_{INC} * \zeta_{INC} = \zeta_{INC} * \mu_{INC} = \delta_{INC}.$$

We will use the following product formula in [7, Section 6].

Proposition 3.7. Let $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$, $\pi = \{V_1, \dots, V_p\} \in INC(\chi)$ and $\sigma \in INC(\chi)$ such that $\sigma \leq \pi$. Suppose that $l_0 = 1, l_m = n$ and $\chi^{-1}(\mathcal{B}) \cap [2, n-1] = \{l_1 < \dots < l_{m-1}\}$, then

$$\begin{aligned} \mu_{INC(\chi)}(\sigma, \pi) &= \prod_{i=1}^m \mu_{INC(\chi|_{V_i})}(\sigma_i, \pi_i) \\ &= \prod_{s=1}^p \mu_{INC}(\sigma|_{V_s}, 1_{V_s}) \\ &= \prod_{i=1}^m \prod_{s=1}^p \mu_{NC}(\sigma_i|_{\tilde{\alpha}_i(V_s)}, 1_{\tilde{\alpha}_i(V_s)}), \end{aligned}$$

where $\sigma_i = \alpha_i(\sigma)$, $\pi_i = \alpha_i(\pi)$, $1 \leq i \leq m$ and $\tilde{\alpha}_i(V_s)$ is the restriction of V_s to the set $[l_{i-1}, l_i]$.

Corollary 3.8. Let $\pi \in INC(\chi)$ and $V \in \pi$. Denote by $[0_n, \pi] = \{\sigma \in INC(\chi) : 0_n \leq \sigma \leq \pi\}$, and $V' = \{1, \dots, n\} \setminus V$. Then,

$$[0_n, \pi] \cong INC(\chi|_V) \times INC(\chi|_{V'}).$$

In particular, we have

$$\mu_{INC}(\sigma, \pi) = \mu_{INC}(\sigma|_V, 1_V) \mu_{INC}(\sigma|_{V'}, \pi|_{V'})$$

for $\sigma \in INC(\chi)$ and $\sigma \leq \pi$.

4. VANISHING CUMULANTS CONDITION FOR FREE-BOOLEAN INDEPENDENCE

In this section, we introduce the notion of operator-valued free-Boolean cumulants for pairs of random variables and give an alternative characterization of free-Boolean independence by using the free-Boolean cumulants.

4.1. Free-Boolean cumulants. Let $(\mathcal{A}, \mathbb{E})$ be a B -valued probability space. Let $\Phi^{(n)}$ be the n - B -linear map from $\underbrace{\mathcal{A} \otimes_B \cdots \otimes_B \mathcal{A}}_{n \text{ times}}$ to B defined as

$$\Phi^{(n)}(a_1, \dots, a_n) = \mathbb{E}(a_1 \cdots a_n).$$

Then, for each noncrossing partition $\pi \in NC(n)$, we can write $\pi = \pi_1 \cup \{V\}$, where $V = (l+1, l+2, \dots, l+s)$ is an interval block of π and $\pi_1 = \pi \setminus \{V\}$. We define an n - B -linear map $\Phi_\pi : \underbrace{\mathcal{A} \otimes_B \cdots \otimes_B \mathcal{A}}_{n \text{ times}} \rightarrow B$ recursively as follows:

$$(1) \quad \Phi_\pi(a_1, \dots, a_n) = \Phi_{\pi_1}(a_1, \dots, a_l, \Phi^{(s)}(a_{l+1}, \dots, a_{l+s})a_{l+s+1}, \dots, a_n).$$

For example, let $\pi = \{\{1, 5, 8\}, \{2, 3, 4\}, \{6, 7\}\}$ be a noncrossing partition of $\{1, \dots, 8\}$. Then,

$$\Phi_\pi(a_1, \dots, a_8) = \Phi_{\{\{1,5,8\}, \{6,7\}\}}(a_1, \mathbb{E}(a_2 a_3 a_4) a_5, a_6, a_7, a_8) = \mathbb{E}(a_1 \mathbb{E}(a_2 a_3 a_4) a_5 \mathbb{E}(a_6 a_7) a_8).$$

Definition 4.1. Given any $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$, $\pi \in INC(\chi)$ and a tuple of elements $(a_1, \dots, a_n)$ in $(\mathcal{A}, \mathbb{E})$, the free-Boolean cumulant $\kappa_{\chi, \pi}$ is an n - B -linear map defined as follows:

$$\kappa_{\chi, \pi}(a_1, \dots, a_n) = \sum_{\substack{\sigma \leq \pi \\ \sigma \in INC(\chi)}} \mu_{INC}(\sigma, \pi) \Phi_\sigma(a_1, \dots, a_n).$$

We start to show that the operator-valued free-Boolean cumulants have the following multiplicative property.

Theorem 4.2. Let $\pi \in INC(\chi)$ and $a_1, \dots, a_n$ be noncommutative random variables in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Suppose that $V = (l+1, l+2, \dots, l+s)$ is an interval block of π , then

$$\begin{aligned} \kappa_{\chi, \pi}(a_1, \dots, a_n) &= \kappa_{\chi|_{V'}, \pi|_{V'}}(a_1, \dots, a_l, \kappa_{\chi|_V, 1_V}(a_{l+1}, \dots, a_{l+s})a_{l+s+1}, \dots, a_n) \\ &= \kappa_{\chi|_{V'}, \pi|_{V'}}(a_1, \dots, (a_l \kappa_{\chi|_V, 1_V}(a_{l+1}, \dots, a_{l+s})), a_{l+s+1}, \dots, a_n), \end{aligned}$$

where $V' = \{1, \dots, n\} \setminus V$.

Proof. For any $\sigma \leq \pi$, $\sigma \in INC(\chi)$, one can decompose it into a union of two interval-noncrossing partitions $\sigma = \sigma_1 \cup \sigma_2$, where $\sigma_1 \leq \pi|_{V'}$, $\sigma_1 \in INC(\chi|_{V'})$ and $\sigma_2 \leq \pi|_V = 1_V$, $\sigma_2 \in INC(\chi|_V)$. By Proposition 3.7 and Corollary 3.8, we have

$$\begin{aligned}
\kappa_{\chi, \pi}(a_1, \dots, a_n) &= \sum_{\substack{\sigma \leq \pi \\ \sigma \in INC(\chi)}} \mu_{INC}(\sigma, \pi) \Phi_{\sigma}(a_1, \dots, a_n) \\
&= \sum_{\substack{\sigma \leq \pi \\ \sigma \in INC(\chi)}} \mu_{INC}(\sigma, \pi) \Phi_{\sigma|_{V'}}(a_1, \dots, a_l, \Phi_{\sigma|_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n) \\
&= \sum_{\substack{\sigma \leq \pi \\ \sigma \in INC(\chi)}} \mu_{INC}(\sigma|_{V'}, \pi|_{V'}) \mu_{INC}(\sigma|_V, 1_V) \Phi_{\sigma|_{V'}}(a_1, \dots, a_l, \Phi_{\sigma|_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n) \\
&= \sum_{\substack{\sigma_1 \leq \pi|_{V'} \\ \sigma_1 \in INC(\chi|_{V'}) \\ \sigma_2 \in INC(\chi|_V)}} \mu_{INC}(\sigma_1, \pi|_{V'}) \mu_{INC}(\sigma_2, 1_V) \Phi_{\sigma|_{V'}}(a_1, \dots, a_l, \Phi_{\sigma|_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n) \\
&= \sum_{\substack{\sigma_1 \leq \pi|_{V'} \\ \sigma_1 \in INC(\chi|_{V'})}} \mu_{INC}(\sigma_1, \pi|_{V'}) \Phi_{\sigma|_{V'}}(a_1, \dots, a_l, \\
&\quad \sum_{\sigma_2 \in INC(\chi|_V)} \mu_{INC}(\sigma_2, 1_V) \Phi_{\sigma|_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n) \\
&= \sum_{\substack{\sigma_1 \leq \pi|_{V'} \\ \sigma_1 \in INC(\chi|_{V'})}} \mu_{INC}(\sigma_1, \pi|_{V'}) \Phi_{\sigma|_{V'}}(a_1, \dots, a_l, \kappa_{\chi|_V, 1_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n) \\
&= \kappa_{\chi|_{V'}, \pi|_{V'}}(a_1, \dots, a_l, \kappa_{\chi|_V, 1_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n).
\end{aligned}$$

The other part follows from the bi-module property. This finishes the proof. $\square$

The preceding theorem shows that $\kappa_{\chi, \pi}(a_1, \dots, a_n)$ is completely determined by cumulant functionals of the form $\kappa_{\chi', 1_{[m]}}$ for $m \in \mathbb{N}$ and $\chi' : \{1, \dots, m\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$.

Definition 4.3. Let $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ be a family of pairs of B -faces of $\mathcal{A}$ in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. We say that the family $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ is combinatorially free-Boolean independent with amalgamation over B if

$$\kappa_{\chi, 1_n}(a_1, \dots, a_n) = 0$$

whenever $\omega : \{1, \dots, n\} \rightarrow \mathcal{I}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$, $a_k \in \mathcal{A}_{\omega(k), \chi(k)}$ and ω is not a constant.

Proposition 4.4. Let $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ be a family of pairs of B -faces in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Then $\kappa_{\chi, 1_n}$ has the following additivity property:

$$\kappa_{\chi, 1_n}(a_{1,1} + a_{2,1}, \dots, a_{1,n} + a_{2,n}) = \kappa_{\chi, 1_n}(a_{1,1}, \dots, a_{1,n}) + \kappa_{\chi, 1_n}(a_{2,1}, \dots, a_{2,n})$$

whenever $\omega_1, \omega_2 : [n] = \{1, \dots, n\} \rightarrow \mathcal{I}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$, $a_{1,k} \in \mathcal{A}_{\omega_1(k), \chi(k)}$, $a_{2,k} \in \mathcal{A}_{\omega_2(k), \chi(k)}$ and $\omega_1([n]) \cap \omega_2([n]) = \emptyset$.

Proof. By a direct calculation, we have

$$\kappa_{\chi, 1_n}(a_{1,1} + a_{2,1}, \dots, a_{1,n} + a_{2,n}) = \sum_{i_1, \dots, i_n \in \{1, 2\}} \kappa_{\chi, 1_n}(a_{i_1, 1}, \dots, a_{i_n, n}).$$

Since $\{(\mathcal{A}_{i,\mathcal{F}}, \mathcal{A}_{i,\mathcal{B}})\}_{i \in I}$ are combinatorially free-Boolean independent, by Definition 4.3, we have

$$\kappa_{\chi, 1_n}(a_{i_1, 1}, \dots, a_{i_n, n}) = 0$$

if $i_j \neq i_k$ for some $j, k \in \{1, \dots, n\}$. The result follows. $\square$

Proposition 4.5. Let $\{(\mathcal{A}_{i,\mathcal{F}}, \mathcal{A}_{i,\mathcal{B}})\}_{i \in \mathcal{I}}$ be a combinatorially free-Boolean independent family of pairs of B -faces in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Assume that $\pi = \{V_1, \dots, V_p\} \in INC(\chi)$. Then

$$\kappa_{\chi, \pi}(a_1, \dots, a_n) = 0$$

whenever $\omega : \{1, \dots, n\} \rightarrow \mathcal{I}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$, $a_k \in \mathcal{A}_{\omega(k), \chi(k)}$ and ω is not a constant on a block W of π .

Proof. We prove the statement by induction on the number of blocks of π .

When $p = 1$, then the statement follows from Definition 4.3. Suppose now that $p > 1$, let $V = (l+1, l+2, \dots, l+s)$ be an interval block of π . By Proposition 4.2, we have

$$\kappa_{\chi, \pi}(a_1, \dots, a_n) = \kappa_{\chi|_{V'}, \pi|_{V'}}(a_1, \dots, a_l, \kappa_{\chi|_V, 1_V}(a_{l+1}, \dots, a_{l+s}) a_{l+s+1}, \dots, a_n),$$

where $V' = \{1, \dots, n\} \setminus V$. If ω is not a constant on V , then $\kappa_{\chi, \pi}(a_1, \dots, a_n) = 0$. Otherwise, $\omega|_{V'}$ is not a constant on a block of $\pi|_{V'}$. The statement follows from an induction argument. $\square$

4.2. Free-Boolean independence is equivalent to combinatorially free-Boolean independence. In this subsection, we will prove that free-Boolean independence defined in Definition 2.5 is equivalent to the combinatorially free-Boolean independence given in Definition 4.3. We will show that mixed moments are uniquely determined by lower order mixed moments in the same way for both free-Boolean independence and combinatorially free-Boolean independence.

The proof for following result is essentially the same as the proof of in [11, Proposition 10.6] in free probability context and we thus leave the details to the reader. Applying Theorem 4.2, we have the following result.

Lemma 4.6. Let $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$ and $a_1, \dots, a_n$ be noncommutative random variables in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Then

$$\mathbb{E}(a_1 \cdots a_n) = \sum_{\pi \in INC(\chi)} \kappa_{\chi, \pi}(a_1 \cdots a_n).$$

For combinatorially free-Boolean independent random variables, we have the following result.

Lemma 4.7. Let $\{(\mathcal{A}_{i,\mathcal{F}}, \mathcal{A}_{i,\mathcal{B}})\}_{i \in \mathcal{I}}$ be a family of combinatorially free-Boolean independent pairs of B -faces in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Assume that $a_k \in \mathcal{A}_{\omega(k), \chi(k)}$, where $\omega : \{1, \dots, n\} \rightarrow \mathcal{I}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$. Let $\epsilon = \ker(\omega)$. Then,

$$(\star) \quad \mathbb{E}(a_1 \cdots a_n) = \sum_{\sigma \in INC(\chi)} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_{\sigma}(a_1 \cdots a_n).$$

Proof. By Lemma 4.6, we have

$$\mathbb{E}(a_1 \cdots a_n) = \sum_{\pi \in INC(\chi)} \kappa_{\chi, \pi}(a_1 \cdots a_n).$$

For each $\pi \in INC(\chi)$, write its blocks as $\pi = \{V_1, \dots, V_p\}$. Since $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ are combinatorially free-Boolean independent, by Lemma 4.5, we have

$$\kappa_{\chi, \pi}(a_1 \cdots a_n) = 0,$$

if ω is not a constant on some block V_s of π . In other words, $\kappa_{\chi, \pi}(a_1, \dots, a_n) \neq 0$ only if ω is a constant on V_s for all s , which implies that V_s is contained in a block of ϵ for all s , i.e., $\pi \leq \epsilon$. Therefore, we have

$$\begin{aligned} \mathbb{E}(a_1 \cdots a_n) &= \sum_{\pi \in INC(\chi), \pi \leq \epsilon} \kappa_{\chi, \pi}(a_1, \dots, a_n) \\ &= \sum_{\pi \in INC(\chi), \pi \leq \epsilon} \left(\sum_{\substack{\sigma \in INC(\chi) \\ \sigma \leq \pi}} \mu_{INC}(\sigma, \pi) \Phi_{\sigma}(a_1, \dots, a_n) \right) \\ &= \sum_{\sigma \in INC(\chi)} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_{\sigma}(a_1, \dots, a_n). \end{aligned}$$

This finishes the proof. $\square$

We now turn to consider the case that the family $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ is free-Boolean independent in $(\mathcal{A}, B, \mathbb{E})$ in the sense of Definition 2.5. In what follows, we assume that $a_k \in \mathcal{A}_{\omega(k), \chi(k)}$, where $\omega : \{1, \dots, n\} \rightarrow \mathcal{I}$, $\chi : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$. Let $\epsilon = \ker(\omega)$, the kernel of ω . Recall that $\chi^{-1}(\mathcal{B}) \cap [2, n-1] = \{l_1, \dots, l_{m-1}\}$. Let χ_1 (or ϵ_1) be the restriction of χ (or ϵ) to $\{1, \dots, l_1\}$ respectively. Let χ'_1 (or ϵ'_1) be the restriction of χ (or ϵ) to the interval $\{l_1, \dots, n\}$ respectively. We need to show that the mixed moments $\mathbb{E}(a_1 \cdots a_n)$ can be determined in the same way as in Lemma 4.7.

To this end, it is enough to consider the case that $\mathcal{A} = \mathcal{L}(\mathcal{X})$, $\mathcal{A}_{i, \mathcal{F}} = \lambda_i(\mathcal{L}(\mathcal{X}_i))$ and $\mathcal{A}_{i, \mathcal{B}} = P_i \lambda_i(\mathcal{L}(\mathcal{X}_i)) P_i$, where $\{(\mathcal{X}_i, \mathcal{X}_i, p_i)\}_{i \in \mathcal{I}}$ is a family of vector spaces with specified vectors and $(\mathcal{X}, \mathcal{X}, p)$ is the reduced free product of them.

We will prove the mixed moments formula (★) in Lemma 4.7 by induction on the number of elements of $\chi^{-1}(\mathcal{B}) \cap [2, n-1]$.

Lemma 4.8. *If $\chi(n) = \mathcal{B}$, then there exists an operator $T \in \mathcal{A}_{\omega(n), \mathcal{F}}$ such that*

$$\mathbb{E}(a_1 \cdots a_n) = \mathbb{E}(a_1 \cdots a_{n-1} T).$$

Proof. If $n \in \chi^{-1}(\mathcal{B})$, then $a_n \in \mathcal{A}_{\omega(n), \mathcal{B}} = P_{\omega(n)} \lambda_{\omega(n)}(\mathcal{L}(\mathcal{X}_{\omega(n)})) P_{\omega(n)}$. Assume that $a_n = P_{\omega(n)} T P_{\omega(n)}$ for some $T \in \lambda_{\omega(n)}(\mathcal{L}(\mathcal{X}_{\omega(n)}))$. Then

$$a_1 \cdots a_n 1_B = a_1 \cdots a_{n-1} P_{\omega(n)} T P_{\omega(n)} 1_B = a_1 \cdots T 1_B = a_1 \cdots a_{n-1} T 1_B$$

since $T 1_B \in P_{\omega(n)} \mathcal{X}$. Thus, the mixed moments are the same if we replace a_n by the element $T \in \lambda_{\omega(n)}(\mathcal{L}(\mathcal{X}_{\omega(n)}))$. $\square$

Lemma 4.9. *If $\chi(1) = \mathcal{B}$, then there exists an operator $T \in \mathcal{A}_{\omega(n), \mathcal{F}}$ such that*

$$\mathbb{E}(a_1 \cdots a_n) = \mathbb{E}(Ta_2 \cdots a_n).$$

Proof. If $1 \in \chi^{-1}(\mathcal{B})$, then $a_1 \in \mathcal{A}_{\omega(1), \mathcal{B}} = P_{\omega(1)}\lambda_{\omega(1)}(\mathcal{L}(\mathcal{X}_{\omega(1)}))P_{\omega(1)}$. Assume that $a_1 = P_{\omega(1)}TP_{\omega(1)}$ for some $T \in \lambda_{\omega(1)}(\mathcal{L}(\mathcal{X}_{\omega(1)}))$. Recall that p is the projection $p : \mathcal{X} \rightarrow B$. Notice that $pP_{\omega(1)} = p$ and

$$\begin{aligned} \mathbb{E}(a_1 \cdots a_n) &= pP_{\omega(1)}TP_{\omega(1)}a_2 \cdots a_n1_B \\ &= pP_{\omega(1)}TP_{\omega(1)}a_2 \cdots a_n1_B \\ &= pTP_{\omega(1)}a_2 \cdots a_n1_B \\ &= pTa_2 \cdots a_n1_B - pT(1_{\mathcal{X}} - P_{\omega(1)})a_2 \cdots a_n1_B, \end{aligned}$$

where $1_{\mathcal{X}}$ is the identity operator in $\mathcal{L}(\mathcal{X})$. Notice that

$$(1_{\mathcal{X}} - P_{\omega(1)})a_2 \cdots a_n1_B \in \bigoplus_{i \neq \omega(1)} \mathring{\mathcal{X}}_i \oplus \bigoplus_{n \geq 2} \left(\bigoplus_{i_1 \neq i_2 \neq \cdots \neq i_n} \mathring{\mathcal{X}}_{i_1} \otimes_B \cdots \otimes_B \mathring{\mathcal{X}}_{i_n} \right)$$

and

$$\begin{aligned} &\bigoplus_{i \neq \omega(1)} \mathring{\mathcal{X}}_i \oplus \bigoplus_{n \geq 2} \left(\bigoplus_{i_1 \neq i_2 \neq \cdots \neq i_n} \mathring{\mathcal{X}}_{i_1} \otimes_B \cdots \otimes_B \mathring{\mathcal{X}}_{i_n} \right) \\ &= V_{\omega(1)} \left(\mathcal{X}_{\omega(1)} \otimes_B \left(\bigoplus_{\substack{i_1 \neq i_2 \neq \cdots \neq i_n \\ i_1 \neq \omega(1), n \geq 1}} \mathring{\mathcal{X}}_{i_1} \otimes_B \cdots \otimes_B \mathring{\mathcal{X}}_{i_n} \right) \right). \end{aligned}$$

Therefore, $\bigoplus_{i \neq \omega(1)} \mathring{\mathcal{X}}_i \oplus \bigoplus_{n \geq 2} \left(\bigoplus_{i_1 \neq i_2 \neq \cdots \neq i_n} \mathring{\mathcal{X}}_{i_1} \otimes_B \cdots \otimes_B \mathring{\mathcal{X}}_{i_n} \right)$ is an invariant subspace of T and

$$pT(I_{\mathcal{X}} - P_{\omega(1)})a_2 \cdots a_n1_B = p(I_{\mathcal{X}} - P_{\omega(1)})T(I_{\mathcal{X}} - P_{\omega(1)})a_2 \cdots a_n1_B = 0,$$

where the last equality follows from the fact that $p(I_{\mathcal{X}} - P_{\omega(1)}) = 0$. We thus proved that the mixed moments $\mathbb{E}(a_1 \cdots a_n)$ will be the same if we replace a_1 by the element $T \in \lambda_{\omega(1)}(\mathcal{L}(\mathcal{X}_{\omega(1)}))$. $\square$

We start with the following result.

Lemma 4.10. *When $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| = 0$, we have*

$$\mathbb{E}(a_1 \cdots a_n) = \sum_{\sigma \in INC(\chi)} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_{\sigma}(a_1 \cdots a_n).$$

Proof. By Lemma 4.8 and 4.9, if $a_1 \in \mathcal{A}_{\omega(1), \mathcal{B}}$ or $a_n \in \mathcal{A}_{\omega(n), \mathcal{B}}$, we may replace a_1 by $T_1 \in \mathcal{A}_{\omega(1), \mathcal{F}}$ and a_n by $T_2 \in \mathcal{A}_{\omega(n), \mathcal{F}}$, we still have

$$\Phi(a_1 \cdots a_n) = \Phi(T_1 a_2 \cdots a_{n-1} T_2).$$

Hence, when $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| = 0$, we may assume that $T_1, a_2, \dots, a_{n-1}, T_2$ are from the left faces of algebras $\mathcal{A}_{\omega(k), \mathcal{F}}$. Notice that the family $\{(\mathcal{A}_{i, \mathcal{F}})\}_{i \in \mathcal{I}}$ is freely independent with amalgamation in $(\mathcal{A}, \mathbb{E})$ (see [11, 16]), we have

$$\begin{aligned}
\mathbb{E}(a_1 \cdots a_n) &= \mathbb{E}(T_1 a_2 \cdots a_{n-1} T_2) \\
&= \sum_{\pi \in NC(n), \pi \leq \epsilon} \kappa_\pi(T_1, a_2, \dots, a_{n-1}, T_2) \\
&= \sum_{\sigma \in NC(n)} \left(\sum_{\substack{\pi \in NC(n) \\ \sigma \leq \pi \leq \epsilon}} \mu(\sigma, \pi) \right) \Phi_\sigma(T_1, a_2, \dots, a_{n-1}, T_2) \\
&= \sum_{\sigma \in INC(\chi)} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_\sigma(T_1, a_2, \dots, a_{n-1}, T_2) \\
&= \sum_{\sigma \in INC(\chi)} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_\sigma(a_1, a_2, \dots, a_{n-1}, a_2),
\end{aligned}$$

where we used the fact that $INC(\chi) = NC(n)$ when $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| = 0$. □

Now, we are now ready to prove our main theorem.

Theorem 4.11. Let $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ be a family of pairs of B -faces in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. The family $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ is free-Boolean independent with amalgamation over B if and only if they are combinatorially free-Boolean independent with amalgamation over B .

Proof. It suffices to show that Equation (★) holds by assuming that $\{(\mathcal{A}_{i, \mathcal{F}}, \mathcal{A}_{i, \mathcal{B}})\}_{i \in \mathcal{I}}$ is free-Boolean independent with amalgamation over B . When $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| = 0$, it is Lemma 4.10. Assume now that Equation (★) in Lemma 4.7 holds whenever $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| \leq m-2$. We shall prove it holds when $|\chi^{-1}(\mathcal{B}) \cap [2, n-1]| = m-1$. Set $\chi^{-1}(\mathcal{B}) = \{l_1 < \dots < l_{m-1}\}$ and $l_0 = 1, l_m = n$.


Let $A_1 = \prod_{i=l_1}^n a_i$. Then $A_1(1_B) \in B \oplus \mathring{\mathcal{X}}_{\omega(l_1)}$. Since the range of A_{l_1} is $B \oplus \mathring{\mathcal{X}}_{\omega(l_1)}$, we can view $A_1 : B \oplus \mathring{\mathcal{X}}_{\omega(l_1)} \rightarrow B \oplus \mathring{\mathcal{X}}_{\omega(l_1)}$ as a linear operator. In this way, A_1 is considered as an element in $\lambda_{\omega(l_1)}(\mathcal{L}(\mathring{\mathcal{X}}_{\omega(l_1)}))$.

Apply the induction for the l_1 -tuple $(a_1, \dots, a_{l_1-1}, A_1)$. Recall that ϵ_1 is the restriction of ϵ to the interval $[1, l_1]$, we have

$$\begin{aligned}
\mathbb{E}(a_1 \cdots a_n) &= \mathbb{E}(a_1 \cdots a_{l_1-1} A_1) \\
(2) \quad &= \sum_{\substack{\sigma_1 \in NC([l_1]) \\ \sigma_1 \leq \epsilon_1}} \left(\sum_{\substack{\pi_1 \in NC([l_1]) \\ \sigma_1 \leq \pi_1 \leq \epsilon_1}} \mu(\sigma_1, \pi_1) \right) \Phi_{\sigma_1}(a_1, \dots, a_{l_1-1} A_1).
\end{aligned}$$

We now fix $\sigma_1 \in NC(l_1), \sigma_1 \leq \epsilon_1$. We shall express $\Phi_{\sigma_1}(a_1, \dots, a_{l_1-1} A_1)$ according to the definition given by (1). We need to know how Φ_{σ_1} is decomposed. To this end, suppose that V is

the block of σ_1 which contains the element l_1 . Denote that $V = \{p_1, p_2, \dots, p_{k_1}\}$, where $p_{k_1} = l_1$. Set $W_1 = [1, p_1 - 1]$, $W_2 = [p_1 + 1, p_2 - 1], \dots, W_{k_1} = [p_{k_1-1} + 1, p_{k_1} - 1] = [p_{k_1-1} + 1, l_1 - 1]$ (W_i will be the empty set if $p_{i-1} + 1 = p_i$), as illustrated in the picture below.



Note that $l_1 \notin W_i$ for all $1 \leq i \leq k_1$, we have

(3)

$$\begin{aligned}
 & \Phi_{\sigma_1}(a_1, \dots, a_{l_1-1}, A_1) \\
 = & \mathbb{E} \left\{ \left(\prod_{k=1}^{k_1-1} [\Phi_{\sigma_1|W_k}(a_1, \dots, a_{l_1-1}, A_1)|_{W_k}] a_{p_k} \right) [\Phi_{\sigma_1|W_{k_1}}(a_1, \dots, a_{l_1-1}, A_1)|_{W_{k_1}}] A_1 \right\} \\
 = & \mathbb{E} \left\{ \left(\prod_{k=1}^{k_1-1} [\Phi_{\sigma_1|W_k}(a_1, \dots, a_{l_1-1})|_{W_k}] a_{p_k} \right) [\Phi_{\sigma_1|W_{k_1}}(a_1, \dots, a_{l_1-1})|_{W_{k_1}}] A_1 \right\} \\
 = & \mathbb{E} \left\{ \left(\prod_{k=1}^{k_1-1} [\Phi_{\sigma_1|W_k}(a_1, \dots, a_{l_1-1})|_{W_k}] a_{p_k} \right) [\Phi_{\sigma_1|W_{k_1}}(a_1, \dots, a_{l_1-1})|_{W_{k_1}}] a_{l_1} \dots a_m \right\}.
 \end{aligned}$$

Denote by $A_2 = \left(\prod_{k=1}^{k_1-1} [\Phi_{\sigma_1|W_k}(a_1, \dots, a_{l_1-1})|_{W_k}] a_{p_k} \right) [\Phi_{\sigma_1|W_{k_1}}(a_1, \dots, a_{l_1-1})|_{W_{k_1}}] a_{l_1}$. Notice that $|\chi^{-1}(\mathcal{B}) \cap [l_1 + 1, n - 1]| = m - 2$. We now apply the induction formula for the tuple $(A_2, a_{l_1+1}, \dots, a_n)$, recall that χ' is the restriction of χ to the interval $[l_1, n]$ and ϵ' is the restriction of ϵ to the interval $[l_1, n]$, we deduce that

$$\begin{aligned}
 & \Phi_{\sigma_1}(a_1, \dots, a_{l_1-1}, A_1) = \mathbb{E}(A_2 a_{l_1+1} a_{l_1+2} \dots a_n) \\
 (4) \quad & = \sum_{\sigma' \in INC(\chi')} \left(\sum_{\substack{\pi' \in INC(\chi') \\ \sigma' \leq \pi' \leq \epsilon'}} \mu_{INC}(\sigma', \pi') \right) \Phi_{\sigma'}(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n).
 \end{aligned}$$


We now fix $\sigma' \in INC(\chi'), \sigma' \leq \epsilon'$. We need to express $\Phi_{\sigma'}(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n)$ according to the definition given in (1). To this end, suppose that V' is the block of σ' which contains the element l_1 . Suppose that $V' = \{q_1, q_2, \dots, q_{k_2}\}$, where $q_1 = l_1$. Let $W'_1 = [l_1, q_2 - 1]$, $W'_2 = [q_2 + 1, q_3 - 1], \dots, W'_{k_2} = [q_{k_2} + 1, n]$ ($W'_j = \emptyset$ if $q_j + 1 = q_{j+1}$), as shown in the picture below



Notice that $l_1 \notin W'_j$ for all $1 \leq j \leq k_2$, we apply the induction assumption to the tuple $(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n)$ to obtain the following:

$$\begin{aligned}
& \Phi_{\sigma'}(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n) \\
&= \mathbb{E} \left\{ A_2 \left[\Phi_{\sigma'|_{W'_1}}(A_2, a_{l_1+1}, \dots, a_n)|_{W'_1} \right] \left(\prod_{k=2}^{k_2} a_{q_k} \left[\Phi_{\sigma'|_{W'_k}}(A_2, a_{l_1+1}, \dots, a_n)|_{W'_k} \right] \right) \right\} \\
(5) \quad &= \mathbb{E} \left\{ A_2 \left[\Phi_{\sigma'|_{W'_1}}(a_{l_1+1}, \dots, a_n)|_{W'_1} \right] \left(\prod_{k=2}^{k_2} a_{q_k} \left[\Phi_{\sigma'|_{W'_k}}(a_{l_1+1}, \dots, a_n)|_{W'_k} \right] \right) \right\} \\
&= \mathbb{E} \left\{ \left(\prod_{k=1}^{k_1-1} \left[\Phi_{\sigma_1|_{W_k}}(a_1, \dots, a_{l_1-1})|_{W_k} \right] a_{p_k} \right) \left[\Phi_{\sigma_1|_{W_{k_1}}}(a_1, \dots, a_{l_1-1})|_{W_{k_1}} \right] \right. \\
&\quad \left. a_{l_1} \left[\Phi_{\sigma'|_{W'_1}}(a_{l_1+1}, \dots, a_n)|_{W'_1} \right] \left(\prod_{k=2}^{k_2} a_{q_k} \left[\Phi_{\sigma'|_{W'_k}}(a_{l_1+1}, \dots, a_n)|_{W'_k} \right] \right) \right\}
\end{aligned}$$

Recall that $\alpha'_1(\pi) := (\alpha_1(\pi), \alpha'(\pi))$ defined in Proposition 3.4. Let $\sigma = \alpha'^{-1}(\sigma_1, \sigma')$. We draw the picture below to show the block $V \in \sigma_1$ and the block $V' \in \sigma_2$ which contain l_1 .



Then,

$$\begin{aligned}
& \Phi_{\sigma'}(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n) \\
&= \mathbb{E} \left\{ \left(\prod_{k=1}^{k_1-1} \left[\Phi_{\sigma_1|_{W_k}}(a_1, \dots, a_{l_1-1})|_{W_k} \right] a_{p_k} \right) \Phi_{\sigma_1|_{W_{k_1}}}(a_1, \dots, a_{l_1-1})|_{W_{k_1}} \right. \\
&\quad \left. a_{l_1} \left[\Phi_{\sigma'|_{W'_1}}(a_{l_1+1}, \dots, a_n)|_{W'_1} \right] \left(\prod_{k=2}^{k_2} a_{q_k} \Phi_{\sigma'|_{W'_k}}(a_{l_1+1}, \dots, a_n)|_{W'_k} \right) \right\} \\
&= \Phi_{\sigma}(a_1, \dots, a_n).
\end{aligned}$$

Putting (2), (3), (4), (5) together, we have

$$\begin{aligned}
& \mathbb{E}(a_1 \cdots a_n) \\
&= \sum_{\substack{\sigma_1 \in NC([l_1]) \\ \sigma_1 \leq \epsilon_1}} \left(\sum_{\substack{\pi_1 \in NC([l_1]) \\ \sigma_1 \leq \pi_1 \leq \epsilon_1}} \mu(\sigma_1, \pi_1) \right) \Phi_{\sigma_1}(a_1, \dots, a_{l_1-1} A_1). \\
&= \sum_{\substack{\sigma_1 \in NC([l_1]) \\ \sigma_1 \leq \epsilon_1}} \left(\sum_{\substack{\pi_1 \in NC([l_1]) \\ \sigma_1 \leq \pi_1 \leq \epsilon_1}} \mu(\sigma_1, \pi_1) \right) \sum_{\sigma' \in INC(\chi')} \left(\sum_{\substack{\pi' \in INC(\chi') \\ \sigma' \leq \pi' \leq \epsilon'}} \mu_{INC}(\sigma', \pi') \right) \Phi_{\sigma'}(A_2, a_{l_1+1}, a_{l_1+2}, \dots, a_n) \\
&= \sum_{\substack{\sigma_1 \in NC([l_1]) \\ \sigma_1 \leq \epsilon_1}} \left(\sum_{\substack{\pi_1 \in NC([l_1]) \\ \sigma_1 \leq \pi_1 \leq \epsilon_1}} \mu(\sigma_1, \pi_1) \right) \sum_{\sigma' \in INC(\chi')} \left(\sum_{\substack{\pi' \in INC(\chi') \\ \sigma' \leq \pi' \leq \epsilon'}} \mu_{INC}(\sigma', \pi') \right) \Phi_{\alpha'^{-1}(\sigma_1, \sigma')}(a_1, \dots, a_n) \\
&= \sum_{\substack{\sigma \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \left(\sum_{\substack{\pi \in INC(\chi) \\ \sigma \leq \pi \leq \epsilon}} \mu_{INC}(\sigma, \pi) \right) \Phi_{\sigma}(a_1, \dots, a_n),
\end{aligned}$$

where we used Corollary 3.8 in the last identity and thus we obtained our desired equation. $\square$

5. LIMIT THEOREMS

Let $A = ((a_i)_{i \in \mathcal{I}}, (a_j)_{j \in \mathcal{J}})$ be a two faced family of noncommutative random variables in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Let $\omega : \{1, \dots, n\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$ and denote by $\chi_\omega : \{1, \dots, n\} \rightarrow \{\mathcal{F}, \mathcal{B}\}$ the map such that $\chi_\omega(k) = \mathcal{B}$ if and only if $\omega(k) \in \mathcal{J}$.

Definition 5.1. A two-faced family $A = ((a_i)_{i \in \mathcal{I}}, (a_j)_{j \in \mathcal{J}})$ in a B -valued probability space $(\mathcal{A}, \mathbb{E})$ is said to have a centered free-Boolean limit if, for all $n \neq 2$,

$$\kappa_{\chi_\omega, 1_n}(a_{\omega(1)}b_1, \dots, a_{\omega(n-1)}b_{n-1}, a_{\omega(n)}) = 0,$$

for all $\omega : \{1, \dots, n\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$ and $b_1, \dots, b_{n-1} \in B$.

The distribution defined by the two faced family A is called an operator-valued free-Boolean Gaussian distribution with covariance $C = (c_{i,j})_{i,j \in \mathcal{I} \sqcup \mathcal{J}}$, where C is defined by $c_{\omega(1), \omega(2)}(b) := \kappa_{\chi_\omega, 1_2}(a_{\omega(1)}, ba_{\omega(2)})$ for all $\omega : \{1, 2\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$ and $b \in B$.

Proposition 5.2. Let $A = ((a_i)_{i \in \mathcal{I}}, (a_j)_{j \in \mathcal{J}})$ be a two faced family of noncommutative random variables in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Let $\omega : \{1, 2\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$. Then,

$$\kappa_{\chi_\omega, 1_2}(a_{\omega(1)}, a_{\omega(2)}) = \mathbb{E}(a_{\omega(1)}a_{\omega(2)}) - \mathbb{E}(a_{\omega(1)})\mathbb{E}(a_{\omega(2)}).$$

Theorem 5.3. Let $\{A_m = ((a_{m,i})_{i \in \mathcal{I}}, (a_{m,j})_{j \in \mathcal{J}})\}_{m=1}^\infty$ be a free-Boolean sequence of families in a B -valued probability space $(\mathcal{A}, \mathbb{E})$ where B is a Banach space, such that

- (1) $\mathbb{E}(a_{m,k}) = 0$ for all $m \in \mathbb{N}$ and $k \in \mathcal{I} \sqcup \mathcal{J}$.
- (2) $\sup_{m \in \mathbb{N}} \|\mathbb{E}(a_{m, \omega(1)}b_1 \cdots a_{m, \omega(n-1)}b_{n-1}a_{m, \omega(n)})\| \leq D_\omega < \infty$ for all $n \in \mathbb{N}$, $\omega : \{1, \dots, n\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$, and $b_1, \dots, b_{n-1} \in B$.
- (3) $\lim_{N \rightarrow \infty} N^{-1} \sum_{1 \leq m \leq N} \mathbb{E}(a_{m, \omega(1)}ba_{m, \omega(2)}) = c_{\omega(1), \omega(2)}(b) \in B$, for all $\omega : \{1, 2\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$ and $b \in B$.

Let $S_{N,k} = N^{-1/2} \sum_{1 \leq m \leq N} a_{m,k}$ for $k \in \mathcal{I} \sqcup \mathcal{J}$ and $S_N = ((S_{N,i})_{i \in \mathcal{I}}, (S_{N,j})_{j \in \mathcal{J}})$. Denote by γ_C the free-Boolean limit distribution in Definition ***, with $C = (c_{i,j})_{i,j \in \mathcal{I} \sqcup \mathcal{J}}$. We have

$$\lim_{N \rightarrow \infty} \mu_{S_N}(P) = \gamma_C(P),$$

for all $P \in \mathbb{C}\langle a_k | k \in \mathcal{I} \sqcup \mathcal{J} \rangle$.

Proof. Since the joint distributions are determined by free-Boolean cumulants uniquely, it is enough to show that

$$\lim_{N \rightarrow \infty} \kappa_{\chi_\omega, 1_n}(S_{N, \omega(1)}b_1, \dots, S_{N, \omega(n-1)}b_{n-1}, S_{N, \omega(n)}) = \kappa_{\chi_\omega, 1_n}(S_{\omega(1)}b_1, \dots, S_{\omega(n-1)}b_{n-1}, S_{\omega(n)}),$$

where the two faced family $S = ((S_i)_{i \in \mathcal{I}}, (S_j)_{j \in \mathcal{J}})$ has a centered B -valued free-Boolean Gaussian distribution with covariance matrix C , for all $n \in \mathbb{N}$, $\omega : \{1, \dots, n\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$, and $b_1, \dots, b_{n-1} \in B$.

By the additivity property of free-Boolean cumulants, we have

$$\begin{aligned} & \kappa_{\chi_{\omega,1n}}(S_{N,\omega(1)}b_1, \dots, S_{N,\omega(n-1)}b_{n-1}, S_{N,\omega(n)}) \\ &= \frac{1}{N^{n/2}} \sum_{1 \leq m \leq N} \kappa_{\chi_{\omega,1n}}(a_{m,\omega(1)}b_1, \dots, a_{m,\omega(n-1)}b_{n-1}, a_{m,\omega(n)}). \end{aligned}$$

Since the free-Boolean cumulant are universal polynomial of mixed moments, we deduce from assumption (2) that

$$\sup_{m \in \mathbb{N}} \|\kappa_{\chi_{\omega,1n}}(a_{m,\omega(1)}b_1, \dots, a_{m,\omega(n-1)}b_{n-1}, a_{m,\omega(n)})\| < \infty$$

and hence

$$\lim_{N \rightarrow \infty} \kappa_{\chi_{\omega,1n}}(S_{N,\omega(1)}b_1, \dots, S_{N,\omega(n-1)}b_{n-1}, S_{N,\omega(n)}) = 0$$

for $n \geq 3$.

As $\kappa_{\chi_{\omega,1n}}(a_{m,\omega(1)}) = \mathbb{E}(a_{m,\omega(1)}) = 0$ for all $m \in \mathbb{N}$ and $\omega : \{1\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$, we have $\kappa_{\chi_{\omega,1n}}(S_{N,\omega(1)}) = 0$ for all $N \in \mathbb{N}$ and $\omega : \{1\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$. Finally, by assumption (3), and Proposition 5.2, we have

$$\begin{aligned} & \kappa_{\chi_{\omega,12}}(S_{N,\omega(1)}b, S_{N,\omega(2)}) \\ &= \frac{1}{N} \sum_{1 \leq m \leq N} \kappa_{\chi_{\omega,12}}(a_{m,\omega(1)}b, a_{m,\omega(2)}) \\ &= \frac{1}{N} \sum_{1 \leq m \leq N} \mathbb{E}(a_{m,\omega(1)}ba_{m,\omega(2)}) \rightarrow C_{\omega(1),\omega(2)}(b), \end{aligned}$$

as $N \rightarrow \infty$, for all $\omega : \{1, 2\} \rightarrow \mathcal{I} \sqcup \mathcal{J}$ and $b \in B$. This finishes the proof. $\square$

6. MOMENT-CONDITIONS FOR FREE-BOOLEAN INDEPENDENCE

Let $\{(\mathcal{X}_i, \mathring{\mathcal{X}}_i, p_i)\}_{i \in \mathcal{I}}$ be B - B -bimodules with specified projectin, and $\{\mathcal{X}, \mathring{\mathcal{X}}, p\}$ be the reduced free product with amalgamation over B . For each $i \in \mathcal{I}$, denote $\mathcal{A}_{i,\mathcal{F}} = \lambda_i(\mathcal{L}(\mathcal{X}_i))$ and $\mathcal{A}_{i,\mathcal{B}} = \beta_i(\mathcal{L}(\mathcal{X}_i)) = P_i \lambda_i(\mathcal{L}(\mathcal{X}_i)) P_i$, where P_i is the projection onto the subspace $B \oplus \mathring{\mathcal{X}}_i$. We also denote by $\mathcal{A}_i$ the algebra generated by $\mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$.

Given a family $\{C_i, D_i\}_{i \in \mathcal{I}}$ of free-Boolean pair of B -faces in a B -valued probability space $(\mathcal{A}, \mathbb{E})$, to study the mixed moments of the family, one can identify C_i with $\lambda_i(\gamma_i(C_i))$ and identify D_i with $\beta_i(\delta_i(D_i))$ following Definition 2.5. In this way, we regard C_i as a subalgebra of $\mathcal{A}_{i,\mathcal{F}}$ and D_i as a subalgebra of $\mathcal{A}_{i,\mathcal{B}}$ throughout this section.

Definition 6.1. Given a set $S_i \subset \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$ and $a_1, \dots, a_m \in S_i$ their product $A = a_1 \cdots a_m$ is called a *simple product* of elements from S_i . It is called a *Boolean product* of elements from S_i if $a_k \in S_i \cap \mathcal{A}_{i,\mathcal{B}}$ for some $1 \leq k \leq m$.

If $A = a_1 \cdots a_m$ is a simple product, but not a Boolean product, then each $a_i \in S_i \cap \mathcal{A}_{i,\mathcal{F}}$. A Boolean product of elements from $\mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$ has a very simple form.

Proposition 6.2. Let $A \in \mathcal{A}_i$ ($i \in \mathcal{I}$) be a Boolean product of elements from $\mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$. Then $A \in \mathcal{A}_{i,\mathcal{B}}$, in particular, $A(\mathcal{X}) \subset \mathcal{X}_i$.

Proof. Write $A = a_1 \cdots a_m$ where $a_k \in \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$ ($1 \leq k \leq m$). For an element $a_k \in \mathcal{A}_{i,\mathcal{B}}$, it can be written as $a_k = P_i a'_k P_i$, where $a'_k \in \mathcal{A}_{k,\mathcal{F}}$ by definition. Hence $A = P_i b_1 \cdots b_m P_i$, where

$$b_k = \begin{cases} a'_k, & \text{if } a_k \in \mathcal{A}_{i,\mathcal{B}} \\ a_k, & \text{if } a_k \in \mathcal{A}_{i,\mathcal{F}} \end{cases}$$

by Proposition 2.2. The assertion follows. $\square$

To state our result in general, from now on, we will let $S_i \subset \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$ for each $i \in \mathcal{I}$. Typically, $S_i = C_i \cup D_i$ or $S_i = \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$.

Lemma 6.3. Let $A_1 \in \mathcal{A}_i$ and $A_2 \in \mathcal{A}_j$ be two simple product of elements from S_i and S_j respectively and $i \neq j$. If A_1 is a Boolean product, then

$$A_1 A_2 1_B = A_1 \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_2).$$

Proof. Write $A_2(1_B) = \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_2) \oplus \mathring{A}_2$, where $\mathring{A}_2 \in \mathring{\mathcal{X}}_j$. As A_1 is a Boolean product, at least one of the factor is from $\mathcal{A}_{i,\mathcal{B}}$, we thus can write $A_1 = a_1 b a_2$, where a_2 is a simple product of elements from $\mathcal{A}_{i,\mathcal{F}}$ and $b \in \mathcal{A}_{i,\mathcal{B}}$. We can express $a_2 = \lambda_i(T)$, where $T \in \mathcal{L}_B(\mathcal{X}_i)$.

Observe that

$$\begin{aligned} \lambda_i(T)(\mathring{A}_2) &= V_i(T \otimes I_{\mathcal{X}(i)}) V_i^{-1} \mathring{A}_2 \\ &= V_i(T \otimes I_{\mathcal{X}(i)})(1_B \otimes \mathring{A}_2) \\ &= V_i[\mathbb{E}_{\mathcal{L}(\mathcal{X}_i)}(T) \otimes \mathring{A}_2 + (T - \mathbb{E}_{\mathcal{L}(\mathcal{X}_i)}(T)) \otimes \mathring{A}_2]. \end{aligned}$$

Hence, $\lambda_i(T)\mathring{A}_2 \in \mathring{\mathcal{X}}_j \oplus (\mathring{\mathcal{X}}_i \otimes \mathring{\mathcal{X}}_j)$. As $b = P_i b P_i$, where P_i is the projection onto $B \oplus \mathring{\mathcal{X}}_i$, we deduce that $b\mathring{A}_2 = 0$. We then have

$$\begin{aligned} A_1 A_2 1_B &= (a_1 b a_2) (\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_2) + \mathring{A}_2) \\ &= (a_1 b a_2) (\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_2)) \\ &= A_1 (\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_2)). \end{aligned}$$

This finishes the proof. $\square$

An application of the preceeding lemman and the bimodule property of expectation $\mathbb{E}_{\mathcal{L}(\mathcal{X})}$ implies the following result.

Corollary 6.4. For $1 \leq i \leq m$, let $A_i \in \mathcal{A}_{k(i)}$ be Boolean product of elements from $\mathcal{A}_{k(i),\mathcal{F}} \cup \mathcal{A}_{k(i),\mathcal{B}}$ and $k(1) \neq k(2) \neq \cdots \neq k(m)$. Then, we have

$$\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_1 \cdots A_m) = \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_1) \cdots \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_m).$$

Lemma 6.5. Let $B \in \mathcal{A}_i$ be a Boolean product of elements from $S_i \subset \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$, then $B(\mathcal{X}) \subset \mathring{\mathcal{X}}_i$.

Proof. Observe that B can be written as $B = a_1 b a_2$, where $b \in \mathcal{A}_{i,\mathcal{B}}$ and a_1, a_2 are simple products of elements from $\mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$. Since $b = P_i b P_i$, where P_i is the projection onto $\mathcal{X}_i = B \oplus \mathring{\mathcal{X}}_i$, we deduce that $b a_2(\mathcal{X}) \subset \mathring{\mathcal{X}}_i$. Hence the assertion follows. $\square$

Lemma 6.6. Let $A_i \in \mathcal{A}_{k(i)} (1 \leq i \leq m)$ be simple products of elements from $S_i \subset \mathcal{A}_{k(i),\mathcal{F}} \cup \mathcal{A}_{k(i),\mathcal{B}}$. If the following conditions hold:

- (1) There exist $1 < l_1 < l_2 < m$ such that $A_{l_1}, A_{l_1+1}, \dots, A_{l_2}$ are not Boolean products.
- (2) A_{l_1-1} and A_{l_2+1} are Boolean products.
- (3) $k(1) \neq k(2) \neq \dots \neq k(m)$.
- (4) $\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_{l_1}) = \dots = \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_{l_2}) = 0$.

Then, the product of operators $A_1 \cdots A_m = 0$.

Proof. Since A_{l_2+1} is a Boolean product, it follows from Lemma 6.5 that

$$A_{l_2+1} \cdots A_m(\mathcal{X}) \subset B \oplus \mathring{\mathcal{X}}_{k(l_2+1)}.$$

As $\mathbb{E}_{\mathcal{L}(X)}(A_{l_1}) = \dots = \mathbb{E}_{\mathcal{L}(X)}(A_{l_2}) = 0$ and each A_j ($l_1 \leq j \leq l_2$) is a simple product of elements from $\mathcal{A}_{k(j),\mathcal{F}}$, in this case, we then have

$$\begin{aligned} A_{l_2}(A_{l_2+1} \cdots A_m)(\mathcal{X}) &\subset \mathring{\mathcal{X}}_{k(l_2)} \otimes (B \oplus \mathring{\mathcal{X}}_{k(l_2+1)}) \\ &\cong \mathring{\mathcal{X}}_{k(l_2)} \oplus (\mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)}). \end{aligned}$$

By induction, we then have

$$\begin{aligned} (6) \quad &(A_{l_1} \cdots A_{l_2})(A_{l_2+1} \cdots A_m)(\mathcal{X}) \\ &\subset (\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)}). \end{aligned}$$

The operator A_{l_1-1} is a Boolean product. We may write $A_{l_1-1} = a_1 b a_2$ where a_2 is a simple product of elements from $\mathcal{A}_{k(l_1-1),\mathcal{F}}$ and $b \in \mathcal{A}_{k(l_1-1),\mathcal{B}}$. It follows that

$$\begin{aligned} a_2(A_{l_1} \cdots A_m)(\mathcal{X}) &\subset \mathcal{X}_{k(l_1-1)} \otimes [(\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)})] \\ &\cong (\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)}) \\ &\oplus (\mathring{\mathcal{X}}_{k(l_1-1)} \otimes \mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1-1)} \otimes \mathring{\mathcal{X}}_{k(l_1)} \otimes \dots \otimes \mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)}). \end{aligned}$$

As $b = P_{k(l_1-1)} b P_{k(l_1-1)}$, where $P_{k(l_1-1)}$ is the projection onto $B \oplus \mathring{\mathcal{X}}_{k(l_1-1)}$. We then have

$$b a_2(A_{l_1} \cdots A_m)(x) = 0 \quad \text{for all } x \in \mathcal{X}.$$

which implies that $A_1 \cdots A_m = 0$. This finishes the proof. $\square$

Remark 6.7. When $S_i = \mathcal{A}_{i,\mathcal{F}} \cup \mathcal{A}_{i,\mathcal{B}}$, note that $\mathcal{A}_{i,\mathcal{B}} = P_i \mathcal{A}_{i,\mathcal{F}} P_i$ and Proposition 2.2, the Boolean product A_{l_1-1} can be written as $A_{l_1-1} = P_{k(l_1-1)} a P_{k(l_1-1)}$, where $a \in \mathcal{A}_{k(l_1-1),\mathcal{F}}$ following Proposition 6.2. The proof for Proposition 6.6 and the proof for Proposition 6.8 can be simplified.

Proposition 6.8. Let $A_i \in \mathcal{A}_{k(i)} (1 \leq i \leq m)$ be simple products of elements from $S_i \subset \mathcal{A}_{k(i),\mathcal{F}} \cup \mathcal{A}_{k(i),\mathcal{B}}$. If the following conditions hold:

- (1) There exist $1 \leq l_1 < l_2 \leq m$ such that $A_{l_1}, A_{l_1+1}, \dots, A_{l_2}$ are not Boolean products.
- (2) Either $l_1 = 1$ or A_{l_1-1} is a Boolean product.
- (3) Either $l_2 = m$ or A_{l_2+1} is a Boolean product.
- (4) $k(1) \neq k(2) \neq \dots \neq k(m)$.

$$(5) \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_{l_1}) = \mathbb{E}_{\mathcal{L}(X)}(A_{l_1+1}) = \cdots = \mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_{l_2}) = 0.$$

Then, we have $\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_1 \cdots A_m) = 0$.

Proof. If $1 < l_1 < l_2 < m$, the assertion follows immediately from Lemma 6.6.

If $l_2 < m, l_1 = 1$, from the proof of Lemma 6.6, we see that

$$(A_1 \cdots A_m)1_B = (A_{l_1} \cdots A_{l_2})(A_{l_2+1} \cdots A_m)1_B \\ \in (\mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)} \otimes \mathring{\mathcal{X}}_{k(l_2+1)}).$$

Hence the assertion holds in this case as well.

If $l_2 = m$, then the assumptions (4) and (5) imply that

$$(7) \quad (A_{l_1} \cdots A_{l_2})1_B \in (\mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)}).$$

We now have two cases: (i) If further $l_1 = 1$, it is clear that $\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_1 \cdots A_m) = 0$, thanks to (7). (ii) If $l_1 > 1$, then A_{l_1-1} is a Boolean product. We may write $A_{l_1-1} = a_1 b a_2$ where a_2 is a simple product of elements from $\mathcal{A}_{k(l_1-1), \mathcal{F}}$ and $b \in \mathcal{A}_{k(l_1-1), \mathcal{B}}$. It follows that

$$a_2(A_{l_1} \cdots A_m)1_B \in \mathcal{X}_{k(l_1-1)} \otimes (\mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \\ \cong (\mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)}) \oplus (\mathring{\mathcal{X}}_{k(l_1-1)} \otimes \mathring{\mathcal{X}}_{k(l_1)} \otimes \cdots \otimes \mathring{\mathcal{X}}_{k(l_2)}).$$

As $b = P_{k(l_1-1)} b P_{k(l_1-1)}$, where $P_{k(l_1-1)}$ is the projection onto $B \oplus \mathring{\mathcal{X}}_{k(l_1-1)}$. We have

$$b a_2(A_{l_1} \cdots A_m)1_B = 0,$$

which implies that $\mathbb{E}_{\mathcal{L}(\mathcal{X})}(A_1 \cdots A_m) = 0$ as well. This finishes the proof. $\square$

Proposition-Definition 6.9. Proposition 6.4, Lemma 6.6 and Proposition 6.8 provide us an algorithm for computing mixed moments of free-Boolean independent pairs of random variables and a canonical way to simplify an arbitrary element as follows. Denote by the algebra $\mathcal{A}$ generated by $\bigcup_{i \in \mathcal{I}} \{C_i \cup D_i\}$, where $\{C_i, D_i\}_{i \in \mathcal{I}}$ is a family of free-Boolean pair of B -faces in the B -valued probability space $(\mathcal{L}(X), \mathbb{E}_{\mathcal{L}(\mathcal{X})})$. That is, $C_i \subset \mathcal{A}_{i, \mathcal{F}}, D_i \subset \mathcal{A}_{i, \mathcal{B}}$ are subalgebras. Then

$$\mathcal{A} = \text{span} \left\{ A_1 \cdots A_m \left| \begin{array}{l} \text{each } A_i \text{ is a simple product of element from } \mathcal{A}_{\omega(i)}, \\ \text{and } \omega(1) \neq \omega(2) \neq \cdots \neq \omega(m) \end{array} \right. \right\}.$$

Let $X = A_1 \cdots A_m$, where each A_i is a simple product from $\mathcal{A}_{\omega(i)}$ and $\omega(1) \neq \omega(2) \neq \cdots \neq \omega(m)$. Whenever some A_k ($1 \leq k \leq m$) is a simple product of element from $\mathcal{A}_{\omega(k), \mathcal{F}}$, we replace it by

$$A_k = (A_k - \mathbb{E}(A_k)1_A) + \mathbb{E}(A_k)1_A.$$

Set $S_i = C_i \cup D_i$ ($i \in \mathcal{I}$). In viewing Lemma 6.6, Proposition 6.2 and the fact that $\mathcal{A}_{i, \mathcal{F}}$ is an algebra, the operator $A_1 \cdots A_m$ can be expressed as the sum of following types of products:

- (1) $Z_{(0)} = b 1_A$, where $b \in B$.
- (2) $Z_f = F_1 \cdots F_k$, where $k \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$, $\mathbb{E}(F_i)$ and $\omega(1) \neq \cdots \neq \omega(k)$.
- (3) $Z_b = B_1 \cdots B_k$, where $k \in \mathbb{N}$, each
 - each B_i is a Boolean product of elements from $S_{\omega(i)}$
 - and $\omega(1) \neq \cdots \neq \omega(k)$.

- (4) $Z_{fb} = F_1 \cdots F_{k_1} B_1 \cdots B_{k_2}$, where $k_1, k_2 \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_1$, each B_j is a Boolean product of elements from $S_{\omega(k_1+j)}$ for $1 \leq j \leq k_2$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2)$.
- (5) $Z_{bf} = B_1 \cdots B_{k_1} F_1 \cdots F_{k_2}$, where $k_1, k_2 \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_2$, each B_j is a Boolean product of elements from $S_{\omega(k_1+j)}$ for $1 \leq j \leq k_1$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2)$.
- (6) $Z_{fbf} = F_1 \cdots F_{k_1} B_1 \cdots B_{k_2} F_{k_1+1} \cdots F_{k_1+k_3}$, where $k_1, k_2, k_3 \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$ for $1 \leq i \leq k_1$, each $F_{k_1+j} \in C_{\omega(k_1+k_2+j)} \subset \mathcal{A}_{\omega(k_1+k_2+j), \mathcal{F}}$ for $1 \leq j \leq k_3$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_1 + k_3$, B_j is a Boolean product of elements from $S_{\omega(k_1+j)}$ for $1 \leq j \leq k_2$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2 + k_3)$.

Furthermore, $\mathbb{E}(Z_f) = 0$ by the definition of free independence, $\mathbb{E}(Z_{bf}) = \mathbb{E}(Z_{fb}) = \mathbb{E}(Z_{fbf}) = 0$ by Proposition 6.8.

Corollary 6.10. Given an operator Z_{fb} and an operator Z_{bf} of the form in Definition-Proposition 6.9 (4) and (5) respectively, for any $A, B \in \mathcal{A}$, we have

$$\mathbb{E}(Z_{fb}A) = \mathbb{E}(BZ_{bf}) = 0.$$

Proof. By linearity, it is enough to consider the case when A is the product of simple products. Let $Z_{fb} = F_1 \cdots F_{k_1} B_1 \cdots B_{k_2}$ by definition. The Boolean product B_{k_2} is concatenated with some factors in A to be a Boolean product. It then follows from Proposition 6.8 that $\mathbb{E}(Z_{fb}A) = 0$. The other case can be proved in the same way. $\square$

Corollary 6.11. Given an operator Z_{fbf} of the form in Definition-Proposition 6.9 (6), for any $A, B \in \mathcal{A}$, we have $\mathbb{E}(Z_{fbf}A) = \mathbb{E}(BZ_{fbf}) = 0$.

Proof. It is enough to consider the case when A and B are product of simple products. Noticing Proposition 6.6 and applying the simplification method described in Definition-Proposition 6.9, it is easy to see that $Z_{fbf}A$ can be written as the summation of element of the types (3) and (5), whose expectations are zero. Hence $\mathbb{E}(Z_{fbf}A) = 0$. Similarly, BZ_{fbf} can be written as the summation of elements of the types (4) and (5). Hence $\mathbb{E}(BZ_{fbf}) = 0$. $\square$

Lemma 6.12. Let $B_1 \in \mathcal{A}_i$ and $B_2 \in \mathcal{A}_j$ be two Boolean products and $i \neq j$. Then for any $A \in \mathcal{A}$, we have

$$\mathbb{E}(B_1 B_2 A) = \mathbb{E}(B_1) \mathbb{E}(B_2 A).$$

Proof. Write $B_2 A(1_B) = \mathbb{E}(B_2 A) + (B_2 A 1_B - \mathbb{E}(B_2 A)) \in B \oplus \mathring{\mathcal{X}}_j$. We can express the Boolean product B_1 as $B_1 = P_i a P_i$ following Proposition 6.2, and therefore $B_1(\mathring{\mathcal{X}}_j) = 0$, which shows that

$$\mathbb{E}(B_1 B_2 A) = \mathbb{E}(B_1(\mathbb{E}(B_2 A))) = \mathbb{E}(B_1) \mathbb{E}(B_2 A).$$

$\square$

The mixed moments $\mathbb{E}(A_1 \cdots A_n)$ can be expressed as a universal polynomial of moments of elements in individual algebras $\mathcal{A}_i$. Thus we have the following equivalent definition for free-Boolean independence under Moments conditions.

Proposition 6.13. Let $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ be a family of pairs of algebras in a B -valued probability space $(\mathcal{A}, \mathbb{E})$. Set $S_i = C_i \cup D_i$. The family $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ is free-Boolean independent if and only if

(1) whenever $B_1, \dots, B_m$ are operators such that:

- for each $1 \leq k \leq m$, B_k is a simple product of elements from $S_{\omega(k)}$, at least one of them is from $D_{\omega(k)}$;
- $\omega(1) \neq \dots \neq \omega(m)$.

then

$$\mathbb{E}(B_1 \cdots B_m) = \mathbb{E}(B_1) \cdots \mathbb{E}(B_m).$$

(2) whenever $A_1, \dots, A_m$ are operators such that:

- For each $1 \leq k \leq m$, A_k is a product of elements from $S_{\omega(k)}$.
- There exist $1 \leq l_1 < l_2 \leq m$ such that A_k is a product of elements from $C_{\omega(k)}$ for all $l_1 \leq k \leq l_2$.
- Either $l_1 = 1$ or A_{l_1-1} is a product of element from $S_{\omega(l_1-1)}$, at least one of them is in $D_{\omega(l_1-1)}$.
- Either $l_2 = m$ or A_{l_1+1} is a product of element from $S_{\omega(l_1+1)}$, at least one of them is in $D_{\omega(l_1+1)}$.
- $\omega(1) \neq \dots \neq \omega(m)$.
- $\mathbb{E}(A_{l_1}) = \mathbb{E}(A_{l_1+1}) = \dots = \mathbb{E}(A_{l_2}) = 0$.

Then, we have $\mathbb{E}(A_1 \cdots A_m) = 0$.

7. POSITIVITY OF THE AMALGAMATED FREE-BOOLEAN PRODUCT

In this section, we deal with B -functionals with positivity property. For the notion of positivity, we need a $*$ -structure on our algebras. We assume the algebra B has a nice positivity structure, i.e. we demand it to be a unital C^* -algebra. For $*$ -algebra $\mathcal{A}$, no such restriction is required.

Definition 7.1. Let $\mathcal{A}$ be a unital $*$ -algebra, element $a \in \mathcal{A}$ is said to be positive if there exists a $b \in \mathcal{A}$ such that $a = bb^*$. A B -linear functional $\mathbb{E}$ is said to be positive if $\mathbb{E}(a)$ is positive for all positive element $a \in \mathcal{A}$. A B -linear functional $\mathbb{E}$ is said to be unital if $\mathbb{E}(1_{\mathcal{A}}) = 1_B$.

In the rest of this section, we always assume that $\mathcal{A}$ is a unital $*$ -algebra and $\mathbb{E}$ is unital. Let $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ be a family of B -faces in a B -probability space $(\mathcal{A}, \mathbb{E})$, which generates $\mathcal{A}$. Suppose that the family $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ is free-Boolean independent with amalgamation over B and C_i, D_i are $*$ -subalgebras of $\mathcal{A}$ for all $i \in \mathcal{I}$. For each i , let $\mathcal{A}_i$ be the unital $*$ -algebras generated by C_i, D_i . Let $\mathbb{E}_i$ be the restriction of $\mathbb{E}$ to $\mathcal{A}_i$. Then $(\mathcal{A}_i, \mathbb{E}_i)$ is a B -valued probability space. We assume that $\mathbb{E}_i$ is unital and positive and unital for all i .

For convenience, we also introduce the following definition. Then, results in Proposition-Definition 6.9 hold in this abstract framework.

Definition 7.2. Given a set $S_i \subset C_i \cup D_i$ and $a_1, \dots, a_m \in S_i$ their product $A = a_1 \cdots a_m$ is called a *simple product* of elements from S_i . It is called a *Boolean product* of elements from S_i if $a_k \in S_i \cap D_i$ for some $1 \leq k \leq m$.

Recall that in the Section(moments-condition), the algebra $\mathcal{A}$ is the linear span of simple products of type $Z_0, Z_f, Z_b, Z_{bf}, Z_{fs}$ and Z_{fbf} , as shown in Proposition-Definition 6.9. Given $Z \in \mathcal{A}$, then Z can be written as

$$Z = Z_0 + \sum_{i_1} Z_f^{(i_1)} + \sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)} + \sum_{i_4} Z_{fb}^{(i_4)} + \sum_{i_5} Z_{fbf}^{(i_5)}.$$

We will show that $\mathbb{E}[ZZ^*]$ is positive.

Remark 7.3. $(Z_{bf}^{(i_3)})^*$ is a Z_{fb} type element and $(Z_{fb}^{(i_4)})^*$ is a Z_{bf} type element.

We first note that, by Corollary 6.10 and Corollary 6.11, we have that

$$\mathbb{E}[ZZ^*] = \mathbb{E}[Z_1 Z_1^*],$$

where $Z_1 = Z_0 + \sum_{i_1} Z_f^{(i_1)} + \sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)}$.

To simplify the notation, we introduce the following notations:

- For operator of the form $Z_b = B_1 \cdots B_k$, where $k \in \mathbb{N}$, each B_i is a Boolean product of elemnts from $S_{\omega(i)}$ and $\omega(1) \neq \cdots \neq \omega(k)$. Set

$$\begin{aligned} \Psi(Z_b) &= \mathbb{E}(B_1 \cdots B_{k-1}) B_k, \quad \text{when } k \geq 2, \\ \Psi^*(Z_b) &= B_1 \mathbb{E}(B_2 \cdots B_{k-1}), \quad \text{when } k \geq 2, \end{aligned}$$

and $\Psi(Z_b) = \Psi^*(Z_b) = Z_b$ when $k = 1$.

- For operators of the form $Z_{bf} = B_1 \cdots B_{k_1} F_1 \cdots F_{k_2}$, where $k_1, k_2 \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_2$, each B_j is a Boolean product of elements from $S_{\omega(k_1+j)}$ for $1 \leq j \leq k_1$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2)$. Set

$$\Psi(Z_{bf}) = \mathbb{E}(B_1 \cdots B_{k_1-1}) (B_{k_1} F_1 \cdots F_{k_2}), \quad \text{when } k_1 \geq 2,$$

and $\Psi(Z_{bf}) = Z_{bf} = B_1 (F_1 \cdots F_{k_2})$ when $k_1 = 1$.

- For operator of the form $Z_{fb} = F_1 \cdots F_{k_1} B_1 \cdots B_{k_2}$, where $k_1, k_2 \in \mathbb{N}$, each $F_i \in C_{\omega(i)} \subset \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_1$, each B_j is a Boolean product of elemnts from $S_{\omega(k_1+j)}$ for $1 \leq j \leq k_2$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2)$. Set

$$\Psi^*(Z_{fb}) = (F_1 \cdots F_{k_1} B_1) \mathbb{E}(B_2 \cdots B_{k_2}), \quad \text{when } k_2 \geq 2,$$

and $\Psi^*(Z_{fb}) = Z_{fb}$ when $k_2 = 1$.

Note that Z_{bf}^* is of the same type as Z_{fb} following Proposition-Defintion 6.9 (4), (5). It is easy to check that the following relation holds:

$$\Psi(Z_{bf})^* = \Psi^*(Z_{bf}^*).$$

Lemma 7.4. Let $Z_{bf} = B_1 \cdots B_{k_1} F_1 \cdots F_{k_2}$, where each $F_i \in \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_2$, each B_j is a Boolean product from $\mathcal{A}_{\omega(k_1+j)}$ for $1 \leq j \leq k_1$, and $\omega(1) \neq \cdots \neq \omega(k_1 + k_2)$. Then, $\mathbb{E}[Z_{bf} Z'] = \mathbb{E}[\Psi(Z_{bf}) Z']$ for all simple products Z' .

Proof. It follows by applying Lemma 6.12 inductively. $\square$

Lemma 7.5. Let $Z_{fb} = F_1 \cdots F_{k_1} B_1 \cdots B_{k_2}$, where each $F_i \in \mathcal{A}_{\omega(i), \mathcal{F}}$ such that $\mathbb{E}(F_i) = 0$ for $1 \leq i \leq k_1$, $k_1 \geq 0$, $k_2 \geq 1$ each B_j is a Boolean product from $\mathcal{A}_{\omega(k_1+j)}$ for $k_1 < j \leq k_1 + k_2$, and $\omega(1) \leq \omega(k_1 + k_2)$. Then, $\mathbb{E}[Z' Z_{fb}] = \mathbb{E}[Z' \Psi^*(Z_{fb})]$ for all simple products Z' .

Proof. It follows by applying Lemma 6.3 inductively. $\square$

Lemma 7.6. $\mathbb{E}[Z_1 Z_1^*] = \mathbb{E}[Z_2 Z_2^*]$, where $Z_2 = Z_0 + \sum_{i_1} Z_f^{(i_1)} + \Psi\left(\sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)}\right)$.

Proof. Apply Lemma 7.4 and Lemma 7.5, we have

$$\begin{aligned}
& \mathbb{E}[Z_1 Z_1^*] \\
&= \mathbb{E}\left[(Z_0 + \sum_{i_1} Z_f^{(i_1)} + \sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)})(Z_0^* + \sum_{i_1} Z_f^{(i_1)*} + \sum_{i_2} Z_b^{(i_2)*} + \sum_{i_3} Z_{bf}^{(i_3)*})\right] \\
&= \mathbb{E}\left[(Z_0 + \sum_{i_1} Z_f^{(i_1)} + \sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)})(Z_0^* + \sum_{i_1} Z_f^{(i_1)*} + \sum_{i_2} Z_b^{(i_2)*} + \sum_{i_3} Z_{bf}^{(i_3)*})\right] \\
&= \mathbb{E}\left[(Z_0 + \sum_{i_1} Z_f^{(i_1)} + \Psi(\sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)}))(Z_0^* + \sum_{i_1} Z_f^{(i_1)*} + (\sum_{i_2} Z_b^{(i_2)*} + \sum_{i_3} Z_{bf}^{(i_3)*}))\right] \\
&= \mathbb{E}\left[(Z_0 + \sum_{i_1} Z_f^{(i_1)} + \Psi(\sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)}))(Z_0^* + \sum_{i_1} Z_f^{(i_1)*} + \Psi^*(\sum_{i_2} Z_b^{(i_2)*} + \sum_{i_3} Z_{bf}^{(i_3)*}))\right] \\
&= \mathbb{E}\left[(Z_0 + \sum_{i_1} Z_f^{(i_1)} + \Psi(\sum_{i_2} Z_b^{(i_2)} + \sum_{i_3} Z_{bf}^{(i_3)}))(Z_0^* + \sum_{i_1} Z_f^{(i_1)*} + (\Psi^*(\sum_{i_2} Z_b^{(i_2)*} + \sum_{i_3} Z_{bf}^{(i_3)*})))\right] \\
&= \mathbb{E}[Z_2 Z_2^*]
\end{aligned}$$

$\square$

Notice that $\Psi(Z_b^{(i_2)})$ is a simple Boolean product and $\Psi(Z_{bf}^{(i_3)})$ can be written as $B_{i_3} Z_{f'}^{(i_3)}$ where $Z_{f'}^{(i_3)}$ is the type (3) of product in Proposition-Definition 6.9. According to the length of the word appearing in the expression of Z_2 defined in Lemma 7.6, we can then rewrite Z_2 as

$$Z_2 = b1_{\mathcal{A}} + \sum_{k=1}^n \sum_{r=1}^{m_k} a_{k,r,1} a_{k,r,2} \cdots a_{k,r,k},$$

where $a_{k,r,p} \in D_{\omega_{k,r}(p)}$ (the right face) for $p \geq 2$, and $a_{k,r,1} \in S_{\omega_{k,r}(p)}$ (either left face or right face), $\mathbb{E}(a_{k,r,p}) = 0$ and $\omega_{k,r} : \{1, \dots, k\} \rightarrow \mathcal{I}$ such that $\omega_{k,r}(1) \neq \dots \neq \omega_{k,r}(k)$.

We now have the following result.

Lemma 7.7. Let $a_1 \in \mathcal{A}_{\omega_1(1)}$, $a_i \in \mathcal{A}_{\omega_1(i), \mathcal{F}}$ for $i = 2, \dots, n$ such that $\mathbb{E}(a_i) = \mathbb{E}(\tilde{a}_i) = 0$ and $\omega_1 : \{1, \dots, n\} \rightarrow \mathcal{I}$, $\omega_1(1) \neq \dots \neq \omega_1(n)$. Let $\tilde{a}_1 \in \mathcal{A}_{\omega_2(1)}$, $\tilde{a}_j \in \mathcal{A}_{\omega_2(j), \mathcal{F}}$ for $j = 2, \dots, m$ such that $\omega_2 : \{1, \dots, m\} \rightarrow \mathcal{I}$, $\omega_2(1) \neq \dots \neq \omega_2(m)$. We then have

$$\mathbb{E}[a_1 a_2 \cdots a_{n-1} a_n \tilde{a}_m \tilde{a}_{m-1} \cdots \tilde{a}_1] = \delta_{n,m} \mathbb{E}[a_1 a_2 \cdots \mathbb{E}[a_{n-1} \mathbb{E}[a_n \tilde{a}_m] \tilde{a}_{m-1}] \cdots \tilde{a}_1].$$

Proof. When $n = 0$ or $m = 0$, there is nothing to prove. It is sufficient to prove that

$$\mathbb{E}[a_1 a_2 \cdots a_{n-1} a_n \tilde{a}_m \tilde{a}_{m-1} \cdots \tilde{a}_1] = \delta_{n,m} \mathbb{E}[a_1 a_2 \cdots a_{n-1} \mathbb{E}[a_n \tilde{a}_m] \tilde{a}_{m-1} \cdots \tilde{a}_1].$$

Notice that if $\omega_1(n) \neq \omega_2(m)$, then by Proposition 6.8 and the definition of freeness, then, by Proposition 6.8, we have

$$\mathbb{E}[a_1 a_2 \cdots a_{n-1} a_n \tilde{a}_m \tilde{a}_{m-1} \cdots \tilde{a}_1] = \mathbb{E}[a_n \tilde{a}_m] = 0.$$

On the other hand, if $\omega_1(n) = \omega_2(m)$, notice that $a_n \tilde{a}_m - \mathbb{E}[a_n \tilde{a}_m] \in \mathcal{A}_{\omega_1(n)}$, $\omega_1(n) \neq \omega_1(n-1)$ and $\omega_1(n) \neq \omega_2(m-1)$, then we have

$$\mathbb{E}[a_1 a_2 \cdots a_{n-1} (a_n \tilde{a}_m - \mathbb{E}[a_n \tilde{a}_m]) \tilde{a}_{m-1} \cdots \tilde{a}_1] = 0,$$

which is the desired equation. $\square$

Therefore, we have the following equation.

$$\mathbb{E}[Z_2 Z_2^*] = bb^* + \sum_{k=1}^n \mathbb{E} \left[\left(\sum_{r=1}^{m_k} a_{k,r,1} a_{k,r,2} \cdots a_{k,r',k} \right) \left(\sum_{r'=1}^{m_k} a_{k,r',1} a_{k,r',2} \cdots a_{k,r',k} \right)^* \right].$$

Moreover, Lemma 7.7 implies that the term $\mathbb{E}[(a_{k,r,1} a_{k,r,2} \cdots a_{k,r',k})(a_{k,r',1} a_{k,r',2} \cdots a_{k,r',k})^*]$ is not vanishing only if $\omega_{k,r} = \omega_{k,r'}$. Let $\sim$ be the equivalence relation on $\{1, \dots, m_k\}$ such that $l_1 \sim l_2$ if and only if $\omega_{k,l_1,k} = \omega_{k,l_2,k}$. Let $\{V_1, \dots, V_s\}$ be the family of equivalence classes of $\{1, \dots, m_k\}$. Then

$$\begin{aligned} & \mathbb{E} \left[\left(\sum_{r=1}^{m_k} a_{k,r,1} a_{k,r,2} \cdots a_{k,r',k} \right) \left(\sum_{r'=1}^{m_k} a_{k,r',1} a_{k,r',2} \cdots a_{k,r',k} \right)^* \right] \\ = & \mathbb{E} \left[\left(\sum_{r \in V_l} a_{k,r,1} a_{k,r,2} \cdots a_{k,r',k} \right) \left(\sum_{r' \in V_l} a_{k,r',1} a_{k,r',2} \cdots a_{k,r',k} \right)^* \right] \end{aligned}$$

To show that $E[Z_2 Z_2^*]$ is positive, we just need to show that

$$\mathbb{E} \left[\left(\sum_{r \in V_l} a_{k,r,1} a_{k,r,2} \cdots a_{k,r',k} \right) \left(\sum_{r' \in V_l} a_{k,r',1} a_{k,r',2} \cdots a_{k,r',k} \right)^* \right] \geq 0$$

for all k .

This is exactly the circumstance in the proof of [16, Proposition 3.5.6]. Therefore, we have reduced the positivity question of $\mathbb{E}[ZZ^*]$ to a known result in free probability context. Since Z is arbitrary, we thus have the following theorem.

Theorem 7.8. Let $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ be a family of B -faces in a B -probability space $(\mathcal{A}, \mathbb{E})$, which generates $\mathcal{A}$. We assume that C_i, D_i are $*$ -subalgebras of $\mathcal{A}$ for all $\mathcal{A}$. For each i , let $\mathcal{A}_i$ be the $*$ -algebras generated by C_i, D_i . We assume that the restriction of $\mathbb{E}$ to $\mathcal{A}_i$ is positive. If the family $\{(C_i, D_i)\}_{i \in \mathcal{I}}$ is free-Boolean independent with amalgamation over B , then $\mathbb{E}$ is positive.

Acknowledgement The second-named author would like to thank Professor Hari Bercovici for the invitation to visit Indiana University, where the authors can meet and part of this work was done. He also wants to thank Professor Alexandru Nica for his continued support. This project was partially supported by NSFC No. 11501423, 11431011.

REFERENCES

- [1] Marek Bożejko, Michael Leinert, and Roland Speicher. Convolution and limit theorems for conditionally free random variables. *Pacific J. Math.*, 175(2):357–388, 1996.
- [2] Ian Charlesworth, Brent Nelson, and Paul Skoufranis. Combinatorics of bi-freeness with amalgamation. *Comm. Math. Phys.*, 338(2):801–847, 2015.
- [3] Yinzhen Gu, Takahiro Hasebe, and Paul Skoufranis. Bi-monotonic independence for pairs of algebras. *arXiv:1708.05334*, 2017.
- [4] Yinzhen Gu and Paul Skoufranis. Bi-boolean independence for pairs of algebras. *arXiv:1703.03072*, to appear in *Complex Anal. Oper. Theory*, 2017.
- [5] Yinzhen Gu and Paul Skoufranis. Conditional bi-free independence with amalgamation, *arXiv:1609.07820*, to appear in *Int. Math. Res. Not. IMRN*, 2017.
- [6] Yinzhen Gu and Paul Skoufranis. Conditionally bi-free independence for pairs of faces. *J. Funct. Anal.*, 273(5):1663–1733, 2017.
- [7] Weihua Liu. A noncommutative de Finetti theorem for boolean independence. *J. Funct. Anal.*, 269(7):1950–1994, 2015.
- [8] Weihua Liu. Free-boolean independence for pairs of algebras. *arXiv:1710.01374*, 2017.
- [9] Wojciech Młotkowski. Operator-valued version of conditionally free product. *Studia Math.*, 153(1):13–30, 2002.
- [10] Naofumi Muraki. The five independences as natural products. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 6(3):337–371, 2003.
- [11] Alexandru Nica and Roland Speicher. *Lectures on the combinatorics of free probability*, volume 335 of *London Mathematical Society Lecture Note Series*. Cambridge University Press, Cambridge, 2006.
- [12] Mihai Popa. A combinatorial approach to monotonic independence over a C^* -algebra. *Pacific J. Math.*, 237(2):299–325, 2008.
- [13] Mihai Popa. A new proof for the multiplicative property of the Boolean cumulants with applications to the operator-valued case. *Colloq. Math.*, 117(1):81–93, 2009.
- [14] Gian-Carlo Rota. On the foundations of combinatorial theory. I. Theory of Möbius functions. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 2:340–368 (1964), 1964.
- [15] Roland Speicher. On universal products. In *Free probability theory (Waterloo, ON, 1995)*, volume 12 of *Fields Inst. Commun.*, pages 257–266. Amer. Math. Soc., Providence, RI, 1997.
- [16] Roland Speicher. Combinatorial theory of the free product with amalgamation and operator-valued free probability theory. *Mem. Amer. Math. Soc.*, 132(627):x+88, 1998.
- [17] Roland Speicher and Reza Woroudi. Boolean convolution. In *Free probability theory (Waterloo, ON, 1995)*, volume 12 of *Fields Inst. Commun.*, pages 267–279. Amer. Math. Soc., Providence, RI, 1997.
- [18] Dan Voiculescu. Operations on certain non-commutative operator-valued random variables. *Astérisque*, (232):243–275, 1995. Recent advances in operator algebras (Orléans, 1992).
- [19] Dan-Virgil Voiculescu. Free probability for pairs of faces I. *Comm. Math. Phys.*, 332(3):955–980, 2014.

WEIHUA LIU: DEPARTMENT OF MATHEMATICS, INDIANA UNIVERSITY, BOOMINGTON, IN 47401, USA.,
E-mail address: liuweih@indiana.edu

PING ZHONG: DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, 200 UNIVERSITY AVENUE WEST, WATERLOO, ON, N2L 3G1, CANADA.
E-mail address: ping.zhong@uwaterloo.ca