The Complexity of Maximum *k*-Order Bounded Component Set Problem

Sounaka Mishra¹

Department of Mathematics, Indian Institute of Technology Madras, Chennai 600 036, India

Shijin Rajakrishnan

Department of Computer Science, Cornell University, New York, USA

Abstract

Given a graph G = (V, E) and a positive integer k, in MAXIMUM k-ORDER BOUNDED COMPONENT SET (MAX-k-OBCS), it is required to find a vertex set $S \subseteq V$ of maximum size such that each component in the induced graph G[S] has at most k vertices. We prove that for constant k, MAX-k-OBCS is hard to approximate within a factor of $n^{1-\epsilon}$, for any $\epsilon > 0$, unless P = NP. This is an improvement on the previous lower bound of \sqrt{n} for MAX-2-OBCS due to Orlovich et. el. [1]. We provide lower bounds on the approximability when k is not a constant as well. MAX-k-OBCS can be seen as a generalization of MAXIMUM INDEPENDENT SET (MAX-IS). We generalize Turán's greedy algorithm for MAX-IS and prove that it approximates MAX-k-OBCS within a factor of $(2k-1)\overline{d} + k$, where \overline{d} is the average degree of the input graph G. This approximation factor is a generalization of Turán's approximation factor for MAX-IS.

Keywords: Approximation algorithms, Graph algorithm, Dissociation set, *k*-order bounded component

1. Introduction

In this paper, we consider the computational complexity of an optimization problem called the MAXIMUM *k*-ORDER BOUNDED COMPONENT SET problem (MAX-*k*-OBCS). In MAX-*k*-OBCS, given a graph G = (V, E) and a positive integer *k*, it is required to find a vertex set $S \subseteq V$ of maximum size such that order of each component in G[S] is at most *k*.

This is a generalization of several important problems in combinatorics. For k = 1, this problem is the same as the MAXIMUM INDEPENDENT SET problem (MAX-IS), where given a graph G = (V, E), the objective is to find a subset of vertices $S \subseteq V$ of maximum size such that there is no edge between any pair of vertices in G[S]. The case

Email addresses: sounak@iitm.ac.in (Sounaka Mishra), sr986@cornell.edu (Shijin Rajakrishnan) 1 Corresponding author

Preprint submitted to Elsevier

of k = 2 gives rise to the MAXIMUM DISSOCIATION SET problem (MAX-DISSO-SET), where given a graph G, the objective is to find a vertex set $S \subseteq V$ of maximum size such that the degree of each vertex in G[S] is at most 1. Both these problems are NP-hard and thus MAX-k-OBCS is NP-hard as well.

Finding a maximum sized independent set is one of the widely studied problems in combinatorics and its hardness to compute has prompted several attempts at developing approximate solutions. Håstad [2] proved that , for any $\epsilon > 0$, MAX-IS cannot be approximated within a factor of $n^{1-\epsilon}$, unless NP = ZPP. Later, Zukerman [3] proved that MAX-IS cannot be approximated within a factor of $n^{1-\epsilon}$, unless P = NP, for any $\epsilon > 0$. This lower bound result rules out the existence of a polynomial time efficient algorithm on general graphs. However, the situation is not quite as grim when restricted to special graph classes. It is polynomial time solvable to find a maximum independent set in claw-free graphs [4], P₅-free graphs [5], and perfect graphs [6]. For other classes, although finding a polynomial time optimal solution might be out of the question unless P = NP, we can nevertheless provide good approximation algorithms. In planar graphs, there exists a PTAS for MAX-IS [7] whereas for graphs with bounded degree, the problem can be approximated within a constant factor [8]. One such algorithm greedily includes a vertex of minimum degree into the solution set, deletes that vertex along with its neighbors from the graph and then repeats this selection process until there are no more vertices to process. This simple greedy algorithm returns an independent set which is within a factor of $(\overline{d} + 1)$ from the optimal solution (Turán's Theorem) [9]. A tighter analysis has improved the approximation factor to $\frac{1}{5}(2d+3)$ [10].

In [11], Yannakakis defined dissociation set and proved that it is NP-complete to find a vertex set *S* of minimum size in a given bipartite graph *G* such that $G[V \setminus S]$ has maximum degree 1. From this result it follows that MAX-DISSO-SET is NP-complete even when restricted to bipartite graphs. It is also known to be NP-complete for $K_{1,4}$ -free bipartite graphs, C_4 -free bipartite graphs of maximum degree 3 [12], planar graphs with maximum degree 4 [13] and planar line graphs of planar bipartite graphs [1]. However, MAX-DISSO-SET is polynomial time solvable when the input graph is restricted to some hereditary classes of graphs, such as {*chair*, *bull*}-free graphs, {*chair*, K_3 }-free graphs, {*mK*₂}-free graphs with $m \ge 2$ [1].

Approximation algorithms for the complementary problem of MAX-*k*-OBCS have been designed by various researchers. In this minimization problem, we are asked to find a vertex set *S* of minimum size in a given graph *G* such that each component in $G[V \setminus S]$ has at most *k* vertices. When k = 1 and 2, these minimization problems are known as MINIMUM VERTEX COVER and MINIMUM DISSOCIATION SET, respectively. In MINIMUM DISSOCIATION SET it is required to find a vertex set *S* of minimum cardinality in a given graph G = (V, E) such that degree of each vertex in $G[V \setminus S]$ is at most 1. It is interesting to observe that MINIMUM DISSOCIATION SET is equivalent to MINIMUM *P*₃ VERTEX COVER in which it is required to find a vertex set *S* of minimum size such that $G[V \setminus S]$ does not have a *P*₃ (path on 3 vertices) as a subgraph. Both MINI-MUM VERTEX COVER and MINIMUM DISSOCIATION SET are known to be approximable within a factor of 2 [14]. For larger values of *k*, the deletion problem is known to be approximable within a factor of *k* and approximable within a factor of (k - 1) when the input graph has girth at least *k* [15]. However, to the best of our knowledge, there is no known result about the approximability of MAX-*k*-OBCS for arbitrary graphs. **Our Contribution.** In this paper, we explore the approximability and inapproximability of MAX-k-OBCS, for different integral values of k. In Section 3, we prove inapproximability results by establishing an approximation preserving reduction from MAX-IS, which shows that for constant k, this problem has no efficient approximation algorithms for general graphs. The reduction used can be composed to prove hardness results even for non-constant k. In particular, we show that for $k = O(\frac{n}{\log n})$, it is not possible to develop an algorithm with a better approximation guarantee than $O(\log n)$, unless P = NP. Having established the hardness of computing efficient solutions to this problem, in the subsequent sections we concentrate on developing upper bounds. In Section 4, we prove that MAX-WEIGHTED-k-OBCS can be approximated within a factor of Δ , where Δ is the maximum degree of the input graph G. This also proves that the problem can be approximated within a constant factor for degree bounded graphs. It is important to note that this bound holds not just for the unweighted case but also for the weighted version of the problem, where there is a weight function $w: V \to \mathbb{R}^+$ as an additional input and the objective is to find a vertex set *S* of maximum weight such that G[S] has no component whose order exceeds k. The technique used in the algorithm is the Local Ratio Method [16]. In Section 5, we build upon the algorithm for MAX-IS to develop a greedy algorithm for MAX-DISSO-SET, with a performance guarantee of (3d+2), where d is the average degree of the graph. We then generalize the algorithm and the analysis to the larger problem with arbitrary values of k and prove that the extension of the algorithm yields a solution which is within a factor of (2k-1)d + kfrom the optimal solution. This can be observed as a generalization of Turán's bound mentioned earlier for MAX-IS. Finally, in Section 6, we establish a reduction to MAX-IS and use existing upper bounds for MAX-IS [17] to prove that MAX-k-OBCS can be approximated within a factor of $O(\frac{k\Delta \log \log \Delta}{\log \Delta})$, for sufficiently large values of Δ .

2. Notations

Throughout this paper, we assume that any graph G = (V, E) mentioned is simple and undirected, with |V| = n vertices and |E| = m edges. Given a graph G = (V, E), the open neighborhood $N_G(v)$ of a vertex v is the set of vertices those are adjacent to vin G and the closed neighborhood $N_G[v]$ of v is defined as $N_G[v] = N_G(v) \cup \{v\}$. For a given set $P \subseteq V$, we define $N_P(v) = N_G(v) \cap P$, for each $v \in V$. The degree of a vertex v in the graph G is defined as $d_G(v) = |N_G(v)|$, the maximum degree of the graph is $\Delta_G = \max_{v \in V} d_G(v)$. When the underlying graph G is unambiguous, we drop the subscript, for example we use just Δ to denote the maximum degree in G instead of Δ_G . The average degree of the graph G is $\overline{d} = \frac{1}{n} \sum_{v \in V} d_G(v)$. There is an interesting relationship between the average degree of a graph and the total number of edges in the graph, which we use in this paper, viz. the sum of the degrees of all the vertices in a graph is twice the number of edges. So $\sum_{v \in V} d_G(v) = 2|E|$, which gives us $|V|\overline{d} = 2|E|$.

Given a subset of vertices, $S \subseteq V$, the graph induced on this set, G[S], is the subgraph whose vertex set is *S* and the edge set is the subset of edges *E* such that both endpoints are in *S*, i.e, G[S] = (V', E') where V' = S and $E' = \{(u, v) \in E \mid u \in S \text{ and } v \in S\}$.

A vertex set $S \subseteq V$ is called an independent set if G[S] has no edges. We shall denote $\alpha(G)$ as the size of a maximum independent set in *G*. An independent set *S* is *maximal*

if it is not a proper subset of another independent set of *G*. A set of vertices *S* of *G* is called a dissociation set if degree of each vertex in G[S] is at most 1 (equivalently, each component in G[S] has at most two verices). In the maximum dissociation set problem (MAX-DISSO-SET), the objective is to find a dissociation set of maximum size in a given graph *G*. A set $S \subseteq V$ is called a *k*-component set in *G* if each component in G[S] has at most *k* vertices. We shall denote the size of a maximum *k*-component set in *G* by comp_k(*G*).

3. Hardness of approximation

In this section, we shall prove inapproximability results for MAX-*k*-OBCS by establishing an AP-reduction [18] from MAX-IS. The following lower bound result for MAX-IS makes this reduction useful.

Theorem 3.1. [3] Unless P = NP, for any $\epsilon > 0$, MAX-IS is not approximable within a factor of $n^{1-\epsilon}$.

The hardness result mentioned in Theorem 3.1 establishes the hardness of approximation for MAX-1-OBCS. Next, we establish the hardness of approximation for MAX-k-OBCS for any constant positive integer k. We make use of the following reduction for this purpose.

Lemma 3.2. For any k > 0, MAX-k-OBCS \leq_{AP} MAX-2k-OBCS.

Proof. Given an instance G = (V, E) of MAX-*k*-OBCS, we construct an instance G' = (V', E') of MAX-2*k*-OBCS as follows. First we make two copies $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ of the graph G with $V_1 = \{v_1 \mid v \in V\}$, $V_2 = \{v_2 \mid v \in V\}$ and E_1, E_2 are defined accordingly. Then, for each edge $(u, v) \in E$, we add two edges (u_1, v_2) and (u_2, v_1) . Finally, we introduce the edge (u_1, u_2) for each $u \in V$. Thus we have $V' = V_1 \cup V_2$ and $E' = E_1 \cup E_2 \cup \{(u_1, v_2), (u_2, v_1) \mid (u, v) \in E\} \cup \{(u_1, u_2) \mid u \in V\}$. It is easy to observe that |V'| = 2|V|. For an illustration we refer to Figure 1.




Figure 1: Sketch of the construction of G' from G

Let S^* be a maximum *k*-component set in *G* and S'^* be a maximum 2*k*-component set in *G'*. From the construction of *G'* from *G*, it follows that the set $S' = \{v_1, v_2 \mid v \in S\}$ is a 2*k*-component set in *G'*, for every *k*-component set *S* in *G*. Also |S'| = 2|S|. Since these two problems are maximization problems we have $2|S^*| \le |S'^*|$.

Let *S'* be a 2*k*-component set in *G'*, consisting of *t* components. From *S'* we will construct a *k*-component set *S* in *G* with $|S| \ge \frac{1}{2}|S'|$.

First, we prove a property of the components in G'[S'] which is as follows. For any two distinct components T_1 and T_2 , if we look at the vertices in G induced by the components then there are no common vertices. In other words, the sets $\{v \in V \mid v_1 \in$ T_1 or $v_2 \in T_1\}$ and $\{v \in V \mid v_1 \in T_2 \text{ or } v_2 \in T_2\}$ are disjoint. Suppose that these two sets have a common vertex, say u. Then by construction, $u_1 \in T_1$ and $u_2 \in T_2$ and $(u_1, u_2) \in E'$. This implies that T_1 and T_2 are connected, which is a contradiction.

Now, we construct a set $S \subseteq V$ from S' as follows. Let T_1, T_2, \ldots, T_t be the components in G'[S']. For each component T_i in G'[S'], if $|T_i \cap V_1| \ge |T_i \cap V_2|$ and $|T_i \cap V_1| \ge k$ choose a set J_i of any k vertices from $T_i \cap V_1$; otherwise if $|T_i \cap V_1| < k$, take $J_i = T_i \cap V_1$. Finally, we update $S = S \cup \{u \in V \mid u_1 \in J_i\}$. If $|T_i \cap V_2| > |T_i \cap V_1|$, then we take vertices from V_2 into the set J_i and update $S = S \cup \{u \in V \mid u_1 \in J_i\}$.

Let $S = J_1 \cup J_2 \cup ... \cup J_t$, where J_i is the vertex set selected from the component T_i of G'[S']. We claim that S is a k-component set in G. It is easy to observe that $G[J_i]$ can have at most k vertices. Next, we need to prove that there is no edge between a vertex u in J_i and a vertex v in J_j , for $i \neq j$. The existence of such an edge would imply that the corresponding components T_i and T_j are not distinct (because of the edges $(u_1, v_1), (u_1, v_2), (u_2, v_1), (u_2, v_2)$).

From the construction of vertex set J_i from the component T_i , it follows that cardinality of J_i is at least half of the vertices in T_i . Therefore, $|S| \ge \frac{1}{2}|S'|$.

Therefore, for any given 2*k*-component set *S*' in *G*', one can construct a *k*-component set *S* in *G* in polynomial time, such that $\frac{|S^*|}{|S|} \leq \frac{|S'^*|}{|S'|}$.

Corollary 3.3. For k > 0, MAX-IS is AP-reducible to MAX-k-OBCS with size amplification of kn, where n is the number of vertices in the input instance of MAX-IS and $k = 2^i$, for some nonnegative integer i.

Proof. We reduce MAX-IS to MAX-*k*-OBCS by composing the reduction in Lemma 3.2 $\lceil \log k \rceil$ times to get an instance of MAX-*k*-OBCS. Since the size of the new instance in the reduction of Lemma 3.2 doubles, on applying the reduction $\lceil \log k \rceil$ times, the size of the MAX-*k*-OBCS instance is $2^{\lceil \log k \rceil} n \approx kn$. Since the reduction in Lemma 3.2 is a ratio preserving reduction, the composite reduction from MAX-IS to MAX-*k*-OBCS is a ratio preserving reduction. Therefore, MAX-IS \leq_{AP} MAX-*k*-OBCS.

Using Theorem 3.1 and Corollary 3.3 we have the following result for MAX-k-OBCS.

Corollary 3.4. For any $\epsilon > 0$, MAX-2^{*i*}-OBCS is hard to approximate within a factor of $n^{1-\epsilon}$, unless P = NP, where *i* is a constant nonnegative integer.

Next we will extend this lower bound result for MAX-*k*-OBCS, where *k* is a constant nonnegative integer.

Theorem 3.5. For any $\epsilon > 0$, MAX-k-OBCS is hard to approximate within a factor of $n^{1-\epsilon}$, unless P = NP, where k is a constant nonnegative integer.

Proof. Now consider a positive integer *k* such that $2^i < k < 2^{i+1}$ for some non-negative integer *i*. Suppose that MAX-*k*-OBCS can be approximated within a factor of β , to obtain

a subset S_k such that $|S_k| \ge \frac{1}{\beta}|S_k^*|$. We can use this subset S_k to produce a solution for MAX-2^{*i*}-OBCS, by considering each component in S_k and dropping vertices arbitrarily from each component until its size is at most 2^{*i*}. Notice that the number of components in S_k with size more than 2^{*i*} can be at most $\frac{|S_k|}{2^i}$, and the number of vertices we drop from each such component is at most $k - 2^i$. Thus the size of the constructed solution to MAX-2^{*i*}-OBCS, S_{2^i} is at least $|S_k| - \frac{|S_k|}{2^i}(k - 2^i) = |S_k|\frac{2^{i+1}-k}{2^i}$. We also know that $|S_{2^i}^*| \le |S_k^*|$, and thus the approximation ratio of this constructed set is

$$\frac{|S_{2^i}|}{|S_{2^i}^*|} \ge |S_k| \frac{2^{i+1}-k}{2^i} \frac{1}{|S_k^*|} \ge \frac{2^{i+1}-k}{2^i} \frac{1}{\beta}$$

Now the hardness result of MAX-2^{*i*}-OBCS implies that $\beta \in \omega(n^{1-\varepsilon})$, and thus MAX*k*-OBCS cannot be approximated within a factor of $n^{1-\varepsilon}$ for any constant *k*, unless P = NP.

The authors in [1] have proved that, for any $\epsilon > 0$, MAX-DISSO-SET is hard to approximate within a factor of $n^{\frac{1}{2}-\epsilon}$, unless P = NP. When i = 1, MAX- 2^i -OBCS is same as MAX-DISSO-SET and we have the following result which is a stronger negative result for MAX-DISSO-SET.

Theorem 3.6. For any $\epsilon > 0$, MAX-DISSO-SET is hard to approximate within a factor of $n^{1-\epsilon}$, unless P = NP.

Next, we consider the inapproximability of MAX-*k*-OBCS when *k* is not a constant and our result is as follows.

Theorem 3.7. If k is not a constant, then for any $\epsilon > 0$, MAX-k-OBCS cannot be approximated in polynomial time with a better factor than $n \frac{\log(n/k)}{\log(n)} - \epsilon$, unless P = NP.

Proof. Note that in the previous theorem, the graph G' constructed in the reduction of MAX-*k*-OBCS from MAX-IS has size $n' = n2^{\log k} = nk$, where *n* is the number of vertices in an instance *G* of MAX-IS.

Now suppose that MAX-*k*-OBCS can be approximated within a factor of $n'^{\frac{\log(n'/k)}{\log(n')}} - \epsilon$ in polynomial time, then

$$\frac{\alpha(G)}{|S|} \le \frac{\operatorname{comp}_k(G')}{|S'|} \le (nk)^{\frac{\log(n)}{\log(nk)} - \epsilon}$$
$$= \frac{n}{(nk)^{\epsilon}} \cdot n^{-\frac{\log k}{\log nk}} k^{\frac{\log n}{\log nk}}$$
$$= \frac{n}{(nk)^{\epsilon}} \cdot \left(n^{-\log k} n^{\log k}\right)^{\frac{1}{\log nk}}$$
$$= \frac{n}{(n)^{\epsilon'}} = n^{1-\epsilon'} < n^{1-\epsilon},$$

where $\epsilon' = \epsilon \left[1 + \frac{\log k}{\log n} \right]$. The last inequality holds as $\epsilon' > \epsilon$. This contradicts the lower bound for MAX-IS available in Theorem 3.1.

From Theorem 3.7, we have the following corollary,

Corollary 3.8. If $k = O(\frac{n}{\log n})$, then MAX-k-OBCS is hard to approximate within a factor of $O(\log n)$, unless P = NP.

4. Approximation algorithms for MAX-WEIGHTED-k-OBCS

We now turn to the weighted version of the problem, MAX-WEIGHTED-*k*-OBCS. The input, in addition to a graph G = (V, E), consists of a weight function on the vertices of the graph, $w : V \to \mathbb{R}^+$. Given a subset of vertices $A \subseteq V$, the weight of the subset is defined as $w(A) = \sum_{v \in A} w(v)$. In MAX-WEIGHTED-*k*-OBCS, the objective is to find a subset of vertices $S \subseteq V$ of maximum weight w(S) such that the size of the largest component in G[S] is at most *k*. Since the unweighted case looked at thus far is a special case of the weighted case (with the weight function being uniform), the lower bounds derived in the previous section hold for the weighted version as well.

In this section, we look at an approximation algorithm for this problem and prove an approximation guarantee of Δ , the maximum degree of the input graph.

Algorithm 4.1: *k*-Weighted-OBCS(*G*,*w*)

Input: An undirected simple graph G = (V, E) and $w : V \to \mathbb{R}^+$ **Output:** $S \subseteq V$ such that each component in G[S] has at most *k* vertices 1 if $G = \emptyset$ then 2 Return \emptyset ; 3 end 4 **if** there exists a vertex u such that $w(u) \leq 0$ **then** Return *k*-Weighted-OBCS($G \setminus \{u\}, w$); 5 6 else Let (s, t) be the edge in *G* that maximizes w(s) + w(t); 7 Define the weight functions $w_1(u) = \begin{cases} w(u), & u \in N[\{s,t\}] \\ 0, & o.w. \end{cases}$, and 8 $w_2 = w - w_1;$ $S' \leftarrow k$ -Weighted-OBCS (G, w_2) ; 9 10 Return $S' \cup \{s, t\}$; 11 end

The algorithm is described in Algorithm 4.1 and the analysis makes use of the local ratio theorem.

Theorem 4.1. [16] Let (G = (V, E), w) be an instance of MAX-WEIGHTED-k-OBCS having n vertices. Let $w_1, w_2 \in \mathbb{R}^n$ such that $w = w_1 + w_2$. Let $S \subseteq V$ be an r-approximate feasible solution of MAX-WEIGHTED-k-OBCS for G with respect to w_1 and w_2 . Then S is an r-approximate solution of MAX-WEIGHTED-k-OBCS for G with respect to w.

By using this theorem we have the following result.

Theorem 4.2. MAX-WEIGHTED-*k*-OBCS can be approximated within a factor of Δ , where Δ is the maximum degree of the input graph *G*.

Proof. We prove the correctness by induction. In the base case, Line 2 is executed, and it is Δ -approximate. In the inductive step, when Line 9 is executed, S' is Δ -approximate with respect to w_2 . Now, the only vertices with a non zero weight in w_1 are the vertices s, t, and their neighbors. We will now prove that the weight $w_1(s) + w_1(t)$ is at least a $1/\Delta$ fraction of the total weight of $N[\{s,t\}]$ (with respect to w_1), which will prove the theorem. First note that since (s, t) was the edge with the maximum combined weight, we have that for any neighbour u of s, $w_1(u) = w(u) \le w(t) = w_1(t)$, and likewise for any neighbour v of t, we have $w_1(v) = w(v) \le w(s) = w_1(s)$. Now we have,

$$\frac{w_1(\{s,t\})}{w_1(N[\{s,t\}])} \ge \frac{w_1(s) + w_1(t)}{w_1(s)d_G(s) + w_1(t)d_G(t)} \ge \frac{1}{\Delta}$$

where the first inequality follows from our observation above, and the second inequality holds since the degree of each vertex is at most Δ .

Thus using the local ratio theorem, we get a Δ -factor approximation algorithm for MAX-WEIGHTED-*k*-OBCS.

It should be noted that due to the relation between the weighted and unweighted cases, the upper bound derived here holds even for the unweighted case.

5. Approximation algorithms for MAX-DISSO-SET

A subset of vertices $S \subseteq V$ in a graph G = (V, E) is called a dissociation set if it induces a subgraph with vertex degree of at most 1. In Theorem 3.6 we show that MAX-DISSO-SET is hard to approximate within a factor of $n^{1-\epsilon}$, unless P = NP. In this section, we prove that a greedy algorithm approximates MAX-DISSO-SET within a factor of $3\overline{d} + 2$.

In this algorithm, we keep track of the components of size 1 and 2 in G[S]. S_1 is the set of vertices in S which form components of size 1 and S_2 is the set of vertices which form components of size 2 in G[S]. Each time we include a vertex into S, we update the set X such that each component in $G[S \cup X]$ has at least 3 vertices.

We first prove the correctness of the Algorithm 5.1 and then turn to its approximation guarantee. Assume that the algorithm runs for *q* iterations. Since we include one vertex in each iteration, it is clear that the size of the solution set is *q*. Let v_i be the vertex included in the *i*th iteration and so the solution set returned is $S = \{v_1, \ldots, v_q\}$.

Lemma 5.1. At any iteration of the algorithm, for each vertex $v \in S_2$, $N_{V'}(v) \subseteq X$.

Proof. We prove this statement by induction on the iteration number. For the base case $S_2 = \emptyset$ and the statement trivially holds. Assume that the statement is true for the *i*th iteration. Let v_{i+1} be the vertex added to S in (i + 1)th iteration. If v_{i+1} has no neighbor in S_1 then S_2 remains unchanged and the statement holds. Otherwise, a new pair of vertices is added to S_2 and line 9 ensures that the neighbors of these two vertices are included in X. Thus the lemma holds in this case as well.

Algorithm 5.1: MAX-DISSO-SET

Input: An undirected simple graph G = (V, E)**Output:** $S \subseteq V$ such that each component in G[S] has at most 2 vertices 1 $S_1 = \emptyset; S_2 = \emptyset; X = \emptyset; S = \emptyset; V' = V; i = 1;$ ² while $[V' \neq \emptyset]$ do choose a vertex $v \in V'$ of minimum degree in G[V']; 3 $d_i = d_{G[V']}(v)$; 4 $S = S \cup \{v\};$ 5 if there exists a vertex $p \in S_1$ such that $(p, v) \in E$ then 6 $S_2 = S_2 \cup \{v, p\};$ 7 $S_1 = S_1 \setminus \{p\};$ $X = X \cup N_{V'}(p) \cup N_{V'}(v);$ 8 9 $V' = V \setminus (S \cup X);$ 10 else 11 $S_1 = S_1 \cup \{v\};$ $X = X \cup [\bigcup_{t \in S_1} [N_{V'}(v) \cap N_{V'}(t)]];$ $V' = V \setminus (S \cup X);$ 12 13 14 end 15 i = i + 1;16 17 end 18 Return S;

The correctness of the algorithm can be observed easily using the above lemma. Consider a solution set $S = S_1 \cup S_2$ returned by this algorithm. From Lemma 5.1, we can see that each of the vertices in S_2 are in a component of size exactly 2. Now consider a vertex $v \in S_1$. It cannot be a neighbor to a vertex in S_2 , again by lemma 5.1. In addition, it cannot have a neighbor in S_1 , since if it did, Line 7 would have moved these two vertices to the set S_2 during the run of the algorithm. Thus we see that the set S_1 forms an independent set in G[S] and thus the maximum size of a component in G[S] is 2, implying that S is a feasible solution.

Lemma 5.2. $n \leq \sum_{i=1}^{q} (d_i + 1) \leq 2n$, where *n* is the number of vertices in the input graph.

Proof. When we choose a vertex $v \in G[V']$ of minimum degree, we denote it as a *chosen* vertex and denote its neighbors in G[V'] as *looked-at* vertices. From the algorithm it is clear that a *chosen* vertex is always included in *S*. Every time a vertex is *looked-at*, exactly one of its neighbours is included in *S*. Once a vertex is *looked-at* twice, the algorithm includes it in the set *X*. Therefore, a vertex can be *looked-at* at most twice and can be *chosen* at most once.

Based on these notions, it follows that *X* is the set of vertices which are marked as *looked-at* twice by the algorithm. In the *i*th step of the algorithm a minimum degree vertex can be a *looked-at* vertex in *V'*. A chosen vertex v_i can make at most d_i vertices as *looked-at* twice. Thus inclusion of v_i into *S* can add at most d_i vertices into the set *X*. Also the set *X* is constructed in *q* steps. Therefore, $\sum_{i=1}^{q} d_i \ge |X|$. From this observation it follows that $\sum_{i=1}^{q} (d_i + 1) \ge |X| + |S| = n$.

The greedy algorithm finally partitions the vertex set *V* as $S \cup X$. Each vertex in *V* is probed at most twice (as *chosen*, or *looked-at* and *chosen*, or *looked-at* and *looked-at*) before

it is put in *S* or *X*. At the *i*th step $(d_i + 1)$ represents the number of *looked-at* vertices plus the *chosen* vertex. Therefore, $\sum_{i=1}^{q} (d_i + 1) \leq 2n$.

Lemma 5.3. $\sum_{i=1}^{q} (d_i + 1) d_i \leq 3n\overline{d}$.

Proof. Every time we choose a vertex $v \in V' = [V \setminus (S \cup X)]$, we assign a token to each edge *e* incident on a vertex of $N_{V'}[v]$.

Since the degree of vertex v in the *i*th step is d_i and it is of minimum degree, we assign at least $\frac{1}{2}d_i(d_i + 1)$ tokens in *i*th step. We claim that at the end of the algorithm, an edge can have at most 3 tokens. Let v_i be the vertex chosen in *i*th step. Let (p,q) be an edge in G[V'] such that $p \in N_{V'}(v_i)$ and $q \notin N_{V'}(v_i)$. While including v_i into S, we are assigning a token to (p,q) and it remain in the subgraph G[V'] with updated vertex set V'. If a neighbor of p or p is chosen as a vertex of smallest degree in any of the subsequent steps then (p,q) will get one more token and the updated graph G[V'] will not have this edge. So we assume that in a subsequent *j*th step let v_j be chosen as smallest degree vertex and it a neighbor of q and not a neighbor of p. While including v_j in S we are assigning the 2nd token to (p,q) and it remains as an edge in the updated subgraph G[V']. Till now both p and q are *looked-at* once each. In a subsequent *k*th step when a vertex $u \in N_{V'}[p] \cup N_{V'}[q]$ is chosen and included in S, the edge (p,q) receives the 3rd token and it will no more in the updated subgraph G[V'] (as p or q will be *looked-at* twice and will not be present in the updated set V').

Therefore, we have $\sum_{i=1}^{q} \frac{1}{2}(d_i+1)d_i \leq 3|\vec{E}| = \frac{3}{2}n\vec{d}$, and thus $\sum_{i=1}^{q}(d_i+1)d_i \leq 3n\vec{d}$.

Lemma 5.4. The solution set returned by Algorithm 5.1 has size at least $\frac{n}{3\overline{d}+2}$.

Proof. By using the inequalities in Lemma 5.2 and Lemma 5.3, we have

$$\sum_{i=1}^{q} (d_i + 1)^2 = \sum_{i=1}^{q} d_i (d_i + 1) + \sum_{i=1}^{q} (d_i + 1)$$

$$\leq 3n\overline{d} + 2n.$$

From the Cauchy-Schwarz inequality, we get

$$\sum_{i=1}^{q} (d_i + 1)^2 \ge \frac{\left(\sum_{i=1}^{q} (d_i + 1)\right)^2}{\sum_{i=1}^{q} 1^2}$$
$$\ge \frac{n^2}{q}.$$

Combining these two inequalities yields $q \ge \frac{n}{3\overline{d}+2}$.

From the above lemma, we have the following result.

Theorem 5.5. MAX-DISSO-SET can be approximated within a factor of $(3\overline{d} + 2)$, where \overline{d} is the average degree of the input graph *G*.

5.1. Approximation algorithm for MAX-k-OBCS

In this section, we will generalize the $(3\overline{d} + 2)$ factor approximation algorithm for MAX-DISSO-SET to get an algorithm for MAX-k-OBCS which is a $(2k - 1)\overline{d} + k$ factor approximation algorithm. The following is a generalization of Algorithm 5.1.

Algorithm 5.2: Algorithm for MAX- <i>k</i> -OBCS
Input: An undirected simple graph $G = (V, E)$
Output: $S \subseteq V$ such that each component in $G[S]$ has at most <i>k</i> vertices
1 $S = \emptyset; X = \emptyset; V' = V; i = 1;$
2 while $[V' \neq \emptyset]$ do
3 choose a vertex $v \in V'$ of minimum degree in $G[V']$;
$ 4 d_i = d_{G[V']}(v);$
$ 5 S = S \cup \{v\};$
6 for each vertex $p \in V'$ and $p \neq v$ do
7 if $G[S \cup \{p\}]$ has a component of size at least $k + 1$ then
$ \mathbf{s} X = X \cup \{p\};$
9 end
10 end
$11 V' = V \setminus (S \cup X);$
12 $i = i + 1;$
13 end
14 Return S;

By using a similar argument we have the following results.

Lemma 5.6. Let q be the number of iterations made by the while loop in the Algorithm 5.2. (a) $n \leq \sum_{i=1}^{q} (d_i + 1) \leq nk$ (b) $\sum_{i=1}^{q} d_i (d_i + 1) \leq \overline{d}(2k - 1)n$.

Proof. (a) By assigning token to the vertices, as given in the proof of Lemma 5.2, it can be proved that $\sum_{i=1}^{q} d_i \ge |X|$. Hence, we get $n \le \sum_{i=1}^{q} (d_i + 1)$. For the other inequality, consider the following argument. In each iteration, when a vertex v is picked, assign one token to that vertex and each of its neighbors in the graph G[V']. Thus, in the *i*th iteration, the number of tokens assigned is exactly $d_i + 1$, and the total number of tokens assigned is $\sum_{i=1}^{q} (d_i + 1)$. Now, we look at the maximum number of tokens that any particular vertex can get. Note that a vertex gets a token if it is either *chosen* or *looked-at*, and once a vertex is *chosen* or included in the set X, it gets no more tokens. Suppose that at the end of the algorithm, a vertex u has (k + 1) or more tokens. This implies that either it was *looked-at* at least k + 1 times and then moved to the set X or it was *looked-at* at least k times and then chosen. Neither of these cases is possible, since when the vertex u is *looked-at* k times k of its neighbors have been *chosen*, and thus the Line 6 will remove u after k of its neighbors have been *chosen*. Thus, any vertex can have at most k tokens, and thus the total number of tokens available is at most nk. This proves the second inequality.

(b) In the *i*th iteration, assume that vertex *v* is *chosen*. Assign a token to all the edges incident on *v* and its neighbors from the set *V'* of the *i*th iteration. As argued in the case of the dissociation set problem, due to minimality of d_i , we assign at least $\sum_{i=1}^{q} \frac{1}{2}d_i(d_i + 1)$

1) tokens. Now, we produce an upper bound on the total number of tokens by seeing the maximum number of tokens that can be assigned to any edge. Consider an edge e = (u, v) - it gets a token whenever at least one of its endpoints is either *looked-at* or *chosen*. Now, from the proof of (a), it follows that any vertex is *looked-at* at most k times and put into the set X, or *looked-at* at most k - 1 times and then *chosen*. Thus the edge e can have at most (2k - 1) tokens, since if it receives that amount, then at least one of its endpoints has been moved to either the set X or the set S and the edge no longer receives any tokens. Thus the total number of tokens that any edge can receive is (2k - 1) and the total number of tokens given out is at most |E|(2k - 1). Thus we get $\sum_{i=1}^{q} d_i(d_i + 1) \leq 2|E|(2k - 1) = \overline{d}(2k - 1)n$.

By using the inequalities in Lemma 5.6 and Cauchy-Schwarz inequality, we have

Theorem 5.7. MAX-*k*-OBCS can be approximated within a factor of $(2k - 1)\overline{d} + k$, where \overline{d} is the average degree of the input graph *G*.

6. Upper bound for MAX-k-OBCS with large Δ

In this section, we provide a reduction from the MAX-DISSO-SET problem to the MAX-IS problem, and use the upper bound result for MAX-IS from [17] to provide a $O(\frac{\Delta \log \log \Delta}{\log \Delta})$ approximation factor, for sufficiently large values of Δ .

Lemma 6.1. MAX-DISSO-SET \leq_{AP} MAX-IS.

Proof. The reduction is a straight forward one, where we solve the MAX-IS on the input instance. Formally, the reduction is the identity mapping, and given an instance G = (V, E) of MAX-DISSO-SET, it maps G = (V, E) as an instance of MAX-IS.

Given an independent set I in G, we construct a dissociation set S = I from I with |S| = |I|. Given a maximum cardinality dissociation set S^* in G, we construct an independent set I with $|I| \ge \frac{1}{2}|S^*|$ as follows. The induced graph $G[S^*]$ will have maximum degree of at most 1, and we can partition the set S^* into two sets $S_0 = \{v \in S^* | d_{G[S^*]}(v) = 0\}$ and $S_1 = \{v \in S^* | d_{G[S^*]}(v) = 1\}$. It is important to observe that $G[S_1]$ forms an induced matching. We construct an independent set I in G by taking all the vertices in S_0 and exactly one vertex from each edge in $G[S_1]$. This set I of vertices is an independent set and $|I| = |S_0| + \frac{1}{2}|S_1| \ge \frac{1}{2}(|S_0| + |S_1|) = \frac{1}{2}|S^*|$. Thus we have $|S^*| \le 2|I| \le 2\alpha(G)$, yielding

$$\frac{|S^*|}{|S|} \le 2\frac{\alpha(G)}{|I|}.$$

Theorem 6.2. [17] MAX-IS can be approximated within a factor of $O(\frac{\Delta \log \log \Delta}{\log \Delta})$ for sufficiently large values of Δ , where Δ is the maximum degree of the input graph.

From Lemma 6.1 and Theorem 6.2, we get the following result,

Theorem 6.3. MAX-DISSO-SET is approximable within a factor of $O(\frac{\Delta \log \log \Delta}{\log \Delta})$ for sufficiently large values of Δ , where Δ is the maximum degree of the input graph.

The reduction in the proof of Lemma 6.1 is also a reduction from MAX-*k*-OBCS to MAX-IS. Here, given an instance G = (V, E) of MAX-*k*-OBCS, we take G = (V, E) as an instance of MAX-IS. If *I* is an independent set in *G* then S = I is also a *k*-component set in *G*. From a maximum cardinality *k*-component set S^* in the graph *G*, we can construct an independent set *I* in *G* with $|I| \ge \frac{1}{k}|S^*|$. Therefore, we have $|S^*| \le k|I| \le k\alpha(G)$. From these inequalities we have

$$\frac{|S^*|}{|S|} \le k \frac{\alpha(G)}{|I|}$$

and the following result.

Theorem 6.4. For k > 0, MAX-k-OBCS is approximable within a factor of $O(\frac{k\Delta \log \log \Delta}{\log \Delta})$ for sufficiently large values of Δ , where Δ is the maximum degree of the input graph.

7. Conclusion

We have proved that MAX-*k*-OBCS is hard to approximate like MAX-IS. We also show that for $k = O(\frac{n}{\log n})$, MAX-*k*-OBCS is hard to approximate within a factor of $O(\log n)$. We generalize the greedy algorithm for MAX-IS to obtain an algorithm for MAX-*k*-OBCS with an approximation factor of $(2k - 1)\overline{d} + k$. This approximation factor is a generalization of Turán's bound for MAX-IS.

References

- Y. Orlovich, A. Dolgui, G. Finke, V. Gordon, F. Werner, The complexity of dissociation set problems in graphs, Discrete Applied Mathematics 159 (13) (2011) 1352–1366.
- [2] J. Håstad, Clique is hard to approximate within $n^{1-\epsilon}$, in: Foundations of Computer Science, 1996. Proceedings., 37th Annual Symposium on, IEEE, 1996, pp. 627–636.
- [3] D. Zuckerman, Linear degree extractors and the inapproximability of max clique and chromatic number, in: Proceedings of the thirty-eighth annual ACM symposium on Theory of computing, ACM, 2006, pp. 681–690.
- [4] D. Nakamura, A. Tamura, A revision of minty's algorithm for finding a maximum weight stable set of a claw-free graph, Journal of the Operations Research Society of Japan 44 (2) (2001) 194–204.
- [5] D. Lokshantov, M. Vatshelle, Y. Villanger, Independent set in P₅-free graphs in polynomial time, in: Proceedings of the Twenty-Fifth Annual ACM-SIAM Symposium on Discrete Algorithms, Society for Industrial and Applied Mathematics, 2014, pp. 570–581.
- [6] M. Grötschel, L. Lovász, A. Schrijver, Geometric algorithms and combinatorial optimization, Vol. 2, Springer Science & Business Media, 2012, pp. 296–298.
- [7] B. S. Baker, Approximation algorithms for np-complete problems on planar graphs, Journal of the ACM (JACM) 41 (1) (1994) 153–180.
- [8] C. H. Papadimitriou, M. Yannakakis, Optimization, approximation, and complexity classes, Journal of computer and system sciences 43 (3) (1991) 425–440.
- [9] D. S. Hochbaum, Efficient bounds for the stable set, vertex cover and set packing problems, Discrete Applied Mathematics 6 (3) (1983) 243–254.
- [10] M. M. Halldórsson, J. Radhakrishnan, Greed is good: Approximating independent sets in sparse and bounded-degree graphs, Algorithmica 18 (1) (1997) 145–163.
- [11] M. Yannakakis, Node-deletion problems on bipartite graphs, SIAM Journal on Computing 10 (2) (1981) 310–327.
- [12] R. Boliac, K. Cameron, V. V. Lozin, On computing the dissociation number and the induced matching number of bipartite graphs, Ars Comb. 72.

- [13] C. H. Papadimitriou, M. Yannakakis, The complexity of restricted spanning tree problems, Journal of the ACM (JACM) 29 (2) (1982) 285–309.
- [14] T. Fujito, A unified approximation algorithm for node-deletion problems, Discrete applied mathematics 86 (2) (1998) 213–231.
- [15] Y. Zhang, Y. Shi, Z. Zhang, Approximation algorithm for the minimum weight connected k-subgraph cover problem, Theoretical Computer Science 535 (2014) 54–58.
- [16] R. Bar-Yehuda, K. Bendel, A. Freund, D. Rawitz, Local ratio: A unified framework for approximation algorithms. in memoriam: Shimon even 1935-2004, ACM Computing Surveys (CSUR) 36 (4) (2004) 422– 463.
- [17] M. M. Halldórsson, Approximations of weighted independent set and hereditary subset problems, in: Graph Algorithms And Applications 2, World Scientific, 2004, pp. 3–18.
- [18] G. Ausiello, Complexity and Approximability Properties: Combinatorial Optimization Problems and Their Approximability Properties, Springer, 1999.