

BOUNDING PERIODS OF SUBVARIETIES OF $(\mathbb{P}^1)^n$

J. P. BELL, D. GHIOCA, AND T. J. TUCKER

ABSTRACT. Using methods of p -adic analysis, along with the powerful result of Medvedev-Scanlon [MS14] for the classification of periodic subvarieties of $(\mathbb{P}^1)^n$, we bound the length of the orbit of a periodic subvariety $Y \subset (\mathbb{P}^1)^n$ under the action of a dominant endomorphism.

1. INTRODUCTION

In [MS94], Morton and Silverman conjecture that there is a constant $C(N, d, D)$ such that for any morphism $f : \mathbb{P}^N \rightarrow \mathbb{P}^N$ of degree d defined over a number field K with $[K : \mathbb{Q}] \leq D$, the number of preperiodic points of f over K is less than or equal to $C(N, d, D)$. This conjecture remains very much open, but in the case where f has good reduction at a prime \mathfrak{p} , a great deal has been proved about bounds depending on \mathfrak{p} , N , d , D (see [Zie96, Pez05, Hut09]).

In this paper, we study the more general problem of bounding length of the orbit of periodic subvarieties of any dimension; we prove the following results.

Theorem 1.1. *Let $n \in \mathbb{N}$, let K be a finite extension of \mathbb{Q}_p , let κ be its residue field, and let e be the ramification index of K/\mathbb{Q}_p . Let $f_1, \dots, f_n \in K(x)$ be rational functions of good reduction modulo p and let $\Phi : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^n$ be the endomorphism given by*

$$(1.1.1) \quad (x_1, \dots, x_n) \mapsto (f_1(x_1), \dots, f_n(x_n)).$$

Let Y be a subvariety of $(\mathbb{P}^1)^n$, and assume there is a nonsingular point $(\alpha_1, \dots, \alpha_n) \in Y(K)$. Then there exists a constant C depending on n , p , e , $[\kappa : \mathbb{F}_p]$ and the degree of the rational functions f_i (but independent of Y) such that if Y is periodic under the action of Φ , then the length of its orbit is bounded by C .

We prove Theorem 1.1 as a consequence of the next result.

Theorem 1.2. *Let K be a finite extension of \mathbb{Q}_p , let κ be its residue field and let e be the ramification index of K/\mathbb{Q}_p . Let $f_1, \dots, f_n \in K(x)$ be rational functions of good reduction modulo p and let $\Phi : (\mathbb{P}^1)^n \rightarrow (\mathbb{P}^1)^n$ be the endomorphism given by*

$$(1.2.1) \quad (x_1, \dots, x_n) \mapsto (f_1(x_1), \dots, f_n(x_n)).$$

Let Y be a subvariety of $(\mathbb{P}^1)^n$, and assume there is a point $(\alpha_1, \dots, \alpha_n) \in Y(K)$ such that for each $i = 1, \dots, n$, we have that α_i is a nonpreperiodic point for f_i . Then there exists a constant C depending on n , p , e , $[\kappa : \mathbb{F}_p]$, and the degree of the rational functions f_i (but independent of Y) such that if Y is periodic under the action of Φ , then the length of its orbit is bounded by C .

We note that any dominant (regular) endomorphism Φ of $(\mathbb{P}^1)^n$ is split, i.e., it is of the form

$$(x_1, x_2, \dots, x_n) \mapsto (f_1(x_{\sigma(1)}), f_2(x_{\sigma(2)}), \dots, f_n(x_{\sigma(n)})),$$

where the f_i 's are rational functions, while σ is a permutation of $\{1, \dots, n\}$. So, at the expense of replacing Φ by an iterate Φ^k (for some positive integer k dividing $n!$), each dominant endomorphism of $(\mathbb{P}^1)^n$ is of the form (1.2.1).

In [BGT15], a similar result was obtained bounding the length of the orbit of any periodic subvariety (with a distinguished smooth p -adic point of good reduction) under the action of an étale endomorphism of an arbitrary variety (not only $(\mathbb{P}^1)^n$). However, the presence of ramification as it is in our case adds new complications. Also, we note that in [GN17], using results regarding the decomposition of polynomials, it was shown that the length of the orbit of a periodic subvariety of \mathbb{A}^n under the coordinatewise action of n one-variable polynomials is uniformly bounded by a constant depending only on n and on the degrees of the polynomials. It is important to point out that the method from [GN17] cannot be extended to treat the case of rational functions; essentially, for rational functions, one lacks all the sharper results regarding their decomposition, as established in [MS14] for the case of polynomials.

In our Theorem 1.2, the constant C bounding the length of the orbit of a periodic subvariety of $(\mathbb{P}^1)^n$ depends on the degrees of the rational functions f_i and also on the good reduction data associated to our problem, which refers to the knowledge about a single point on Y . So, from this point of view, our result is less restrictive than the one obtained in [Hut], where a similar bound as the one from our Theorem 1.2 is established under the additional assumption that also the periodic subvariety Y (along with its iterates under Φ) have good reduction modulo p , i.e., reducing modulo p yields a subvariety \bar{Y} , which is an irreducible subvariety defined over $\overline{\mathbb{F}_p}$ of same degree as Y (and a similar property holds for each iterate of Y under Φ). Even though the result from [Hut] is valid for more general dynamical systems (i.e., endomorphisms of \mathbb{P}^n as opposed to endomorphisms of $(\mathbb{P}^1)^n$), the additional hypothesis regarding the good behaviour of the periodic subvariety modulo p makes it more restrictive than our Theorem 1.2.

Theorem 1.2 is proved using a p -adic analytic parametrization of forward orbits under the action of a rational function (such as in [BGKT10]), combined with the description of periodic subvarieties of $(\mathbb{P}^1)^n$ (obtained by Medvedev-Scanlon [MS14]).

We start by describing in Section 2 the various useful results regarding the p -adic parametrization of orbits of points under the action of a rational function. We continue by proving in Section 3 the special case of Theorem 1.2 when the periodic subvariety is a curve contained in $(\mathbb{P}^1)^2$ (see Theorem 3.1). We conclude our paper by reducing the general Theorem 1.2 to the special Theorem 3.1, using the description of periodic subvarieties of $(\mathbb{P}^1)^n$ provided by Medvedev-Scanlon in [MS14]; then we derive the conclusion of Theorem 1.1 as a consequence of Theorem 1.2.

2. USEFUL RESULTS

In this Section, we follow the setup and the results from [BGKT10, Section 2]; see also [BGT16, Chapter 6]. For a prime p , we let \mathbb{Z}_p denote the ring of p -adic integers, \mathbb{Q}_p will denote the field of p -adic rationals, and \mathbb{C}_p will denote the completion of an algebraic closure of \mathbb{Q}_p with respect to the p -adic absolute value $|\cdot|_p$. Given a point $y \in \mathbb{C}_p$ and a real number $r > 0$, write

$$D(y, r) = \{x \in \mathbb{C}_p : |x - y|_p < r\}, \quad \overline{D}(y, r) = \{x \in \mathbb{C}_p : |x - y|_p \leq r\}$$

for the open and closed disks, respectively, of radius r about y in \mathbb{C}_p . We write $[y] \subseteq \mathbb{P}^1(\mathbb{C}_p)$ for the residue class of a point $y \in \mathbb{P}^1(\mathbb{C}_p)$. That is, $[y] = D(y, 1)$ if $|y| \leq 1$, or else $[y] = \mathbb{P}^1(\mathbb{C}_p) \setminus \overline{D}(0, 1)$ if $|y| > 1$.

Let K be a finite extension of \mathbb{Q}_p , let \mathfrak{o}_K be its ring of p -adic integers, let π be a uniformizer for \mathfrak{o}_K and let κ be its corresponding residue field. Let $f \in K(x)$ be a rational function of degree $d \geq 1$. We say that f has good reduction modulo π (or equivalently modulo p) if writing the rational function f as

$$f([X_0 : X_1]) = [f_0(X_0, X_1) : f_1(X_0, X_1)]$$

for some coprime homogeneous polynomials $f_0, f_1 \in \mathfrak{o}_K[X_0, X_1]$ of degree d for which at least one coefficient is a π -adic unit, then reducing the coefficients of both f_0 and f_1 modulo π (or equivalently, modulo p) yields two coprime homogeneous polynomials in $\kappa[X_0, X_1]$ of same degree d . Note that our choice of coordinates here induces a choice of model $\mathbb{P}_{\mathfrak{o}_K}^1$ and a morphism $F : \mathbb{P}_{\mathfrak{o}_K}^1 \rightarrow \mathbb{P}_{\mathfrak{o}_K}^1$ of \mathfrak{o}_K -schemes such that $(F)_K = f$ when f has good reduction at π .

The action of a p -adic power series $f \in \mathfrak{o}_K[[z]]$ on $D(0, 1)$ is either attracting (i.e., f contracts distances) or quasiperiodic (i.e., f is distance-preserving), depending on its linear coefficient. Rivera-Letelier gives a more precise description of this dichotomy in [RL03, Sections 3.1 and 3.2]. The following two Lemmas follow from [RL03, Propositions 3.3 and 3.16] and they were stated in [BGKT10, Lemmas 2.1 and 2.2] in the case $K = \mathbb{Q}_p$; the general case follows by an identical argument (for more details regarding the applications to arithmetic dynamics of the p -adic parametrization of orbits, see [BGT16]).

Lemma 2.1. *Let K be a finite extension of \mathbb{Q}_p , let \mathfrak{o}_K be its ring of p -adic integers, and let π be a uniformizer for \mathfrak{o}_K . We let $f(z) = a_0 + a_1z + a_2z^2 + \dots \in \mathfrak{o}_K[[z]]$ be a nonconstant power series with $|a_0|_p, |a_1|_p < 1$. Then there is a point $y \in \pi \cdot \mathfrak{o}_K$ such that $f(y) = y$, and $\lim_{n \rightarrow \infty} f^n(z) = y$ for all $z \in D(0, 1)$. Write $\lambda = f'(y)$; then $|\lambda|_p < 1$, and:*

- (i) *(Attracting). If $\lambda \neq 0$, then there is a radius $0 < r < 1$ and a power series $u \in K[[z]]$ mapping $\overline{D}(0, r)$ bijectively onto $\overline{D}(y, r)$ with $u(0) = y$, such that for all $z \in D(y, r)$ and $n \geq 0$,*

$$f^n(z) = u(\lambda^n u^{-1}(z)).$$

- (ii) *(Superattracting). If $\lambda = 0$, then write f as*

$$f(z) = y + c_m(z - y)^m + c_{m+1}(z - y)^{m+1} + \dots \in \mathfrak{o}_K[[z - y]]$$

with $m \geq 2$ and $c_m \neq 0$. If c_m has an $(m-1)$ -st root in K (or equivalently, in \mathfrak{o}_K), then there are radii $0 < r, s < 1$ and a power series $u \in K[[z]]$ mapping $\overline{D}(0, s)$ bijectively onto $\overline{D}(y, r)$ with $u(0) = y$, such that for all $z \in D(y, r)$ and $n \geq 0$,

$$f^n(z) = u\left((u^{-1}(z))^{m^n}\right).$$

Proof. The proof is identical to the proof of [BGKT10, Lemma 2.1]; see also [SS] for a similar result in the superattracting case. \square

The next result is a slight generalization of [BGKT10, Lemma 2.2].

Lemma 2.2. *Let K , \mathfrak{o}_K and π be as in Lemma 2.1 and let κ be the residue field of K . Let $f(z) = a_0 + a_1z + a_2z^2 + \dots \in \mathfrak{o}_K[[z]]$ be a nonconstant power series*

with $|a_0|_p < 1$ but $|a_1|_p = 1$. Then for any nonperiodic $x \in \pi \cdot \mathfrak{o}_K$, there are: an integer $k \geq 1$ depending only on p , e and $[\kappa : \mathbb{F}_p]$, radii $0 < r < 1$ and $s \geq |k|_p$, and a power series $u \in K[[z]]$ mapping $\overline{D}(0, s)$ bijectively onto $\overline{D}(x, r)$ with $u(0) = x$, such that for all $z \in \overline{D}(x, r)$ and $n \geq 0$,

$$f^{nk}(z) = u(nk + u^{-1}(z)).$$

Proof. The conclusion of Lemma 2.2 (including the fact that k is bounded solely in terms of p , e and $[\kappa : \mathbb{F}_p]$) follows by combining [BGT15, Proposition 2.1] with [Poo13]. \square

3. THE CASE OF CURVES

We first prove Theorem 1.2 when Y is a curve in $\mathbb{P}^1 \times \mathbb{P}^1$.

Theorem 3.1. *Let K be a finite extension of \mathbb{Q}_p , let κ be its residue field and let e be the ramification index of K/\mathbb{Q}_p . Let $f_1, f_2 \in K(x)$ be rational functions of good reduction modulo p and let $\Phi : (\mathbb{P}^1)^2 \rightarrow (\mathbb{P}^1)^2$ be the endomorphism given by $(x_1, x_2) \mapsto (f_1(x_1), f_2(x_2))$. Let Y be an irreducible, periodic curve of $(\mathbb{P}^1)^2$ defined over K , and assume there is a point α on $Y(K)$ which is not preperiodic under the action of Φ . Then there exists a constant c depending on the degrees of the rational functions f_i and also depending on p , e and $[\kappa : \mathbb{F}_p]$, but independent of Y , such that the length of the orbit of Y under Φ is bounded by c .*

Proof. We let $\alpha := (\alpha_1, \alpha_2) \in \mathbb{P}_K^1 \times \mathbb{P}_K^1$. If for some $i = 1, 2$, we have that α_i is preperiodic for f_i , then since α is not preperiodic under the action of Φ but Y is periodic under the action of Φ , we conclude that the curve Y projects to α_i on the i -th coordinate axis. Therefore, the length of the orbit of Y is the length of the orbit of α_i (under the action of f_i). Because $\alpha_i \in \mathbb{P}_K^1$ is a periodic point for f_i and moreover, f_i has good reduction modulo p , then the length of the orbit of α_i is bounded only in terms of p , e , $[\kappa : \mathbb{F}_p]$, as proven in [Zie96, Hut09]; hence the length of the orbit of Y is bounded by a constant which is independent of Y .

So, from now on, we may assume that each α_i (for $i = 1, 2$) is not preperiodic.

At the expense of replacing K by a finite extension of degree bounded solely in terms of the degrees of the rational functions f_1 and f_2 (and therefore replacing e and $[\kappa : \mathbb{F}_p]$ by multiples of them bounded by a quantity depending again only in terms of the degrees of the f_i 's), we may assume the following. For some $i = 1, 2$ if α_i modulo π is in the cycle of a periodic critical point Q of f_i of period ℓ , given a local coordinate $x_{i,Q}$ at Q , we write

$$(3.1.1) \quad f_i^\ell(x_{i,Q}) = c_{i,Q} x_{i,Q}^{m_{i,Q}} + O(x_{i,Q}^{m_{i,Q}+1}),$$

where $m_{i,Q} \geq 2$ and $c_{i,Q} \neq 0$; then K contains all the $(m_{i,Q} - 1)$ -st roots of $c_{i,Q}$. Since there are at most finitely many superattracting periodic points, then we are guaranteed that a finite extension of K suffices. If there is no superattracting periodic point modulo p in the orbit of α_i under the action of f_i modulo π , then we do not need to extend K .

We let \mathfrak{o}_K be the subring of p -adic integers in K , and we let π be a uniformizer for \mathfrak{o}_K .

We proceed as in the proof of [BGKT10, Theorem 1.4] (see also [BGT16, Chapter 11]); our goal is to show that there exists a positive integer k depending only on data coming from K (i.e, its ramification index and also the size of its residue

field) and also there exists a positive integer ℓ (which may depend on the α_i 's) such that for each $i = 1, 2$ and for each $0 \leq j \leq k - 1$ there exist positive integers m_i and p -adic analytic power series $F_{i,j} \in K[[z_0, z_1, z_{m_i}]]$ such that (modulo linear transformations) we have p -adic analytic parametrizations:

$$(3.1.2) \quad f_i^{\ell+j+nk}(\alpha_i) = F_{i,j} \left(n, \pi^n, \pi^{m_i^n} \right) \text{ for each } n \geq 0.$$

The integers m_i from (3.1.2) are the same as the integers $m_{i,Q}$ from equation (3.1.1); if α_i is not superattracting, then actually $F_{i,j}$ is a p -adic analytic power series only in the variables z_0 and z_1 , or alternatively, $m_i = 1$ in that case.

In order to derive the p -adic parametrization (3.1.2), we proceed as in the proof of [BGKT10, Theorem 1.4]; in particular, we split our analysis into various steps.

Step (i). First we note that there exist positive integers ℓ_0 and k_0 such that the residue class of $f_i^{\ell_0}(\alpha_i)$ (for $i = 1, 2$) is fixed under the induced action on \mathbb{P}_κ^1 of the reduction of $f_i^{k_0}$ modulo π ; here we use the fact that each f_i has good reduction modulo π . Furthermore, note that both integers k_0 and ℓ_0 depend only on p and on $[\kappa : \mathbb{F}_p]$; for more details, see [BGKT10, Step (ii), p. 1063]. By a $\text{PGL}(2, \mathfrak{o}_K)$ -change of coordinates for each $i = 1, 2$, we may assume that $f_i^{\ell_0}(\alpha_i) \in \pi \mathfrak{o}_K$, and therefore $f_i^{k_0}$ may be written as a nonconstant power series in $\mathfrak{o}_K[[z]]$ mapping $D(0, 1)$ into itself.

Step (ii). Let $i \in \{1, 2\}$. If $\left| f_i^{k_0} \left(f_i^{\ell_0}(\alpha_i) \right) \right|_p < 1$, then Lemma 2.1 yields that there is a point $y_i \in D(0, 1)$ fixed by $f_i^{k_0}$ along with radii r_i and s_i (where $s_i := r_i$ in the non-superattracting case), and an associated power series $u_i \in K[[z]]$. Set $k_{1,i} = k_0$ and $\ell_{1,i} = \ell_0 + n_i k_{1,i}$ for a suitable integer $n_i \geq 0$ so that

$$(3.1.3) \quad f_i^{\ell_{1,i}}(\alpha_i) \in \overline{D}(y_i, r_i).$$

Define $\lambda_i := (f_i^{k_0})'(y_i)$ to be the multiplier of the point y_i ; then in this case, $|\lambda_i|_p < 1$. Define $\mu_i := u_i^{-1}(f_i^{\ell_{1,i}}(\alpha_i))$; note that $\mu_i \in \pi \cdot \mathfrak{o}_K$, because $s_i < 1$. In addition, $\mu_i \neq 0$, because u_i is bijective and $u_i(0) = y_i$ is fixed by $f_i^{k_0}$, while $u_i(\mu_i) = f_i^{\ell_{1,i}}(\alpha_i)$ is not fixed by $f_i^{k_0}$.

If $|(f_i^{k_0})'(f_i^{\ell_0}(\alpha_i))|_p = 1$, we apply Lemma 2.2 to $f_i^{k_0}$ and the point $f_i^{\ell_0}(\alpha_i)$ to obtain radii r_i and s_i and a power series u_i . Define $\mu_i := u_i^{-1}(f_i^{\ell_0}(\alpha_i))$, and set $\ell_1 = \ell_0$ and $k_{3,i} = n_i k_0$, for a suitable integer $n_i \geq 1$ depending only on p , e and $[\kappa : \mathbb{F}_p]$ (see Lemma 2.2 in which the integer k from its conclusion depends only on p , e and $[\kappa : \mathbb{F}_p]$) so that $f_i^{k_{3,i} + \ell_1}(\alpha_i) \in \overline{D}(f_i^{\ell_1}(\alpha_i), r_i)$. (Following the notation from [BGKT10], the jump from a subscript of 0 to 3 is because certain complications, to be addressed in Steps (iii) and (iv), do not arise in the quasiperiodic case.) Note that $f_i^{\ell_{1,i} + n k_{3,i}}(\alpha_i)$ may be expressed as a power series in the integer $n \geq 0$; specifically, $f_i^{\ell_{1,i} + n k_{3,i}}(\alpha_i) = u_i(n k_{3,i} + \mu_i)$.

Step (iii). In this step, we consider only the case that $0 < |\lambda_i|_p < 1$ (i.e., attracting but not superattracting). We will express certain functions of n as power series in n and π^n and thus obtain a p -adic parametrization of the orbit of α_i under f_i as in (3.1.2), where the corresponding p -adic analytic function $F_{i,j}$ is actually a function of only 2 variables (i.e., $m_i = 1$ with the notation as in (3.1.2)).

Write $\lambda_i = \alpha_i \pi^{e_i}$, where $e_i \geq 1$ and $\alpha_i \in \mathfrak{o}_K^*$. If α_i is a root of unity, we can choose an integer $M_{1,i} \geq 1$ such that $\alpha_i^{M_{1,i}} = 1$. If α_i is not a root of unity, it is well known that there is an integer $M_{1,i} \geq 1$ such that $\alpha_i^{n M_{1,i}}$ can be written as a

power series in n with coefficients in \mathfrak{o}_K . Furthermore, $M_{1,i}$ depends only on p , e and $[\kappa : \mathbb{F}_p]$, which can be seen also applying Lemma 2.2 to the function $z \mapsto \alpha_i z$; for more details, see [BGKT10, Step (iii)]. We set

$$(3.1.4) \quad k_{3,i} := M_{1,i} k_{1,i};$$

also, we replace λ_i by $\lambda_i^{M_{1,i}}$ and replace e_i by $M_{1,i} \cdot e_i$. (The subscript again jumps to 3 in (3.1.4) because of the complications of Step (iv).) Thus, we can write

$$(3.1.5) \quad \lambda_i^n = (\pi^n)^{e_i} g_{1,i}(n) \quad \text{for all integers } n \geq 0,$$

for some power series $g_{1,i}(z) \in \mathfrak{o}_K[[z]]$.

Step (iv). In this step, we consider only the superattracting case, that $\lambda_i = 0$, and we will express certain functions of n as power series in n , π^n , and $\pi^{m_i^n}$, where $m_i \geq 2$ is a certain integer (see (3.1.1)).

We let first the integer $m_{i,0}$ be the order of the unique superattracting point of f_i in $D(0,1)$, as in Lemma 2.1(ii). Then we write $m_{i,0} := a_i \pi^{b_i}$, for some $a_i \in \mathfrak{o}_K^*$ and some integer $b_i \geq 0$. Then as in Step (iii), we can find a positive integer $M_{1,i}$ depending only on p , e and $[\kappa : \mathbb{F}_p]$ such that $a_i^{n M_{1,i}}$ can be written as a power series in n with coefficients in \mathfrak{o}_K . Set

$$k_{2,i} := M_{1,i} \cdot k_{1,i} \quad \text{and} \quad m_{i,1} := m_{i,0}^{M_{1,i}}.$$

Then $m_{i,1}^n$ can be written as a power series in n and π^n , with coefficients in \mathfrak{o}_K .

In addition, recall that $\mu_i = u_i^{-1}(f_i^{\ell_i}(\alpha_i))$ satisfies $0 < |\mu_i|_p < 1$; thus, we can write $\mu_i = \beta_i \pi^{e_i}$, where $e_i \geq 1$ and $\beta_i \in \mathfrak{o}_K^*$. If β_i is a root of unity with, say, $\beta_i^{M_{2,i}} = 1$ for some positive integer $M_{2,i}$ (depending only on p , e and $[\kappa : \mathbb{F}_p]$, since the size of the group of roots of unity contained in K depends only on p , e and $[\kappa : \mathbb{F}_p]$), choose a positive integer $M_{3,i}$ (again depending only on p , e and $[\kappa : \mathbb{F}_p]$) so that $M_{2,i} | (m_{i,1}^{2M_{3,i}} - m_{i,1}^{M_{3,i}})$. Set

$$k_{3,i} := M_{3,i} k_{2,i} \quad \text{and} \quad m_{i,2} := m_{i,1}^{M_{3,i}}$$

and note that $\beta_i^{m_{i,2}^n}$ is constant in n .

On the other hand, if β_i is not a root of unity, then as in Step (iii), there is an integer $M'_{2,i}$ depending only on p , e and $[\kappa : \mathbb{F}_p]$ such that $\beta_i^{n M'_{2,i}}$ can be written as a power series in n with coefficients in \mathfrak{o}_K . As above, choose a positive integer $M'_{3,i}$ (depending only on p , e and $[\kappa : \mathbb{F}_p]$) such that $M'_{2,i} | (m_{i,1}^{2M'_{3,i}} - m_{i,1}^{M'_{3,i}})$, and set

$$k_{3,i} := M'_{3,i} k_{2,i} \quad \text{and also,} \quad m_{i,2} := m_{i,1}^{M'_{3,i}}.$$

Then $m_{i,2}^n \equiv m_{i,2} \pmod{M'_{2,i}}$ for all positive integers n , and therefore

$$\beta_i^{m_{i,2}^n} = \beta_i^{m_{i,2}} \cdot \beta_i^{m_{i,2}^n - m_{i,2}} = \beta_i^{m_{i,2}} \cdot \left(\beta_i^{M'_{2,i}} \right)^{(m_{i,2}^n - m_{i,2})/M'_{2,i}}$$

can be written as a power series in $(m_{i,2}^n - m_{i,2})/M'_{2,i}$ with coefficients in \mathfrak{o}_K . Using the fact that $p \nmid M'_{2,i}$, and expressing $m_{i,2}^n = (m_{i,1}^n)^{M'_{3,i}}$ as a power series in n and π^n with coefficients in \mathfrak{o}_K , we conclude that $\beta_i^{m_{i,2}^n}$ can in fact be written as a power series in n and π^n , with coefficients in \mathfrak{o}_K .

Thus, whether or not β_j is a root of unity, we can write

$$(3.1.6) \quad \mu_i^{m_{i,2}^n} = \left(\pi^{m_{i,2}^n} \right)^{e_i} g_{1,i}(n, \pi^n) \quad \text{for all integers } n \geq 0,$$

for some power series $g_{1,i}(z_0, z_1) \in \mathfrak{o}_K[[z_0, z_1]]$.

Step (v). Let k be the least common multiple of $k_{3,1}$ and of $k_{3,2}$; then k is bounded by a constant depending only on p , e and $[\kappa : \mathbb{F}_p]$.

Note that replacing each $k_{3,i}$ by k does not change the conclusions of our previous Steps since the radii r_i , s_i , the power series u_i , the integer $\ell_{1,i}$, and the point μ_i are unaffected when we replace k by a multiple of it. Moreover, in the quasiperiodic case, $f_i^{k+\ell_{1,i}}(\alpha_i)$ still lies in $\overline{D}(f_i^{\ell_{1,i}}(\alpha_i), r_i)$.

In the attracting case, we replace λ_i by $\lambda_i^{k/k_{3,i}}$, also replace e_i by $\frac{k}{k_{3,i}}e_i$, and let

$$g_{2,i}(z) := (g_{1,i}(z))^{k/k_{3,i}} \in \mathfrak{o}_K[[z]];$$

and in the superattracting case, we replace m_i by $m_i^{k/k_{3,i}}$ and let

$$g_{2,i}(z_0, z_1) := g_{1,i}\left(\frac{k}{k_{3,i}}z_0, z_1^{k/k_{3,i}}\right) \in \mathfrak{o}_K[[z_0, z_1]].$$

With this new notation, it follows from Steps (i)–(iv) that for any integer $n \geq 0$,

- (1) $f_i^{\ell_{1,i}+nk}(\alpha_i) = u_i(nk + \mu_i)$, if $f_i^{\ell_{1,i}}(\alpha_i)$ lies in a quasiperiodic residue class;
- (2) $f_i^{\ell_{1,i}+nk}(\alpha_i) = u_i(\lambda_i^n \mu_i) = u_i((\pi^n)^{e_i} g_{2,i}(n) \mu_i)$, if $f_i^{\ell_{1,i}}(\alpha_i)$ lies in an attracting residue class; and
- (3) $f_i^{\ell_{1,i}+nk}(\alpha_i) = u_i(\mu_i^{m_i^n}) = u_i((\pi^{m_i^n})^{e_i} g_{2,i}(n, \pi^n))$, if $f_i^{\ell_{1,i}}(\alpha_i)$ lies in a superattracting residue class,

where $\mu_i = u_i^{-1}(f_i^{\ell_{1,i}}(\alpha_i))$ is as in Step (ii). In particular, in all three cases, we have expressed $f_i^{\ell_{1,i}+nk}(\alpha_i)$ as a power series in n , π^n , and, if needed, $\pi^{m_i^n}$.

Let $L = \max\{\ell_{1,1}, \ell_{1,2}\}$. For each $\ell = L, \dots, L+k-1$ and each $i = 1, 2$, choose a linear fractional transformation $\eta_{i,\ell} \in \mathrm{PGL}(2, \mathfrak{o}_K)$ so that $\eta_{i,\ell} \circ f_i^\ell(\alpha_i) \in D(0, 1)$. Then $\eta_{i,\ell} \circ f_i^{\ell-\ell_{1,i}}(D(0, 1)) \subseteq D(0, 1)$, because f_i has good reduction. Finally, define $E_{i,\ell} = \eta_{i,\ell} \circ f_i^{\ell-\ell_{1,i}} \circ u_i$, so that $E_{i,\ell} \in K[[z]]$ maps $\overline{D}(0, s_i)$ into $D(0, 1)$.

Step (vi). In this step, we will write down power series $F_{i,\ell}$ for $f_i^{\ell+nk}$ in terms of n , π^n , and $\pi^{m_i^n}$. We will also produce bounds $B_{i,\ell}$ to be used in applying [BGKT10, Lemma 3.1]. For each $i = 1, 2$, we consider the three cases that $f_i^\ell(\alpha_i)$ lies in a quasiperiodic, attracting (but not superattracting), or superattracting residue class for the function f_i^k .

In the quasiperiodic case, for each $\ell = L, \dots, L+k-1$, define the power series

$$F_{i,\ell}(z_0) = E_{i,\ell}(kz_0 + \mu_i) \in K[[z_0]],$$

so that $F_{i,\ell}(n) = \eta_{i,\ell} \circ f_i^{\ell+nk}(\alpha_i)$ for all $n \geq 0$. All coefficients of $F_{i,\ell}$ have absolute value at most 1, because $|k|_p, |\mu_i|_p \leq s_i$ and $E_{i,\ell}$ maps $\overline{D}(0, s_i)$ into $D(0, 1)$. Hence, we set our bound $B_{i,\ell}$ to be $B_{i,\ell} := 0$.

Second, in the attracting (but not superattracting) case, for each $\ell = L, \dots, L+k-1$, define the power series

$$F_{i,\ell}(z_0, z_1) = E_{i,\ell}(z_1^{e_i} g_{2,i}(z_0) \mu_i) \in K[[z_0, z_1]],$$

where $E_{i,\ell}$ and $g_{2,i}$ are as in Step (v), so that $F_{i,\ell}(n, \pi^n) = \eta_{i,\ell} \circ f_i^{\ell+nk}(\alpha_i)$ for all $n \geq 0$.

Still in the attracting (but not superattracting) case, because $E_{i,\ell}$ maps $\overline{D}(0, s_i)$ into $D(0, 1)$, there is some $B_{i,\ell} > 0$ such that for every $j \geq 0$, the coefficient of z^j in $E_{i,\ell}(z)$ has absolute value at most $p^{jB_{i,\ell}}$. Recalling also that $g_{2,i} \in \mathfrak{o}_K[[z]]$

and $|\mu_i|_p < 1$, it follows that if we write $F_{i,\ell}(z_0, z_1) = \sum_{j=0}^{\infty} h_j(z_0) z_1^j$ (where each $h_j \in K[[z]]$), then for each $j \geq 0$, all coefficients of h_j have absolute value at most $p^{jB_{i,\ell}}$.

Third, in the superattracting case, for each $\ell = L, \dots, L+k-1$, define the power series

$$F_{i,\ell}(z_0, z_1, z_{m_i}) = E_{i,\ell}(g_{2,i}(z_0, z_1) z_{m_i}^{e_i}) \in K[[z_0, z_1, z_{m_i}]],$$

where $E_{i,\ell}$ and $g_{2,i}$ are as in Step (v), so that $F_{i,\ell}(n, \pi^n, \pi^{m_i^n}) = \eta_{i,\ell} \circ f_i^{\ell+nk}(\alpha_i)$ for all $n \geq 0$.

Still in the superattracting case, because $E_{i,\ell}$ maps $\overline{D}(0, s_i)$ into $D(0, 1)$, there is some $B_{i,\ell} > 0$ such that for every $j \geq 0$, the coefficient of z^j in $E_{i,\ell}(z)$ has absolute value at most $p^{jB_{i,\ell}}$. Hence, if we write $F_{i,\ell}(z_0, z_1, z_{m_i}) = \sum_{j_1, j_2 \geq 0} h_{j_1, j_2}(z_0) z_1^{j_1} z_{m_i}^{j_2}$ (where each $h_{j_1, j_2} \in K[[z]]$), then as before, since $g_{2,i} \in \mathfrak{o}_K[[z_0, z_1]]$, all coefficients of h_{j_1, j_2} have absolute value at most $p^{j_2 B_{i,\ell}} \leq p^{B_{i,\ell}(j_1 + j_2)}$. Finally, set

$$B := \max_{L \leq \ell \leq L+k-1} (B_{1,\ell} + B_{2,\ell}).$$

Step (vii). For each $\ell = L, \dots, L+k-1$, let $\mathcal{H}_\ell \in \mathfrak{o}_K[t_1, t_2]$ be the polynomial generating the vanishing ideal of the curve $Y \subset (\mathbb{P}^1)^2$ dehomogenized with respect to the coordinates determined by $(\eta_{1,\ell}, \eta_{2,\ell})$. Then we define

$$(3.1.7) \quad G_\ell(z_0, z_1, z_{m_1}, z_{m_2}) = \mathcal{H}_\ell(F_{1,\ell}, F_{2,\ell}) \in K[[z_0, z_1, z_{m_1}, z_{m_2}]].$$

If there is no superattracting point for f_i lying in the orbit of α_i modulo π , then $m_i = 1$, i.e., the above power series G_ℓ has one less variable. Also, if it happens that $m_1 = m_2$, then again we need one variable less in (3.1.7). From now on, we assume both α_1 and α_2 land in superattracting cycles modulo π (i.e., $m_1, m_2 > 1$) and also, assume $m_1 \neq m_2$; this would be the most general case in (3.1.7) since we would deal with a p -adic analytic function of 4 variables (the other cases are similar, only simpler). So, without loss of generality, we may assume $m_2 > m_1 > 1$.

In all cases, by construction, $G_\ell(n, \pi^n, \pi^{m_1^n}, \pi^{m_2^n})$ is defined for all integers $n \geq 0$, and moreover,

$$(3.1.8) \quad G_\ell(n, \pi^n, \pi^{m_1^n}, \pi^{m_2^n}) = 0 \text{ if and only if } \Phi^{\ell+nk}(\alpha) \in Y.$$

Now, if $G_\ell = 0$ for some $\ell = L, \dots, L+k-1$, then we get that $\Phi^{\ell+nk}(\alpha) \in Y$ for all $n \geq 0$ and since α is not preperiodic, while Y is a curve, we conclude that Y must be fixed under the action of Φ^k , as desired (note that k is bounded solely in terms of the degrees of the f_i 's and also on p, e and $[\kappa : \mathbb{F}_p]$).

So, from now on, we assume that $G_\ell \neq 0$ for each $\ell = L, \dots, L+k-1$. Then for each G_ℓ , we write

$$G_\ell(z_0, \pi^n, \pi^{m_1^n}, \pi^{m_2^n}) = \sum_{w \in \mathbb{N}^3} g_w(z_0) \pi^{f_w(n)},$$

where for each $w := (w_1, w_2, w_3) \in \mathbb{N}^3$, we have that $f_w(n) := w_1 n + w_2 m_1^n + w_3 m_2^n$. Then we let v be the smallest element of \mathbb{N}^3 with respect to the usual lexicographic order on \mathbb{N}^3 such that $g_v \neq 0$. Also, for each element $w := (w_1, w_2, w_3) \in \mathbb{N}^3$, we define $|w| := w_1 + w_2 + w_3$; for more details, see [BGKT10, Section 3].

By our choice of the bound B in Step (vi), and because all coefficients of \mathcal{H}_ℓ lie in \mathfrak{o}_K , all coefficients of g_w have absolute value at most $p^{B|w|}$, for every $w \in \mathbb{N}^3$. Since $G_\ell(n, \pi^n, \pi^{m_1^n}, \pi^{m_2^n})$ is defined at every $n \geq 0$, then g_v must converge on $\overline{D}(0, 1)$; therefore, we may choose a radius $0 < s_\ell \leq 1$ for g_v satisfying the hypotheses of

[BGKT10, Lemma 3.1]. More precisely, g_v has only finitely many zeros in $\overline{D}(0, 1)$ and so, we let $0 < s_\ell \leq 1$ be the minimum distance between any two distinct such zeros.

Let s be the minimum of all the s_ℓ as we vary $\ell \in \{L, \dots, L + k - 1\}$. The set \mathfrak{o}_K may be covered by finitely many disks $D(\gamma, s)$ (for some points $\gamma \in \mathfrak{o}_K$). We let $N := k \cdot p^M$ for some sufficiently large integer M such that $|p^M|_p < s$.

Since each G_ℓ is nontrivial, we may apply [BGKT10, Lemma 3.1] (with the bound B from Step (vi) and radius s from the previous paragraph) and conclude that for each $j \in \{1, \dots, N\}$, the set

$$(3.1.9) \quad \{n \in \mathbb{N}: \Phi^{j+nN}(\alpha) \in Y\} \text{ has natural density } 0.$$

In order to derive (3.1.9), we use (3.1.8) and [BGKT10, Lemma 3.1 and Corollary 1.5]. However, since Y is periodic, the set

$$\{n \in \mathbb{N}: \Phi^n(\alpha) \in Y\} \text{ has positive natural density,}$$

which contradicts (3.1.9). In conclusion, it must be that some G_ℓ is trivial and therefore Y must be fixed by Φ^k .

This concludes the proof of Theorem 3.1. \square

4. PROOF OF OUR MAIN RESULTS

We first prove Theorem 1.2. Our strategy is to prove separately its conclusion assuming that either

- (1) all rational functions f_i are *disintegrated*, using the terminology from [MS14] (or *non-special*, using the terminology from [GN16, GN17]), i.e., for each $i = 1, \dots, n$, we have that $\deg(f_i) \geq 2$ and also, $f_i(x)$ is not conjugated (through a linear transformation) to $x^{\pm \deg(f_i)}$, or to $\pm C_{\deg(f_i)}$ (where, for each $m \geq 2$, $\pm C_m(x)$ is the m -th Chebyshev polynomial, which satisfies $C_m(x + \frac{1}{x}) = x^m + \frac{1}{x^m}$), or to a Lattés map (i.e., the quotient of an endomorphism of an elliptic curve); or
- (2) all rational functions f_i are special, i.e., for each $i = 1, \dots, n$, we have that $\deg(f_i) \geq 2$ and also, $f_i(x)$ is conjugated either to $x^{\pm \deg(f_i)}$, or to $\pm C_{\deg(f_i)}$, or to a Lattés map; or
- (3) each rational function f_i has degree equal to 1.

In each of the above cases (1)-(3), we show that the length of the period of a periodic subvariety (satisfying the hypotheses of Theorem 1.2) is uniformly bounded. Case (1) follows from combining our Theorem 3.1 along with the description from [MS14] of the periodic subvarieties of $(\mathbb{P}^1)^n$ under the coordinatewise action of n disintegrated rational functions (for more details, see Theorem 4.1). Case (2) above follows using classical results regarding algebraic subgroups of product of tori and of elliptic curves (for more details, see Theorem 4.3). Finally, case (3) follows using a direct analysis since for linear maps, one can find explicit formulas for the points in any of their orbits (for more details, see Theorem 4.2). Then we finish the proof of Theorem 1.2 by employing one more time the results of [MS14] which allows us to prove the conclusion of Theorem 1.2 after splitting the action of the f_i 's in the above 3 cases and using Theorems 4.1, 4.2 and 4.3.

Theorem 4.1. *Theorem 1.2 holds under the additional assumption that each rational function f_i has degree greater than 1, and moreover, no $f_i(x)$ is conjugated to either $x^{\pm \deg(f_i)}$, or to $\pm C_{\deg(f_i)}$, or to a Lattés map.*

Proof. Since each f_i is disintegrated then, according to [MS14, Proposition 2.21], we know that each (irreducible) periodic subvariety $Y \subset (\mathbb{P}^1)^n$ is an irreducible component of the intersection of finitely many hypersurfaces $Y_{i,j}$ of $(\mathbb{P}^1)^n$, which are of the form $Y_{i,j} := \pi_{i,j}^{-1}(C_{i,j})$, where $C_{i,j}$ is an irreducible periodic curve defined over K under the action of (f_i, f_j) on $\mathbb{P}^1 \times \mathbb{P}^1$ and $\pi_{i,j} : (\mathbb{P}^1)^n \rightarrow \mathbb{P}^1 \times \mathbb{P}^1$ is the projection on the (i, j) -th coordinate axes; note that $C_{i,j}$ is also defined over K since the projection of the irreducible K -subscheme Y of $(\mathbb{P}^1)^n$ on any two coordinates would still be an irreducible K -subscheme (of $(\mathbb{P}^1)^2$). By Theorem 3.1, we know that the length of the orbit of $C_{i,j}$ under the induced action of (f_i, f_j) on $\mathbb{P}^1 \times \mathbb{P}^1$ is bounded independently of $C_{i,j}$ (and only depending on the degrees of f_i and f_j and also on p , e and $[\kappa : \mathbb{F}_p]$); note that the point $(\alpha_i, \alpha_j) \in C_{i,j}(K)$ satisfies the hypotheses from Theorem 3.1. Therefore, each hypersurface $Y_{i,j}$ is periodic and moreover, the length of its period under the coordinatewise action on $(\mathbb{P}^1)^n$ of the rational functions f_i is bounded solely in terms of the degrees of the rational functions f_i and also in terms of p , e and $[\kappa : \mathbb{F}_p]$. Since Y is an irreducible component of $\bigcap_{i,j} Y_{i,j}$, this concludes the proof of Theorem 4.1. \square

Theorem 4.2. *Theorem 1.2 holds under the assumption that each rational function f_i has degree equal to 1.*

Proof. We use Poonen's [Poo13] result on p -adic interpolation of iterates. Since each f_i is a degree one map with good reduction at π , there is an M depending only on p , e , and $[\kappa : \mathbb{F}_p]$ such that $f_i^M(x) \equiv x \pmod{p^c}$, where c is the smallest integer greater than $1/(p-1)$. Now, let $\gamma = (\gamma_1, \dots, \gamma_n)$ be any point in $Y(\overline{K})$, and let R be the ring of p -adic integers in $K(\gamma)$. Then taking the local power series expansion for f_i^M at γ_i gives a power series $g_i \in R[[x]]$, convergent on R , such that $g_i(z) = f_i^M(z)$ for all $z \in R$. Since $f_i^M(x) \equiv x \pmod{p^c}$, we have $g_i(x) \equiv x \pmod{p^c}$.

By [Poo13], for each i , there is therefore a power series $\theta_i \in R[[x]]$, convergent on R , such that $\theta_i(k) = (f_i^M)^k(\gamma_i)$ for all positive integers k . Now, let H be any polynomial in the vanishing ideal of Y , dehomogenized at γ . Since there are infinitely many k such that $\Phi^{kM}(\gamma) \in Y$ (because Y is periodic under Φ), there are infinitely many k such that $H(\theta_1(k), \dots, \theta_n(k)) = 0$. Since a convergent p -adic power series has finitely many zeros unless the series is identically zero, it follows that $H(\theta_1(k), \dots, \theta_n(k)) = 0$ for all k , so $\Phi^{Mk}(\gamma) \in Y$ for all k . Since this is true for all $\gamma \in Y(\overline{K})$, we must have that Y has period dividing M .

As an aside, we could argue also directly, using the fact that each f_i is a linear map to find explicit formulas for $f_i^n(\alpha_i)$ and thus construct explicit p -adic analytic functions as above, thus proving the desired conclusion in Theorem 4.2. \square

Theorem 4.3. *Theorem 1.2 holds under the assumption that each rational function f_i has degree larger than 1 and is conjugate either to $x^{\pm \deg(f_i)}$, or to $\pm C_{\deg(f_i)}$, or to a Lattés map.*

Proof. Arguing as in the proof of Theorem 4.2, at the expense of replacing each f_i by a conjugate (which does not change the conclusion of our result), we may assume each $f_i(x)$ is either a Lattés map, or equal to $\pm C_{\deg(f_i)}$, or equal to $x^{\pm \deg(f_i)}$.

In this case there exist elliptic curves E_1, \dots, E_k (for some $0 \leq k \leq n$) and there exists an endomorphism Ψ of $S := \mathbb{G}_m^{n-k} \times \prod_{i=1}^k E_i$ along with a finite morphism

$\eta : S \longrightarrow (\mathbb{P}^1)^n$ such that

$$(4.3.1) \quad \eta \circ \Psi = \Phi \circ \eta.$$

Indeed, for each Lattés map f_i (without loss of generality, we assume f_{n-k+1}, \dots, f_n are all the Lattés maps), we know there exists an elliptic curve E_i along with some endomorphism g_i and also there exists some morphism $\eta_i : E_i \longrightarrow \mathbb{P}^1$ of degree 2 (identifying each point P of E_i with $-P$) such that $\eta_i \circ g_i = f_i \circ \eta_i$ for each $i = n - k + 1, \dots, n$. Furthermore, assuming f_1, \dots, f_ℓ (for some $0 \leq \ell \leq n - k$) are all the rational functions conjugated to $\pm C_{\deg(f_i)}$, then for each $i = 1, \dots, \ell$, we let $\eta_i : \mathbb{G}_m \longrightarrow \mathbb{P}^1$ be defined by $\eta_i(x) = x + \frac{1}{x}$; also, we let $g_i(x) = \pm x^{\deg(f_i)}$ for each such i . Finally, for each $i = \ell + 1, \dots, n - k$, we let $g_i = f_i$ (which is thus equal to $x^{\pm \deg(f_i)}$) and also let $\eta_i(x) = x$ for each such $i = \ell + 1, \dots, n - k$. Then (4.3.1) holds with $\eta := (\eta_1, \dots, \eta_n)$ and also with $\Psi := (g_1, \dots, g_n) : S \longrightarrow S$ (acting coordinatewise).

Now, $Y \subset (\mathbb{P}^1)^n$ is periodic under the action of Φ if and only if an irreducible component of $Z := \eta^{-1}(Y)$ is periodic under the action of Ψ ; moreover, the two periodic varieties would then have the same length for their corresponding orbits. Hence, it suffices to bound the length of the period for any periodic subvariety of S defined over some given local field. Note that if Y is defined over K , then Z is defined over another local field L whose degree over K is bounded by 2^n since for each $i = 1, \dots, \ell$ and also for each $i = n - k + 1, \dots, n$, the degree of η_i is 2 (while $\deg(\eta_i) = 1$ if $i = \ell + 1, \dots, n - k$). Therefore, replacing K by L simply increases the size of the residue field of the new local field and also increases its ramification index (over \mathbb{Q}_p), but both the size of the residue field of K and the ramification index of K increase by a scalar factor which depends only on n .

Furthermore, at the expense of conjugating each $g_i(x)$ (for $1 \leq i \leq \ell$) by a suitable linear transformation $x \mapsto \zeta_i x$, we may actually assume each $g_i(x) = x^{\deg(f_i)}$ (i.e., modulo a conjugation by a linear scaling map given by a root of unity, we may assume the sign in front of each monomial from the definition of $g_i(x)$ is positive). In order to replace the original maps by the new maps, we might need to replace K by a finite extension, which depends only on the degrees of the original rational functions f_i .

We observe that S is a split semiabelian variety (whose abelian part is a product of elliptic curves). Thus the periodic subvarieties of S (under the coordinatewise action of group homomorphisms on each of its 1-dimensional factors) are torsion translates of algebraic subgroups (a more general result holds inside any semiabelian variety, as proven in [Hin88, Lemme 10]). Moreover, noting that there are no nontrivial morphisms between a torus and an elliptic curve, then each periodic subvariety Z of S is the zero locus of finitely many equations of the form

$$(4.3.2) \quad \prod_{i=1}^{n-k} x_i^{c_i} = \zeta$$

for some root of unity $\zeta \in K$ and some integers c_i , or

$$(4.3.3) \quad \sum_{i=1}^k \psi_i(y_i) = Q,$$

where E is one of the elliptic curves E_j (for some $j = 1, \dots, k$) and $\psi_i : E_i \rightarrow E$ are group endomorphisms, while Q is a torsion point of $E(K)$. Clearly, if E_i and E are not isogenous, then ψ_i is the trivial map.

Furthermore, the algebraic subgroup of \mathbb{G}_m^{n-k} defined by the equations

$$(4.3.4) \quad \prod_{i=1}^{n-k} x_i^{c_i} = 1$$

(obtained by replacing each ζ by 1 in (4.3.2)) must be periodic under the coordinatewise action of the g_i 's. Then we note that for each equation (4.3.4), letting I be the set consisting of all indices i such that $c_i \neq 0$, we have that $\deg(g_i)$ is the same for each $i \in I$, or otherwise the hypersurface given by the equation (4.3.4) is not periodic under the coordinatewise action of the g_i 's for $i \in I$. So, each hypersurface of the form (4.3.4) is invariant under the coordinatewise action of g_1^2, \dots, g_{n-k}^2 on \mathbb{G}_m^{n-k} since each g_i is a monomial $x^{\pm \deg(g_i)}$ and therefore $g_i^2(x) = x^{\deg(f_i)^2}$; in particular, the algebraic subgroup which is the zero locus of all equations (4.3.4) is fixed by the coordinatewise action of the g_i^2 's.

Similarly, the algebraic subgroup of $E_1 \times \dots \times E_k$ given by the equations

$$(4.3.5) \quad \sum_{i=1}^k \psi_i(y_i) = 0,$$

(obtained by replacing each Q by 0 in (4.3.3)) must be periodic under the coordinatewise action of g_{n-k+1}, \dots, g_n . We claim that the length of its period is bounded by the number of roots of unity contained in the endomorphism ring of E . Indeed, whenever ψ_i is nontrivial, then E_i and E have isomorphic endomorphism rings and moreover, the endomorphism g_i of E_i descends to an endomorphism of E . Note that if ψ_i and ψ_j are nontrivial and the endomorphisms g_{n-k+i} and g_{n-k+j} of E_i , respectively of E_j , correspond to elements ω_i and ω_j in the endomorphism ring R of E whose quotient ω_i/ω_j is not a root of unity, then the algebraic subgroup given by equation (4.3.5) is not periodic under the action of (g_{n-k+1}, \dots, g_n) . Therefore, the length of the period of the algebraic group given by equation (4.3.5) is absolutely bounded because there are at most 6 roots of unity in an order of an imaginary quadratic number ring (such as the ring R).

Using the fact that Z is the zero locus of finitely many equations of the form (4.3.2) and (4.3.3), while the algebraic groups given by equations (4.3.4) and (4.3.5) have the size of their orbit under Ψ bounded by 6, we obtain that the length of the orbit of Z under Ψ is bounded in terms of the orders of the roots of unity ζ appearing in equations of the form (4.3.2) and also in terms of the orders of the torsion points Q appearing in equations of the form (4.3.3). So, we conclude that the length of the orbit under the action of Ψ of any periodic subvariety of S (defined over the local field K) is bounded solely in terms of the size of the group of roots of unity contained in K and also in terms of the size of the torsion subgroups of $E_i(K)$, for $i = 1, \dots, k$. Since any local field contains finitely many roots of unity (the bound depending solely on the size of its residue field and also depending on its ramification index over \mathbb{Q}_p), and also, any elliptic curve has finitely many torsion points over the local field K , with an upper bound for the size of its torsion depending solely on the size of the residue field of K and on the ramification index for K/\mathbb{Q}_p (see [Sil86, Chapter VII]), then we obtain the desired conclusion in Theorem 4.3. \square

Finally, we can prove Theorem 1.2

Proof of Theorem 1.2. As proven in [Med07] (see also [MS14, Theorem 2.30] for the case when each f_i is a polynomial), each periodic subvariety Y of the dynamical system $((\mathbb{P}^1)^n, \Phi)$ may be written (after a suitable re-ordering of the coordinate axes, and thus of the rational functions acting on them) as $Y_1 \times Y_2 \times Y_3$, where $Y_i \subset (\mathbb{P}^1)^{n_i}$ for some integers n_i satisfying $n_1 + n_2 + n_3 = n$ and moreover, the following holds:

- the rational functions f_1, \dots, f_{n_1} are all disintegrated;
- each rational function $f_i(x)$ from the list $f_{n_1+1}, \dots, f_{n_1+n_2}$ is conjugate either to $x^{\pm \deg(f_i)}$, or to $\pm C_{\deg(f_i)}$, or to a Lattés function (and its degree is greater than 1);
- the rational functions $f_{n_1+n_2+1}, \dots, f_{n_1+n_2+n_3}$ have degree equal to 1;
- Y_1 is periodic under the action of $\Phi_1 := (f_1, \dots, f_{n_1})$ on $(\mathbb{P}^1)^{n_1}$, while Y_2 is periodic under the action of $\Phi_2 := (f_{n_1+1}, \dots, f_{n_1+n_2})$ and Y_3 is periodic under the action of $\Phi_3 := (f_{n_1+n_2+1}, \dots, f_{n_1+n_2+n_3})$.

Therefore, the uniform bound on the length of the orbit of Y under the action of Φ follows using Theorems 4.1, 4.2 and 4.3, which provide uniform bounds for the length of the orbits of each of the periodic subvarieties Y_j (for $1 \leq j \leq 3$). \square

We conclude this section by proving Theorem 1.1.

Proof of Theorem 1.1. We argue by induction on n ; the case $n = 1$ follows from [Zie96]. So, we assume now that $n \geq 2$ and that Theorem 1.1 holds for endomorphisms of $(\mathbb{P}^1)^{n-1}$. In particular, we may assume that Y projects dominantly onto each coordinate axes, since otherwise $Y = Y_0 \times \{\gamma\}$, where $\gamma \in \mathbb{P}_K^1$ is a periodic point (of bounded period, according to [Zie96]) and then the inductive hypothesis yields the desired conclusion.

Let \mathcal{Y} be a model for Y over \mathfrak{o}_K such that \mathcal{Y} is closed in $(\mathbb{P}_{\mathfrak{o}_K}^1)^n$. Then \mathcal{Y} is a projective \mathfrak{o}_K -scheme, so every point in $Y(K)$ extends to a unique point in $\mathcal{Y}(\mathfrak{o}_K)$. We let α denote the point in $\mathcal{Y}(\mathfrak{o}_K)$ to which $(\alpha_1, \dots, \alpha_n)$ extends.

Let $\bar{\alpha}$ be the reduction of α modulo p , i.e., the intersection of α with the special fiber of \mathcal{Y} . Since the intersection of α with the generic fiber Y of \mathcal{Y} is a smooth point x , then [BGT15, Proposition 2.2] yields the existence of a Zariski dense, uncountable set $U \subset \mathcal{Y}(\mathfrak{o}_v)$ consisting of sections β whose intersection with the special fiber of \mathcal{Y} equals $\bar{\alpha}$. The fact that U is uncountable follows from the p -adic implicit function theorem used in the proof of [BGT15, Proposition 2.2], which yields the existence of a p -adic submanifold of \mathcal{Y} whose points all have the same reduction modulo p . More precisely, as shown in [BGT15, Proposition 2.2], letting $d := \dim(Y)$, at the expense of relabelling the coordinate axes, we know that for a sufficiently small p -adic neighborhood \mathcal{U} of $(\alpha_{n-d+1}, \dots, \alpha_n)$ (where $\alpha := (\alpha_1, \dots, \alpha_n)$), we have p -adic analytic functions $\mathcal{F}_1, \dots, \mathcal{F}_n$ on \mathcal{U} such that for each $\gamma \in \mathcal{U}$, the point $(\mathcal{F}_1(\gamma), \dots, \mathcal{F}_n(\gamma))$ is on \mathcal{Y} .

Each one of the points $\beta \in U$ is of the form $(\beta_1, \dots, \beta_n)$ and if for some point β , we have that each β_i is not preperiodic for the action of f_i , then the result follows from Theorem 1.2. So, assume for each $\beta \in U$, there exists some $i \in \{1, \dots, n\}$ such that β_i is preperiodic for f_i . Therefore there must exist some $i_0 \in \{1, \dots, n\}$ such that there exists an uncountable set $U_0 \subset \mathcal{Y}(\mathfrak{o}_v)$ consisting of points β with the property that the corresponding β_{i_0} is preperiodic for the action

of f_{i_0} . Furthermore, we may assume that there exists no submanifold \mathcal{U}_0 of \mathcal{U} of dimension less than d such that for each $\beta \in U_0$, there exists some $\gamma \in \mathcal{U}_0$ such that $\beta = (\mathcal{F}_1(\gamma), \dots, \mathcal{F}_n(\gamma))$.

For the sake of simplifying the notation, we identify each β_{i_0} with its intersection with the generic fiber of $\mathbb{P}_{\mathfrak{o}_v}^1$ and therefore, view each β_{i_0} as a point in $\mathbb{P}^1(K)$ whose reduction modulo p is $\overline{\alpha_{i_0}}$. There are two cases.

Case 1. The set $\{\beta_{i_0} : \beta \in U_0\}$ is uncountable.

Then, according to our assumption, we obtain that f_{i_0} has an uncountable set of preperiodic points, which automatically yields that f_{i_0} itself must be preperiodic; more precisely, $\deg f_{i_0} = 1$ (i.e. f_{i_0} is an automorphism) and $f_{i_0}^\ell$ is the identity, for some positive integer ℓ . So, at the expense of replacing \mathcal{Y} by \mathcal{Y}^ℓ , we may assume \mathcal{Y} is fixed under the action of f_{i_0} on the i_0 -th coordinate axis. Therefore, the length of the period of \mathcal{Y} equals the length of the period of \mathcal{Y}_0 under the coordinatewise action of the rational functions f_i for $i \neq i_0$, where \mathcal{Y}_0 is the projection of \mathcal{Y} on the remaining $(n-1)$ coordinate axes, other than the i_0 -th coordinate axis. In this case, we are done by the inductive hypothesis; also, note that ℓ depends solely on p , e and $[\kappa : \mathbb{F}_p]$. Indeed, the linear map $f_{i_0}(x)$ is conjugated over a quadratic extension L of K to one of these two maps $x \mapsto x+1$ or $x \mapsto ax$; furthermore, since f_{i_0} has finite order, then $f_{i_0}(x)$ must be conjugated to a map $x \mapsto ax$, where a is a root of unity contained in L . Because the number of roots of unity in a local field L is bounded solely in terms of p and of the degree $[L : \mathbb{Q}_p]$, our claim follows.

Case 2. The set $\{\beta_{i_0} : \beta \in U_0\}$ is countable.

Let $\mathcal{U}_0 \subset \mathcal{U}$ be the set consisting of all points $\gamma \in \mathcal{U}$ such that $(\mathcal{F}_1(\gamma), \dots, \mathcal{F}_n(\gamma)) \in U_0$. Then we know that the set $\{\mathcal{F}_{i_0}(\gamma) : \gamma \in \mathcal{U}_0\}$ is countable; furthermore, by our assumption, we know that \mathcal{U}_0 is an uncountable set, which is not contained in a submanifold of \mathcal{U} of dimension less than d . Therefore, \mathcal{F}_{i_0} must be constant, and thus the projection of \mathcal{Y} on the i_0 -th coordinate axis must be constant, contrary to our hypothesis from the beginning of our proof. This concludes the proof of Theorem 1.1. \square

REFERENCES

- [BGT15] J. P. Bell, D. Ghioca, and T. J. Tucker, *Applications of p -adic analysis for bounding periods for subvarieties under étale maps*, Int. Math. Res. Not. **IMRN** **2015**, no. 11, 3576-3597.
- [BGT16] J. P. Bell, D. Ghioca, and T. J. Tucker, *The dynamical Mordell-Lang conjecture*, Mathematical Surveys and Monographs **210**, American Mathematical Society, Providence, RI, 2016, xiii+280 pp.
- [BGKT10] R. L. Benedetto, D. Ghioca, P. Kurlberg, and T. J. Tucker, *A gap principle for dynamics*, Compos. Math. **146** (2010), no. 4, 1056-1072.
- [GN16] D. Ghioca and K. D. Nguyen, *Dynamical anomalous subvarieties: structure and bounded height theorems*, Adv. Math. **288** (2016), 1433-1462.
- [GN17] D. Ghioca and K. D. Nguyen, *Dynamics of split polynomial maps: uniform bounds for periods and applications*, Int. Math. Res. Not. **IMRN** **2017**, no. 1, 213-231.
- [Hin88] M. Hindry, *Autour d'une conjecture de Serge Lang*, Invent. Math. **94** (1988), no. 3, 575-603.
- [Hut09] B. Hutz, *Good reduction of periodic points on projective varieties*, Illinois J. Math. **53** (2009), no. 4, 1109-1126.
- [Hut] B. Hutz, *Good reduction and canonical heights of subvarieties*, arxiv:1603.00860.
- [Med07] A. Medvedev, *Minimal sets in ACFA*, Thesis (Ph.D.), University of California, Berkeley, 2007, 96 pp.
- [MS14] A. Medvedev and T. Scanlon, *Invariant varieties for polynomial dynamical systems*, Ann. of Math. (2) **179** (2014), no. 1, 81-177.

- [MS94] P. Morton and J. H. Silverman, *Rational periodic points of rational functions*, Int. Math. Res. Not. IMRN **1994**, no. 2, 97-110.
- [Pez05] T. Pezda, *Cycles of polynomial mappings in several variables over rings of integers in finite extensions of the rationals. II*, Monatsh. Math. **145** (2005), no. 4, 321-331.
- [Poo13] B. Poonen, *p-adic interpolation of iterates*, Bull. Lond. Math. Soc. **46** (2014), no. 3, 525-527.
- [RL03] J. Rivera-Letelier, *Dynamique des fonctions rationnelles sur des corps locaux*, Astérisque (2003), no. 287, 147-230, Geometric methods in dynamics. II.
- [Sch03] W. Schmidt, *Linear recurrence sequences*, Diophantine Approximation (Cetraro, Italy, 2000), Lecture Notes in Math. 1819, Springer-Verlag Berlin Heidelberg, 2003, pp. 171-247.
- [SS] A. Salerno and J. H. Silverman, *Integrality properties of Böttcher coordinates for one-dimensional superattracting germs*, arXiv:1708.09275.
- [Sil86] J. H. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics **106**, Springer-Verlag, New York, 1986, xii+400 pp.
- [Zie96] M. E. Zieve, *Cycles of polynomial mappings*, PhD thesis, University of California, Berkeley, 1996, 58 pp.

JASON BELL, DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF WATERLOO, WATERLOO, ON N2L 3G1, CANADA
E-mail address: `jpbell@uwaterloo.ca`

DRAGOS GHIOCA, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER, BC V6T 1Z2, CANADA
E-mail address: `dghioca@math.ubc.ca`

THOMAS TUCKER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY 14627, USA
E-mail address: `ttucker@math.rochester.edu`