

General Kastler-Kalau-Walze Type Theorems for Manifolds with Boundary II *

Yong Wang

School of Mathematics and Statistics, Northeast Normal University, Changchun Jilin 130024, China ;

E-mail: wangy581@nenu.edu.cn

Abstract In this paper, we establish some general Kastler-Kalau-Walze type theorems for any dimensional manifolds with boundary which generalize the results in [WW1].

MSC: 58G20; 53A30; 46L87

Keywords: Noncommutative residue for manifolds with boundary; Dirac operators

1 Introduction

The noncommutative residue found in [Gu],[Wo] plays a prominent role in noncommutative geometry. For one-dimensional manifolds, the noncommutative residue was discovered by Adler [Ad] in connection with geometric aspects of nonlinear partial differential equations. For arbitrary closed compact n -dimensional manifolds, the noncommutative residue was introduced by Wodzicki in [Wo] using the theory of zeta functions of elliptic pseudodifferential operators. In [Co1], Connes used the noncommutative residue to derive a conformal 4-dimensional Polyakov action analogue. Furthermore, Connes made a challenging observation that the noncommutative residue of the square of the inverse of the Dirac operator was proportional to the Einstein-Hilbert action in [Co2]. Let s be the scalar curvature and Wres denote the noncommutative residue. Then the Kastler-Kalau-Walze theorem gives an operator-theoretic explanation of the gravitational action and says that for a 4-dimensional closed spin manifold, there exists a constant c_0 , such that

$$\text{Wres}(D^{-2}) = c_0 \int_M s \text{dvol}_M. \quad (1.1)$$

In [Ka], Kastler gave a brute-force proof of this theorem. In [KW], Kalau and Walze proved this theorem in the normal coordinates system simultaneously. And then, Ackermann proved that the Wodzicki residue $\text{Wres}(D^{-2})$ in turn is essentially the second coefficient of the heat kernel expansion of D^2 in [Ac].

On the other hand, Fedosov etc. defined a noncommutative residue on Boutet de Monvel's algebra and proved that it was a unique continuous trace in [FGLS]. In [Sc],

*partially supported by NSFC 11771070.

Schrohe gave the relation between the Dixmier trace and the noncommutative residue for manifolds with boundary. We [Wa2] proved a Kastler-Kalau-Walze type theorem for 4-dimensional spin manifolds with boundary. Furthermore, We [Wa3] generalized the definition of lower dimensional volumes in [Po] to manifolds with boundary and found a Kastler-Kalau-Walze type theorem for 6-dimensional manifolds with boundary. In [WW2], we established a general Kastler-Kalau-Walze type theorem for any dimensional manifolds with boundary which generalized the results in [Wa3]. In [WW1], we computed the lower dimensional volume $\text{Vol}_6^{(1,3)}$ for 6-dimensional spin manifolds with boundary and got another Kastler-Kalau-Walze type theorem for 6-dimensional manifolds with boundary. **The motivation of this paper** is to establish a general Kastler-Kalau-Walze type theorem for any dimensional manifolds with boundary which extends the theorem in [WW1]. Our main result is as follows.

Theorem 1.1 *Let M be a $n = 2m + 2$ -dimensional compact spin manifold with the boundary ∂M , then*

$$\begin{aligned} \widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(-2m+1)}] &= \frac{(2-n)(2\pi)^{n/2}}{12\Gamma(n/2)} \int_M s \, \text{dvol}_M \\ &\quad + \frac{2}{1-n} L_0 \text{Vol}(S_{n-2}) \int_{\partial M} K \, \text{dvol}_{\partial M}, \end{aligned} \quad (1.2)$$

where $\text{Vol}(S_{n-2})$ is the canonical volume of the $n-2$ dimensional sphere S_{n-2} and s is the scalar curvature on M and K is the extrinsic curvature on ∂M as well as L_0 is a constant (see (3.42)).

This paper is organized as follows: In Section 2, we recall the definition of the lower dimensional volumes of compact Riemannian manifolds with boundary. In Section 3, we establish some general Kastler-Kalau-Walze type theorems for any dimensional manifolds with boundary which extend the theorem in [WW1].

2 Lower-Dimensional Volumes of Spin Manifolds with boundary

In this section we consider an $n = 2m + 2$ -dimensional oriented Riemannian manifold (M, g^M) with boundary ∂M equipped with a fixed spin structure. We assume that the metric g^M on M has the following form near the boundary

$$g^M = \frac{1}{h(x_n)} g^{\partial M} + \text{d}x_n^2, \quad (2.1)$$

where $g^{\partial M}$ is the metric on ∂M . Let $U \subset M$ be a collar neighborhood of ∂M which is diffeomorphic $\partial M \times [0, 1)$. By the definition of $h(x_n) \in C^\infty([0, 1))$ and $h(x_n) > 0$, there exists $\tilde{h} \in C^\infty((-\varepsilon, 1))$ such that $\tilde{h}|_{[0, 1)} = h$ and $\tilde{h} > 0$ for some sufficiently small $\varepsilon > 0$. Then there exists a metric \hat{g} on $\widehat{M} = M \cup_{\partial M} \partial M \times (-\varepsilon, 0]$ which has

the form on $U \cup_{\partial M} \partial M \times (-\varepsilon, 0]$

$$\hat{g} = \frac{1}{\tilde{h}(x_n)} g^{\partial M} + \mathbf{d}x_n^2, \quad (2.2)$$

such that $\hat{g}|_M = g$. We fix a metric \hat{g} on the \widehat{M} such that $\hat{g}|_M = g$. We can get the spin structure on \widehat{M} by extending the spin structure on M . Let D be the Dirac operator associated to \hat{g} on the spinors bundle $S(T\widehat{M})$. We want to compute

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(-2m+1)}]$$

(for the related definitions, see , Section 2, 3 [Wa1]).

To define the lower dimensional volume, some basic facts and formulae about Boutet de Monvel's calculus which can be found in Sec.2 in [Wa1] are needed. Let

$$F : L^2(\mathbf{R}_t) \rightarrow L^2(\mathbf{R}_v); F(u)(v) = \int e^{-ivt} u(t) dt$$

denote the Fourier transformation and $\Phi(\overline{\mathbf{R}^+}) = r^+ \Phi(\mathbf{R})$ (similarly define $\Phi(\overline{\mathbf{R}^-})$), where $\Phi(\mathbf{R})$ denotes the Schwartz space and

$$r^+ : C^\infty(\mathbf{R}) \rightarrow C^\infty(\overline{\mathbf{R}^+}); f \rightarrow f|_{\overline{\mathbf{R}^+}}; \overline{\mathbf{R}^+} = \{x \geq 0; x \in \mathbf{R}\}.$$

We define $H^+ = F(\Phi(\overline{\mathbf{R}^+}))$; $H_0^- = F(\Phi(\overline{\mathbf{R}^-}))$ which are orthogonal to each other. We have the following property: $h \in H^+$ (H_0^-) iff $h \in C^\infty(\mathbf{R})$ which has an analytic extension to the lower (upper) complex half-plane $\{\text{Im}\xi < 0\}$ ($\{\text{Im}\xi > 0\}$) such that for all nonnegative integer l ,

$$\frac{d^l h}{d\xi^l}(\xi) \sim \sum_{k=1}^{\infty} \frac{d^l}{d\xi^l} \left(\frac{c_k}{\xi^k} \right)$$

as $|\xi| \rightarrow +\infty, \text{Im}\xi \leq 0$ ($\text{Im}\xi \geq 0$).

Let H' be the space of all polynomials and $H^- = H_0^- \oplus H'$; $H = H^+ \oplus H^-$. Denote by π^+ (π^-) respectively the projection on H^+ (H^-). For calculations, we take $H = \tilde{H} = \{\text{rational functions having no poles on the real axis}\}$ (\tilde{H} is a dense set in the topology of H). Then on \tilde{H} ,

$$\pi^+ h(\xi_0) = \frac{1}{2\pi i} \lim_{u \rightarrow 0^-} \int_{\Gamma^+} \frac{h(\xi)}{\xi_0 + iu - \xi} d\xi, \quad (2.3)$$

where Γ^+ is a Jordan close curve included $\text{Im}(\xi) > 0$ surrounding all the singularities of h in the upper half-plane and $\xi_0 \in \mathbf{R}$. Similarly, define π' on \tilde{H} ,

$$\pi' h = \frac{1}{2\pi} \int_{\Gamma^+} h(\xi) d\xi. \quad (2.4)$$

So, $\pi'(H^-) = 0$. For $h \in H \cap L^1(\mathbf{R})$, $\pi' h = \frac{1}{2\pi} \int_{\mathbf{R}} h(v) dv$ and for $h \in H^+ \cap L^1(\mathbf{R})$, $\pi' h = 0$.

Let M be an n -dimensional compact oriented manifold with boundary ∂M . Denote by \mathcal{B} Boutet de Monvel's algebra, we recall the main theorem in [FGLS].

Theorem 2.1 (Fedosov-Golse-Leichtnam-Schrohe) *Let X and ∂X be connected, $\dim X = n \geq 3$, $A = \begin{pmatrix} \pi^+ P + G & K \\ T & S \end{pmatrix} \in \mathcal{B}$, and denote by p , b and s the local symbols of P , G and S respectively. Define:*

$$\begin{aligned} \widetilde{\text{Wres}}(A) &= \int_X \int_{\mathbf{S}} \text{tr}_E [p_{-n}(x, \xi)] \sigma(\xi) dx \\ &\quad + 2\pi \int_{\partial X} \int_{\mathbf{S}'} \{ \text{tr}_E [(\text{tr} b_{-n})(x', \xi')] + \text{tr}_F [s_{1-n}(x', \xi')] \} \sigma(\xi') dx', \end{aligned} \quad (2.5)$$

Then a) $\widetilde{\text{Wres}}([A, B]) = 0$, for any $A, B \in \mathcal{B}$; b) It is a unique continuous trace on $\mathcal{B}/\mathcal{B}^{-\infty}$.

Let p_1, p_2 be nonnegative integers and $p_1 + p_2 \leq n$. Then by Sec 2.1 of [Wa2], we have

Definition 2.2. Lower-dimensional volumes of spin manifolds with boundary are defined by

$$\text{Vol}_n^{(p_1, p_2)} M := \widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}]. \quad (2.6)$$

Denote by $\sigma_l(A)$ the l -order symbol of an operator A . Similar to (2.1.7) in [Wa2], we have that

$$\widetilde{\text{Wres}}[\pi^+ D^{-p_1} \circ \pi^+ D^{-p_2}] = \int_M \int_{|\xi|=1} \text{trace}_{S(TM)} [\sigma_{-n}(D^{-p_1-p_2})] \sigma(\xi) d\xi + \int_{\partial M} \Phi, \quad (2.7)$$

where

$$\begin{aligned} \Phi &= \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \text{trace}_{S(TM)} \left[\partial_{x_n}^j \partial_{\xi'}^\alpha \partial_{\xi_n}^k \sigma_r^+(D^{-p_1})(x', 0, \xi', \xi_n) \right. \\ &\quad \left. \times \partial_{x'}^\alpha \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l(D^{-p_2})(x', 0, \xi', \xi_n) \right] d\xi_n \sigma(\xi') d\xi', \end{aligned} \quad (2.8)$$

and the sum is taken over $r - k + |\alpha| + \ell - j - 1 = -n, r \leq -p_1, \ell \leq -p_2$. Since $[\sigma_{-n}(D^{-n+2})]_M$ has the same expression as $\sigma_{-n}(D^{-n+2})$ in the case of manifolds without boundary, by (2.5) in [Ac], we have

$$\text{Wres}(D^{-n+2}) = \frac{(2-n)(2\pi)^{n/2}}{12\Gamma(n/2)} \int_M \text{sdvol}_M. \quad (2.9)$$

So we only need to compute $\int_{\partial M} \Phi$.

3 Kastler-Kalau-Walze type theorems for any dimensional manifolds with boundary

In this section, we compute the lower dimensional volume for any dimensional compact manifolds with boundary and get a Kastler-Kalau-Walze type formula in this case. From now on we always assume that M carries a spin structure so that the spinor bundle and the Dirac operator are defined on M .

Set $\tilde{E}_n = \frac{\partial}{\partial x_n}$, $\tilde{E}_j = \sqrt{h(x_n)}E_j$ ($1 \leq j \leq n-1$), where $\{E_1, \dots, E_{n-1}\}$ are orthonormal basis of $T\partial_M$. Let ∇^L denote the Levi-civita connection about g^M . In the local coordinates $\{x_i; 1 \leq i \leq n\}$ and the fixed orthonormal frame $\{\tilde{E}_1, \dots, \tilde{E}_n\}$, the connection matrix $(\omega_{s,t})$ is defined by

$$\nabla^L(\tilde{E}_1, \dots, \tilde{E}_n)^t = (\omega_{s,t})(\tilde{E}_1, \dots, \tilde{E}_n)^t.$$

The Dirac operator is defined by

$$D = \sum_{j=1}^n c(\tilde{E}_j) \left[\tilde{E}_j + \frac{1}{4} \sum_{s,t} \omega_{s,t}(\tilde{E}_j) c(\tilde{E}_s) c(\tilde{E}_t) \right]. \quad (3.1)$$

Let $g^{ij} = g(dx_i, dx_j)$ and

$$\nabla_{\partial_i}^L \partial_j = \sum_k \Gamma_{ij}^k \partial_k; \quad \Gamma^k = g^{ij} \Gamma_{ij}^k,$$

Let the cotangent vector $\xi = \sum \xi_j dx_j$ and $\xi^j = g^{ij} \xi_i$. Let the symbols of D^2 and D be

$$\sigma(D^2) = \sigma_2(D^2) + \sigma_1(D^2) + \sigma_0(D^2); \quad \sigma(D^1) = \sigma_1(D^1) + \sigma_0(D^1). \quad (3.2)$$

By the composition formula of pseudodifferential operators, then we have

$$\sigma_{-2}(D^{-2}) = |\xi|^{-2}, \quad \sigma_{-(2m-2)}(D^{-(2m-2)}) = (|\xi|^2)^{1-m}, \quad \sigma_{-2m+1}(D^{-2m+1}) = \frac{\sqrt{-1}c(\xi)}{|\xi|^{2m}}. \quad (3.3)$$

By (3.8) in [WW2], we have:

$$\begin{aligned} \sigma_{-1-2m}(D^{-2m}) &= m\sigma_2(D^2)^{(-m+1)}\sigma_{-3}(D^{-2}) \\ &- \sqrt{-1} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m+2} \partial_{\xi_\mu} \sigma_2^{-m+k+1}(D^2) \partial_{x_\mu} \sigma_2^{-1}(D^2) (\sigma_2(D^2))^{-k}. \end{aligned} \quad (3.4)$$

By (3.4), we know that

$$\begin{aligned} \sigma_{-2m}(D^{1-2m}) &= \sigma_{-2m}(D^{-2m} \cdot D) = \left\{ \sum_{|\alpha|=0}^{+\infty} (-\sqrt{-1})^{|\alpha|} \frac{1}{\alpha!} \partial_\xi^\alpha [\sigma(D^{-2m})] \partial_x^\alpha [\sigma(D)] \right\}_{-2m} \\ &= \sigma_{-2m}(D^{-2m})\sigma_0(D) + \sigma_{-2m-1}(D^{-2m})\sigma_1(D) + \sum_{|\alpha|=1} (-\sqrt{-1}) \partial_\xi^\alpha [\sigma_{-2m}(D^{-2m})] \partial_x^\alpha [\sigma_1(D)] \end{aligned}$$

$$\begin{aligned}
&= |\xi|^{-2m} \sigma_0(D) + \sum_{j=1}^{2m+2} \partial_{\xi_j} (|\xi|^{-2m}) \partial_{x_j} c(\xi) + \left[m \sigma_2(D^2)^{(-m+1)} \sigma_{-3}(D^{-2}) \right. \\
&\quad \left. - \sqrt{-1} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m+2} \partial_{\xi_\mu} \sigma_2^{-m+k+1}(D^2) \partial_{x_\mu} \sigma_2^{-1}(D^2) (\sigma_2(D^2))^{-k} \right] \sqrt{-1} c(\xi). \quad (3.5)
\end{aligned}$$

Since Φ is a global form on ∂M , so for any fixed point $x_0 \in \partial M$, we can choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\tilde{U} = U \times [0, 1)$ and the metric $\frac{1}{h(x_n)} g^{\partial M} + dx_n^2$. The dual metric of g^M on \tilde{U} is $h(x_n) g^{\partial M} + dx_n^2$. Now we can compute Φ (see formula (2.8) for the definition of Φ). Since the sum is taken over $r + \ell - k - j - |\alpha| - 1 = -n$, $r \leq -1, \ell \leq 1 - 2m$, then we have the $\int_{\partial M} \Phi$ is the sum of the following five cases:

case a) I) $r = -1, \ell = 1 - 2m, k = j = 0, |\alpha| = 1$

$$\begin{aligned}
\text{case a) I)} &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{trace} \left[\partial_{\xi'}^\alpha \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \right. \\
&\quad \left. \times \partial_{x'}^\alpha \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \quad (3.6)
\end{aligned}$$

By Lemma 2.2 in [Wa2] and (3.12) in [WW2], we have for $j < n$

$$\begin{aligned}
&\partial_{x_j} \sigma_{1-2m}(D^{1-2m})(x_0) = \partial_{x_j} [\sqrt{-1} c(\xi) |\xi|^{-2m}] \\
&= \sqrt{-1} [\partial_{x_j} c(\xi)](x_0) |\xi|^{-2m} + \sqrt{-1} c(\xi) \partial_{x_j} (|\xi|^{-2m})(x_0) = 0. \quad (3.7)
\end{aligned}$$

so **case a) I)** vanishes.

case a) II) $r = -1, \ell = 1 - 2m, k = |\alpha| = 0, j = 1$

By (2.8), we get

$$\begin{aligned}
\text{case a) II)} &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \right. \\
&\quad \left. \times \partial_{\xi_n}^2 \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\
&= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{\xi_n}^2 \partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \right. \\
&\quad \left. \times \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \quad (3.8)
\end{aligned}$$

By (2.2.23) in [Wa2], we have

$$\pi_{\xi_n}^+ \partial_{x_n} \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = \frac{\partial_{x_n} [c(\xi')](x_0)}{2(\xi_n - i)} + \sqrt{-1} h'(0) \left[\frac{ic(\xi')}{4(\xi_n - i)} + \frac{c(\xi') + ic(dx_n)}{4(\xi_n - i)^2} \right]. \quad (3.9)$$

So

$$\partial_{\xi_n}^2 \pi_{\xi_n}^+ \partial_{x_n} \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = \frac{\partial_{x_n}[c(\xi')](x_0)}{4(\xi_n - i)^3} + \sqrt{-1}h'(0) \left[\frac{ic(\xi')}{8(\xi_n - i)^3} + \frac{c(\xi') + ic(dx_n)}{24(\xi_n - i)^4} \right]. \quad (3.10)$$

We know that

$$\sigma_{1-2m}(D^{1-2m}) = \frac{\sqrt{-1}[c(\xi') + \xi_n c(dx_n)]}{(1 + \xi_n^2)^m}, \quad (3.11)$$

By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, then we have the equalities:

$$\begin{aligned} \text{tr}[c(\xi')c(dx_n)] &= 0; \quad \text{tr}[c(dx_n)^2] = -2^{m+1}; \quad \text{tr}[c(\xi')^2](x_0)|_{|\xi'|=1} = -2^{m+1}; \\ \text{tr}[\partial_{x_n}c(\xi')c(dx_n)] &= 0; \quad \text{tr}[\partial_{x_n}c(\xi')c(\xi')](x_0)|_{|\xi'|=1} = -2^m h'(0). \end{aligned} \quad (3.12)$$

By (3.10), (3.11) and (3.12), we have

$$\begin{aligned} & \text{trace} \left[\partial_{\xi_n}^2 \partial_{x_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{1-2m}(D^{1-2m}) \right] (x_0)|_{|\xi'|=1} \\ &= \text{trace} \left\{ \left\{ \frac{\partial_{x_n}[c(\xi')](x_0)}{4(\xi_n - i)^3} + \sqrt{-1}h'(0) \left[\frac{ic(\xi')}{8(\xi_n - i)^3} + \frac{c(\xi') + ic(dx_n)}{24(\xi_n - i)^4} \right] \right\} \right. \\ & \quad \left. \times \frac{\sqrt{-1}[c(\xi') + \xi_n c(dx_n)]}{(1 + \xi_n^2)^m} \right\} \\ &= \text{trace} \left\{ \left[\frac{\partial_{x_n}[c(\xi')](x_0)}{4(\xi_n - i)^3} + \frac{(4i - 3\xi_n)h'(0)}{24(\xi_n - i)^4} c(\xi') - \frac{h'(0)c(dx_n)}{24(\xi_n - i)^4} \right] \right. \\ & \quad \left. \times \frac{\sqrt{-1}[c(\xi') + \xi_n c(dx_n)]}{(1 + \xi_n^2)^m} \right\} \\ &= \frac{2^m h'(0)i}{12(\xi_n + i)^m (\xi_n - i)^{m+3}}. \end{aligned} \quad (3.13)$$

By (3.13) and the Cauchy integral formula, we have

$$\text{case a) II} = \text{Vol}(S_{n-2}) \frac{2^m h'(0)\pi}{12(m+2)!} [(\xi_n + i)^{-m}]^{(m+2)}|_{\xi_n=i} dx', \quad (3.14)$$

where $\text{Vol}(S_{n-2})$ is the canonical volume of S_{n-2} and denote the p -th derivative of $f(\xi_n)$ by $[f(\xi_n)]^{(p)}$.

case a) III) $r = -1$, $l = 1 - 2m$, $j = |\alpha| = 0$, $k = 1$

By (2.8) and an integration by parts, we get

$$\begin{aligned} \text{case a) III)} &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \right. \\ & \quad \left. \times \partial_{\xi_n} \partial_{x_n} \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \right. \\ & \quad \left. \times \partial_{x_n} \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.15)$$

By (2.2.29) in [Wa2], we have

$$\partial_{\xi_n}^2 \pi_{\xi_n}^+ \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3}. \quad (3.16)$$

By (3.3), direct computations show that

$$\partial_{x_n} \sigma_{1-2m}(D^{1-2m})(x_0)|_{|\xi'|=1} = \frac{\sqrt{-1} \partial_{x_n} [c(\xi')](x_0)}{(1 + \xi_n^2)^m} - \frac{\sqrt{-1} m h'(0) c(\xi)}{(1 + \xi_n^2)^{m+1}}. \quad (3.17)$$

So by (3.16), (3.17) and (3.12) and the Cauchy integral formula, we have

$$\begin{aligned} \text{case a) III)} &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left\{ \frac{c(\xi') + ic(dx_n)}{(\xi_n - i)^3} \right. \\ &\quad \times \left[\frac{\sqrt{-1} \partial_{x_n} [c(\xi')](x_0)}{(1 + \xi_n^2)^m} - \frac{\sqrt{-1} m h'(0) c(\xi)}{(1 + \xi_n^2)^{m+1}} \right] \Big\} (x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} 2^m h'(0) \times \frac{-i\xi_n^2 - 2m\xi_n + 2mi - i}{(\xi_n + i)^{m+1} (\xi_n - i)^{m+4}} d\xi_n \sigma(\xi') dx' \\ &= \frac{\pi i h'(0) 2^m \text{Vol}(S^{n-2}) dx'}{(m+3)!} \left[\frac{-i\xi_n^2 - 2m\xi_n + 2mi - i}{(\xi_n + i)^{m+1}} \right]^{(m+3)} \Big|_{\xi_n=i}. \end{aligned} \quad (3.18)$$

case b) $r = -2$, $l = 1 - 2m$, $k = j = |\alpha| = 0$

By (2.8), we get

$$\text{case b)} = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} [\pi_{\xi_n}^+ \sigma_{-2}(D^{-1}) \times \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})] (x_0) d\xi_n \sigma(\xi') dx'. \quad (3.19)$$

By (2.2.34)-(2.2.37) in [Wa2], we have

$$\pi_{\xi_n}^+ \sigma_{-2}(D^{-1})(x_0)|_{|\xi'|=1} = B_1 - B_2, \quad (3.20)$$

where

$$B_1 = -\frac{A_1}{4(\xi_n - i)} - \frac{A_2}{4(\xi_n - i)^2}, \quad (3.21)$$

and

$$A_1 = ic(\xi') \sigma_0(D) c(\xi') + ic(dx_n) \sigma_0(D) c(dx_n) + ic(\xi') c(dx_n) \partial_{x_n} [c(\xi')];$$

$$A_2 = [c(\xi') + ic(dx_n)] \sigma_0(D) [c(\xi') + ic(dx_n)] + c(\xi') c(dx_n) \partial_{x_n} c(\xi') - i \partial_{x_n} [c(\xi')]. \quad (3.22)$$

$$B_2 = \frac{h'(0)}{2} \left[\frac{c(dx_n)}{4i(\xi_n - i)} + \frac{c(dx_n) - ic(\xi')}{8(\xi_n - i)^2} + \frac{3\xi_n - 7i}{8(\xi_n - i)^3} [ic(\xi') - c(dx_n)] \right]. \quad (3.23)$$

Similar to (2.2.38) in [Wa2], we have

$$\partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})(x_0)|_{|\xi'|=1} = \sqrt{-1} \left[\frac{c(dx_n)}{(1 + \xi_n^2)^m} - m \times \frac{2\xi_n c(\xi') + 2\xi_n^2 c(dx_n)}{(1 + \xi_n^2)^{m+1}} \right]. \quad (3.24)$$

By (3.23), (3.24) and (3.12), we have

$$\begin{aligned} \text{tr}[B_2 \times \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})(x_0)]|_{|\xi'|=1} &= \frac{\sqrt{-1}}{2} h'(0) \text{trace} \\ &\left\{ \left[\frac{1}{4i(\xi_n - i)} + \frac{1}{8(\xi_n - i)^2} - \frac{3\xi_n - 7i}{8(\xi_n - i)^3} \right] c(dx_n) + \left[\frac{-1}{8(\xi_n - i)^2} + \frac{3\xi_n - 7i}{8(\xi_n - i)^3} \right] ic(\xi') \right\} \\ &\times \left\{ \left[\frac{1}{(1 + \xi_n^2)^m} - \frac{2m\xi_n^2}{(1 + \xi_n^2)^{m+1}} \right] c(dx_n) - \frac{2m\xi_n}{(1 + \xi_n^2)^{m+1}} c(\xi') \right\} \\ &= h'(0) 2^{m-2} \times \frac{(2m-1)\xi_n^3 - 2i(2m-1)\xi_n^2 - (6m-1)\xi_n + 4i}{(\xi_n - i)^2(1 + \xi_n^2)^{m+1}}. \end{aligned} \quad (3.25)$$

By (2.2.40) in [Wa2], we have

$$\begin{aligned} B_1 &= \frac{-1}{4(\xi_n - i)^2} [(2 + i\xi_n)c(\xi')\sigma_0(D)c(\xi') + i\xi_n c(dx_n)\sigma_0(D)c(dx_n) \\ &+ (2 + i\xi_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi') + ic(dx_n)\sigma_0(D)c(\xi') + ic(\xi')\sigma_0(D)c(dx_n) - i\partial_{x_n}c(\xi')]. \end{aligned} \quad (3.26)$$

By (3.24), we have

$$\partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})(x_0)|_{|\xi'|=1} = \sqrt{-1} \left[\frac{1 + (1-2m)\xi_n^2}{(1 + \xi_n^2)^{m+1}} c(dx_n) - \frac{2m\xi_n}{(1 + \xi_n^2)^{m+1}} c(\xi') \right]. \quad (3.27)$$

Similar to Lemma 2.4 in [Wa2], we have

$$\sigma_0(D)(x_0) = c_0 c(dx_n), \quad \text{where } c_0 = \frac{1-n}{4} h'(0). \quad (3.28)$$

By the relation of the Clifford action and $\text{tr}AB = \text{tr}BA$, then we have the equalities:

$$\begin{aligned} \text{tr}[c(\xi')\sigma_0(D)c(\xi')c(dx_n)] &= -c_0 2^{m+1}; \quad \text{tr}[c(dx_n)\sigma_0(D)c(dx_n)^2] = c_0 2^{m+1}; \\ \text{tr}[c(\xi')c(dx_n)\partial_{x_n}c(\xi')c(dx_n)](x_0)|_{|\xi'|=1} &= -2^m h'(0); \quad \text{tr}[c(dx_n)\sigma_0(D)c(\xi')^2] = c_0 2^{m+1}. \end{aligned} \quad (3.29)$$

By (3.26)-(3.29), considering for $i < n$, $\int_{|\xi'|=1} \{\text{odd number product of } \xi_i\} \sigma(\xi') = 0$, then

$$\begin{aligned} \text{tr}[B_1 \times \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})(x_0)]|_{|\xi'|=1} &= \frac{2^m i h'(0)}{4(\xi_n - i)^2(1 + \xi_n^2)^{m+1}} \\ &\cdot \left\{ (n-1)[(2m-1)\xi_n^2 - 2im\xi_n - 1] \right\} \end{aligned}$$

$$+[(1-2m)i\xi_n^3 + 2(1-2m)\xi_n^2 + (2m+1)i\xi_n + 2]\}. \quad (3.30)$$

By (3.25) and (3.30) and the Cauchy integral formula, we have

$$\begin{aligned} \text{case b)} &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[(B_1 - B_2) \times \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})](x_0) d\xi_n \sigma(\xi') dx' \\ &= 2^m h'(0) \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \frac{(4m^2 - 1)\xi_n^2 + (-4m^2 - 6m + 2)i\xi_n - (n+1)}{4(\xi_n - i)^{m+3}(\xi_n + i)^{m+1}} d\xi_n \sigma(\xi') dx' \\ &= \frac{2^{m+1} h'(0) \text{Vol}(S^{n-2}) \pi i dx'}{(m+2)!} \left[\frac{(4m^2 - 1)\xi_n^2 + (-4m^2 - 6m + 2)i\xi_n - (n+1)}{4(\xi_n + i)^{m+1}} \right]^{(m+2)} \Big|_{\xi_n=i}. \end{aligned} \quad (3.31)$$

case c) $r = -1$, $l = -2m$, $k = j = |\alpha| = 0$

By (2.8) and an integration by parts, we get

$$\begin{aligned} \text{case c)} &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{-2m}(D^{-2m+1})](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{-2m}(D^{-2m+1})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.32)$$

By (2.2.44) in [Wa2] and similar to (3.29) in [WW2], we have

$$\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = -\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2}; \quad (3.33)$$

By Lemma 2.2 in [Wa2], we have

$$\sum_{j=1}^{2m+2} \partial_{\xi_j} (|\xi|^{-2m}) \partial_{x_j} (c(\xi))(x_0)|_{|\xi'|=1} = -2m \xi_n (1 + \xi_n^2)^{-m-1} \partial_{x_n} [c(\xi')](x_0). \quad (3.34)$$

$$\begin{aligned} &\left[-\sqrt{-1} \sum_{k=0}^{m-2} \sum_{\mu=1}^{2m+2} \partial_{\xi_\mu} \sigma_2^{-m+k+1}(D^2) \partial_{x_\mu} \sigma_2^{-1}(D^2) (\sigma_2(D^2))^{-k} \right] \sqrt{-1} c(\xi)(x_0)|_{|\xi'|=1} \\ &= -2c(\xi) h'(0) \xi_n \left(-\frac{m^2}{2} + \frac{m}{2} \right) (1 + \xi_n^2)^{-m-2}. \end{aligned} \quad (3.35)$$

By (3.26) in [WW2], we have

$$\begin{aligned} \sigma_{-3}(D^{-2})(x_0)|_{|\xi'|=1} &= \frac{-i}{(1 + \xi_n^2)^2} \left(-\frac{1}{2} h'(0) \sum_{k < n} \xi_k c(\widetilde{e_k}) c(\widetilde{e_n}) + \frac{n-1}{2} h'(0) \xi_n \right) - \frac{2ih'(0)\xi_n}{(1 + \xi_n^2)^3} \\ &= \frac{-i}{(1 + \xi_n^2)^2} \left(-\frac{1}{2} h'(0) c(\xi') c(dx_n) + \frac{n-1}{2} h'(0) \xi_n \right) - \frac{2ih'(0)\xi_n}{(1 + \xi_n^2)^3}. \end{aligned} \quad (3.36)$$

So by (3.5), (3.34)-(3.36), we have

$$\begin{aligned} \sigma_{-2m}(D^{1-2m})(x_0)|_{|\xi'|=1} &= \frac{(1-n)h'(0)c(dx_n)}{4(1+\xi_n^2)^m} - 2m\xi_n(1+\xi_n^2)^{-m-1}\partial_{x_n}[c(\xi')](x_0) \\ &+ mi(1+\xi_n^2)^{-m+1}[c(\xi')+\xi_nc(dx_n)] \times \left[\frac{-ih'(0)c(\xi')c(dx_n)}{2(1+\xi_n^2)^2} + \frac{(-n+1)h'(0)i\xi_n}{2(1+\xi_n^2)^2} - \frac{2ih'(0)\xi_n}{(1+\xi_n^2)^3} \right] \\ &- [c(\xi')+\xi_nc(dx_n)]h'(0)\xi_n(-m^2+m)(1+\xi_n^2)^{-m-2}. \end{aligned} \quad (3.37)$$

By (3.33) and (3.37), we have

$$\begin{aligned} \text{trace}[\partial_{\xi_n}\pi_{\xi_n}^+\sigma_{-1}(D^{-1}) \times \sigma_{-2m}(D^{-2m+1})](x_0)|_{|\xi'|=1} &= \frac{2^mh'(0)}{4(\xi_n-i)^{m+3}(\xi_n+i)^{m+2}} \\ &\times \left[(2nm-2m-n+1)i\xi_n^3 + (-2m+1+2\pi)\xi_n^2 \right. \\ &\left. + (2nm-2m+4m^2-n+1)i\xi_n + (n-1+2\pi) \right]. \end{aligned} \quad (3.38)$$

Then by the Cauchy integral formula, we get

$$\begin{aligned} \text{case c)} &= \frac{-\pi 2^{m-1}h'(0)\text{Vol}(S^{n-2})dx'}{(m+2)!} \left\{ \frac{1}{(\xi_n+i)^{m+2}} \left[(2nm-2m-n+1)i\xi_n^3 \right. \right. \\ &\left. \left. + (-2m+1+2\pi)\xi_n^2 + (2nm-2m+4m^2-n+1)i\xi_n + (n-1+2\pi) \right] \right\} \Big|_{\xi_n=i}^{(m+2)}. \end{aligned} \quad (3.39)$$

Similar to (3.41) in [WW2], we have for the extrinsic curvature

$$K(x_0) = \frac{1-n}{2}h'(0). \quad (3.40)$$

By (3.14),(3.18),(3.31), (3.39) and (3.40), we have

$$\Phi = \frac{2}{1-n}\text{Vol}(S^{n-2})L_0 \int_{\partial M} K \text{dvol}_{\partial M}, \quad (3.41)$$

where

$$\begin{aligned} L_0 &= \frac{2^m\pi}{12(m+2)!}[(\xi_n+i)^{-m}]^{(m+2)}\Big|_{\xi_n=i} + \frac{\pi i 2^m}{(m+3)!} \left[\frac{-i\xi_n^2 - 2m\xi_n + 2mi - i}{(\xi_n+i)^{m+1}} \right]^{(m+3)} \Big|_{\xi_n=i} \\ &+ \frac{2^{m+1}\pi i}{(m+2)!} \left[\frac{(4m^2-1)\xi_n^2 + (-4m^2-6m+2)i\xi_n - (n+1)}{4(\xi_n+i)^{m+1}} \right]^{(m+2)} \Big|_{\xi_n=i} \\ &+ \frac{-\pi 2^{m-1}}{(m+2)!} \left\{ \frac{1}{(\xi_n+i)^{m+2}} \left[(2nm-2m-n+1)i\xi_n^3 \right. \right. \\ &\left. \left. + (-2m+1+2\pi)\xi_n^2 + (2nm-2m+4m^2-n+1)i\xi_n + (n-1+2\pi) \right] \right\} \Big|_{\xi_n=i}^{(m+2)}. \end{aligned} \quad (3.42)$$

By (2.9), (3.41) and (3.42), we prove Theorem 1.1.

Nextly, for $n = 2m + 1$ -dimensional spin manifolds with boundary, we compute $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(1-2m)}]$. By Proposition 3.5 in [WW2], we have

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(1-2m)}] = \int_{\partial M} \Phi. \quad (3.43)$$

From the formula (2.8) for the definition of Φ , and the sum is taken over $r - k + |\alpha| + \ell - j - 1 = -(2m + 1)$, $r \leq -1$, $\ell \leq 1 - 2m$, then $r = -1$, $\ell = 1 - 2m$, $k = |\alpha| = j = 0$,

$$\begin{aligned} \Phi &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}_{S(TM)} [\pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{1-2m}(D^{1-2m})] d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}_{S(TM)} [\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{1-2m}(D^{1-2m})] d\xi_n \sigma(\xi') dx' \end{aligned} \quad (3.44)$$

By (5.3) in [Wa2], we have

$$\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1})(x_0)|_{|\xi'|=1} = -\frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)^2}. \quad (3.45)$$

So

$$\begin{aligned} &\text{trace}_{S(TM)} [\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{1-2m}(D^{1-2m})](x_0)|_{|\xi'|=1} \\ &= \frac{-i}{2(\xi_n - i)^2(1 + \xi_n^2)^m} \text{trace}_{S(TM)} \{ [c(\xi') + ic(dx_n)] [c(\xi') + \xi_n c(dx_n)] \} \\ &= \frac{-2^{m-1}}{(\xi_n - i)^{m+1}(\xi_n + i)^m}. \end{aligned} \quad (3.46)$$

Then

$$\int_{\partial M} \Phi = \frac{2^{-m} m(m+1) \cdots (2m-1) \text{Vol}_{\partial M} \text{Vol}(S^{n-2}) \pi}{m!}. \quad (3.47)$$

So we have

Theorem 3.1 *Let M be a $n = 2m + 1$ -dimensional compact spin manifold with the boundary ∂M , then*

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(1-2m)}] = \frac{2^{-m} m(m+1) \cdots (2m-1) \text{Vol}_{\partial M} \text{Vol}(S^{n-2}) \pi}{m!}. \quad (3.48)$$

Nextly for $n = 2m + 1$ -dimensional spin manifolds with boundary, we compute $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(2-2m)}]$. Now we can compute Φ (see formula (2.8) for the definition of Φ). Since the sum is taken over $r + \ell - k - j - |\alpha| - 1 = -(2m + 1)$, $r \leq -1$, $\ell \leq 2 - 2m$, then we have the $\int_{\partial M} \Phi$ is the sum of the following five cases:

case a) I) $r = -1$, $l = 2 - 2m$, $k = j = 0$, $|\alpha| = 1$

$$\begin{aligned} \text{case a) I)} &= - \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{|\alpha|=1} \text{trace} \left[\partial_{\xi'}^{\alpha} \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \right. \\ &\quad \left. \times \partial_{x'}^{\alpha} \partial_{\xi_n} \sigma_{2-2m}(D^{2-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.49)$$

By (3.12) in [WW2], we have for $j < n$, $\partial_{x_j} \sigma_{2-2m}(D^{2-2m})(x_0) = 0$. So case a) I) = 0.

case a) II) $r = -1$, $l = 2 - 2m$, $k = |\alpha| = 0$, $j = 1$

By (2.8), we get

$$\begin{aligned} \text{case a) II)} &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{x_n} \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \right. \\ &\quad \left. \times \partial_{\xi_n}^2 \sigma_{1-2m}(D^{1-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.50)$$

By (3.9) and (3.16) in [WW2] and

$$\text{tr}[c(\xi')] = \text{tr}[c(dx_n)] = \text{tr}[\partial_{x_n} c(\xi')] = 0, \quad (3.51)$$

we have case a) II) = 0.

case a) III) $r = -1$, $l = 2 - 2m$, $j = |\alpha| = 0$, $k = 1$

By (2.8) and an integration by parts, we get

$$\begin{aligned} \text{case a) III)} &= -\frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{\xi_n} \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \right. \\ &\quad \left. \times \partial_{\xi_n} \partial_{x_n} \sigma_{2-2m}(D^{2-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx' \\ &= \frac{1}{2} \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} \left[\partial_{\xi_n}^2 \pi_{\xi_n}^{+} \sigma_{-1}(D^{-1}) \right. \\ &\quad \left. \times \partial_{x_n} \sigma_{2-2m}(D^{2-2m}) \right] (x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.52)$$

By (3.16) and (3.21) in [WW2] and (3.51), we have case a) III) = 0.

case b) $r = -2$, $l = 2 - 2m$, $k = j = |\alpha| = 0$

By (2.8), we get

$$\begin{aligned} \text{case b)} &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace} [\pi_{\xi_n}^{+} \sigma_{-2}(D^{-1}) \times \partial_{\xi_n} \sigma_{2-2m}(D^{2-2m})] (x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.53)$$

By (3.33) in [WW2] and (3.20)-(3.23), (3.51), (3.53) and

$$\text{tr}[c(\xi') \sigma_0(D) c(\xi')] = \text{tr}[c(dx_n) \sigma_0(D) c(dx_n)] = \text{tr}[c(\xi') c(dx_n) \partial_{x_n} c(\xi')] = 0, \quad (3.54)$$

we have case b) = 0.

case c) $r = -1, l = 1 - 2m, k = j = |\alpha| = 0$

By (2.8) and an integration by parts, we get

$$\begin{aligned} \text{case c)} &= -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{1-2m}(D^{-2m+2})](x_0) d\xi_n \sigma(\xi') dx' \\ &= i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{trace}[\partial_{\xi_n} \pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \sigma_{1-2m}(D^{-2m+2})](x_0) d\xi_n \sigma(\xi') dx'. \end{aligned} \quad (3.55)$$

By (3.33) and (3.8),(3.26),(3.29) in [WW2] and (3.51),(3.54), (3.55), we have case c) = 0. So we get $\int_{\partial M} \Phi = 0$.

Theorem 3.2 *Let M be a $n = 2m + 1$ -dimensional compact spin manifold with the boundary ∂M , then*

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(-2m+2)}] = \frac{(2-n)(2\pi)^{n/2}}{12\Gamma(n/2)} \int_M \text{sdvol}_M. \quad (3.56)$$

Nextly, for $n = 2m$ -dimensional spin manifolds with boundary, we compute $\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(2-2m)}]$. By Proposition 3.5 in [WW2], we have

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(2-2m)}] = \int_{\partial M} \Phi. \quad (3.57)$$

From the formula (2.8) for the definition of Φ , and the sum is taken over $r - k + |\alpha| + \ell - j - 1 = -2m, r \leq -1, \ell \leq 2 - 2m$, then $r = -1, \ell = 2 - 2m, k = |\alpha| = j = 0$,

$$\Phi = -i \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \text{traces}_{S(TM)}[\pi_{\xi_n}^+ \sigma_{-1}(D^{-1}) \times \partial_{\xi_n} \sigma_{2-2m}(D^{2-2m})] d\xi_n \sigma(\xi') dx'. \quad (3.58)$$

By (2.2.44) in [Wa2] and (3.47) in [WW2] and (3.51),(3.58), we get $\Phi = 0$.

Theorem 3.3 *Let M be a $n = 2m$ -dimensional compact spin manifold with the boundary ∂M , then*

$$\widetilde{\text{Wres}}[\pi^+ D^{-1} \circ \pi^+ D^{(-2m+2)}] = 0. \quad (3.59)$$

Similar to Theorem 3.2, we can get

Theorem 3.4 *Let M be a $n = 2m + 3$ -dimensional compact spin manifold with the boundary ∂M , then*

$$\widetilde{\text{Wres}}[\pi^+ D^{-2} \circ \pi^+ D^{(-2m+1)}] = \frac{(2-n)(2\pi)^{n/2}}{12\Gamma(n/2)} \int_M \text{sdvol}_M. \quad (3.60)$$

References

- [Ac] T. Ackermann, A note on the Wodzicki residue, *J. Geom. Phys.*, 20, 404-406, 1996.
- [Ad] M. Adler.: On a trace functional for formal pseudo-differential operators and the symplectic structure of Korteweg-de Vries type equations, *Invent. Math.* 50, 219-248, 1979.
- [Co1] A. Connes, Quantized calculus and applications, XIth International Congress of Mathematical Physics (paris,1994), 15-36, Internat Press, Cambridge, MA, 1995.
- [Co2] A. Connes. The action functional in noncommutative geometry, *Comm. Math. Phys.*, 117:673-683, 1998.
- [FGLS] B. V. Fedosov, F. Golse, E. Leichtnam, and E. Schrohe. The noncommutative residue for manifolds with boundary, *J. Funct. Anal*, 142:1-31,1996.
- [Gu] V.W. Guillemin, A new proof of Weyl's formula on the asymptotic distribution of eigenvalues, *Adv. Math.* 55 no.2, 131-160, 1985.
- [Ka] D. Kastler, The Dirac operator and gravitation, *Commun. Math. Phys*, 166:633-643, 1995.
- [KW] W. Kalau and M.Walze, Gravity, non-commutative geometry, and the Wodzicki residue, *J. Geom. Phys.*, 16:327-344, 1995.
- [Po] R. Ponge, Noncommutative Geometry and lower dimensional volumes in Riemannian geometry, *Lett. Math. Phys.* 83, 1-19, 2008.
- [Sc] E. Schrohe, Noncommutative residue, Dixmier's trace, and heat trace expansions on manifolds with boundary, *Contemp. Math.* 242, 161-186, 1999.
- [WW1] J. Wang and Y. Wang, The Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary, *J. Math. Phys.* 56, no. 5, 052501, 14 pp, 2015
- [WW2] J. Wang and Y. Wang, A general Kastler-Kalau-Walze type theorem for manifolds with boundary, *Int. J. Geom. Methods Mod. Phys.* 13, no. 1, 1650003, 16 pp, 2016.
- [Wa1] Y. Wang, Differential forms and the Wodzicki residue for manifolds with boundary. *J. Geom. Phys.* 56, 731-753, 2006.
- [Wa2] Y. Wang, Gravity and the noncommutative residue for manifolds with boundary. *Lett. Math. Phys.* 80, 37-56, 2007.
- [Wa3] Y. Wang, Lower-dimensional volumes and Kastler-kalau-Walze type theorem for manifolds with boundary. *Commun. Theor. Phys.* Vol 54, 38-42, 2010.
- [Wo] M. Wodzicki.: Local invariants of spectral asymmetry. *Invent. Math.* 75(1), 143-178, 1995.