

# LOCAL $k$ -CONNECTEDNESS OF AN INVERSE LIMIT OF POLYHEDRA

G. C. BELL AND A. NAGÓRKO

ABSTRACT. We provide an easily verifiable condition for local  $k$ -connectedness of an inverse limit of polyhedra.

## 1. INTRODUCTION

The Nöbeling space characterization theorem [3, 1, 2, 10, 12] states that if a space is strongly universal in the class of  $n$ -dimensional separable complete metric spaces and is  $k$ -connected and locally  $k$ -connected for each  $k < n$ , then it is homeomorphic to the  $n$ -dimensional Nöbeling space  $N_n^{2n+1}$ .

In geometric group theory many spaces arise naturally as inverse limits of polyhedra (simplicial complexes endowed with the metric topology). In particular, boundaries at infinity of hyperbolic spaces can be expressed as such. Striking examples of applications of the Nöbeling space characterization theorem are proofs that the boundary at infinity of the curve complex of the  $(n + 5)$ -punctured 2-dimensional sphere is homeomorphic to the  $n$ -dimensional Nöbeling space  $N_n^{2n+1}$  [8, 7].

In the present paper we prove a condition for local  $k$ -connectedness of an inverse limit of polyhedra that is easy to verify. It is designed to aid detection of local  $k$ -connectedness in many examples of spaces arising in geometric group theory. We prove the following theorem.

**Definition 1.1.** Let  $K$  and  $L$  be simplicial complexes. We say that a map  $p: K \rightarrow L$  is  $n$ -**regular** if it is quasi-simplicial (i.e. it is simplicial into the first barycentric subdivision  $\beta L$  of  $L$ ) and if for each simplex  $\delta$  of  $\beta L$  the inverse image  $p^{-1}(\delta)$  has vanishing homotopy groups in dimensions less than  $n$  (regardless of the choice of basepoint).

**Theorem 1.2.** *Let*

$$X = \varprojlim (K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \cdots).$$

*Assume that for each  $i$  the following conditions are satisfied.*

---

2010 *Mathematics Subject Classification.* Primary 55M15, 54C55; Secondary 55M10.

*Key words and phrases.* local  $k$ -connectedness, Nöbeling space.

This research was supported by the NCN (Narodowe Centrum Nauki) grant no. 2011/01/D/ST1/04144.

(I)  $K_i$  is a locally finite-dimensional simplicial complex endowed with the metric topology.

(II)  $p_i$  is a quasi-simplicial map that is surjective and  $n$ -regular.

Then  $X$  is locally  $k$ -connected for each  $k < n$ , and each short projection  $\pi_i^m: K_m \rightarrow K_i$  and each long projection  $\pi_i: X \rightarrow K_i$  is a weak  $n$ -homotopy equivalence (it induces homomorphisms on homotopy groups in dimensions less than  $n$ , regardless of the choice of basepoint).

Note that there is no assumption of local finiteness of complexes in the statement of Theorem 1.2.

It is known that (under suitable assumptions) if the bonding maps in the inverse limit are  $n$ -soft, then the inverse limit is  $k$ -connected for  $k < n$ , [5]. The condition for local  $k$ -connectedness given in the present paper may be regarded as a combinatorial analog of this statement for inverse limits of polyhedra. Note that  $n$ -regular bonding maps need not be  $n$ -soft.

## 2. PRELIMINARIES

In this section we set the basic definitions and reference known results that will be used in the later sections.

### 2.1. Absolute extensors in dimension $n$ .

**Definition 2.1.** We say that a space  $X$  is  **$k$ -connected** if each map  $\varphi: S^k \rightarrow X$  from a  $k$ -dimensional sphere into  $X$  is null-homotopic in  $X$ .

**Definition 2.2.** We say that a space  $X$  is **locally  $k$ -connected** if for each point  $x \in X$  and each open neighborhood  $U \subset X$  of  $x$  there exists an open neighborhood  $V$  of  $x$  such that each map  $\varphi: S^k \rightarrow V$  from a  $k$ -dimensional sphere into  $V$  is null-homotopic in  $U$ .

**Definition 2.3.** We say that a metric space  $X$  is an **absolute neighborhood extensor in dimension  $n$**  if every map into  $X$  from a closed subset  $A$  of an  $n$ -dimensional metric space extends over an open neighborhood of  $A$ . The class of absolute neighborhood extensors in dimension  $n$  is denoted by  $ANE(n)$  and its elements are called  $ANE(n)$ -spaces.

**Definition 2.4.** We say that a metric space  $X$  is an **absolute extensor in dimension  $n$**  if every map into  $X$  from a closed subset of an  $n$ -dimensional metric space  $Y$  extends over the entire space  $Y$ . The class of absolute extensors in dimension  $n$  is denoted by  $AE(n)$  and its elements are called  $AE(n)$ -spaces.

**Definition 2.5.** Let  $\mathcal{C}$  be a class of topological spaces. We let  $AE(\mathcal{C})$  denote the class of **absolute extensors for all spaces from the class  $\mathcal{C}$** . We write  $AE(X)$  for  $AE(\{X\})$ .

Absolute extensors and absolute neighborhood extensors in dimension  $n$  were characterized by Dugundji in the following theorem.

**Theorem 2.6** ([6]). *Let  $X$  be a metric space. Then,*

- (1)  $X \in ANE(n) \iff X$  is locally  $k$ -connected for all  $k < n$ ; and
- (2)  $X \in AE(n) \iff X \in ANE(n)$  and  $X$  is  $k$ -connected for all  $k < n$ .

**Lemma 2.7** (cf. [15, Theorem 6.1.8]). *Assume that  $A_1 \subset A_2 \subset \dots$  is a sequence of subsets of a metric space such that each  $A_i$  is closed and for each  $i$ ,  $A_i \subset \text{Int } A_{i+1}$ . If for each  $i$ ,  $A_i$  is  $AE(n)$ , then  $A = \bigcup_i A_i$  is  $AE(n)$ .*

**2.2. Polyhedra.** For a simplicial complex, the underlying polyhedron has two topologies, the Whitehead (weak) topology and the metric topology. The metric topology is the topology of point-wise convergence of barycentric coordinates [9, p. 100]. The weak topology is metrizable if and only if the complex is locally finite, and it coincides with the metric topology in this case. Since we work in the metric category with complexes that are not locally finite, **we always assume the metric topology on simplicial complexes** [11, 9].

**Definition 2.8.** Let  $K$  be a simplicial complex. We let  $\tau(K)$  denote the **triangulation of  $K$**  (the set of simplices of  $K$ ). We let  $V(K)$  denote the **vertex set of  $K$** . We let  $\beta K$  denote the **barycentric subdivision of  $K$**  (i.e., the same space but with a finer triangulation  $\tau(\beta K)$ ).

**Definition 2.9.** Let  $K$  be a simplicial complex. Let  $\kappa > 0$ . Let  $\ell_1(V(K))$  denote the Banach space  $\{x \in \mathbb{R}^{V(K)} : \sum_{v \in V(K)} |(x)_v| < \infty\}$ , where  $(x)_v$  denotes the  $v$ -th coordinate, equipped with the standard  $\|\cdot\|_1$  norm. For  $v \in v(K)$  define  $e_v : V(K) \rightarrow \mathbb{R}$  by the formula  $e_v(w) = 0$  for  $w \neq v$  and  $e_v(v) = \kappa$ . We embed each vertex  $v$  of  $K$  as  $e_v \in \ell_1(V(K))$  and extend this embedding to  $K$  to be affine on each simplex of  $K$ . We consider  $K$  to be a subspace of  $\ell_1(V(K))$ . We call the induced metric on  $K$  the **metric of scale  $\kappa$  on  $K$** . For  $\kappa = 1$  it is the standard metric, as defined in [9, p. 100]. The topology induced by this metric is called **the metric topology**.

**Definition 2.10.** A **polyhedron** is a simplicial complex endowed with the metric topology.

**Definition 2.11.** Let  $K$  and  $L$  be polyhedra. We say that  $K$  is a **full subpolyhedron** of  $L$  if whenever the vertices  $v_0, \dots, v_n$  span a simplex in  $K$  and each  $v_i$  is a vertex in  $L$ , then the  $v_i$  span a simplex in  $L$ .

**Lemma 2.12.** *A locally finite-dimensional polyhedron is a complete metric  $ANE(\infty)$ -space.*

*Proof.* It is complete by [9, Lemma 11.5]. It is an  $ANE(\infty)$ -space by [9, Theorem 11.3].  $\square$

**Definition 2.13.** Let  $K$  and  $L$  be simplicial complexes and let  $p: K \rightarrow L$ . We say that  $p$  is **quasi-simplicial** if it is a simplicial map into  $\beta L$ .

**Lemma 2.14** (cf. [4]). *Assume that  $K$  and  $L$  are polyhedra endowed with metrics of scale  $\kappa$  and  $\lambda$  respectively. If  $p: K \rightarrow L$  is quasi-simplicial, then it is  $\frac{\lambda}{2\kappa}$ -Lipschitz.*

### 2.3. Weak $n$ -homotopy.

**Definition 2.15.** We say that a map is a **weak  $n$ -homotopy equivalence** if it induces isomorphisms on homotopy groups of dimensions less than  $n$ , regardless of the choice of basepoint.

**Definition 2.16.** Let  $\mathcal{F}$  be a cover of a space  $X$ . We say that the two maps  $f, g: Y \rightarrow X$  are  **$\mathcal{F}$ -close** if for each  $y \in Y$  there exists  $F \in \mathcal{F}$  such that  $f(y), g(y) \in F$ .

**Definition 2.17.** Let  $\mathcal{U}$  be a cover of a space  $X$ . We say that two maps  $f, g: Y \rightarrow X$  are  **$\mathcal{U}$ -homotopic** if there exists a homotopy  $H: Y \times [0, 1] \rightarrow X$  whose paths refine  $\mathcal{U}$ , i.e. for each  $y \in Y$  there exists  $U \in \mathcal{U}$  such that  $H(\{y\} \times [0, 1]) \subset U$ .

### 2.4. Carrier Theorem.

**Definition 2.18.** Let  $\mathcal{C}$  be a class of topological spaces. We say that a cover  $\mathcal{F}$  of a topological space is a  **$\mathcal{C}$ -cover**, if for each  $\mathcal{A} \subset \mathcal{F}$  the intersection  $\bigcap \mathcal{A}$  is either empty or belongs to  $\mathcal{C}$ .

**Definition 2.19.** We say that a cover is **locally finite-dimensional** if its nerve is locally finite-dimensional.

**Definition 2.20.** A **carrier** is a function  $C: \mathcal{F} \rightarrow \mathcal{G}$  from a cover  $\mathcal{F}$  of a space  $X$  into a collection  $\mathcal{G}$  of subsets of a topological space such that for each  $\mathcal{A} \subset \mathcal{F}$  if  $\bigcap_{A \in \mathcal{A}} A \neq \emptyset$ , then  $\bigcap_{A \in \mathcal{A}} C(A) \neq \emptyset$ . We say that a map  $f$  is **carried by  $C$**  if it is defined on a closed subset of  $X$  and  $f(F) \subset C(F)$  for each  $F \in \mathcal{F}$ . Here we write  $f(F)$  to mean  $f(F \cap \text{dom}(f))$ .

**Carrier Theorem** ([13]). *Assume that  $C: \mathcal{F} \rightarrow \mathcal{G}$  is a carrier such that  $\mathcal{F}$  is a cover of a space  $X$  and  $\mathcal{G}$  is an  $AE(X)$ -cover of another space. If  $\mathcal{F}$  is closed, locally finite, and locally finite-dimensional, then each map carried by  $C$  extends to a map of the entire space  $X$ , also carried by  $C$ .*

**Corollary 2.21** ([13]). *If  $\mathcal{F}$  is a closed locally finite locally finite-dimensional  $AE(n)$ -cover of a space  $Y$ , then any two  $\mathcal{F}$ -close maps from a metric space of dimension less than  $n$  into  $Y$  are  $\mathcal{F}$ -homotopic.*

**2.5. Covers.** We regard covers as indexed collections of sets and use the usual notation,  $\mathcal{F} = \{F_i\}_{i \in I}$ , where  $I$  denotes the indexing set.

**Definition 2.22.** Let  $K$  be a polyhedron. Let  $L \subset K$  be a subcomplex of  $K$ . The **open star**  $\text{ost}_K L$  **of  $L$  in  $K$**  is the complement of the union of all simplices of  $K$  that do not intersect  $L$ :

$$\text{ost}_K L = K \setminus \bigcup \{\delta \in \tau(K) : \delta \cap L = \emptyset\}.$$

The **barycentric star**  $\text{bst}_K L$  **of  $L$  in  $K$**  is the union of all simplices of  $\beta K$  that intersect  $L$ :

$$\text{bst}_K L = \bigcup \{\delta \in \tau(\beta K) : \delta \cap L \neq \emptyset\}.$$

We let

$$\mathcal{O}_K = \{\text{ost}_K \{v\} : v \in V(K)\}$$

denote the **cover of  $K$  by open stars of vertices**. We let

$$\mathcal{B}_K = \{\text{bst}_K \{v\} : v \in V(K)\}$$

denote a **cover of  $K$  by barycentric stars of vertices**.

**Lemma 2.23** ([14]). *Let  $K$  be a polyhedron. The cover  $\mathcal{B}_K$  by barycentric stars of vertices is a closed locally finite  $AE(\infty)$ -cover of  $K$ . Moreover, if  $K$  is locally finite-dimensional, then  $\mathcal{B}_K$  is locally finite-dimensional. The cover  $\mathcal{O}_K$  by open stars of vertices is an open  $AE(\infty)$ -cover of  $K$ .*

**Definition 2.24.** Let

$$X = \varprojlim \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \cdots \right)$$

Let  $v(K_i)$  denote the set of vertices of  $K_i$ . We let

$$\mathcal{O}_{K_i} = \{O_v = \text{ost}_{K_i} v\}_{v \in v(K_i)}$$

be the cover of  $K_i$  by open stars of vertices of  $K_i$  (see Definition 2.22) and

$$\mathcal{O}_i = \{\pi_i^{-1}(\text{ost}_{K_i} v)\}_{v \in v(K_i)}$$

be the cover of  $X$  by sets of threads that pass through elements of  $\mathcal{O}_{K_i}$ .

**Definition 2.25.** Let

$$X = \lim_{\leftarrow} \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \cdots \right)$$

Let  $v(K_i)$  denote the set of vertices of  $K_i$ . We let

$$\mathcal{B}_{K_i} = \{\text{bst}_{K_i} v\}_{v \in v(K_i)}$$

be the cover of  $K_i$  by barycentric stars of vertices of  $K_i$  (see Definition 2.22) and

$$\mathcal{B}_i = \{\pi_i^{-1}(\text{bst}_{K_i} v)\}_{v \in v(K_i)}$$

be the cover of  $X$  by sets of threads that pass through elements of  $\mathcal{B}_{K_i}$ .

**Definition 2.26.** We say that a cover  $\mathcal{F} = \{F_i\}_{i \in I}$  is **isomorphic** to a cover  $\mathcal{G} = \{G_i\}_{i \in I}$  if for each  $J \subset I$  we have

$$\bigcap_{j \in J} F_j \neq \emptyset \iff \bigcap_{j \in J} G_j \neq \emptyset.$$

Note the identical indexing set of  $\mathcal{F}$  and  $\mathcal{G}$ .

**Definition 2.27.** Let  $p: Y \rightarrow Z$  be a map. Let  $\mathcal{F} = \{F_i\}_{i \in I}$  be a cover of  $Z$ . A **pull-back of  $\mathcal{F}$**  is a cover  $p^{-1}(\mathcal{F})$  of  $Y$  defined by the formula

$$p^{-1}(\mathcal{F}) = \{p^{-1}(F_i)\}_{i \in I}.$$

The pull-back  $p^{-1}(\mathcal{F})$  retains the indexing set  $I$  of  $\mathcal{F}$ .

**Lemma 2.28.** *Let  $f: X \rightarrow Y$  be a map, let  $\mathcal{G}$  be a cover of  $Y$  and let  $f^{-1}(\mathcal{G})$  be a pull-back of  $\mathcal{G}$ .*

- (1) *If  $f$  is surjective, then the covers  $f^{-1}(\mathcal{G})$  and  $\mathcal{G}$  are isomorphic.*
- (2) *If  $\mathcal{G}$  is open / closed / locally finite / locally finite-dimensional, then so is  $f^{-1}(\mathcal{G})$ .*

### 3. NERVE THEOREM

A Nerve Theorem in its abstract form states that if two spaces admit isomorphic  $AE(n)$ -covers, then they are weak  $n$ -homotopy equivalent. For general spaces, some local finiteness and dimension restrictions are placed on the covers [13]. In our case we need such a theorem without these restrictions, which we are able to prove for polyhedra covered by subcomplexes.

The goal of this section is to prove the following theorem.

**Theorem 3.1.** *If a quasi-simplicial map  $p: K \rightarrow L$  of polyhedra is surjective and  $n$ -regular, then it is a weak  $n$ -homotopy equivalence and the pull-back  $p^{-1}(\mathcal{B}_L)$  is an  $AE(n)$ -cover of  $K$ .*

The main tool used in the proof is Theorem 3.5, which when applied to a canonical map into the nerve of a cover [13] implies the usual Nerve Theorem.

### 3.1. Nerve Theorem for non locally finite covers.

**Lemma 3.2.** *If  $K$  is a full subpolyhedron of a polyhedron  $L$ , then*

- (1)  $K$  is a deformation retract of  $\text{bst}_L K$ ; and
- (2)  $K$  is a deformation retract of  $\text{ost}_L K$ .

*Proof.* We let  $V_L$  denote the set of vertices of  $L$  and  $V_K \subset V_L$  denote the set of vertices of  $K$ . We regard  $L$  as a subspace of  $\ell_1(V_L)$  (see Definition 2.9). Let  $p: L \rightarrow \ell_1(V_L)$  be the map that sends all coordinates that do not belong to  $K$  to 0, defined by the formula

$$(p(x))_v = \begin{cases} 0 & v \in V_L \setminus V_K \\ (x)_v & v \in V_K \end{cases}.$$

We have

$$\text{ost}_L K = \{x \in L: p(x) \neq 0\}.$$

Let  $q: \text{ost}_L K \rightarrow L$  be defined by the formula

$$q(x) = \frac{p(x)}{\|p(x)\|_1}.$$

We define a map  $\Phi: \text{ost}_L K \times [0, 1] \rightarrow L$  by the formula

$$\Phi(x, t) = t \cdot q(x) + (1 - t) \cdot x.$$

Observe that  $\text{supp } \Phi(x, t) \subset \text{supp } (p(x))$  and  $V_K \cap \text{supp } \Phi(x, t) \neq \emptyset$ , hence  $\Phi(x, t) \in \text{ost}_L K$ . Note that  $\Phi(x, 1) \in K$  because  $K$  is a full subcomplex. Hence  $\Phi$  is a deformation retraction of  $\text{ost}_L K$  to  $K$ .

Observe that

$$\text{bst}_L K = \{x \in L: \exists_{v \in V_K} (x)_v \geq \max\{(x)_w: w \in V_L\}\}.$$

Since  $\Phi$  preserves the relation  $\exists_{v \in V_K} (x)_v \geq \max\{(x)_w: w \in V_L\}$ ,  $\Phi$  restricted to  $\text{bst}_L K \times [0, 1]$  is into  $\text{bst}_L K$ , hence it is a deformation retraction of  $\text{bst}_L K$  to  $K$ .  $\square$

**Corollary 3.3.** *If  $K$  is a polyhedron and  $\mathcal{K} = \{K_i\}_{i \in I}$  is an  $AE(n)$ -cover of  $K$  by subcomplexes, then the collection*

$$\text{ost}_{\beta K} \mathcal{K} = \{\text{ost}_{\beta K} K_i\}_{i \in I}$$

*is an open  $AE(n)$ -cover of  $K$ . Moreover,  $\text{ost}_{\beta K} \mathcal{K}$  and  $\mathcal{K}$  are isomorphic covers of  $K$ .*

*Proof.* Observe that if  $\{A_j\}_{j \in J}$  is a collection of subcomplexes of a polyhedron  $K$ , then

$$\bigcap_{j \in J} \text{ost}_{\beta K} A_j = \text{ost}_{\beta K} \bigcap_{j \in J} A_j.$$

An application of Lemma 3.2 finishes the proof.  $\square$

**Lemma 3.4.** *If  $\mathcal{U}$  is an open  $AE(n)$ -cover of a space  $Y$ , then any two  $\mathcal{U}$ -close maps from a metric space of dimension less than  $n$  into  $Y$  are  $\mathcal{U}$ -homotopic.*

*Proof.* Let  $\mathcal{U}$  be an open  $AE(n)$ -cover of a metric space  $Y$ . Let  $X$  be a metric space of dimension less than  $n$ . Let  $f, g: X \rightarrow Y$  be  $\mathcal{U}$ -close. We have to show that  $f$  and  $g$  are homotopic by a homotopy whose paths refine  $\mathcal{U}$ .

Let  $\mathcal{V} = \{V_U\}_{U \in \mathcal{U}}$  be the collection of subsets of  $X$  defined by  $V_U = f^{-1}(U) \cap g^{-1}(U)$ . Since  $f$  and  $g$  are  $\mathcal{U}$ -close,  $\mathcal{V}$  is an open cover of  $X$ .

Let  $\mathcal{W}$  be a closed, locally finite cover of  $X$  with multiplicity at most  $n$  that refines  $\mathcal{V}$ . Let  $\mathcal{F} = \{W \times [0, 1] : W \in \mathcal{W}\}$  be a cover of  $X \times [0, 1]$ . Note that  $\mathcal{F}$  is a closed, locally finite, locally finite-dimensional cover of  $X \times [0, 1]$ .

Let  $H_0: K \times \{0, 1\} \rightarrow Y$  be the map defined by  $H_0(x, 0) = f(x)$  and  $H_0(x, 1) = g(x)$ .

Let  $C: \mathcal{V} \rightarrow \mathcal{U}$  be the map defined by  $C(V_U) = U$ . It is a carrier and both  $f$  and  $g$  are carried by  $C$ , directly from the definition of  $\mathcal{V}$ . Let  $D: \mathcal{W} \rightarrow \mathcal{V}$  be a map such that for each  $W \in \mathcal{W}$  we have  $W \subset D(W)$ . The map  $D$  exists because  $\mathcal{W}$  refines  $\mathcal{V}$ . Let  $E: \mathcal{F} \rightarrow \mathcal{W}$  be a map defined by  $D(W \times [0, 1]) = W$ . The composition  $C \circ D \circ E$  is a carrier and  $H_0$  is carried by it. Observe that  $\mathcal{U}$  is an  $AE(X \times [0, 1])$ -cover of  $Y$  (see Definition 2.5). By the Carrier Theorem,  $H_0$  can be extended to a map  $H: X \times [0, 1]$  that is carried by  $C \circ D \circ E$ . Clearly,  $H$  is a homotopy between  $f$  and  $g$ . For each path  $\{x\} \times [0, 1] \subset X \times [0, 1]$  there exists element  $W \times [0, 1] \in \mathcal{F}$  such that  $\{x\} \times [0, 1] \subset W \times [0, 1]$ , as  $\mathcal{W}$  is a cover of  $X$ . Since  $H$  is carried into  $\mathcal{U}$ , the whole path lies in an element of  $\mathcal{U}$ . Therefore  $H$  is a  $\mathcal{U}$ -homotopy.  $\square$

**Theorem 3.5.** *Let  $\mathcal{K} = \{K_i\}_{i \in I}$  be a cover of a polyhedron  $K$  by subcomplexes. Let  $\mathcal{L} = \{L_i\}_{i \in I}$  be a cover of a polyhedron  $L$  by subcomplexes. Let  $p: K \rightarrow L$  be a surjective simplicial map that maps elements of  $\mathcal{K}$  into the corresponding elements of  $\mathcal{L}$  (i.e.  $p(K_i) \subset L_i$  for each  $i \in I$ ). If  $\mathcal{K}$  and  $\mathcal{L}$  are isomorphic  $AE(n)$ -covers, then  $p$  is a weak  $n$ -homotopy equivalence.*

*Proof.* To begin, we will show that  $p$  induces monomorphisms on homotopy groups of dimensions less than  $n$ . Let  $m < n$ . Let  $\varphi: S^m \rightarrow K$ . Assume that

$p \circ \varphi$  is null-homotopic in  $L$ . Let  $\Phi: B^{m+1} \rightarrow L$  denote such a null-homotopy ( $\Phi|_{S^m} = p \circ \varphi$ ). We have to show that  $\varphi$  is null-homotopic in  $K$ .

Let  $\mathcal{F}$  be a finite closed cover of  $B^{m+1}$  with mesh small enough so that for each  $F \in \mathcal{F}$  we can pick  $i_F \in I$  such that

$$(*) \quad \varphi(F \cap S^m) \subset \text{ost}_{\beta K} K_{i_F} \text{ and } (**) \quad \Phi(F) \subset \text{ost}_{\beta L} L_{i_F}.$$

By  $(**)$ , a map  $C: \mathcal{F} \rightarrow \text{ost}_{\beta L} \mathcal{L}$  defined by  $C(F) = \text{ost}_{\beta L} L_{i_F}$  is a carrier. Since  $\mathcal{K}$  and  $\mathcal{L}$  are isomorphic, the map  $C': \mathcal{F} \rightarrow \text{ost}_{\beta K} \mathcal{K}$  defined by  $C'(F) = \text{ost}_{\beta K} K_{i_F}$  is a carrier as well. By  $(*)$ ,  $\varphi$  is carried by  $C'$ . By Corollary 3.3,  $\text{ost}_{\beta K} \mathcal{K}$  is an  $AE(n)$ -cover. By the Carrier Theorem,  $\varphi$  extends to a map  $\tilde{\varphi}: B^{m+1} \rightarrow K$  that is carried by  $C'$ . This is a null-homotopy of  $\varphi$  in  $K$  and we are done with the proof that  $p$  induces monomorphisms on homotopy groups of dimensions less than  $n$ .

Next we will show that  $p$  induces epimorphisms on homotopy groups of dimensions less than  $n$ . Let  $m < n$ . Let  $\psi: S^m \rightarrow L$ . Let  $\mathcal{G}$  be a closed finite cover of  $S^m$  that refines  $\psi^{-1}(\text{ost}_{\beta L} \mathcal{L})$ . For each  $G \in \mathcal{G}$  pick  $i_G \in I$  such that  $\psi(G) \subset \text{ost}_{\beta L} L_{i_G}$ . Then the map  $D: \mathcal{G} \rightarrow \text{ost}_{\beta L} \mathcal{L}$  defined by  $D(G) = \text{ost}_{\beta L} L_{i_G}$  is a carrier and  $\psi$  is carried by  $D$ . Since  $\mathcal{K}$  and  $\mathcal{L}$  are isomorphic, the map  $D'(G) = \text{ost}_{\beta K} K_{i_G}$  is a carrier as well. By Corollary 3.3,  $\text{ost}_{\beta K} \mathcal{K}$  is an  $AE(n)$ -cover. By the Carrier Theorem, there exists a map  $\tilde{\psi}: S^m \rightarrow K$  that is carried by  $D'$ . Observe that if  $x \in G \in \mathcal{G}$ , then  $\tilde{\psi}(x) \in \text{ost}_{\beta K} K_{i_G}$  and  $\psi(x) \in \text{ost}_{\beta L} L_{i_G}$ . Since  $p(K_{i_G}) \subset L_{i_G}$ , the maps  $\psi$  and  $p \circ \tilde{\psi}$  are  $\text{ost}_{\beta L} \mathcal{L}$ -close. By Corollary 3.3,  $\text{ost}_{\beta L} \mathcal{L}$  is an  $AE(n)$ -cover and by Lemma 3.4,  $\psi$  and  $p \circ \tilde{\psi}$  are homotopic. Hence  $p$  induces epimorphisms on homotopy groups of dimensions less than  $n$ . This concludes the second part of the proof.  $\square$

### 3.2. Proof of Theorem 3.1.

**Lemma 3.6.** *Let  $K$  and  $L$  be polyhedra. If  $p: K \rightarrow L$  is a simplicial, surjective map such that for each  $\delta \in \tau(L)$  the inverse image  $p^{-1}(\delta)$  is an  $AE(n)$ , then  $p$  is a weak  $n$ -homotopy equivalence.*

*Proof.* Apply Theorem 3.5 with the cover  $\mathcal{L} = \tau(L)$  of  $L$  and the cover  $\mathcal{K} = p^{-1}(\mathcal{L})$  of  $K$ .  $\square$

*Proof.* (Proof of Theorem 3.1) The map  $p$  is simplicial onto  $\beta K$  and satisfies the conditions of Lemma 3.6, hence it is a weak  $n$ -homotopy equivalence.

Let  $\mathcal{A} \subset \mathcal{B}_L$  such that  $\bigcap_{A \in \mathcal{A}} A \neq \emptyset$ . By Lemma 2.23,  $\bigcap_{A \in \mathcal{A}} A$  is an absolute extensor in dimension  $n$ . We have  $B = \bigcap_{A \in \mathcal{A}} p^{-1}(A) = p^{-1}(\bigcap_{A \in \mathcal{A}} A)$ . The map  $p|_B: B \rightarrow \bigcap_{A \in \mathcal{A}} A$  satisfies the conditions of Lemma 3.6, hence it is a weak  $n$ -homotopy equivalence. Therefore  $B$  has vanishing homotopy

groups in dimensions less than  $n$  and by Theorem 2.6, it is an absolute extensor in dimension  $n$ . Hence  $p^{-1}(\mathcal{B}_L)$  is an  $AE(n)$ -cover.  $\square$

#### 4. A LIFTING CONDITION

**Lemma 4.1.** *Let  $p: Y \rightarrow Z$  be a surjective map of metric spaces. Let  $\mathcal{F}$  be a cover of  $Z$ . Assume that either  $\mathcal{F}$  is an open cover or  $\mathcal{F}$  is a closed, locally finite, locally finite-dimensional cover. Assume that  $p^{-1}(\mathcal{F})$  is an  $AE(n)$ -cover.*

*Let  $X$  be an at most  $n$ -dimensional metric space. Let  $f: X \rightarrow Z$ . Let  $A$  be a closed subset of  $X$ . Let  $g_0: A \rightarrow Y$  be a map such that  $p \circ g_0 = f|_A$  (a lift of  $f$  on  $A$ ).*

*Then there exists a map  $g: X \rightarrow Y$  that satisfies the following conditions:*

- (1)  $g|_A = g_0$  ( $g$  extends  $g_0$ ); and
- (2)  $p \circ g$  is  $\mathcal{F}$ -close to  $f$ .

*Proof.* We have the following commutative diagram.

$$\begin{array}{ccc} Y & \xrightarrow{p} & Z \\ \uparrow g_0 & \nearrow g & \uparrow f \\ A & \hookrightarrow & X \end{array}$$

Let  $\mathcal{P} = p^{-1}(\mathcal{F})$  and  $\mathcal{G} = f^{-1}(\mathcal{F})$ . Let  $C: \mathcal{G} \rightarrow \mathcal{P}$  be defined by the formula  $C(f^{-1}(F)) = p^{-1}(F)$ . Since  $p$  is surjective,  $C$  is a carrier. For each  $F \in \mathcal{F}$  and each  $x \in f^{-1}(F) \cap A$  we have  $g_0(x) \in p^{-1}(f(x)) \in p^{-1}(F)$ ; hence  $g_0$  is carried by  $C$ .

Let  $\mathcal{H}$  be a closed locally finite locally finite-dimensional cover of  $X$  such that  $\mathcal{H}$  refines  $\mathcal{G}$ . If  $\mathcal{F}$  is an open cover, then  $\mathcal{H}$  exists as a refinement of an open cover  $\mathcal{G}$ . If  $\mathcal{F}$  is closed locally finite locally finite-dimensional, then we can take  $\mathcal{H} = \mathcal{G}$ , which has the required properties by Lemma 2.28. Let  $D: \mathcal{H} \rightarrow \mathcal{G}$  be any map such that  $H \subset D(H)$  for each  $H \in \mathcal{H}$ .

Observe that  $C \circ D: \mathcal{H} \rightarrow \mathcal{P}$  is a carrier,  $g_0$  is carried by  $C \circ D$ ,  $\mathcal{P}$  is an  $AE(n)$ -cover,  $X$  is at most  $n$ -dimensional metric space and  $\mathcal{H}$  is closed, locally finite and locally finite-dimensional. By the Carrier Theorem, there exists a map  $g: X \rightarrow Y$  that extends  $g_0$  and that is carried by  $C \circ D$ . This implies that for each  $x \in X$  there exists  $F \in \mathcal{F}$  such that  $x \in f^{-1}(F)$  and  $g(x) \in p^{-1}(F)$ . Therefore  $p(g(x)) \in F$  so  $p \circ g$  is  $\mathcal{F}$ -close to  $f$ .  $\square$

##### 4.1. A lifting condition for inverse limits.

**Definition 4.2.** Let

$$Z = \varprojlim \left( Z_1 \xleftarrow{p_1} Z_2 \xleftarrow{p_2} \cdots \right).$$

Let  $\mathcal{F}_i$  be a cover of  $Z_i$ . We define the following conditions.

- (A) For each  $i$ ,  $Z_i$  is a complete metric space.
- (B) For each  $i$ , the bonding map  $p_i: Z_{i+1} \rightarrow Z_i$  is surjective and 1-Lipschitz.
- (C) For each  $i$ ,  $\mathcal{F}_i$  is either an open cover or a closed, locally finite, locally finite-dimensional cover.
- (D) For each  $i$ , the pull-back  $p_i^{-1}(\mathcal{F}_i)$  is an  $AE(n)$ -cover.
- (E)  $\sum_i \text{mesh } \mathcal{F}_i < \infty$ .

We denote short projections in the inverse limit by  $\pi_i^k: Z_k \rightarrow Z_i (k > i)$  and long projections in the inverse limit by  $\pi_k: Z \rightarrow Z_k$ .

**Lemma 4.3.** *Let*

$$Z = \varprojlim \left( Z_1 \xleftarrow{p_1} Z_2 \xleftarrow{p_2} \cdots \right).$$

*Let  $\mathcal{F}_i$  be a cover of  $Z_i$ . Assume that conditions of Definition 4.2 are satisfied.*

*Let  $X$  be an at most  $n$ -dimensional metric space. Let  $f: X \rightarrow Z_1$ . Let  $A$  be a closed subset of  $X$ . Let  $g_0: A \rightarrow Z$  be a map such that  $\pi_1 \circ g_0 = f|_A$  (a lift of  $f$  on  $A$ ).*

*Then there exists a map  $g: X \rightarrow Z$  such that  $g|_A = g_0$  ( $g$  extends  $g_0$ ).*

*Proof.* Let  $f_1 = f$ . Applying Lemma 4.1 for each  $i > 1$  we construct a map  $f_i: X \rightarrow Z_i$  that satisfies the following conditions.

- (i)  $p_{i-1} \circ f_i$  is  $\mathcal{F}_{i-1}$ -close to  $f_{i-1}$ .
- (ii)  $\pi_i \circ g_0 = f_i|_A$ .

For each  $k$  and each  $m > k$  we let

$$a_m^k = \pi_k^m \circ f_m: X \rightarrow Z_k.$$

By (i), we have  $d_{\text{sup}}(f_m, p_m \circ f_{m+1}) \leq \text{mesh } \mathcal{F}_m$ . By (B) the short projection  $\pi_k^m$  is 1-Lipschitz, hence  $d_{\text{sup}}(a_m^k, a_{m+1}^k) = d_{\text{sup}}(\pi_k^m \circ f_m, \pi_k^m \circ p_m \circ f_{m+1}) \leq \text{mesh } \mathcal{F}_m$ . Therefore for  $l > m$  we have

$$\begin{aligned} d_{\text{sup}}(a_l^k, a_m^k) &\leq d_{\text{sup}}(a_l^k, a_{l-1}^k) + d_{\text{sup}}(a_{l-1}^k, a_{l-2}^k) + \cdots + d_{\text{sup}}(a_{m+1}^k, a_m^k) \leq \\ &\leq \text{mesh } \mathcal{F}_{l-1} + \text{mesh } \mathcal{F}_{l-2} + \cdots + \text{mesh } \mathcal{F}_{m+1} + \text{mesh } \mathcal{F}_m. \end{aligned}$$

By (E), the sequence  $a_m^k$  is uniformly convergent. Let

$$a^k = \lim_{m \rightarrow \infty} a_m^k.$$

By (A),  $Z_k$  is complete, hence  $a^k: X \rightarrow Z_k$  is well defined.

It follows from the definition that  $p_k \circ a_m^{k+1} = a_m^k$ . Passing to the limit we have

$$p_k \circ a^{k+1} = \lim_{m \rightarrow \infty} p_k \circ a_m^{k+1} = \lim_{m \rightarrow \infty} a_m^k = a^k.$$

Therefore, for each  $x$  the sequence  $(a^k(x))_k$  is a thread in  $Z$  and we can define a map  $g: X \rightarrow Z$  by the formula

$$(g(x))_k = a^k(x).$$

Observe that by (ii) and by the definition of  $a_m^k$  we have

$$a_m^k|_A = \pi_k^m \circ f_m|_A = \pi_k^m \circ \pi_m \circ g_0|_A = \pi_k \circ g_0|_A.$$

Therefore  $a^k|_A = \pi_k \circ g_0$  for each  $k$ , hence  $g|_A = g_0|_A$ .  $\square$

**Theorem 4.4.** *Let*

$$Z = \varprojlim \left( Z_1 \xleftarrow{p_1} Z_2 \xleftarrow{p_2} \cdots \right).$$

*Let  $\mathcal{F}_i$  be a cover of  $Z_i$ . If the conditions of Definition 4.2 are satisfied and  $Z_1$  is an  $ANE(n)$ , then  $Z$  is an  $ANE(n)$ .*

*Proof.* We will show that  $Z$  is an  $ANE(n)$  directly from the definition. Let  $X$  be an at most  $n$ -dimensional metric space and let  $A \subset X$  be a closed subset. Take  $g_0: A \rightarrow Z$ . By assumption,  $Z_1$  is an  $ANE(n)$  so we can extend  $\pi_1 \circ g_0: A \rightarrow Z_1$  to a map  $f_1: U \rightarrow Z_1$ , where  $U$  is an open neighborhood of  $A$  in  $X$ . By Lemma 4.3,  $f_1$  can be lifted to a map  $g: U \rightarrow Z$  such that  $g|_A = g_0$ . This is the extension of  $f$  we sought.  $\square$

## 5. LOCAL $k$ -CONNECTEDNESS OF INVERSE LIMITS OF POLYHEDRA

**Definition 5.1.** Let

$$(L) \quad X = \varprojlim \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \cdots \right)$$

Fix  $m$  and let  $A \subset K_m$  be a subcomplex of  $K_m$ . A **restriction of (L) to  $A$**  is the inverse limit

$$(R) \quad X' = \varprojlim \left( K'_m \xleftarrow{p'_m} K'_{m+1} \xleftarrow{p'_{m+1}} \cdots \right),$$

where  $K'_m = A$  and for each  $j \geq m$ ,  $K'_{j+1} = p_j^{-1}(K'_j)$  and  $p'_j = p_j|_{K'_{j+1}}$ .

**Lemma 5.2.** *Let (R) be the restriction of (L) to a subcomplex  $A \subset K_m$ .*

*If, for each  $i$ , (L) satisfies the conditions:*

*(I)  $K_i$  is a polyhedron and*

*(II)  $p_i$  is a quasi-simplicial map that is surjective and  $n$ -regular,*

*then so does (R). The inverse limit  $X'$  is homeomorphic to  $\pi_m^{-1}(A)$ , where  $\pi_m: X \rightarrow X_m$  denotes the long projection.*

*Proof.* We have

$$\pi_m^{-1}(A) = \lim_{\leftarrow} \left( K'_1 \xleftarrow{p'_1} K'_2 \xleftarrow{p'_2} \dots \right),$$

where  $K'_j = p_j(K'_{j+1})$  and  $p'_j = p_j|_{K'_{j+1}}$  for  $j < m$ . The restriction (R) is the same sequence with first  $m - 1$  elements removed. This changes the metric on the limit, but not the topology, hence  $X'$  is homeomorphic to  $\pi_m^{-1}(A)$ .

The other conditions are trivial to verify.  $\square$

**Theorem 5.3.** *Let*

$$X = \lim_{\leftarrow} \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \dots \right).$$

*Assume that for each  $i$  the following conditions are satisfied:*

- (I)  $K_i$  is a locally finite-dimensional polyhedron; and
- (II)  $p_i$  is a quasi-simplicial map that is surjective and  $n$ -regular.

*Then*

- (1)  $X$  is an  $ANE(n)$ ;
- (2) each short projection  $\pi_i^k: K_k \rightarrow K_i$  and each long projection  $\pi_i: X \rightarrow K_i$  is a weak  $n$ -homotopy equivalence;
- (3) for each  $i$ , the covers  $\mathcal{O}_i$  and  $\mathcal{B}_i$  are  $AE(n)$ -covers of  $X$ .

*Proof.* Fix a metric of scale  $2^{-i}$  on  $K_i$  (see Definition 2.9). Let  $(*)$  denote the inverse limit

$$X = \lim_{\leftarrow} \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \dots \right)$$

along with a sequence  $\mathcal{B}_{K_i}$  of covers, where  $\mathcal{B}_{K_i}$  is a cover of  $K_i$  by barycentric stars of vertices.

We will verify that  $(*)$  satisfies conditions (A)–(E) of Definition 4.2.

Condition (A): By Lemma 2.12, every  $K_i$  is a complete metric space.

Condition (B): By condition (II),  $p_i$  is surjective. By the choice of scale on  $K_i$  and Lemma 2.14, each bonding map is 1-Lipschitz.

Condition (C): By Lemma 2.23, each  $\mathcal{B}_{K_i}$  is a closed, locally finite, locally finite-dimensional cover.

Condition (D): By assumption  $p_i$  is quasi-simplicial and  $n$ -regular. Hence by Theorem 3.1, the pull-back  $p_i^{-1}(\mathcal{B}_{K_i})$  is an  $AE(n)$ -cover.

Condition (E): By the choice of the metric on  $K_i$ , we have  $\sum_i \text{mesh } \mathcal{B}_{K_i} \leq \sum_i \text{diam } K_i < \infty$ .

To prove (1) it is enough to verify the conditions of Theorem 4.4 for  $(*)$ . We just verified conditions of Definition 4.2. By Lemma 2.12,  $K_1$  is an  $ANE(n)$ . We are done.

By Theorem 3.1 each  $p_i$  is a weak  $n$ -homotopy equivalence and therefore all the short projections  $\pi_i^k$  are weak  $n$ -homotopy equivalences. To finish the proof of (2), we must show that

- (mono) for fixed  $i > 0$  and  $m < n$ , the long projection  $\pi_i : X \rightarrow K_i$  induces a monomorphism on the homotopy group of dimension  $m$ , regardless of the choice of base point; and
- (epi) for fixed  $i > 0$  and  $m < n$ , the long projection  $\pi_i : X \rightarrow K_i$  induces an epimorphism on the homotopy group of dimension  $m$ , regardless of the choice of base point.

We prove (mono). Fix  $i > 0$  and  $m < n$ . Let  $\varphi : S^m \rightarrow X$ . Assume that  $\pi_i \circ \varphi$  is null-homotopic in  $K_i$ . Let  $\tilde{\Phi} : B^{m+1} \rightarrow K_i$  denote the null-homotopy ( $B^{m+1}$  denotes the  $m + 1$ -dimensional unit ball in  $\mathbb{R}^{m+1}$ ). We have the following commutative diagram:

$$\begin{array}{ccc} X & \xrightarrow{\pi_i} & K_i \\ \uparrow \varphi & \nwarrow \tilde{\Phi} & \uparrow \Phi \\ S^m & \hookrightarrow & B^{m+1} \end{array}$$

The inverse limit

$$X = \varprojlim \left( K_i \xleftarrow{p_i} K_{i+1} \xleftarrow{p_{i+1}} \dots \right)$$

along with the sequence of covers by barycentric stars of vertices (which we obtain by truncating  $(*)$ ) satisfies the conditions of Definition 4.2. Since  $B^{m+1}$  is at most  $n$ -dimensional, by Lemma 4.3, there exists a lift  $\tilde{\Phi}$  such that the diagram is commutative. This lift is a null-homotopy of  $\varphi$  in  $X$ , hence  $\pi_i$  induces a monomorphism on the homotopy group of dimension  $m$ . We will show (epi) at the end of the proof.

Next we prove (3). To begin, we will show that for each  $i$ ,  $\mathcal{B}_i$  is an  $AE(n)$ -cover. Let  $\mathcal{A}$  be a collection of elements of  $\mathcal{B}_i$  such that the intersection  $\bigcap \mathcal{A}$  is non-empty. By the definition of  $\mathcal{B}_i$ , we have  $\mathcal{A} = \{\pi_i^{-1}(\text{bst}_{K_i} v)\}_{v \in V}$  for some set of vertices  $V$  of  $K_i$ . Let  $A = \bigcap_{v \in V} p_i^{-1}(\text{bst}_{K_i} v)$ . Since  $p_i$  is quasi-simplicial,  $A$  is a subcomplex of  $K_{i+1}$ . By Theorem 3.1,  $A$  has vanishing homotopy groups in dimensions less than  $n$ . Let

$$(**) \quad X' = \varprojlim \left( K'_{i+1} \xleftarrow{p'_{i+1}} K'_{i+2} \xleftarrow{p'_{i+2}} \dots \right),$$

where  $K'_{i+1} = A$  and for each  $j \geq i + 1$ ,  $K'_{j+1} = p_j^{-1}(K'_j)$  and  $p'_j = p_j|_{K'_{j+1}}$ , be the restriction of  $(*)$  to  $A$ . By Lemma 5.2 it satisfies assumptions (I) and (II) and is homeomorphic to  $\bigcap \mathcal{A}$ . Hence from what we have already proven,  $X' = \bigcap \mathcal{A}$  is an  $ANE(n)$  and the long projection  $\pi_{i+1} : \bigcap \mathcal{A} \rightarrow A$

induces monomorphisms on homotopy groups of dimensions less than  $n$ . Since  $A$  has vanishing homotopy groups in these dimensions, so does  $\bigcap \mathcal{A}$ . By Theorem 2.6,  $\bigcap \mathcal{A}$  is an  $AE(n)$  hence  $\mathcal{B}_i$  is an  $AE(n)$ -cover.

It follows from Lemma 2.7 that  $\mathcal{O}_i$  is an  $AE(n)$ -cover, as open stars are (infinite) unions of iterated barycentric stars.

Finally, we prove (epi). Fix  $i > 0$  and  $m < n$ . Let  $\varphi: S^m \rightarrow K_i$ . Let  $\mathcal{O}$  be a cover of  $K_i$  by open stars of vertices. We have just shown that  $\pi_i^{-1}(\mathcal{O})$  is an  $AE(n)$ -cover. Hence by Lemma 4.1, there exists a map  $\psi: S^m \rightarrow X$  such that  $\pi_i \circ \psi$  and  $\varphi$  are  $\mathcal{O}$ -close. By Lemma 3.4, these maps are homotopic. This shows that  $\pi_i$  induces epimorphisms on homotopy groups of dimensions less than  $n$ . Hence  $\pi_i$  is a weak  $n$ -homotopy equivalence.  $\square$

**Corollary 5.4.** *Let*

$$X = \varprojlim \left( K_1 \xleftarrow{p_1} K_2 \xleftarrow{p_2} \cdots \right).$$

*Assume that for each  $i$ , (I)  $K_i$  is a locally finite-dimensional polyhedron; (II)  $p_i$  is a quasi-simplicial map that is surjective and  $n$ -regular. Let  $Y$  be a metric space of dimension less than  $n$  and let  $f, g: Y \rightarrow X$ . Then if  $f$  is  $\mathcal{O}_i$ -close to  $g$ , then  $f$  is  $\mathcal{O}_i$ -homotopic to  $g$ .*

*Proof.* By Theorem 5.3(3),  $\mathcal{O}_i$  is an  $AE(n)$ -cover of  $X$ . By Lemma 3.4,  $f$  and  $g$  are  $\mathcal{O}_i$ -homotopic.  $\square$

## REFERENCES

- [1] S. M. Ageev. Axiomatic method of partitions in the theory of Nöbeling spaces. II. An unknotting theorem. *Mat. Sb.*, 198(5):3–32, 2007.
- [2] S. M. Ageev. Axiomatic method of partitions in the theory of Nöbeling spaces. III. Consistency of the system of axioms. *Mat. Sb.*, 198(7):3–30, 2007.
- [3] S. M. Ageev. The axiomatic partition method in the theory of Nöbeling spaces. I. Improving partition connectivity. *Mat. Sb.*, 198(3):3–50, 2007.
- [4] G. C. Bell and A. Nagórko. A new construction of universal spaces for asymptotic dimension. *Topology Appl.*, 160(1):159–169, 2013.
- [5] A. C. Chigogidze.  $n$ -soft mappings of  $n$ -dimensional spaces. *Mat. Zametki*, 46(1):88–95, 124, 1989.
- [6] J. Dugundji. Absolute neighborhood retracts and local connectedness in arbitrary metric spaces. *Compositio Math.*, 13:229–246 (1958), 1958.
- [7] D. Gabai. On the topology of ending lamination space. *Geom. Topol.*, 18(5):2683–2745, 2014.
- [8] S. Hensel and P. Przytycki. The ending lamination space of the five-punctured sphere is the Nöbeling curve. *J. Lond. Math. Soc. (2)*, 84(1):103–119, 2011.
- [9] S.-t. Hu. *Theory of retracts*. Wayne State University Press, Detroit, 1965.
- [10] M. Levin. Characterizing Nöbeling spaces, 2006. Available online from <http://www.arxiv.org/pdf/math.GT/0602361>.
- [11] A. T. Lundell and S. Weingram. *The Topology of CW Complexes*. The University Series in Higher Mathematics. Van Nostrand Reinhold Company, 1969.
- [12] A. Nagórko. *Characterization and topological rigidity of Nöbeling manifolds*. PhD thesis, University of Warsaw, 2006.

- [13] A. Nagórko. Carrier and nerve theorems in the extension theory. *Proc. Amer. Math. Soc.*, 135(2):551–558, 2007. <http://arxiv.org/abs/math.GN/0506525>.
- [14] A. Nagórko. Characterization and topological rigidity of Nöbeling manifolds. *Mem. Amer. Math. Soc.*, 223(1048):viii+92, 2013.
- [15] K. Sakai. *Geometric aspects of general topology*. Springer Monographs in Mathematics. Springer, Tokyo, 2013.

(G. C. Bell) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NORTH CAROLINA AT GREENSBORO, GREENSBORO, NC 27412, USA

*E-mail address:* `gcbell@uncg.edu`

(A. Nagórko) FACULTY OF MATHEMATICS, INFORMATICS, AND MECHANICS, UNIVERSITY OF WARSAW, BANACHA 2, 02-097 WARSZAWA, POLAND

*E-mail address:* `amn@mimuw.edu.pl`