

INEQUALITIES OF CHERN CLASSES ON NONSINGULAR PROJECTIVE n -FOLDS OF FANO AND GENERAL TYPE WITH AMPLE CANONICAL BUNDLE

RONG DU[†] AND HAO SUN^{††}

ABSTRACT. Let X be a nonsingular projective n -fold ($n \geq 2$) of Fano or of general type with ample canonical bundle K_X over an algebraic closed field κ of any characteristic. We produce a new method to give a bunch of inequalities in terms of all the Chern classes c_1, c_2, \dots, c_n by pulling back Schubert classes in the Chow group of Grassmannian under the Gauss map. Moreover, we show that if the characteristic of κ is 0, then the Chern ratios $(\frac{c_{2,1n-2}}{c_1^n}, \frac{c_{2,2,1n-4}}{c_1^n}, \dots, \frac{c_n}{c_1^n})$ are contained in a convex polyhedron for all X . So we give an affirmative answer to a generalized open question, that whether the region described by the Chern ratios is bounded, posted by Hunt ([Hun]) to all dimensions. As a corollary, we can get that there exist constants d_1, d_2, d_3 and d_4 depending only on n such that $d_1 K_X^n \leq \chi_{top}(X) \leq d_2 K_X^n$ and $d_3 K_X^n \leq \chi(X, \mathcal{O}_X) \leq d_4 K_X^n$. If the characteristic of κ is positive, K_X (or $-K_X$) is ample and $\mathcal{O}_X(K_X)$ ($\mathcal{O}_X(-K_X)$, respectively) is globally generated, then the same results hold.

1. INTRODUCTION

One of the fundamental questions in algebraic geometry is the classification of algebraic varieties. The classical method is by considering the numerical invariants first. Among all kinds of numerical invariants, Chern numbers are the most natural and important ones. The terminology geography which was first introduced by Persson in 1981 (see [Per]) is used to describe the distribution of Chern numbers of nonsingular projective varieties of general type, i.e. whether there exists nonsingular projective varieties X of dimension n for every given set of numbers such that X has precisely those Chern numbers. So the first thing is to determine if the quotients of

[†] The Research is Sponsored by the National Natural Science Foundation of China (Grant No. 11471116, 11531007) and Science and Technology Commission of Shanghai Municipality (Grant No. 18dz2271000).

^{††} The Research is Sponsored by the National Natural Science Foundation of China (Grant No. 11771294, 11301201).

those numbers are in a bounded set or not. For $n = 2$ and X is minimal over the complex numbers, we have Noether's inequalities:

$$K_X^2 \geq 2p_g - 4,$$

here $p_g = h^0(X, K_X)$. From Noether's formula, we can have

$$5c_1^2 + 36 \geq c_2.$$

On the other hand, we have the famous Bogomolov-Miyaoka-Yau inequality

$$c_2 \geq \frac{1}{3}c_1^2.$$

So c_2/c_1^2 is bounded.

For $n = 2$ and over a field of positive characteristic, Noether's inequality (see [Lie]) and Noether's formula (see [Bad] Chap. 5) remain true, while Lang ([Lan]) and Easton ([Eas]) gave examples of surfaces, such as generalized Raynaud surfaces, for which the Bogomolov-Miyaoka-Yau inequality no longer holds (cf. [Szp], Sec. 3.4). In fact, from Raynaud's example ([Ray]) even the weaker inequality $c_2 \geq 0$ due to Castelnuovo and de Franchis fails (cf. [Gu], Sec. 3). So it is natural to formulate an inequality in positive characteristic bounding c_2 from below by c_1^2 . Shepherd-Barron has already considered a similar question and proved that $\chi > 0$ (equivalently, $c_2 > -c_1^2$) with a few possible exceptional cases when the characteristic of the field is not greater than 7 ([S-B], Theorem 8). Later, Gu solved Shepherd-Barron's question and got an optimal lower bound of χ/c_1^2 .

For $n = 3$ and over the complex numbers, Hunt initiated the study of geography for threefolds ([Hun]). Later, Chang, Kim and Nollet gave a bound of c_3 by quadratic forms in term of c_1c_2 and c_1^3 of threefolds with ample canonical bundle ([C-K-N]). Later, Chang and Lopez obtained a linear bound of c_3 of threefolds with ample canonical bundle, i.e. boundedness for the region described by the Chern ratios c_3/c_1c_2 , c_1^3/c_1c_2 ([C-L]). Their idea is to control the Euler number by the Riemann-Roch formula and cohomology calculations. Their result relies on the Bogomolov-Miyaoka-Yau inequality heavily for dimension 3. M. Chen-Hacon and J. Chen-Hacon (see [CM-H], [CJ-H]) also considered the geography of Gorenstein minimal complex 3-folds of general type after 2000. For positive characteristic, as far as the authors know that there are no such kind of considerations.

For $n \geq 4$, such problem seems unknown even over the field of complex numbers. Except for the famous Bogomolov-Miyaoka-Yau inequality which is in the case of characteristic 0, many other mathematicians studied inequalities of Chern classes (see [F-L], [B-S-S], [Ca-Sc], [Ch-Le], [Ko], [Mi], [Sun]). In this paper, we show that there are similar boundedness result as the dimension 3 for characteristic 0 if K_X (or $-K_X$) is ample

and for positive characteristic if moreover $\mathcal{O}_X(K_X)$ ($\mathcal{O}_X(-K_X)$, respectively) is globally generated. So we give an affirmative answer to a generalized open question, fore that whether the region described by the Chern ratios is bounded, posted by Hunt ([Hun]) to all dimensions. In particular, we generalize Chang-Lopez's result to all dimensions. (For the notations, please see the paragraph before Theorem 3.7.)

Main Theorem:(see Theorem 3.7)

Let X be a nonsingular projective variety of dimension n over an algebraic closed field κ with any characteristic. Suppose K_X (or $-K_X$) is ample. If the characteristic of κ is 0 or the characteristic of κ is positive and $\mathcal{O}_X(K_X)$ ($\mathcal{O}_X(-K_X)$, respectively) is globally generated, then

$$\left(\frac{c_{2,1^{n-2}}}{c_1^n}, \frac{c_{2,2,1^{n-4}}}{c_1^n}, \dots, \frac{c_n}{c_1^n} \right) \in \mathbb{A}^{p(n)} \quad (1.1)$$

is contained in a convex polyhedron in $\mathbb{A}^{p(n)}$ independent of X , where $p(n)$ is the partition number and the elements in the parentheses arranged from small to big in terms of the alphabet order of the lower indices of the numerators.

In particular, we show that the Euler number $\chi_{top}(X)$ and Euler characteristic of the structure sheaf $\chi(X, \mathcal{O}_X)$ can be controlled linearly by K_X^n . More precisely, there exist constants d_1, d_2, d_3 and d_4 , which do not depend on X but only on n , such that

$$d_1 K_X^n \leq \chi_{top}(X) \leq d_2 K_X^n$$

and

$$d_3 K_X^n \leq \chi(X, \mathcal{O}_X) \leq d_4 K_X^n.$$

Our results can also infer the classical boundedness result for dimension 2 and Chang-Lopez's result for dimension 3 easily without using the Bogomolov-Miyaoka-Yau inequality. Furthermore, we can deduce a bunch of inequalities of Chern classes not only Chern numbers.

Remark 1.1. Fulton-Lazarsfeld, Demailly-Peternell-Schneider, and Catanese-Schneider's results can give many inequalities of Chern classes. Moreover, combining their results and Bogomolov-Miyaoka-Yau inequalities (for characteristic 0), one gets immediately bounds for $\frac{c_{2,1^{n-2}}}{c_1^n}$, which can also start our induction in the proof of the main theorem (see Theorem 3.7). However, their methods depend on the characteristic 0 and Bogomolov-Miyaoka-Yau inequality. Our method is totally different and doesn't rely on Bogomolov-Miyaoka-Yau inequality. We can deal with any characteristic of the algebraic closed field simultaneously.

In Section 2, we introduce Schubert cycles and Schubert classes of Grassmannian. Pieri's formula and Giambelli's formula are also mentioned in this section for Schubert calculus later. In Section 3, we recall Fujita conjecture and known relevant very ampleness results in any characteristic first. Then we solve our main theorem by estimating Chern classes from Schubert calculus and the help of Zak's theorem. In Section 4, we produce a new algorithm to give a bunch of inequalities in terms of all the Chern classes c_1, c_2, \dots, c_n by pulling back the Schubert classes in the Chow group of the Grassmannian under the Gauss map.

2. SCHUBERT CYCLES AND SCHUBERT CLASSES OF GRASSMANNIAN

We will recall the basic definition Schubert cycles and classes of the Chow group of $G(n, k)$, the Grassmannian of k -dimensional subspaces in an n -dimensional vector space V , and analyze their intersections, a subject that goes by the name of the *Schubert calculus*. Of course we may also consider $G(n, k)$ in its projective guise as $\mathbb{G}(n-1, k-1)$, the Grassmannian of projective $(k-1)$ -planes in \mathbb{P}^{n-1} . We recommend excellent books [G-H] and [E-H] for details.

Schubert cycles are defined in terms of a chosen complete flag \mathcal{V} in V , i.e., a nested sequence of subspaces

$$0 \subset V_1 \subset \dots \subset V_{n-1} \subset V_n = V$$

with $\dim V_i = i$. The Schubert cycles are indexed by sequences $\vec{a} = (a_1, a_2, \dots, a_k)$ of integers with

$$n - k \geq a_1 \geq a_2 \geq \dots \geq a_k \geq 0$$

We define $|\vec{a}| := \sum_{i=1}^k a_i$ and $l(\vec{a}) := k$.

For such a sequence \vec{a} , we define the *Schubert cycle* $\Sigma_{\vec{a}}(\mathcal{V}) \subset G(n, k)$ to be the closed subset

$$\Sigma_{\vec{a}}(\mathcal{V}) = \{\Lambda \in G(n, k) \mid \dim(V_{n-k+i-a_i} \cap \Lambda) \geq i \text{ for all } i\}.$$

We know that the class $[\Sigma_{\vec{a}}(\mathcal{V})] \in A(G(n, k))$ does not depend on the choice of the flag, since any two flags differ by an element of GL_n , where $A(G(n, k))$ is the Chow group of $G(n, k)$. So we shorten the notation to $\Sigma_{\vec{a}}$ and define *Schubert classes*

$$\sigma_{\vec{a}} := [\Sigma_{\vec{a}}] \in A(G(n, k)).$$

The following theorem shows that $A(G(n, k))$ is a free abelian group and that the classes $\sigma_{\vec{a}}$ form a basis.

Theorem 2.1. ([E-H] Corollary 4.7) The Schubert classes form a free basis for $A(G(n, k))$, and the intersection form

$$A^m(G(n, k)) \times A^{\dim G(n, k) - m}(G(n, k)) \rightarrow \mathbb{Z}$$

have the Schubert classes as dual bases.

To simplify notation, we generally suppress trailing zeros in the indices, writing σ_{a_1, \dots, a_s} in place of $\sigma_{(a_1, \dots, a_s, 0, \dots, 0)}$. Also, we use the shorthand σ_{p^r} to denote $\sigma_{p, \dots, p}$ with r indices equal to p .

Let $\mathcal{V} := G(n, k) \times V$ be the trivial vector bundle of rank n on $G(n, k)$ whose fiber at every point is the vector space V . We write S for the rank- k subbundle of \mathcal{V} whose fiber at a point $\Lambda \in G(n, k)$ is the subspace Λ itself; that is,

$$S_{[\Lambda]} = \Lambda \subset V = \mathcal{V}_{[\Lambda]}.$$

S is called the *universal subbundle* on $G(n, k)$; the quotient $Q = \mathcal{V}/S$ is called the *universal quotient bundle*, i.e.,

$$0 \rightarrow S \rightarrow k^n \rightarrow Q \rightarrow 0. \quad (2.2)$$

Proposition 2.2. (see [E-H] Sec. 5.6.2 Grassmannians or [G-H] Sec. 3.3)

$$c_p S = (-1)^p \sigma_{1^p}.$$

Next we will talk about the intersection of Schubert classes. One situation in which we can give a simple formula for the product of Schubert classes is when one of the classes has the special form σ_b with integer b . Such classes are called *special Schubert classes*.

Proposition 2.3. (Pieri's Formula) For any Schubert class $\sigma_{\vec{a}}$ and any integer b ,

$$\sigma_b \sigma_{\vec{a}} = \sum_{\substack{|\vec{c}| = |\vec{a}| + b \\ a_i \leq c_i \leq a_{i-1}, \forall i}} \sigma_{\vec{c}}.$$

Pieri's formula tells us how to intersect an arbitrary Schubert class with one of the special Schubert classes σ_b with integer b . Giambelli's formula is complementary, in that it tells us how to express an arbitrary Schubert class in terms of special ones; the two together give us a way of calculating the product of two arbitrary Schubert classes.

Proposition 2.4. (Giambelli's Formula)

$$\sigma_{a_1, a_2, \dots, a_q} = \begin{vmatrix} \sigma_{a_1} & \sigma_{a_1+1} & \sigma_{a_1+2} & \cdots & \sigma_{a_1+q-1} \\ \sigma_{a_2-1} & \sigma_{a_2} & \sigma_{a_2+1} & \cdots & \sigma_{a_2+q-2} \\ \sigma_{a_3-2} & \sigma_{a_3-1} & \sigma_{a_3} & \cdots & \sigma_{a_3+q-3} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \sigma_{a_q-q+1} & \sigma_{a_q-q+2} & \sigma_{a_q-q+3} & \cdots & \sigma_{a_q} \end{vmatrix} \quad (2.3)$$

3. A LINEAR BOUND ON THE CHERN RATIOS

Let X be a nonsingular projective variety of dimension n ($n \geq 2$) over an algebraic closed field κ with any characteristic. Suppose K_X or $-K_X$ is ample and mK_X is very ample (m can be negative if $-K_X$ is ample, i.e. X is Fano).

When X is a complex surface and L is an ample line bundle on X , Reider ([Rei]) showed that $K_X + 3L$ is always generated by global sections and $K_X + 4L$ very ample. Around the same period, Fujita ([Fuj]) raised the following interesting conjecture.

Fujita's Conjecture: Let X be a smooth n -dimensional complex projective algebraic variety and let L be an ample divisor on X .

- (1) For $t \geq n + 1$, $tL + K_X$ is base point free.
- (2) For $t \geq n + 2$, $tL + K_X$ is very ample.

For the very ampleness conjecture, one of the first results proved in dimension $n \geq 3$ is the very ampleness of $2K_X + 12n^nL$, using an analytic method based on the solution of a Monge-Ampère equation by Demailly (see [Dem1]). Other related works are ([Dem2], [E-L-N], [Siu1], [Siu2], [Siu3], [Yeu]) to improve the effective bound.

Since those proofs rely on the Kodaira Vanishing Theorem and its generalizations, it seems to us that we don't have such results for positive characteristic. However, Smith proved another version of the Fujita conjecture in arbitrary characteristic if L is ample and generated by global sections via tight closure theory (see [Smi1], [Smi2]). Later, Keeler used the method of positive characteristic to show another version of Fujita's Conjecture (see [Kee]).

Theorem 3.1. (see [Kee], Theorem 1.1) Let X be a projective scheme of pure dimension n , smooth over a field κ of arbitrary characteristic. Let L be an ample, globally generated line bundle and let H be an ample line bundle. Then

- (1) $K_X + nL + H$ is base point free.
- (2) $K_X + (n + 1)L + H$ is very ample.

We will use the results of Fujita's very ampleness conjecture to get a bound which depends only on the dimension n for the Chern ratios and will use Van de Ven's idea first (cf. [Hun] Introduction).

Assume $i : X \hookrightarrow \mathbb{P}^N$ is the canonical embedding defined by the linear system $|mK_X|$ (i.e. $mK_X = i^* \mathcal{O}_{\mathbb{P}^N}(1)$). Let γ be the Gauss map:

$$\begin{aligned} \gamma : X &\longrightarrow G(\mathbb{P}^N, \mathbb{P}^n) = G(N + 1, n + 1) \\ x &\longmapsto T_{X,x}, \end{aligned} \tag{3.4}$$

where $T_{X,x}$ is the tangent space to X at x . There is an usual bundle sequence on $G(N+1, n+1)$:

$$0 \rightarrow S \rightarrow \kappa^{N+1} \rightarrow Q \rightarrow 0,$$

where S is the universal bundle (see [G-H] Chapter I or [E-H] Chapter 3), which pulls back to an exact sequence on X

$$0 \rightarrow \gamma^* S \rightarrow \gamma^* \kappa^{N+1} \rightarrow \gamma^* Q \rightarrow 0. \quad (3.5)$$

On the other hand we have the Euler exact sequence on \mathbb{P}^N :

$$0 \rightarrow \mathcal{O}_{\mathbb{P}^N}(-1) \rightarrow \mathcal{O}_{\mathbb{P}^N}^{N+1} \rightarrow T_{\mathbb{P}^N}(-1) \rightarrow 0,$$

which pulls back to an exact sequence on X

$$0 \rightarrow i^* \mathcal{O}_{\mathbb{P}^N}(-1) \rightarrow i^* \mathcal{O}_{\mathbb{P}^N}^{N+1} \rightarrow i^* T_{\mathbb{P}^N}(-1) \rightarrow 0. \quad (3.6)$$

Moreover we have the twisted adjunction sequence on X :

$$0 \rightarrow T_X(-1) \rightarrow i^* T_{\mathbb{P}^N}(-1) \rightarrow N_{X/\mathbb{P}^N}(-1) \rightarrow 0. \quad (3.7)$$

These three sequences (3.5), (3.6) and (3.7) fit together in a diagram

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & \gamma^* S & \longrightarrow & T_X(-1) & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & i^* \mathcal{O}_{\mathbb{P}^N}(-1) & \longrightarrow & i^* \mathcal{O}_{\mathbb{P}^N}^{N+1} & \longrightarrow & i^* T_{\mathbb{P}^N}(-1) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & \gamma^* Q & & N_{X/\mathbb{P}^N}(-1) & & \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & & .
 \end{array}$$

By the snake lemma, we have the exact sequence

$$0 \rightarrow i^* \mathcal{O}_{\mathbb{P}^N}(-1) \rightarrow \gamma^* S \rightarrow T_X(-1) \rightarrow 0,$$

i.e.

$$0 \rightarrow \mathcal{O}_X(-mK_X) \rightarrow \gamma^* S \rightarrow T_X(-mK_X) \rightarrow 0.$$

So

$$c(\gamma^* S) = c(\mathcal{O}_X(-mK_X))c(T_X(-mK_X)). \quad (3.8)$$

Lemma 3.2. Let E be a vector bundle of rank r and L be a line bundle on a scheme X over an algebraic closed field k . Then for all $p \geq 0$,

$$c_p(E \otimes L) = \sum_{i=0}^p \binom{r-i}{p-i} c_i(E) c_1(L)^{p-i}. \quad (3.9)$$

From the above lemma, we have the following result.

Lemma 3.3.

$$c(T_X(-mK_X)) = \sum_{p=0}^n \sum_{i=0}^p \binom{n-i}{p-i} m^{p-i} c_i c_1^{p-i}. \quad (3.10)$$

Next, we will prove our main theorem. The following two lemmas are needed in the proof.

Lemma 3.4. Let $D_0 = 1$, and for any positive integer n let

$$D_n = \begin{vmatrix} a_1 & a_2 & a_3 & \cdots & a_n \\ 1 & a_1 & a_2 & \cdots & a_{n-1} \\ & 1 & a_1 & \cdots & a_{n-2} \\ \mathbf{0} & \ddots & \ddots & \ddots & \vdots \\ & & & 1 & a_1 \end{vmatrix},$$

then one has

$$\sum_{i=0}^n (-1)^i D_i a_{n-i} = 0,$$

here $a_0 = 1$.

Proof. Using the expansion D_n along the first row, one sees that

$$\begin{aligned} D_n &= \sum_{i=1}^n (-1)^{1+i} a_i \begin{vmatrix} 1 & a_1 & \cdots & a_{i-2} & a_i & a_{i+1} & \cdots & a_{n-1} \\ & \ddots & \ddots & \vdots & \vdots & \vdots & & \vdots \\ & & 1 & a_1 & a_3 & a_4 & \cdots & a_{n-i+2} \\ & & & 1 & a_2 & a_3 & \cdots & a_{n-i+1} \\ & & & & a_1 & a_2 & \cdots & a_{n-i} \\ & & & & 1 & a_1 & \cdots & a_{n-i-1} \\ \mathbf{0} & & & & & \ddots & \ddots & \vdots \\ & & & & & & 1 & a_1 \end{vmatrix} \\ &= \sum_{i=1}^n (-1)^{1+i} a_i D_{n-i}. \end{aligned}$$

Hence

$$\sum_{i=0}^n (-1)^i D_i a_{n-i} = 0.$$

Q.E.D.

We say cycles δ_1 and δ_2 satisfying $\delta_1 \leq \delta_2$ if and only if $\delta_2 - \delta_1$ is a nonnegative cycle.

Lemma 3.5. $\sigma_{1^t} \leq \sigma_1^t$, where t is a positive integer.

Proof. By Pieri's Formula and induction on t , we have

$$\sigma_{1^t} \leq \sigma_{1^{t-1}} \sigma_1 \leq \sigma_1^t.$$

Q.E.D.

The crucial idea for proving the main theorem is pulling back effective Schubert classes of the Chow group of the Grassmannian under the Gauss map after Schubert calculating in the Grassmannian. So we need to guarantee the intersection theory of Schubert cycles can be kept under the Gauss map which is true by Zak's theorem.

Theorem 3.6. ([Zak], Corollary 2.8) Let X be a nonsingular projective variety of dimension n and $X \neq \mathbb{P}^n$ over an algebraically closed field κ . Then the Gauss map is finite. If in addition the characteristic of κ is 0, then the Gauss map is the normalization morphism.

Let n be a positive integer. Denote $p(n)$ to be the partition number of n , i.e. the number of the way to express n as the summation of positive integers without considering the orders of them. We can define alphabet order of all vectors as follows. Given any two vectors $\vec{a} = (a_1, a_2, \dots, a_r)$ and $\vec{b} = (b_1, b_2, \dots, b_s)$ such that $a_1 \geq a_2 \geq \dots \geq a_r > 0$ and $b_1 \geq b_2 \geq \dots \geq b_s > 0$. Suppose $r = s$ otherwise we just put 0's after the short one such that $l(\vec{a}) = l(\vec{b})$. If $a_1 > b_1$ then we denote $\vec{a} > \vec{b}$. Otherwise if $a_1 = b_1$, then we compare a_2 and b_2 . Without loss of generality, suppose $a_2 > b_2$ then we denote $\vec{a} > \vec{b}$. Otherwise if $a_1 = b_1$ and $a_2 = b_2$, then we compare a_3 and b_3 and keep going. So we can compare the order of any two vectors \vec{a} and \vec{b} .

Suppose $a_1 \geq a_2 \geq \dots \geq a_r > 0$. Denote $c_{\vec{a}} = c_{a_1, a_2, \dots, a_r} := c_{a_1} c_{a_2} \dots c_{a_r}$, $c_{\vec{a}} S = c_{a_1, a_2, \dots, a_r} S := c_{a_1} S c_{a_2} S \dots c_{a_r} S$ and $c_i^t S := (c_i S)^t$.

Theorem 3.7. Let X be a nonsingular projective variety of dimension n over an algebraically closed field κ with any characteristic. Suppose K_X (or $-K_X$) is ample. If the characteristic of κ is 0 or the characteristic of κ is positive and $\mathcal{O}_X(K_X)$ ($\mathcal{O}_X(-K_X)$, respectively) is globally generated, then

$$\left(\frac{c_{2, 1^{n-2}}}{c_1^n}, \frac{c_{2, 2, 1^{n-4}}}{c_1^n}, \dots, \frac{c_n}{c_1^n} \right) \in \mathbb{A}^{p(n)} \quad (3.11)$$

is contained in a convex polyhedron in $\mathbb{A}^{p(n)}$ independent of X , where the elements in the parentheses are arranged from small to big in terms of the alphabet order of the lower indices of the numerators.

Proof. Assume $X \neq \mathbb{P}^n$ without loss of generality, because finite objects will not affect the result. Suppose mK_X is very ample (m can be negative if $-K_X$ is ample, i.e. X is Fano), where m only depends on n by the known results of Fujita's very ampleness conjecture. By Proposition 2.2 and Lemma 3.5, we have

$$0 \leq (-1)^n c_{\vec{a}} S = (-1)^n c_{a_1, a_2, \dots, a_r} S \leq (-1)^n c_{1^n} S, \quad (3.12)$$

for any \vec{a} with $a_1 \geq a_2 \geq \dots \geq a_r > 0$ and $|\vec{a}| = \sum_{i=1}^r a_i = n$. Let γ be the Gauss map defined in (3.4). From Zak's Theorem 3.6, γ is finite. Then

$$0 \leq (-1)^n c_{\vec{a}}(\gamma^* S) \leq (-1)^n (c_{1^n}(\gamma^* S)). \quad (3.13)$$

We only need to show that each element in the q -th coordinate is bounded independent of X . We use induction on q . For $q = 1$,

$$0 \leq (-1)^n c_{2, 1^{n-2}}(\gamma^* S) \leq (-1)^n (c_{1^n}(\gamma^* S)). \quad (3.14)$$

Form (3.8) and Lemma 3.3, we have

$$\begin{aligned} c_1(\gamma^* S) &= c_1(T_X(-mK_X)) + mc_1 \\ &= (nm + 1)c_1 + mc_1 \\ &= ((n + 1)m + 1)c_1. \end{aligned} \quad (3.15)$$

$$\begin{aligned} c_2(\gamma^* S) &= c_2(T_X(-mK_X)) + mc_1 c_1(T_X(-mK_X)) \\ &= \left(\frac{1}{2}n(n-1)m^2 + (n-1)m\right)c_1^2 + c_2 + mc_1(nm + 1)c_1 \\ &= \left(\frac{1}{2}n(n+1)m^2 + nm\right)c_1^2 + c_2. \end{aligned} \quad (3.16)$$

So

$$\begin{aligned} 0 \leq (-1)^n \left(\left(\frac{1}{2}n(n+1)m^2 + nm \right) c_1^2 + c_2 \right) ((n+1)m + 1)^{n-2} c_1^{n-2} \\ \leq (-1)^n ((n+1)m + 1)^n c_1^n, \end{aligned}$$

and $\frac{c_{2, 1^{n-2}}}{c_{1^n}}$ is bounded independent of X , i.e., $q = 1$ is correct.

Now for any \vec{a} with $a_1 \geq a_2 \geq \dots \geq a_r > 0$ and $\sum_{i=1}^r a_i = n$, by Lemma 3.3 and (3.8), one sees

$$c_p(T_X(-mK_X)) = \sum_{i=0}^p \binom{n-i}{p-i} m^{p-i} c_i c_1^{p-i} \quad (3.17)$$

and

$$c_p(\gamma^* S) = c_p(T_X(-mK_X)) + c_{p-1}(T_X(-mK_X))mc_1. \quad (3.18)$$

So

$$c_p(\gamma^*S) = \sum_{i=0}^p \binom{n-i}{p-i} m^{p-i} c_i c_1^{p-i} + \sum_{i=0}^{p-1} \binom{n-i}{p-1-i} m^{p-i} c_i c_1^{p-i}. \quad (3.19)$$

From (3.13), it follows

$$0 \leq (-1)^n c_{\vec{a}}(\gamma^*S) \leq (-1)^n ((n+1)m+1)^n c_1^n.$$

We can see that the biggest lower index of Chern class in the left hand side of (3.19) is \vec{a} , so we can control the value of $\frac{c_{\vec{a}}}{c_1^n}$ in terms of $\frac{c_{\vec{b}}}{c_1^n}$, for every $\vec{b} < \vec{a}$. By induction, each $\frac{c_{\vec{b}}}{c_1^n}$ is independent of X , so we are done. Q.E.D.

By the proof of the above theorem, we can have the following result easily.

Corollary 3.8. Let X be a nonsingular projective variety of dimension n over an algebraically closed field κ with any characteristic. Suppose K_X (or $-K_X$) is ample. If the characteristic of κ is 0 or the characteristic of κ is positive and $\mathcal{O}_X(K_X)$ ($\mathcal{O}_X(-K_X)$, respectively) is globally generated, then the Euler number $\chi_{top}(X)$ and Euler characteristic of the structure sheaf $\chi(X, \mathcal{O}_X)$ quotient by K_X^n is bounded, i.e. there exist constants d_1, d_2, d_3 and d_4 , which do not depend on X but only on n , such that

$$d_1 K_X^n \leq \chi_{top}(X) \leq d_2 K_X^n \quad (3.20)$$

and

$$d_3 K_X^n \leq \chi(X, \mathcal{O}_X) \leq d_4 K_X^n. \quad (3.21)$$

Proof. Since $\chi_{top}(X) = c_n$, (3.20) holds by Theorem 3.7.

For (3.21), it is direct result from Theorem 3.7 and Grothendieck-Hirzebruch-Riemann-Roch theorem since the Todd classs of X can be express as the linear combination of Chern classes. Q.E.D.

4. EQUALITIES OF CHERN CLASSES

Let X be a nonsingular projective n -fold ($n \geq 2$) of Fano or of general type with ample canonical bundle K_X over an algebraically closed field κ of any characteristic. We will produce a new method to give a bunch of inequalities in terms of all the Chern classes c_1, c_2, \dots, c_n by pulling back Schubert classes in the Grassmannian under Gauss map. Hold the notations in Section 2 and 3.

For nonsingular surface ($n = 2$):

By (3.8) and Lemma 3.3, we have

$$c_1(\gamma^*S) = c_1(T_X(-mK_X)) + mc_1$$

$$\begin{aligned}
&= (2m+1)c_1 + mc_1 \\
&= (3m+1)c_1,
\end{aligned}$$

and

$$\begin{aligned}
c_2(\gamma^*S) &= c_2(T_X(-mK_X)) + mc_1c_1(T_X(-mK_X)) \\
&= (m^2+m)c_1^2 + c_2 + mc_1(2m+1)c_1 \\
&= (3m^2+2m)c_1^2 + c_2.
\end{aligned}$$

From Proposition 2.2, it follows

$$\begin{aligned}
-c_1S &= \sigma_1, \\
c_2S = \sigma_{1,1} &= \begin{vmatrix} \sigma_1 & \sigma_2 \\ 1 & \sigma_1 \end{vmatrix} = \sigma_1^2 - \sigma_2,
\end{aligned}$$

So

$$0 \leq c_2S = \sigma_{1,1} = \sigma_1^2 - \sigma_2 \leq \sigma_1^2 = (c_1S)^2.$$

Thank for Theorem 3.6, one sees

$$0 \leq c_2(\gamma^*S) \leq (c_1(\gamma^*S))^2,$$

i.e.

$$0 \leq (3m^2+2m)c_1^2 + c_2 \leq (3m+1)^2c_1^2,$$

i.e.

$$-(3m^2+2m)c_1^2 \leq c_2 \leq (6m^2+4m+1)c_1^2.$$

Remark 4.1. From Bombieri's ([Bom]) or Reider's ([Rei]) result, we know that if characteristic of the field κ is 0 and K_X is ample, then $5K_X$ is very ample. If characteristic of the field κ is positive, by Ekedahl's result ([Eke] or cf. [Ca-Fr]), then we also have $5K_X$ is very ample. So the result is not new. However, if K_X is very ample, we have a uniform formula

$$-5c_1^2 \leq c_2 \leq 11c_1^2$$

independent of the characteristic of κ .

For nonsingular 3-fold ($n = 3$):

By (3.8) and Lemma 3.3, we have

$$\begin{aligned}
c_1(\gamma^*S) &= c_1(T_X(-mK_X)) + mc_1 \\
&= (3m+1)c_1 + mc_1 \\
&= (4m+1)c_1,
\end{aligned}$$

$$\begin{aligned}
c_2(\gamma^*S) &= c_2(T_X(-mK_X)) + mc_1c_1(T_X(-mK_X)) \\
&= (3m^2+2m)c_1^2 + c_2 + mc_1(3m+1)c_1 \\
&= (6m^2+3m)c_1^2 + c_2
\end{aligned}$$

and

$$\begin{aligned} c_3(\gamma^* S) &= c_3(T_X(-mK_X)) + mc_1c_2(T_X(-mK_X)) \\ &= (4m^3 + 3m^2)c_1^3 + 2mc_1c_2 + c_3 \\ &= m^2(4m + 3)c_1^3 + 2mc_1c_2 + c_3. \end{aligned}$$

From Proposition 2.2, one obtains

$$\begin{aligned} -c_1S &= \sigma_1, \\ c_2S &= \sigma_{1,1} = \begin{vmatrix} \sigma_1 & \sigma_2 \\ 1 & \sigma_1 \end{vmatrix} = \sigma_1^2 - \sigma_2 \end{aligned}$$

and

$$-c_3S = \sigma_{1,1,1} = \begin{vmatrix} \sigma_1 & \sigma_2 & \sigma_3 \\ 1 & \sigma_1 & \sigma_2 \\ 0 & 1 & \sigma_1 \end{vmatrix} = \sigma_1^3 + \sigma_3 - 2\sigma_1\sigma_2,$$

So

$$-c_1Sc_2S = \sigma_1\sigma_{1,1} = \sigma_1^3 - \sigma_1\sigma_2 \leq \sigma_1^3 = -(c_1S)^3.$$

By Zak's theorem, we can get an inequality in terms of c_1c_2 and c_1^3 :

$$c_1(\gamma^* S)c_2(\gamma^* S) \geq (c_1(\gamma^* S))^3,$$

i.e.,

$$(4m + 1)c_1((6m^2 + 3m)c_1^2 + c_2) \geq (4m + 1)^3c_1^3,$$

i.e.

$$(4m + 1)c_1c_2 \geq (4m + 1)(10m^2 + 5m + 1)c_1^3.$$

Moreover,

$$-c_1Sc_2S = \sigma_1\sigma_{1,1} = \sigma_{2,1} + \sigma_{1,1,1} \geq \sigma_{1,1,1} = -c_3S \geq 0.$$

So

$$c_1(\gamma^* S)c_2(\gamma^* S) \leq c_3(\gamma^* S) \leq 0,$$

i.e.,

$$(4m + 1)c_1((6m^2 + 3m)c_1^2 + c_2) \leq m^2(4m + 3)c_1^3 + 2mc_1c_2 + c_3 \leq 0,$$

i.e.

$$m(20m^2 + 15m + 3)c_1^3 + (2m + 1)c_1c_2 \leq c_3 \leq -(m^2(4m + 3)c_1^3 + 2mc_1c_2).$$

Remark 4.2. The right hand side of the above inequality is deduced by Hunt in Section 1 of [Hun] by using Gauss-Bonnet Theorem I (see [G-H], Chapter 3.3). But the left hand side of the above inequality is new.

For nonsingular 4-folds:

By (3.8) and Lemma 3.3, we obtain

$$\begin{aligned} c(T_X(-mK_X)) &= 1 + (4m+1)c_1 + ((6m^2+3m)c_1^2 + c_2) \\ &\quad + ((4m^3+3m^2)c_1^3 + 2mc_1c_2 + c_3) \\ &\quad + ((m^4+m^3)c_1^4 + m^2c_1^2c_2 + mc_1c_3 + c_4). \end{aligned}$$

So

$$\begin{aligned} c_1(\gamma^*S) &= c_1(T_X(-mK_X)) + mc_1 \\ &= (4m+1)c_1 + mc_1 \\ &= (5m+1)c_1, \end{aligned}$$

$$\begin{aligned} c_2(\gamma^*S) &= c_2(T_X(-mK_X)) + mc_1c_1(T_X(-mK_X)) \\ &= (6m^2+3m)c_1^2 + c_2 + mc_1(4m+1)c_1 \\ &= (10m^2+4m)c_1^2 + c_2, \end{aligned}$$

$$\begin{aligned} c_3(\gamma^*S) &= c_3(T_X(-mK_X)) + mc_1c_2(T_X(-mK_X)) \\ &= (4m^3+3m^2)c_1^3 + 2mc_1c_2 + c_3 \\ &\quad + mc_1((6m^2+3m)c_1^2 + c_2) \\ &= (10m^3+6m^2)c_1^3 + 3mc_1c_2 + c_3, \end{aligned}$$

and

$$\begin{aligned} c_4(\gamma^*S) &= c_4(T_X(-mK_X)) + mc_1c_3(T_X(-mK_X)) \\ &= (m^4+m^3)c_1^4 + m^2c_1^2c_2 + mc_1c_3 + c_4 \\ &\quad + mc_1((4m^3+3m^2)c_1^3 + 2mc_1c_2 + c_3) \\ &= (5m^4+4m^3)c_1^4 + 3m^2c_1^2c_2 + 2mc_1c_3 + c_4. \end{aligned}$$

From Proposition 2.2, it follows

$$-c_1S = \sigma_1,$$

$$c_2S = \sigma_{1,1} = \begin{vmatrix} \sigma_1 & \sigma_2 \\ 1 & \sigma_1 \end{vmatrix} = \sigma_1^2 - \sigma_2,$$

$$-c_3S = \sigma_{1,1,1} = \begin{vmatrix} \sigma_1 & \sigma_2 & \sigma_3 \\ 1 & \sigma_1 & \sigma_2 \\ 0 & 1 & \sigma_1 \end{vmatrix} = \sigma_1^3 + \sigma_3 - 2\sigma_1\sigma_2,$$

and

$$c_4 S = \sigma_{1,1,1,1} = \begin{vmatrix} \sigma_1 & \sigma_2 & \sigma_3 & \sigma_4 \\ 1 & \sigma_1 & \sigma_2 & \sigma_3 \\ 0 & 1 & \sigma_1 & \sigma_2 \\ 0 & 0 & 1 & \sigma_1 \end{vmatrix} = \sigma_1^4 - 3\sigma_1^2\sigma_2 + 2\sigma_1\sigma_3 + \sigma_2^2 - \sigma_4.$$

Similarly, we can have several inequalities.

(1) We can get an inequality in terms of $c_1^2 c_2$ and c_1^4 :

$$0 \leq (c_1 S)^2 c_2 S = \sigma_1^2 \sigma_{1,1} = \sigma_1^4 - \sigma_1^2 \sigma_2 \leq \sigma_1^4 = (c_1 S)^4,$$

so

$$0 \leq (c_1(\gamma^* S))^2 c_2(\gamma^* S) \leq (c_1(\gamma^* S))^4,$$

i.e.,

$$0 \leq (5m+1)^2 c_1^2 ((10m^2 + 4m)c_1^2 + c_2) \leq (5m+1)^4 c_1^4,$$

i.e.,

$$-(5m+2)c_1^4 \leq c_1^2 c_2 \leq (15m^2 + 6m + 1)c_1^4.$$

(2) We can also get an inequality in terms of $c_1^2 c_2$, $c_1 c_3$ and c_1^4 :

$$\begin{aligned} 0 \leq c_1 S c_3 S &= \sigma_1 \sigma_{1,1,1} \\ &= \sigma_1^4 + \sigma_1 \sigma_3 - 2\sigma_1^2 \sigma_2 \\ &= \sigma_1^4 + \sigma_1 \sigma_3 - 2\sigma_1(\sigma_3 + \sigma_{2,1}) \\ &= \sigma_1^4 - \sigma_1 \sigma_3 - 2\sigma_1 \sigma_{2,1} \leq \sigma_1^4 = (c_1 S)^4, \end{aligned}$$

so

$$0 \leq c_1(\gamma^* S) c_3(\gamma^* S) \leq (c_1(\gamma^* S))^4,$$

i.e.

$$0 \leq (5m+1)c_1((10m^3 + 6m^2)c_1^3 + 3mc_1 c_2 + c_3) \leq (5m+1)^4 c_1^4,$$

i.e.

$$-2m^2(5m+3)c_1^4 - 3mc_1^2 c_2 \leq c_1 c_3 \leq (115m^3 + 69m^2 + 15m + 1)c_1^4 - 3mc_1^2 c_2.$$

(3) We can also get an inequality in terms of $c_1^2 c_2$, c_2^2 and c_1^4 :

$$\begin{aligned} 0 \leq (c_2 S)^2 &= \sigma_{1,1}^2 \\ &= (\sigma_1^2 - \sigma_2)^2 \\ &= \sigma_1^4 - 2\sigma_1^2 \sigma_2 + \sigma_2^2 \\ &= \sigma_1^4 - 2(\sigma_{1,1} + \sigma_2)\sigma_2 + \sigma_2^2 \\ &= \sigma_1^4 - 2\sigma_{1,1} - \sigma_2^2 \leq \sigma_1^4 = (c_1 S)^4, \end{aligned}$$

so

$$0 \leq (c_2(\gamma^* S))^2 \leq (c_1(\gamma^* S))^4,$$

i.e.

$$0 \leq ((10m^2 + 4m)c_1^2 + c_2)^2 \leq (5m+1)^4 c_1^4,$$

i.e.

$$\begin{aligned} & -4m^2(5m+2)^2c_1^4 - 4m(5m+2)c_1^2c_2 \leq c_2^2 \\ & \leq ((5m+1)^4 - 4m^2(5m+2)^2)c_1^4 - 4m(5m+2)c_1^2c_2. \end{aligned}$$

(4) We can also get an inequality in terms of $c_1^2c_2$, c_1c_3 , c_4 and c_1^4 :

$$\begin{aligned} 0 \leq c_4S &= \sigma_{1,1,1,1} \\ &= \sigma_1^4 - 3\sigma_1^2\sigma_2 + 2\sigma_1\sigma_3 + \sigma_2^2 - \sigma_4 \\ &= \sigma_1^4 - 2(\sigma_1\sigma_3 + \sigma_{2,1}\sigma_1) - (\sigma_2^2 + \sigma_{1,1}\sigma_2) \\ &\quad + 2\sigma_1\sigma_3 + \sigma_2^2 - \sigma_4 \\ &= \sigma_1^4 - 2\sigma_{2,1}\sigma_1 - \sigma_{1,1}\sigma_2 - \sigma_4 \leq \sigma_1^4 = (c_1S)^4, \end{aligned}$$

so

$$0 \leq c_4(\gamma^*S) \leq (c_1(\gamma^*S))^4,$$

i.e.,

$$0 \leq (5m^4 + 4m^3)c_1^4 + 3m^2c_1^2c_2 + 2mc_1c_3 + c_4 \leq (5m+1)^4c_1^4,$$

i.e.,

$$-(5m^4 + 4m^3)c_1^4 \leq m^2c_1^2c_2 + 2mc_1c_3 + c_4 \leq ((5m+1)^4 - (5m^4 + 4m^3))c_1^4.$$

(5) We can also get an inequality in terms of $c_1^2c_2$, c_2^2 and c_1^4 :

$$\begin{aligned} (c_1S)^2c_2S &= \sigma_1^2\sigma_{1,1} \\ &= \sigma_1^4 - \sigma_1^2\sigma_2 \\ &= \sigma_1^4 - 2\sigma_1^2\sigma_2 + (\sigma_2 + \sigma_{1,1})\sigma_2 \\ &\geq \sigma_1^4 - 2\sigma_1^2\sigma_2 + \sigma_2^2 \\ &= (\sigma_1^2 - \sigma_2)^2 = (c_2S)^2, \end{aligned}$$

so

$$(c_2(\gamma^*S))^2 \leq (c_1(\gamma^*S))^2c_2(\gamma^*S),$$

i.e.

$$((10m^2 + 4m)c_1^2 + c_2)^2 \leq (5m+1)^2c_1^2((10m^2 + 4m)c_1^2 + c_2),$$

i.e.

$$c_2^2 \leq 2m(5m+2)(15m^2 + 6m + 1)c_1^4 + (5m^2 + 2m + 1)c_1^2c_2.$$

(6) We can also get an inequality in terms of $c_1^2c_2$, c_2^2 , c_1c_3 and c_1^4 :

$$\begin{aligned} (c_2S)^2 - c_1Sc_3S &= (\sigma_1^2 - \sigma_2)^2 - \sigma_1(\sigma_1^3 + \sigma_3 - 2\sigma_1\sigma_2) \\ &= \sigma_2^2 - \sigma_1\sigma_3 \\ &= (\sigma_4 + \sigma_{3,1} + \sigma_{2,2}) - (\sigma_4 + \sigma_{3,1}) \\ &= \sigma_{2,2} \geq 0, \end{aligned}$$

so

$$c_1(\gamma^*S)c_3(\gamma^*S) \leq (c_2(\gamma^*S))^2.$$

i.e.

$$(5m+1)c_1((10m^3+6m^2)c_1^3+3mc_1c_2+c_3) \leq ((10m^2+4m)c_1^2+c_2)^2,$$

i.e.

$$c_1c_3 \leq 10m^2(5m^2+4m+1)c_1^4+5m(4m+1)c_1^2c_2+c_2^2.$$

(7) We can also get an inequality in terms of $c_1^2c_2$, c_4 , c_1c_3 and c_1^4 :

$$\begin{aligned} c_1Sc_3S - c_4S &= \sigma_1(\sigma_1^3 + \sigma_3 - 2\sigma_1\sigma_2) \\ &\quad - (\sigma_1^4 - 3\sigma_1^2\sigma_2 + 2\sigma_1\sigma_3 + \sigma_2^2 - \sigma_4) \\ &= \sigma_1^2\sigma_2 - \sigma_1\sigma_3 - \sigma_2^2 + \sigma_4 \\ &= \sigma_1\sigma_{2,1} + \sigma_4 - (\sigma_4 + \sigma_{3,1} + \sigma_{2,2}) \\ &= \sigma_{2,1,1} \geq 0, \end{aligned}$$

so

$$c_4(\gamma^*S) \leq c_1(\gamma^*S)c_3(\gamma^*S),$$

i.e.,

$$(5m^4+4m^3)c_1^4+3m^2c_1^2c_2+2mc_1c_3+c_4 \leq (5m+1)c_1((10m^3+6m^2)c_1^3+3mc_1c_2+c_3),$$

i.e.

$$c_4 \leq 3m^2(15m^2+12m+2)c_1^4+3m(4m+1)c_1^2c_2+(3m+1)c_1c_3.$$

For general dimension n :

Let $c_0S = \sigma_0 = 1$, by Lemma 3.4, Giambelli's formula and Proposition 2.2, one obtains

$$c_nS\sigma_0 + c_{n-1}S\sigma_1 + c_{n-2}S\sigma_2 + \cdots + c_1S\sigma_{n-1} + c_0S\sigma_n = 0,$$

for any positive integer n . This implies

$$\sum_{i=0}^{\infty} c_iSx^i \sum_{i=0}^{\infty} \sigma_i x^i = 1 + \sum_{j=1}^{\infty} \left(\sum_{i=0}^j c_iS\sigma_{j-i} \right) x^j = 1,$$

where x is an indeterminate. It follows that

$$\begin{aligned} 1 + \sum_{i=1}^{\infty} \sigma_i x^i &= \frac{1}{1 + \sum_{i=1}^{\infty} c_iSx^i} \\ &= 1 + \sum_{k=1}^{\infty} (-1)^k \left(\sum_{i=1}^{\infty} c_iSx^i \right)^k \end{aligned}$$

$$\begin{aligned}
&= 1 + \sum_{k=1}^{\infty} (-1)^k \left(\sum_{m=1}^{\infty} \sum_{\substack{i_1+\dots+i_k=m \\ i_1, \dots, i_k \geq 1}} c_{i_1, \dots, i_k} S x^m \right) \\
&= 1 + \sum_{m=1}^{\infty} \sum_{k=1}^m \left((-1)^k \sum_{\substack{i_1+\dots+i_k=m \\ i_1, \dots, i_k \geq 1}} c_{i_1, \dots, i_k} S \right) x^m
\end{aligned}$$

Therefore, one sees

$$\begin{aligned}
\sigma_m &= \sum_{k=1}^m \left((-1)^k \sum_{\substack{i_1+\dots+i_k=m \\ i_1, \dots, i_k \geq 1}} c_{i_1, \dots, i_k} S \right) \\
&= \sum_{\substack{j_1+2j_2+\dots+mj_m=m \\ j_1, \dots, j_m \geq 0}} (-1)^{j_1+\dots+j_m} \frac{(j_1+\dots+j_m)!}{j_1! \dots j_m!} c_{1j_1} S \dots c_{mj_m} S.
\end{aligned} \tag{4.22}$$

In particular, we have

$$-c_1 S = \sigma_1 \tag{4.23}$$

$$c_1^2 S - c_2 S = \sigma_2 \tag{4.24}$$

$$-c_1^3 S + 2c_1 S c_2 S - c_3 S = \sigma_3 \tag{4.25}$$

$$c_1^4 S - 3c_1^2 S c_2 S + 2c_1 S c_3 S + c_2^2 S - c_4 S = \sigma_4 \tag{4.26}$$

.....

We can have a bunch of inequalities of Chern classes.

Step 1: Express n as the summation of positive integers without considering the orders of them, say $a_1 \geq a_2 \geq \dots \geq a_r > 0$, such that $n \geq \sum_{i=1}^r a_i$. Let $\vec{a} = (a_1, a_2, \dots, a_r)$.

Step 2: By Giambelli's Formula, write $\sigma_{\vec{a}}$ in terms of $\sigma_1, \sigma_2, \dots, \sigma_n$.

Step 3: By (4.22), express $\sigma_{\vec{a}}$ in terms of $c_1 S, \dots, c_n S$.

Step 4: Since $\sigma_{\vec{a}} > 0$, by Zak's Theorem 3.6, pull back of the $\sigma_{\vec{a}}$ under Gauss map γ , we can express $\gamma^*(\sigma_{\vec{a}})$ in terms of $\gamma^*(c_1 S), \dots, \gamma^*(c_n S)$ which is great than or equal to 0.

Step 5: Express $\gamma^*(c_1 S), \dots, \gamma^*(c_n S)$ in terms of c_1, c_2, \dots, c_n by (3.8) and Lemma 3.3 and get a inequality finally.

Example 4.1. Suppose $n = 5$.

Step 1: Consider $\vec{a} = (3, 2)$;

Step 2: Calculate

$$\sigma_{3,2} = \begin{vmatrix} \sigma_3 & \sigma_4 \\ \sigma_1 & \sigma_2 \end{vmatrix} = \sigma_3 \sigma_2 - \sigma_1 \sigma_4;$$

Step 3: We have

$$\begin{aligned} 0 \leq \sigma_{3,2} &= (-c_1^3 S + 2c_1 S c_2 S - c_3 S)(c_1^2 S - c_2 S) \\ &\quad + c_1 S(c_1^4 S - 3c_1^2 S c_2 S + 2c_1 S c_3 S + c_2^2 S - c_4 S) \\ &= c_1^2 S c_3 S - c_1 S c_2^2 S + c_3 S c_2 S - c_1 S c_4 S; \end{aligned}$$

Step 4:

$$(\gamma^*(c_1 S))^2 \gamma^*(c_3 S) - \gamma^*(c_1 S)(\gamma^*(c_2 S))^2 + \gamma^*(c_3 S) \gamma^*(c_2 S) - \gamma^*(c_1 S) \gamma^*(c_4 S) \geq 0.$$

Step 5: By (3.8) and Lemma 3.3,

$$\begin{aligned} c_1(\gamma^* S) &= c_1(T_X(-mK_X)) + mc_1 \\ &= (5m + 1)c_1 + mc_1 \\ &= (6m + 1)c_1, \end{aligned}$$

$$\begin{aligned} c_2(\gamma^* S) &= c_2(T_X(-mK_X)) + mc_1 c_1(T_X(-mK_X)) \\ &= (10m^2 + 4m)c_1^2 + c_2 + mc_1(5m + 1)c_1 \\ &= (15m^2 + 5m)c_1^2 + c_2, \end{aligned}$$

$$\begin{aligned} c_3(\gamma^* S) &= c_3(T_X(-mK_X)) + mc_1 c_2(T_X(-mK_X)) \\ &= (10m^3 + 6m^2)c_1^3 + 3mc_1 c_2 + c_3 \\ &\quad + mc_1((10m^2 + 4m)c_1^2 + c_2) \\ &= (20m^3 + 10m^2)c_1^3 + 4mc_1 c_2 + c_3, \end{aligned}$$

and

$$\begin{aligned} c_4(\gamma^* S) &= c_4(T_X(-mK_X)) + mc_1 c_3(T_X(-mK_X)) \\ &= (5m^4 + 4m^3)c_1^4 + 3m^2 c_1^2 c_2 + 2mc_1 c_3 + c_4 \\ &\quad + mc_1((10m^3 + 6m^2)c_1^3 + 3mc_1 c_2 + c_3) \\ &= (15m^4 + 10m^3)c_1^4 + 6m^2 c_1^2 c_2 + 3mc_1 c_3 + c_4. \end{aligned}$$

Finally, we get

$$\begin{aligned} &-(420m^5 + 350m^4 + 120m^3 + 15m^2)c_1^5 + (8m^3 - 18m^2 - 6m)c_1^3 c_2 \\ &\quad + (33m^2 + 14m + 1)c_1^2 c_3 - (2m + 1)c_1 c_2^2 \\ &\quad - (6m + 1)c_1 c_4 + c_2 c_3 \geq 0. \end{aligned}$$

If K_X is very ample, i.e. $m = 1$, then we have

$$-905c_1^5 - 16c_1^3 c_2 + 48c_1^2 c_3 - 3c_1 c_2^2 - 7c_1 c_4 + c_2 c_3 \geq 0$$

Remark 4.3. For fixed dimension n , the number of the inequalities of Chern classes is $\sum_{i=2}^n p(n)$, where $p(n)$ is the partition number of n . Partitions can be graphically visualized with Young diagrams or Ferrers diagrams. They occur in a number of branches of mathematics and physics, including the study of symmetric polynomials, the symmetric group and in group representation theory in general. It is known that (cf. [And]) an asymptotic expression for $p(n)$ is given by

$$p(n) \sim \frac{1}{4n\sqrt{3}} \exp\left(\pi\sqrt{\frac{2n}{3}}\right) \text{ as } n \rightarrow \infty.$$

ACKNOWLEDGEMENTS

The first author would like to thank N. Mok for providing excellent research environment in the University of Hong Kong while part of this research was done there. Both authors would like to thank for the reviewers for pointing out some typos and useful suggestions in the original version.

REFERENCES

- [And] G. Andrews: *The Theory of Partitions*, Cambridge University Press, 1976.
- [Bad] L. Badescu: *Algebraic surfaces*, Universitext Vol. 207, Springer (2001).
- [Bom] E. Bombieri: *Canonical models of surfaces of general type*, Publ. Math. IHES, 42 (1973), 171-219.
- [B-S-S] M. Beltrametti, M. Schneider, A. Sommese: *Chern inequalities and spannedness of adjoint bundles*, Proceedings of the Hirzebruch 65 Conference on Algebraic Geometry (Ramat Gan, 1993), 97-107, Israel Math. Conf. Proc., 9, Bar-Ilan Univ., Ramat Gan, 1996.
- [Ca-Fr] F. Catanese and M. Franciosi, *Divisors of small genus on algebraic surfaces and projective embeddings*, Proceedings of the conference Hirzebruch 65, Tel Aviv 1993, Contemp. Math., A.M.S. (1994), subseries Israel Mathematical Conference Proceedings Vol. 9 (1996), 109-140.
- [Ca-Sc] F. Catanese, M. Schneider: *Bounds for stable bundles and degrees of Weierstrass schemes*, Math. Ann. 293 (1992), 579-594.
- [Ch-Le] K. Chan, N. Leung: *Miyaoka-Yau-type inequalities for Kähler-Einstein manifolds*, Comm. Anal. Geom. 15 (2007), no. 2, 359-379.
- [CJ-H] J. Chen, C. Hacon: *On the geography of threefolds of general type*, J. Algebra 321 (2009), no. 9, 2500-2507.
- [C-K-N] M.C. Chang, H. Kim, S. Nollot: *Bounds on c_3 for threefolds*. Manuscripta Math. 97, 135-141 (1998).
- [C-L] M. Chang, A. Lopez: *A linear bound on the Euler number of threefolds of Calabi-Yau and of general type*, Manuscripta Math. 105 (2001), no. 1, 47-67.
- [CM-H] M. Chen, C. Hacon: *On the geography of Gorenstein minimal 3-folds of general type*, Asian J. Math. 10 (2006), no. 4, 757-763.
- [Dem1] J.-P. Demailly: *A numerical criterion for very ample line bundles*, J. Differential Geom., 37 (1993), 323-374.

- [Dem2] J.-P. Demailly: *Effective bounds for very ample line bundles*, Invent. Math. 124 (1996), 243-261.
- [D-P-S] J.-P. Demailly, T. Peternell, M. Schneider: *Compact complex manifolds with numerically effective tangent bundles*, J. Algebraic Geom. 3 (1994), no. 2, 295-345.
- [Eas] R. Easton: *Surfaces violating Bogomolov-Miyaoka-Yau in positive characteristic*, Proceedings of the American Mathematical Society, (7) 136 (2008), 2271-2278.
- [E-H] D. Eisenbud and J. Harris: *3264 and All That: A Second Course in Algebraic Geometry*, Cambridge University Press, Cambridge, 2016. xiv+616 pp.
- [Eke] T. Ekedahl: *Canonical models of surfaces of general type in positive characteristic*, Publ. Math. IHES, 67 (1988), 97-144.
- [E-L-N] L. Ein, R. Lazarsfeld, M. Nakamaye: *Zero-estimates, intersection theory, and a theorem of Demailly*. Higher-dimensional complex varieties (Trento, 1994), 183-207, de Gruyter, Berlin, 1996.
- [F-L] W. Fulton and R. Lazarsfeld: *Positive polynomials for ample vector bundles*, Ann. of Math. (2) 118 (1983), no. 1, 35-60.
- [Fuj] T. Fujita: *On polarized manifolds whose adjoint bundles are not semipositive*, Algebraic Geometry, Sendai, 1985, Adv. Stud. in Pure Math., North Holland, T. Oda (ed.), 10 (1987), 167-178.
- [G-H] P. Griffiths, J. Harris: *Principles of algebraic geometry*, Wiley, New York, 1978.
- [Gu] Y. Gu: *On algebraic surfaces of general type with negative c_2* , Compos. Math. 152 (2016), no. 9, 1966-1998.
- [Hun] B. Hunt: *Complex manifold geography in dimension 2 and 3*, J. Differential Geom. 30 (1989), no. 1, 51-153.
- [Kee] D. Keeler, *Fujita's conjecture and Frobenius amplitude*, Amer. J. Math. 130 (2008), no. 5, 1327-1336.
- [Ko] D. Kotschick: *Chern numbers and diffeomorphism types of projective varieties*, J. Topol. 1 (2008), no. 2, 518-526.
- [Lan] W. Lang: *Examples of surfaces of general type with vector fields*, Arithmetic and geometry, Vol. II, Progr. Math., 36, Boston, MA: Birkhäuser Boston, pp. (1983), 167-173.
- [Lie] C. Liedtke: *Algebraic surfaces of general type with small c_2 in positive characteristic*, Nagoya Math. J., 191 (2008), 111-134.
- [Mi] Y. Miyaoka: *Themes and variations on inequalities of Chern classes*. (Japanese) Sūgaku 41 (1989), no. 3, 193-207.
- [Ray] M. Raynaud: *Contre-exemple au vanishing theorem en caractéristique $p > 0$* , Tata Inst. Fund. Res. Studies in Math., 8, Berlin, New York: Springer-Verlag (1978), 273-278.
- [Rei] I. Reider: *Vector bundles of rank 2 and linear systems on algebraic surfaces*, Ann. of Math., 127 (1988), 309-316.
- [Per] U. Persson: *Chern invariants of surfaces of general type*, Compositio Math. 43 (1981) 3-58.
- [S-B] N. I. Shepherd-Barron: *Geography for surfaces of general type in positive characteristic*, Invent. Math., 106(1) (1991), 263-274.
- [Siu1] Y.-T. Siu: *An effective Matsusaka big theorem*, Ann. Inst. Fourier (Grenoble) 43 (1993), no. 5, 1387-1405.

- [Siu2] Y.-T. Siu: *Very ampleness criterion of double adjoints of ample line bundles*, Modern methods in complex analysis (Princeton, NJ, 1992), 291-318, Ann. of Math. Stud., 137, Princeton Univ. Press, Princeton, NJ, 1995.
- [Siu3] Y.-T. Siu: *Effective very ampleness*, Invent. Math. 124 (1996), 563-571.
- [Smi1] K. Smith: *Fujita's freeness conjecture in terms of local cohomology*, J. Algebraic Geom. 6 (1997), no. 3, 417-429.
- [Smi2] K. Smith: *A tight closure proof of Fujita's freeness conjecture for very ample line bundles*, Math. Ann. 317 (2000), no. 2, 285-293.
- [Sun] H. Sun: *Tilt-stability, vanishing theorems and Bogomolov-Gieseker type inequalities*, arXiv:1609.03245.
- [Szp] L. Szpiro: *Sur le théorème de rigidité de Parsin et Arakelov*.
- [Yeu] S.-K. Yeung: *Very ampleness of line bundles and canonical embedding of coverings of manifolds*, Compositio Math. 123 (2000), no. 2, 209-223.
- [Zak] F. L. Zak: *Tangents and secants of algebraic varieties*, (English summary) Translated from the Russian manuscript by the author. Translations of Mathematical Monographs, 127. American Mathematical Society, Providence, RI, 1993. viii+164 pp.

DEPARTMENT OF MATHEMATICS, SHANGHAI KEY LABORATORY OF PMMP, EAST CHINA NORMAL UNIVERSITY, RM. 312, MATH. BLDG, NO. 500, DONGCHUAN ROAD, SHANGHAI, 200241, P. R. CHINA

E-mail address: rdu@math.ecnu.edu.cn

DEPARTMENT OF MATHEMATICS, SHANGHAI NORMAL UNIVERSITY, SHANGHAI 200234, P. R. OF CHINA

E-mail address: hsun@shnu.edu.cn