

Equivalence of Intuitionistic Inductive Definitions and Intuitionistic Cyclic Proofs under Arithmetic

Stefano Berardi (Torino University)
Makoto Tatsuta (National Institute of Informatics)

Abstract

A cyclic proof system gives us another way of representing inductive definitions and efficient proof search. In 2011 Brotherston and Simpson conjectured the equivalence between the provability of the classical cyclic proof system and that of the classical system of Martin-Löf’s inductive definitions. This paper studies the conjecture for intuitionistic logic. This paper first points out that the countermodel of FOSSACS 2017 paper by the same authors shows the conjecture for intuitionistic logic is false in general. Then this paper shows the conjecture for intuitionistic logic is true under arithmetic, namely, the provability of the intuitionistic cyclic proof system is the same as that of the intuitionistic system of Martin-Löf’s inductive definitions when both systems contain Heyting arithmetic HA. For this purpose, this paper also shows that HA proves Podelski-Rybalchenko theorem for induction and Kleene-Brouwer theorem for induction. These results immediately give another proof to the conjecture under arithmetic for classical logic shown in LICS 2017 paper by the same authors.

1 Introduction

An inductive definition is a way to define a predicate by an expression which may contain the predicate itself. The predicate is interpreted by the least fixed point of the defining equation. Inductive definitions are important in computer science, since they can define useful recursive data structures such as lists and trees. Inductive definitions are important also in mathematical logic, since they increase the proof theoretic strength. Martin-Löf’s system of inductive definitions given in [11] is one of the most popular systems of inductive definitions. This system has production rules for an inductive predicate, and the production rule determines the introduction rules and the elimination rules for the predicate.

Brotherston and Simpson [5, 8] proposed an alternative formalization of inductive definitions, called a cyclic proof system. A proof, called a *cyclic proof*, is defined by proof search, going upwardly in a proof figure. If we encounter the same sequent (called a *bud*) as some sequent we already passed (called a *companion*) we can stop. The induction rule is replaced by a case rule, for this purpose. The soundness is guaranteed by some additional condition, called a *global trace condition*, which can show the case rule decreases some measure of a bud from that of the companion. In general, for proof search, a cyclic proof system can find an induction formula in a more efficient way than Martin-Löf’s system, since a cyclic proof system does not have to choose fixed induction formulas in advance. A cyclic proof system enables us efficient implementation of theorem provers with inductive definitions [4, 6, 7, 9].

Brotherston and Simpson [8] investigated Martin-Löf’s system **LKID** of inductive definitions in classical logic for the first-order language, and the cyclic proof system **CLKID**^ω for the same language, showed the provability of **CLKID**^ω includes that of **LKID**, and conjectured the equivalence.

By 2017, the equivalence was left an open question. In general, the conjecture was proved to be false in [2], by showing a counterexample. However, if we restrict both systems to only the natural number inductive predicate and add Peano arithmetic to both systems, the conjecture was proved to be true in [15], by internalizing a cyclic proof in ACA₀ and using some results in reverse mathematics. [3] proved that if we add Peano arithmetic to both systems, **CLKID**^ω and **LKID** are equivalent, namely the conjecture is true under arithmetic, by showing arithmetical Ramsey theorem and Podelski-Rybalchenko theorem for induction.

This paper studies the conjecture for intuitionistic logic, namely, the provability of the intuitionistic cyclic proof system, called **CLJID**^ω, is the same as that of the intuitionistic system of Martin-Löf’s inductive definitions, called **LJID**. This question is theoretically interesting, and answers will potentially

give new techniques of theorem proving by cyclic proofs to type theories with inductive types and program extraction by constructive proofs.

This paper first points out that the countermodel of [2] also shows the conjecture for intuitionistic logic is false in general. Then this paper shows the conjecture for intuitionistic logic is true under arithmetic, namely, the provability of \mathbf{CLJID}^ω is the same as that of \mathbf{LJID} , when both systems contain Heyting arithmetic \mathbf{HA} . Note that a counterexample in [2] does not work for a system that contains \mathbf{HA} .

We explain main ideas of this paper. The proof transformation given in [3] is intuitionistic, so we can use it for our purpose. Only thing is to intuitionistically show $\text{Ind}(>_\Pi)$. [3] uses classical logic to show arithmetical Ramsey theorem and Podelski-Rybalchenko theorem for induction, and we cannot use this technique in \mathbf{HA} . In order to show it we will take the following steps:

- (1) For each $\pi \in B$, there is n such that $\text{Ind}(>_\pi^n)$. ($()^n$ denotes the n -time composition.)
- (2) Finiteness of path relations $\{>_\pi \mid \pi \in B\}$.
- (3) Kleene-Brouwer theorem for induction.
- (4) Podelski-Rybalchenko theorem for induction.

The global trace condition gives (1). (4) is proved by (3). Combining (2) and (4), we will obtain $\text{Ind}(>_\Pi)$. The places we need arithmetic are the proofs of (3) and (4), since they use sequences of numbers. The claims (1) and (2) can be easily shown in almost the same way as [3]. We will show the claim (3) by refining an ordinary proof of Kleene-Brouwer theorem for orders. We will show the claim (4) by using Erdős trees and (3).

The results of this paper immediately give another proof to the conjecture under arithmetic for classical logic shown in [3] by using the fact $\Gamma \vdash_{\mathbf{CLKID}^\omega + \mathbf{PA}} \Delta$ implies $E, \Gamma, \neg \Delta \vdash_{\mathbf{CLJID}^\omega + \mathbf{HA}}$ for some finite set E of excluded middles.

There are not papers that study the conjecture for intuitionistic logic or Kleene-Brouwer theorem for induction in intuitionistic first-order logic. For Podelski-Rybalchenko theorem for induction, [1] intuitionistically showed it but they used second-order logic.

Section 2 describes Brotherston-Simpson conjecture. Section 3 defines $\mathbf{CLJID}^\omega + \mathbf{HA}$ and $\mathbf{LJID} + \mathbf{HA}$. Section 4 explains main ideas. Section 5 proves Kleene-Brouwer theorem for induction and Podelski-Rybalchenko theorem for induction. Section 6 discusses proof transformation and the main theorem. We conclude in Section 7.

2 Brotherston-Simpson Conjecture for Intuitionistic Logic

2.1 Intuitionistic Martin-Löf's Inductive Definition System LJID

An intuitionistic Martin-Löf's inductive definition system, called \mathbf{LJID} , is defined as the system obtained from classical Martin-Löf's inductive definition system \mathbf{LKID} defined in [8] by restricting every sequent to intuitionistic sequents and replacing $(\rightarrow L)$, $(\vee R)$, and $(\text{Ind } P_j)$ by

$$\frac{\Gamma \vdash F \quad \Gamma, G \vdash \Delta}{\Gamma, F \rightarrow G \vdash \Delta} (\rightarrow L) \quad \frac{\Gamma \vdash F}{\Gamma \vdash F \vee G} (\vee R_l) \quad \frac{\Gamma \vdash G}{\Gamma \vdash F \vee G} (\vee R_r)$$

$$\frac{\text{minor premises}}{\Gamma, P_j \vec{u} \vdash F_j \vec{u}} (\text{Ind } P_j)$$

where the minor premises are the same as the minor premises of $(\text{Ind } P_j)$ in \mathbf{LKID} (page 9 of [8]). Note that we replace these rules because their formalization in \mathbf{LKID} does not work for intuitionistic logic.

2.2 Cyclic Proof System \mathbf{CLJID}^ω

An intuitionistic cyclic proof system, called \mathbf{CLJID}^ω , is defined as the system obtained from classical cyclic proof system \mathbf{CLKID}^ω defined in [8] by restricting every sequent to intuitionistic sequents and replacing $(\rightarrow L)$ and $(\vee R)$ in the same way as \mathbf{LJID} . Note that the *global trace condition* in \mathbf{CLJID}^ω is the same as that in \mathbf{CLKID}^ω (Definition 5.5 of [8]).

2.3 Brotherston-Simpson Conjecture

Brotherston-Simpson conjecture (the conjecture 7.7 in [8]) is that the provability of **LKID** is the same as that of **CLKID**^ω. In general, the conjecture was proved to be false in [2], by showing a counterexample. [3] proved that the conjecture is true for any inductive predicates with their stage-number inductive predicates, if we add arithmetic to both systems.

This paper studies an intuitionistic version of the conjecture, namely the equivalence between **CLJID**^ω and **LJID**.

The counterexample given in [2] also shows that the equivalence between **CLJID**^ω and **LJID** does not hold in general, because the proof of the statement H in [2] is actually in **CLJID**^ω, and **LJID** does not prove H since **LKID** does not prove H . This gives us the following theorem.

Theorem 2.1 *There are some signature and some set of production rules such that the provability of **CLJID**^ω is not the same as that of **LJID**.*

This paper will show that the same results as [3] for intuitionistic logic, namely, the provability of **LJID** is the same as that of **CLJID**^ω if we add Heyting arithmetic to both systems. This means that the conjecture is true for intuitionistic logic under arithmetic.

3 Addition of Heyting Arithmetic

In this section, we define systems **CLJID**^ω + **HA** and **LJID** + **HA**.

Definition 3.1 **CLJID**^ω + **HA** and **LJID** + **HA** are defined to be obtained from **CLJID**^ω and **LJID** by adding Heyting arithmetic. Namely, we add constants and function symbols $0, s, +, \times$, the inductive predicate symbol N , the productions for N , and Heyting axioms:

$$\begin{array}{l} \frac{}{N0} \quad \frac{Nx}{Nsx} \quad \vdash Nx \rightarrow sx \neq 0, \quad \vdash Nx \wedge Ny \rightarrow sx = sy \rightarrow x = y, \\ \vdash Nx \rightarrow x + 0 = x, \quad \vdash Nx \wedge Ny \rightarrow x + sy = s(x + y), \\ \vdash Nx \rightarrow x \times 0 = 0, \quad \vdash Nx \wedge Ny \rightarrow x \times sy = x \times y + x. \end{array}$$

We define $x < y$ by $\exists z. x + sz = y$ and $x \leq y$ by $x = y \vee x < y$.

We can assume some coding of a sequence of numbers by a number in Heyting arithmetic, because the definitions on pages 115–117 of [14] work also in **HA**. We write $\langle t_0, \dots, t_n \rangle$ for the sequence of t_0, \dots, t_n . In particular, we write $\langle \rangle$ for the empty sequence. We define the i -th element of t_0, \dots, t_n as t_i . We also write $|t|$, and $(t)_u$ for the length of the sequence t , and the u -th element of the sequence t respectively. Note that $|\langle t_0, \dots, t_n \rangle| = n + 1$ and $(\langle t_0, \dots, t_n \rangle)_i = t_i$. We write $*$ for the concatenation operation of sequences.

We call an atomic formula an *inductive atomic formula* when its predicate symbol is an inductive predicate symbol. For a predicate P , we sometimes write $t \in P$ for $P(t)$.

4 Main Ideas

In [3], a given proof in **CLKID**^ω + **PA** is transformed into a proof in **LKID** + **PA**. The main construction of the proof in **LKID** + **PA** is summarized by

$$\frac{\frac{\frac{(1)\text{Ineq}}{\langle x_0, x \rangle >_{\Pi} \langle y_0, y \rangle} \quad (2)Hx_0x}{Gy_0y}}{J_1} \quad (1) \quad \begin{array}{c} \vdots \\ \text{each subproof} \\ J_2 \end{array} \quad \frac{\text{Ind}(>_{\Pi})}{Gx_0x} \quad (2)}{J}$$

One of the key points in this proof construction is the proof of $\text{Ind}(>_{\Pi})$.

The same construction works for $\mathbf{CLJID}^{\omega} + \mathbf{HA}$ and $\mathbf{LJID} + \mathbf{HA}$ except $\text{Ind}(>_{\Pi})$, since the proof uses only intuitionistic sequents when the goal sequent is an intuitionistic sequent.

In order to intuitionistically show $\text{Ind}(>_{\Pi})$, we will take the following steps:

- (1) For each $\pi \in B$, there is n such that $\text{Ind}(>_{\pi}^n)$.
- (2) Finiteness of path relations $\{>_{\pi} \mid \pi \in B\}$.
- (3) Kleene-Brouwer theorem for induction.
- (4) Podelski-Rybalchenko theorem for induction.

The global trace condition gives (1). (4) is proved by (3). Combining (2) and (4), we will obtain $\text{Ind}(>_{\Pi})$. The places we need arithmetic are the proofs of (3) and (4), since they use sequences of numbers.

The claim (1) can be shown in the same way as [3] since that paper did not use classical logic for proving (1). The claim (2) can be easily shown also in intuitionistic logic by using iteration to the least point.

For (3), we will show Kleene-Brouwer theorem for induction, which states that if we have both induction principle for a lifted tree (namely $\langle u \rangle * T$ for some tree T) with respect to the one-step extension relation and induction principle for relations on children, then we have induction principle for the Kleene-Brouwer relation. We can prove it by refining an ordinary proof of Kleene-Brouwer theorem for orders.

For (4), we will show Podelski-Rybalchenko theorem for induction stating that if transition invariant $>_{\Pi}$ is a finite union of relations $>_{\pi}$ such that each $\text{Ind}(>_{\pi}^n)$ is provable for some n , and each $(>_{\pi})$ is decidable, then $\text{Ind}(>_{\Pi})$ is provable.

First each $\text{Ind}(>_{\pi})$ is obtained by $\text{Ind}(>_{\pi}^n)$. Next by the decidability of each $(>_{\pi})$, we can assume all of $(>_{\pi})$ are disjoint to each other. For simplicity, we explain the idea of our proof for well-foundedness instead of induction principle.

Assume the relation $>_{\Pi}$ has some decreasing transitive sequence

$$u_1 >_{\Pi} u_2 >_{\Pi} u_3 >_{\Pi} \dots$$

in order to show contradiction, where a sequence is called *transitive* when $u_i >_{\Pi} u_j$ for any $i < j$.

We say the edge from u to v is of color R_i when $u >_{R_i} v$. A sequence is called *monotonically-colored* when for any element there is a color such that the edge from the element to any element after it in the sequence has the same color. Define MS as the set of monotonically-colored finite sequences. It will be shown to be well-founded with the one-step extension relation. A set of sequences beginning with the same element in some tree is called a *lifted tree*. For a decreasing transitive sequence x of U , a lifted tree $T \in U^{<\omega}$ is called an *Erdős tree* of x when the elements of x are the same as elements of elements of T , every element of T is monotonically-colored, and the edges from a parent to its children have different colors. Let ET be a function that returns an Erdős tree. Then we consider

$$\text{ET}(\langle u_1 \rangle), \text{ET}(\langle u_1, u_2 \rangle), \text{ET}(\langle u_1, u_2, u_3 \rangle), \dots$$

Define $\text{MS}_{\langle r \rangle}$ as the set of sequences beginning with r in MS. Define $>_{\text{KB1},r}$ as the Kleene-Brouwer relation for the lifted tree $\text{MS}_{\langle r \rangle}$ and some left-to-right-decreasing relation on children of the lifted tree. Define $>_{\text{KB2},r}$ as the Kleene-Brouwer relation for the lifted tree $\text{MS}_{\langle r \rangle}$ and some right-to-left-decreasing relation on children of the lifted tree. By Kleene-Brouwer theorem, $(>_{\text{KB1},r})$ and $(>_{\text{KB2},r})$ are well-founded. Define $\text{ET2}(\langle u_1, \dots, u_n \rangle)$ as the $(>_{\text{KB2},u_1})$ -sorted sequence of elements in $\text{ET}(\langle u_1, \dots, u_n \rangle)$. Then consider

$$\text{ET2}(\langle u_1 \rangle), \text{ET2}(\langle u_1, u_2 \rangle), \text{ET2}(\langle u_1, u_2, u_3 \rangle), \dots$$

Define $>_{\text{KB},r}$ as the Kleene-Brouwer relation for $>_{\text{KB1},r}$ and the set of $(>_{\text{KB2},r})$ -sorted finite sequences of elements in $\text{MS}_{\langle r \rangle}$. This definition is a key idea. By this definition, we have

$$\text{ET2}(\langle u_1 \rangle) >_{\text{KB},u_1} \text{ET2}(\langle u_1, u_2 \rangle) >_{\text{KB},u_1} \text{ET2}(\langle u_1, u_2, u_3 \rangle) >_{\text{KB},u_1} \dots$$

Since $(>_{\text{KB},u_1})$ is well-founded by Kleene-Brouwer theorem, we have contradiction.

In general we need classical logic to derive induction principle from well-foundedness, but the idea we have explained will work well for showing induction principle in intuitionistic logic.

5 HA-Provable Podelski-Rybalchenko Theorem for Induction

This section will prove Podelski-Rybalchenko theorem for induction, inside Heyting arithmetic **HA**. First we will prove Kleene-Brouwer theorem for induction, inside **HA**. Next we will show induction for the set MS of monotonically-colored subsequences. Then by applying Kleene-Brouwer theorem to a part of MS and some orders $>_{u, \text{Left}}$ and $>_{u, \text{Right}}$, we will obtain two Kleene-Brouwer relations $>_{\text{KB1}, r}$ and $>_{\text{KB2}, r}$ and show their induction principle. Then by applying Kleene-Brouwer theorem to some lifted tree determined by $>_{\text{KB2}, r}$ and the relation $>_{\text{KB1}, r}$, we will obtain a Kleene-Brouwer relation $>_{\text{KB}, r}$ and show its induction principle. Then we will show induction for decreasing transitive sequences is reduced to induction for Erdős trees with the relation $>_{\text{KB}, r}$. Since Erdős trees are in the lifted tree, by combining them, we will prove Podelski-Rybalchenko theorem for induction.

We write $>_R$ or $>$ for a binary relation. We write $<_R$ for the binary relation of the inverse of $>_R$. We write $y <_R x \in X$ for $y <_R x \wedge y \in X$. We write $x \in \sigma$ when x is an element of the sequence σ . We write $U^{<\omega}$ for the set of finite sequences of elements in U . For a set S of sequences, we write $\langle u \rangle * S$ for $\{\langle u \rangle * \sigma \mid \sigma \in S\}$. For a set U and a binary relation $>_R$ for U , the *induction principle* for $(U, >_R)$ is defined as

$$\begin{aligned} \text{Ind}(U, >_R, F) &\equiv \forall x \in U ((\forall y <_R x \in U. Fy) \rightarrow Fx) \rightarrow \forall x \in U. Fx, \\ \text{Ind}(U, >_R) &\equiv \text{Ind}(U, >_R, F) \text{ (for every formula } Fx). \end{aligned}$$

For a set U a set T is called a *tree* of U if $T \subseteq U^{<\omega}$ and T is nonempty and closed under prefix operations. Note that the empty sequence is a prefix of any sequence. As a graph, the set of nodes is T and the set of edges is $\{(x, y) \in T^2 \mid y = x * \langle u \rangle\}$. We call a set $T \subseteq U^{<\omega}$ a *lifted tree* of U when there are a tree $T' \subseteq U^{<\omega}$ and $r \in U$ such that $T = \langle r \rangle * T'$. We define $\text{LiftedTree}(T, U)$ as a first-order formula that means T is a lifted tree of U .

For $x, y \in U^{<\omega}$ we define the one-step extension relation $x >_{\text{ext}} y$ if $y = x * \langle u \rangle$ for some u . For a set $T \subseteq U^{<\omega}$ and $\sigma \in U^{<\omega}$, we define T_σ as $\{\rho \in T \mid \rho = \sigma * \rho'\}$. Note that T_σ is a subset of T . For a nonempty sequence σ , we define $\text{first}(\sigma)$ and $\text{last}(\sigma)$ as the first and the last element of σ respectively.

The next lemma shows induction implies $x \not> x$.

Lemma 5.1 *If $\text{HA} \vdash \text{Ind}(U, >)$, then $\text{HA} \vdash \forall x, y \in U (y < x \rightarrow y \neq x)$.*

Proof. Fix $x, y \in U$ and assume $y < x$ and $y = x$ in order to show contradiction.

Define Fz be $z \neq x$. Then we can show

$$\text{HA} \vdash \forall z \in U ((\forall w < z \in U. Fw) \rightarrow Fz)$$

by case analysis for $z \neq x \vee z = x$ as follows. In the first case $z \neq x$, Fz . In the second case, by taking w to be x in $\forall w < z \in U. Fw$, we have Fx .

By $\text{Ind}(U, >)$, $\forall z \in U. Fz$. By taking z to be x , we have contradiction. \square

Definition 5.2 (Kleene-Brouwer Relation) For a set U , a lifted tree T of U , and the set of binary relations $>_u$ on U for every $u \in U$, we define the *Kleene-Brouwer relation* $>_{\text{KB}}$ for T and $\{(>_u) \mid u \in U\}$ as follows: for $x, y \in T$, $x >_{\text{KB}} y$ if (1) $x = z * \langle u, u_1 \rangle * w_1$, $y = z * \langle u, u_2 \rangle * w_2$, and $u_1 >_u u_2$ for some z, u, u_1, w_1, u_2, w_2 , or (2) $y = x * z$ for some $z \neq \langle \rangle$.

When $(>_u)$ is some fixed $(>)$ for all u , for simplicity we call the relation $(>_{\text{KB}})$ the Kleene-Brouwer relation for T and $>$.

Note that $(>_{\text{KB}})$ is a relation on T . This Kleene-Brouwer relation is slightly different from ordinary Kleene-Brouwer order for the following points: it creates a relation instead of an order, it uses a set of relations indexed by an element, and it is defined for a lifted tree instead of a tree (in order to use indexed relations).

The next theorem shows Kleene-Brouwer theorem for induction, which states that if we have both induction principle for a lifted tree with respect to the extension relation and induction principle for relations on children, then we have induction principle for the Kleene-Brouwer relation.

Theorem 5.3 (Kleene-Brouwer Theorem for Induction) *If $\text{HA} \vdash \text{LiftedTree}(T, U)$, $\text{HA} \vdash \text{Ind}(T, >_{\text{ext}})$ and $\text{HA} \vdash \forall u \in U. \text{Ind}(U, >_u)$, then $\text{HA} \vdash \text{Ind}(T, >_{\text{KB}})$.*

Proof. By induction on $(T, >_{\text{ext}})$, we will show $\forall \sigma \in T. \text{Ind}(T_\sigma, >_{\text{KB}})$. After we prove it, we can take σ to be $\langle \rangle$ to show the theorem, since $T_{\langle \rangle} = T$.

Fix $\sigma \in T$ in order to show $\text{Ind}(T_\sigma, >_{\text{KB}})$. Note that we can use induction hypothesis for every $\sigma * \langle u \rangle \in T$:

$$\text{Ind}(T_{\sigma * \langle u \rangle}, >_{\text{KB}}). \quad (1)$$

Assume

$$\forall x \in T_\sigma ((\forall y <_{\text{KB}} x \in T_\sigma.Fy) \rightarrow Fx) \quad (2)$$

in order to show $\forall x \in T_\sigma.Fx$. For simplicity we write $F(X)$ for $\forall x \in X.Fx$. Let $Gu \equiv F(T_{\sigma * \langle u \rangle})$. By $\text{Ind}(U, >_{\text{last}(\sigma)})$ we will show the following claim.

Claim: $\forall u \in U.Gu$.

Fix $u \in U$ in order to show Gu .

By IH for v with $>_{\text{last}(\sigma)}$ we have

$$v <_{\text{last}(\sigma)} u \rightarrow F(T_{\sigma * \langle v \rangle}). \quad (3)$$

We can show

$$\forall x \in T_{\sigma * \langle u \rangle} ((\forall y <_{\text{KB}} x \in T_{\sigma * \langle u \rangle}.Fy) \rightarrow (\forall y <_{\text{KB}} x \in T_\sigma.Fy)) \quad (4)$$

as follows. Fix $x \in T_{\sigma * \langle u \rangle}$, assume

$$\forall y <_{\text{KB}} x \in T_{\sigma * \langle u \rangle}.Fy \quad (5)$$

and assume $y <_{\text{KB}} x \in T_\sigma$ in order to show Fy . By definition of $>_{\text{KB}}$, we have $y \in T_{\sigma * \langle v \rangle}$ for some $v <_{\text{last}(\sigma)} u$, or $y \in T_{\sigma * \langle u \rangle}$. In the first case, Fy by (3). In the second case, Fy by (5). Hence we have shown (4).

Combining (4) with (2), we have

$$\forall x \in T_{\sigma * \langle u \rangle} ((\forall y <_{\text{KB}} x \in T_{\sigma * \langle u \rangle}.Fy) \rightarrow F(x)). \quad (6)$$

By IH (1) for $\sigma * \langle u \rangle$, we have $\text{Ind}(T_{\sigma * \langle u \rangle}, >_{\text{KB}})$, namely,

$$\forall x \in T_{\sigma * \langle u \rangle} ((\forall y <_{\text{KB}} x \in T_{\sigma * \langle u \rangle}.Fy) \rightarrow Fx) \rightarrow \forall x \in T_{\sigma * \langle u \rangle}.Fx. \quad (7)$$

By (7)(6), $F(T_{\sigma * \langle u \rangle})$. Hence we have shown the claim.

If $y <_{\text{KB}} \sigma \in T_\sigma$, we have $y \in T_{\sigma * \langle u \rangle}$ for some u , since $y <_{\text{KB}} \sigma$ implies $y \neq \sigma$ by definition of KB and Lemma 5.1 for $>_u$. By the claim, Fy . Hence

$$\forall y <_{\text{KB}} \sigma \in T_\sigma.Fy. \quad (8)$$

By letting $x := \sigma$ in (2), we have $(\forall y <_{\text{KB}} \sigma \in T_\sigma.Fy) \rightarrow F\sigma$. By (8), $F\sigma$. Combining it with the claim, $\forall x \in T_\sigma.Fx$. \square

Definition 5.4 For a set U and a relation $>$ for U , we define the set $\text{DS}(U, >)$ of decreasing sequences as $\{\langle x_0, \dots, x_{n-1} \rangle \mid n \geq 0, x_i \in U, \forall i < n-1. (x_i > x_{i+1})\}$.

We define the set $\text{DT}(U, >)$ of decreasing transitive sequences by $\{\langle x_0, \dots, x_{n-1} \rangle \mid n \geq 0, x_i \in U, \forall i (\forall j \leq n-1. (i < j \rightarrow x_i > x_j))\}$.

We define $>_{R_1 \cup \dots \cup R_k}$ as the union of $>_{R_i}$ for all $1 \leq i \leq k$. We define $>_{R_1 + \dots + R_k}$ as the disjoint union of $>_{R_i}$ for all $1 \leq i \leq k$. (Whenever we use it, we implicitly assume the disjointness is provable in **HA**.)

We define $\text{Monoseq}_{R_1, \dots, R_k}(x)$ to hold when $x = \langle x_0, \dots, x_{n-1} \rangle \in \text{DT}(U, >_{R_1 + \dots + R_k})$ and $\forall i < n-1. (\forall j \leq n-1. (i < j \rightarrow \bigwedge_{1 \leq l \leq k} (x_i >_{R_l} x_{i+1} \rightarrow x_i >_{R_l} x_j)))$. Note that n may be 0.

We define MS as $\{x \in \text{DT}(U, >_{R_1 + \dots + R_k}) \mid \text{Monoseq}_{R_1, \dots, R_k}(x)\}$.

MS is the set of monotonically-colored finite sequences. Note that $\text{MS}_{\langle r \rangle}$ is a subset of MS and a lifted tree of U for any $r \in U$.

Definition 5.5 For a relation $>_{R_i}$ on U_i for $1 \leq i \leq k$, we define a relation $>_{R_1 \times \dots \times R_k}$ on $U_1 \times \dots \times U_k$ by: $(x_1, \dots, x_k) >_{R_1 \times \dots \times R_k} (y_1, \dots, y_k)$ if there is some i such that $x_i >_{R_i} y_i$ and $x_j = y_j$ for all $j \neq i$.

The next lemma shows induction for cartesian product.

Lemma 5.6 If $\mathbf{HA} \vdash \text{Ind}(U_i, >_{R_i})$ for $1 \leq i \leq k$, then $\mathbf{HA} \vdash \text{Ind}(U_1 \times \dots \times U_k, >_{R_1 \times \dots \times R_k})$.

Proof. First we will show the case $k = 2$.

For simplicity, we write U for $U_1 \times U_2$ and $>_{\times}$ for $>_{R_1 \times R_2}$.

Assume

$$\forall x \in U ((\forall y <_{\times} x \in U.Fy) \rightarrow Fx) \quad (9)$$

in order to show $\forall x \in U.Fx$.

Define $Gx_1 \equiv \forall x_2 \in U_2.F(x_1, x_2)$. We will show

$$\forall x_1 \in U_1 ((\forall y_1 <_{R_1} x_1 \in U_1.Gy_1) \rightarrow Gx_1). \quad (10)$$

Fix $x_1 \in U_1$ and assume

$$\forall y_1 <_{R_1} x_1 \in U_1.Gy_1 \quad (11)$$

in order to show Gx_1 . We will show

$$\forall x_2 \in U_2 ((\forall y_2 <_{R_2} x_2.F(x_1, y_2)) \rightarrow F(x_1, x_2)). \quad (12)$$

Fix $x_2 \in U_2$ and assume

$$\forall y_2 <_{R_2} x_2.F(x_1, y_2) \quad (13)$$

in order to show $F(x_1, x_2)$. We will show

$$\forall y <_{\times} (x_1, x_2) \in U.Fy. \quad (14)$$

Fix $y <_{\times} (x_1, x_2) \in U$ in order to show Fy . Let (y_1, y_2) be y . Consider cases by $y <_{\times} (x_1, x_2)$.

Case 1. $y_1 = x_1$ and $y_2 <_{R_2} x_2$.

By taking y_2 to be y_2 in (13), $F(x_1, y_2)$. Hence Fy .

Case 2. $y_1 <_{R_1} x_1$ and $y_2 = x_2$.

By taking y_1 to be y_1 in (11), Gy_1 . Hence $\forall x_2 \in U_2.F(y_1, x_2)$. By taking x_2 to be x_2 in it, $F(y_1, x_2)$. Hence Fy .

In both cases, Fy . Hence we have shown (14). By taking x to be (x_1, x_2) in (9), we have $F(x_1, x_2)$. Hence we have shown (12). By $\text{Ind}(U_2, >_{R_2})$ for $\lambda x_2.F(x_1, x_2)$, we have $\forall x_2 \in U_2.F(x_1, x_2)$. Hence Gx_1 . Hence we have shown (10). By $\text{Ind}(U_1, >_{R_1})$ for G , we have $\forall x_1 \in U_1.Gx_1$. Hence $\forall x \in U.Fx$.

We have shown the case $k = 2$.

Next we will show the case $k > 2$. We use induction on k to show the claim. By IH, we have $\text{Ind}(U_1 \times \dots \times U_{k-1}, >_{R_1 \times \dots \times R_{k-1}})$. By using the case $k = 2$ for it and $\text{Ind}(U_k, >_{R_k})$, we have $\text{Ind}(U_1 \times \dots \times U_k, >_{(R_1 \times \dots \times R_{k-1}) \times R_k})$. Since $(>_{(R_1 \times \dots \times R_{k-1}) \times R_k})$ is $(>_{R_1 \times \dots \times R_k})$, we have the claim. \square

The next lemma shows that induction principle for each relation implies induction principle for monotonically-colored sequences. This lemma can be proved by refining Lemma 6.4 (1) of [1] from second-order logic to first-order logic.

Lemma 5.7 If $\mathbf{HA} \vdash \text{Ind}(\text{DT}(U, >_{R_i}), >_{\text{ext}})$ for all $1 \leq i \leq k$, then $\mathbf{HA} \vdash \forall r \in U. \text{Ind}(\text{MS}_{\langle r \rangle}, >_{\text{ext}})$.

Proof. Fix $r \in U$ in order to show $\text{Ind}(\text{MS}_{\langle r \rangle}, >_{\text{ext}})$.

Assume

$$\forall \sigma \in \text{MS}_{\langle r \rangle} ((\forall \rho <_{\text{ext}} \sigma \in \text{MS}_{\langle r \rangle}. F\rho) \rightarrow F\sigma) \quad (15)$$

in order to show $\forall \sigma \in \text{MS}_{\langle r \rangle}. F\sigma$.

For $1 \leq i \leq k$, define

$$\text{Seq}_i(\sigma) = \langle x_{n_1}, \dots, x_{n_m} \rangle$$

where $\sigma = \langle x_1, \dots, x_n \rangle$, $\{x_{n_1}, \dots, x_{n_m}\} = \{x_j \in \sigma \mid x_j >_{R_i} x_{j+1}\}$, and $n_1 <_N \dots <_N n_m$ for the natural number order $>_N$. Formally $\text{Seq}_i(\sigma) = \rho$ is an abbreviation for some **HA**-formula $F(\sigma, \rho)$. Note that $\text{Seq}_i(\sigma)$ may be $\langle \rangle$.

For simplicity we write DT_k for $\text{DT}(U, >_{R_1}) \times \dots \times \text{DT}(U, >_{R_k})$.

We define ext^k for DT_k by

$$(x_1, \dots, x_k) >_{\text{ext}^k} (y_1, \dots, y_k)$$

where for some $1 \leq i \leq k$, $x_i >_{\text{ext}} y_i$ and $x_j = y_j$ for all $j \neq i$. Note that the set of elements of σ is the union of the sets of $\text{Seq}_i(\sigma)$ ($1 \leq i \leq k$) and $\{\text{last}(\sigma)\}$.

Define

$$G(x_1, \dots, x_k) \equiv \forall \sigma \in \text{MS}_{\langle r \rangle} ((1 \leq \forall i \leq k. \text{Seq}_i(\sigma) = x_i) \rightarrow F\sigma).$$

We write \vec{x} for (x_1, \dots, x_k) . We will show

$$\forall \vec{x} \in \text{DT}_k ((\forall \vec{y} <_{\text{ext}^k} \vec{x} \in \text{DT}_k. G(\vec{y})) \rightarrow G(\vec{x})). \quad (16)$$

Fix \vec{x} and assume

$$\forall \vec{y} <_{\text{ext}^k} \vec{x} \in \text{DT}_k. G(\vec{y}) \quad (17)$$

in order to show $G\vec{x}$. Fix $\sigma \in \text{MS}_{\langle r \rangle}$ and assume $\text{Seq}_i(\sigma) = x_i$ for all $1 \leq i \leq k$ in order to show $F\sigma$.

We can show $\forall \rho <_{\text{ext}} \sigma \in \text{MS}_{\langle r \rangle}. F\rho$ as follows. Assume $\rho <_{\text{ext}} \sigma$. Let $\rho = \sigma * \langle u \rangle$. Then $\text{last}(\sigma) >_{R_i} u$ for some $1 \leq i \leq k$. Then

$$\text{Seq}_i(\rho) = \text{Seq}_i(\sigma) * \langle \text{last}(\sigma) \rangle, \quad \text{Seq}_j(\rho) = \text{Seq}_j(\sigma) \quad (\forall j \neq i).$$

Hence $(\text{Seq}_1(\rho), \dots, \text{Seq}_k(\rho)) <_{\text{ext}^k} (\text{Seq}_1(\sigma), \dots, \text{Seq}_k(\sigma))$, namely, $(\text{Seq}_1(\rho), \dots, \text{Seq}_k(\rho)) <_{\text{ext}^k} \vec{x}$. By (17), $G(\text{Seq}_1(\rho), \dots, \text{Seq}_k(\rho))$. Hence $F\rho$. Hence we have shown $\forall \rho <_{\text{ext}} \sigma \in \text{MS}_{\langle r \rangle}. F\rho$.

By (15), $F\sigma$. Hence we have shown $G\vec{x}$. Hence we have shown (16).

By Lemma 5.6 for $\text{Ind}(\text{DT}(U, >_{R_i}), >_{\text{ext}})$ for all $1 \leq i \leq k$, we have $\text{Ind}(\text{DT}_k, >_{\text{ext}^k})$. By it and (16), we have

$$\forall \vec{x} \in \text{DT}_k. G\vec{x}.$$

For every $\sigma \in \text{MS}_{\langle r \rangle}$, by letting $x_i = \text{Seq}_i(\sigma)$ for all $1 \leq i \leq k$, we have $G\vec{x}$, and hence we have $F\sigma$. \square

Next we create Kleene-Brouwer relations $>_{\text{KB1},r}$ and $>_{\text{KB2},r}$ for monotonically-colored sequences beginning with r . Then we consider the set of $(>_{\text{KB2},r})$ -sorted finite sequences of monotonically-colored finite sequences beginning with r . It is a lifted tree. Then, by induction principle for MS, the lifted tree is well-founded with the one-step extension relation. The Kleene-Brouwer relation for the lifted tree and $>_{\text{KB1},r}$ gives us $>_{\text{KB},r}$ for the lifted tree. Since an Erdős tree is in the lifted tree, this will later show induction principle for Erdős trees.

Definition 5.8 For $u \in U$, we define $>_{u,\text{Left}}$ for U by: $u_1 >_{u,\text{Left}} u_2$ if $u >_{R_j} u_1$, $u >_{R_l} u_2$, and $j < l$ for some j, l .

We define $>_{\text{KB1},r}$ for $\text{MS}_{\langle r \rangle}$ as the KB relation for $\text{MS}_{\langle r \rangle} \subseteq U^{<\omega}$ and $(>_{u,\text{Left}}) \subseteq U^2$ for all $u \in U$.

For $u \in U$, we define $>_{u,\text{Right}}$ for U by: $u_1 >_{u,\text{Right}} u_2$ if $u_1 <_{u,\text{Left}} u_2$.

We define $>_{\text{KB2},r}$ for $\text{MS}_{\langle r \rangle}$ as the KB relation for $\text{MS}_{\langle r \rangle} \subseteq U^{<\omega}$ and $(>_{u,\text{Right}}) \subseteq U^2$ for all $u \in U$.

We define $>_{\text{KB},r}$ for $\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2},r})_{\langle \langle r \rangle \rangle}$ as the KB relation for $\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2},r})_{\langle \langle r \rangle \rangle} \subseteq \text{MS}_{\langle r \rangle}^{<\omega}$ and $>_{\text{KB1},r}$.

$>_{u,\text{Left}}$ is the left-to-right-decreasing order of children of u in some ordered tree of U in which the edge label R_i is put to an edge (x, y) such that $x >_{R_i} y$, each parent has at most one child of the same edge label, and children are ordered by their edge labels with $R_1 < \dots < R_k$. Similarly $>_{u,\text{Right}}$ is the right-to-left-decreasing order of children of u in the ordered tree.

Definition 5.9 For $u \in U \subseteq N$, finite $T \subseteq \text{MS}$ such that $\forall \rho \in T. \forall v \in \rho. (v >_{R_1+\dots+R_k} u)$, and for $\sigma \in T$, we define the function insert by:

$$\begin{aligned} \text{insert}(u, T, \sigma) = \\ \text{insert}(u, T, \sigma * \langle v \rangle) \text{ if } \text{last}(\sigma) >_{R_i} u, v = \mu v. (\sigma * \langle v \rangle \in T \wedge \text{last}(\sigma) >_{R_i} v), \\ T \cup \{\sigma * \langle u \rangle\} \text{ otherwise,} \end{aligned}$$

where $\mu v.F(v)$ denotes the least element v with the natural number order such that $F(v)$. Formally $\text{insert}(u, T, \sigma) = T'$ is an abbreviation of some **HA**-formula $G(u, T, \sigma, T')$. It is the same for ET below.

For $x \in \text{DT}(U, >_{R_1+\dots+R_k}) - \{\langle \rangle\}$, we define $\text{ET}(x) \subseteq \text{MS}$ by

$$\begin{aligned} \text{ET}(\langle u \rangle) &= \{\langle u \rangle\}, \\ \text{ET}(x * \langle u \rangle) &= \text{insert}(u, \text{ET}(x), \langle \text{first}(x) \rangle) \text{ if } x \neq \langle \rangle. \end{aligned}$$

Note that $\text{insert}(u, T, \sigma)$ adds a new element u to the set T at some position below σ to obtain a new set. $\text{ET}(x)$ is an Erdős tree obtained from the decreasing transitive sequence x .

The next lemma (1) states a new element is inserted at a leaf. It is proved by induction on the number of elements in T . The claim (2) states that edges from a parent to its children have different colors. It is proved by induction on the length of x .

Lemma 5.10 (1) For $u \in U$, $T \subseteq \text{MS}$, and $\sigma \in T$, if $u \notin \rho$ for all $\rho \in T$, $\sigma = \langle x_0, \dots, x_{n-1} \rangle$, $x_i >_{R_j} x_{i+1}$ implies $x_i >_{R_j} u$ for all $i < n-1$, and $\text{insert}(u, T, \sigma) = T'$, then there is some $\rho \in T_\sigma$ such that $\rho * \langle u \rangle \in \text{MS}$, $T' = T + \{\rho * \langle u \rangle\}$, and $\rho * \langle u \rangle$ is a maximal sequence in T' .

(2) If $\sigma * \langle u, u_1 \rangle * \rho_1, \sigma * \langle u, u_2 \rangle * \rho_2 \in \text{ET}(x)$, $u >_{R_i} u_1$, and $u >_{R_i} u_2$, then $u_1 = u_2$.

Definition 5.11 For $x \in \text{DT}(U, >_{R_1+\dots+R_k}) - \{\langle \rangle\}$, we define

$$\text{ET2}(x) \equiv \langle x_0, \dots, x_{n-1} \rangle$$

where $\{x_0, \dots, x_{n-1}\} = \text{ET}(x)$ and $\forall i < n-1. (x_i >_{\text{KB2, first}(x)} x_{i+1})$.

Note that $>_{\text{KB2, first}(x)}$ is a total order on $\text{ET}(x)$ by Lemma 5.10 (2). $\text{ET2}(x)$ is the decreasing sequence of all nodes in the Erdős tree $\text{ET}(x)$ ordered by $>_{\text{KB2, first}(x)}$.

The next lemma shows ET2 is monotone. It is the key property of reduction in Lemma 5.13.

Lemma 5.12 **HA** $\vdash \forall r \in U. \forall x, y \in \text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle}. (x >_{\text{ext}} y \rightarrow \text{ET2}(x) >_{\text{KB}, r} \text{ET2}(y))$.

Proof. Fix $r \in U$ and $x, y \in \text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle}$ and assume $x >_{\text{ext}} y$. Let $y = x * \langle u \rangle$. Then $\text{ET}(y) = \text{insert}(u, \text{ET}(x), \langle r \rangle)$. By Lemma 5.10 (1), we have σ such that $\text{ET}(y) = \text{ET}(x) + \{\sigma * \langle u \rangle\}$. Then we have two cases:

Case 1. $\text{last}(\text{ET2}(x)) >_{\text{KB2}, r} \sigma * \langle u \rangle$.

Then $\text{ET2}(y) = \text{ET2}(x) * \langle \sigma * \langle u \rangle \rangle$. By definition, $\text{ET2}(x) >_{\text{KB}, r} \text{ET2}(y)$.

Case 2. $\sigma * \langle u \rangle >_{\text{KB2}, r} \tau$ for some $\tau \in \text{ET2}(x)$.

Let ρ be the next element of $\sigma * \langle u \rangle$ in $\text{ET2}(y)$. Then $\text{ET2}(x) = \alpha * \langle \rho \rangle * \beta$ and $\text{ET2}(y) = \alpha * \langle \sigma * \langle u \rangle, \rho \rangle * \beta$. By definition of ET2, $\sigma * \langle u \rangle >_{\text{KB2}, r} \rho$. Since $\sigma * \langle u \rangle$ is maximal in $\text{ET}(y)$ by Lemma 5.10 (1), there is no $\alpha \neq \langle \rangle$ such that $\sigma * \langle u \rangle * \alpha = \rho$. Hence $\sigma * \langle u \rangle <_{\text{KB1}, r} \rho$. Hence $\text{ET2}(x) >_{\text{KB}, r} \text{ET2}(y)$. \square

The next lemma shows that induction for decreasing transitive sequences is reduced to induction for Erdős trees with $>_{\text{KB}, r}$.

Lemma 5.13 **HA** $\vdash \forall r \in U. \text{Ind}(\text{ET2}(\text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle}), >_{\text{KB}, r}) \text{ implies } \text{HA} \vdash \text{Ind}(\text{DT}(U, >_{R_1+\dots+R_k}), >_{\text{ext}})$.

Proof. In this proof, for simplicity, we write DT for $\text{DT}(U, >_{R_1+\dots+R_k})$ and we also write ET_r for $\text{ET2}(\text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle})$.

Assume **HA** $\vdash \forall r \in U. \text{Ind}(\text{ET}_r, >_{\text{KB}, r})$. Assume

$$\forall x \in \text{DT}((\forall y <_{\text{ext}} x \in \text{DT}. Fy) \rightarrow Fx). \quad (18)$$

in order to show

$$\forall x \in \text{DT}. Fx. \quad (19)$$

Define $Gy \equiv \forall z \in \text{DT}(z \neq \langle \rangle \rightarrow \text{ET2}(z) = y \rightarrow Fz)$. We will show

$$\forall r \in U. \forall x \in \text{ET}_r((\forall y <_{\text{KB}, r} x \in \text{ET}_r. Gy) \rightarrow Gx). \quad (20)$$

Fix $r \in U$ and $x \in \text{ET}_r$ and assume

$$\forall y <_{\text{KB}, r} x \in \text{ET}_r. Gy \quad (21)$$

in order to show Gx . Fix $x_0 \in \text{DT}$ and assume $x_0 \neq \langle \rangle$ and $\text{ET2}(x_0) = x$ in order to show Fx_0 .

We can show

$$\forall y_0 <_{\text{ext}} x_0 \in \text{DT}. Fy_0$$

as follows. Let r be $\text{first}(x_0)$. Fix $y_0 <_{\text{ext}} x_0 \in \text{DT}$. Then $x_0, y_0 \in \text{DT}_{\langle r \rangle}$. Let $y = \text{ET2}(y_0)$. By Lemma 5.12, $y <_{\text{KB}, r} x$. By (21) and $y \in \text{ET}_r$, Gy . Hence Fy_0 .

By taking x to be x_0 in (18), Fx_0 . Hence $\forall x_0 \in \text{DT}(x_0 \neq \langle \rangle \rightarrow x = \text{ET2}(x_0) \rightarrow Fx_0)$, namely, Gx . We have shown (20).

By (20) and $\forall r \in U. \text{Ind}(\text{ET}_r, >_{\text{KB}, r})$, we have $\forall r \in U. \forall x \in \text{ET}_r. Gx$.

For any $x \in \text{DT}$ such that $x \neq \langle \rangle$, by taking r to be $\text{first}(x)$ and x to be $\text{ET2}(x)$ in it we have $G(\text{ET2}(x))$. Hence Fx . For $x = \langle \rangle$, by taking x to be $\langle \rangle$ in (18), $F\langle \rangle$. Combining them, we have (19). \square

The next lemma shows induction holds when we restrict the universe.

Lemma 5.14 $\mathbf{HA} \vdash \text{Ind}(U, >)$ and $\mathbf{HA} \vdash V \subseteq U$ imply $\mathbf{HA} \vdash \text{Ind}(V, >)$.

Proof. We will show $\text{Ind}(V, >)$ for F , namely,

$$\forall x \in V((\forall y < x \in V. Fy) \rightarrow Fx) \rightarrow \forall x \in V. Fx. \quad (22)$$

Let Gx be $x \in V \rightarrow Fx$. By $\text{Ind}(U, >)$ for G , we have

$$\forall x \in U((\forall y < x \in U. Gy) \rightarrow Gx) \rightarrow \forall x \in U. Gx.$$

By predicate logic, it is equivalent to (22). \square

The next lemma shows induction is implied from induction for decreasing sequences.

Lemma 5.15 $\mathbf{HA} \vdash \text{Ind}(\text{DS}(U, >), >_{\text{ext}})$ implies $\mathbf{HA} \vdash \text{Ind}(U, >)$.

Proof. In this proof, for simplicity, we write DS for $\text{DS}(U, >)$.

Assume

$$\forall x \in U((\forall y < x \in U. Fy) \rightarrow Fx) \quad (23)$$

in order to show $\forall x \in U. Fx$.

Define $Gx \equiv F(\text{last}(x))$. We will show

$$\forall x \in \text{DS}((\forall z <_{\text{ext}} x \in \text{DS}. Gz) \rightarrow Gx). \quad (24)$$

Fix $x \in \text{DS}$ and assume

$$\forall z <_{\text{ext}} x \in \text{DS}. Gz \quad (25)$$

in order to show Gx .

We can show

$$\forall y < \text{last}(x) \in U. Fy \quad (26)$$

as follows. Assume $y < \text{last}(x)$ in order to show Fy . Then $x * \langle y \rangle \in \text{DS}$ and $x >_{\text{ext}} x * \langle y \rangle$. By taking z to be $x * \langle y \rangle$ in (25), we have $G(x * \langle y \rangle)$. By definition of G , $F(\text{last}(x * \langle y \rangle))$. Hence Fy . We have shown (26).

By taking x to be $\text{last}(x)$ in (23), we have $F(\text{last}(x))$. Hence Gx . Hence we have shown (24).

By $\text{Ind}(\text{DS}, >_{\text{ext}})$ with (24), we have $\forall x \in \text{DS}. Gx$. By taking x to be $\langle x \rangle$ in it, we have $G(\langle x \rangle)$. By definition of G , we have Fx . \square

The next lemma shows induction for a power of a relation implies induction for the relation.

Lemma 5.16 $\mathbf{HA} \vdash \text{Ind}(U, >^n)$ implies $\mathbf{HA} \vdash \text{Ind}(U, >)$.

Proof. We can assume $n > 1$.

Assume

$$\forall x \in U((\forall y < x \in U.Fy) \rightarrow Fx) \quad (27)$$

in order to show $\forall x \in U.Fx$.

We will show

$$\forall x \in U((\forall y <^n x \in U.Fy) \rightarrow Fx). \quad (28)$$

Fix $x \in U$ and assume

$$\forall y <^n x \in U.Fy \quad (29)$$

in order to show Fx .

By induction on m , we will show

$$\forall m \leq n. \forall w <^{n-m} x \in U.Fw \quad (30)$$

Case 1. $m = 0$.

Assume $w <^n x \in U$. By taking y to be w in (29), Fw .

Case 2. $m > 0$.

Assume $w <^{n-m} x \in U$ in order to show Fw . We can show $\forall y < w \in U.Fy$ as follows. Assume $y < w$. Then $y <^{n-(m-1)} x$. By IH for $m-1$, Fy .

By taking x to be w in (27), Fw .

We have shown (30). By taking m to be n and w to be x in it, Fx . Hence we have shown (28). By $\text{Ind}(U, >^n)$, $\forall x \in U.Fx$. \square

Define

$$\begin{aligned} \text{Trans}(U, >_R) &\equiv \forall xyz \in U(x >_R y \wedge y >_R z \rightarrow x >_R z), \\ \text{Decide}(U, >_R) &\equiv \forall xy \in U(x >_R y \vee \neg(x >_R y)). \end{aligned}$$

The next theorem states that if some powers of relations $>_{R_i}$ have induction principle, $>_{R_i}$ are decidable and their union is transitive, then the union has induction principle. This theorem is the same as Theorem 6.1 in [3] except **HA** and the decidability condition $\text{Decide}(U, >_{R_i})$.

Theorem 5.17 (Podelski-Rybalchenko Theorem for Induction) *If $\mathbf{HA} \vdash \text{Ind}(U, >_{R_1}^{n_1})$, $\mathbf{HA} \vdash \text{Decide}(U, >_{R_1})$, \dots , $\mathbf{HA} \vdash \text{Ind}(U, >_{R_k}^{n_k})$, $\mathbf{HA} \vdash \text{Decide}(U, >_{R_k})$, and $\mathbf{HA} \vdash \text{Trans}(U, >_{R_1+\dots+R_k})$, then $\text{Ind}(U, >_{R_1+\dots+R_k})$.*

Proof. We will discuss in **HA**.

By Lemma 5.16, $\text{Ind}(U, >_{R_i})$. Define $>_{R'_1}$ as $>_{R_1}$ and $>_{R'_{i+1}}$ as $(>_{R_{i+1}}) - (>_{R'_1}) - \dots - (>_{R'_i})$. Then $(>_{R'_1}), \dots, (>_{R'_k})$ are disjoint and $\forall xy \in U(x >_{R_1 \cup \dots \cup R_k} y \rightarrow x >_{R'_1+\dots+R'_k} y)$ by $\text{Decide}(U, >_{R_i})$ for $1 \leq i \leq k$. Since $(>_{R'_i}) \subseteq (>_{R_i})$, $\text{Ind}(U, >_{R'_i})$. For simplicity, from now on we write $>_{R_i}$ for $>_{R'_i}$ in this proof. We will show $\text{Ind}(U, >_{R_1+\dots+R_k})$.

From $\text{Ind}(U, >_{R_i})$, we have $\text{Ind}(\text{DT}(U, >_{R_i}), >_{\text{ext}})$ for $1 \leq i \leq k$. By Lemma 5.7, we have $\forall r \in U. \text{Ind}(\text{MS}_{\langle r \rangle}, >_{\text{ext}})$. Apparently $\forall u \in U. \text{Ind}(U, >_{u, \text{Left}})$. By taking U to be U , T to be $\text{MS}_{\langle r \rangle}$, and $>_u$ to be $>_{u, \text{Left}}$ in Theorem 5.3 for $>_{\text{KB1}, r}$, we have $\forall r \in U. \text{Ind}(\text{MS}_{\langle r \rangle}, >_{\text{KB1}, r})$. By Theorem 5.3 for $>_{\text{KB2}, r}$, we have $\forall r \in U. \text{Ind}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r})$ similarly. Hence $\forall r \in U. \text{Ind}(\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r}), >_{\text{ext}})$. From Lemma 5.14, we have $\forall r \in U. \text{Ind}(\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r})_{\langle \langle r \rangle \rangle}, >_{\text{ext}})$. By taking T to be $\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r})_{\langle \langle r \rangle \rangle}$, U to be $\text{MS}_{\langle r \rangle}$, and $(>_u)$ to be $(>_{\text{KB1}, r})$ in Theorem 5.3 for $>_{\text{KB}, r}$, we have $\forall r \in U. \text{Ind}(\text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r})_{\langle \langle r \rangle \rangle}, >_{\text{KB}, r})$. Since $\text{ET2}(\text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle}) \subseteq \text{DS}(\text{MS}_{\langle r \rangle}, >_{\text{KB2}, r})_{\langle \langle r \rangle \rangle}$, by Lemma 5.14, we have $\forall r \in U. \text{Ind}(\text{ET2}(\text{DT}(U, >_{R_1+\dots+R_k})_{\langle r \rangle}), >_{\text{KB}, r})$. By Lemma 5.13, $\text{Ind}(\text{DT}(U, >_{R_1+\dots+R_k}), >_{\text{ext}})$. By $\text{Trans}(U, >_{R_1+\dots+R_k})$, $\text{DT}(U, >_{R_1+\dots+R_k})$ is $\text{DS}(U, >_{R_1+\dots+R_k})$. Hence we have $\text{Ind}(\text{DS}(U, >_{R_1+\dots+R_k}), >_{\text{ext}})$. By Lemma 5.15, we have $\text{Ind}(U, >_{R_1+\dots+R_k})$. \square

6 Proof Transformation

This section defines main proof transformation from $\mathbf{CLJID}^\omega + \mathbf{HA}$ to $\mathbf{LJID} + \mathbf{HA}$. The proof is the same as [3] except Theorem 5.17 and Lemma 6.6. First we will define stage numbers and path relations, and then define proof transformation using them.

For notational convenience, we assume a cyclic proof Π in this section. Let the buds in Π be J_{1i} ($i \in I$) and the companions be J_{2j} ($j \in K$). Assume $f : I \rightarrow K$ such that the companion of a bud J_{1i} is $J_{2, f(i)}$.

$$\begin{array}{c}
\frac{Q_1 \vec{u}_1 \quad \dots \quad Q_n \vec{u}_n \quad P_1 \vec{t}_1 \quad \dots \quad P_m \vec{t}_m}{P \vec{t}} \\
\\
\frac{Q_1 \vec{u}_1 \quad \dots \quad Q_n \vec{u}_n \quad v > v_1 \quad P'_1 \vec{t}_1 v_1 \quad \dots \quad v > v_m \quad P'_m \vec{t}_m v_m \quad Nv}{P' \vec{t} v}
\end{array}$$

Figure 1: Production Rules

6.1 Stage Numbers for Inductive Definitions

In this subsection, we define and discuss stage transformation.

We introduce a stage number to each inductive atomic formula so that the argument of the formula comes into the inductive predicate at the stage of the stage number. This stage number will decrease by a progressing trace. A proof in **LJID** + **HA** will be constructed by using the induction on stage numbers.

First we give stage transformation of an inductive atomic formula. We assume a fresh inductive predicate symbol P' for each inductive predicate symbol P and we call it a *stage-number inductive predicate symbol*. $P'(\vec{t}, v)$ means that the element \vec{t} comes into P at the stage v . We transform $P(\vec{t})$ into $\exists v P'(\vec{t}, v)$. We call a variable v a *stage number* of \vec{t} when $P'(\vec{t}, v)$. $P(\vec{t})$ and $\exists v P'(\vec{t}, v)$ will become equivalent by inference rules introduced by the transformation of production rules. We call $P'(\vec{t}, v)$ a *stage-number inductive atomic formula*.

Secondly we give stage transformation of a production rule. We transform the production of P (the first rule) in Figure 1 into the production of P' (the second rule) in Figure 1, where v, v_1, \dots, v_m are fresh variables.

Next we give stage transformation of a sequent. For given fresh variables \vec{v} , we transform a sequent J into $J_{\vec{v}}^{\circ}$ defined as follows. We define Γ^{\bullet} as the sequent obtained from Γ by replacing $P(\vec{t})$ by $\exists v P'(\vec{t}, v)$. For fresh variables \vec{v} , we define $(\Gamma)_{\vec{v}}^{\circ}$ as the sequent obtained from Γ^{\bullet} by replacing the i -th element of the form $\exists v P'(\vec{t}, v)$ in the sequent Γ^{\bullet} by $P'(\vec{t}, v_i)$. We define $(\Gamma \vdash \Delta)^{\bullet}$ by $\Gamma^{\bullet} \vdash \Delta^{\bullet}$, and define $(\Gamma \vdash \Delta)_{\vec{v}}^{\circ}$ by $(\Gamma)_{\vec{v}}^{\circ} \vdash \Delta^{\bullet}$.

We write $(a_i)_{i \in I}$ for the sequence of elements a_i where i varies in I . We extend the notion of proofs by allowing open assumptions. We write $\Gamma \vdash_{\text{CLJID}^{\omega} + \text{HA}} \Delta$ with assumptions $(J_i)_{i \in I}$ when there is a proof with assumptions $(J_i)_{i \in I}$ and the conclusion $\Gamma \vdash \Delta$ in **CLJID**^ω + **HA**.

Definition 6.1 In a path π in a proof, we define $\text{Ineq}(\pi)$ as the set of the forms $v > v'$ and $v = v'$ for any stage numbers v, v' eliminated by every case distinction in π .

The next lemma shows a stage number is a number.

Lemma 6.2 (1) $P' \vec{t} v \vdash_{\text{CLJID}^{\omega} + \text{HA}} Nv$.
(2) $P' \vec{t} v \vdash_{\text{LJID} + \text{HA}} Nv$.

Proof. (1) and (2) are proved by (Case P') and (Ind P') respectively. \square

An example for $\text{Ineq}(\pi)$ is as follows. When the path π has the case distinction

$$\begin{array}{c}
\Gamma^{\circ}, u = t(y), \hat{v} = v, Q_1 u_1(y), \dots, Q_n u_n(y), \\
v > v_1, P'_1 t_1(y) v_1, \dots, v > v_m, P'_m t_m(y) v_m, \\
Nv \vdash \Delta^{\bullet}
\end{array}$$

we take $\hat{v} = v, v > v_1, \dots, v > v_m$ for $\text{Ineq}(\pi)$.

The proof of the next proposition gives *stage transformation* of a proof into a proof of the stage transformation of the conclusion of the original proof. We write Π° for the stage transformation of Π .

Proposition 6.3 (Stage Transformation) For any fresh variables \vec{v} , $\Gamma \vdash_{\text{CLJID}^{\omega} + \text{HA}} \Delta$ with assumptions $(\Gamma_i \vdash \Delta_i)_{i \in I}$ without any buds implies $(\Gamma)_{\vec{v}}^{\circ} \vdash_{\text{CLJID}^{\omega} + \text{HA}} \Delta^{\bullet}$ with assumptions $(\text{Ineq}(\pi_i), (\Gamma_i)_{\vec{v}_i}^{\circ} \vdash \Delta_i^{\bullet})_{i \in I}$ without any buds for some fresh variables $(\vec{v}_i)_{i \in I}$, where π_i is the path from the conclusion to the assumption $(\Gamma_i)_{\vec{v}_i}^{\circ} \vdash \Delta_i^{\bullet}$.

Proof. By induction on the proof. We will transform the proof Π of $\Gamma \vdash_{\mathbf{CLJID}^\omega + \mathbf{HA}} \Delta$ with assumptions $(J_i)_{i \in I}$ into the proof of $\Gamma \xrightarrow{\circ} \vdash_{\mathbf{CLJID}^\omega + \mathbf{HA}} \Delta^\bullet$ with assumptions $(\text{Ineq}(\pi_i), (J_i)_{\xrightarrow{\circ}}^{\circ})_{i \in I}$ by transforming each rule as follows.

Case 1. The rule is not $(P_i \text{ R})$, (Case), (Cut), (Axiom) with a common inductive atomic formula, or logical rules with some of the main formula and the auxiliary formulas being an inductive atomic formula in the antecedent.

Since the main formula and the auxiliary formulas are not inductive atomic formulas in the antecedent, we can just replace each sequent J by $(J)_{\xrightarrow{\circ}}^{\circ}$. If the rule is an assumption, since $I = \{1\}$, we take \vec{v}_1 to be \vec{v} .

For example,

$$\frac{\Gamma \vdash P_1 t_1}{\Gamma \vdash P_1 t_1 \vee P_2 t_2} (\vee R_l)$$

is transformed into

$$\frac{(\Gamma)_{\xrightarrow{\circ}}^{\circ} \vdash \exists v_1 P'_1 t_1 v_1}{(\Gamma)_{\xrightarrow{\circ}}^{\circ} \vdash \exists v_1 P'_1 t_1 v_1 \vee \exists v_2 P'_2 t_2 v_2} (\vee R_l)$$

Case 2. The rule is (Axiom) with a common inductive atomic formula.

We transform

$$\frac{}{\Gamma, P(\vec{t}) \vdash P(\vec{t})} (\text{Axiom})$$

into

$$\frac{\frac{}{\Gamma \xrightarrow{\circ} \vdash, P'(\vec{t}, v) \vdash P'(\vec{t}, v)} (\text{Axiom})}{\Gamma \xrightarrow{\circ} \vdash, P'(\vec{t}, v) \vdash \exists v P'(\vec{t}, v)} (\exists R)$$

Case 3. The rule is (Cut), or logical rules with some of the main formula and the auxiliary formulas being an inductive atomic formula in the antecedent.

We replace each sequent J by $(J)_{\xrightarrow{\circ}}^{\circ}$ and use

$$\frac{\Gamma \xrightarrow{\circ} \vdash, P'(\vec{t}, v) \vdash \Delta^\bullet}{\Gamma \xrightarrow{\circ} \vdash, \exists v P'(\vec{t}, v) \vdash \Delta^\bullet} (\exists L)$$

Note that by IH we can assume each $P'(\vec{t}, v)$ has a fresh v , so we can use $(\exists L)$. For example, we transform

$$\frac{\Gamma, P t \vdash}{\Gamma \vdash \neg P t} (\neg R)$$

into

$$\frac{\frac{\Gamma \xrightarrow{\circ} \vdash, P' t v \vdash}{\Gamma \xrightarrow{\circ} \vdash, \exists v P' t v \vdash} (\exists L)}{\Gamma \xrightarrow{\circ} \vdash \neg \exists v P' t v} (\neg R)$$

Case 4. The rule is $(P \text{ R})$. Assume the production rule of P and its stage transformation in Figure 1. Let Σ_0 be the set of all assumptions of this production rule of P' and t_0 be $v_1 + \dots + v_n + 1$. By $(P' \text{ R})$ for this production rule,

$$\Sigma_0 \vdash P' \vec{t} v.$$

Since $P_i \vec{t}_i v_i \vdash N v_i$, $(\Sigma_1)_{\xrightarrow{\circ}}^{\circ} \vdash N v_i$ where Σ_1 is $Q_1 \vec{u}_1, \dots, Q_m \vec{u}_m, P_1 \vec{t}_1, \dots, P_n \vec{t}_n$ and \vec{v} is $v_1 \dots v_n$. By (Subst) with $v := t_0$, and (Cut) with $\vdash t_0 > v_i$ and $(\Sigma_1)_{\xrightarrow{\circ}}^{\circ} \vdash N t_0$,

$$(\Sigma_1)_{\xrightarrow{\circ}}^{\circ} \vdash P' \vec{t} t_0.$$

By $(\exists R)$, then by $(\exists L)$, $(\Sigma_1)^\bullet \vdash \exists v P' \vec{t} v$.

By IH,

$$\begin{aligned} \Gamma_{\vec{v}}^\circ &\vdash Q_j \vec{u}_j, \\ \Gamma_{\vec{v}}^\circ &\vdash \exists v_i P'_i \vec{t}_i v_i. \end{aligned}$$

By (Cut), $\Gamma_{\vec{v}}^\circ \vdash \exists v P' \vec{t} v$.

Case 5. The rule is (Case P). Assume the production rule of P and its stage transformation in Figure

1. Let \vec{v} be $\vec{v}'\hat{v}$. Let v_1, \dots, v_m be fresh variables. Let the rule be

$$\frac{\text{case distinctions}}{\Gamma, P \vec{u} \vdash \Delta} \text{ (Case } P\text{)}$$

with the case distinctions

$$\begin{aligned} \Gamma, \vec{u} = \vec{t}(\vec{y}), Q_1 \vec{u}_1(\vec{y}), \dots, Q_n \vec{u}_n(\vec{y}), \\ P_1 \vec{t}_1(\vec{y}), \dots, P_m \vec{t}_m(\vec{y}) \vdash \Delta. \end{aligned}$$

Let

$$\begin{aligned} \Sigma &\equiv (\vec{u} = \vec{t}(\vec{y}), Q_1 \vec{u}_1(\vec{y}), \dots, Q_n \vec{u}_n(\vec{y}), \\ &\quad P_1 \vec{t}_1(\vec{y})v_1, \dots, P_m \vec{t}_m(\vec{y})v_m), \\ \Sigma_1 &\equiv (\hat{v} = v, v > v_1, \dots, v > v_m). \end{aligned}$$

By IH with $\vec{v}'v_1 \dots v_m$ for the case distinction we obtain a proof of

$$\Gamma_{\vec{v}'}^\circ, \Sigma \vdash \Delta^\bullet$$

with assumptions $(\text{Ineq}(\pi'_i), (J_i)_{\vec{v}'_i}^\circ)_{i \in I}$ for some $(\vec{v}'_i)_{i \in I}$ where π'_i is the path from the transformation of the case distinction to $(J_i)_{\vec{v}'_i}^\circ$. Add Σ_1 to the antecedent of every sequent in the path π'_i , by renaming fresh variables to keep freshness of all fresh variables in the path. Then we have a proof of

$$\Sigma_1, \Gamma_{\vec{v}'}^\circ, \Sigma \vdash \Delta^\bullet$$

with assumptions $(\Sigma_1, \text{Ineq}(\pi'_i), (J_i)_{\vec{v}'_i}^\circ)_{i \in I}$. We have

$$\frac{\text{case distinctions}}{\Gamma_{\vec{v}'}^\circ, P' \vec{u} \hat{v} \vdash \Delta^\bullet} \text{ (Case } P'\text{)}$$

with the case distinctions

$$\Gamma_{\vec{v}'}^\circ, \Sigma_1, \Sigma, Nv \vdash \Delta^\bullet.$$

By (Wk) with Nv we obtain the case distinction. By (Case P'),

$$\Gamma_{\vec{v}'}^\circ, P' \vec{u} \hat{v} \vdash \Delta^\bullet.$$

Let π_i be the path from $\Gamma^\circ, P' \vec{u} \hat{v} \vdash \Delta^\bullet$ to $(J_i)_{\vec{v}'_i}^\circ$. Then

$$\text{Ineq}(\pi_i) = (\Sigma_1, \text{Ineq}(\pi'_i)).$$

Hence we have a proof of

$$\Gamma_{\vec{v}'}^\circ, P' \vec{u} \hat{v} \vdash \Delta^\bullet$$

with assumptions $(\text{Ineq}(\pi_i), (J_i)_{\vec{v}'_i}^\circ)_{i \in I}$. \square

6.2 Path Relation

In this section, we will introduce path relations and discuss them.

We assume a subproof Π_j of Π such that it does not have buds, its conclusion is J_{2j} and its assumptions are J_{1i} ($i \in I_j$).

For J in Π_j° , we define \tilde{J} as $\langle v_1, \dots, v_k \rangle$ where J is $\Gamma_{v_1 \dots v_k}^\circ \vdash \Delta^\bullet$.

For a path π from the conclusion to an assumption in Π_j° , we write $\tilde{\pi}$ for the corresponding path in Π . We extend this notation to a finite composition of π 's. By the correspondence (\sim), a stage-number inductive atomic formula in Π_j° corresponds to an inductive atomic formula in Π , and a path, a trace, and a progressing trace in Π_j° correspond to the same kind of objects in Π .

Definition 6.4 For a finite composition π of paths in $\{\Pi_j^\circ \mid j \in K\}$ such that $\tilde{\pi}$ is a path in the infinite unfolding of Π , we define the *path relation* $\tilde{>}_\pi$ by

$$x \tilde{>}_\pi y \equiv |x| = |\tilde{J}_2| \wedge |y| = |\tilde{J}_1| \wedge \bigwedge_{F(q_1, q_2)} (x)_{q_2} > (y)_{q_1} \wedge \bigwedge_{G(q_1, q_2)} (x)_{q_2} = (y)_{q_1}$$

where J_1 and J_2 are the top and bottom sequents of π respectively, \tilde{J}_1 and \tilde{J}_2 are those of the path $\tilde{\pi}$, $F(q_1, q_2)$ is that there is some progressing trace from the q_2 -th atomic formula in \tilde{J}_2 to the q_1 -th atomic formula in \tilde{J}_1 , $G(q_1, q_2)$ is that there is some non-progressing trace from the q_2 -th atomic formula in \tilde{J}_2 to the q_1 -th atomic formula in \tilde{J}_1 .

We define B_1 as the set of paths from conclusions to assumptions in Π_j° ($j \in K$). We define B as the set of finite compositions of elements in B_1 such that if $\pi \in B$ then $\tilde{\pi}$ is a path in the infinite unfolding of Π .

Definition 6.5 For $\pi \in B$, define $x >_\pi y$ by

$$x >_\pi y \equiv (x)_0 = j \wedge (y)_0 = f(i) \wedge (x)_1 \tilde{>}_\pi (y)_1,$$

where J_{1i} is the top sequent of $\tilde{\pi}$, and J_{2j} is the bottom sequent of $\tilde{\pi}$.

Note that $(\)_0$ and $(\)_1$ are operations for a number that represents a sequence of numbers defined in Section 3. The first element is a companion number.

Lemma 6.6 $\{>_\pi \mid \pi \in B\}$ is finite.

Proof. Define C_n as $\{>_{\pi_1 \dots \pi_m} \mid m \leq n, \pi_i \in B_1\}$. Since $>_\pi$ is a relation on $N \times N^{\leq p}$ where p is the maximum number of inductive atomic formulas in the antecedents of Π , there is L such that $|C_n| \leq L$ for all n . Then we have the least n such that $C_{n+1} = C_n$. Then $|\{>_\pi \mid \pi \in B\}| = |C_n|$. \square

The next proposition shows a sequent is implied from its stage-number transformation.

Proposition 6.7 For any fresh variables \vec{v} , $\Gamma_{\vec{v}}^\circ \vdash_{\text{LJID+HA}} \Delta^\bullet$ implies $\Gamma \vdash_{\text{LJID+HA}} \Delta$.

Proof. First we can show $\Gamma^\bullet \vdash_{\text{LJID+HA}} \Delta^\bullet$ by

$$\frac{P' \vec{t} v, \Gamma^\bullet \vdash \Delta^\bullet}{\exists v P' \vec{t} v, \Gamma^\bullet \vdash \Delta^\bullet} (\exists L)$$

Secondly by

$$\vdash_{\text{LJID+HA}} P \vec{t} \leftrightarrow \exists v P'(\vec{t}, v)$$

we have

$$\Gamma \vdash_{\text{LJID+HA}} \Delta.$$

\square

The next lemma shows $\tilde{>}_\pi$ is an abstraction of $\text{Ineq}(\pi)$.

Lemma 6.8 For a proof Π without any buds, if π is a path from $(J_2)_{\vec{x}}^\circ$ to $(J_1)_{\vec{y}}^\circ$ in Π° ,

$$\text{Ineq}(\pi) \vdash_{\text{HA}} \langle \vec{x} \rangle \tilde{>}_\pi \langle \vec{y} \rangle.$$

Proof. By induction on $|\pi|$.

Case 1. $|\pi| = 0$. $J_1 = J_2$. $x \tilde{>}_\pi y$ is $|x| = m \wedge |y| = m \wedge (x)_0 = (y)_0 \wedge \dots \wedge (x)_{m-1} = (y)_{m-1}$ where m is the number of stage-number inductive atomic formulas in $(J_1)^\circ_{\vec{y}}$. Hence $\vdash_{\mathbf{HA}} x \tilde{>}_\pi y$.

Case 2. $|\pi| > 0$. Consider cases according to the last rule of π . Let $\pi = \pi_1 \pi_2$ such that $|\pi_1| = 1$. Let the top sequent of π_1 be $(J_3)^\circ_{\vec{z}}$. Let x, y, z be $\langle \vec{x} \rangle, \langle \vec{y} \rangle, \langle \vec{z} \rangle$ respectively.

By IH, $\text{Ineq}(\pi_2) \vdash_{\mathbf{HA}} z \tilde{>}_{\pi_2} y$.

We will show $\text{Ineq}(\pi_1) \vdash_{\mathbf{HA}} x \tilde{>}_{\pi_1} z$. Since the rule that changes the stage number is only (Case), we will show only the case (Case). Assume the production rule of P and its stage transformation in Figure 1. Let the path for the rule (Case P') be

$$\frac{\Gamma^\circ_{\vec{y}}, \Sigma_1, \Sigma, Nv \vdash \Delta^\bullet}{\Gamma^\circ_{\vec{y}}, P' \vec{u} \hat{v} \vdash \Delta^\bullet}$$

where

$$\begin{aligned} \Sigma &\equiv (\vec{u} = \vec{t}(\vec{y}), Q_1 \vec{u}_1(\vec{y}), \dots, Q_n \vec{u}_n(\vec{y}), \\ &\quad P'_1 \vec{t}_1(\vec{y})v_1, \dots, P'_m \vec{t}_m(\vec{y})v_m), \\ \Sigma_1 &\equiv (\hat{v} = v, v > v_1, \dots, v > v_m). \end{aligned}$$

Then $x = \langle \vec{v}, \hat{v} \rangle$ and $z = \langle \vec{v}, v_1, \dots, v_m \rangle$, and

$$x \tilde{>}_{\pi_1} z \leftrightarrow v_1 < \hat{v} \wedge \dots \wedge v_m < \hat{v}.$$

Since $\text{Ineq}(\pi_1) \equiv \Sigma_1$, $\text{Ineq}(\pi_1) \vdash_{\mathbf{HA}} x \tilde{>}_{\pi_1} z$. Since $\text{Ineq}(\pi) = (\text{Ineq}(\pi_1), \text{Ineq}(\pi_2))$, $\text{Ineq}(\pi) \vdash_{\mathbf{HA}} x \tilde{>}_{\pi_1} z \tilde{>}_{\pi_2} y$. Since $(\tilde{>}_{\pi_1}) \circ (\tilde{>}_{\pi_2}) \subseteq (\tilde{>}_{\pi_1 \pi_2})$, $\text{Ineq}(\pi) \vdash_{\mathbf{HA}} x \tilde{>}_{\pi_1 \pi_2} y$. \square

The next lemma is the only lemma that uses the global trace condition.

Lemma 6.9 *For all $\pi \in B$, there is $n > 0$ such that $\vdash_{\mathbf{HA}} \text{Ind}(U, >_\pi^n)$.*

Proof. Let the bottom sequent of π be J_{2j} and the top sequent be J_{1i} . Let the companion of J_{1i} be J_{2k} .

Case 1. $j \neq k$.

Assume

$$H \equiv \forall x \in U. (\forall y <_\pi x \in U. Fy) \rightarrow Fx$$

and fix $x \in U$.

Assume $y <_\pi x \in U$. By taking x to be y in H ,

$$(\forall z <_\pi y \in U. Fz) \rightarrow Fy.$$

By $\neg(z <_\pi y)$ from $y <_\pi x$ and $j \neq k$, we have Fy . Hence $H \vdash_{\mathbf{HA}} \forall y <_\pi x \in U. Fy$.

By taking x to be x in H , we have Fx . Hence $H \vdash_{\mathbf{HA}} \forall x \in U. Fx$. Hence $\vdash_{\mathbf{HA}} \text{Ind}(U, >_\pi, F)$. We can take n to be 1.

Case 2. $j = k$.

By applying the global trace condition to the infinite path $\tilde{\pi}^\omega$, there is a progressing trace in the path. Hence there are n, m, q such that the trace passes the q -th stage-number inductive atomic formula in the top sequent of π^m and the q -th stage-number inductive atomic formula in the top sequent of π^{m+n} .

Define $x <_q y$ by $((x)_1)_q < ((y)_1)_q$. By mathematical induction we can easily show

$$(\forall x \in U. (\forall y <_q x \in U. Fy) \rightarrow Fx) \rightarrow \forall x \in U. Fx.$$

If $y <_\pi^n x$, then $y <_{\pi^n} x$, and hence $((y)_1)_q < ((x)_1)_q$, which implies $y <_q x$. Therefore

$$(\forall x \in U. (\forall y <_{\pi^n} x \in U. Fy) \rightarrow Fx) \rightarrow \forall x \in U. Fx,$$

which is $\text{Ind}(U, >_{\pi^n}, F)$. \square

We define $>_\Pi$ as $\bigcup \{>_\pi \mid \pi \in B\}$. Note that $>_\Pi$ is transitive, since the top sequent of π_1 is the bottom sequent of π_2 by the first element, and $((>_{\pi_1}) \circ (>_{\pi_2})) \subseteq (>_{\pi_1 \pi_2})$.

6.3 Proof Transformation

This section gives main proof transformation.

The next lemma shows we can replace (Case) rules of $\mathbf{CLJID}^\omega + \mathbf{HA}$ by (Ind) rules of $\mathbf{LJID} + \mathbf{HA}$.

Lemma 6.10 *If there is a proof with some assumptions and without any buds in $\mathbf{CLJID}^\omega + \mathbf{HA}$, then there is a proof of the same conclusions with the same assumptions in $\mathbf{LJID} + \mathbf{HA}$.*

Proof. It is sufficient to replace the rule (Case) by the rule (Ind). This is straightforward and has been proved by Lemma 4.1.4 in [5]. We give only a proof idea here.

Assume the production rule of P in Figure 1. We can replace

$$\frac{\begin{array}{c} \vdots \Pi_i \\ \Gamma, \vec{u} = \vec{t}(\vec{y}), \Sigma \vdash \Delta \quad (\forall i \in I) \end{array}}{\Gamma, P\vec{u} \vdash \Delta} \text{ (Case)}$$

by

$$\frac{\begin{array}{c} \vdots \Pi'_i \\ \Sigma' \vdash F\vec{t}(\vec{y}) \quad (\forall i \in I) \end{array}}{\frac{P\vec{u} \vdash F\vec{u}}{\Gamma, P\vec{u} \vdash \Delta} \text{ (Ind } P)} \text{ (}\rightarrow L\text{)}(\forall L)(Cut)$$

where

$$\begin{aligned} F'\vec{z} &\equiv \forall \vec{x}(\vec{z} = \vec{u} \rightarrow \Gamma \rightarrow \Delta), \\ F\vec{z} &\equiv P\vec{z} \wedge F'\vec{z}. \end{aligned}$$

□

The next is the main lemma and shows each bud in a cyclic proof is provable in $\mathbf{LJID} + \mathbf{HA}$.

Lemma 6.11 *For every bud J of a proof in $\mathbf{CLJID}^\omega + \mathbf{HA}$ and fresh variables \vec{v} , $(J)_{\vec{v}}^\circ$ is provable in $\mathbf{LJID} + \mathbf{HA}$.*

Proof. For $j \in K$, define

$$G_j x \equiv \forall \vec{z}(x = \langle \vec{v} \rangle \rightarrow (\Gamma_{2j})_{\vec{v}}^\circ \rightarrow \Delta_{2j}^\bullet)$$

where \vec{z} is $\text{FV}((\Gamma_{2j})_{\vec{v}}^\circ, \Delta_{2j}^\bullet)$.

Define

$$\begin{aligned} Gx_0x &\equiv \bigwedge_{j \in K} (x_0 = j \rightarrow G_j x), \\ Hx_0x &\equiv \forall y_0 y. \langle y_0, y \rangle <_\Pi \langle x_0, x \rangle \rightarrow Gx_0y. \end{aligned}$$

We will show that for every companion J in Π and fresh variables \vec{v} , $(J)_{\vec{v}}^\circ$ is provable in $\mathbf{LJID} + \mathbf{HA}$. Fix a companion in Π and fresh variables \vec{v} . Let the companion be J_{2j} . We will construct a proof of $(J_{2j})_{\vec{v}}^\circ$ in $\mathbf{LJID} + \mathbf{HA}$. We have a subproof Π_j of Π such that it does not have buds, its conclusion is J_{2j} and its assumptions are J_{1i} ($i \in I_j$). By Proposition 6.3,

$$\begin{array}{c} (\text{Ineq}(\pi_{ji}), (\Gamma_{1i})_{\vec{v}}^\circ \vdash \Delta_{1i}^\bullet)_{i \in I_j} \\ \vdots \Pi_j^\circ \\ (\Gamma_{2j})_{\vec{v}}^\circ \vdash \Delta_{2j}^\bullet \end{array}$$

for some fresh variables $(\vec{v}_i)_{i \in I_j}$, where π_{ji} is the path from J_{2j} to J_{1i} in Π_j . Next we transform Π_j° into the $(\mathbf{LJID} + \mathbf{HA})$ -proof Π'_j with the same conclusion and the same assumptions by Lemma 6.10.

Next we add

$$x_0 = j, x = \langle \vec{v} \rangle, Hx_0x$$

$$\begin{array}{c}
\text{Lemma 6.8} \\
\hline
x = \langle \vec{v} \rangle, y = \langle \vec{v}_i \rangle, \text{Ineq}(\pi_{ji}) \vdash x \succ_{\pi_{ji}} y \quad (Wk)(= L) \quad \frac{x_0 = j, y_0 = f(i), \langle x_0, x \rangle \succ_{\pi_{ji}} \langle y_0, y \rangle, Hx_0x \vdash Gy_0y}{x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, \text{Ineq}(\pi_{ji}) \vdash Gf(i)\langle \vec{v}_i \rangle} \quad \frac{(\rightarrow L)(\forall L)(Cut)}{(Cut)(Subst)} \\
\hline
\frac{x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, \text{Ineq}(\pi_{ji}) \vdash Gf(i)\langle \vec{v}_i \rangle}{x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, \text{Ineq}(\pi_{ji}), (\Gamma_{1i})_{\vec{v}_i}^{\circ} \vdash \Delta_{1i}^{\bullet}} \quad (\rightarrow L)(\forall L)(Cut)
\end{array}$$

Figure 2: Subproof Π_{ij}

$$\begin{array}{c}
\left(\begin{array}{c} \vdots \Pi_{ij} \\ x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, \text{Ineq}(\pi_{ji}), (\Gamma_{1i})_{\vec{v}_i}^{\circ} \vdash \Delta_{1i}^{\bullet} \end{array} \right)_{i \in I_j} \\
\vdots \Pi_j'' \\
\frac{x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, (\Gamma_{2j})_{\vec{v}}^{\circ} \vdash \Delta_{2j}^{\bullet}}{Hx_0x \vdash x_0 = j \rightarrow G_jx} \quad (\rightarrow R)(\forall R)(\rightarrow R)(\rightarrow R) \\
\frac{(\forall j \in K) \quad \frac{(\rightarrow R)(\forall R)(Cut)}{Hx_0x \vdash Gx_0x} \quad (\wedge R)}{\vdash \text{Ind}(>_{\Pi}, G)} \quad \frac{(\rightarrow R)(\forall R)(Cut)}{\vdash \forall x_0x. Gx_0x}
\end{array}$$

Figure 3: Proof Π'''

to every antecedent in Π'_j to obtain

$$\begin{array}{c}
(x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, \text{Ineq}(\pi_{ji}), (\Gamma_{1i})_{\vec{v}_i}^{\circ} \vdash \Delta_{1i}^{\bullet})_{i \in I_j} \\
\vdots \Pi_j'' \\
x_0 = j, x = \langle \vec{v} \rangle, Hx_0x, (\Gamma_{2j})_{\vec{v}}^{\circ} \vdash \Delta_{2j}^{\bullet}
\end{array}$$

Let Π_{ij} be the proof in Figure 2. We have a proof Π''' of $\forall x_0x. Gx_0x$ with the assumption $\text{Ind}(>_{\Pi}, G)$ in Figure 3.

By applying Theorem 5.17 to $U = N \times N^{\leq p}$ and $\{>_{\pi} \mid \pi \in B\}$,

$$\mathbf{HA} \vdash \text{Ind}(U, >_{\pi}^{n_{\pi}}) \quad (\forall \pi \in B)$$

implies

$$\mathbf{HA} \vdash \text{Ind}(U, >_{\Pi}).$$

By Lemma 6.9, we have

$$\vdash_{\mathbf{HA}} \text{Ind}(U, >_{\pi}^{n_{\pi}}) \quad (\forall \pi \in B).$$

By the definition of $>_{\pi}$,

$$\vdash_{\mathbf{HA}} \text{Decide}(U, >_{\pi})$$

for all $\pi \in B$.

Then

$$\begin{array}{c}
\text{Lemma 6.9} \quad \vdash \text{Decide}(U, >_{\pi}) \quad \text{Theorem 5.17} \\
\hline
\vdash \text{Ind}(U, >_{\Pi}, G) \\
\vdots \Pi''' \\
\vdash \forall x_0x. Gx_0x \quad (\forall L)(Cut) \\
\vdash Gj\langle \vec{v} \rangle \quad (\rightarrow L)(\wedge L)(Cut) \\
\vdash Gj\langle \vec{v} \rangle \quad (\rightarrow L)(\forall L)(Cut) \\
(J_{2j})_{\vec{v}}^{\circ}
\end{array}$$

We have shown that for every companion J in Π and fresh variables \vec{v} , $(J)_{\vec{v}}^{\circ}$ is provable in $\mathbf{LJID} + \mathbf{HA}$.

Fix a bud be J_{1k} in Π and fresh variables \vec{v} . Let J_{2j} be the companion of the bud. Since $(J_{2j})_{\vec{v}}^{\circ}$ is provable, $(J_{1k})_{\vec{v}}^{\circ}$ is provable in **LJID** + **HA**. \square

We write **LJID** + **HA** + (Σ, Φ) for the system **LJID** + **HA** with the signature Σ and the set Φ of production rules. Similarly we write **CLJID** $^{\omega}$ + **HA** + (Σ, Φ) . For simplicity, in Φ we write only P for the set of production rules for P . We define $\Sigma_N = \{0, s, +, \times, <, N\}$ and $\Phi_N = \{N\}$. We write P'' for $(P')'$.

The next is the main proposition stating a cyclic proof is transformed into an (**LJID** + **HA**)-proof with stage-number inductive predicates.

Proposition 6.12 *If a sequent J is provable in **CLJID** $^{\omega}$ + **HA** + $(\Sigma_N \cup \{\vec{P}\}, \Phi_N \cup \{\vec{P}\})$, then J is provable in **LJID** + **HA** + $(\Sigma_N \cup \{N', \vec{P}, \vec{P}'\}, \Phi_N \cup \{\vec{P}, \vec{P}'\})$ where N', \vec{P}' are the stage-number inductive predicates of N, \vec{P} .*

Proof. Let Π_1 be the cyclic proof of $\Gamma \vdash_{\mathbf{CLJID}^{\omega} + \mathbf{HA}} \Delta$. Let $(J_{1i})_{i \in I}$ be all the buds in Π_1 . Define Π_2 be a proof obtained from Π_1 by removing all bud-companion relations. Then Π_2 is a proof of $\Gamma \vdash \Delta$ with assumptions $(J_{1i})_{i \in I}$ and without buds in **CLJID** $^{\omega}$ + **HA**. Choose fresh variables \vec{v} . By Proposition 6.3, there is $(\vec{v}_i)_{i \in I}$ such that $\Gamma_{\vec{v}}^{\circ} \vdash \Delta^{\bullet}$ with assumptions $(\text{Ineq}(\pi_i), (J_{1i})_{\vec{v}_i}^{\circ})_{i \in I}$ and without buds in **CLJID** $^{\omega}$ + **HA**. By Lemma 6.10, $\Gamma_{\vec{v}}^{\circ} \vdash \Delta^{\bullet}$ with assumptions $(\text{Ineq}(\pi_i), (J_{1i})_{\vec{v}_i}^{\circ})_{i \in I}$ and without buds is provable in **LJID** + **HA**. By (Wk), there is a proof Π_3 of $\Gamma_{\vec{v}}^{\circ} \vdash \Delta^{\bullet}$ with assumptions $((J_{1i})_{\vec{v}_i}^{\circ})_{i \in I}$ in **LJID** + **HA**. By Lemma 6.11, $(J_{1i})_{\vec{v}_i}^{\circ}$ is provable in **LJID** + **HA**. Combining them with Π_3 , $\Gamma_{\vec{v}}^{\circ} \vdash \Delta^{\bullet}$ is provable in **LJID** + **HA**. By Proposition 6.7, $\Gamma \vdash \Delta$ is provable in **LJID** + **HA**. \square

The next shows conservativity for stage-number inductive predicates.

Proposition 6.13 (Conservativity of N' and P'') *Let $\Sigma = \Sigma_N \cup \{\vec{Q}, \vec{P}, \vec{P}'\}$, $\Phi = \Phi_N \cup \{\vec{P}, \vec{P}'\}$, $\Sigma' = \Sigma \cup \{N', \vec{P}''\}$, and $\Phi' = \Phi \cup \{N', \vec{P}''\}$. Then **LJID** + **HA** + (Σ', Φ') is conservative over **LJID** + **HA** + (Σ, Φ) .*

Proof. Define $\widetilde{(\quad)}$ by replacing N' by $\lambda xy. Nx \wedge Ny \wedge x \leq y$ and replacing P_j'' by $\lambda \vec{x} yz. P_j' \vec{x} y \wedge Nz \wedge y < z$ for all j .

By induction on a proof, we will show that **LJID** + **HA** + $(\Sigma', \Phi') \vdash J$ implies **LJID** + **HA** + $(\Sigma, \Phi) \vdash \widetilde{J}$.

Use case analysis by the last rule. If the last rule is not (Ind N') or (Ind P_j''), the claim follows immediately from IH.

Case 1. The last rule is (Ind N').

Let the rule be

$$\frac{\Gamma, Nv \vdash F0v \quad \Gamma, Fxv_1, v_1 < v, Nv \vdash Fsxv}{\Gamma, N'ab \vdash Fab} \text{ (Ind } N')$$

By IH for these two premises, we have

$$\widetilde{\Gamma}, Nv \vdash \widetilde{F0v} \tag{31}$$

and

$$\widetilde{\Gamma}, \widetilde{F}xv_1, v_1 < v, Nv \vdash \widetilde{F}sxv. \tag{32}$$

Define $F'a$ be $\forall z \in N. (a \leq z \rightarrow \widetilde{F}az)$. By (Ind N) we will show $\widetilde{\Gamma}, Na \vdash F'a$.

We can show the first premise $\widetilde{\Gamma} \vdash F'0$, since it immediately follow from (31).

We can show the second premise $\widetilde{\Gamma}, Nx, F'x \vdash F'sx$ as follows. Assume $\widetilde{\Gamma}, Nx, F'x$ and fix $z \in N$ and assume $sx \leq z$ in order to show $\widetilde{F}sxz$. Then there is $z' \in N$ such that $z = sz'$. Then $x \leq z'$. By $F'x$, $\widetilde{F}xz'$. By taking v_1 to be z' and v to be z in (32), $\widetilde{F}sxz$.

By (Ind N), we have $\widetilde{\Gamma}, Na \vdash F'a$. Hence $\widetilde{\Gamma}, Na, Nb, a \leq b \vdash \widetilde{F}ab$. Hence $(\widetilde{\Gamma}, \widetilde{N'ab}) \vdash \widetilde{F'ab}$.

Case 2. The last rule is (Ind P_j'').

Let the rule be

$$\frac{J_i \quad (\forall i \in I)}{\Gamma, P_j'' \vec{a}bc \vdash F_j \vec{a}bc} \text{ (Ind } P_j'')$$

Define $F'_k \vec{\alpha} b$ be $Nb \wedge \forall z \in N(b < z \rightarrow \tilde{F}_k \vec{\alpha} bz)$ for all k .

We will show $\tilde{\Gamma}, P'_j \vec{\alpha} b \vdash F'_j \vec{\alpha} b$ by

$$\frac{J'_i \quad (\forall i \in I)}{\tilde{\Gamma}, P'_j \vec{\alpha} b \vdash F'_j \vec{\alpha} b} \text{ (Ind } P'_j)$$

We will show each J'_i . Let J'_i be

$$\tilde{\Gamma}, Q_1 \vec{u}_1, \dots, Q_n \vec{u}_n, F'_1 \vec{t}_1 v_1, v_1 < v, \dots, F'_m \vec{t}_m v_m, v_m < v, Nv \vdash F' \vec{t} v.$$

and the production rules be

$$\begin{array}{c} \frac{Q_1 \vec{u}_1 \quad \dots \quad Q_n \vec{u}_n \quad P_1 \vec{t}_1 \quad \dots \quad P_m \vec{t}_m}{P \vec{t}} \\[10pt] \frac{Q_1 \vec{u}_1 \quad \dots \quad Q_n \vec{u}_n \quad P'_1 \vec{t}_1 v_1 \quad v_1 < v \quad \dots \quad P'_m \vec{t}_m v_m \quad v_m < v \quad Nv}{P' \vec{t} v} \\[10pt] \frac{Q_1 \vec{u}_1 \quad \dots \quad Q_n \vec{u}_n \quad P''_1 \vec{t}_1 v_1 w_1 \quad w_1 < w \quad v_1 < v \quad \dots \quad P''_m \vec{t}_m v_m w_m \quad w_m < w \quad v_m < v \quad N' v w_0 \quad w_0 < w \quad Nw}{P'' \vec{t} vw} \end{array}$$

We can assume the production rule for J'_i is the stage-number transformation of the production rule for J_i . Then J_i is

$$\Gamma, Q_1 \vec{u}_1, \dots, Q_n \vec{u}_n, F_1 \vec{t}_1 v_1 w_1, w_1 < w, v_1 < v, \dots, F_m \vec{t}_m v_m w_m, w_m < w, v_m < v, N' v w_0, w_0 < w, Nw \vdash F \vec{t} vw$$

By IH for J_i we have \tilde{J}_i , namely

$$\tilde{\Gamma}, Q_1 \vec{u}_1, \dots, Q_n \vec{u}_n, \tilde{F}_1 \vec{t}_1 v_1 w_1, w_1 < w, v_1 < v, \dots, \tilde{F}_m \vec{t}_m v_m w_m, w_m < w, v_m < v,$$

$$Nv, Nw_0, v \leq w_0, w_0 < w, Nw \vdash \tilde{F} \vec{t} vw. \quad (33)$$

We can show J'_i as follows. Assume the antecedent, fix z , and assume Nz and $v < z$ in order to show $\tilde{F} \vec{t} vz$. Take w to be z , w_l to be sv_l , and w_0 to be v in (33). Then $w_l < w$ by $v_l < v$ and $v < z$. We also have $w_0 < w$ by $v < z$. Moreover $\tilde{F}_l \vec{t}_l v_l w_l$ by $F'_l \vec{t}_l v_l$. Hence $\tilde{F}_l \vec{t}_l vz$ by (33). Hence we have shown J'_i .

By (Ind P'), we have $\tilde{\Gamma}, P'_j \vec{\alpha} b \vdash F'_j \vec{\alpha} b$. Hence $\tilde{\Gamma}, P'_j \vec{\alpha} b, Nc, b < c \vdash \tilde{F}_j \vec{\alpha} bc$. Hence we have $(\Gamma, \widetilde{P''_j \vec{\alpha} bc}) \vdash \widetilde{F_j \vec{\alpha} bc}$. \square

The next main theorem shows Brotherston-Simpson conjecture under arithmetic for intuitionistic logic.

Theorem 6.14 (Equivalence of LJID + HA and CLJID $^\omega$ + HA) *Let $\Sigma = \Sigma_N \cup \{\vec{Q}, \vec{P}, \vec{P}'\}$ and $\Phi = \Phi_N \cup \{\vec{P}, \vec{P}'\}$. Then the provability of CLJID $^\omega$ + HA + (Σ, Φ) is the same as that of LJID + HA + (Σ, Φ) .*

Proof. (1) **LJID + HA + (Σ, Φ) to CLJID $^\omega$ + HA + (Σ, Φ) .**

For this claim, we can obtain a proof from the proof of Lemma 7.5 in [8] by restricting every sequent into intuitionistic sequents and replacing **LKID** + (Σ, Φ) and **CLKID** $^\omega$ + (Σ, Φ) by **LJID** + (Σ, Φ) and **CLJID** $^\omega$ + (Σ, Φ) respectively.

(2) **CLJID** $^\omega$ + **HA** + (Σ, Φ) to **LJID** + **HA** + (Σ, Φ) .

Let $\Sigma' = \Sigma \cup \{N', \vec{P}''\}$ and $\Phi' = \Phi \cup \{N', \vec{P}''\}$. Assume J is provable in **CLJID** $^\omega$ + **HA** + (Σ, Φ) . By Proposition 6.12, J is provable in **LJID** + **HA** + (Σ', Φ') . By Proposition 6.13, J is provable in **LJID** + **HA** + (Σ, Φ) . \square

7 Conclusion

We have studied Brotherston-Simpson conjecture for intuitionistic logic. We have pointed out that the countermodel of [2] shows the Brotherston-Simpson conjecture for intuitionistic logic is false in general. We have shown **HA**-provability of Kleene-Brouwer theorem for induction and Podelski-Rybalchenko theorem for induction. By using them, we have shown the conjecture for intuitionistic logic is true under arithmetic, namely, the provability of the intuitionistic cyclic proof system is the same as that of the intuitionistic system of Martin-Lof's inductive definitions when both systems contain Heyting arithmetic.

References

- [1] S. Berardi and S. Steila, An intuitionistic version of Ramseys Theorem and its use in Program Termination, *Annals of Pure and Applied Logic* **166** (2015) 1382-1406.
- [2] S. Berardi and M. Tatsuta, Classical System of Martin-Lof's Inductive Definitions is not Equivalent to Cyclic Proof System, In: *Proceedings of FoSSaCS 2017*, LNCS 10203 (2017) 301–317.
- [3] S. Berardi and M. Tatsuta, Equivalence of Inductive Definitions and Cyclic Proofs under Arithmetic, In: *Proceedings of Thirty-Second Annual IEEE Symposium on Logic in Computer Science (LICS2017)* (2017) 1–12.
- [4] J. Brotherston, Cyclic Proofs for First-Order Logic with Inductive Definitions, In: *Proceedings of TABLEAUX 2005*, 2005.
- [5] J. Brotherston, Sequent calculus proof systems for inductive definitions, PhD. thesis, University of Edinburgh, 2006.
- [6] J. Brotherston, R. Bornat and C. Calcagno, Cyclic Proofs of Program Termination in Separation Logic, In: *Proceedings of POPL 2008*, 2008.
- [7] J. Brotherston, D. Distefano and R.L. Petersen, Automated Cyclic Entailment Proofs in Separation Logic, In: *Proceedings of CADE-23*, 2011.
- [8] J. Brotherston and A. Simpson, Sequent calculi for induction and infinite descent, *Journal of Logic and Computation* **21** (6) (2011) 1177–1216.
- [9] J. Brotherston, N. Gorogiannis and R.L. Petersen, A Generic Cyclic Theorem Prover, In: *Proceedings of APLAS 2012*, 2012.
- [10] C.G. Jockusch, Jr, Ramsey's theorem and recursion theory, *Journal of Symbolic Logic* **37**(2) (1972) 268–280.
- [11] P. Martin-Löf, Hauptatz for the intuitionistic theory of iterated inductive definitions, In: *Proceedings of the Second Scandinavian Logic Symposium*, North-Holland, (1971) 179-216.
- [12] A. Podelski and A. Rybalchenko, Transition Invariants, In: *Proceedings of 19th IEEE Symposium on Logic in Computer Science (LICS 2004)*, 32-41 (2004).
- [13] F.P. Ramsey, On a problem in formal logic, In: *Proceedings of London Mathematical Society*, vol. 30 (1930) 264-286.
- [14] J.R. Shoenfield, *Mathematical Logic*, Addison-Wesley, 1967.
- [15] A. Simpson, Cyclic Arithmetic is Equivalent to Peano Arithmetic, In: *Proceedings of FoSSaCS 2017* (2017) 283–300.