

Binary quadratic forms and ray class groups

ICK SUN EUM, JA KYUNG KOO AND DONG HWA SHIN

Abstract

Let K be an imaginary quadratic field different from $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$. For a positive integer N , let $K_{\mathfrak{n}}$ be the ray class field of K modulo $\mathfrak{n} = N\mathcal{O}_K$. By using the congruence subgroup $\pm\Gamma_1(N)$, we construct an extended form class group whose operation is basically the Dirichlet composition, and explicitly show that this group is isomorphic to the Galois group $\text{Gal}(K_{\mathfrak{n}}/K)$. We also present algorithms to find all form classes and show how to multiply two form classes. As an application, we describe $\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$ in terms of these extended form class groups for which $K_{\mathfrak{n}}^{\text{ab}}$ is the maximal abelian extension of K unramified outside prime ideals dividing \mathfrak{n} .

1 Introduction

Let K be an imaginary quadratic field of discriminant d_K with ring of integers \mathcal{O}_K . Let $\mathcal{Q}(d_K)$ be the set of primitive positive definite binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2$ ($\in \mathbb{Z}[x, y]$) of discriminant $b^2 - 4ac = d_K$. Define an equivalence relation on $\mathcal{Q}(d_K)$, called the *proper equivalence*, by

$$Q' \sim Q \iff Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \text{ for some } \sigma \in \text{SL}_2(\mathbb{Z}).$$

Then, the set $C(d_K) = \mathcal{Q}(d_K)/\sim$ of equivalence classes under Dirichlet composition becomes a group, called the *form class group* of discriminant d_K ([1, Theorem 3.9]).

Let I_K be the group of fractional ideals of K and P_K be its subgroup of principal fractional ideals. It is a classical fact that the form class group $C(d_K)$ is isomorphic to the ideal class group $\text{Cl}(\mathcal{O}_K) = I_K/P_K$ as follows: For each $Q \in \mathcal{Q}(d_K)$, let ω_Q be the zero of $Q(x, 1)$ in the complex upper half-plane \mathbb{H} .

2010 *Mathematics Subject Classification.* Primary 11R29; Secondary 11E16, 11G15, 11R37.

Key words and phrases. binary quadratic forms, class field theory, class groups, complex multiplication.

The first author was supported by the Dongguk University Research Fund of 2016 and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2017R1C1B5017567). The third (corresponding) author was supported by the Hankuk University of Foreign Studies Research Fund of 2018 and the National Research Foundation of Korea (NRF) grant funded by the Korea government (MSIT) (NRF-2017R1A2B1006578).

THEOREM 1.1. *We have an isomorphism of groups*

$$\begin{aligned} \phi : C(d_K) &\rightarrow \text{Cl}(\mathcal{O}_K) \\ \text{form class containing } Q = ax^2 + bxy + cy^2 &\mapsto \text{ideal class containing } a[\omega_Q, 1]. \end{aligned}$$

PROOF. See [1, Theorem 7.7]. □

REMARK 1.2. In Theorem 1.1, one can replace the integral ideal $a[\omega_Q, 1] = [(-b + \sqrt{d_K})/2, a]$ by the fractional ideal $[\omega_Q, 1]$.

On the other hand, let H_K be the Hilbert class field of K whose Galois group is isomorphic to $\text{Cl}(\mathcal{O}_K)$ ([1, Theorem 8.10] or [4, Theorem 9.9 in Chapter V]). The following theorem is a consequence of the theory of complex multiplication and Theorem 1.1.

THEOREM 1.3. *We have an isomorphism of groups*

$$\begin{aligned} C(d_K) &\rightarrow \text{Gal}(H_K/K) \\ \text{form class containing } Q &\mapsto (j(\tau_K) \mapsto j(\omega_Q)), \end{aligned}$$

where $j(\tau)$ is the elliptic modular function and τ_K is an element of \mathbb{H} such that $\mathcal{O}_K = [\tau_K, 1]$.

PROOF. See [2], [3] or [6, Theorem 1 in Chapter 10]. □

Now, for a finite abelian extension L of K such that $L \supseteq H_K$, it is natural to ask whether there is some form class group that is isomorphic to $\text{Gal}(L/K)$. Since $\text{Gal}(H_K/K) (\simeq C(d_K))$ is a quotient group of $\text{Gal}(L/K)$, if we loosen the proper equivalence on $C(d_K)$ induced from $\text{SL}_2(\mathbb{Z})$, then we would expect to get a certain new form class group isomorphic to $\text{Gal}(L/K)$. Here we note that L is contained in some ray class field $K_{\mathfrak{n}}$ modulo $\mathfrak{n} = N\mathcal{O}_K$ for a positive integer N ([1, p. 149]).

PROPOSITION 1.4. *Let \mathcal{F}_N be the field of meromorphic modular functions of level N whose Fourier coefficients lie in the N th cyclotomic field. Then we have*

$$K_{\mathfrak{n}} = K(h(\tau_K) \mid h(\tau) \in \mathcal{F}_N \text{ is finite at } \tau_K).$$

PROOF. See [6, Corollary to Theorem 2 in Chapter 10]. □

In this paper, we shall first construct a newly extended form class group $C_N(d_K)$ isomorphic to the ray class group $\text{Cl}(\mathfrak{n})$, through the equivalence relation induced from $\pm\Gamma_1(N)$ (Theorem 2.9). It turns out that the binary operation on $C_N(d_K)$ is essentially the Dirichlet composition on $C(d_K)$ (Remark 2.10 (iv)).

In view of Theorem 1.3 and Proposition 1.4 we shall further establish an isomorphism

$$\begin{aligned} C_N(d_K) &\rightarrow \text{Gal}(K_{\mathfrak{n}}/K) \\ \text{form class containing } Q = ax^2 + bxy + cy^2 &\mapsto \left(h(\tau_K) \mapsto h \left[\begin{smallmatrix} a & (b-b_K)/2 \\ 0 & 1 \end{smallmatrix} \right] (\omega_Q) \mid h(\tau) \in \mathcal{F}_N \text{ is finite at } \tau_K \right), \end{aligned}$$

where $\min(\tau_K, \mathbb{Q}) = x^2 + b_K x + c_K \in \mathbb{Z}[x]$ (Theorem 3.8). This indicates that a form class $[ax^2 + bxy + cy^2]$ in $C_N(d_K)$ has perfect information on an element of $\text{Gal}(K_{\mathfrak{n}}/K)$. Of course, we shall present an algorithm in order to list all representatives of form classes in $C_N(d_K)$ (Proposition 4.4) and give some examples.

Let $K_{\mathfrak{n}}^{\text{ab}}$ be the maximal abelian extension of K unramified outside prime ideals dividing \mathfrak{n} . As an application, we shall construct a dense subset of $\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$, equipped with Krull topology, in terms of extended form class groups (Theorem 5.4).

2 Extended form class groups as ray class groups

Throughout this paper, let K be an imaginary quadratic field of discriminant d_K other than $\mathbb{Q}(\sqrt{-1})$ and $\mathbb{Q}(\sqrt{-3})$. For a positive integer N , let $\mathcal{Q}_N(d_K)$ be the set of primitive positive definite binary quadratic forms $Q(x, y) = ax^2 + bxy + cy^2$ of discriminant d_K such that $\gcd(N, a) = 1$, that is,

$$\mathcal{Q}_N(d_K) = \{ax^2 + bxy + cy^2 \in \mathcal{Q}(d_K) \mid \gcd(N, a) = 1\}.$$

By $\pm\Gamma_1(N)$ we mean the congruence subgroup of $\text{SL}_2(\mathbb{Z})$ given by

$$\pm\Gamma_1(N) = \left\{ \sigma \in \text{SL}_2(\mathbb{Z}) \mid \sigma \equiv \pm \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \pmod{N} \right\}.$$

PROPOSITION 2.1. *The group $\pm\Gamma_1(N)$ acts on the set $\mathcal{Q}_N(d_K)$ on the right by*

$$Q^\sigma = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \quad (\sigma \in \pm\Gamma_1(N), Q \in \mathcal{Q}_N(d_K)).$$

PROOF. Since $\text{SL}_2(\mathbb{Z})$ acts on $\mathcal{Q}(d_K)$, it suffices to show that $\pm\Gamma_1(N)$ preserves the set $\mathcal{Q}_N(d_K)$. Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$ and $\sigma \in \pm\Gamma_1(N)$. We then see that

$$\begin{aligned} Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) &\equiv Q \left(\pm \begin{bmatrix} 1 & s \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \right) \pmod{N\mathbb{Z}[x, y]} \quad \text{for some } s \in \mathbb{Z} \\ &\equiv ax^2 + (2as + b)xy + (as^2 + bs + c)y^2 \pmod{N\mathbb{Z}[x, y]}. \end{aligned}$$

This claims that $Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right)$ belongs to $\mathcal{Q}_N(d_K)$, as desired. □

DEFINITION 2.2. Define an equivalence relation \sim_N on the set $\mathcal{Q}_N(d_K)$ by

$$Q \sim_N Q' \iff Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \text{ for some } \sigma \in \pm\Gamma_1(N).$$

Denote by $C_N(d_K)$ the set of equivalence classes, namely,

$$C_N(d_K) = \mathcal{Q}_N(d_K) / \sim_N.$$

Now, we are in need of the following basic lemma for later use.

LEMMA 2.3. Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}(d_K)$ and $\sigma = \begin{bmatrix} r & s \\ u & v \end{bmatrix} \in \mathrm{SL}_2(\mathbb{Z})$.

(i) If $\omega \in \mathbb{H}$, then

$$[\sigma(\omega), 1] = \frac{1}{j(\sigma, \omega)}[\omega, 1] \quad \text{where } j(\sigma, \omega) = u\omega + v.$$

(ii) If $Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right)$, then

$$\omega_Q = \sigma(\omega_{Q'}).$$

(iii) We have

$$N_{K/\mathbb{Q}}([\omega_Q, 1]) = \frac{1}{a},$$

where $N_{K/\mathbb{Q}}(\cdot)$ is applied to fractional ideals of K .

PROOF. (i) It follows from the fact $\sigma \in \mathrm{SL}_2(\mathbb{Z})$ that

$$[\sigma(\omega), 1] = \left[\frac{r\omega + s}{u\omega + v}, 1 \right] = \frac{1}{u\omega + v} [r\omega + s, u\omega + v] = \frac{1}{j(\sigma, \omega)} [\omega, 1].$$

(ii)

$$Q \left(\begin{bmatrix} \omega_Q \\ 1 \end{bmatrix} \right) = 0 = Q' \left(\begin{bmatrix} \omega_{Q'} \\ 1 \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} \omega_{Q'} \\ 1 \end{bmatrix} \right) = j(\sigma, \omega_{Q'})^2 Q \left(\begin{bmatrix} \sigma(\omega_{Q'}) \\ 1 \end{bmatrix} \right).$$

Since $\omega_Q, \omega_{Q'} \in \mathbb{H}$, we conclude $\omega_Q = \sigma(\omega_{Q'})$.

(iii)

$$\mathrm{disc}_{K/\mathbb{Q}}([\omega_Q, 1]) = \begin{vmatrix} (-b + \sqrt{d_K})/2a & 1 \\ (-b - \sqrt{d_K})/2a & 1 \end{vmatrix}^2 = \frac{d_K}{a^2}.$$

On the other hand, since

$$\mathrm{disc}_{K/\mathbb{Q}}([\omega_Q, 1]) = N_{K/\mathbb{Q}}([\omega_Q, 1])^2 d_K$$

([7, Proposition 13 in Chapter III]), we achieve

$$N_{K/\mathbb{Q}}([\omega_Q, 1]) = \frac{1}{a}.$$

□

From now on, we let $\mathfrak{n} = N\mathcal{O}_K$ and

$$\mathrm{Cl}(\mathfrak{n}) = I_K(\mathfrak{n})/P_{K,1}(\mathfrak{n})$$

be the ray class group of K modulo \mathfrak{n} , where $I_K(\mathfrak{n})$ is the subgroup of I_K consisting of fractional ideals of K prime to \mathfrak{n} and $P_{K,1}(\mathfrak{n})$ is its subgroup consisting of principal fractional ideals $\lambda\mathcal{O}_K$ with $\lambda \in K^*$ such that $\lambda \equiv^* 1 \pmod{\mathfrak{n}}$ ([4, pp. 136–137]).

DEFINITION 2.4. Define a map

$$\begin{aligned}\phi_N : \mathcal{C}_N(d_K) &\rightarrow \text{Cl}(\mathfrak{n}) \\ [Q] &\mapsto \text{ray class containing } [\omega_Q, 1].\end{aligned}$$

Here, $[Q]$ stands for the form class containing $Q \in \mathcal{Q}_N(d_K)$.

REMARK 2.5. By Remark 1.2, we see that $\phi_1 = \phi$, the classical isomorphism described in Theorem 1.1.

PROPOSITION 2.6. *The map ϕ_N is well defined.*

PROOF. First, we shall show that if $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$, then the fractional ideal $[\omega_Q, 1]$ is prime to \mathfrak{n} . Observe that $a[\omega_Q, 1] = [(-b + \sqrt{d_K})/2, a]$ is an integral ideal of K with

$$N_{K/\mathbb{Q}}(a[\omega_Q, 1]) = a$$

by Lemma 2.3 (iii). This, together with the fact $\gcd(N, a) = 1$, implies that $[\omega_Q, 1]$ is prime to \mathfrak{n} .

Second, we shall show that if $Q, Q' \in \mathcal{Q}_N(d_K)$ such that $[Q] = [Q']$, then $[\omega_Q, 1]$ and $[\omega_{Q'}, 1]$ belong to the same ray class in $\text{Cl}(\mathfrak{n})$. Let

$$Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = a'x^2 + b'xy + c'y^2 = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \quad \text{for some } \sigma = \begin{bmatrix} * & * \\ u & v \end{bmatrix} \in \pm\Gamma_1(N).$$

We then derive by Lemma 2.3 (i) and (ii) that

$$[\omega_{Q'}, 1] = (u\omega_{Q'} + v)[\sigma(\omega_{Q'}), 1] = (u\omega_{Q'} + v)[\omega_Q, 1].$$

Since $\sigma \equiv \pm \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \pmod{N}$ and $\gcd(N, a') = 1$, we attain

$$u\omega_{Q'} + v \equiv^* u \frac{-b' + \sqrt{d_K}}{2a'} + v \equiv^* \pm 1 \pmod{\mathfrak{n}}.$$

This yields that $[\omega_Q, 1]$ and $[\omega_{Q'}, 1]$ belong to the same ray class in $\text{Cl}(\mathfrak{n})$. □

PROPOSITION 2.7. *The map ϕ_N is injective.*

PROOF. Suppose that

$$\phi_N([Q]) = \phi_N([Q']) \quad \text{for some } Q, Q' \in \mathcal{Q}_N(d_K),$$

and so

$$[\omega_Q, 1] = \lambda[\omega_{Q'}, 1] \quad \text{for some } \lambda \in K^* \text{ such that } \lambda \equiv^* 1 \pmod{\mathfrak{n}}. \quad (1)$$

Then, we get by Theorem 1.1 that

$$Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \quad \text{for some } \sigma = \begin{bmatrix} * & * \\ u & v \end{bmatrix} \in \text{SL}_2(\mathbb{Z}).$$

And, it follows from Lemma 2.3 (i), (ii) and (1) that

$$[\omega_{Q'}, 1] = j(\sigma, \omega_{Q'})[\sigma(\omega_{Q'}), 1] = (u\omega_{Q'} + v)[\omega_Q, 1] = \lambda(u\omega_{Q'} + v)[\omega_{Q'}, 1],$$

and hence

$$\lambda(u\omega_{Q'} + v) \in \mathcal{O}_K^* = \{1, -1\}.$$

Now that $\lambda \equiv^* 1 \pmod{\mathfrak{n}}$, we deduce

$$u\omega_{Q'} + v \equiv^* \pm 1 \pmod{\mathfrak{n}}. \quad (2)$$

If we let $Q'(x, y) = a'x^2 + b'xy + c'y^2$, then we have $\mathcal{O}_K = [(-b' + \sqrt{d_K})/2, 1]$ and

$$u\omega_{Q'} + v \pm 1 = \frac{1}{a'} \left(u \frac{-b' + \sqrt{d_K}}{2} + a'(v \pm 1) \right).$$

Thus, it follows from the fact $\gcd(N, a') = 1$ and (2) that

$$u \equiv 0, \quad v \equiv \pm 1 \pmod{N}.$$

Moreover, since $\det(\sigma) = 1$, we obtain

$$\sigma \equiv \pm \begin{bmatrix} 1 & * \\ 0 & 1 \end{bmatrix} \pmod{N}.$$

Therefore, Q and Q' belong to the same class in $C_N(d_K)$, namely, $[Q] = [Q']$. This proves the proposition. \square

PROPOSITION 2.8. *The map ϕ_N is surjective.*

PROOF. Let $C \in \text{Cl}(\mathfrak{n})$. Take an integral ideal \mathfrak{a} in C^{-1} , and let $\xi_1, \xi_2 \in K^*$ such that

$$\mathfrak{a}^{-1} = [\xi_1, \xi_2] \quad \text{and} \quad \xi = \frac{\xi_1}{\xi_2} \in \mathbb{H}.$$

Since $1 \in \mathfrak{a}^{-1}$, one can write

$$1 = u\xi_1 + v\xi_2 \quad \text{for some } u, v \in \mathbb{Z}. \quad (3)$$

We then claim $\gcd(N, u, v) = 1$. Otherwise, $d = \gcd(N, u, v) > 1$, and so $d\mathfrak{a}^{-1} = [d\xi_1, d\xi_2]$ contains 1 by (3), which implies $d\mathcal{O}_K \supseteq \mathfrak{a}$. But, this contradicts the fact that \mathfrak{a} is prime to \mathfrak{n} .

Thus we may take a matrix $\sigma = \begin{bmatrix} * & * \\ \tilde{u} & \tilde{v} \end{bmatrix}$ in $\text{SL}_2(\mathbb{Z})$ such that

$$\sigma \equiv \begin{bmatrix} * & * \\ u & v \end{bmatrix} \pmod{N} \quad (4)$$

by the surjectivity of $\text{SL}_2(\mathbb{Z}) \rightarrow \text{SL}_2(\mathbb{Z}/N\mathbb{Z})$ ([10, Lemma 1.38]). If we set $\omega = \sigma(\xi)$, then we derive that

$$[\omega, 1] = [\sigma(\omega), 1]$$

$$\begin{aligned}
&= \frac{1}{\tilde{u}\xi + \tilde{v}}[\xi, 1] \quad \text{by Lemma 2.3 (i)} \\
&= \frac{\xi_2}{\tilde{u}\xi_1 + \tilde{v}\xi_2}[\xi_1/\xi_2, 1] \quad \text{by the fact } \xi = \xi_1/\xi_2 \\
&= \frac{1}{\tilde{u}\xi_1 + \tilde{v}\xi_2}[\xi_1, \xi_2] \\
&= \frac{1}{\tilde{u}\xi_1 + \tilde{v}\xi_2}\mathfrak{a}^{-1}.
\end{aligned}$$

Here, we note that

$$\begin{aligned}
\tilde{u}\xi_1 + \tilde{v}\xi_2 - 1 &= \tilde{u}\xi_1 + \tilde{v}\xi_2 - (u\xi_1 + v\xi_2) \quad \text{by (3)} \\
&= (\tilde{u} - u)\xi_1 + (\tilde{v} - v)\xi_2 \\
&\in N\mathfrak{a}^{-1} \quad \text{by (4),}
\end{aligned}$$

from which we see that

$$\tilde{u}\xi + \tilde{v} \equiv^* 1 \pmod{\mathfrak{n}}.$$

Therefore, $[\omega, 1]$ and \mathfrak{a}^{-1} belong to the same ray class C . Thus, if we let Q be the element of $\mathcal{Q}_N(d_K)$ satisfying $\omega_Q = \omega$, then we conclude

$$\phi_N([Q]) = C.$$

□

THEOREM 2.9. *The set $C_N(d_K)$ can be regarded as an abelian group isomorphic to the ray class group $\text{Cl}(\mathfrak{n})$.*

PROOF. Define a binary operation \cdot on $C_N(d_K)$ by

$$[Q] \cdot [Q'] = \phi_N^{-1}(\phi_N([Q])\phi_N([Q'])),$$

where $\phi_N([Q])\phi_N([Q'])$ is the product of ray classes in $\text{Cl}(\mathfrak{n})$. This binary operation makes $C_N(d_K)$ an abelian group isomorphic to $\text{Cl}(\mathfrak{n})$. We shall describe the group operation on $C_N(d_K)$ explicitly in the following Remark 2.10 (iv). □

REMARK 2.10. (i) If M is a positive divisor of N and $\mathfrak{m} = M\mathcal{O}_K$, then we have by Definition 2.4 a commutative diagram of homomorphisms

$$\begin{array}{ccc}
C_N(d_K) & \xrightarrow{\phi_N} & \text{Cl}(\mathfrak{n}) \\
\downarrow & & \downarrow \\
C_M(d_K) & \xrightarrow{\phi_M} & \text{Cl}(\mathfrak{m})
\end{array}$$

The vertical maps are natural projections.

(ii) Let τ_K be the element of \mathbb{H} induced by the principal form

$$\begin{cases} x^2 + xy + \frac{1-d_K}{4}y^2 & \text{if } d_K \equiv 1 \pmod{4}, \\ x^2 - \frac{d_K}{4}y^2 & \text{if } d_K \equiv 0 \pmod{4}. \end{cases}$$

Since $[\tau_K, 1] = \mathcal{O}_K$, the principal form gives rise to the identity element of $C_N(d_K)$.

- (iii) For a quadratic form $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$, we want to find its inverse $[Q]^{-1}$ in $C_N(d_K)$. Let $\mathfrak{c} = a^{\varphi(N)}[\omega_Q, 1]$, where φ is the Euler function. Then, \mathfrak{c} is an integral ideal of K which belongs to the same ray class as $[\omega_Q, 1]$ because $a^{\varphi(N)} \equiv 1 \pmod{N}$. Since $\mathfrak{c}\bar{\mathfrak{c}} = N_{K/\mathbb{Q}}(\mathfrak{c})\mathcal{O}_K$ and $N_{K/\mathbb{Q}}([\omega_Q, 1]) = 1/a$ by Lemma 2.3 (iii), we get

$$\mathfrak{c}^{-1} = \frac{1}{N_{K/\mathbb{Q}}(\mathfrak{c})}\bar{\mathfrak{c}} = \frac{1}{a^{\varphi(N)-1}}[-\bar{\omega}_Q, 1];$$

and hence we obtain

$$1 = \frac{1}{a^{\varphi(N)-1}}(0 \cdot (-\bar{\omega}_Q) + a^{\varphi(N)-1} \cdot 1).$$

Take an element σ in $\text{SL}_2(\mathbb{Z})$ such that

$$\sigma \equiv \begin{bmatrix} * & * \\ 0 & a^{\varphi(N)-1} \end{bmatrix} \pmod{N}.$$

Now, if we let $Q' \in \mathcal{Q}_N(d_K)$ satisfying $\omega_{Q'} = \sigma(-\bar{\omega}_Q)$, then we achieve by the proof of Proposition 2.8 that Q' and \mathfrak{c}^{-1} give the same ray class. Therefore, $[Q']$ is the inverse of $[Q]$ in $C_N(d_K)$.

- (iv) Let $Q_1(x, y) = a_1x^2 + b_1xy + c_1y^2$, $Q_2(x, y) = a_2x^2 + b_2xy + c_2y^2 \in \mathcal{Q}_N(d_K)$. We will describe an algorithm how to find $[Q_1] \cdot [Q_2]$ explicitly. One may take a matrix ρ in $\text{SL}_2(\mathbb{Z})$ so that $Q_3(x, y) = a_3x^2 + b_3xy + c_3y^2$ defined by

$$Q_3 \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q_2 \left(\rho \begin{bmatrix} x \\ y \end{bmatrix} \right) \quad (5)$$

satisfies $\gcd(a_1, a_3, (b_1 + b_3)/2) = 1$ ([1, Lemmas 2.3 and 2.25]). We then attain

$$[\omega_{Q_1}, 1][\omega_{Q_3}, 1] = \left[\frac{-B + \sqrt{d_K}}{2a_1a_3}, 1 \right], \quad (6)$$

where B is an integer for which

$$B \equiv b_1 \pmod{2a_1}, \quad B \equiv b_3 \pmod{2a_3} \quad \text{and} \quad B^2 \equiv d_K \pmod{4a_1a_3}$$

([1, Lemma 3.2 and (7.13)]). (This ideal multiplication gives us the Dirichlet composition on $C_1(d_K) = C(d_K)$.) On the other hand, we know by Definition 2.4 that $\phi_N([Q_1])\phi_N([Q_2])$ is the ray class containing the fractional ideal

$$\mathfrak{c} = [\omega_{Q_1}, 1][\omega_{Q_2}, 1].$$

Thus, we get that

$$\begin{aligned} \mathfrak{c} &= [\omega_{Q_1}, 1][\rho(\omega_{Q_3}), 1] \quad \text{by (5) and Lemma 2.3 (ii)} \\ &= \frac{1}{j(\rho, \omega_{Q_3})}[\omega_{Q_1}, 1][\omega_{Q_3}, 1] \quad \text{by Lemma 2.3 (i)} \\ &= \frac{1}{j(\rho, \omega_{Q_3})} \left[\frac{-B + \sqrt{d_K}}{2a_1a_3}, 1 \right] \quad \text{by (6).} \end{aligned}$$

By the fact $\mathfrak{c}\bar{\mathfrak{c}} = N_{K/\mathbb{Q}}(\mathfrak{c})\mathcal{O}_K$ and Lemma 2.3 (iii) we see that

$$\mathfrak{a} = \mathfrak{c}^{-1} = \frac{1}{N_{K/\mathbb{Q}}(\mathfrak{c})}\bar{\mathfrak{c}} = a_1a_2\bar{\mathfrak{c}} = [-a_1\bar{\omega}_{Q_1}, a_1][-a_2\bar{\omega}_{Q_2}, a_2]$$

is an integral ideal in the ray class $(\phi_N([Q_1])\phi([Q_2]))^{-1}$. Now, by using the argument in the proof of Proposition 2.8 one can have $Q_4 \in \mathcal{Q}_N(d_K)$ so that $\phi_N([Q_4])$ is the ray class containing $\mathfrak{a}^{-1} = \mathfrak{c}$. Therefore, we achieve

$$[Q_4] = [Q_1] \cdot [Q_2].$$

3 Extended form class groups as Galois groups

Let $K_{\mathfrak{n}}$ be the ray class field of K modulo \mathfrak{n} ($= N\mathcal{O}_K$), that is, $K_{\mathfrak{n}}$ is the unique abelian extension of K whose Galois group $\text{Gal}(K_{\mathfrak{n}}/K)$ corresponds to $\text{Cl}(\mathfrak{n})$ via the Artin reciprocity map for \mathfrak{n} . In this section, we shall establish an isomorphism of $C_N(d_K)$ onto $\text{Gal}(K_{\mathfrak{n}}/K)$ in a concrete way.

Let \mathcal{F}_N be the field of meromorphic modular functions of level N with Fourier coefficients in $\mathbb{Q}(\zeta_N)$, where $\zeta_N = e^{2\pi i/N}$. It is well known that \mathcal{F}_N is a Galois extension of \mathcal{F}_1 with

$$\text{Gal}(\mathcal{F}_N/\mathcal{F}_1) \simeq \text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$$

([10, Theorem 6.6]). In particular, the subgroup $\text{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$ of $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$ acts on the field \mathcal{F}_N as follows: Let $h(\tau) \in \mathcal{F}_N$ and $\alpha \in \text{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$. Then we have

$$h(\tau)^\alpha = h(\tilde{\alpha}(\tau)),$$

where $\tilde{\alpha}$ is any matrix in $\text{SL}_2(\mathbb{Z})$ that reduces to α via $\text{SL}_2(\mathbb{Z}) \rightarrow \text{SL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$.

DEFINITION 3.1. We call a Family

$$\{h_\alpha(\tau)\}_{\alpha \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}}$$

of functions in \mathcal{F}_N a *Fricke family* of level N if

$$h_\alpha(\tau)^\beta = h_{\alpha\beta}(\tau) \quad \text{for all } \alpha, \beta \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}.$$

REMARK 3.2. For a Fricke family $\{h_\alpha(\tau)\}_\alpha$, let $h(\tau) = h_{I_2}(\tau)$. Then we get

$$h(\tau)^\alpha = h_{I_2}(\tau)^\alpha = h_{I_2\alpha}(\tau) = h_\alpha(\tau) \quad (\alpha \in \text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}).$$

This shows that $\{h_\alpha(\tau)\}_\alpha$ is a family of Galois conjugates of $h(\tau) = h_{I_2}(\tau)$ under $\text{Gal}(\mathcal{F}_N/\mathcal{F}_1)$.

For a class $C \in \text{Cl}(\mathfrak{n})$ take an integral ideal \mathfrak{a} in C^{-1} , and let $\xi_1, \xi_2 \in K^*$ such that

$$\mathfrak{a}^{-1} = [\xi_1, \xi_2] \quad \text{and} \quad \xi = \frac{\xi_1}{\xi_2} \in \mathbb{H}.$$

Let τ_K be the element of \mathbb{H} stated in Remark 2.10 (ii). Since $\mathcal{O}_K \subseteq \mathfrak{a}^{-1}$ and $\xi \in \mathbb{H}$, one can write

$$\begin{bmatrix} \tau_K \\ 1 \end{bmatrix} = \begin{bmatrix} r & s \\ u & v \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \text{for some } A = \begin{bmatrix} r & s \\ u & v \end{bmatrix} \in M_2^+(\mathbb{Z}). \quad (7)$$

It then follows that

$$\begin{bmatrix} \tau_K & \bar{\tau}_K \\ 1 & 1 \end{bmatrix} = \begin{bmatrix} r & s \\ u & v \end{bmatrix} \begin{bmatrix} \xi_1 & \bar{\xi}_1 \\ \xi_2 & \bar{\xi}_2 \end{bmatrix}.$$

Taking determinant and squaring, we obtain

$$d_K = \det(A)^2 \text{disc}_{K/\mathbb{Q}}(\mathfrak{a}^{-1}) = \det(A)^2 N_{K/\mathbb{Q}}(\mathfrak{a})^{-2} d_K$$

([7, Proposition 13 in Chapter III]). Thus, we deduce $\det(A) = N_{K/\mathbb{Q}}(\mathfrak{a})$ which is prime to N .

DEFINITION 3.3. Let $\{h_\alpha(\tau)\}_\alpha$ be a Fricke family of level N , and let $C \in \text{Cl}(\mathfrak{n})$. Following the above notations, we define

$$h_{\mathfrak{n}}(C) = h_A(\xi).$$

Here, we regard A as an element of $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\}$.

PROPOSITION 3.4. *The value $h_{\mathfrak{n}}(C)$ depends only on the ray class C , not on the choices of \mathfrak{a} and ξ_1, ξ_2 .*

PROOF. First, let \mathfrak{a}' be another integral ideal in C^{-1} . Then we have

$$\mathfrak{a}' = \lambda \mathfrak{a} \quad \text{for some } \lambda \in K^* \text{ such that } \lambda \equiv^* 1 \pmod{\mathfrak{n}},$$

and so

$$\mathfrak{a}'^{-1} = \lambda^{-1} \mathfrak{a}^{-1} = [\lambda^{-1} \xi_1, \lambda^{-1} \xi_2] \quad \text{and} \quad \frac{\lambda^{-1} \xi_1}{\lambda^{-1} \xi_2} = \frac{\xi_1}{\xi_2} = \xi \in \mathbb{H}.$$

We see from the fact $\mathfrak{a}, \mathfrak{a}' = \lambda \mathfrak{a} \subseteq \mathcal{O}_K$ that

$$(\lambda - 1)\mathfrak{a} \subseteq \mathcal{O}_K.$$

Moreover, since $\lambda \equiv^* 1 \pmod{\mathfrak{n}}$ and \mathfrak{a} is prime to \mathfrak{n} , we attain

$$(\lambda - 1)\mathfrak{a} \subseteq \mathfrak{n},$$

and hence

$$(\lambda - 1)\mathcal{O}_K \subseteq N\mathfrak{a}^{-1}.$$

Thus we obtain by the fact $\mathcal{O}_K = [\tau_K, 1]$ that

$$(\lambda - 1)\tau_K = N(a\xi_1 + b\xi_2) \quad \text{and} \quad \lambda - 1 = N(c\xi_1 + d\xi_2) \quad \text{for some } a, b, c, d \in \mathbb{Z}. \quad (8)$$

On the other hand, since $\lambda\mathcal{O}_K \subseteq \lambda\mathfrak{a}'^{-1} = \mathfrak{a}^{-1} = [\xi_1, \xi_2]$, we may write

$$\begin{bmatrix} \lambda\tau_K \\ \lambda \end{bmatrix} = \begin{bmatrix} r' & s' \\ u' & v' \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \text{for some } \begin{bmatrix} r' & s' \\ u' & v' \end{bmatrix} \in M_2^+(\mathbb{Z}). \quad (9)$$

One can then derive by (7), (8) and (9) that

$$N \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} = \begin{bmatrix} r' & s' \\ u' & v' \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} - \begin{bmatrix} r & s \\ u & v \end{bmatrix} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix},$$

which yields

$$\begin{bmatrix} r' & s' \\ u' & v' \end{bmatrix} \equiv \begin{bmatrix} r & s \\ u & v \end{bmatrix} \pmod{N}.$$

Second, let $\xi'_1, \xi'_2 \in \mathbb{H}$ such that

$$\mathfrak{a}^{-1} = [\xi_1, \xi_2] = [\xi'_1, \xi'_2] \quad \text{and} \quad \xi' = \frac{\xi'_1}{\xi'_2} \in \mathbb{H}.$$

We then express

$$\begin{bmatrix} \tau_K \\ 1 \end{bmatrix} = A' \begin{bmatrix} \xi'_1 \\ \xi'_2 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} \xi'_1 \\ \xi'_2 \end{bmatrix} = B \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \text{for some } A' \in M_2^+(\mathbb{Z}) \text{ and } B \in \mathrm{SL}_2(\mathbb{Z}),$$

and so by (7) we deduce

$$A' \begin{bmatrix} \xi'_1 \\ \xi'_2 \end{bmatrix} = A \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} = AB^{-1} \begin{bmatrix} \xi'_1 \\ \xi'_2 \end{bmatrix}.$$

Hence we achieve

$$\xi' = B(\xi) \quad \text{and} \quad A' = AB^{-1}.$$

Therefore we get that

$$h_{A'}(\xi') = h_{AB^{-1}}(B(\xi)) = h_{AB^{-1}}(\tau)^B|_{\tau=\xi} = h_{AB^{-1}B}(\tau)|_{\tau=\xi} = h_A(\xi),$$

which proves the proposition. \square

REMARK 3.5. (i) If C_0 denotes the identity class in $\mathrm{Cl}(\mathfrak{n})$, namely, C_0 is the ray class containing $\mathcal{O}_K = [\tau_K, 1]$, then

$$h_{\mathfrak{n}}(C_0) = h_{I_2}(\tau_K).$$

(ii) The invariant $h_{\mathfrak{n}}(C)$ is an analogue of the Siegel-Ramachandra invariant given in [5, p. 235] and [9].

Let

$$\mathrm{GL}_2(\mathbb{A}_f) = \prod'_{p: \text{primes}} \mathrm{GL}_2(\mathbb{Q}_p),$$

where $'$ denotes the restricted product, that is, for almost all p the p -component of an element of $\mathrm{GL}_2(\mathbb{A}_f)$ lies in $\mathrm{GL}_2(\mathbb{Z}_p)$. One may express $\mathrm{GL}_2(\mathbb{A}_f)$ as

$$\mathrm{GL}_2(\mathbb{A}_f) = U\mathrm{GL}_2^+(\mathbb{Q}) = \mathrm{GL}_2^+(\mathbb{Q})U, \quad \text{where } U = \prod_{p: \text{primes}} \mathrm{GL}_2(\mathbb{Z}_p)$$

([6, Theorem 1 in Chapter 7]). Let

$$\mathcal{F} = \bigcup_{M=1}^{\infty} \mathcal{F}_M.$$

Then, we have a surjective homomorphism

$$\sigma_f : \mathrm{GL}_2(\mathbb{A}_f) \rightarrow \mathrm{Aut}(\mathcal{F})$$

with $\text{Ker}(\sigma_f) = \mathbb{Q}^*$ ([6, Theorems 4 and 6 in Chapter 7] or [10, Theorem 6.23]). More precisely, let $h(\tau) \in \mathcal{F}_N$ and $\gamma = \alpha\beta \in \text{GL}_2(\mathbb{A}_f)$ with $\alpha = (\alpha_p)_p \in U$ and $\beta \in \text{GL}_2^+(\mathbb{Q})$. By using the Chinese remainder theorem, one can find a unique matrix $\tilde{\alpha}$ in $\text{GL}_2(\mathbb{Z}/N\mathbb{Z})$ satisfying $\tilde{\alpha} \equiv \alpha_p \pmod{N}$ for all primes p such that $p \mid N$. We then obtain

$$h(\tau)^{\sigma_f(\gamma)} = h^{\tilde{\alpha}}(\beta(\tau)) \quad (10)$$

([6, Theorem 2 in Chapter 7 and p. 79]).

For $\omega \in K \cap \mathbb{H}$, we have an embedding

$$q_\omega : K^* \rightarrow \text{GL}_2^+(\mathbb{Q})$$

defined by

$$\xi \begin{bmatrix} \omega \\ 1 \end{bmatrix} = q_\omega(\xi) \begin{bmatrix} \omega \\ 1 \end{bmatrix} \quad (\xi \in K^*).$$

By continuity one can extend q_ω to an embedding

$$q_{\omega,p} : K_p^* \rightarrow \text{GL}_2(\mathbb{Q}_p)$$

for each prime p , and hence to an embedding of idele groups

$$q_\omega : \mathbb{A}_K^* \rightarrow \text{GL}_2(\mathbb{A}_f)$$

([6, p. 149]). Let K^{ab} be the maximal abelian extension of K .

PROPOSITION 3.6 (Shimura's reciprocity law). *Let s be a idele of K and (s^{-1}, K) be the Artin symbol for s^{-1} on K^{ab} . Let $\omega \in K \cap \mathbb{H}$ and $h(\tau) \in \mathcal{F}$ which is finite at ω . Then, $h(\omega)$ lies in K^{ab} and satisfies*

$$h(\omega)^{(s^{-1}, K)} = h(\tau)^{\sigma_f(q_\omega(s))}|_{\tau=\omega}.$$

PROOF. See [6, Theorem 1 in Chapter 11] or [10, Theorem 6.31 (i)]. □

THEOREM 3.7. *Let $\{h_\alpha(\tau)\}_\alpha$ be a Fricke family of level N , and let $C \in \text{Cl}(\mathfrak{n})$. If $h_\mathfrak{n}(C)$ is finite, then it belongs to $K_\mathfrak{n}$ and satisfies*

$$h_\mathfrak{n}(C)^{\sigma_\mathfrak{n}(C'^{-1})} = h_\mathfrak{n}(CC') \quad \text{for all } C' \in \text{Cl}(\mathfrak{n})$$

where $\sigma_\mathfrak{n} : \text{Cl}(\mathfrak{n}) \rightarrow \text{Gal}(K_\mathfrak{n}/K)$ is the Artin reciprocity map for \mathfrak{n} .

PROOF. Let \mathfrak{a} and \mathfrak{a}' be integral ideals in C^{-1} and C'^{-1} , respectively. Take $\xi_1, \xi_2, \xi_1'', \xi_2'' \in K^*$ so that

$$\mathfrak{a}^{-1} = [\xi_1, \xi_2] \quad \text{with } \xi = \frac{\xi_1}{\xi_2} \in \mathbb{H},$$

and

$$(\mathfrak{a}\mathfrak{a}')^{-1} = [\xi_1'', \xi_2''] \quad \text{with } \xi'' = \frac{\xi_1''}{\xi_2''} \in \mathbb{H}.$$

Since $\mathcal{O}_K \subseteq \mathfrak{a}^{-1} \subseteq (\mathfrak{a}\mathfrak{a}')^{-1}$, we may write

$$\begin{bmatrix} \tau_K \\ 1 \end{bmatrix} = A \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \text{for some } A \in M_2^+(\mathbb{Z}) \quad (11)$$

and

$$\begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} = B \begin{bmatrix} \xi_1'' \\ \xi_2'' \end{bmatrix} \quad \text{for some } B \in M_2^+(\mathbb{Z}). \quad (12)$$

Let s be an idele of K such that for every prime p

$$\begin{cases} s_p = 1 & \text{if } p \mid N, \\ s_p \mathcal{O}_p = \mathfrak{a}'_p & \text{if } p \nmid N. \end{cases} \quad (13)$$

Now that \mathfrak{a}' is prime to \mathfrak{n} , we get

$$s_p^{-1} \mathcal{O}_p = \mathfrak{a}'_p^{-1} \quad \text{for all primes } p. \quad (14)$$

Observe that for every prime p

$$q_{\xi,p}(s_p^{-1}) \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} = \xi_2 q_{\xi,p}(s_p^{-1}) \begin{bmatrix} \xi \\ 1 \end{bmatrix} = \xi_2 s_p^{-1} \begin{bmatrix} \xi \\ 1 \end{bmatrix} = s_p^{-1} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix}.$$

Thus,

$$B^{-1} \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix} \quad \text{and} \quad q_{\xi,p}(s_p^{-1}) \begin{bmatrix} \xi_1 \\ \xi_2 \end{bmatrix}$$

are bases for the \mathbb{Z}_p -module $(\mathfrak{a}\mathfrak{a}')^{-1}$ by (12) and (14). So, there exists $u_p \in \text{GL}_2(\mathbb{Z}_p)$ such that

$$q_{\xi,p}(s_p^{-1}) = u_p B^{-1}. \quad (15)$$

If we let

$$u = (u_p)_p \in \prod_{p: \text{primes}} \text{GL}_2(\mathbb{Z}_p),$$

then we obtain

$$q_{\xi}(s^{-1}) = u B^{-1}. \quad (16)$$

Now, we derive that

$$\begin{aligned} h_{\mathfrak{n}}(C)^{(s,K)} &= h_A(\xi)^{(s,K)} \quad \text{by Definition 3.3} \\ &= h_A(\tau)^{\sigma_{\mathfrak{f}}(q_{\xi}(s^{-1}))}|_{\tau=\xi} \quad \text{by Proposition 3.6} \\ &= h_A(\tau)^{\sigma_{\mathfrak{f}}(uB^{-1})}|_{\tau=\xi} \quad \text{by (16)} \\ &= h_{Au}(B^{-1}(\tau))|_{\tau=\xi} \quad \text{by (10),} \\ &\quad \text{where } u \text{ is regarded as an element of } \text{GL}_2(\mathbb{Z}/N\mathbb{Z})/\{\pm I_2\} \\ &= h_{AB}(B^{-1}(\tau))|_{\tau=\xi} \quad \text{because for every prime divisor } p \text{ of } N \\ &\quad \text{we have } s_p = 1 \text{ by (13), and so } u_p B^{-1} = I_2 \text{ by (15)} \\ &= h_{AB}(B^{-1}(\xi)) \\ &= h_{AB}(\xi'') \quad \text{by (12)} \\ &= h_{\mathfrak{n}}(CC') \quad \text{since } \begin{bmatrix} \tau_K \\ 1 \end{bmatrix} = AB \begin{bmatrix} \xi_1'' \\ \xi_2'' \end{bmatrix} \text{ by (11) and (12).} \end{aligned}$$

In particular, if $C' = C^{-1}$, then we see that

$$h_{\mathfrak{n}}(C) = h_{\mathfrak{n}}(CC')^{(s^{-1}, K)} = h_{\mathfrak{n}}(C_0)^{(s^{-1}, K)} = h_{I_2}(\tau_K)^{(s^{-1}, K)}.$$

This implies that $h_{\mathfrak{n}}(C)$ belongs to $K_{\mathfrak{n}}$ because $h_{I_2}(\tau_K)$ lies in $K_{\mathfrak{n}}$ by Proposition 1.4. Since $\text{ord}_{\mathfrak{p}} s_p = \text{ord}_{\mathfrak{p}} \mathfrak{a}'$ for all primes p such that $p \nmid N$ and prime ideals lying above p by (13), we achieve

$$(s, K)|_{K_{\mathfrak{n}}} = \sigma_{\mathfrak{n}}(C'^{-1}).$$

Therefore, we conclude

$$h_{\mathfrak{n}}(C)^{\sigma_{\mathfrak{n}}(C'^{-1})} = h_{\mathfrak{n}}(CC').$$

□

Let $\min(\tau_K, \mathbb{Q}) = x^2 + b_K x + c_K \in \mathbb{Z}[x]$, and so

$$\tau_K = \frac{-b_K + \sqrt{d_K}}{2}.$$

THEOREM 3.8. *We have an isomorphism of groups*

$$\begin{aligned} \text{C}_N(d_K) &\rightarrow \text{Gal}(K_{\mathfrak{n}}/K) \\ [ax^2 + bxy + cy^2] &\mapsto \left(h(\tau_K) \mapsto h \left[\begin{smallmatrix} a & (b-b_K)/2 \\ 0 & 1 \end{smallmatrix} \right] \left(\frac{-b+\sqrt{d_K}}{2a} \right) \mid h(\tau) \in \mathcal{F}_N \text{ is finite at } \tau_K \right). \end{aligned}$$

PROOF. Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$. Then, $C = \phi_N([Q])$ is the ray class containing the fractional ideal $\mathfrak{c} = [\omega_Q, 1]$. Since

$$\mathfrak{c}^{-1} = \frac{1}{N_{K/\mathbb{Q}}(\mathfrak{c})} \bar{\mathfrak{c}} = \frac{1}{a} [-\bar{\omega}_Q, 1]$$

by Lemma 2.3 (iii), $\mathfrak{a} = a^{\varphi(N)} \mathfrak{c}^{-1}$ is an integral ideal in C^{-1} . It then follows that

$$\mathfrak{a}^{-1} = \frac{1}{a^{\varphi(N)}} \mathfrak{c} = \frac{1}{a^{\varphi(N)}} [\omega_Q, 1]$$

and

$$\begin{bmatrix} \tau_K \\ 1 \end{bmatrix} = \begin{bmatrix} a^{\varphi(N)+1} & a^{\varphi(N)}(b-b_K)/2 \\ 0 & a^{\varphi(N)} \end{bmatrix} \begin{bmatrix} \omega_Q/a^{\varphi(N)} \\ 1/a^{\varphi(N)} \end{bmatrix}.$$

Since $a^{\varphi(N)} \equiv 1 \pmod{N}$, we have

$$h_{\mathfrak{n}}(C) = h \left[\begin{smallmatrix} a & (b-b_K)/2 \\ 0 & 1 \end{smallmatrix} \right] (\omega_Q).$$

Now, by composing two isomorphisms

$$\begin{aligned} \text{C}_N(d_K) &\rightarrow \text{Cl}(\mathfrak{n}) \\ [ax^2 + bxy + cy^2] &\mapsto \text{ray class containing } [(-b + \sqrt{d_K})/2a, 1] \end{aligned}$$

given in Theorem 2.9 and

$$\begin{aligned} \text{Cl}(\mathfrak{n}) &\rightarrow \text{Gal}(K_{\mathfrak{n}}/K) \\ C &\mapsto \left(h(\tau_K) = h_{\mathfrak{n}}(C_0) \mapsto h_{\mathfrak{n}}(C_0)^{\sigma_{\mathfrak{n}}(C^{-1})} = h_{\mathfrak{n}}(C) \mid h(\tau) \in \mathcal{F}_N \text{ is finite at } \tau_K \right) \end{aligned}$$

obtained by Theorem 3.7, we establish the theorem. □

4 Explicit construction of extended form class groups

In this section, we shall explain how to find representatives of forms classes in $C_N(d_K)$.

LEMMA 4.1. *Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$ and $u, v \in \mathbb{Z}$. Then, the fractional ideal $(u\omega_Q + v)\mathcal{O}_K$ is prime to \mathfrak{n} if and only if $Q(v, -u)$ is prime to N .*

PROOF. We deduce from the facts $\gcd(N, a) = 1$ and $\mathfrak{n} = N\mathcal{O}_K$ that

$$\begin{aligned} & (u\omega_Q + v)\mathcal{O}_K \text{ is prime to } \mathfrak{n} \\ \iff & \text{ the integral ideal } a(u\omega_Q + v)\mathcal{O}_K \text{ is prime to } \mathfrak{n} \\ \iff & N_{K/\mathbb{Q}}(a(u\omega_Q + v)) \text{ is prime to } N. \end{aligned}$$

Hence, we obtain that

$$\begin{aligned} N_{K/\mathbb{Q}}(a(u\omega_Q + v)) &= a^2(u\omega_Q + v)(u\bar{\omega}_Q + v) \\ &= a^2(u^2\omega_Q\bar{\omega}_Q + uv(\omega_Q + \bar{\omega}_Q) + v^2) \\ &= a^2(u^2(c/a) + uv(-b/a) + v^2) \\ &= a(cu^2 - buv + av^2) \\ &= aQ(v, -u). \end{aligned}$$

This proves the lemma. \square

Let $P_K(\mathfrak{n})$ be the subgroup of $I_K(\mathfrak{n})$ consisting of principal fractional ideals prime to \mathfrak{n} .

LEMMA 4.2. *Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$ and $C \in P_K(\mathfrak{n})/P_{K,1}(\mathfrak{n}) \subseteq \text{Cl}(\mathfrak{n})$. Then we have*

$$C = [(u\omega_Q + v)\mathcal{O}_K] \quad \text{for some } u, v \in \mathbb{Z} \text{ such that } \gcd(N, Q(v, -u)) = 1.$$

PROOF. Take an integral ideal \mathfrak{c} in C . Since $\mathcal{O}_K = [a\omega_Q, 1]$, we get

$$\mathfrak{c} = (t\omega_Q + v)\mathcal{O}_K \quad \text{for some } t, v \in \mathbb{Z}.$$

Set $u = ta$. Then, the lemma follows from Lemma 4.1. \square

Define an equivalence relation \equiv_N on \mathbb{Z}^2 by

$$\begin{bmatrix} r \\ s \end{bmatrix} \equiv_N \begin{bmatrix} u \\ v \end{bmatrix} \iff \begin{bmatrix} r \\ s \end{bmatrix} \equiv \pm \begin{bmatrix} u \\ v \end{bmatrix} \pmod{N}.$$

LEMMA 4.3. *Let $Q(x, y) = ax^2 + bxy + cy^2 \in \mathcal{Q}_N(d_K)$, and let $\begin{bmatrix} r \\ s \end{bmatrix}, \begin{bmatrix} u \\ v \end{bmatrix} \in \mathbb{Z}^2$ such that $\gcd(N, Q(s, -r)) = \gcd(N, Q(v, -u)) = 1$. Then, $(r\omega_Q + s)\mathcal{O}_K$ and $(u\omega_Q + v)\mathcal{O}_K$ represent the same ray class in $\text{Cl}(\mathfrak{n})$ if and only if*

$$\begin{bmatrix} r \\ s \end{bmatrix} \equiv_N \begin{bmatrix} u \\ v \end{bmatrix}.$$

PROOF. By Lemma 4.1, both $(r\omega_Q + s)\mathcal{O}_K$ and $(u\omega_Q + v)\mathcal{O}_K$ are prime to \mathfrak{n} . Then we see that

$$\begin{aligned}
& (r\omega_Q + s)\mathcal{O}_K \text{ and } (u\omega_Q + v)\mathcal{O}_K \text{ represent the same ray class in } \text{Cl}(\mathfrak{n}) \\
\iff & \left(\frac{r\omega_Q + s}{u\omega_Q + v} \right) \mathcal{O}_K \in P_{K,1}(\mathfrak{n}) \\
\iff & \frac{r\omega_Q + s}{u\omega_Q + v} \equiv^* \pm 1 \pmod{\mathfrak{n}} \quad \text{because } \mathcal{O}_K^* = \{1, -1\} \\
\iff & a(r\omega_Q + s) \equiv^* \pm a(u\omega_Q + v) \pmod{\mathfrak{n}} \\
\iff & (r \pm u)(a\omega_Q) + (s \pm v)a \in \mathfrak{n} \quad \text{since } a\omega_Q \in \mathcal{O}_K \\
\iff & r \pm u \equiv (s \pm v)a \equiv 0 \pmod{N} \quad \text{due to } \mathfrak{n} = N\mathcal{O}_K = [Na\omega_Q, N] \\
\iff & \begin{bmatrix} r \\ s \end{bmatrix} \equiv \pm \begin{bmatrix} u \\ v \end{bmatrix} \pmod{N} \quad \text{by the fact } \gcd(N, a) = 1 \\
\iff & \begin{bmatrix} r \\ s \end{bmatrix} \equiv_N \begin{bmatrix} u \\ v \end{bmatrix}.
\end{aligned}$$

□

PROPOSITION 4.4. *One can list elements of $C_N(d_K)$ explicitly.*

PROOF. Note first that

$$C(d_K) \simeq \text{Gal}(K_{\mathfrak{n}}/K)/\text{Gal}(K_{\mathfrak{n}}/H_K) \quad \text{and} \quad P_K(\mathfrak{n})/P_{K,1}(\mathfrak{n}) \simeq \text{Gal}(K_{\mathfrak{n}}/H_K). \quad (17)$$

One can readily find reduced forms Q_1, Q_2, \dots, Q_n in $\mathcal{Q}(d_K)$ which represent all classes in $C(d_K)$ ([1, Theorem 2.8]). Take $\sigma_1, \sigma_2, \dots, \sigma_n \in \text{SL}_2(\mathbb{Z})$ for which

$$Q'_i \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q_i \left(\sigma_i \begin{bmatrix} x \\ y \end{bmatrix} \right) \quad (i = 1, 2, \dots, n)$$

belong to $\mathcal{Q}_N(d_K)$ ([1, Lemmas 2.3 and 2.25]). Then,

$$\left\{ \left[[\omega_{Q'_i}, 1] \right] \in \text{Cl}(\mathfrak{n}) \mid i = 1, 2, \dots, n \right\}$$

is a subset of $\text{Cl}(\mathfrak{n})$ whose image under $\text{Cl}(\mathfrak{n}) \rightarrow \text{Cl}(\mathcal{O}_K)$ is all of $\text{Cl}(\mathcal{O}_K)$. Furthermore, for each $i \in \{1, 2, \dots, n\}$, we attain by Lemmas 4.1, 4.2 and 4.3 that

$$P_K(\mathfrak{n})/P_{K,1}(\mathfrak{n}) = \left\{ \left[(u\omega_{Q'_i} + v)\mathcal{O}_K \right] \mid \begin{bmatrix} u \\ v \end{bmatrix} \in \mathbb{Z}^2 / \equiv_N \text{ such that } \gcd(N, Q'_i(v, -u)) = 1 \right\}. \quad (18)$$

Now, let $C \in \text{Cl}(\mathfrak{n})$. By (17) and (18), there is one and only one pair $i \in \{1, 2, \dots, n\}$ and $\begin{bmatrix} u \\ v \end{bmatrix} \in \mathbb{Z}^2 / \equiv_N$ with $\gcd(N, Q'_i(v, -u)) = 1$ so that

$$C = \left[\frac{1}{u\omega_{Q'_i} + v} [\omega_{Q'_i}, 1] \right].$$

Take a matrix ρ in $\text{SL}_2(\mathbb{Z})$ satisfying

$$\rho \equiv \begin{bmatrix} * & * \\ u & v \end{bmatrix} \pmod{N}.$$

Since

$$\frac{j(\rho, \omega_{Q'_i})}{u\omega_{Q'_i} + v} \equiv^* 1 \pmod{\mathfrak{n}},$$

we get by Lemma 2.3 (i) that

$$C = \left[\frac{1}{j(\rho, \omega_{Q'_i})} [\omega_{Q'_i}, 1] \right] = [\rho(\omega_{Q'_i}), 1].$$

If $\tilde{Q} \in \mathcal{Q}_N(d_K)$ such that $\omega_{\tilde{Q}} = \rho(\omega_{Q'_i})$, then we attain

$$C = \phi_N([\tilde{Q}]) = \phi_N \left(\left[Q'_i \left(\rho^{-1} \begin{bmatrix} x \\ y \end{bmatrix} \right) \right] \right).$$

This completes the proof. □

EXAMPLE 4.5. Let $K = \mathbb{Q}(\sqrt{-2})$ and $N = 3$. There is only one reduced form

$$Q_1 = x^2 + 2y^2$$

of discriminant $d_K = -8$. Set $Q'_1 = Q_1$. By the proof of Proposition 4.4 one can find two representatives \tilde{Q}_1, \tilde{Q}_2 of form classes in $\text{C}_3(-8)$ whose zeros are

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (\omega_{Q'_1}) \quad \text{and} \quad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} (\omega_{Q'_1}).$$

And, we obtain

$$\text{C}_3(-8) = \{[x^2 + 2y^2], [2x^2 + y^2]\},$$

and hence $\text{C}_3(-8) \simeq \mathbb{Z}/2\mathbb{Z}$.

EXAMPLE 4.6. Let $K = \mathbb{Q}(\sqrt{-5})$ and $N = 2$. Then there are two reduced forms of discriminant $d_K = -20$, namely,

$$Q_1 = x^2 + 5y^2 \quad \text{and} \quad Q_2 = 2x^2 + 2xy + 3y^2.$$

Let

$$Q'_1 = Q_1 \quad \text{and} \quad Q'_2 = Q_2 \left(\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \right) = 3x^2 - 2xy + 2y^2$$

with

$$\omega_{Q'_1} = \sqrt{-5} \quad \text{and} \quad \omega_{Q'_2} = \frac{1 + \sqrt{-5}}{3}.$$

By the proof of Proposition 4.4 we have four representatives $\tilde{Q}_1, \tilde{Q}_2, \tilde{Q}_3, \tilde{Q}_4$ of form classes in $\text{C}_2(-20)$ with zeros

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (\omega_{Q'_1}), \quad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} (\omega_{Q'_1}), \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (\omega_{Q'_2}), \quad \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} (\omega_{Q'_2}), \quad \text{respectively.}$$

Thus, we achieve

$$C_2(-20) = \{[x^2 + 5y^2], [5x^2 + y^2], [3x^2 - 2xy + 2y^2], [7x^2 - 6xy + 2y^2]\}.$$

Note that

$$\tilde{Q}'_4 = \tilde{Q}_4 \left(\begin{bmatrix} 1 & 0 \\ 2 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \right) = 3x^2 + 2xy + 2y^2 \sim_2 \tilde{Q}_4.$$

We then see by using the argument in Remark 2.10 (iii) that

$$[\tilde{Q}_4]^{-1} = [\tilde{Q}'_4]^{-1} = [\tilde{Q}_3].$$

Therefore, we conclude

$$C_2(-20) \simeq \mathbb{Z}/4\mathbb{Z}.$$

EXAMPLE 4.7. Let $K = \mathbb{Q}(\sqrt{-5})$ and $N = 6$. Let Q_1 and Q_2 be reduced forms of discriminant $d_K = -20$ stated in Example 4.6. In this case, we let

$$Q'_1 = Q_1 \quad \text{and} \quad Q'_2 = Q_2 \left(\begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \right) = 7x^2 - 6xy + 2y^2$$

with

$$\omega_{Q'_1} = \sqrt{-5} \quad \text{and} \quad \omega_{Q'_2} = \frac{3 + \sqrt{-5}}{7}.$$

There are eight representatives of form classes in $C_6(-20)$ with zeros

$$\begin{aligned} & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (\omega_{Q'_1}), \quad \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} (\omega_{Q'_1}), \quad \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix} (\omega_{Q'_1}), \quad \begin{bmatrix} -1 & -1 \\ 3 & 2 \end{bmatrix} (\omega_{Q'_1}), \\ & \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} (\omega_{Q'_2}), \quad \begin{bmatrix} 0 & -1 \\ 1 & 3 \end{bmatrix} (\omega_{Q'_2}), \quad \begin{bmatrix} 1 & 1 \\ 2 & 3 \end{bmatrix} (\omega_{Q'_2}), \quad \begin{bmatrix} 1 & 0 \\ 3 & 1 \end{bmatrix} (\omega_{Q'_2}) \end{aligned}$$

by Proposition 4.4. Hence we attain

$$\begin{aligned} C_6(-20) = \{ & [x^2 + 5y^2], [5x^2 + y^2], [29x^2 - 26xy + 6y^2], [49x^2 + 34xy + 6y^2], \\ & [7x^2 - 6xy + 2y^2], [83x^2 + 48xy + 7y^2], [107x^2 - 80xy + 15y^2], [43x^2 - 18xy + 2y^2]\}. \end{aligned}$$

We close this section by mentioning that the equivalence relation on $\mathcal{Q}_N(d_K)$ induced from the congruence subgroup

$$\Gamma_0(N) = \left\{ \sigma \in \text{SL}_2(\mathbb{Z}) \mid \sigma \equiv \begin{bmatrix} * & * \\ 0 & * \end{bmatrix} \pmod{N} \right\}$$

gives us a new form class group as a quotient group of $C_N(d_K)$.

REMARK 4.8. Let $\mathcal{O} = [N\tau_K, 1]$ be an order of conductor N in K and $\text{Cl}(\mathcal{O})$ be the ideal class group of the order \mathcal{O} ([1, p. 123]). Then we know that

$$\text{Cl}(\mathcal{O}) \simeq I_K(\mathfrak{n})/P_{K, \mathbb{Z}}(\mathfrak{n}),$$

where $P_{K, \mathbb{Z}}(\mathfrak{n})$ is the subgroup of $I_K(\mathfrak{n})$ consisting of principal fractional ideals $\lambda \mathcal{O}_K$ such that

$$\lambda \equiv^* m \pmod{\mathfrak{n}} \quad \text{for some } m \in \mathbb{Z} \text{ with } \gcd(N, m) = 1$$

([1, Proposition 7.22]). And, by the existence theorem of class field theory, there is a unique abelian extension $H_{\mathcal{O}}$ of K for which

$$\text{Gal}(H_{\mathcal{O}}/K) \simeq I_K(\mathfrak{n})/P_{K, \mathbb{Z}}(\mathfrak{n})$$

via the Artin reciprocity map for \mathfrak{n} . Define an equivalence relation $\sim_{\mathcal{O}}$ on $\mathcal{Q}_N(d_K)$ by

$$Q \sim_{\mathcal{O}} Q' \iff Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \text{ for some } \sigma \in \Gamma_0(N).$$

Let

$$C_{\mathcal{O}}(d_K) = \mathcal{Q}_N(d_K) / \sim_{\mathcal{O}},$$

and define a map

$$\begin{aligned} \phi_{\mathcal{O}} : C_{\mathcal{O}}(d_K) &\rightarrow I_K(\mathfrak{n})/P_{K, \mathbb{Z}}(\mathfrak{n}) \\ [Q] &\mapsto \text{class containing } [\omega_Q, 1]. \end{aligned}$$

Then, in a similar way as we did in Propositions 2.6, 2.7 and 2.8 one can readily show that $\phi_{\mathcal{O}}$ is well defined and bijective. Therefore, we may consider the set $C_{\mathcal{O}}(d_K)$ as a quotient group of $C_N(d_K)$, which is isomorphic to $\text{Gal}(H_{\mathcal{O}}/K)$.

5 The maximal abelian extension unramified outside prime ideals dividing \mathfrak{n}

Let $K_{\mathfrak{n}}^{\text{ab}}$ be the maximal abelian extension of K unramified outside prime ideals dividing \mathfrak{n} ($= N\mathcal{O}_K$). If $N = 1$, then $K_{\mathfrak{n}}^{\text{ab}}$ is nothing but the Hilbert class field H_K of K . So, we assume $N \geq 2$. As an application, we shall describe $\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$ in view of extended form class groups. Here we shall regard $\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$ as a topological group equipped with Krull topology: For each $\rho \in \text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$, we take the cosets

$$\rho \text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/F)$$

as a basis of open neighborhoods of ρ , where F runs through all finite (abelian) subextensions of $K_{\mathfrak{n}}^{\text{ab}}/K$ ([8, §I.1]).

If L is a finite abelian extension of K unramified outside prime ideals dividing \mathfrak{n} , then its conductor also divides \mathfrak{n}^{ℓ} for some $\ell \geq 1$. Thus L is contained in the ray class field $K_{\mathfrak{n}^{\ell}}$ ([10, p. 116]), and hence we get

$$K_{\mathfrak{n}}^{\text{ab}} = \bigcup_{\ell \geq 1} K_{\mathfrak{n}^{\ell}}.$$

Furthermore, now that

$$K_{\mathfrak{n}} \subseteq K_{\mathfrak{n}^2} \subseteq \cdots \subseteq K_{\mathfrak{n}^{\ell}} \subseteq \cdots,$$

we attain the isomorphisms

$$\mathrm{Gal}(K_n^{\mathrm{ab}}/K) \simeq \varprojlim_{\ell} \mathrm{Gal}(K_n^{\ell}/K) \simeq \varprojlim_{\ell} C_{N^{\ell}}(d_K) \quad (19)$$

of topological groups by Theorem 3.8 ([11, §2 in Appendix]). Here, the inverse system $\{C_{N^{\ell}}(d_K)\}_{\ell}$ is given by the natural surjections $C_{N^n}(d_K) \leftarrow C_{N^m}(d_K)$ ($1 \leq n \leq m$). And we observe

$$\mathcal{Q}_{N^{\ell}}(d_K) = \mathcal{Q}_N(d_K) \quad \text{for all } \ell \geq 1.$$

For each $Q \in \mathcal{Q}_N(d_K)$ and $\ell \geq 1$, denote by

$$[Q]_{N^{\ell}} = \text{the form class containing } Q \text{ in } C_{N^{\ell}}(d_K).$$

Then we have

$$\varprojlim_{\ell} C_{N^{\ell}}(d_K) = \left\{ ([Q_1]_N, [Q_2]_{N^2}, \dots, [Q_{\ell}]_{N^{\ell}}, \dots) \in \prod_{\ell} C_{N^{\ell}}(d_K) \mid [Q_{\ell+1}]_{N^{\ell}} = [Q_{\ell}]_{N^{\ell}} \text{ for all } \ell \geq 1 \right\}.$$

Now, define an equivalence relation $\sim_{N^{\infty}}$ on the set $\mathcal{Q}_N(d_K)$ by

$$Q \sim_{N^{\infty}} Q' \iff Q \sim_{N^{\ell}} Q' \text{ for all } \ell \geq 1.$$

For each $Q \in \mathcal{Q}_N(d_K)$, let $[Q]_{N^{\infty}}$ be the form class containing Q in $\mathcal{Q}_N(d_K)/\sim_{N^{\infty}}$. We also define a map

$$\begin{aligned} \iota : \mathcal{Q}_N(d_K)/\sim_{N^{\infty}} &\rightarrow \varprojlim_{\ell} C_{N^{\ell}}(d_K) \\ [Q]_{N^{\infty}} &\mapsto ([Q]_N, [Q]_{N^2}, \dots, [Q]_{N^{\ell}}, \dots). \end{aligned}$$

Then it is straightforward that ι is well defined and injective.

LEMMA 5.1. *We derive*

$$\varprojlim_{\ell} C_{N^{\ell}}(d_K) = \overline{\iota(\mathcal{Q}_N(d_K)/\sim_{N^{\infty}})}.$$

PROOF. Let $([Q_1]_N, [Q_2]_{N^2}, \dots, [Q_{\ell}]_{N^{\ell}}, \dots) \in \varprojlim_{\ell} C_{N^{\ell}}(d_K)$ be given. For every $\ell \geq 1$, we see that

$$\begin{aligned} \iota([Q_{\ell}]_{N^{\infty}}) &= ([Q_{\ell}]_N, [Q_{\ell}]_{N^2}, \dots, [Q_{\ell}]_{N^{\ell}}, [Q_{\ell}]_{N^{\ell+1}}, \dots) \\ &= ([Q_1]_N, [Q_2]_{N^2}, \dots, [Q_{\ell}]_{N^{\ell}}, [Q_{\ell}]_{N^{\ell+1}}, \dots). \end{aligned}$$

Considering the Krull topology on $\mathrm{Gal}(K_n^{\mathrm{ab}}/K)$ we conclude that $\iota(\mathcal{Q}_N(d_K)/\sim_{N^{\infty}})$ is a dense subset of $\varprojlim_{\ell} C_{N^{\ell}}(d_K)$. \square

For $T = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, let us define a new equivalence relation \sim_T on $\mathcal{Q}_N(d_K)$ by

$$Q \sim_T Q' \iff Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma \begin{bmatrix} x \\ y \end{bmatrix} \right) \text{ for some } \sigma \in \langle -I_2, T \rangle.$$

LEMMA 5.2. *Two equivalence relations \sim_{N^∞} and \sim_T are the same.*

PROOF. Let $Q(x, y) = ax^2 + bxy + cy^2$ and $Q'(x, y) = a'x^2 + b'xy + c'y^2$ be two elements of $\mathcal{Q}_N(d_K)$. Since $\langle -I_2, T \rangle$ is contained in $\pm\Gamma_1(N^\ell)$ for all $\ell \geq 1$, it is immediate that if $Q \sim_T Q'$, then $Q \sim_{N^\infty} Q'$.

Conversely, assume that $Q \sim_{N^\infty} Q'$. Then, for each $\ell \geq 1$, there is $\sigma_\ell \in \pm\Gamma_1(N^\ell)$ such that

$$Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma_\ell \begin{bmatrix} x \\ y \end{bmatrix} \right).$$

Hence it follows from

$$Q \left(\sigma_1 \begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\sigma_\ell \begin{bmatrix} x \\ y \end{bmatrix} \right)$$

that

$$Q \left(\sigma_1 \sigma_\ell^{-1} \begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\begin{bmatrix} x \\ y \end{bmatrix} \right),$$

which yields that $\sigma_1 \sigma_\ell^{-1}$ belongs to the stabilizer subgroup $\text{Stab}(Q) (\subseteq \text{SL}_2(\mathbb{Z}))$ of Q . Since we are assuming $K \neq \mathbb{Q}(\sqrt{-1}), \mathbb{Q}(\sqrt{-3})$, $\text{Stab}(Q) = \{I_2, -I_2\}$; and hence $\sigma_1 = \sigma_\ell$ or $\sigma_1 = -\sigma_\ell$. Owing to the assumption $N \geq 2$ we achieve

$$\sigma_1 \in \bigcap_{\ell \geq 1} \pm\Gamma_1(N^\ell) = \langle -I_2, T \rangle.$$

Therefore, we conclude $Q \sim_T Q'$. □

LEMMA 5.3. *Let $Q(x, y) = ax^2 + bxy + cy^2$ and $Q'(x, y) = a'x^2 + b'xy + c'y^2$ be two elements of $\mathcal{Q}_N(d_K)$. Then,*

$$Q \sim_T Q' \iff a = a' \text{ and } a \text{ divides } \frac{b - b'}{2}.$$

PROOF. Observe that b and b' have the same parity by the discriminant condition

$$b^2 - 4ac = b'^2 - 4a'c' = d_K. \tag{20}$$

We then see that

$$\begin{aligned} Q \sim_T Q' &\iff Q' \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = Q \left(\begin{bmatrix} 1 & s \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} \right) \text{ for some } s \in \mathbb{Z} \\ &\iff a'x^2 + b'xy + c'y^2 = ax^2 + (2ax + b)xy + (a^2s + bs + c)y^2 \text{ for some } s \in \mathbb{Z} \\ &\iff a' = a \text{ and } b' = 2as + b \text{ for some } s \in \mathbb{Z} \text{ by (20)} \\ &\iff a = a' \text{ and } a \text{ divides } (b - b')/2. \end{aligned}$$

□

THEOREM 5.4. *The set $\mathcal{Q}_N(d_K)/\sim_T$ can be viewed as a dense subset of $\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$.*

PROOF. Let

$$\phi : \varprojlim_{\ell} C_{N^{\ell}}(d_K) \rightarrow \text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K)$$

be the isomorphism obtained in (19). Then we get by Lemmas 5.1 and 5.2

$$\text{Gal}(K_{\mathfrak{n}}^{\text{ab}}/K) = \overline{(\phi \circ \iota)(\mathcal{Q}_N(d_K)/\sim_T)}.$$

Moreover, Lemma 5.3 enables us to distinguish different classes in $\mathcal{Q}_N(d_K)/\sim_T$ from one another. \square

References

- [1] D. A. Cox, *Primes of the Form $x^2 + ny^2$: Fermat, Class field theory, and Complex Multiplication*, 2nd edn, Pure and Applied Mathematics (Hoboken), John Wiley & Sons, Inc., Hoboken, NJ, 2013.
- [2] M. Deuring, *Die Klassenkörper der komplexen Multiplikation*, Enzyklopädie der mathematischen Wissenschaften: Mit Einschluss ihrer Anwendungen, Band I 2, Heft 10, Teil II (Article I 2, 23), B. G. Teubner Verlagsgesellschaft, Stuttgart, 1958.
- [3] H. Hasse, *Neue Begründung der Komplexen Multiplikation I, II*, J. für die Reine und Angewandte Math. 157, 165 (1927, 1931), 115–139, 64–88.
- [4] G. J. Janusz, *Algebraic Number Fields*, 2nd edn, Grad. Studies in Math. 7, Amer. Math. Soc., Providence, R. I., 1996.
- [5] D. Kubert and S. Lang, *Modular Units*, Grundlehren der mathematischen Wissenschaften 244, Springer-Verlag, New York-Berlin, 1981.
- [6] S. Lang, *Elliptic Functions*, With an appendix by J. Tate, 2nd edn, Grad. Texts in Math. 112, Springer-Verlag, New York, 1987.
- [7] S. Lang, *Algebraic Number Theory*, 2nd edn, Grad. Texts in Math. 110, Springer-Verlag, New York, 1994.
- [8] J. Neukirch, *Class Field Theory*, Grundlehren der mathematischen Wissenschaften 280, Springer-Verlag, Berlin-Heidelberg, 1986.
- [9] K. Ramachandra, *Some applications of Kronecker's limit formula*, Ann. of Math. (2) 80 (1964), 104–148.
- [10] G. Shimura, *Introduction to the Arithmetic Theory of Automorphic Functions*, Iwanami Shoten and Princeton University Press, Princeton, N. J., 1971.
- [11] L. C. Washington, *Introduction to Cyclotomic Fields*, 2nd edn, Grad. Texts in Math. 83, Springer-Verlag, New York, 1997.

DEPARTMENT OF MATHEMATICS EDUCATION
DONGGUK UNIVERSITY-GYEONGJU
GYEONGJU-SI, GYEONGSANGBUK-DO 38066
REPUBLIC OF KOREA

E-mail address: zandc@dongguk.ac.kr

DEPARTMENT OF MATHEMATICS
HANKUK UNIVERSITY OF FOREIGN STUDIES
YONGIN-SI, GYEONGGI-DO 17035
REPUBLIC OF KOREA

E-mail address: dhshin@hufs.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES
KAIST
DAEJEON 34141
REPUBLIC OF KOREA

E-mail address: jkkoo@math.kaist.ac.kr