arXiv:1712.04253v1 [math.OC] 12 Dec 2017

Upper bounds for Z_1-eigenvalues of generalized Hilbert tensors *

Juan Meng†, Yisheng Song‡

Abstract

In this paper, we introduce the concept of Z_1-eigenvalue to infinite dimensional generalized Hilbert tensors (hypermatrix) $\mathcal{H}_\lambda^\infty = (\mathcal{H}_{i_1 i_2 \cdots i_m})$,

$$\mathcal{H}_{i_1 i_2 \cdots i_m} = \frac{1}{i_1 + i_2 + \cdots i_m + \lambda},\ \lambda \in \mathbb{R} \setminus \mathbb{Z}^-;\ i_1, i_2, \cdots, i_m = 0, 1, 2, \cdots, n, \cdots,$$

and proved that its Z_1-spectral radius is not larger than π for $\lambda > \frac{1}{2}$, and is at most $\frac{\pi}{\sin \lambda\pi}$ for $\frac{1}{2} \geq \lambda > 0$. Besides, the upper bound of Z_1-spectral radius of an mth-order n-dimensional generalized Hilbert tensor $\mathcal{H}_\lambda^n$ is obtained also, and such a bound only depends on n and λ.

Key words: Infinite-dimensional generalized Hilbert tensor, Z_1-eigenvalue, Spectral radius, Hilbert inqualities.
AMS subject classifications (2010): 47H15, 47H12, 34B10, 47A52, 47J10, 47H09, 15A48, 47H07

1 Introduction

A generalized Hilbert matrix has the form [13]:

$$H_\lambda^\infty = \left(\frac{1}{i + j + \lambda}\right)_{i,j \in \mathbb{Z}^+} \tag{1.1}$$

where $\mathbb{Z}^+$ ($\mathbb{Z}^-$) is the set of all non-negative (non-positive) integers and $\lambda \in \mathbb{R} \setminus \mathbb{Z}^-$. Denote such a Hilbert matrix with $i, j \in I_n = \{0, 1, 2, \cdots, n\}$ by H_λ^n. When $\lambda = 1$, such a matrix is called Hilbert matrix, which was introduced by Hilbert [12]. Choi [6] and Ingham [14] proved that Hilbert matrix H_1^∞ is a bounded linear operator (but not compact operator)

*This work was supported by the National Natural Science Foundation of P.R. China (Grant No. 11571095, 11601134, 11701154).

†School of Mathematics and Information Science, Henan Normal University, XinXiang HeNan, P.R. China, 453007. Email: 1015791785@qq.com

‡Corresponding author. School of Mathematics and Information Science and Henan Engineering Laboratory for Big Data Statistical Analysis and Optimal Control, Henan Normal University, XinXiang HeNan, P.R. China, 453007. Email: songyisheng@htu.cn.

from Hilbert space l^2 into itself. Magnus [18] and Kato [15] studied the spectral properties of H_1^∞. Frazer [7] and Taussky [29] discussed some nice properties of n-dimensional Hilbert matrix H_1^n. Rosenblum [23] showed that for a real $\lambda < 1$, H_λ^∞ defines a bounded operator on l^p for $2 < p < \infty$ and that $\pi \sec \pi u$ is an eigenvalue of H_λ^∞ for $|\Re u| < \frac{1}{2} - \frac{1}{p}$. For each non-integer complex number λ, Aleman, Montes-Rodríguez, Sarafoleanu [1] showed that H_λ^∞ defines a bounded linear operator on the Hardy spaces H^p $(1 < p < \infty)$.

As a natural extension of a generalized Hilbert matrix, the generalized Hilbert tensor (hypermatrix) was introduced by Mei and Song [24]. For each $\lambda \in \mathbb{R} \setminus \mathbb{Z}^-$, the entries of an mth-order infinite dimensional generalized Hilbert tensor $\mathcal{H}_\lambda^\infty = (\mathcal{H}_{i_1 i_2 \cdots i_m})$ are defined by

$$\mathcal{H}_{i_1 i_2 \cdots i_m} = \frac{1}{i_1 + i_2 + \cdots i_m + \lambda},\ i_1, i_2, \cdots, i_m = 0, 1, 2, \cdots, n, \cdots. \tag{1.2}$$

They showed $\mathcal{H}_\lambda^\infty$ defines a bounded and positively $(m-1)$-homogeneous operator from l^1 into l^p $(1 < p < \infty)$. Song and Qi [25] studied the operator properties of Hilbert tensors $\mathcal{H}_1^\infty$ and the spectral properties of $\mathcal{H}_1^n$. Such a tensor, $\mathcal{H}_\lambda^\infty$ may be refered to as a Hankel tensor with $v = (1, \frac{1}{2}, \frac{1}{3}, \cdots, \frac{1}{n}, \cdots)$. The concept of Hankel tensor was introduced by Qi [22]. For more further research of Hankel tensors, see Qi [22], Chen and Qi [5], Xu [31]. Denote such an mth-order n-dimensional generalized Hilbert tensor by $\mathcal{H}_\lambda^n$.

For a real vector $x = (x_1, x_2, \cdots, x_n, x_{n+1}, \cdots) \in l^1$, $\mathcal{H}_\lambda^\infty x^{m-1}$ is an infinite dimensional vector with its ith component defined by

$$(\mathcal{H}_\lambda^\infty x^{m-1})_i = \sum_{i_2, \cdots, i_m = 0}^{\infty} \frac{x_{i_2} \cdots x_{i_m}}{i + i_2 + \cdots + i_m + \lambda}, \lambda \in \mathbb{R} \setminus \mathbb{Z}^-;\ i = 0, 1, 2, \cdots. \tag{1.3}$$

Accordingly, $\mathcal{H}_\lambda^\infty x^m$ is given by

$$\mathcal{H}_\lambda^\infty x^m = \sum_{i_1, i_2, \cdots, i_m = 0}^{\infty} \frac{x_{i_1} x_{i_2} \cdots x_{i_m}}{i_1 + i_2 + \cdots + i_m + \lambda}, \lambda \in \mathbb{R} \setminus \mathbb{Z}^-. \tag{1.4}$$

Mei and Song [24] proved that $\mathcal{H}_\lambda^\infty x^m < \infty$ and $\mathcal{H}_\lambda^\infty x^{m-1} \in l^p$ $(1 < p < \infty)$ for all real vector $x \in l^1$.

In this paper, we will introduce the concept of Z_1-eigenvalue μ for an mth-order infinite dimensional generalized Hilbert tensor $\mathcal{H}_\lambda^\infty$ and will study some upper bounds of Z_1-spectral radius for infinite dimensional generalized Hilbert tensor $\mathcal{H}_\lambda^\infty$ and n-dimensional generalized Hilbert tensor $\mathcal{H}_\lambda^n$.

In Section 2, we will give some Lemmas and basic conclusions, and introduce the concept of Z_1-eigenvalue. In Section 3, with the help of the Hilbert type inequalities, the upper bound of Z_1–spectral radius of $\mathcal{H}_\lambda^\infty$ with $\lambda > 0$ is at most π when $\lambda > \frac{1}{2}$, and is not larger than $\frac{\pi}{\sin \lambda\pi}$ when $0 < \lambda \leq \frac{1}{2}$. Furthermore, for each Z_1-eigenvalue μ of $\mathcal{H}_\lambda^n$, $|\mu|$ is smaller than or equal to $C(n, \lambda)$, where $C(n, \lambda)$ only depends on the structured coefficient λ of generalized Hilbert tensor and the dimensionality n of European space.

2 Preliminaries and Basic Results

For $0 < p < \infty$, l^p is a space consisting of all real number sequences $x = (x_i)_{i=1}^{+\infty}$ satisfying $\sum\limits_{i=1}^{+\infty} |x_i|^p < \infty$. If $p \geq 1$, then a norm on l^p is defined by

$$\|x\|_{l^p} = \left(\sum_{i=1}^{+\infty} |x_i|^p\right)^{\frac{1}{p}}.$$

It is well known that l^2 is a Hilbert space with the inner product

$$\langle x, y\rangle = \sum_{i=0}^{+\infty} x_i y_i.$$

Clearly, $\|x\|_{l^2} = \sqrt{\langle x, x\rangle}$.

For $p \geq 1$, a norm $\mathbb{R}^n$ can be defined by

$$\|x\|_p = \left(\sum_{i=1}^{n} |x_i|^p\right)^{\frac{1}{p}}.$$

It is well known that

$$\|x\|_2 \leq \|x\|_1 \leq \sqrt{n}\|x\|_2. \tag{2.1}$$

The following Hilbert type inequalities were proved by Frazer [7] on $\mathbb{R}^n$ and Ingham [14] on l^2, respectively.

Lemma 2.1. *(Frazer [7]) Let $x = (x_1, x_2, \cdots, x_n)^\top \in \mathbb{R}^n$. Then*

$$\sum_{i=0}^{n}\sum_{j=0}^{n} \frac{|x_i||x_j|}{i+j+1} \leq (n \sin\frac{\pi}{n}) \sum_{k=0}^{n} x_k^2 = \|x\|_2^2 n \sin\frac{\pi}{n}, \tag{2.2}$$

Lemma 2.2. *(Ingham [14]) Let $x = (x_1, x_2, \cdots, x_n, \cdots)^\top \in l^2$ and $a > 0$. Then*

$$\sum_{i=0}^{\infty}\sum_{j=0}^{\infty} \frac{|x_i||x_j|}{i+j+a} \leq M(a) \sum_{k=0}^{\infty} x_k^2 = M(a)\|x\|_{l^2}^2, \tag{2.3}$$

where

$$M(a) = \begin{cases} \frac{\pi}{\sin a\pi}, & 0 < a \leq \frac{1}{2}; \\ \pi, & a > \frac{1}{2}. \end{cases}$$

An m-order n-dimensional tensor (hypermatrix) $\mathcal{A} = (a_{i_1\cdots i_m})$ is a multi-array of real entries $a_{i_1\cdots i_m} \in \mathbb{R}$, where $i_j \in I_n = \{1, 2, \cdots, n\}$ for $j \in [m] = \{1, 2, \cdots, m\}$. We use $T_{m,n}$ denote the set of all real mth-order n-dimensional tensors. Then $\mathcal{A} \in T_{m,n}$ is called a symmetric tensor if the entries $a_{i_1\cdots i_m}$are invariant under any permutation of their indices. $\mathcal{A} \in T_{m,n}$ is called nonnegative (positive) if $a_{i_1 i_2\cdots i_m} \geq 0 (a_{i_1 i_2\cdots i_m} > 0)$ for all $i_1, i_2, \cdots, i_m$.

Definition 2.1. (Chang and Zhang [2]) Let $\mathcal{A} \in T_{m,n}$. A number $\mu \in \mathbb{R}$ is called Z_1-eigenvalue of $\mathcal{A}$ if there is a real vector x such that

$$\begin{cases} \mathcal{A}x^{m-1} = \mu x \\ \|x\|_1 = 1 \end{cases} \tag{2.4}$$

and call such a vector x an Z_1-eigenvector associated with μ.

For the concepts of eigenvalues of higher order tensors, Qi [19, 20] first used and introduced them for symmetric tensors, and Lim [17] independently introduced this notion but restricted x to be a real vector and λ to be a real number. Subsequently, the spectral properties of nonnegative matrices had been generalized to n-dimensional nonnegative tensors under various conditions by Chang et al. [3, 4], He and Huang [9], He [10], He et al. [11], Li et al. [16], Qi [21], Song and Qi [26, 27], Wang et al. [30], Yang and Yang [32, 33] and references therein. The notion of Z_1-eigenvalue was introduced by Chang and Zhang [2] for higher Markov chains. Now we introduce it to infinite dimensional generalized Hilbert tensors.

Let

$$T_\infty x = \begin{cases} \|x\|_{l^1}^{2-m} \mathcal{H}_\lambda^\infty x^{m-1}, & x \neq \theta \\ \theta, & x = \theta, \end{cases} \tag{2.5}$$

where $\theta = (0, 0, \cdots, 0, \cdots)$. Mei and Song [24] first used the concept of the operator T_∞ induced by a generalized Hilbert tensor $\mathcal{H}_\lambda^\infty$ and showed T_∞ is a bounded and positively homogeneous operator from l^1 into l^p $(1 < p < \infty)$. Then T_∞ is refered to as a bounded and positively homogeneous operator from l^2 into l^2. So, the concept of Z_1-eigenvalue may be introduced to the infinite dimensional Hilbert tensor $\mathcal{H}_\lambda^\infty$.

Definition 2.2. Let $\mathcal{H}_\lambda^\infty$ be an mth-order infinite dimensional generalized Hilbert tensor. A real number μ is called a Z_1-eigenvalue of $\mathcal{H}_\lambda^\infty$ if there exists a nonzero vector $x \in l^2$ satisfying

$$T_\infty x = \|x\|_{l^1}^{2-m} \mathcal{H}_\lambda^\infty x^{m-1} = \mu x. \tag{2.6}$$

Such a vector x is called an Z_1-eigenvector associated with μ.

3 Main Results

Theorem 3.1. *Let $\mathcal{H}_\lambda^n$ be an mth-order n-dimensional generalized Hilbert tensor. Then*

$$|\mu| \leq C(n, \lambda) \text{ for all } Z_1\text{-eigenvalue } \mu \text{ of } \mathcal{H}_\lambda^n,$$

where $[\lambda]$ is the largest integer not exceeding λ and

$$C(n,\lambda) = \begin{cases} n \sin \frac{\pi}{n}, & \lambda \geq 1; \\ \frac{n}{\lambda}, & 1 > \lambda > 0; \\ \frac{n}{\min\{\lambda - [\lambda], 1 + [\lambda] - \lambda\}}, & -mn < \lambda < 0; \\ \frac{n}{-mn - a}, & \lambda < -mn. \end{cases}$$

Proof. For $\lambda \geq 1$, it follows from Lemma 2.1 that for all nonzero vector $x \in \mathbb{R}^n$,

$$\begin{aligned}
|\mathcal{H}_\lambda^n x^m| &= \left| \sum_{i_1,i_2,\cdots,i_m=0}^{n} \frac{x_{i_1}x_{i_2}\cdots x_{i_m}}{i_1+i_2+\cdots+i_m+\lambda} \right| \\
&\leq \sum_{i_1,\cdots,i_m=0}^{n} \frac{|x_{i_1}x_{i_2}\cdots x_{i_m}|}{i_1+i_2+\underbrace{0+\cdots+0}_{m-2}+\lambda} \\
&= \sum_{i_1,i_2,\cdots,i_m=0}^{n} \frac{|x_{i_1}||x_{i_2}|\cdots|x_{i_m}|}{i_1+i_2+\lambda} \\
&= \left(\sum_{i_1=0}^{n}\sum_{i_2=0}^{n} \frac{|x_{i_1}||x_{i_2}|}{i_1+i_2+\lambda} \right) \sum_{i_3,i_4,\cdots,i_m=0}^{n} |x_{i_3}||x_{i_4}|\cdots|x_{i_m}| \\
&\leq \left(\sum_{i_1=0}^{n}\sum_{i_2=0}^{n} \frac{|x_{i_1}||x_{i_2}|}{i_1+i_2+1} \right) \sum_{i_3,i_4,\cdots,i_m=0}^{n} |x_{i_3}||x_{i_4}|\cdots|x_{i_m}| \\
&\leq (\|x\|_2^2 n \sin\frac{\pi}{n}) \left(\sum_{i=0}^{n} |x_i| \right)^{m-2} \\
&= \|x\|_2^2 \|x\|_1^{m-2} n \sin\frac{\pi}{n}.
\end{aligned}$$

That is,

$$|\mathcal{H}_\lambda^n x^m| \leq \|x\|_2^2 \|x\|_1^{m-2} n \sin\frac{\pi}{n}. \tag{3.1}$$

Since μ is a Z_1-eigenvalue of $\mathcal{H}_\lambda^n$, then there exists a nonzero vector x such that

$$\mathcal{H}_\lambda^n x^{m-1} = \mu x \text{ and } \|x\|_1 = 1. \tag{3.2}$$

Thus, we have,

$$|\mu x^\top x| = |x^\top(\mathcal{H}_\lambda^n x^{m-1})| = |\mathcal{H}_\lambda^n x^m| \leq \|x\|_2^2 \|x\|_1^{m-2} n \sin\frac{\pi}{n},$$

and then,

$$|\mu| \|x\|_2^2 \leq \|x\|_2^2 \|x\|_1^{m-2} n \sin\frac{\pi}{n}.$$

As a result,

$$|\mu| \leq n \sin\frac{\pi}{n}. \tag{3.3}$$

For all $\lambda \in \mathbb{R} \setminus \mathbb{Z}^-$ with $\lambda < 1$, it is obvious that for $1 > \lambda > 0$,

$$\min_{i_1,\cdots,i_m \in I_n} |i_1+i_2+\cdots+i_m+\lambda| = \lambda.$$

For $-mn < \lambda < 0$, there exist some positive integers $i_1', i_2', \cdots, i_m'$ and $i_1'', i_2'', \cdots, i_m''$ such that

$$i_1'+i_2'+\cdots+i_m' = -[\lambda] \text{ and } i_1''+i_2''+\cdots+i_m'' = -[\lambda]-1,$$

and hence,

$$\min_{i_1,\cdots,i_m\in I_n} |i_1+i_2+\cdots+i_m+\lambda| = \min\{\lambda-[\lambda], \lambda-(-[\lambda]-1)\}.$$

For $\lambda < -mn$, we also have,

$$\min_{i_1,\cdots,i_m\in I_n} |i_1+i_2+\cdots+i_m+\lambda| = |mn+\lambda| = -mn-\lambda.$$

Therefore, we have for $\lambda \in \mathbb{R}\setminus\mathbb{Z}^-$ with $\lambda < 1$,

$$\frac{1}{|i_1+i_2+\cdots+i_m+\lambda|} \le N(\lambda) = \begin{cases} \frac{1}{\lambda}, & 1>\lambda>0; \\ \frac{1}{\min\{\lambda-[\lambda],1+[\lambda]-\lambda\}}, & -mn<\lambda<0; \\ \frac{1}{-mn-a}, & \lambda<-mn. \end{cases}$$

Then, for all nonzero vector $x\in\mathbb{R}^n$, we have

$$\begin{aligned} |\mathcal{H}_\lambda^n x^m| &= \left| \sum_{i_1,i_2,\cdots,i_m=0}^{n} \frac{x_{i_1}x_{i_2}\cdots x_{i_m}}{i_1+i_2+\cdots+i_m+\lambda} \right| \\ &\le \sum_{i_1,\cdots,i_m=0}^{n} \frac{|x_{i_1}x_{i_2}\cdots x_{i_m}|}{|i_1+i_2+\cdots+i_m+\lambda|} \\ &\le N(\lambda) \sum_{i_1,i_2,\cdots,i_m=0}^{n} |x_{i_1}||x_{i_2}|\cdots|x_{i_m}| \\ &= N(\lambda)\left(\sum_{i=0}^{n}|x_i|\right)^m = N(\lambda)\|x\|_1^m. \end{aligned}$$

For each Z_1-eigenvalue μ of $\mathcal{H}_\lambda^n$ with its eigenvector x, from (3.2) and $\|x\|_1 \le \sqrt{n}\|x\|_2$, it follows taht

$$|\mu|(\frac{1}{n}\|x\|_1^2) \le |\mu|\|x\|_2^2 = |\mathcal{H}_\lambda^n x^m| \le N(\lambda)\|x\|_1^m,$$

and hence,

$$|\mu| \le nN(\lambda).$$

This completes the proof. □

When $\lambda = 1$, the following conclusion of Hilbert tensor is easily obtained. Also see Song and Qi [25] for the conclusions about H-eigenvalue and Z-eigenvalue of such a tensor.

Corollary 3.2. *Let $\mathcal{H}$ be an mth-order n-dimensional Hilbert tensor. Then for all Z_1-eigenvalue μ of $\mathcal{H}$,*

$$|\mu| \le n\sin\frac{\pi}{n}.$$

Theorem 3.3. *Let $\mathcal{H}_\lambda^\infty$ be an mth-order infinite dimensional generalized Hilbert tensor. Assume $\lambda > 0$, then for Z_1-eigenvalue μ of $\mathcal{H}_\lambda^\infty$,*

$$|\mu| \le M(\lambda) = \begin{cases} \frac{\pi}{\sin\lambda\pi}, & 0<\lambda\le\frac{1}{2}; \\ \pi, & \lambda>\frac{1}{2}. \end{cases}$$

Proof. For $x \in l^2$, it follows from Lemma 2.2 that

$$\begin{aligned}
|\langle x, \mathcal{H}_\lambda^\infty x^{m-1}\rangle| = |\mathcal{H}_\lambda^\infty x^m| &= \left| \sum_{i_1,i_2,\cdots,i_m=0}^{+\infty} \frac{x_{i_1}x_{i_2}\cdots x_{i_m}}{i_1+i_2+\cdots+i_m+\lambda} \right| \\
&\leq \sum_{i_1,\cdots,i_m=0}^{+\infty} \frac{|x_{i_1}x_{i_2}\cdots x_{i_m}|}{i_1+i_2+\underbrace{0+\cdots+0}_{m-2}+\lambda} \\
&= \sum_{i_1,i_2,\cdots,i_m=0}^{+\infty} \frac{|x_{i_1}||x_{i_2}|\cdots|x_{i_m}|}{i_1+i_2+\lambda} \\
&= \left(\sum_{i_1=0}^{+\infty}\sum_{i_2=0}^{+\infty} \frac{|x_{i_1}||x_{i_2}|}{i_1+i_2+\lambda} \right) \sum_{i_3,i_4,\cdots,i_m=0}^{+\infty} |x_{i_3}||x_{i_4}|\cdots|x_{i_m}| \\
&= \left(\sum_{i_1=0}^{+\infty}\sum_{i_2=0}^{\infty} \frac{|x_{i_1}||x_{i_2}|}{i_1+i_2+\lambda} \right) \left(\sum_{i=0}^{+\infty} |x_i| \right)^{m-2} \\
&\leq M(\lambda)\|x\|_{l^2}^2 \|x\|_{l^1}^{m-2},
\end{aligned}$$

and so,

$$|\langle x, T_\infty x\rangle| = |\langle x, \|x\|_1^{2-m}\mathcal{H}_\lambda^\infty x^{m-1}\rangle| = \|x\|_{l^1}^{2-m}|\mathcal{H}_\lambda^\infty x^m| \leq M(\lambda)\|x\|_{l^2}^2. \tag{3.4}$$

For each Z_1-eigenvalue μ of $\mathcal{H}_\lambda^\infty$, there exists a nonzero vector $x \in l^2$ such that

$$T_\infty x = \|x\|_{l^1}^{2-m}\mathcal{H}_\lambda^\infty x^{m-1} = \mu x,$$

and so,

$$\mu\|x\|_{l^2}^2 = \mu\langle x, x\rangle = \langle x, \|x\|_{l^1}^{2-m}\mathcal{H}_\lambda^\infty x^{m-1}\rangle = \|x\|_{l^1}^{2-m}\mathcal{H}_\lambda^\infty x^m.$$

Therefore, we have

$$|\mu|\|x\|_{l^2}^2 = \|x\|_{l^1}^{2-m}|\mathcal{H}_\lambda^\infty x^m| \leq M(\lambda)\|x\|_{l^2}^2,$$

and then,

$$|\mu| \leq M(\lambda).$$

This completes the proof. □

When $\lambda = 1$, the following conclusion of infinite dimensional Hilbert tensor is easily obtained.

Corollary 3.4. *Let $\mathcal{H}_\infty$ be an mth-order infinite dimensional Hilbert tensor. Then for all Z_1-eigenvalue μ of $\mathcal{H}_\infty$,*

$$|\mu| \leq \pi.$$

Remark 3.1. *(i) In Theorem 3.1, the upper bound of Z_1-eigenvalue of $\mathcal{H}_\lambda^n$ are showed. However the upper bound may not be the best. Then which number is its best upper bounds?*

(ii) In Theorem 3.3, the upper bound of Z_1-eigenvalue of $\mathcal{H}_\lambda^\infty$ are showed for $\lambda > 0$, then for $\lambda < 0$ with $\lambda \in \mathbb{R} \setminus \mathbb{Z}^-$, it is unknown whether have similar conclusions or not. And it is not clear whether the upper bound may be attained or cannot be attained.

References

[1] A. Aleman, A. Montes-Rodríguez, A. Sarafoleanu, *The eigenfunctions of the Hilbert matrix*, Constr Approx. 36(2012), 353–374.

[2] K.C. Chang, K. Pearson, and T. Zhang, *On the uniqueness and non-uniqueness of the positive Z-eigenvector for transition probability tensors*, J. Math. Anal. Appl., 408(2013), 525-540.

[3] K.C. Chang, K. Pearson, and T. Zhang, *Some variational principles for Z-eigenvalues of nonnegative tensors*, Linear Algebra Appl. 438(2013) 4166-4182.

[4] K.C. Chang, K. Pearson, and T. Zhang, *Perron-Frobenius theorem for nonnegative tensors*, Commun. Math. Sci. 6(2008) 507-520.

[5] H. Chen, L. Qi, *Positive definiteness and semi-definiteness of even order symmetric Cauchy tensors* J. Ind. Manag. Optim. 11(4)(2015), 1263–1274.

[6] Man-Duen Choi, *Tricks for Treats with the Hilbert Matrix*, Amer. Math. Monthly 90(1983),301-312.

[7] H. Frazer, *Note on Hilbert's Inequality*, J. London Math. Soc. (1946) s1-21 (1): 7-9.

[8] S. Fucik, J. Necas, J. Soucek and V. Soucek, *Spectral Analysis of Nonlinear Operators*, Lecture Notes in Mathematics 346, Springer-Verlag, Berlin, Heidelberg, New York, 1973.

[9] J. He, T. Huang, *Upper bound for the largest Z-eigenvalue of positive tensors*, Appl. Math. Lett. 38(2014), 110-114.

[10] J. He, *Bounds for the largest eigenvalue of nonnegative tensors*, J. Comput. Anal. Appl. 20(7) (2016) 1290-1301.

[11] J. He, Y. M. Liu, H. Ke, J. K. Tian, X. Li, *Bounds for the Z-spectral radius of nonnegative tensors*, SpringerPlus, 5(1) (2016), 1727.

[12] D. Hilbert, *Ein Beitrag zur Theorie des Legendre'schen Polynoms*, Acta Mathematica (Springer Netherlands) 18(1894) 155-159.

[13] C.K.Hill, *On the singly-infinite Hilbert matrix*, J. London Math. Soc. 35(1960), 17-29.

[14] A. E. Ingham, *A Note on Hilbert's Inequality*, J. London Math. Soc. (1936) s1-11 (3): 237-240.

[15] T. Kato, *On the Hilbert Matrix*, Proc. American Math. Soc. 8(1)(1957) 73-81.

[16] W. Li, D. Liu, S.W. Vong. *Z-eigenpair bounds for an irreducible nonnegative tensor*, Linear Algebra Appl. 483 (2015) 182-199.

[17] L.H. Lim, *Singular values and eigenvalues of tensors: A variational approach*, in: Proc. 1st IEEE International workshop on computational advances of multi-tensor adaptive processing, Dec. 13-15, 2005, pp. 129-132.

[18] W. Magnus *On the Spectrum of Hilbert's Matrix*, American J. Math. 72(4)(1950) 699-704.

[19] L. Qi, *Eigenvalues of a real supersymmetric tensor*, J. Symbolic Comput. 40(2005) 1302-1324.

[20] L. Qi, *Rank and eigenvalues of a supersymmetric tensor, the multivariate homogeneous polynomial and the algebraic hypersurface it defines*, J. Symbolic Comput. 41(2006) 1309-1327.

[21] L. Qi, *Symmetric nonnegative tensors and copositive tensors*, Linear Algebra Appl., 439(2013) 228-238.

[22] L. Qi, *Hankel Tensors: Associated Hankel Matrices and Vandermonde Decomposition*, Commun. Math. Sci. 13(1)(2015) 113-125.

[23] M. Rosenblum, *On the Hilbert matrix I*, Proc.Am. Math. Soc. 9(1958), 137-140.

[24] W. Mei, Y. Song, *Infinite and finite dimensional generalized Hilbert tensors*, Linear Algebra Appl. 532(2017) 8-24.

[25] Y. Song, L. Qi, *Infinite and finite dimensional Hilbert tensors*, Linear Algebra Appl. 451 (2014) 1-14.

[26] Y. Song and L. Qi, *Positive eigenvalue-eigenvector of nonlinear positive mappings*, Front. Math. China, 9(1)(2014) 181-199.

[27] Y. Song and L. Qi, *Spectral properties of positively homogeneous operators induced by higher order tensors*, SIAM. J. Matrix Anal. Appl., 34(4)(2013) 1581-1595.

[28] Y. Song, L. Qi, *Necessary and sufficient conditions for copositive tensors*, Linear Multilinear Algebra 63(1)(2015), 120–131.

[29] O. Taussky, *A remark concerning the characteristic roots of the finite segments of the Hilbert matrix,* Quarterly J. Math. Oxford ser., vol. 20 (1949) 80-83.

[30] Y. Wang, K. Zhang, H. Sun, *Criteria for strong H-tensors*, Front. Math. China 11.3 (2016) 577-592.

[31] C. Xu, *Hankel tensors, Vandermonde tensors and their positivities*, Linear Algebra Appl. 491 (2016) 56-72.

[32] Q. Yang and Y. Yang, *Further Results for Perron-Frobenius Theorem for Nonnegative Tensors II*, SIAM. J. Matrix Anal. Appl. 32(4)(2011) 1236-1250.

[33] Y. Yang and Q. Yang, *Further Results for Perron-Frobenius Theorem for Nonnegative Tensors*, SIAM. J. Matrix Anal. Appl., 31(5)(2010) 2517-2530.