

# Strongly order continuous operators on Riesz spaces

by

AKBAR BAHRAMNEZHAD AND KAZEM HAGHNEJAD AZAR

Department of Mathematic, University of Mohaghegh Ardabili, Ardabil, Iran

E-mail: bahramnezhad@uma.ac.ir

## Abstract

In this paper we introduce two new classes of operators that we call strongly order continuous and strongly  $\sigma$ -order continuous operators. An operator  $T : E \rightarrow F$  between two Riesz spaces is said to be strongly order continuous (resp. strongly  $\sigma$ -order continuous), if  $x_\alpha \xrightarrow{uo} 0$  (resp.  $x_n \xrightarrow{uo} 0$ ) in  $E$  implies  $Tx_\alpha \xrightarrow{o} 0$  (resp.  $Tx_n \xrightarrow{o} 0$ ) in  $F$ . We give some conditions under which order continuity will be equivalent to strongly order continuity of operators on Riesz spaces. We show that the collection of all *so*-continuous linear functionals on a Riesz space  $E$  is a band of  $E^\sim$ .

**Key Words:** Riesz space, order convergence, unbounded order convergence, strongly order continuous operator.

**2010 Mathematics Subject Classification:** Primary 46B42, 47B60.

## 1 Introduction

The concept of unbounded order convergence or *uo*-convergence was introduced in [8] and is proposed firstly in [3]. It has recently been intensively studied in several papers [4, 5, 6]. Recall that a net  $(x_\alpha)_{\alpha \in \mathcal{A}}$  in a Riesz space  $E$  is *order convergent* (or, *o*-convergent for short) to  $x \in E$ , denoted by  $x_\alpha \xrightarrow{o} x$  whenever there exists another net  $(y_\beta)_{\beta \in \mathcal{B}}$  in  $E$  such that  $y_\beta \downarrow 0$  and that for every  $\beta \in \mathcal{B}$ , there exists  $\alpha_0 \in \mathcal{A}$  such that  $|x_\alpha - x| \leq y_\beta$  for all  $\alpha \geq \alpha_0$ . A net  $(x_\alpha)$  in a Riesz space  $E$  is *unbounded order convergent* (or, *uo*-convergent for short) to  $x \in E$  if  $|x_\alpha - x| \wedge u \xrightarrow{o} 0$  for all  $u \in E^+$ . We denote this convergence by  $x_\alpha \xrightarrow{uo} x$  and write that  $x_\alpha$  *uo*-convergent to  $x$ . This is an analogue of pointwise convergence in function spaces. Let  $\mathbb{R}^A$  be the Riesz space of all real-valued functions on a non-empty set  $A$ , equipped with the pointwise order. It is easily seen that a net  $(x_\alpha)$  in  $\mathbb{R}^A$  *uo*-converges to  $x \in \mathbb{R}^A$  if and only if it converges pointwise to  $x$ . For instance in  $c_0$  and  $\ell_p(1 \leq p \leq \infty)$ , *uo*-convergence of nets is the same as coordinate-wise convergence. Assume that  $(\Omega, \Sigma, \mu)$  is a measure space and let  $E = L_p(\mu)$  for some  $1 \leq p < \infty$ . Then *uo*-convergence of sequences in  $L_p(\mu)$  is the same as almost everywhere convergence. In [9], Wickstead characterized the spaces in which weak convergence of nets implies *uo*-convergence and vice versa. In [4], Gao characterized the space  $E$  such that in its dual space  $E^*$ , *uo*-convergence implies  $w^*$ -convergence and vice versa. He also characterized the spaces in whose dual space simultaneous *uo*- and  $w^*$ -convergence imply weak/norm convergence. We show that the collection of all order bounded strongly order continuous linear functionals on a Riesz space  $E$  is a band of  $E^\sim$  [Theorem 2.7]. For unexplained terminology and facts on Banach lattices and positive operators, we refer the reader to [1, 2]. Let us start with the definition.

**Definition 1.** An operator  $T : E \rightarrow F$  between two Riesz spaces is said to be:

1. *Strongly order continuous (or, so-continuous for short), if  $x_\alpha \xrightarrow{uo} 0$  in  $E$  implies  $Tx_\alpha \xrightarrow{o} 0$  in  $F$ .*
2. *Strongly  $\sigma$ -order continuous (or,  $s\sigma o$ -continuous for short), if  $x_n \xrightarrow{uo} 0$  in  $E$  implies  $Tx_n \xrightarrow{o} 0$  in  $F$ .*

The collection of all *so*-continuous operators of  $L_b(E, F)$  will be denoted by  $L_{so}(E, F)$ , that is

$$L_{so}(E, F) := \{T \in L_b(E, F) : T \text{ is } so\text{-continuous}\}.$$

Similarly,  $L_{s\sigma o}(E, F)$  will denote the collection of all order bounded operators from  $E$  to  $F$  that are *s $\sigma o$* -continuous. That is,

$$L_{s\sigma o}(E, F) := \{T \in L_b(E, F) : T \text{ is } s\sigma o\text{-continuous}\}.$$

Clearly, we have  $L_{so}(E, F) \subset L_{s\sigma o}(E, F)$  and  $L_{so}(E, F)$  and  $L_{s\sigma o}(E, F)$  are both vector subspaces of  $L_b(E, F)$ .

Recall that an operator  $T : E \rightarrow F$  between two Riesz spaces is said to be order continuous (resp.  $\sigma$ -order continuous) if  $x_\alpha \xrightarrow{o} 0$  (resp.  $x_n \xrightarrow{o} 0$ ) in  $E$  implies  $Tx_\alpha \xrightarrow{o} 0$  (resp.  $Tx_n \xrightarrow{o} 0$ ) in  $F$ . The collection of all order continuous operators of  $L_b(E, F)$  will be denoted by  $L_n(E, F)$ , that is

$$L_n(E, F) := \{T \in L_b(E, F) : T \text{ is order continuous}\}.$$

Similarly,  $L_c(E, F)$  will denote the collection of all order bounded operators from  $E$  to  $F$  that are  $\sigma$ -order continuous. That is,

$$L_c(E, F) := \{T \in L_b(E, F) : T \text{ is } \sigma\text{-order continuous}\}.$$

Note that every *so*-continuous (resp. *s $\sigma o$* -continuous) operator is order (resp.  $\sigma$ -order) continuous. The converse is not true in general. For example the identity operator  $I : c_0 \rightarrow c_0$  is order continuous but is not *so*-continuous. Indeed, the standard basis sequence of  $c_0$  is *uo*-converges to 0 but is not order convergent.

## 2 Main Results

**Lemma 1.** ([5, Lemma 3.1]). *In a Riesz space we have the following:*

1. *If  $x_\alpha \xrightarrow{uo} x$  and  $x_\alpha \xrightarrow{uo} y$ , then  $x = y$ . In other hands, unbounded order limits are uniquely determined.*
2. *If  $x_\alpha \xrightarrow{uo} x$ ,  $y_\alpha \xrightarrow{uo} y$  and  $k, r$  are real numbers, then  $kx_\alpha + ry_\alpha \xrightarrow{uo} kx + ry$ . Furthermore  $x_\alpha \vee y_\alpha \xrightarrow{uo} x \vee y$  and  $x_\alpha \wedge y_\alpha \xrightarrow{uo} x \wedge y$ . In particular  $x_\alpha^+ \xrightarrow{uo} x^+$ ,  $x_\alpha^- \xrightarrow{uo} x^-$  and  $|x_\alpha| \xrightarrow{uo} |x|$ .*
3. *If  $x_\alpha \xrightarrow{uo} x$  and  $x_\alpha \geq y$  holds for all  $\alpha$ , then  $x \geq y$ .*

Note that the  $uo$ -convergence in a Riesz space  $E$  does not necessarily correspond to a topology on  $E$ . For example, let  $E = c$ , the Banach lattice of real valued convergent sequences. Put  $x_n = \sum_{k=1}^n e_k$ , where  $(e_n)$  is the standard basis. Then  $(x_n)$  is  $uo$ -convergent to  $x = (1, 1, 1, \dots)$ , but it is not norm convergent.

**Proposition 1.** *Let  $E, F$  be Riesz spaces such that  $E$  is finite-dimensional. Then  $L_{so}(E, F) = L_n(E, F)$  and  $L_{s\sigma o}(E, F) = L_c(E, F)$ .*

*Proof.* Follows immediately if we observe that in a finite-dimensional Riesz space order convergence is equivalent to  $uo$ -convergence.  $\square$

Recall that a Riesz space is said to be  $\sigma$ -laterally complete if every disjoint sequence has a supremum. For a set  $A$ ,  $\mathbb{R}^A$  is an example of  $\sigma$ -laterally complete Riesz space.

**Proposition 2.** *Let  $E, F, G$  be Riesz spaces. Then we have the following:*

1. *If  $T \in L_{so}(E, F)$  and  $S \in L_n(F, G)$ , then  $ST \in L_{so}(E, G)$ . As a consequence,  $L_{so}(E)$  is a left ideal for  $L_n(E)$ . Similarly,  $L_{s\sigma o}(E)$  is a left ideal for  $L_c(E)$ .*
2. *If  $E$  is  $\sigma$ -Dedekind complete and  $\sigma$ -laterally complete and  $S \in L_c(E, F)$  and  $T \in L_{s\sigma o}(F, G)$ , then  $TS \in L_{s\sigma o}(E, G)$ . In this case,  $L_{s\sigma o}(E, F) = L_c(E, F)$ .*

*Proof.* 1. Let  $(x_\alpha)$  be a net in  $E$  such that  $x_\alpha \xrightarrow{uo} 0$ . By assumption,  $Tx_\alpha \xrightarrow{o} 0$ . So,  $STx_\alpha \xrightarrow{o} 0$ . Hence,  $ST \in L_{so}(E, G)$ .

2. Let  $E$  be a  $\sigma$ -Dedekind complete and  $\sigma$ -laterally complete Riesz space. By Theorem 3.9 of [6], we see that a sequence  $(x_n)$  in  $E$  is  $uo$ -null if and only if it is order null. So, if  $(x_n)$  be a sequence in  $E$  such that  $x_n \xrightarrow{uo} 0$ , then  $x_n \xrightarrow{o} 0$ . Thus,  $Sx_n \xrightarrow{o} 0$  and then  $TSx_n \xrightarrow{o} 0$ . Hence,  $TS \in L_{s\sigma o}(E, G)$ . Clearly, we have  $L_{s\sigma o}(E, F) = L_c(E, F)$ . This ends the proof.  $\square$

Let  $T : E \rightarrow F$  be a positive operator between Riesz spaces. We say that an operator  $S : E \rightarrow F$  is dominated by  $T$  (or that  $T$  dominates  $S$ ) whenever  $|Sx| \leq T|x|$  holds for each  $x \in E$ .

**Proposition 3.** *If a positive  $so$ -continuous operator  $T : E \rightarrow F$  dominates  $S$ , then  $S$  is  $so$ -continuous.*

*Proof.* Let  $T : E \rightarrow F$  be a positive  $so$ -continuous operator between Riesz spaces such that  $T$  dominates  $S : E \rightarrow F$  and let  $x_\alpha \xrightarrow{uo} 0$  in  $E$ . By part (2) of Lemma 1,  $|x_\alpha| \xrightarrow{uo} 0$ . So, by assumption,  $T|x_\alpha| \xrightarrow{o} 0$  and from the inequality  $|Sx| \leq T|x|$  and part (2) of Lemma 1 again, we have  $Sx_\alpha \xrightarrow{o} 0$ . Hence,  $S$  is  $so$ -continuous.  $\square$

For an operator  $T : E \rightarrow F$  between two Riesz spaces we shall say that its modulus  $|T|$  exists (or that  $T$  possesses a modulus) whenever  $|T| := T \vee (-T)$  exists in the sense that  $|T|$  is the supremum of the set  $\{-T, T\}$  in  $L(E, F)$ . If  $E$  and  $F$  are Riesz spaces with  $F$  Dedekind complete, then every order bounded operator  $T : E \rightarrow F$  possesses a modulus [2, Theorem 1.18]. From this discussion it follows that when  $E$  and  $F$  are Riesz spaces with  $F$  Dedekind complete, then each order bounded operator  $T : E \rightarrow F$  satisfies

$$T^+(x) = \sup\{Ty : 0 \leq y \leq x\}, \text{ and}$$

$$T^-(x) = \sup\{-Ty : 0 \leq y \leq x\}$$

for each  $x \in E^+$ .

**Theorem 1.** *For an order bounded linear functional  $f$  on a Riesz space  $E$  the following statements are equivalent.*

1.  $f$  is so-continuous.
2.  $f^+$  and  $f^-$  are both so-continuous.
3.  $|f|$  is so-continuous.

*Proof.* (1)  $\Rightarrow$  (2) By Lemma 1, we may assume that  $(x_\alpha) \subset E^+$ . Let  $x_\alpha \xrightarrow{uo} 0$  and let  $(r_\alpha)$  be a net in  $\mathbb{R}$  such that  $r_\alpha \downarrow 0$ . In view of  $f^+x = \sup\{fy : 0 \leq y \leq x\}$ , for each  $\alpha$  there exists a net  $(y_\alpha)$  in  $E$  with  $0 \leq y_\alpha \leq x_\alpha$  such that  $f^+x_\alpha - r_\alpha \leq fy_\alpha$ . So,  $f^+x_\alpha \leq fy_\alpha + r_\alpha$ . Since  $x_\alpha \xrightarrow{uo} 0$ , we have  $y_\alpha \xrightarrow{uo} 0$ . Thus, by assumption,  $fy_\alpha \xrightarrow{o} 0$ . It follows from  $f^+x_\alpha \leq (fy_\alpha + r_\alpha) \xrightarrow{o} 0$  that  $f^+x_\alpha \xrightarrow{o} 0$ . Hence,  $f^+$  is so-continuous. Now, as  $f^- = (-f)^+$ , we conclude that  $f^-$  is also so-continuous.

(2)  $\Rightarrow$  (3) Follows from the identity  $|f| = f^+ + f^-$ .

(3)  $\Rightarrow$  (1) Follows immediately from Proposition 3 by observing that  $|f|$  dominates  $f$ .  $\square$

**Remark 1.** *One can easily formulate by himself the analogue of Theorem 1 for  $s\sigma o$ -continuous operators.*

Recall that a subset  $A$  of a Riesz space is said to be order closed whenever  $(x_\alpha) \subset A$  and  $x_\alpha \xrightarrow{o} x$  imply  $x \in A$ . An order closed ideal is referred to as a band. Thus, an ideal  $A$  is a band if and only if  $(x_\alpha) \subset A$  and  $0 \leq x_\alpha \uparrow x$  imply  $x \in A$ . In the next theorem we show that  $L_{so}(E, \mathbb{R})$  and  $L_{s\sigma o}(E, \mathbb{R})$  are both bands of  $E^\sim$ . The details follow.

**Theorem 2.** *If  $E$  is a Riesz space, then  $L_{so}(E, \mathbb{R})$  and  $L_{s\sigma o}(E, \mathbb{R})$  are both bands of  $E^\sim$ .*

*Proof.* We only show that  $L_{so}(E, \mathbb{R})$  is a band of  $E^\sim$ . That  $L_{s\sigma o}(E, \mathbb{R})$  is a band can be proven in a similar manner. Note first that if  $|g| \leq |f|$  holds in  $E^\sim$  with  $f \in L_{so}(E, \mathbb{R})$ , then from Theorem 1 it follows that  $g \in L_{so}(E, \mathbb{R})$ . That is  $L_{so}(E, \mathbb{R})$  is an ideal of  $E^\sim$ . To see that the ideal  $L_{so}(E, \mathbb{R})$  is a band, let  $0 \leq f_\lambda \uparrow f$  in  $E^\sim$  with  $(f_\lambda) \subset L_{so}(E, \mathbb{R})$ , and let  $0 \leq x_\alpha \xrightarrow{uo} 0$  in  $E$ . Then for each fixed  $\lambda$  we have

$$0 \leq f(x_\alpha) = ((f - f_\lambda)(x_\alpha) + f_\lambda(x_\alpha)) \xrightarrow{o} 0.$$

So,  $f(x_\alpha) \xrightarrow{o} 0$ . Thus,  $f \in L_{so}(E, \mathbb{R})$ , and the proof is finished.  $\square$

**Problem 1.** *Can we find a so-continuous operator  $T : E \rightarrow F$  between Riesz spaces whose modulus is not so-continuous?*

## References

- [1] Y. A. Abramovich, C. D. Aliprantis, *An Invitation to Operator Theory*, vol. **50**, American Mathematical Society, Providence (2002).
- [2] C.D. Aliprantis, and O. Burkinshaw, *Positive Operators*, Springer, Berlin, (2006).
- [3] R. Demarr, *Partially ordered linear spaces and locally convex linear topological spaces*, Illinois J. Math., **8**, 601-606 (1964).
- [4] N. Gao, *Unbounded order convergence in dual spaces*, J. Math. Anal. Appl. **419**(1), 347-354 (2014).
- [5] N. Gao and F. Xanthos, *Unbounded order convergence and application to martingales without probability*, J. Math. Anal. Appl., **415**, 931-947 (2014).
- [6] N. Gao, V.G. Troitsky, and F. Xanthos, *Uo-convergence and its applications to cesaro means in Banach lattices*, Israel J. Math., appear. arXiv:1509.07914 [math.FA].
- [7] S. Kaplan, *On unbounded order convergence*, Real Analysis exchange, **23** (1), 75-184 (1998-99).
- [8] H. Nakano, *Ergodic theorems in semi-ordered linear spaces*, Ann. of Math. (2), **49**, 538-556 (1948).
- [9] A. W. Wickstead, *Weak and unbounded order convergence in Banach lattices*, J. Austral. Math. Soc. Ser. A **24**, 312-319 (1977).