

Stochastic Thermodynamic Interpretation of Information Geometry

Sosuke Ito

RIES, Hokkaido University, N20 W10, Kita-ku, Sapporo, Hokkaido 001-0020, Japan

(Dated: March 2, 2022)

In recent years, the unified theory of information and thermodynamics has been intensively discussed in the context of stochastic thermodynamics. The unified theory reveals that information theory would be useful to understand non-stationary dynamics of systems far from equilibrium. In this letter, we have found a new link between stochastic thermodynamics and information theory well known as information geometry. By applying this link, information geometric inequality can be interpreted as thermodynamic uncertainty relationship between speed and thermodynamic cost. We have numerically applied information geometric inequality to a thermodynamic model of biochemical enzyme reaction.

PACS numbers: 02.40.-k, 05.20.-y, 05.40.-a, 05.70.Ln, 89.70.-a

The crucial relationship between thermodynamics and information theory has been well studied in last decades [1]. Historically, thermodynamic-informational links had been discussed in the context of the second law of thermodynamics and the paradox of Maxwell's demon [2]. Recently, several studies have newly revealed thermodynamic interpretation of informational quantities such as the Kullback-Leibler divergence [3], mutual information [4–6], the transfer entropy and information flow [7–19]. The above interpretations of informational quantities are based on the theory of stochastic thermodynamics [20, 21], which mainly focus on the entropy production in stochastic dynamics of systems far from equilibrium.

Information thermodynamic relationship has been attracted not only in terms of Maxwell's demon, but also in terms of geometry [22, 23, 25–28]. Indeed, differential geometric interpretations of thermodynamics have been discussed especially in the near-equilibrium system [22, 29–34]. Moreover, the technique of differential geometry in information theory, well known as information geometry [35], has received remarkable attention in the field of neuroscience, signal processing, quantum mechanics, and machine learning [36–38]. In spite of the deep link between information and thermodynamics, the direct connection of thermodynamics and information geometry has been elusive especially for non-stationary and non-equilibrium dynamics. For example, G. E. Crooks discovered the link of thermodynamics and information geometry [22, 32] based on the Gibbs ensemble, and then his discussion is only valid for a near-equilibrium system.

In this letter, we discover the fundamental link between information geometry and thermodynamics based on stochastic thermodynamics for the master equation. We obtain thermodynamic interpretations of information geometric quantities, and derive thermodynamic inequalities based on information geometry. Our result can be interpreted as a kind of thermodynamic uncertainty relationships or thermodynamic trade-off relationships [39–50] between thermodynamic cost and speed. We numer-

ically test our trade-off relationship by using a model of biochemical enzyme reaction.

Stochastic thermodynamics.– To clarify the link between stochastic thermodynamics and information geometry, we here start with a formalism of stochastic thermodynamics for the master equation, that is known as the Schnakenberg network theory [51, 52].

We here consider $(n + 1)$ -states system. We assume that transitions between states are induced by n_{bath} -multiple thermal baths. The master equation for the probability p_x (≥ 0 , $\sum_{x=0}^n p_x = 1$) to find the state at $x = \{0, 1, \dots, n\}$ is given by

$$\frac{d}{dt}p_x = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x' \rightarrow x}^{(\nu)} p_{x'}, \quad (1)$$

where $W_{x' \rightarrow x}^{(\nu)}$ is the transition rate from x' to x induced by ν -th thermal bath. We assume that any transition rates has a finite value $W_{x' \rightarrow x}^{(\nu)} > 0$ for $x \neq x'$. We also assume the condition

$$\sum_{x=0}^n W_{x' \rightarrow x}^{(\nu)} = 0, \quad (2)$$

or equivalently $W_{x' \rightarrow x'}^{(\nu)} = -\sum_{x \neq x'} W_{x' \rightarrow x}^{(\nu)} < 0$, which leads to the conservation of probability $\sum_{x=0}^n dp_x/dt = 0$. This equation (2) indicates that the master equation is then given by the thermodynamic flux from the state x' to x [51],

$$J_{x' \rightarrow x}^{(\nu)} := W_{x' \rightarrow x}^{(\nu)} p_{x'} - W_{x \rightarrow x'}^{(\nu)} p_x, \quad (3)$$

$$\frac{d}{dt}p_x = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n J_{x' \rightarrow x}^{(\nu)}. \quad (4)$$

In a stationary state, the sum of the thermodynamic flux is zero for any x , i.e., $dp_x/dt = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n J_{x' \rightarrow x}^{(\nu)} = 0$. Especially, in a stationary state where the detailed balance (or the reversibility) holds (i.e., $J_{x' \rightarrow x}^{(\nu)} = 0$ for any

x , x' and ν), the system is said to be in thermodynamic equilibrium. If we consider the conjugated thermodynamic force

$$F_{x' \rightarrow x}^{(\nu)} := \ln[W_{x' \rightarrow x}^{(\nu)} p_{x'}] - \ln[W_{x \rightarrow x'}^{(\nu)} p_x], \quad (5)$$

thermodynamic equilibrium is equivalently given by $F_{x' \rightarrow x}^{(\nu)} = 0$ for any x , x' and ν .

In stochastic thermodynamics [21], we treat the entropy change of thermal bath and the system in a stochastic way. In the transition from x' to x , the stochastic entropy change of ν -th thermal bath is defined as

$$\dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} := \ln \frac{W_{x' \rightarrow x}^{(\nu)}}{W_{x \rightarrow x'}^{(\nu)}}, \quad (6)$$

and the stochastic entropy change of the system is defined as the stochastic Shannon entropy change

$$\dot{s}_{x' \rightarrow x}^{\text{sys}} := \ln p_{x'} - \ln p_x, \quad (7)$$

respectively. The thermodynamic force is then given by the total stochastic entropy change in the transition from x' to x induced by ν -th thermal bath $F_{x' \rightarrow x}^{(\nu)} = \dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} + \dot{s}_{x' \rightarrow x}^{\text{sys}}$.

To simplify notations, we introduce the set of directed edges $E = \{(x' \rightarrow x, \nu) | 0 \leq x' < x \leq n, 1 \leq \nu \leq n_{\text{bath}}\}$ which implies the set of all possible transitions between two states. The total entropy change rate is given by the sum of the products of thermodynamic forces and fluxes over possible edges

$$\dot{S}^{\text{tot}} := \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} F_{x' \rightarrow x}^{(\nu)} = \langle F \rangle, \quad (8)$$

where a parenthesis $\langle \dots \rangle$ is defined as $\langle A \rangle := \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} A_{x' \rightarrow x}^{(\nu)}$ for any function of edge $A_{x' \rightarrow x}^{(\nu)}$. Because the signs of thermodynamic force $F_{x' \rightarrow x}^{(\nu)}$ and flux $J_{x' \rightarrow x}^{(\nu)}$ are same, we obtain the non-negativity of the total entropy change rate,

$$\dot{S}^{\text{tot}} = \langle F \rangle = \langle \dot{s}^{\text{bath}} \rangle + \langle \dot{s}^{\text{sys}} \rangle \geq 0, \quad (9)$$

that is well known as the second law of thermodynamics.

Information geometry.— Next, we introduce the theory of information geometry [35]. In information geometry, we consider the set of probabilities that satisfies the normalization. Due to the constraint of the normalization, the set of probability gives a Riemannian manifold, and its natural Riemannian metric is given by the Fisher metric [53]. In this letter, we only consider the discrete distribution group.

The discrete distribution group $\mathbf{p} = (p_0, p_1, \dots, p_n)$, $p_x \geq 0$, and $\sum_{x=0}^n p_x = 1$ gives the n -dimensional manifold S_n . To introduce the metric on the manifold S_n ,




FIG. 1: Schematic of information geometry on the manifold S_2 . The manifold S_2 leads to the sphere surface of radius 2 (see also Supplementary Information (SI)). The statistical length \mathcal{L} is bounded by the shortest length $D = 2\theta = 2\cos^{-1}(\mathbf{r}_{\text{ini}} \cdot \mathbf{r}_{\text{fin}})$.

we use the Kullback-Leibler divergence [54] between two distributions \mathbf{p} and $\mathbf{p}' = (p'_0, p'_1, \dots, p'_n)$ defined as

$$D_{\text{KL}}(\mathbf{p}||\mathbf{p}') := \sum_{x=0}^n p_x \ln \frac{p_x}{p'_x}. \quad (10)$$

The Kullback-Leibler divergence is non-negative $D_{\text{KL}}(\mathbf{p}||\mathbf{p}') \geq 0$, and zero if and only if $\mathbf{p} = \mathbf{p}'$. The square of the line element ds is given by the divergence

$$ds^2 := 2D_{\text{KL}}(\mathbf{p}||\mathbf{p} + d\mathbf{p}), \quad (11)$$

where $d\mathbf{p} = (dp_0, dp_1, \dots, dp_n)$ is the infinitesimal displacement that satisfies $\sum_{x=0}^n dp_x = 0$. The second-order Taylor series of Eq. (11) gives the metric of the manifold

$$ds^2 := \sum_{x=0}^n \frac{(dp_x)^2}{p_x} = \sum_{x=0}^n p_x (d \ln p_x)^2. \quad (12)$$

Here, we define the metric g_{ij} as

$$ds^2 := \sum_{i=1}^{n'} \sum_{j=1}^{n'} g_{ij} d\lambda_i d\lambda_j, \quad (13)$$

where $(\lambda_1, \dots, \lambda_{n'})$ is the set of parameters such as control parameters. From Eqs. (12) and (13), the metric g_{ij} is given by the Fisher information $g_{ij} = \mathbb{E}[\partial_{\lambda_i} \ln p][\partial_{\lambda_j} \ln p]$ [54], where $\mathbb{E}[A] := \sum_{x=0}^n p_x A(x)$ denotes the expected value of any function $A(x)$. The square of the line element is also given by $ds^2 = \sum_{x=0}^n (2d\sqrt{p_x})^2$ under the constraint $\sum_{x=0}^n (\sqrt{p_x})^2 = 1$. Thus, the manifold S_n leads to the geometry of the n -sphere surface of radius 2 (see also Fig. 1).

We define the statistical length \mathcal{L} [55, 56] as

$$\mathcal{L} := \int ds = \int \frac{ds}{dt} dt. \quad (14)$$

Let $\mathbf{r} = (\sqrt{p_0}, \sqrt{p_1}, \dots, \sqrt{p_n})$ be the unit vector that satisfies $\mathbf{r} \cdot \mathbf{r} = \sum_{x=0}^n (\sqrt{p_x})^2 = 1$. The statistical length \mathcal{L} from the state \mathbf{r}_{ini} to the final state \mathbf{r}_{fin} is bounded by

$$\mathcal{L} \geq 2 \cos^{-1}(\mathbf{r}_{\text{ini}} \cdot \mathbf{r}_{\text{fin}}) := \mathcal{D}(\mathbf{r}_{\text{ini}}; \mathbf{r}_{\text{fin}}), \quad (15)$$

because $\mathcal{D}(\mathbf{r}_{\text{ini}}; \mathbf{r}_{\text{fin}}) = 2\theta$ gives the shortest length between \mathbf{r}_{ini} and \mathbf{r}_{fin} on the n -sphere surface S_n , where θ is the angle between \mathbf{r}_{ini} and \mathbf{r}_{fin} given by the inner product $\mathbf{r}_{\text{ini}} \cdot \mathbf{r}_{\text{fin}} = \cos \theta$.

Thermodynamics of information geometry.— We here discuss the main result of the letter. To treat the theory of information geometry as stochastic thermodynamics, we consider a stochastic thermodynamic expression of ds^2 as follows.

By using the definition of the thermodynamic force (5), the master equation is given by

$$\frac{d}{dt}p_x = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} p_x e^{-F_{x \rightarrow x'}^{(\nu)}}. \quad (16)$$

From Eqs. (12), (16) and $\sum_{x=0}^n d^2 p_x / dt^2 = 0$, we obtain an expression of the quantity ds^2/dt^2 in terms of thermodynamic quantities

$$\begin{aligned} \frac{ds^2}{dt^2} &= \sum_{x=0}^n p_x \frac{d}{dt} \left(-\frac{1}{p_x} \frac{dp_x}{dt} \right) \\ &= -\sum_{x=0}^n p_x \frac{d}{dt} \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} e^{-F_{x \rightarrow x'}^{(\nu)}} \right) \\ &= -\mathbb{E} \left[\frac{d}{dt} \overline{e^{-F}} \right], \end{aligned} \quad (17)$$

where $\overline{A}(x) := \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} A_{x \rightarrow x'}^{(\nu)}$ denotes the rate of any function of edge $A_{x \rightarrow x'}^{(\nu)}$ for the fixed initial state x . We mention that the total entropy change rate is given by $\dot{S}^{\text{tot}} = \mathbb{E}[\overline{F}]$ because of the anti-symmetry of the thermodynamic force $F_{x \rightarrow x'}^{(\nu)} = -F_{x' \rightarrow x}^{(\nu)}$. The equation (17) implies that geometric dynamics on the manifold S_n is given by the thermodynamic factor e^{-F} , that is well discussed in the context of stochastic thermodynamics (especially in the context of the fluctuation theorem [57–62]). We carefully mention that the expected value of $\overline{e^{-F}}$ gives $\mathbb{E}[\overline{e^{-F}}] = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n p_x W_{x \rightarrow x'}^{(\nu)} = 0$ compared to the integral fluctuation theorem $\langle e^{-F_{\text{traj}}} \rangle_{\text{traj}} = 1$ with the entropy production and the ensemble average of trajectories, F_{traj} and $\langle \dots \rangle_{\text{traj}}$, respectively [60, 62].

Due to the non-negativity $ds^2/dt^2 \geq 0$, we have a thermodynamic inequality

$$0 \geq \mathbb{E} \left[\frac{d}{dt} \overline{e^{-F}} \right]. \quad (18)$$

The equality holds if the system is in a stationary state, i.e., $ds/dt = 0$. To use Eqs. (2), (5) and (6) in the middle

line of Eq. (17), we obtain another expression of ds^2/dt^2 (see SI),

$$\frac{ds^2}{dt^2} = \left\langle \frac{d\dot{s}^{\text{bath}}}{dt} \right\rangle - \left\langle \frac{dF}{dt} \right\rangle = - \left\langle \frac{d\dot{s}^{\text{sys}}}{dt} \right\rangle. \quad (19)$$

Thus, we obtain another expression of Eq. (18)

$$\left\langle \frac{d\dot{s}^{\text{bath}}}{dt} \right\rangle \geq \left\langle \frac{dF}{dt} \right\rangle. \quad (20)$$

This result (20) implies another aspect of the second law of thermodynamics $\langle F \rangle \geq 0$. In a relaxation process where the entropy change of bath does not change in time (i.e., $d\dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)}/dt = 0$ for any x' and x), the thermodynamic force tends to decrease in time (i.e., $\langle dF/dt \rangle \leq 0$). This fact is consistent with the minimal entropy production principle in a stationary state.

In case of a near-equilibrium system, the detailed balance approximately holds, i.e., $F_{x' \rightarrow x}^{(\nu)} \simeq 0$ for any x' , x and ν . Then, the square of the line element is given by the entropy change in thermal baths $ds^2 \simeq \langle d\dot{s}^{\text{bath}} \rangle dt$. For example, if we assume the canonical distribution $p_x = \exp(\beta(\phi - H_x))$ where $\phi = -\beta^{-1} \ln[\sum_{x=0}^n \exp(-\beta H_x)]$ is the free energy that satisfies $\sum_{x=0}^n p_x = 1$, β is the inverse temperature and H_x is the Hamiltonian of the system in the state x , we obtain $\dot{s}_{x' \rightarrow x}^{\text{sys}} = \beta(H_x - H_{x'}) := \beta \dot{H}_{x' \rightarrow x}$ and $ds^2 = -\langle d\dot{s}^{\text{sys}} \rangle dt = -\langle d(\beta \dot{H}) \rangle dt$. We also mention that the linear irreversible thermodynamics [51] implies that the Fisher metric g_{ij} for thermodynamic forces $F_{x' \rightarrow x}^{(\nu)}$ is related to the Onsager coefficients (see SI). Due to the Cramér-Rao bound [53, 54], the Onsager coefficients relates the lower bound of the variance of unbiased estimator for thermodynamic forces.

The statistical length $\mathcal{L} = \int_0^\tau dt(ds/dt)$ from time $t = 0$ to $t = \tau$ is given by

$$\mathcal{L} = \int_{t=0}^{t=\tau} dt \left(-\mathbb{E} \left[\frac{d}{dt} \overline{e^{-F}} \right] \right)^{\frac{1}{2}}. \quad (21)$$

or equivalently $\mathcal{L} = \int_{t=0}^{t=\tau} dt \sqrt{[\langle d\dot{s}^{\text{bath}}/dt \rangle - \langle dF/dt \rangle]}$. From the shortest length of the manifold S_n , we obtain the following thermodynamic inequality from Eqs. (15) and (21),

$$\int_{t=0}^{t=\tau} dt \left(-\mathbb{E} \left[\frac{d}{dt} \overline{e^{-F}} \right] \right)^{\frac{1}{2}} \geq \mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau)), \quad (22)$$

or equivalently $\int_{t=0}^{t=\tau} dt \sqrt{[\langle d\dot{s}^{\text{bath}}/dt \rangle - \langle dF/dt \rangle]} \geq \mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))$. The equality holds if the path of transient dynamics is given by a geodesic line on the manifold S_n . This inequality shows a thermodynamic constraint on $\langle d\dot{s}^{\text{bath}}/dt \rangle$ and $\langle dF/dt \rangle$ for transient dynamics between initial and final probability distributions.

Thermodynamic uncertainty.— We finally reach to a thermodynamic uncertainty relationship between the




FIG. 2: Numerical calculation of thermodynamic quantities in the three states model of enzyme reaction. We numerically shows the non-negativity of $ds^2/dt^2 \geq 0$ and $ds^2/dt^2 = -\langle dF/dt \rangle + \langle d\dot{s}^{\text{bath}}/dt \rangle$ in the graph. We also show the total entropy change rate $\langle F \rangle \geq 0$. We note that $d\langle F \rangle/dt$ is not equal to $\langle dF/dt \rangle$.

speed and thermodynamic cost. We here consider the curve in Riemannian geometry from time $t = 0$ to $t = \tau$. The curve \mathcal{C} is defined as

$$\mathcal{C} = \frac{1}{2} \int_{t=0}^{t=\tau} dt \left(\frac{ds}{dt} \right)^2 = -\frac{1}{2} \int_{t=0}^{t=\tau} dt \mathbb{E} \left[\frac{d}{dt} e^{-F} \right], \quad (23)$$

or equivalently $\mathcal{C} = \int_{t=0}^{t=\tau} dt [\langle d\dot{s}^{\text{bath}}/dt \rangle - \langle dF/dt \rangle]/2$. Especially in the case that system is close to be in thermodynamic equilibrium, the curve \mathcal{C} is given by $\mathcal{C} \simeq \int \langle d\dot{s}^{\text{bath}} \rangle / 2$. Therefore, the curve \mathcal{C} can be interpreted as a thermodynamic cost.

Due to the Cauchy-Schwarz inequality $\int_0^\tau dt \int_0^\tau (ds/dt)^2 dt \geq (\int_0^\tau (ds/dt) dt)^2$ [22], we obtain a thermodynamic uncertainty relationship

$$\tau \geq \frac{\mathcal{L}^2}{2\mathcal{C}}. \quad (24)$$

The equality holds if the speed of dynamics $ds^2/dt^2 = -\mathbb{E} [de^{-F}/dt]$ does not depend on time. This result implies the uncertainty relationship between a thermodynamic cost \mathcal{C} and the speed τ . If the transient time τ is small, a thermodynamic cost \mathcal{C} should be large, i.e., $\mathcal{C} \geq [\mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))]^2 / (2\tau)$ or equivalently

$$\tau \geq \frac{[\mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))]^2}{2\mathcal{C}}. \quad (25)$$

From the inequality (25), we can discuss the efficiency $0 \leq \eta \leq 1$ defined as

$$\eta := \frac{[\mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))]^2}{2\tau\mathcal{C}}. \quad (26)$$






FIG. 3: Numerical calculation of thermodynamic uncertainty relationship in the three states model of enzyme reaction. We numerically shows the geometric inequality $\mathcal{L} \geq \mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))$ and thermodynamic uncertainty relationship $\tau \geq \mathcal{L}^2 / (2\mathcal{C}) \geq [\mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))]^2 / (2\mathcal{C})$ in the graph.

In case of a near equilibrium system, the efficiency is given by $\eta := [\mathcal{D}(\mathbf{r}(0); \mathbf{r}(\tau))]^2 / (\tau \int \langle d\dot{s}^{\text{bath}} \rangle)$, which implies the trade off between the entropy change in thermal baths and the geodesic distance from the initial state to the final state.

Three states model of enzyme reaction.— We numerically test thermodynamic inequalities of information geometry by using a thermodynamic model of biochemical reaction. We here consider a three states model (see also SI) that represents a chemical reaction $A + B \rightleftharpoons AB$ with enzyme X ,



We here consider the probability distribution of the states $x = A, AX, AB$. We assume that the system is attached to a single heat bath ($n_{\text{bath}} = 1$) with inverse temperature β . The master equation is given by Eq. (1), where the transition rates are supposed to be

$$\begin{aligned} W_{A \rightarrow AX}^{(1)} &= k_{AX+}[X], & W_{AX \rightarrow A}^{(1)} &= k_{AX+}e^{-\beta\Delta\mu_{AX}}, \\ W_{A \rightarrow AB}^{(1)} &= k_{AB+}[B], & W_{AB \rightarrow A}^{(1)} &= k_{AB+}e^{-\beta\Delta\mu_{AB}}, \\ W_{AX \rightarrow AB}^{(1)} &= k_+[B], & W_{AB \rightarrow AX}^{(1)} &= k_+e^{-\beta\Delta\mu[X]}, \end{aligned} \quad (30)$$

where $[X]$ ($[B]$) is the concentration of X (B), k_{AX+} , k_{AB+} , and k_+ are reaction rate constants, and $\Delta\mu_{AX}$, $\Delta\mu_{AB}$, and $\Delta\mu$ are the chemical potential differences.

In a numerical simulation, we set $k_{AX+} = k_{AB+} = k_+ = 1$, $\beta\Delta\mu_{AX} = 1.0$, $\beta\Delta\mu_{AB} = 0.5$, and $\beta\Delta\mu = 2.0$.

We assume that the time evolution of the concentrations are given by $[X] = \tan^{-1}(\omega_X t)$, $[B] = \tan^{-1}(\omega_B t)$ with $\omega_X = 1$ and $\omega_B = 2$, which means the concentrations $[X]$ and $[B]$ performs as control parameters. At time $t = 0$, we set the initial probability distribution $(p_A, p_{AX}, p_{AB}) = (0.9998, 0.001, 0.001)$.

In Fig. 2, we numerically show the inequality $\langle d\dot{s}^{\text{bath}}/dt \rangle \geq \langle dF/dt \rangle$. We also check that this inequality does not coincide with the second law of thermodynamics $\langle F \rangle \geq 0$. We also check the thermodynamic uncertainty relationship $\mathcal{L}^2/(2\mathcal{C}) \leq \tau$ in Fig. 3. Because the path from initial distribution $(p_A, p_{AX}, p_{AB}) = (0.9998, 0.001, 0.001)$ to the final distribution is close to the geodesic line, the thermodynamic uncertainty relationship gives a tight bound of the transition time τ .

Conclusion.— In this letter, we reveal the link between stochastic thermodynamic quantities (J , F , \dot{s}^{sys} , \dot{s}^{bath}) and information geometric quantities (ds^2 , g_{ij} , \mathcal{L} , \mathcal{C}). Because the theory of information geometry is applicable to various fields of science such as neuroscience, signal processing, machine learning and quantum mechanics, this link would help to understand thermodynamic aspect of such a topic. The trade-off relationship between speed and thermodynamic cost Eq. (25) would be helpful to understand biochemical reactions and gives a new insight into recent studies of the relationship between information and thermodynamics in biochemical processes [7, 41, 63–67].

ACKNOWLEDGEMENT

I am grateful to Shumpei Yamamoto for discussions of stochastic thermodynamics for the master equation, to Naoto Shiraishi, Keiji Saito, Hal Tasaki, and Shin-Ichi Sasa for discussions of thermodynamic uncertainty relationships, to Schuyler B. Nicholson for discussion of information geometry and thermodynamics, and to Pieter rein ten Wolde for discussions of thermodynamics in a chemical reaction. We also thank Tamiki Komatsuzaki to acknowledge my major contribution of this work and allow me to submit this manuscript alone. I mentioned that, after my submission of the first version of this manuscript on arXiv [68], I heard that Schuyler B. Nicholson independently discovered a similar result such as Eq. (19) [69]. I thank Sebastian Goldt, Matteo Poletini, and Taro Toyozumi for valuable comments on the manuscript. This research is supported by JSPS KAKENHI Grant No. JP16K17780.

[1] Parrondo, J. M., Horowitz, J. M., & Sagawa, T. *Thermodynamics of information*. Nature physics, **11**(2), 131-139 (2015).

[2] Leff, H. S., & Rex, A. F. (Eds.). *Maxwell's demon: entropy, information, computing* (Princeton University Press, 2014).

[3] Kawai, R., Parrondo, J. M. R., & Van den Broeck, C. *Dissipation: The phase-space perspective*. Physical review letters, **98**(8), 080602 (2007).

[4] Sagawa, T., & Ueda, M. *Generalized Jarzynski equality under nonequilibrium feedback control*. Physical review letters, **104**(9), 090602 (2010).

[5] Still, S., Sivak, D. A., Bell, A. J., & Crooks, G. E. *Thermodynamics of prediction*. Physical review letters, **109**(12), 120604 (2012).

[6] Sagawa, T., & Ueda, M. *Fluctuation theorem with information exchange: role of correlations in stochastic thermodynamics*. Physical review letters, **109**(18), 180602 (2012).

[7] Ito, S., & Sagawa, T. *Information thermodynamics on causal networks*. Physical review letters, **111**(18), 180603 (2013).

[8] Hartich, D., Barato, A. C., & Seifert, U. *Stochastic thermodynamics of bipartite systems: transfer entropy inequalities and a Maxwells demon interpretation*. Journal of Statistical Mechanics: Theory and Experiment (2014). P02016.

[9] Hartich, D., Barato, A. C., & Seifert, U. *Sensory capacity: An information theoretical measure of the performance of a sensor*. Physical Review E, **93**(2), 022116 (2016).

[10] Spinney, R. E., Lizier, J. T., & Prokopenko, M. *Transfer entropy in physical systems and the arrow of time*. Physical Review E, **94**(2), 022135 (2016).

[11] Ito, S. *Backward transfer entropy: Informational measure for detecting hidden Markov models and its interpretations in thermodynamics, gambling and causality*. Scientific reports, **6** (2016).

[12] Crooks, G. E., & Still, S. E. *Marginal and conditional second laws of thermodynamics*. arXiv preprint arXiv:1611.04628 (2016).

[13] Allahverdyan, A. E., Janzing, D., & Mahler, G. *Thermodynamic efficiency of information and heat flow*. Journal of Statistical Mechanics: Theory and Experiment, (2009) P09011.

[14] Horowitz, J. M., & Esposito, M. *Thermodynamics with continuous information flow*. Physical Review X, **4**(3), 031015 (2014).

[15] Horowitz, J. M., & Sandberg, H. *Second-law-like inequalities with information and their interpretations*. New Journal of Physics, **16**(12), 125007 (2014).

[16] Shiraishi, N., & Sagawa, T. *Fluctuation theorem for partially masked nonequilibrium dynamics*. Physical Review E, **91**(1), 012130 (2015).

[17] Shiraishi, N., Ito, S., Kawaguchi, K., & Sagawa, T. *Role of measurement-feedback separation in autonomous Maxwell's demons*. New Journal of Physics, **17**(4), 045012 (2015).

[18] Yamamoto, S., Ito, S., Shiraishi, N., & Sagawa, T. *Linear irreversible thermodynamics and Onsager reciprocity for information-driven engines*. Physical Review E, **94**(5), 052121 (2016).

[19] Goldt, S., & Seifert, U. *Stochastic thermodynamics of learning*. Physical review letters, **118**(1), 010601 (2017).

[20] Sekimoto, K. *Stochastic energetics*. (Springer, 2010).

[21] Seifert, U. *Stochastic thermodynamics, fluctuation theorems and molecular machines*. Reports on Progress in

- Physics, **75**(12), 126001 (2012).
- [22] Crooks, G. E. *Measuring thermodynamic length*. Physical Review Letters, **99**(10), 100602 (2007).
 - [23] Feng, E. H., & Crooks, G. E. *Length of times arrow*. Physical review letters, **101**(9), 090602 (2008).
 - [24] Poletti, M., & Esposito, M. *Nonconvexity of the relative entropy for Markov dynamics: A Fisher information approach*. Physical Review E, **88**(1), 012112 (2013).
 - [25] Shimazaki, H. *Neurons as an Information-theoretic Engine*. arXiv preprint arXiv:1512.07855 (2015).
 - [26] Nicholson, S. B., & Kim, E. J. *Investigation of the statistical distance to reach stationary distributions*. Physics Letters A, **379**(3), 83-88 (2015).
 - [27] Tajima, H., & Hayashi, M. *Finite-size effect on optimal efficiency of heat engines*. Physical Review E, **96**(1), 012128 (2017). x
 - [28] Lahiri, S., Sohl-Dickstein, J., & Ganguli, S. *A universal tradeoff between power, precision and speed in physical communication*. arXiv preprint arXiv:1603.07758 (2016).
 - [29] Weinhold, F. *Metric geometry of equilibrium thermodynamics*. The Journal of Chemical Physics, **63**(6), 2479-2483 (1975).
 - [30] Ruppeiner, G. *Thermodynamics: A Riemannian geometric model*. Physical Review A, **20**(4), 1608 (1979).
 - [31] Salamon, P., & Berry, R. S. *Thermodynamic length and dissipated availability*. Physical Review Letters, **51**(13), 1127 (1983).
 - [32] Sivak, D. A., & Crooks, G. E. *Thermodynamic metrics and optimal paths*. Physical review letters, **108**(19), 190602 (2012).
 - [33] Machta, B. B. *Dissipation bound for thermodynamic control*. Physical review letters, **115**(26), 260603 (2015).
 - [34] Rotskoff, G. M., Crooks, G. E., & Vanden-Eijnden, E. *Geometric approach to optimal nonequilibrium control: Minimizing dissipation in nanomagnetic spin systems*. Physical Review E, **95**(1), 012148 (2017).
 - [35] Amari, S. I., & Nagaoka, H. *Methods of information geometry*. (American Mathematical Soc., 2007).
 - [36] Oizumi, M., Tsuchiya, N., & Amari, S. I. *Unified framework for information integration based on information geometry*. Proceedings of the National Academy of Sciences, **113**(51), 14817-14822 (2016).
 - [37] Pires, D. P., Cianciaruso, M., Céleri, L. C., Adesso, G., & Soares-Pinto, D. O. *Generalized geometric quantum speed limits*. Physical Review X, **6**(2), 021031 (2016).
 - [38] Amari, S. I. *Information geometry and its applications*. (Springer Japan, 2016).
 - [39] Lan, G., Sartori, P., Neumann, S., Sourjik, V., & Tu, Y. *The energy-speed-accuracy trade-off in sensory adaptation*. Nature physics, **8**(5), 422-428 (2012).
 - [40] Govern, C. C., & ten Wolde, P. R. *Optimal resource allocation in cellular sensing systems*. Proceedings of the National Academy of Sciences, **111**(49), 17486-17491 (2014).
 - [41] Ito, S., & Sagawa, T. *Maxwell's demon in biochemical signal transduction with feedback loop*. Nature communications, **6** (2015).
 - [42] Barato, A. C., & Seifert, U. *Thermodynamic uncertainty relation for biomolecular processes*. Physical review letters, **114**(15), 158101 (2015).
 - [43] Gingrich, T. R., Horowitz, J. M., Perunov, N., & England, J. L. *Dissipation bounds all steady-state current fluctuations*. Physical review letters, **116**(12), 120601 (2016).
 - [44] Shiraishi, N., Saito, K., & Tasaki, H. *Universal trade-off relation between power and efficiency for heat engines*. Physical review letters, **117**(19), 190601 (2016).
 - [45] Pietzonka, P., Barato, A. C., & Seifert, U. *Universal bounds on current fluctuations*. Physical Review E, **93**(5), 052145 (2016).
 - [46] Barato, A. C., & Seifert, U. *Cost and precision of Brownian clocks*. Physical Review X, **6**(4), 041053 (2016).
 - [47] Horowitz, J. M., & Gingrich, T. R. *Proof of the finite-time thermodynamic uncertainty relation for steady-state currents*. Physical Review E, **96**(2), 020103 (2017).
 - [48] Proesmans, K., & Van den Broeck, C. *Discrete-time thermodynamic uncertainty relation*. EPL (Europhysics Letters), **119**(2), 20001 (2017).
 - [49] Maes, C. *Frenetic bounds on the entropy production*. Physical review letters. **119**(16), 160601 (2017).
 - [50] Dechant, A., & Sasa, S. I. *Current fluctuations and transport efficiency for general Langevin systems*. arXiv preprint arXiv:1708.08653 (2017).
 - [51] Schnakenberg, J. *Network theory of microscopic and macroscopic behavior of master equation systems*. Reviews of Modern physics, **48**(4), 571 (1976).
 - [52] Andrieux, D., & Gaspard, P. *Fluctuation theorem for currents and Schnakenberg network theory*. Journal of statistical physics, **127**(1), 107-131 (2007).
 - [53] Rao, C. R. *Information and the accuracy attainable in the estimation of statistical parameters*. In *Breakthroughs in statistics* (pp. 235-247). (Springer New York, 1992).
 - [54] Cover, T. M., & Thomas, J. A. *Elements of information theory*. (John Wiley & Sons, 2012).
 - [55] Wootters, W. K. *Statistical distance and Hilbert space*. Physical Review D, **23**(2), 357 (1981).
 - [56] Braunstein, S. L., & Caves, C. M. *Statistical distance and the geometry of quantum states*. Physical Review Letters, **72**(22), 3439 (1994).
 - [57] Jarzynski, C. *Nonequilibrium equality for free energy differences*. Physical Review Letters, **78**(14), 2690 (1997).
 - [58] Crooks, G. E. *Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences*. Physical Review E, **60**(3), 2721 (1999).
 - [59] Evans, D. J., & Searles, D. J. *The fluctuation theorem*. Advances in Physics, **51**(7), 1529-1585 (2002).
 - [60] Seifert, U. *Entropy production along a stochastic trajectory and an integral fluctuation theorem*. Physical review letters, **95**(4), 040602 (2005).
 - [61] Esposito, M., & Van den Broeck, C. *Three faces of the second law. I. Master equation formulation*. Physical Review E, **82**(1), 011143 (2010).
 - [62] Esposito, M., & Van den Broeck, C. *Three detailed fluctuation theorems*. Physical review letters, **104**(9), 090601 (2010).
 - [63] Barato, A. C., Hartich, D., & Seifert, U. (2014). *Efficiency of cellular information processing*. New Journal of Physics, **16**(10), 103024.
 - [64] Sartori, P., Granger, L., Lee, C. F., & Horowitz, J. M. *Thermodynamic costs of information processing in sensory adaptation*. PLoS computational biology, **10**(12), e1003974 (2014).
 - [65] Bo, S., Del Giudice, M., & Celani, A. *Thermodynamic limits to information harvesting by sensory systems*. Journal of Statistical Mechanics: Theory and Experiment, (2015). P01014.
 - [66] Ouldrige, T. E., Govern, C. C., & ten Wolde, P. R. *Thermodynamics of computational copying in biochemi-*

cal systems. Physical Review X, **7**(2), 021004 (2017).
 [67] McGrath, T., Jones, N. S., ten Wolde, P. R., & Ouldridge, T. E. *Biochemical machines for the interconversion of mutual information and work*. Physical Review Letters, **118**(2), 028101 (2017).

[68] Ito, S. "Stochastic Thermodynamic Interpretation of Information Geometry". arXiv preprint arXiv:1712.04311v1 (2017).
 [69] Nicholson, S. B. *et. al.*, in preparation.

SUPPLEMENTARY INFORMATION

I. Intuitive proof of the fact that the manifold S_2 gives sphere surface of radius 2

We here intuitively show the fact that the manifold S_2 gives sphere surface of radius 2. The probability vector $\mathbf{p} = (p_0, p_1, p_2)$ holds the normalization $\sum_{x=0}^2 p_x = 1$. The square of the line element ds is given by

$$ds^2 = \sum_{x=0}^2 \frac{(dp_x)^2}{p_x}. \quad (31)$$

We here introduce the polar coordinate system (ϕ, ψ) where $p_0 = (\cos \psi)^2$, $p_1 = (\sin \psi)^2 (\cos \phi)^2$, $p_2 = (\sin \psi)^2 (\sin \phi)^2$. We can check that the normalization $\sum_{x=0}^2 p_x = 1$ holds. We obtain $dp_0 = -2(\cos \psi)(\sin \psi)d\psi$, $dp_1 = 2(\cos \psi)(\sin \psi)(\cos \phi)^2 d\psi - 2(\cos \phi)(\sin \phi)(\sin \psi)^2 d\phi$, and $dp_2 = 2(\cos \psi)(\sin \psi)(\sin \phi)^2 d\psi + 2(\cos \phi)(\sin \phi)(\sin \psi)^2 d\phi$. From Eq. (31), we then obtain

$$\begin{aligned} ds^2 &= 4[(\sin \psi)^2 + (\cos \psi)^2 (\cos \phi)^2 + (\cos \psi)^2 (\sin \phi)^2](d\psi)^2 + 0 \times (d\phi)(d\psi) + 4[(\sin \phi)^2 (\sin \psi)^2 + (\cos \phi)^2 (\sin \psi)^2](d\phi)^2 \\ &= 2^2[(d\psi)^2 + (\sin \psi)^2 (d\phi)^2]. \end{aligned} \quad (32)$$

Because the metric of a sphere surface of radius R is given by $ds^2 = R^2[(d\psi)^2 + (\sin \psi)^2 (d\phi)^2]$, the manifold S_2 gives a sphere surface of radius $R = 2$.

II. Detailed derivation of Eq. (19) in the main text

From Eq. (17) in the main text, we obtain

$$\begin{aligned} \frac{ds^2}{dt^2} &= -\mathbb{E} \left[\frac{d}{dt} e^{-F} \right] \\ &= -\sum_{x=0}^n p_x \frac{d}{dt} \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} e^{-F_{x \rightarrow x'}^{(\nu)}} \right) \\ &= -\sum_{x=0}^n p_x \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} \left(-\frac{d}{dt} F_{x \rightarrow x'}^{(\nu)} \right) e^{-F_{x \rightarrow x'}^{(\nu)}} \right) - \sum_{x=0}^n p_x \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n \left(\frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) e^{-F_{x \rightarrow x'}^{(\nu)}} \right). \end{aligned} \quad (33)$$

By using the definition Eqs. (5) and (6), the first term is calculated as follows

$$\begin{aligned} & -\sum_{x=0}^n p_x \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n W_{x \rightarrow x'}^{(\nu)} \left(-\frac{d}{dt} F_{x \rightarrow x'}^{(\nu)} \right) e^{-F_{x \rightarrow x'}^{(\nu)}} \right) \\ &= -\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n p_x W_{x \rightarrow x'}^{(\nu)} \left(\frac{d}{dt} F_{x' \rightarrow x}^{(\nu)} \right) \\ &= -\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x > x'} p_x W_{x \rightarrow x'}^{(\nu)} \left(\frac{d}{dt} F_{x' \rightarrow x}^{(\nu)} \right) - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x' > x} p_{x'} W_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} F_{x' \rightarrow x}^{(\nu)} \right) \\ &= -\sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} F_{x' \rightarrow x}^{(\nu)} \right) = -\left\langle \frac{dF}{dt} \right\rangle, \end{aligned} \quad (34)$$

where we used $F_{x' \rightarrow x}^{(\nu)} = -F_{x \rightarrow x'}^{(\nu)}$ and $F_{x' \rightarrow x'}^{(\nu)} = 0$. The second term is also calculated as follows

$$\begin{aligned}
& - \sum_{x=0}^n p_x \left(\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n \left(\frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) e^{-F_{x \rightarrow x'}^{(\nu)}} \right) \\
&= - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n p_{x'} W_{x' \rightarrow x}^{(\nu)} \frac{1}{W_{x \rightarrow x'}^{(\nu)}} \left(\frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) \\
&= - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x' \neq x} p_{x'} W_{x' \rightarrow x}^{(\nu)} \frac{1}{W_{x \rightarrow x'}^{(\nu)}} \left(\frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x=0}^n p_x W_{x \rightarrow x}^{(\nu)} \frac{1}{W_{x \rightarrow x}^{(\nu)}} \left(\frac{d}{dt} W_{x \rightarrow x}^{(\nu)} \right) \\
&= - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x > x'} p_{x'} W_{x' \rightarrow x}^{(\nu)} \frac{1}{W_{x \rightarrow x'}^{(\nu)}} \left(\frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) + \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x=0}^n p_x \left(\sum_{x' \neq x} \frac{d}{dt} W_{x \rightarrow x'}^{(\nu)} \right) \\
&= - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x \neq x'} p_{x'} W_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} \ln(W_{x \rightarrow x'}^{(\nu)}) \right) + \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x' \neq x} p_{x'} W_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} \ln(W_{x' \rightarrow x}^{(\nu)}) \right) \\
&= \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x' \neq x} p_{x'} W_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} \dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} \right) \\
&= \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x > x'} p_{x'} W_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} \dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} \right) - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x, x' | x > x'} p_x W_{x \rightarrow x'}^{(\nu)} \left(\frac{d}{dt} \dot{s}_{x \rightarrow x'}^{\text{bath}(\nu)} \right) \\
&= \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} \left(\frac{d}{dt} \dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} \right) = \left\langle \frac{d \dot{s}^{\text{bath}}}{dt} \right\rangle, \tag{35}
\end{aligned}$$

where we used Eq. (2) in the main text, $\dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} = -\dot{s}_{x \rightarrow x'}^{\text{bath}(\nu)}$ and $\dot{s}_{x' \rightarrow x'}^{\text{bath}(\nu)} = 0$.

By using $F_{x' \rightarrow x}^{(\nu)} = \dot{s}_{x' \rightarrow x}^{\text{bath}(\nu)} + \dot{s}_{x' \rightarrow x}^{\text{sys}}$, we obtain Eq. (19) in the main text

$$\frac{ds^2}{dt^2} = \left\langle \frac{d \dot{s}^{\text{bath}}}{dt} \right\rangle - \left\langle \frac{dF}{dt} \right\rangle = - \left\langle \frac{d \dot{s}^{\text{sys}}}{dt} \right\rangle. \tag{36}$$

We also obtain the definition of the Fisher metric from the result (36)

$$\begin{aligned}
\frac{ds^2}{dt^2} &= \left\langle \frac{d \dot{s}^{\text{bath}}}{dt} \right\rangle - \left\langle \frac{dF}{dt} \right\rangle \\
&= - \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_{x'}} \frac{dp_{x'}}{dt} - \frac{1}{p_x} \frac{dp_x}{dt} \right] \\
&= - \sum_{\nu=1}^{n_{\text{bath}}} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n \sum_{x''=0}^n p_{x'} W_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_{x'}} W_{x'' \rightarrow x'}^{(\nu')} p_{x''} - \frac{1}{p_x} W_{x'' \rightarrow x}^{(\nu')} p_{x''} \right] \\
&= \sum_{x=0}^n p_x \left[\sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n \frac{p_{x'} W_{x' \rightarrow x}^{(\nu)}}{p_x} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x''=0}^n \frac{p_{x''} W_{x'' \rightarrow x}^{(\nu')}}{p_x} \right] \\
&= \mathbb{E} \left[\left(\frac{d \ln p}{dt} \right)^2 \right] \\
&= \mathbb{E} \left[\left(\sum_{i=1}^{n'} \frac{\partial \ln p}{\partial \lambda_i} \frac{d \lambda_i}{dt} \right)^2 \right] \\
&= \sum_{i=1}^{n'} \sum_{j=1}^{n'} \frac{d \lambda_i}{dt} g_{ij} \frac{d \lambda_j}{dt}, \tag{37}
\end{aligned}$$

where we used $\sum_{x=0}^n W_{x' \rightarrow x}^{(\nu)} = 0$ and the master equation $dp_x/dt = \sum_{\nu=1}^{n_{\text{bath}}} \sum_{x'=0}^n p_{x'} W_{x' \rightarrow x}^{(\nu)}$.

III. Linear irreversible thermodynamic interpretation of information geometry

We here discuss stochastic thermodynamic interpretation of information geometry for a near-equilibrium system. If we assume $F_{x' \rightarrow x}^{(\nu)} = 0$, we have $J_{x' \rightarrow x}^{(\nu)} = 0$. Thus, we have a linear expansion of thermodynamic force $F_{x' \rightarrow x}^{(\nu)}$ in terms of the thermodynamic flow $J_{x' \rightarrow x}^{(\nu)}$ for a near-equilibrium condition (i.e., $F_{x' \rightarrow x}^{(\nu)} \simeq 0$ for any x and x')

$$\begin{aligned} F_{x' \rightarrow x}^{(\nu)} &= \ln \left(1 + \frac{J_{x' \rightarrow x}^{(\nu)}}{W_{x' \rightarrow x}^{(\nu)} p_x} \right) \\ &= \alpha_{x' \rightarrow x}^{(\nu)} J_{x' \rightarrow x}^{(\nu)} + o(J_{x' \rightarrow x}^{(\nu)}), \end{aligned} \quad (38)$$

$$\alpha_{x' \rightarrow x}^{(\nu)} := \left. \frac{1}{W_{x' \rightarrow x}^{(\nu)} p_x} \right|_{F_{x' \rightarrow x}^{(\nu)} = 0}. \quad (39)$$

We call this coefficient $\alpha_{x' \rightarrow x}^{(\nu)}$ as the Onsager coefficient of the edge $(x' \rightarrow x, \nu)$. The symmetry of the coefficient $\alpha_{x' \rightarrow x}^{(\nu)} = \alpha_{x \rightarrow x'}^{(\nu)}$ holds due to the condition $F_{x' \rightarrow x}^{(\nu)} = 0$.

If we consider the Kirchhoff's current law in a stationary state, the linear combination of the coefficient $\alpha_{x \rightarrow x'}^{(\nu)}$ leads to a Onsager coefficient [51]. Let $\{C_1, \dots, C_m\}$ be the cycle basis of the Markov network for the master equation. The thermodynamic force of the cycle $F(C_i)$ is defined as

$$F(C_i) = \sum_{(x' \rightarrow x, \nu) \in E} S(\{x' \rightarrow x, \nu\}, C_i) F_{x' \rightarrow x}^{(\nu)} \quad (40)$$

where

$$S(\{x' \rightarrow x, \nu\}, C_i) = \begin{cases} 1 & (\{x' \rightarrow x, \nu\} \in C_i) \\ -1 & (\{x \rightarrow x', \nu\} \in C_i) \\ 0 & (\text{otherwise}) \end{cases}. \quad (41)$$

The thermodynamic flow of the cycle $J(C_i)$ is defined as

$$J_{x' \rightarrow x}^{(\nu)} = \sum_{i=1}^m S(\{x' \rightarrow x, \nu\}, C_i) J(C_i). \quad (42)$$

We then obtain the linear relationship $J(C_j) = \sum_{i=1}^m L_{ji} F(C_i)$ with the Onsager coefficient

$$L_{ij} = \sum_{(x' \rightarrow x, \nu) \in E} \alpha_{x' \rightarrow x}^{(\nu)} S(\{x' \rightarrow x, \nu\}, C_i) S(\{x' \rightarrow x, \nu\}, C_j), \quad (43)$$

for a near-equilibrium condition, the second law of thermodynamics

$$\begin{aligned} 0 &\leq \dot{S}^{\text{tot}} \\ &= \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} F_{x' \rightarrow x}^{(\nu)} \\ &= \sum_{(x' \rightarrow x, \nu) \in E} \sum_{i=1}^m S(\{x' \rightarrow x, \nu\}, C_i) J(C_i) F_{x' \rightarrow x}^{(\nu)} \\ &= \sum_{i=1}^m J(C_i) F(C_i), \\ &= \sum_{j=1}^m \sum_{i=1}^m L_{ij} F(C_i) F(C_j), \end{aligned} \quad (44)$$

and the Onsager reciprocal relationship $L_{ij} = L_{ji}$. For $m = 2$, the second law of thermodynamics $L_{11} F(C_1)^2 + L_{22} F(C_2)^2 + 2L_{12} F(C_1) F(C_2) \geq 0$ is then given by $L_{11} \geq 0$, $L_{22} \geq 0$, and $L_{11} L_{22} - L_{12}^2 \geq 0$.

For a near-equilibrium system, the square of line element is also given by

$$\begin{aligned}
ds^2 &= - \left\langle \frac{d\dot{s}^{\text{sys}}}{dt} \right\rangle dt^2 \\
&= - \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_{x'}} \frac{dp_{x'}}{dt} - \frac{1}{p_x} \frac{dp_x}{dt} \right] dt^2 \\
&= - \sum_{(x' \rightarrow x, \nu) \in E} J_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_{x'}} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x''=0}^n J_{x'' \rightarrow x'}^{(\nu')} - \frac{1}{p_x} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x''=0}^n J_{x'' \rightarrow x}^{(\nu')} \right] dt^2 \\
&= \sum_{(x' \rightarrow x, \nu) \in E} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x''=0}^n J_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_x} J_{x'' \rightarrow x}^{(\nu')} - \frac{1}{p_{x'}} J_{x'' \rightarrow x'}^{(\nu')} \right] dt^2 \\
&= \sum_{(x' \rightarrow x, \nu) \in E} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x''=0}^n J_{x' \rightarrow x}^{(\nu)} \left[\frac{1}{p_x} J_{x'' \rightarrow x}^{(\nu')} - \frac{1}{p_{x'}} J_{x'' \rightarrow x'}^{(\nu')} \right] dt^2 \\
&= \sum_{\nu=1}^{n_{\text{bath}}} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n \sum_{x''=0}^n \left[\frac{J_{x' \rightarrow x}^{(\nu)} J_{x'' \rightarrow x}^{(\nu')}}{p_x} \right] dt^2 \\
&= \sum_{\nu=1}^{n_{\text{bath}}} \sum_{\nu'=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n \sum_{x''=0}^n \left[\frac{F_{x' \rightarrow x}^{(\nu)} F_{x'' \rightarrow x}^{(\nu')}}{\alpha_{x' \rightarrow x}^{(\nu)} p_x \alpha_{x'' \rightarrow x}^{(\nu')}} \right] dt^2. \tag{45}
\end{aligned}$$

We here consider the situation that thermodynamic forces are assumed to be control parameters $F_{x' \rightarrow x}^{(\nu_x)} dt := d\lambda_{(x', x, \nu_x)}$. Then the square of line element can be written by the following Fisher metric

$$ds^2 = \sum_{\nu_x=1}^{n_{\text{bath}}} \sum_{x=0}^n \sum_{x'=0}^n \sum_{\nu_y=1}^{n_{\text{bath}}} \sum_{y=0}^n \sum_{y'=0}^n g_{(x', x, \nu_x)(y', y, \nu_y)} d\lambda_{(x', x, \nu_x)} d\lambda_{(y', y, \nu_y)} \tag{46}$$

$$g_{(x', x, \nu_x)(y', y, \nu_y)} = \frac{\delta_{xy}}{\alpha_{x' \rightarrow x}^{(\nu_x)} p_x \alpha_{y' \rightarrow y}^{(\nu_y)}}. \tag{47}$$

This result implies that the Fisher metric is given by the Onsager coefficients of the edge for a near equilibrium condition. Because the Cramér-Rao bound [53, 54] implies that the variance of unbiased estimator for thermodynamic forces is bounded by the inverse of this Fisher metric, the Onsager coefficients of the edge gives the lower bound of the variance of unbiased estimator for thermodynamic forces in a near equilibrium condition.

IV. Detail of the three states model of enzyme reaction

The master equation of the three states model of enzyme reaction is given by

$$\begin{aligned}
\frac{dp_A}{dt} &= -(k_{AX+} + k_{AB+})[X]p_A + k_{AB-}p_{AB} + k_{AX-}p_{AX}, \\
\frac{dp_{AB}}{dt} &= k_{AB+}p_A - (k_{AB-} + k_-[X])p_{AB} + k_+[B]p_{AX}, \\
\frac{dp_{AX}}{dt} &= k_{AX+}[X]p_A + k_-p_{AB} - (k_{AX-} + k_+[B])p_{AX},
\end{aligned} \tag{48}$$

where k_- , k_{AB-} , k_{AX-} are given by the local detailed balance

$$\begin{aligned}
\ln \frac{k_{AX+}}{k_{AX-}} &= \beta \Delta \mu_{AX}, \\
\ln \frac{k_{AB+}}{k_{AB-}} &= \beta \Delta \mu_{AB}, \\
\ln \frac{k_+}{k_-} &= \beta \Delta \mu.
\end{aligned} \tag{49}$$

We here assume that the sum of the concentrations $[A] + [AB] + [AX] = n_A$ is a constant. The probabilities distributions p_A , p_{AB} , and p_{AX} are then corresponding to the fractions of $[A]/n_A$, $[AB]/n_A$ and $[AX]/n_A$. From the master equation (51), we obtain the rate equations of enzyme reaction

$$\begin{aligned}\frac{d[A]}{dt} &= -(k_{AX+} + k_{AB+})[X][A] + k_{AB-}[AB] + k_{AX-}[AX], \\ \frac{d[AB]}{dt} &= k_{AB+}[A] - (k_{AB-} + k_-[X])[AB] + k_+[B][AX], \\ \frac{d[AX]}{dt} &= k_{AX+}[X][A] + k_-[AB] - (k_{AX-} + k_+[B])[AX].\end{aligned}\tag{50}$$

In this model, the stochastic entropy changes of heat bath are given by

$$\begin{aligned}\dot{s}_{A \rightarrow AB}^{\text{bath}(1)} &= \beta \Delta \mu_{AB} + \ln[B], \\ \dot{s}_{AB \rightarrow AX}^{\text{bath}(1)} &= -\beta \Delta \mu - \ln[B] + \ln[X], \\ \dot{s}_{AX \rightarrow AB}^{\text{bath}(1)} &= -\beta \Delta \mu_{AX} - \ln[X].\end{aligned}\tag{51}$$

If the entropy change of heat bath in a cycle $A \rightarrow AB \rightarrow AX \rightarrow A$ has non-zero value, i.e., $\dot{s}_{A \rightarrow AB}^{\text{bath}(1)} + \dot{s}_{AB \rightarrow AX}^{\text{bath}(1)} + \dot{s}_{AX \rightarrow AB}^{\text{bath}(1)} = \beta(\Delta \mu_{AB} - \Delta \mu - \Delta \mu_{AX}) \neq 0$, the system is driven by thermodynamic forces in a cycle $F_{A \rightarrow AB}^{(1)} + F_{AB \rightarrow AX}^{(1)} + F_{AX \rightarrow A}^{(1)} \neq 0$ and the system is in nonequilibrium. In numerical calculations, we set $(\Delta \mu_{AB} - \Delta \mu - \Delta \mu_{AX}) \neq 0$.